Melatonin alleviates acid-induced stress in peanut (*Arachis hypogaea* L.) through physiological, antioxidative regulation, and molecular defense mechanisms

Muhammad Hafeez Ullah Khan[1], Ruoyi Fu[1], Ali Muhammad[2], Shuaichao Zheng[1], Daijing Zhang[2], Zhiyong Zhang[1]*, Quanyong Liu[1]*

[1]Xinxiang Key Laboratory of Crop Root Biology and Green Efficient Production/Henan Key Laboratory of Quality and Stress-Resilient Bio-breeding and Safe Production of Cotton and Oil Crops, School of Life Sciences, Henan Institute of Science and Technology, Xinxiang, Henan, 453000, China

[2]College of Life Sciences, Henan Normal University, Xinxiang 453007, Henan, China.

Correspondence Author:

Zhiyong Zhang*

(Email: z_zy123@126.com)

Muhammad Hafeez Ullah Khan; hafeezkhan_88@yahoo.com, Ruoyi Fu; fu_ruoyi0123@163.com, Ali Muhammad; alikhaksar87@gmail.com, Shuaichao Zheng; yndxzsc@126.com, Daijing Zhang; zdjdai@163.com, Zhiyong Zhang; z_zy123@126.com, Quanyong Liu; liuquanyong@hist.edu.cn.

Abstract

Acidic stress severely restricts crop growth by disrupting nutrient uptake, redox homeostasis, and membrane stability, yet mitigation strategies remain limited. Here, we investigated the role of melatonin (MT) in modulating growth, photosynthesis, oxidative stress, antioxidant defense, and proton transport in peanut seedlings grown under acidic (pH 4.0) and near-optimal (pH 6.5) conditions. Acid stress markedly reduced biomass accumulation, chlorophyll content, and redox balance, while enhancing ROS (H_2O_2) and lipid peroxidation (MDA). Exogenous MT application, particularly at 50-100 µM, significantly improved shoot and root fresh and dry weights, restored chlorophyll a, b, total chlorophyll, and carotenoids, and reduced H_2O_2 and MDA accumulation, with more pronounced effects under pH 4.0 than pH 6.5. MT strongly activated antioxidant enzymes (SOD, CAT, APX), while POD activity declined, reflecting melatonin's dual role as both a direct ROS scavenger and a regulator of enzymatic redox networks. Notably, MT induced strong, dose-dependent upregulation of H^+-ATPase genes (AH1 and AH2) in both leaves and roots under acidic conditions, suggesting enhanced proton extrusion, intracellular pH homeostasis, and stress adaptation. Although MT also conferred benefits at pH 6.5, responses were generally moderate compared with acid stress. Collectively, these results demonstrate that MT confers robust acid stress tolerance in peanut by integrating growth regulation, antioxidant defense, and proton homeostasis. This study provides the first integrated evidence that melatonin enhances acid stress tolerance by coordinating antioxidant defense with H^+-ATPase-mediated proton homeostasis, revealing a novel mechanism for crop adaptation to low-pH soils.

Keywords: Melatonin supplementation, Peanut seedling resilience, Acidic soil adaptation, ROS-pH coordination, Low-pH crop physiology

Introduction

Soil acidification has become an escalating global concern, now affecting nearly 30–40% of the world's arable lands and posing a serious threat to agricultural productivity and food security (Huang *et al.*, 2023). Soils with a pH below 5.5 are considered acidic, and such conditions limit the availability of essential macronutrients like phosphorus (P), calcium (Ca^{2+}), and magnesium (Mg^{2+}). At the same time, they increase the solubility of toxic metal ions, especially aluminium (Al^{3+}) and manganese (Mn^{2+}), which can be highly detrimental to plant growth (Delfim *et al.*, 2025). This imbalance in nutrient and ion dynamics often results in inhibited root elongation, reduced nutrient uptake, and disrupted photosynthetic processes. Furthermore, an excess of hydrogen ions (H$^+$) in the rhizosphere disturbs pH stability and alters microbial community structure, compounding physiological stress in plants (Naz *et al.*, 2022; Thapa *et al.*, 2025).

Peanut (*Arachis hypogaea* L.), a major legume widely cultivated in tropical and subtropical regions, is particularly vulnerable to acid soil stress (Li *et al.*, 2024b; Jamir *et al.*, 2025). During early seedling stages, low pH and other abiotic stress conditions restrict root development, damage cell membranes, and interfere with ion transport mechanisms, ultimately impairing water and nutrient absorption. These physiological disruptions often lead to reduced chlorophyll levels, decreased photosynthetic efficiency, and poor biomass accumulation (Liu *et al.*, 2018; Borhannuddin Bhuyan *et al.*, 2019; Li *et al.*, 2021). On a cellular scale, acid stress intensifies the generation of reactive oxygen species (ROS), including superoxide anions (O$_2^-$), hydrogen peroxide (H$_2$O$_2$), and hydroxyl radicals (•OH), which can damage lipids, proteins, and nucleic acids (Ahmad *et al.*, 2023b; Hasan *et al.*, 2025). Although plants possess both enzymatic antioxidants (such as SOD, POD, APX, and CAT) and non-enzymatic compounds (like ascorbate, glutathione, and flavonoids). These defense mechanisms are often inadequate to neutralize the rapid oxidative surge induced by acidic environments, particularly in sensitive crops like peanut.

In recent years, melatonin (N-acetyl-5-methoxytryptamine) has been identified as an effective plant growth regulator and stress alleviator (Zheng *et al.*, 2025; Muhammad *et al.*, 2026). Melatonin was first found in animals, and is now widely recognized in plants to modulate plant growth, development and defence responses (Ahmad *et al.*, 2023a). Exogenous melatonin has been reported to alleviate various

abiotic stresses such as drought, salinity, acid, extreme temperature, nutrient deficiency and heavy metal toxicity by enhancing photosynthetic efficiency, maintaining redox balance and elevating antioxidant capacity (Basharat *et al.*, 2025; Dou *et al.*, 2025; Kong *et al.*, 2025). Melatonin is also involved in alleviating aluminium toxicity under suppressed soil pH, maintaining the integrity of cellular membrane, stabilizing chloroplast structure and activating the antioxidant system (Wang *et al.*, 2025a). In addition to its direct ROS-scavenging ability, melatonin indirectly regulates redox homeostasis by modulating the expression of redox-responsive genes (Hasan et al., 2025; Jiang *et al.*, 2025a).

Proton pumps, particularly plasma membrane and vacuolar H-ATPases, are fundamental molecular systems that enable plants to sustain cytoplasmic pH homeostasis under an acidic environment (Cosse & Seidel, 2021; Zeng *et al.*, 2024; MacDonald *et al.*, 2025). These ATPases actively transport protons from the cytoplasm into the apoplast or vacuole, thereby establishing electrochemical gradients that drive essential physiological processes such as nutrient uptake, ion transport, stomatal regulation and stress adaptation (MacDonald et al., 2025; Sun *et al.*, 2025). In peanut, various ATPase genes have been characterized, among which AH1 and AH2 are key regulators of intracellular pH balance and ion homeostasis under acid stress. The upregulation of H-ATPase genes has been reported in several crops as an adaptive mechanism to low pH, facilitating proton extrusion and protecting cellular metabolism (Li *et al.*, 2022; Li *et al.*, 2024a). However, the regulatory dynamics of these genes under melatonin treatment remain largely unexplored in the peanut seedling.

Melatonin has been shown in other crop species to influence proton pump activity and ion transport. For instance, melatonin-induced activation of H-ATPases can enhance nutrient absorption, stabilize membrane potential, and promote root development under stress conditions (Huang *et al.*, 2022; Tahjib-Ul-Arif *et al.*, 2025). Therefore, integrating physiological, biochemical and transcriptional analysis, particularly through expression profiling of AH1 and AH2, can provide valuable mechanistic insight into how melatonin supports pH regulation and overall homeostasis in peanut seedlings under acidic conditions.

The present study aims to elucidate the integrated physiological, antioxidative, and molecular mechanisms underlying melatonin-mediated acid tolerance in peanut seedlings, with a particular focus on AH1 and AH2 gene expression. We demonstrate

that melatonin enhances peanut tolerance to acid stress by improving growth and

photosynthetic traits, strengthening antioxidant defense, and upregulating key H⁺-

ATPase genes to stabilize cellular pH and ion homeostasis. Unravelling these

mechanisms will contribute to the development of sustainable agronomic strategies

and genetic improvement of peanut for acid soil environments.

Materials and methods

Plant material and experimental design

The experiment was conducted in the greenhouse (25 ± 2°C, 60-70% RH, 16/8 h

light/dark, 200 μmol m⁻² s⁻¹) at the Henan Institute of Science and Technology,

Xinxiang City, Henan Province, China, under controlled environmental conditions. A

paper roll culture system was used in a hydroponic setup to evaluate the effects of

melatonin on peanut growth under acidic stress conditions. The experiment followed a

randomized complete design (RCD) with eight biological replications to ensure

statistical reliability.

Uniform, mature, and healthy seeds of peanut (*Arachis hypogaea* L. cv.

Xinbaihua 25) were selected as experimental material. Seeds were surface-sterilized

with 1% sodium hypochlorite solution for 5 min and rinsed several times with

distilled water to remove any residual disinfectant. To investigate the effects of

melatonin on plant growth, photosynthetic pigments, antioxidant responses, and gene

expression, the seeds were soaked for 10 hours in melatonin solutions of 0.5 μM, 5

μM, 50 μM, and 100 μM. Seeds soaked in distilled water under identical conditions

served as the control (CK).

For each replicate, seven seeds were placed vertically on moistened

germination paper to ensure uniform orientation and root emergence. The papers were

then carefully rolled and secured with rubber bands to maintain contact and prevent

unrolling during incubation. Following treatment, the paper rolls containing the seeds

were placed vertically in plastic containers filled with half-strength Hoagland nutrient

solution prepared according to the modified composition described by Cai et al. (2021)

(Cai *et al.*, 2021). Two nutrient pH regimes were maintained: a normal condition with

pH 6.5 and an acidic condition with pH 4.0. The pH of each solution was carefully

adjusted using 1 M HCl or NaOH and monitored daily to ensure consistent stability

throughout the experiment. The lower ends of the paper rolls were immersed

approximately 2-3 cm into the solution to facilitate the upward capillary movement of

nutrients and water while maintaining adequate aeration around the root zone. To prevent nutrient depletion and pH fluctuations, the nutrient solutions were replaced every two days. During the early germination stage, the paper rolls were kept under dark conditions until radicle emergence to ensure uniform germination. After emergence, seedlings were exposed to normal light conditions for continued growth. After 15 days of cultivation, seedlings were harvested for measurements of growth parameters, photosynthetic pigments, antioxidant enzyme activities, and gene expression analyses.

Plant Germination and Growth Parameters

Seed germination was monitored daily until radicle emergence, which was defined as a visible protrusion of at least 2 mm from the seed coat, and the germination percentage (GP) was calculated at day 7. After 15 days of cultivation, seedlings were carefully removed from the paper rolls and gently rinsed with distilled water to remove residual nutrient solution. Shoot length was measured from the base to the tip of the longest leaf, and root length from the base to the tip of the longest root, using a ruler to the nearest millimeter. Fresh weights of shoots and roots were determined separately after blotting with filter paper to remove surface moisture, and samples were then oven-dried at 70 °C for 72 h or until a constant weight was achieved to determine dry weights. All measurements were performed for each seedling in every replicate, and the mean values of the eight replicates were used for statistical analysis. Data are presented as mean ± standard error (SE) and used for subsequent physiological, biochemical, and molecular analyses.

Photosynthetic Pigment Measurement

Photosynthetic pigments, including chlorophyll a, chlorophyll b, and carotenoids, were extracted from 0.2 g of fresh peanut leaves using 80% (v/v) acetone. The leaf samples were homogenized in chilled acetone, kept on ice for 10-15 min in the dark, and centrifuged at $10,000 \times g$ for 10 min at 4 °C. The clear supernatant was collected, and absorbance was recorded at 663, 645, and 470 nm using a UV–visible spectrophotometer with 80% acetone as blank. Pigment contents were calculated using standard equations and expressed as mg g^{-1} fresh weight. All extractions were performed under low-light conditions to minimize pigment degradation, and six independent replicates were analyzed per treatment.

Antioxidant Enzyme Assays

To assess the antioxidant defense response in peanut seedlings, the activities of catalase (CAT), superoxide dismutase (SOD), peroxidase (POD), and ascorbate peroxidase (APX), content, were determined from fresh leaf tissues. Approximately 0.5 g of fresh leaves was homogenized in 5 mL of ice-cold 50 mM phosphate buffer (pH 7.0) containing 1 mM EDTA and 1% (w/v) polyvinylpyrrolidone (PVP). The homogenate was centrifuged at 12,000 × g for 15 min at 4 °C, and the resulting supernatant was used for all enzyme assays.

Superoxide dismutase (SOD; EC 1.15.1.1) activity was assayed by the inhibition of nitroblue tetrazolium (NBT) reduction under light, with one unit of SOD activity defined as the amount of enzyme required to cause 50% inhibition of NBT photoreduction. Catalase (CAT; EC 1.11.1.6) activity was determined by monitoring the decrease in absorbance at 240 nm due to H_2O_2 decomposition. Peroxidase (POD; EC 1.11.1.7) activity was measured using guaiacol as a substrate, recording the increase in absorbance at 470 nm as guaiacol was oxidized. Ascorbate peroxidase (APX; EC 1.11.1.11) activity was estimated by monitoring the decrease in absorbance at 290 nm as ascorbate was oxidized in the presence of H_2O_2.

Quantification of Malondialdehyde (MDA) and Hydrogen Peroxide (H_2O_2)

The levels of malondialdehyde (MDA) and hydrogen peroxide (H_2O_2) were determined to evaluate lipid peroxidation and oxidative stress in peanut leaves and roots. Lipid peroxidation was determined by quantifying malondialdehyde (MDA) using the thiobarbituric acid (TBA) reaction. For MDA estimation, 1 mL of enzyme extract was mixed with 2 mL of 0.6% (w/v) TBA in 10% (w/v) trichloroacetic acid (TCA) and heated at 95 °C for 15 min, then rapidly cooled and centrifuged at 10,000 × g for 10 min. The absorbance of the supernatant was measured at 532 nm and corrected for nonspecific turbidity by subtracting absorbance at 600 nm. MDA content was calculated using the extinction coefficient ($\varepsilon = 155$ mM^{-1} cm^{-1}) and expressed as nmol g^{-1} fresh weight (FW).

Hydrogen peroxide (H_2O_2) content was determined spectrophotometrically using a colorimetric KI reaction with quantification against a standard curve. Fresh leaf tissue (0.5 g) was ground in 5 mL of cold 0.1% (w/v) TCA and centrifuged at 12,000 × g for 15 min at 4 °C. A reaction mixture was prepared by mixing 0.5 mL of the extract with 0.5 mL of 10 mM potassium phosphate buffer (pH 7.0) and 1.0 mL of 1 M potassium iodide (KI) (all kept on ice and protected from light). The reaction was allowed to

proceed for 1 min at room temperature in the dark, and the absorbance was read at
390 nm. A standard calibration curve was prepared in parallel using known
concentrations of H_2O_2 (for example 0, 10, 25, 50, 100, 200 µM) prepared in 0.1%
TCA; sample H_2O_2 content was interpolated from this curve and expressed as µmol
H□O□ g□¹ FW (adjusting for extract and assay volumes and any dilutions). All
reactions and standards were run in technical triplicate and blank reactions (reagents
without plant extract) were included to correct baseline absorbance.

Gene Expression Analysis

Total RNA was extracted from peanut roots and leaves using the RNAprep Pure Plant
Plus Kit (Tiangen Biotech, Beijing, China) according to the manufacturer's
instructions. The quality and quantity of RNA were determined using a NanoDrop
2000 spectrophotometer (Thermo Fisher Scientific, USA) by measuring the
absorbance ratios at 260/280 nm and 260/230 nm. RNA integrity was confirmed by
1.0% agarose gel electrophoresis, ensuring that samples displayed sharp and intact
28S and 18S rRNA bands without visible degradation. Only high-quality RNA
(A260/A280 ratio between 1.9 and 2.1) was used for cDNA synthesis.

First-strand cDNA was synthesized from 50 µg of total RNA using the
PrimeScript™ RT reagent kit with gDNA Eraser (Takara, Japan) to eliminate potential
genomic DNA contamination. The synthesized cDNA was diluted 10-fold with
nuclease-free water and stored at -20 °C until further analysis.

Quantitative real-time PCR (qRT-PCR) was conducted using TB Green®
Premix Ex Taq™ II (Takara, Japan) on a QuantStudio™ 5 Real-Time PCR System
(Applied Biosystems, USA). Each 20 µL reaction contained 10 µL of 2× SYBR Green
master mix, 0.4 µL each of forward and reverse primers (10 µM), 2 µL of diluted
cDNA, and 7.2 µL of nuclease-free water. The PCR cycling conditions were as
follows: initial denaturation at 95 °C for 30 s, followed by 40 amplification cycles of
95 °C for 5 s and 60 °C for 30 s. A melting curve analysis (60-95 °C) was performed
at the end of each run to confirm the specificity of amplification.

Each sample was analyzed in triplicate, and the mean values of three technical
replicates from three independent biological replicates were used for statistical
analysis. The relative transcript abundance of target genes was calculated using the
2□ΔΔCt method (Livak and Schmittgen, 2001). The AhGAPDH gene was used as an

internal reference due to its stable expression across different tissues and treatments.

Primer sequences used in the study are listed in Supplementary Table 1.

Statistical Analysis

All data were analyzed using SPSS v26.0. Results are presented as mean ± SE. Two-

way ANOVA was used to compare treatments, and significant differences were

determined using Tukey's test ($p < 0.05$). For qRT-PCR, the average Ct values of

technical replicates were used, and fold changes were calculated using the $2^{-\Delta\Delta Ct}$

method.

Table 1. Primer sequences used for qRT-PCR analysis of plasma membrane H$^+$-

ATPase genes in peanut.

Gene Name	Primer Sequence
AH1 H+- ATPase 1-F	GCCTCTGACATAGTTCTAAC
AH1 H+- ATPase 1-R	TAAAGCCAACCACAATACGG
AH2 H+- ATPase 2-F	GAAAAGAATGGGCGTGGTTG
AH2 H+- ATPase 2-R	TCAAGGGACACAATCTCACC

The table lists forward (F) and reverse (R) primer sequences designed for the
amplification of *AhAH1* and *AhAH2* genes.

Results

Effect of melatonin on peanut seed germination under acid condition

Seed germination of peanut was significantly affected by pH and melatonin treatment

(Fig. 1a and b). At acidic pH (4), control seeds exhibited the lowest germination

(~70%). Application of MT markedly improved germination in a dose-dependent

manner: 0.5 µM MT increased germination to ~88% (+18%), 5 µM MT to ~90%

(+20%), 50 µM MT to ~93% (+23%), and 100 µM MT to ~98% (+28%) with related

to control, indicating a strong protective effect of MT under acid stress. Under near-

neutral conditions (pH 6.5), germination was consistently high across treatments, with

control seeds reaching ~85% and MT-treated seeds ranging from 90% to nearly 100%.

The relative improvement of germination by MT was more pronounced under acidic

stress, indicating that MT alleviates pH-induced inhibition. These results demonstrate

that melatonin significantly mitigates the inhibitory effects of low pH on peanut seed

germination, with the largest relative benefit observed at 100 µM MT under acidic

conditions, and that moderate MT concentrations are sufficient to maintain near-maximal germination at near-neutral pH.

Melatonin supplementation enhances peanut plant growth and biomass accumulation under acid stress

To evaluate the role of melatonin in modulating peanut seedling growth under contrasting pH conditions, root and shoot development as well as biomass accumulation were assessed under acidic (pH 4) and near-neutral (pH 6.5) environments (Figs. 1c–h). Acidic stress (pH 4) markedly suppressed seedling growth, as evidenced by substantial reductions in root and shoot elongation in control plants, indicating strong inhibition of early developmental processes. Melatonin treatment markedly alleviated this inhibition, resulting in a dose-dependent enhancement of root length, shoot length, and biomass. Root length under pH 4 increased progressively from approximately 6 cm in control plants to nearly 20 cm at 100 µM MT, while shoot length increased from about 2.5 cm in the control to over 6 cm at the highest MT concentration, demonstrating a strong stimulatory effect of MT on seedling elongation under acid stress (Fig. 1c and d). Under neutral conditions (pH 6.5), root and shoot lengths were generally higher than those at pH 4 in control plants; however, the growth-promoting effect of MT was less pronounced. Low MT concentration (0.5 µM) enhanced root and shoot length, while higher MT doses showed either moderate improvement or stabilization of growth, suggesting that MT exerts a stronger relative effect under stress conditions than under optimal pH (Fig. 1c and d).

Consistent with growth responses, root fresh weight and shoot fresh weight under pH 4 increased substantially with increasing MT concentration. Root fresh weight rose from 1.09 g in control plants to 3.45 g at 100 µM MT, while shoot fresh weight increased from 1.09 g to 2.86 g, indicating enhanced biomass accumulation (Fig. 1e and f). This trend was further supported by dry weight measurements, where root dry weight increased from 0.07 g in control plants to 0.17 g at 100 µM MT, and shoot dry weight increased from 0.06 g to 0.15 g, confirming that MT-induced growth was associated with true biomass gain rather than water accumulation alone (Fig. 1g and h).

Under pH 6.5, overall growth and biomass parameters were higher than those observed under acidic conditions in the control plants; however, the response pattern

to MT differed from that at pH 4. Low MT concentration (0.5 µM) promoted root and

shoot growth and fresh weight accumulation, whereas higher MT concentrations (5-

100 µM) resulted in a gradual decline in root and shoot length and biomass (Fig. 1e–

h). Notably, root dry weight and shoot dry weight at pH 6.5 showed little or no

improvement with increasing MT concentration and tended to decrease at higher

doses, suggesting a concentration-dependent shift in MT responsiveness under non-

stress conditions.

Overall, these results demonstrate that melatonin strongly enhances root and

shoot growth and biomass accumulation under acidic stress (pH 4), with maximal

effects observed at higher MT concentrations, whereas under near-neutral conditions

(pH 6.5), low MT levels are sufficient to promote growth and excessive MT may exert

inhibitory effects. This indicates that the growth-promoting role of melatonin is highly

dependent on environmental pH and stress intensity.

Figure 1. Effect of melatonin (MT) on seed germination and seedling growth under normal (pH 6.5) and acidic (pH 4.0) conditions on peanut. (a) seed germination and seedling growth phenotypes, (b) seed germination (%), (c) shoot length (cm), (d) root length (cm), (e) shoot fresh weight (g), (f) root fresh weight (g), (g) root dry weight (g), (h) shoot dry weight. Statistical differences among treatments were analyzed using appropriate multiple comparison tests. ns indicates not significant, while *, **, ***, and **** denote significant

346   differences at P < 0.05, P < 0.01, P < 0.001, and P < 0.0001, respectively. Data in the figures

347   are means of eight replications (n = 8) ± standard deviation (SD). Bar = 2 cm.

Melatonin modulates photosynthetic pigments in response to acid stress in

peanut

Melatonin application significantly influenced the accumulation of photosynthetic

pigments in peanut seedlings under both acidic (pH 4) and near-neutral (pH 6.5)

conditions (Figs. 2a-d). Under pH 4, control plants exhibited relatively low levels of

chlorophyll a, chlorophyll b, total chlorophyll, and carotenoids, indicating impaired

pigment biosynthesis under acid stress. However, MT treatments progressively

enhanced pigment accumulation, with a clear dose-dependent response.

Chlorophyll a content under pH 4 increased gradually with increasing MT

concentration, showing moderate enhancement at 5 µM MT and a pronounced

increase at 50 and 100 µM MT, where values reached their maximum (Fig. 2a). A

similar trend was observed for chlorophyll b, resulting in a substantial rise in total

chlorophyll content, particularly at 50 and 100 µM MT. These increases were

significantly higher than the control (P < 0.05), demonstrating that MT effectively

mitigates acid-induced inhibition of chlorophyll synthesis (Fig. 2b and c). Under pH

6.5, pigment contents were generally higher than those under pH 4 in control plants,

reflecting more favorable growth conditions. Notably, low MT concentration (0.5 µM)

caused a marked increase in chlorophyll a, chlorophyll b, and total chlorophyll,

reaching peak values under this pH. However, higher MT concentrations (50-100 µM)

resulted in a gradual decline in pigment levels, suggesting that excessive MT under

optimal pH may suppress chlorophyll accumulation (Figs. 2a-c).

Carotenoid content showed a distinct but complementary pattern. Under pH 4,

carotenoid levels increased steadily with rising MT concentration, reaching the

highest values at 100 µM MT, which were significantly greater than the control (P <

0.05). This indicates enhanced antioxidant capacity under acid stress (Fig. 2d). In

contrast, under pH 6.5, carotenoid content peaked sharply at 0.5 µM MT and declined

at higher concentrations, paralleling the response observed for chlorophyll pigments

(Fig. 2d). Overall, these results demonstrate that melatonin enhances photosynthetic

pigment accumulation in peanut seedlings in a pH-dependent and concentration-

dependent manner. Low MT concentrations are optimal for pigment enhancement

under near-neutral conditions, whereas higher MT doses are more effective in

protecting pigment biosynthesis under acidic stress, likely through improved stress

tolerance and antioxidant regulation.

Figure 2. Effect of melatonin (MT) on photosynthetic pigments of peanut under normal

(pH 6.5) and acidic (pH 4.0) conditions. (a) Chlorophyll *a* (mg/g FW), **(b)** chlorophyll *b*

(mg/g FW), **(c)** total chlorophyll content (mg/g FW), **(d)** carotenoids (mg/g FW). Statistical

differences among treatments were analyzed using appropriate multiple comparison tests. ns

indicates not significant, while ***, and **** denote significant differences at P < 0.001, and

P < 0.0001, respectively. Data in the figures are means of eight replications (n = 8) ± standard

deviation (SD).

Melatonin differentially regulates antioxidant defense system in response

to acid stress in peanut

We evaluated the enzymatic antioxidant activity in acid-stressed peanut plants treated

with melatonin to understand how exogenous melatonin mitigates acid stress. Our

findings revealed that the radical scavenging activities of SOD, POD, CAT, and APX,

were markedly affected by both external pH and melatonin treatment in peanut leaves,

reflecting dynamic regulation of reactive oxygen species (ROS) homeostasis under

stress and non-stress conditions (Figs. 3a-h).

Superoxide Dismutase (SOD) Activity

SOD activity showed a strong induction under acidic conditions (pH 4) compared with pH 6.5 across all treatments, indicating enhanced superoxide radical generation under acid stress. In control plants, SOD activity at pH 4 was substantially higher (60.79 U mg^{-1} FW) than at pH 6.5 (18.70 U mg^{-1} FW), reflecting a basal stress response. Application of melatonin significantly increased SOD activity in a dose-dependent manner, with 100 μM MT reaching 137.1 U mg^{-1} FW, representing a ~128% increase over control, suggesting that MT promotes superoxide detoxification capacity under severe acidic stress (Fig. 3a). In contrast, under pH 6.5, SOD activity was relatively low and showed only moderate induction by MT, with a peak at 0.5 μM MT followed by a decline at higher concentrations (Fig. 3a). This differential response indicates that melatonin enhances ROS-scavenging systems primarily under stress conditions, while under near-optimal pH it avoids excessive activation of antioxidant machinery, thereby maintaining redox balance.

In roots, SOD activity exhibited a similar but more pronounced response. At pH 4, melatonin increased SOD activity from 54.34 U mg^{-1} FW (control) up to 119.39 U mg^{-1} FW at 100 μM MT (~126% increase). This strong induction indicates that MT substantially enhances superoxide detoxification in roots exposed to acidic stress (Fig. 3b). At pH 6.5, the increase was smaller but still significant (18.70 U mg^{-1} FW control to 42.29 U mg^{-1} FW at 0.5 μM MT, ~126% increase), however, MT application still increased SOD relative to control, though to a lesser extent than under acidic conditions (Fig. 3b). These results suggest that MT-mediated regulation of SOD is stress-responsive and tissue-specific, with a stronger effect in roots facing direct exposure to low-pH environments.

Peroxidase (POD) Activity

POD activity exhibited an opposite trend compared with SOD, particularly under acidic conditions. At pH 4, control plants showed the highest POD activity (175.53 U mg^{-1} FW), reflecting stress-induced activation of cell wall-associated and vacuolar peroxidases, which are often involved in stress signaling, lignification, and ROS-mediated defense responses. Melatonin application led to a progressive decrease in POD activity, with 100 μM MT reducing activity to 29.24 U mg^{-1} FW (~83% reduction compared to control). This decline suggests that melatonin alleviates oxidative stress, thereby reducing the requirement for POD-mediated hydrogen

peroxide consumption (Fig. 3c). Importantly, this reduction highlights the dual role of melatonin: while it enhances ROS scavenging through SOD, CAT, and APX, it simultaneously suppresses excessive POD activity that is often associated with oxidative damage, rigidification of cell walls, and growth inhibition. Under pH 6.5, POD activity was generally lower than at pH 4 and displayed a distinct pattern, with moderate increases at higher MT concentrations (Fig. 3c). This suggests that under non-stress conditions, POD may participate more in fine-tuning redox signaling rather than bulk ROS detoxification.

In roots, POD activity displayed a similar pattern. At pH 4, POD activity also decreased from 168.59 U mg⁻¹ FW in control to 35.63 U mg⁻¹ FW at 100 µM MT (~79% reduction), mirroring the trend observed in leaves and supporting the role of melatonin in alleviating oxidative stress by reducing H_2O_2 accumulation (Fig. 3d). However, at pH 6.5, POD activity increased progressively with melatonin concentration, reaching near-control or higher levels at 100 µM MT (Fig. 3d). This contrasting behavior highlights the dual role of melatonin: under stress, it suppresses excessive ROS-scavenging enzyme activity by lowering ROS production, whereas under non-stress conditions, it maintains or enhances POD activity to support redox homeostasis and signaling.

Catalase (CAT) Activity

CAT activity was strongly stimulated by melatonin under both pH conditions, though the magnitude and pattern of induction differed. Under acidic pH, CAT activity increased from 139.4 U mg⁻¹ FW in control to 273.2 U mg⁻¹ FW at100 µM MT (~96% increase over control) (Fig. 3e). This indicates that melatonin enhances the rapid decomposition of hydrogen peroxide into water and oxygen, a crucial mechanism for preventing oxidative damage under acid-induced stress. At pH 6.5, CAT activity was generally higher than control and showed a pronounced induction at low to moderate MT concentrations, with the highest activity observed at 0.5 µM MT, followed by a gradual decline at higher doses (Fig. 3e). This suggests that under near-neutral pH, lower MT concentrations are sufficient to optimize hydrogen peroxide homeostasis, while excessive MT may reduce the need for CAT due to improved redox equilibrium.

In roots, CAT activity similarly increased under acidic stress, from 167.6 U mg⁻¹ FW (control) to 331.6 U mg⁻¹ FW at 100 µM MT (~98% increase) (Fig. 3f).

At near-neutral pH, control activity was 147.9 U mg^{-1} FW, and melatonin slightly

increased CAT at low concentrations (0.5-5 µM MT) but decreased at higher doses

(100 µM MT, 147.2 U mg^{-1} FW) (Fig. 3f). These results indicate that melatonin

effectively boosts CAT-mediated H_2O_2 scavenging, especially under acid stress,

enhancing antioxidant defense in both leaves and roots.

Ascorbate Peroxidase (APX) Activity

APX activity increased consistently with melatonin concentration under pH 4,

indicating a strong MT-mediated activation of the ascorbate-glutathione cycle under

acidic stress. At pH 4, control leaves had 28.8 U mg^{-1} FW, which rose to 66.9 U

mg^{-1} FW at 100 µM MT (~132% increase) (Fig. 3g). The highest APX activity at

100 µM MT suggests that APX plays a central role in fine-scale hydrogen peroxide

detoxification in chloroplasts and cytosol when plants experience acid-induced

oxidative pressure. Under pH 6.5, APX activity showed a different pattern, increased

from 26.8 U mg^{-1} FW (control) to 62.9 U mg^{-1} FW at 0.5 µM MT, while higher MT

concentrations showed slightly reduced activity (~34.7–51.8 U mg^{-1} FW), indicating

a dose-dependent modulation (Fig. 3g). This response implies that under optimal

conditions, melatonin primarily enhances APX activity at low doses to maintain

signaling-compatible ROS levels, while higher concentrations may downregulate

APX as oxidative stress is already minimal.

APX activity in roots increased significantly with MT under acidic conditions,

from 18.3 U mg^{-1} FW in control to 63.3 U mg^{-1} FW at 100 µM MT (Fig. 3h). At pH

6.5, APX activity peaked at moderate MT doses (66.2 U mg^{-1} FW at 0.5 µM), while

higher MT doses slightly reduced activity (Fig. 3h), showing that APX is finely

regulated by both MT concentration and pH to balance ROS detoxification with

normal metabolism.

Collectively, these results demonstrate that melatonin orchestrates a

coordinated antioxidant response in peanut leaves by enhancing ROS-scavenging

enzymes (SOD, CAT, APX) while downregulating stress-associated POD activity,

particularly under acidic conditions. This dual regulatory role allows melatonin to

mitigate oxidative damage while preserving ROS signaling necessary for growth and

adaptation. The contrasting responses observed between pH 4 and pH 6.5 further

highlight that melatonin acts as a stress-dependent modulator of redox homeostasis

rather than a constitutive antioxidant, enabling peanut plants to adapt efficiently to acidic stress environments.

Figure 3. Effect of melatonin (MT) on antioxidant contents of peanut under normal (pH 6.5) and acidic (pH 4.0) conditions. (a) leaf SOD (U/g FW), (b) leaf POD ((U/g FW), (c) leaf CAT (U/g FW), (d) leaf APX (U/g FW), (e) root SOD (U/g FW), (f) root POD ((U/g FW), (f) root CAT (U/g FW), (g) root APX (U/g FW). Statistical differences among treatments were analyzed using appropriate multiple comparison tests. ns indicates not significant, while **,

, and * denote significant differences at P < 0.01, P < 0.001, and P < 0.0001, respectively. Data in the figures are means of eight replications (n = 8) ± standard deviation (SD).

Melatonin reduces hydrogen peroxide and lipid peroxidation contents in peanut under acidic condition

Melatonin is widely recognized for its role in mitigating oxidative stress across various biological systems, including plants. We conducted a comprehensive study to assess how melatonin influences peanut seedling growth under acidic condition. Our findings revealed that acid-treated plants exhibited significantly elevated levels of , H_2O_2, and MDA compared to MT treated plants (Figs. 4a-d)

Hydrogen Peroxide (H_2O_2) Content

Under control conditions (CK), H_2O_2 levels were high at pH 4 (79.40 µmol g⁻¹ FW) and pH 6.5 (71.06 µmol g⁻¹ FW), indicating substantial oxidative stress, particularly under acidic conditions. Melatonin application significantly reduced H_2O_2 accumulation at pH 4 in a dose-dependent manner. Compared with the control, 0.5 µM MT reduced H_2O_2 to 57.95 µmol g⁻¹ FW (≈28% decrease), 5 µM MT to 49.28 µmol g⁻¹ FW (≈38% decrease), 50 µM MT to 46.31 µmol g⁻¹ FW (≈43% decrease), and 100 µM MT to 39.99 µmol g⁻¹ FW (≈50% decrease) (Fig. 4a). In contrast, at pH 6.5, low MT concentrations initially decreased H_2O_2, with 0.5 µM MT reducing levels to 44.38 µmol g⁻¹ FW (≈36% decrease) and 5 µM MT to 53.03 µmol g⁻¹ FW (≈26% decrease). However, higher MT doses led to a reversal of this trend, with H_2O_2 increasing to 62.34 µmol g⁻¹ FW at 50 µM MT (≈11% decrease) and 68.71 µmol g⁻¹ FW at 100 µM MT (nearly comparable to control) (Fig. 4a). This suggests a dose and pH-dependent dual role of melatonin, acting as a strong antioxidant under acidic stress but allowing moderate ROS signaling at optimal pH.

Root tissues showed a similar but more pronounced response. Under pH 4, control plants accumulated 85.04 µmol g⁻¹ FW H_2O_2. Melatonin significantly lowered H_2O_2 to 64.25 µmol g⁻¹ FW at 0.5 µM (≈24% reduction), 58.54 µmol g⁻¹ FW at 5 µM (≈32% reduction), 48.76 µmol g⁻¹ FW at 50 µM (≈44% reduction), and 43.81 µmol g⁻¹ FW at 100 µM MT (≈51% reduction) (Fig. 4b). At pH 6.5, H_2O_2 content in roots declined at low MT doses (44.25 µmol g⁻¹ FW at 0.5 µM; ≈46% decrease) but increased at higher concentrations, reaching 74.78 µmol g⁻¹ FW at 100 µM MT (≈9% decrease compared with control) (Fig. 4b). These results indicate that

melatonin efficiently scavenges excessive ROS under acid stress, while at neutral pH high MT may permit ROS accumulation for signaling functions.

Malondialdehyde (MDA) Content

MDA content, an indicator of lipid peroxidation, was markedly higher under acidic conditions. In leaves at pH 4, control plants exhibited 12.35 nmol g^{-1} FW MDA. Melatonin treatments progressively reduced MDA levels to 8.29 nmol g^{-1} FW at 0.5 µM (≈32% reduction), 6.38 nmol g^{-1} FW at 5 µM (≈48% reduction), 4.94 nmol g^{-1} FW at 50 µM (≈60% reduction), and 3.89 nmol g^{-1} FW at 100 µM MT (≈69% reduction) (Fig. 4c). At pH 6.5, control MDA content was lower (9.0 nmol g^{-1} FW). Low MT concentrations further decreased MDA to 3.92 nmol g^{-1} FW at 0.5 µM (≈56% reduction) and 5.52 nmol g^{-1} FW at 5 µM (≈39% reduction). However, higher MT doses increased MDA to 6.66 nmol g^{-1} FW at 50 µM (≈26% reduction) and 7.83 nmol g^{-1} FW at 100 µM MT (≈12% reduction) (Fig. 4c), again reflecting melatonin's concentration-dependent dual behavior.

Root MDA followed trends similar to leaves but with overall lower values. At pH 4, control roots had 11.16 nmol g^{-1} FW MDA. This decreased to 9.04 nmol g^{-1} FW at 0.5 µM (≈19% reduction), 7.15 nmol g^{-1} FW at 5 µM (≈35% reduction), 5.93 nmol g^{-1} FW at 50 µM (≈46% reduction), and 3.65 nmol g^{-1} FW at 100 µM MT (≈68% reduction) (Fig. 4d). At pH 6.5, MDA declined sharply at low MT doses (3.83 nmol g^{-1} FW at 0.5 µM; ≈64% reduction) but increased again at higher concentrations, reaching 8.25 nmol g^{-1} FW at 100 µM MT (≈20% reduction compared with control) (Fig. 4d). This indicates that melatonin effectively protects membrane integrity under acidic stress, while excessive doses at optimal pH may attenuate this protective effect.

Collectively, these results demonstrate that melatonin markedly alleviates oxidative stress and lipid peroxidation in peanut leaves and roots, particularly under acidic (pH 4) conditions. The strong reduction in H_2O_2 and MDA at pH 4 indicates that melatonin primarily functions as an antioxidant under stress. In contrast, under pH 6.5, low melatonin concentrations enhance oxidative protection, whereas higher doses partially restore ROS and lipid peroxidation levels, suggesting a regulated balance between ROS scavenging and signaling. This dual role of melatonin

highlights its importance in fine-tuning redox homeostasis in peanut plants under

contrasting pH environments.

Figure 4. Effect of melatonin (MT) on H$_2$O$_2$ and MDA contents of peanut under normal

(pH 6.5) and acidic (pH 4.0) conditions. (a) leaf H$_2$O$_2$ (µmol/g FW), (b) leaf MDA (nmol/g

FW), (c) root H$_2$O$_2$ (µmol/g FW), (d) root MDA (nmol/g FW). Statistical differences among

treatments were analyzed using appropriate multiple comparison tests. ns indicates not

significant, while *, **, and **** denote significant differences at P < 0.05, P < 0.01, and P <

0.0001, respectively. Data in the figures are means of eight replications (n = 8) ± standard

deviation (SD).

Melatonin induces dose-dependent upregulation of H☐-ATPase genes

under acid stress

To investigate the transcriptional response of plasma membrane H☐-ATPase genes to

melatonin (MT) under acidic conditions, the relative expression levels of *AhAH1* and

AhAH2 were analyzed in peanut leaves and roots under acid stress (Figs. 5a and b).

Both genes are central to proton extrusion and pH homeostasis during environmental

stress.

Melatonin treatment significantly enhanced the expression of plasma

membrane H☐-ATPase genes *AhAH1* and *AhAH2* in peanut seedlings, with a

pronounced induction under acidic conditions. The basal expression of *AhAH1* was

low in control plants, with minimal transcript accumulation observed in both leaves

and roots. MT supplementation resulted in a consistent and progressive increase in

AhAH1 expression at all concentrations tested. At 0.5 and 5 µM MT, *AhAH1* transcript

levels increased by approximately 2.4, and 2.6-fold in leaves and 1.5, and 2.5-fold in

roots relative to the control (Fig. 5a). This induction intensified with increasing MT

concentration, reaching ~3.7-fold in leaves and ~3.2-fold in roots at 50 µM MT. The

strongest response was observed at 100 µM MT, where *AhAH1* expression increased

by ~7.0-fold in leaves and ~5.1-fold in roots compared with control plants, indicating

robust, dose-dependent transcriptional activation (Fig. 5a). The stronger induction in

roots highlights the pivotal role of root-localized H□-ATPase activity in coping with

acidic stress.

A similar positive response was observed for *AhAH2*, which exhibited higher

basal expression than *AhAH1* in control plants, particularly in roots. Nevertheless, MT

treatment further enhanced *AhAH2* expression at all concentrations under acid stress.

At low MT concentrations (0.5 and 5 µM), *AhAH2* transcript abundance increased by

approximately 1.7- and 3.1-fold in leaves and 2.6- and 4.5-fold in roots, respectively,

relative to the control (Fig. 5b). This upregulation became progressively stronger with

increasing MT dosage, reaching approximately 5.1-fold in leaves and 8.2-fold in roots

at 50 µM MT. The maximum induction occurred at 100 µM MT, where *AhAH2*

expression was elevated by approximately 6.6-fold in leaves and 11.6-fold in roots

compared with control plants (Fig. 5b), demonstrating a robust and concentration-

dependent transcriptional response. Importantly, All MT treatments significantly

increased *AhAH2* transcript levels relative to the control, with the highest induction

observed at 100 µM MT. This consistent upregulation under acidic stress suggests that

AhAH2 is positively responsive to melatonin and likely contributes to enhanced

proton extrusion and pH homeostasis. The pronounced upregulation of *AhAH2* at

higher melatonin concentrations suggests that this gene encodes a stress-adaptive H□-

ATPase that remains transcriptionally active under strong melatonin stimulation,

thereby contributing to the maintenance of cytosolic pH homeostasis and sustained

root function under acidic stress conditions.

Overall, these results demonstrate that melatonin induces a consistent,

concentration-dependent increase in H□-ATPase gene expression at all MT

treatments relative to the control, with the strongest fold induction occurring under

acidic conditions. This transcriptional enhancement provides molecular evidence for

melatonin-mediated reinforcement of proton transport capacity, supporting improved

acid tolerance in peanut seedlings.

Figure 5. Melatonin-induced expression of plasma membrane H□-ATPase genes in

peanut leaves and roots. (a) Relative expression of *AhAH1* and (b) relative expression of

AhAH2 in leaves and roots under different melatonin (MT) concentrations (0.5, 5, 50, and 100

μM) compared with the control. Bars represent mean ± SD of biological replicates. Statistical

differences among treatments were analyzed using appropriate multiple comparison tests. ns

indicates not significant, while *, **, ***, and **** denote significant differences at P < 0.05,

P < 0.01, P < 0.001, and P < 0.0001, respectively.

Discussion

Melatonin alleviates acid stress-induced growth inhibition by promoting

seed germination and seedling growth in peanut

Acidic stress conditions severely constrain seed germination and early seedling

establishment by disrupting metabolic activation, reserve mobilization, and cell

expansion (Msimbira & Smith, 2020; Hasan et al., 2025). Seed germination and early

seedling establishment are among the most pH-sensitive developmental stages in

crops, and acidic stress is known to impair metabolic activation, reserve mobilization,

and cell expansion (Shavrukov & Hirai, 2016; Ali *et al.*, 2023; Wang *et al.*, 2025b). In

this study, peanut seeds exposed to acidic pH (4.0) showed a clear reduction in

germination percentage and seedling growth compared with near-neutral pH (6.5),

confirming the high pH sensitivity of early developmental stages. Exogenous

melatonin (MT) markedly alleviated this inhibition, significantly improving

germination and promoting vigorous seedling growth, with a much stronger relative

effect under acidic stress than under near-neutral conditions (Figs.1a-h). The enhanced

germination under acidic pH likely reflects improved metabolic activation during

imbibition, as MT is known to stimulate hydrolytic enzymes such as α-amylases and proteases, thereby facilitating efficient mobilization of stored reserves under stress (Liu *et al.*, 2023; Fu *et al.*, 2025). In addition, MT interacts with key phytohormones, including gibberellins and auxins, which regulate radicle protrusion and early seedling vigor, further supporting rapid germination under low-pH conditions (Arnao & Hernández-Ruiz, 2019; Fu et al., 2025). Consistent with improved germination, MT substantially enhanced root and shoot growth and biomass accumulation under acidic conditions, indicating improved seedling establishment (Figs.1a-h). Similar MT-mediated promotion of early growth under unfavourable soil conditions has been reported in other crops, where MT improved root development and stress resilience by maintaining cellular homeostasis and nutrient acquisition (Muhammad *et al.*, 2024; Chung *et al.*, 2025; Hasan et al., 2025). In contrast, under near-neutral pH, low MT doses were sufficient to slightly enhance seedling performance, whereas higher concentrations often reduced growth and physiological indices, highlighting the role of MT as a stress-responsive regulator rather than a constitutive growth promoter. This inhibitory effect likely results from the disruption of finely tuned redox and hormonal signaling, as melatonin also functions as a regulatory molecule and excessive levels can suppress seedling growth under non-stress conditions (Arnao & Hernández-Ruiz, 2019; Ameen *et al.*, 2024; Hasan et al., 2025). Collectively, these results demonstrate that melatonin effectively mitigates acid-induced inhibition of peanut seed germination and early seedling growth, underscoring its potential for improving crop establishment in acidic soils.

Melatonin enhances photosynthetic pigment stability and efficiency under acid stress in peanut

Acidic stress markedly impaired photosynthetic pigment accumulation in peanut seedlings, as evidenced by reduced chlorophyll a, chlorophyll b, total chlorophyll, and carotenoid contents under pH 4, indicating disrupted chloroplast function and pigment biosynthesis (Figs. 2a-d). Such reductions are well documented consequences of low pH-induced proton toxicity, metal ion imbalance, and oxidative damage, which collectively destabilize chlorophyll synthesis and accelerate pigment degradation during early vegetative growth (Sharma *et al.*, 2020; Yang *et al.*, 2023; Jiang *et al.*, 2025b). Our results showed that exogenous melatonin application substantially alleviated this inhibition, resulting in a dose-dependent restoration of chlorophyll and

carotenoid levels under acidic conditions, with maximal pigment accumulation
observed at higher MT concentrations (Figs. 2a-d). This response suggests that
melatonin preserves pigment integrity under acid stress, likely by protecting
chloroplast membranes and photosynthetic complexes from oxidative injury through
its strong antioxidant and redox-buffering capacity (Dou et al., 2025; Hasan et al.,
2025). Carotenoid accumulation followed a similar pattern, increasing progressively
with MT concentration under pH 4, suggesting enhanced photoprotection and reactive
oxygen species scavenging (Figs. 2a-d). The associated increase in carotenoid content
under acidic conditions further supports a protective role of melatonin, as carotenoids
are essential for quenching excess excitation energy and preventing photo-oxidative
damage under stress. Similar MT-induced stabilization of photosynthetic pigments has
been reported in crops exposed to salinity, drought, and metal toxicity, where
melatonin maintained chlorophyll biosynthesis and delayed stress-induced senescence
(An *et al.*, 2025; Basharat et al., 2025; Chen *et al.*, 2025; Hasan et al., 2025; Li *et al.*,
2025d). In contrast, under near-neutral pH (6.5), pigment levels were inherently high
in control plants, and low MT concentrations were sufficient to enhance chlorophyll
accumulation, whereas higher doses led to pigment stabilization or decline. This
concentration-dependent response indicates that melatonin primarily functions as a
stress-responsive regulator, exerting its strongest effects when pigment homeostasis is
challenged rather than under optimal growth conditions (Ali et al., 2023; Yang et al.,
2023). Overall, these results demonstrate that melatonin enhances photosynthetic
pigment stability in peanut in a pH and dose-dependent manner, with higher MT
concentrations effectively counteracting acid-induced pigment loss. This protective
role of melatonin on chlorophyll and carotenoid metabolism likely contributes to
improved photosynthetic competence and stress resilience of peanut seedlings under
acidic environments, a condition that remains underexplored in crop systems.

Melatonin-regulated antioxidant redox homeostasis confers acid stress
tolerance in peanut

Acidic stress markedly disrupts redox homeostasis in plants by enhancing reactive
oxygen species (ROS) production and impairing antioxidant efficiency, thereby
constraining early growth and physiological stability (Hlihor *et al.*, 2022; Jahan *et al.*,
2022; Hasan et al., 2025). In the present study, peanut seedlings exposed to acidic pH
(4) exhibited strong oxidative pressure, as evidenced by differentially modulated basal

activities of stress-responsive antioxidant enzymes, such as SOD, POD, CAT and APX. Exogenous MT application significantly reprogrammed the antioxidant defense system, resulting in a coordinated and stress-adaptive response. Under acidic conditions, MT markedly enhanced SOD, CAT, and APX activities in both leaves and roots, indicating improved superoxide dismutation and hydrogen peroxide detoxification, while simultaneously suppressing excessive POD activity (Figs. 3a-h). The marked reduction in POD activity observed in melatonin-treated peanut leaves and roots under acidic conditions likely reflects melatonin-mediated alleviation of oxidative stress rather than suppression of antioxidant capacity. Exogenous melatonin significantly reduced POD activity in a dose-dependent manner, indicating that MT effectively lowers ROS production and H_2O_2 availability, thereby reducing the requirement for POD-dependent scavenging (Figs. 3b and c). It has been reported that melatonin has been shown to modulate antioxidant enzyme activities and reduce ROS accumulation under various stress conditions, contributing to redox homeostasis (meta-analysis in rice and other plants) (Yang *et al.*, 2021; Muhammad *et al.*, 2025). Moreover, in other systems melatonin reduced ROS accumulation while lowering peroxidase activity, supporting the interpretation that decreased POD activity can accompany improved oxidative balance (Yang et al., 2021; Hasan et al., 2025). Thus, the reduced POD activity in MT-treated peanut seedlings reflects a functional shift from stress-induced peroxidase activation to a more energy-efficient antioxidant strategy, highlighting melatonin's dual role as both a ROS scavenger and a redox regulator under acidic stress (Figs. 3b and c). This pattern suggests that MT not only strengthens ROS-scavenging capacity but also limits stress-associated peroxidative reactions linked to cell wall rigidification and growth inhibition. Similar MT-mediated enhancement of enzymatic antioxidants and redox balance under acidic or oxidative stress has been reported in many crops, where MT improved stress tolerance by activating ROS-detoxifying enzymes and their associated genes (Ahammed *et al.*, 2020; Ahammed *et al.*, 2024; Chakraborty & Raychaudhuri, 2025; Chung et al., 2025; Li *et al.*, 2025c). Notably, the contrasting responses observed at pH 6.5 further support the stress-dependent role of melatonin. Under near-neutral conditions, MT induced only moderate or dose-limited changes in antioxidant activity, indicating that MT fine-tunes redox signaling rather than overstimulating defense pathways when oxidative stress is minimal (Figs. 3a-h). This dual behavior, strong activation under stress and restrained modulation under optimal conditions, has been recognized as a

748 hallmark of melatonin function in plants (Moustafa-Farag *et al.*, 2020; Menhas *et al.*,
749 2025; Shehzadi *et al.*, 2025). Collectively, these findings demonstrate that melatonin
750 enhances acid-stress tolerance in peanut by orchestrating a balanced antioxidant
751 response, promoting efficient ROS detoxification while preserving redox signaling,
752 thereby enabling physiological stability and growth under low-pH environments.

Melatonin fine-tunes redox homeostasis by limiting hydrogen peroxide accumulation and membrane lipid peroxidation under acidic stress

The pronounced accumulation of hydrogen peroxide (H_2O_2) and malondialdehyde
(MDA) in peanut seedlings under acidic conditions confirms that low pH imposes
severe oxidative stress, leading to excessive ROS generation and membrane lipid
peroxidation (Hlihor et al., 2022; Hasan et al., 2025). Under pH 4, control plants
exhibited markedly elevated H_2O_2 and MDA levels in both leaves and roots, reflecting
impaired redox balance and enhanced oxidative damage. Exogenous melatonin
significantly and dose-dependently reduced H_2O_2 accumulation under acidic
conditions, with the highest MT concentration lowering H_2O_2 by ~50% in leaves and
~51% in roots relative to the control (Figs. 4a and b). This strong suppression of ROS
indicates that melatonin acts primarily as an efficient antioxidant under acid stress,
either through direct free-radical scavenging or by reinforcing enzymatic
detoxification pathways, thereby reducing cellular H_2O_2 availability (Arnao &
Hernández-Ruiz, 2019; Ahammed et al., 2020; Ahammed et al., 2024; Hasan et al.,
2025; Muhammad et al., 2025).

Consistent with reduced ROS levels, MDA content declined sharply in MT-treated
seedlings under pH 4, with up to ~69% and ~68% reductions in leaves and roots,
respectively, indicating effective protection of membrane integrity (Figs. 4c and d).
The tight coupling between decreased H_2O_2 and lower lipid peroxidation suggests that
melatonin limits oxidative damage by stabilizing cellular membranes and preventing
ROS-induced fatty acid oxidation, a response widely associated with improved stress
tolerance (Ali et al., 2023; Ahammed et al., 2024; Hasan et al., 2025). In contrast,
under near-neutral pH (6.5), low MT concentrations enhanced oxidative protection,
whereas higher doses partially restored H_2O_2 and MDA levels (Figs. 4a-d), supporting
the concept that melatonin permits controlled ROS accumulation for signaling under
non-stress conditions (Hasan et al., 2025; Li *et al.*, 2025a). Collectively, these results
demonstrate that melatonin functions as a redox modulator in peanut, acting

predominantly as an antioxidant under acidic stress while maintaining ROS-
dependent signaling at optimal pH, thereby ensuring balanced cellular homeostasis.
783

Melatonin enhances plasma membrane H⁺-ATPase transcription to
maintain pH homeostasis under acidic stress

Plasma membrane H⁺-ATPases are key regulators of proton extrusion, cytosolic pH
stability, and nutrient uptake, and their function becomes especially critical under
acidic stress, where excessive proton influx disrupts cellular metabolism and root
growth (Chao & Chao, 2025; Dong *et al.*, 2025). In the present study, melatonin (MT)
significantly and dose-dependently upregulated the expression of the H⁺-ATPase
genes *AhAH1* and *AhAH2* in peanut seedlings under acidic conditions, with markedly
stronger induction in roots than in leaves. This transcriptional response provides
molecular evidence that MT reinforces proton transport to the apoplast to improve
acid tolerance (Figs. 5a and b).

The strong MT-induced expression of *AhAH1* and *AhAH2* is consistent with
previous reports showing that melatonin enhances H⁺-ATPase activity and
expression under various abiotic stresses (Wang *et al.*, 2025c). For example, MT has
been shown to stimulate plasma membrane H⁺-ATPases in cucumber and tomato
under salinity and nutrient stress, thereby stabilizing intracellular pH and improving
ion homeostasis (Li et al., 2022; Zeng et al., 2024; Li *et al.*, 2025b). These findings
align closely with our observation that *AhAH1* and *AhAH2* expression was most
strongly induced at higher melatonin concentrations under pH 4, highlighting a
conserved role of melatonin in activating proton transport machinery under acidic
stress (Figs. 5a and b). Under acidic or aluminium-stress conditions, enhanced H⁺-
ATPase activity promotes proton efflux and mitigates rhizotoxic effects, supporting
root elongation and function (Zhang *et al.*, 2017; Munyaneza *et al.*, 2024). The robust
upregulation of *AhAH2* at higher melatonin doses indicates that this isoform may
function as a stress-adaptive proton pump, supporting sustained root activity,
membrane potential stability, and nutrient acquisition under acidic conditions. The
pronounced induction of *AhAH2*, particularly in roots, suggests isoform-specific stress
responsiveness, in line with studies in Arabidopsis and rice, where certain H⁺-
ATPase genes are preferentially activated under low-pH stress to maintain root

apoplastic and cytosolic pH balance (Gámez-Arjona *et al.*, 2022; Chakraborty *et al.*,
2024; Jain & Schmidt, 2024).

Mechanistically, melatonin-mediated upregulation of H^+-ATPase genes likely
reflects its integrative role in redox and hormonal signaling. Melatonin modulates
ROS levels and interacts with auxin and calcium-dependent pathways, both of which
are known regulators of H^+-ATPase transcription and activation (Huang et al., 2022;
Yang *et al.*, 2024; Chung et al., 2025). By alleviating oxidative stress while
maintaining controlled ROS signaling, MT may create favorable cellular conditions
for sustained proton pump expression and activity under acidic environments. Overall,
these findings demonstrate that melatonin enhances acid stress tolerance in peanut by
transcriptionally activating plasma membrane H^+-ATPase genes, particularly *AhAH2*
in roots. This response likely strengthens proton extrusion, stabilizes intracellular pH,
and supports root performance under low-pH conditions, highlighting H^+-ATPase
regulation as a key molecular component of melatonin-mediated adaptation to acidic
stress in crops.

Interaction between melatonin concentration and pH regulates peanut
physiological responses

Across all measured parameters, including germination, seedling growth, biomass
accumulation, photosynthetic pigments, oxidative stress markers, antioxidant enzyme
activities, and H^+-ATPase gene expression, a clear interaction between melatonin
concentration and pH level was observed. Acidic conditions (pH 4) consistently
amplified the positive effects of melatonin, whereas near-neutral pH (6.5) markedly
reduced or altered MT responsiveness. This interaction indicates that melatonin does
not act as a simple growth promoter, but rather functions as a stress-dependent
regulator whose efficacy is tightly linked to environmental pH (Fig. 6).

Under acidic stress, increasing MT concentrations progressively improved
germination, root and shoot development, pigment accumulation, redox homeostasis,
and proton pump gene expression. These responses reflect a coordinated mitigation of
proton toxicity, oxidative damage, and metabolic disruption, suggesting that higher
MT availability is required to counteract the elevated ROS burden and pH imbalance
typical of acid stress (Fig. 6). Similar pH or stress-intensity-dependent melatonin
responses have been reported under salinity, metal toxicity, and drought, where
stronger stress necessitated higher MT doses to restore cellular homeostasis (Arnao &

Hernández-Ruiz, 2019; Ahmad et al., 2023b; Ahammed et al., 2024; An et al., 2025;
Basharat et al., 2025).

In contrast, under near-neutral pH, most physiological parameters peaked at
low MT concentrations and declined at higher doses, indicating that excessive
melatonin can disrupt optimal redox and hormonal signaling when stress pressure is
minimal (Fig. 6). Basal ROS are essential signaling molecules for normal growth,
photosynthesis, and development, and over-scavenging by high MT levels may
suppress these signaling pathways, leading to reduced growth and metabolic
efficiency (Back, 2021; Liu et al., 2023; Dzinyela *et al.*, 2025). Thus, the MT and pH
interaction demonstrates that melatonin acts within an optimal concentration window
that shifts upward with increasing stress severity.

Collectively, this interaction highlights melatonin as a conditional bioregulator
whose beneficial range depends on environmental pH, emphasizing the importance of
dose optimization when applying MT in agronomic systems.

Figure 6. Melatonin and pH interaction shows that a higher melatonin level is required to
alleviate seedling growth and yield, oxidative damage, photosynthetic pigments, ROS and
H⬚-ATPase genes expression and pH imbalance under acidic stress, whereas excessive
melatonin can disrupt normal growth under optimal conditions.

Conclusion

The present study provides comprehensive evidence that exogenous melatonin application markedly enhances peanut tolerance to acidic stress by integrating redox regulation, growth maintenance, and pH homeostasis. Acidic conditions severely restricted seed germination, seedling growth, biomass accumulation, pigment synthesis, and differentially affect antioxidants, while inducing excessive H_2O_2 accumulation and membrane lipid peroxidation. Melatonin application significantly reversed these effects, particularly at pH 4, by suppressing oxidative damage, protecting the photosynthetic apparatus, and sustaining biomass production. This protection was associated with a coordinated reprogramming of the antioxidant system, characterized by enhanced SOD, CAT and APX activities and reduced POD activity, indicating alleviation of oxidative pressure and a shift toward more efficient H_2O_2 detoxification. Importantly, melatonin strongly induced the plasma membrane H□-ATPase genes *AhAH1* and *AhAH2*, especially in roots, providing molecular evidence that melatonin reinforces proton extrusion capacity and cellular pH stability under acid stress.

Overall, the strong melatonin and pH interaction observed across all parameters demonstrates that higher melatonin concentrations are required to confer effective protection under acidic stress, whereas excessive melatonin under near-neutral conditions can impair normal physiological regulation. Collectively, these findings establish melatonin as a multifunctional regulator of acid stress adaptation in peanut and reveal a previously underexplored role of melatonin in mechanistically linking redox control with proton transport to support early crop establishment on acidic soils.

Author Contributions

Muhammad Hafeez Ullah Khan: conceptualization, investigation, writing – original draft. **Fu Ruoyi:** methodology and formal analysis. **Ali Muhammad:** visualization and formal analysis. **Shuaichao Zheng:** data curation and formal analysis. **Daijing Zhang:** writing – review and editing. **Zhiyong Zhang:** supervision, funding acquisition, writing – review and editing. **Quanyong Liu:** supervision, funding acquisition, writing – review and editing.

Funding and Acknowledgement

This research was partially supported by the National Natural Science Foundation of China (31571600, 31271648), the Key R&D and Promotion Projects of Henan Province (222102110303, 232102110221, 242102111094, 252102111072), the International Science and Technology Cooperation Project of Henan Province

899 (242102521050), and the Henan Center of Outstanding Overseas Scientists
900 (GZS2024018). We gratefully acknowledge Professor Xiaoping Pan from East
901 Carolina University for her great input, comments, and proofreading of this paper.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

References

Ahammed, G. J., Li, Z., Chen, J., Dong, Y., Qu, K., Guo, T., Wang, F., Liu, A., Chen, S. & Li, X. 2024. Reactive oxygen species signaling in melatonin-mediated plant stress response. *Plant Physiology and Biochemistry,* **207,** 108398.

Ahammed, G. J., Wu, M., Wang, Y., Yan, Y., Mao, Q., Ren, J., Ma, R., Liu, A. & Chen, S. 2020. Melatonin alleviates iron stress by improving iron homeostasis, antioxidant defense and secondary metabolism in cucumber. *Scientia Horticulturae,* **265,** 109205.

Ahmad, I., Song, X., Hussein Ibrahim, M. E., Jamal, Y., Younas, M. U., Zhu, G., Zhou, G. & Adam Ali, A. Y. 2023a. The role of melatonin in plant growth and metabolism, and its interplay with nitric oxide and auxin in plants under different types of abiotic stress. *Front Plant Sci,* **14,** 1108507.

Ahmad, R., Alsahli, A. A., Alansi, S. & Altaf, M. A. 2023b. Exogenous melatonin confers drought stress by promoting plant growth, photosynthetic efficiency and antioxidant defense system of pea (Pisum sativum L.). *Scientia Horticulturae,* **322,** 112431.

Ali, S., Gill, R. A., Ulhassan, Z., Zhang, N., Hussain, S., Zhang, K., Huang, Q., Sagir, M., Tahir, M. B., Gill, M. B., Mwamba, T. M., Ali, B. & Zhou, W. 2023. Exogenously applied melatonin enhanced the tolerance of Brassica napus against cobalt toxicity by modulating antioxidant defense, osmotic adjustment, and expression of stress response genes. *Ecotoxicology and Environmental Safety,* **252,** 114624.

Ameen, M., Zafar, A., Mahmood, A., Zia, M. A., Kamran, K., Javaid, M. M., Yasin, M. & Khan, B. A. 2024. Melatonin as a master regulatory hormone for genetic responses to biotic and abiotic stresses in model plant Arabidopsis thaliana: a comprehensive review. *Functional Plant Biology,* **51**.

An, W., Wang, G., Dou, J., Zhang, Y., Yang, Q., He, Y., Tang, Z. & Yu, J. 2025. Protective mechanisms of exogenous melatonin on chlorophyll metabolism and photosynthesis in tomato seedlings under heat stress. *Frontiers in Plant Science,* **16,** 1519950.

Arnao, M. B. & Hernández-Ruiz, J. 2019. Melatonin: a new plant hormone and/or a plant master regulator? *Trends in Plant Science,* **24,** 38–48.

Back, K. 2021. Melatonin metabolism, signaling and possible roles in plants. *The Plant Journal,* **105,** 376–391.

941   Basharat, S., Saeed, W., Mubeen, S., Khan, L. U., Zhang, S., Liu, P. & Waseem, M. 2025.
942   Exogenous Melatonin Regulates Hormone Signalling and
943   Photosynthesis-Related Genes to Enhance Brassica napus. Yield: A
944   Transcriptomic Perspective. *Journal of Pineal Research,* **77,** e70077.
945   Borhannuddin Bhuyan, M., Hasanuzzaman, M., Nahar, K., Mahmud, J. A., Parvin, K.,
946   Bhuiyan, T. F. & Fujita, M. 2019. Plants behavior under soil acidity stress:
947   Insight into morphophysiological, biochemical, and molecular responses.
948   *Plant abiotic stress tolerance: Agronomic, molecular and biotechnological*
949   *approaches*, 35–82.
950   Cai, B., Vancov, T., Si, H., Yang, W., Tong, K., Chen, W. & Fang, Y. 2021. Isolation and
951   Characterization of Endomycorrhizal Fungi Associated with Growth Promotion
952   of Blueberry Plants. *J Fungi (Basel),* **7**.
953   Chakraborty, N., Das, A., Pal, S., Roy, S., Sil, S. K., Adak, M. K. & Hasanuzzaman, M.
954   2024. Exploring Aluminum tolerance mechanisms in plants with reference to
955   rice and Arabidopsis: A comprehensive review of genetic, metabolic, and
956   physiological adaptations in acidic soils. *Plants,* **13,** 1760.
957   Chakraborty, S. & Raychaudhuri, S. S. 2025. Melatonin improves the lead tolerance in
958   Plantago ovata by modulating ROS homeostasis, phytohormone status and
959   expression of stress-responsive genes. *Plant Cell Reports,* **44,** 39.
960   Chao, Z.-F. & Chao, D.-Y. 2025. Barriers and carriers for transition metal homeostasis
961   in plants. *Plant Communications,* **6**.
962   Chen, T., Xu, L., Yang, P., Tong, J., Liu, Y., Su, Y. & Sun, S. 2025. Plant melatonin: roles
963   and regulatory mechanisms in plant growth, development, and responses to
964   biotic and abiotic stresses. *Plant Signaling & Behavior,* **20,** 2585629.
965   Chung, I.-M., Thiruvengadam, M., Samynathan, R., Lee, S.-B., Choi, H.-J., Jung, B.-S.,
966   Park, Y., Jeon, D., Venkidasamy, B. & Kim, S.-H. 2025. Molecular insights into
967   melatonin-mediated stress tolerance in rice. *Plant Cell Reports,* **44,** 271.
968   Cosse, M. & Seidel, T. 2021. Plant proton pumps and cytosolic pH-homeostasis.
969   *Frontiers in Plant Science,* **12,** 672873.
970   Delfim, J., Moreira, A. & Moraes, L. A. 2025. Wheat growth, yield, nutrient
971   concentration and soil chemical properties influenced by liming in an acid soil.
972   *Journal of Plant Nutrition,* **48,** 87–100.
973   Dong, T., Yan, L., Wang, J., Niu, Y. & Wang, L. 2025. Salt Tolerance in Soybean (Glycine
974   max L.): A Comprehensive Review of Molecular Mechanisms, Key Regulators,
975   and Future Perspectives for Saline Soil Utilization. *Plants,* **14,** 3668.
976   Dou, J., Tang, Z., Yu, J., Wang, G., An, W., Zhang, Y. & Yang, Q. 2025. Effects of
977   exogenous melatonin on the growth and photosynthetic characteristics of
978   tomato seedlings under saline-alkali stress. *Scientific Reports,* **15,** 5172.
979   Dzinyela, R., Manda, T., Hwarari, D., Ramakrishnan, M., Ahmad, Z., Agassin, R. H.,
980   Yang, L. & Movahedi, A. 2025. Melatonin-mediated phytohormonal crosstalk
981   improves salt stress tolerance in plants. *Planta,* **262,** 86.
982   Fu, D., Wu, W., Mustafa, G., Yang, Y. & Yang, P. 2025. Molecular mechanisms of rice
983   seed germination. *New Crops,* **2,** 100051.
984   Gámez-Arjona, F. M., Sánchez-Rodríguez, C. & Montesinos, J. C. 2022. The root
985   apoplastic pH as an integrator of plant signaling. *Frontiers in Plant Science,* **13,**
986   931979.

Hasan, M. K., Imon, M. A. A.-H., Jahan, I., Das, T. K., Mishu, M. S., Shopan, J., Saha, M., Khan, M. S. I., Islam, M. S. & Ahammed, G. J. 2025. Melatonin-mediated adaptation of spinach (Spinacia oleracea L.) in acidic soil. *Plant Physiology and Biochemistry,* **223,** 109816.

Hlihor, R. M., Roşca, M., Hagiu-Zaleschi, L., Simion, I. M., Daraban, G. M. & Stoleru, V. 2022. Medicinal plant growth in heavy metals contaminated soils: Responses to metal stress and induced risks to human health. *Toxics,* **10,** 499.

Huang, K., Li, M., Li, R., Rasul, F., Shahzad, S., Wu, C., Shao, J., Huang, G., Li, R., Almari, S., Hashem, M. & Aamer, M. 2023. Soil acidification and salinity: the importance of biochar application to agricultural soils. *Frontiers in Plant Science,* **Volume 14 - 2023**.

Huang, X., Tanveer, M., Min, Y. & Shabala, S. 2022. Melatonin as a regulator of plant ionic homeostasis: implications for abiotic stress tolerance. *Journal of Experimental Botany,* **73,** 5886–5902.

Jahan, I., Shopan, J., Rahman, M. M., Sarkar, A., Baset, M. A., Zhang, Z., Li, X., Ahammed, G. J. & Hasan, M. K. 2022. Long-term traditional fertilization alters tea garden soil properties and tea leaf quality in Bangladesh. *Agronomy,* **12,** 2128.

Jain, D. & Schmidt, W. 2024. Protein phosphorylation orchestrates acclimations of arabidopsis plants to environmental pH. *Molecular & Cellular Proteomics,* **23**.

Jamir, Z., Tzudir, L., Kumari, S., Solo, V., Dolie, S., Nongmaithem, D., Yadav, R., Kikon, N. & Mollier, M. 2025. Productivity of Groundnut (Arachis hypogaea L.) under Different Sowing Times, Utrient Supply Levels and Planting Geometry in Sub-Humid Tropical Environment. *Current World Environment,* **20,** 121.

Jiang, D., Ou, Y., Jiang, G., Dai, G., Liu, S. & Chen, G. 2025a. Melatonin-priming ameliorates aluminum accumulation and toxicity in rice through enhancing aluminum exclusion and maintaining redox homeostasis. *Plant Physiology and Biochemistry,* **219,** 109433.

Jiang, Z., van Zanten, M. & Sasidharan, R. 2025b. Mechanisms of plant acclimation to multiple abiotic stresses. *Communications Biology,* **8,** 655.

Kong, X., Wang, R., Jia, P., Li, H., Khan, A., Muhammad, A., Fiaz, S., Xing, Q. & Zhang, Z. 2025. Physio-biochemical and molecular mechanisms of low nitrogen stress tolerance in peanut (Arachis hypogaea L.). *Plant Molecular Biology,* **115,** 19.

Li, J., Yang, J., Liang, X., Zhan, S., Bai, Y. & Ruan, L. 2025a. Melatonin-Mediated Regulation of Antioxidant Defense Enhances the Resistance of Tea Plants (Camellia sinensis L.) to Lead-Induced Stress. *Plants,* **14,** 1417.

Li, J., Zhang, Z. & Shi, G. 2024a. Genome-wide identification and expression profiling of heavy metal ATPase (HMA) genes in peanut: Potential roles in heavy metal transport. *International Journal of Molecular Sciences,* **25,** 613.

Li, L., Cheng, X., Zhang, Y., Kohtz, D., Wang, X., Zhang, X., Kong, X., Xue, H., Jia, P., Bai, N., Li, Z., Xiao, P., Pan, X., Zhang, Z. & Zhang, B. 2024b. Exogenous melatonin improves peanut field productivity and quality at reduced nitrogen application. *Field Crops Research,* **319,** 109650.

Li, L., Li, Q., Davis, K. E., Patterson, C., Oo, S., Liu, W., Liu, J., Wang, G., Fontana, J. E., Thornburg, T. E., Pratt, I. S., Li, F., Zhang, Z., Zhou, Y., Pan, X. & Zhang, B. 2021. Response of Root Growth and Development to Nitrogen and Potassium

Deficiency as well as microRNA-Mediated Mechanism in Peanut (Arachis
hypogaea L.). *Frontiers in Plant Science,* **Volume 12 - 2021**.

Li, Q., Zhang, Y., Liu, Y., Li, T., Xu, H., Wei, Q., Zeng, H., Ni, H. & Li, S. 2025b. Multiple
functions of exogenous melatonin in cucumber seed germination, seedling
establishment, and alkali stress resistance. *BMC Plant Biology,* **25,** 359.

Li, W.-Q., Li, J.-Y., Bi, S.-J., Jin, J.-Y., Fan, Z.-L., Shang, Z.-L., Zhang, Y.-F. & Wang, Y.-J.
2025c. Melatonin enhances maize germination, growth, and salt tolerance by
regulating reactive oxygen species accumulation and antioxidant systems.
Plants, **14,** 296.

Li, Y., Zeng, H., Xu, F., Yan, F. & Xu, W. 2022. H+-ATPases in plant growth and stress
responses. *Annual Review of Plant Biology,* **73,** 495–521.

Li, Z., Tang, Z., Song, J., Li, J., Wang, X. & Zhang, X. 2025d. Melatonin stabilizes
photosynthesis and scavenges ROS to enhance salt tolerance in cucumber.
BMC Plant Biology, **25,** 1433.

Liu, H., Ren, X., Zhu, J., Wu, X. & Liang, C. 2018. Effect of exogenous abscisic acid on
morphology, growth and nutrient uptake of rice (Oryza sativa) roots under
simulated acid rain stress. *Planta,* **248,** 647–659.

Liu, Z., Dai, H., Hao, J., Li, R., Pu, X., Guan, M. & Chen, Q. 2023. Current research and
future directions of melatonin's role in seed germination. *Stress Biology,* **3,** 53.

MacDonald, E., Johannes, L. & Wunder, C. 2025. Acidification on the plasma
membrane. *Current Opinion in Cell Biology,* **95,** 102531.

Menhas, S., Chen, M., Jin, H., Xu, J., Zhu, S. & Lin, D. 2025. Plant growth stage and
melatonin concentration dependency together drive the metal-nutrient
dynamics of rice in paddy soil. *International Journal of Phytoremediation,* **27,**
958–971.

Moustafa-Farag, M., Elkelish, A., Dafea, M., Khan, M., Arnao, M. B., Abdelhamid, M.
T., El-Ezz, A. A., Almoneafy, A., Mahmoud, A., Awad, M., Li, L., Wang, Y.,
Hasanuzzaman, M. & Ai, S. 2020. Role of Melatonin in Plant Tolerance to Soil
Stressors: Salinity, pH and Heavy Metals. *Molecules,* **25,** 5359.

Msimbira, L. A. & Smith, D. L. 2020. The roles of plant growth promoting microbes in
enhancing plant tolerance to acidity and alkalinity stresses. *Frontiers in
Sustainable Food Systems,* **4,** 106.

Muhammad, A., Kong, X., Li, L., Khan, M. H. U., Jia, P., Miao, C. & Zhang, Z. 2026.
Melatonin Enhances Peanut Productivity by Enriching Root-Associated
Nitrogen-Fixing Bacteria. *Journal of Pineal Research,* **78,** e70105.

Muhammad, I., Ahmad, S. & Shen, W. 2024. Melatonin-mediated molecular
responses in plants: Enhancing stress tolerance and mitigating environmental
challenges in cereal crop production. *International Journal of Molecular
Sciences,* **25,** 4551.

Muhammad, I., Fahad, S., Khalofah, A., Zheng, B. & Shen, W. 2025. Melatonin
Enhances Antioxidant Defense Systems and Stress Tolerance in Plants under
Variable Environmental Conditions. *Rice (N Y),* **18,** 70.

Munyaneza, V., Zhang, W., Haider, S., Xu, F., Wang, C. & Ding, G. 2024. Strategies for
alleviating aluminum toxicity in soils and plants. *Plant and Soil,* **504,** 167–190.

Naz, M., Dai, Z., Hussain, S., Tariq, M., Danish, S., Khan, I. U., Qi, S. & Du, D. 2022. The
soil pH and heavy metals revealed their impact on soil microbial community.
Journal of Environmental Management, **321,** 115770.

Sharma, A., Kumar, V., Shahzad, B., Ramakrishnan, M., Singh Sidhu, G. P., Bali, A. S.,
Handa, N., Kapoor, D., Yadav, P. & Khanna, K. 2020. Photosynthetic response
of plants under different abiotic stresses: a review. *Journal of Plant Growth*
Regulation, **39,** 509–531.
Shavrukov, Y. & Hirai, Y. 2016. Good and bad protons: genetic aspects of acidity stress
responses in plants. *Journal of Experimental Botany,* **67,** 15–30.
Shehzadi, K., Maqsood, M. F., Kanwal, R., Shahbaz, M., Naqve, M., Zulfiqar, U., Jamil,
M., Khalid, N., Ali, M. F. & Soufan, W. 2025. Enhancing cadmium stress
resilience in chickpea (Cicer arietinum L.) via exogenous melatonin application.
International Journal of Phytoremediation, **27,** 794–809.
Sun, G., Luan, M., Xue, S., Yan, J. & Lan, W. 2025. Vacuolar H+-ATPase Is Required for
Efficient Vacuolar Phosphate Storage and Systemic Pi Homeostasis in
Arabidopsis. *Plant, Cell & Environment,* **48,** 587–598.
Tahjib-Ul-Arif, M., Zahan, I., Hossain, M. S., Imran, S., Hasanuzzaman, M., Dawood, M.
F., Dawood, A. F., Asaduzzaman, M., Rhaman, M. S. & Souri, Z. 2025.
Melatonin-mediated ionic homeostasis in plants: Mitigating nutrient
deficiency and salinity stress. *Discover Plants,* **2,** 143.
Thapa, A., Hasan, M. R. & Kabir, A. H. 2025. Transcriptional reprogramming and
microbiome dynamics in garden pea exposed to high pH stress during
vegetative stage. *Planta,* **261,** 83.
Wang, F., Li, X., Chen, C., Zhao, L. & Wei, Y. 2025a. Mechanism of Melatonin in
Alleviating Aluminum Toxicity in Plants: A Review. *Biology,* **14,** 1316.
Wang, J., Jin, D., Deng, Z., Zheng, L., Guo, P., Ji, Y., Song, Z., Zeng, H. Y., Kinoshita, T. &
Liao, Z. 2025b. The apoplastic pH is a key determinant in the hypocotyl
growth response to auxin dosage and light. *Nature Plants*, 1–16.
Wang, L., Ju, C., Han, C., Yu, Z., Bai, M. Y. & Wang, C. 2025c. The interaction of
nutrient uptake with biotic and abiotic stresses in plants. *Journal of*
Integrative Plant Biology, **67,** 455–487.
Yang, S.-J., Huang, B., Zhao, Y.-Q., Hu, D., Chen, T., Ding, C.-B., Chen, Y.-E., Yuan, S. &
Yuan, M. 2021. Melatonin enhanced the tolerance of Arabidopsis thaliana to
high light through improving anti-oxidative system and photosynthesis.
Frontiers in Plant Science, **12,** 752584.
Yang, X., Liu, D., Liu, C., Li, M., Yan, Z., Zhang, Y. & Feng, G. 2024. Possible melatonin-
induced salt stress tolerance pathway in Phaseolus vulgaris L. using
transcriptomic and metabolomic analyses. *BMC Plant Biology,* **24,** 72.
Yang, Y., Guan, S., Jiang, X., Li, M., Wei, S. & Diao, M. 2023. Exogenous melatonin
alleviates the inhibitory effect of NaHCO3 on tomato growth by regulating the
root pH value and promoting plant photosynthesis. *Agronomy,* **13,** 2777.
Zeng, H., Chen, H., Zhang, M., Ding, M., Xu, F., Yan, F., Kinoshita, T. & Zhu, Y. 2024.
Plasma membrane H+-ATPases in mineral nutrition and crop improvement.
Trends in Plant Science, **29,** 978–994.
Zhang, J., Wei, J., Li, D., Kong, X., Rengel, Z., Chen, L., Yang, Y., Cui, X. & Chen, Q. 2017.
The role of the plasma membrane H+-ATPase in plant responses to aluminum
toxicity. *Frontiers in Plant Science,* **8,** 1757.
Zheng, S., Tang, Z., Li, L., Jia, P., Kong, X., Xue, H., Cheng, X., Wang, X., Zhang, Y. & Bai,
N. 2025. Melatonin reprograms soil microbial community, creates friendly soil

environments, and promotes peanut growth. *Plant Physiology and*
Biochemistry, **218,** 109307.

1128