Beneficial microbes may have favoured the evolution of adaptive immunity

Lucas Mathieu[1], Thomas Pradeu[2], Richard Watson[3], and Kevin Lala[1]

[1]School of Biology, University of St Andrews, St Andrews, UK
[2]University of Bordeaux, CNRS, ImmunoConcept, UMR 5164, Bordeaux, France
[3]Electronics and Computer Science/Institute for Life Sciences, University of Southampton, Southampton, UK

June 2025

Abstract

Vertebrate adaptive immunity has long been considered uniquely advantageous for its capacity to recognise and remember a wide diversity of molecular targets, potentially conferring lasting defence from any threatening microbe. However, over the last two decades, results from comparative immunology have called into question the evolutionary benefits of vertebrate adaptive immunity as a defence mechanism, raising the possibility of alternative evolutionary explanations. For instance, the hypothesis that managing the complex and often beneficial host-microbiota associations characteristic of the vertebrate gut may have favoured the evolution of adaptive immunity. Here, we use individual-based simulations to investigate co-evolutionary interactions between hosts and their microbiota and the implication for the evolution of adaptive immunity. We focus on the capacity of adaptive immunity to produce and modify a diverse repertoire of immune receptors by using somatic variation and selective clonal reinforcement, in combination with a reinforcement bias from the innate immune system. Strikingly, we find that a high diversity of rapidly evolving parasites is insufficient to favour the evolution of adaptive immunity. Instead, our findings suggest that neutral and beneficial microbes can play a crucial role in weakening selection on innate immune defence against parasites in favour of the exploitation of beneficial microbes. In turn, this facilitates the emergence of adaptive immunity as a mechanism reducing the risk associated with infrequent parasite infections while exploiting beneficial microbes. In addition, we show that adaptive immunity can subsequently alter selection affecting both innate immune defence and microbe evolution, such that its loss becomes increasingly unlikely with evolutionary time. Together, these results suggest that simultaneous interactions with diverse mutualists and parasites is a compelling selective explanation for the emergence and maintenance of adaptive immunity.

1 Introduction

Vertebrate adaptive immunity is generally defined as the capacity to generate lineages of cells expressing somatically-diversified receptors used in the binding, recognition and potential elimination

of microbes [1]; and to re-mobilise the receptors acquired during the initial response to a pathogen (e.g., using long-lived cells) to accelerate subsequent responses to similar stimuli [2]. This process is called 'adaptive' because it allows hosts to generate receptors in response to novel immune challenges through stochastic and feedback-dependent phenotypic variation *within lifetime* [3] instead of relying on genetic variation and natural selection across generations. Although vertebrate immune systems vary largely, this core functional pattern is seemingly uniquely conserved across vertebrates but absent in invertebrates, and is thought to have evolved in ancient fish ∼500 Mya [4, 5]. However, generating and updating a reservoir of new receptors is presumably costly because stochastic and pre-emptive variation often produces significant amounts of unhelpful phenotypic variation. For instance, a large proportion (up to 90%) of B lymphocyte precursors are eliminated to prevent auto-immune reactions [6]. In addition, among the staggering diversity of lymphocyte receptors produced within a vertebrate host's lifespan only a fraction is susceptible to react to specific antigens (e.g., in a recent study only ∼ 0.003% of human naive B cells were specific to the receptor-binding domain of SARS-CoV-2; [7]).

For the past few decades, the evolutionary trade-off imposed by generating costly and potentially ineffectual phenotypic variation has been explained by the presumed defensive benefit afforded by the flexibility offered by the under-determination of initial adaptive immune responses by host genomes and by the rapid secondary responses produced subsequently [1]. However, the idea that adaptive immunity is a markedly beneficial form of defence has recently received increased scrutiny [8], for at least three reasons. First, immune memory is broadly spread across invertebrates and plants and is even present in cells considered as part of the innate immune system in vertebrates [8, 9, 10], while the efficacy of long-term immune memory starkly varies across vertebrates [11]. Second, supposing that adaptive immunity is a superior system of defence raises the question of why a similar mechanism did not evolve in invertebrates [12]. Third, adaptive immune responses are often dependent on the initial recognition and capture of microbes by innate immune cells [13]. Together, these objections have motivated further hypotheses to explain why adaptive immunity evolved and was subsequently conserved across jawed and jawless vertebrates but not in invertebrates [14, 15, 16, 12]. Multiple authors have approached this question by arguing that evolutionary constraints rather than selective explanations are responsible for the absence of functional counterparts to adaptive immunity in invertebrates —besides immune memory, often called "trained immunity" in non-vertebrate systems [17]. For instance, by proposing that invertebrates lacked innovations such as the RAG transposons involved in somatic recombination [18, 19] or the available genomic complexity required for the regulatory control of adaptive immune responses [16]. In the following, we focus our analysis on a particular selective explanation, but we consider the influence of developmental and genetic constraints in Discussion.

Here, we investigate the hypothesis that adaptive immunity evolved not solely for defence but rather to regulate interactions between the host and a diverse microbiota. That is, it evolved to "combat pathogens while maintaining a tolerant relationship with symbiotic microorganisms" [20]. This argument was initially proposed by McFall-Ngai (2007) and further discussed by Lee and Mazmanian (2010) and others [23, 24] based on two principal motivations. First, extensive empirical evidence points to the complex and usually beneficial interactions between vertebrates (particularly mammals) and their gut microbiota, which are often mediated by adaptive immunity [25, 26, 27, 28]. Second, invertebrates usually form associations with fewer resident microbes than vertebrates [29, 30]. From the host's perspective, matching a high diversity of gut microbes —both beneficial and threatening— constitutes a challenging evolutionary problem. Innate immunity

is traditionally viewed as constituted of conserved immune pathways such as TLRs that target common microbe-associated molecular patterns (MAMPs), which may limit distinctions between harmful and beneficial microbes and impede the ability to foster a diverse and beneficial microbiota [31] (but see [32, 33]). By contrast, matching a diverse community of microbes with a somatically diversified repertoire of receptors may seem like a more parsimonious evolutionary solution [22]. However, recent discussions of a microbiota-driven adaptive immunity have primarily been motivated by contemporary immunity-microbiota interactions [34, 35], which may not accurately reflect interactions across ancient evolutionary history. Crucially, a mechanistic account of how interactions with the microbiota could facilitate the emergence and subsequent conservation of an ancestral form of adaptive immunity is still lacking, to our knowledge. Below, we propose a minimal mechanistic model to evaluate this hypothesis.

We focus our analysis on a common functional pattern conserved in vertebrate immunity [36, 4] characterised by 1) the generation of somatic variation in receptor genes exclusively expressed by distinct clonal cell lineages, leading to a diverse anticipatory repertoire [37]; and 2) the differential multiplication of clonal receptor-bearing cell lineages mediated by affinity-dependent competition for antigen binding. Assuming that hosts can sample some of the microbes present in their gut, this is sufficient to allow the proportion of receptors with high binding affinity to follow the proportion of their corresponding antigens, potentially leading to highly specific immune responses [1, 38]. The precise mechanisms involved in the initial generation of receptor diversity and subsequent selection differ between jawed and jawless vertebrates [36, 39, 4, 5, 40, 16]. However, in both vertebrate lineages this process of somatic variation and selection generates two archetypes of receptor-expressing cell types (hereafter 'lymphocytes'): a regulatory subset (i.e., VLRA in jawless fish and T lymphocytes in vertebrates) and a secretory subset (respectively VLRB in jawless fish and B cells). We focus our reasoning on the latter archetype, on account of its high conservation and direct role in regulating microbe-host interactions in the gut —sometimes independently from the intervention of the regulatory archetype in the case of mucosa-associated secretory lymphocytes [41, 42, 24, 43]. Admittedly, regulatory lymphocytes also play a key role in host-microbe interactions [44] and might have evolved hand-in-hand with, or shortly following, humoral immunity to reduce the risk of 'self'-directed responses [45, 24]. For simplicity, here we assumed that regulatory control evolved as required by humoral immunity and we focused on interactions at the gut mucosal interface. We refer to this restricted model of humoral adaptive immunity as 'somatically diversified immunity' (henceforth SDI) to stress the importance of somatically diversified and selected receptors. We avoid the term "adaptive immunity" to prevent confusions between this minimal model and contemporary capacities of systemic adaptive immune systems such as affinity maturation by hypermutation [38], long-term memory [46, 47] and complex regulatory control [48, 49, 50], which are not all conserved across vertebrates — see Rimer and colleagues (2014) for a comparative review in cold-blooded vertebrates.

To investigate whether interactions with the microbiota could have favoured the evolution of SDI, we consider an ancestral population of vertebrates with innate immunity co-evolving with a population of microbes. Using agent-based evolutionary simulations, we compare scenarios where the diversity of microbes and the evolutionary constraints affecting their evolution vary. In parallel, we assess the importance of host factors commonly assumed to influence host-microbe associations, including the shape of the dose-response function associating microbe density to host fitness. We show that SDI is favoured in hosts interacting with diverse populations of fast-evolving microbes.

However, a higher diversity of threatening microbes did not increase the frequency of SDI. Rather, we find that diverse and generally beneficial communities of microbes tend to weaken selection for innate immune defence, which in turn favours the emergence of SDI. We also examined the conditions in which SDI can subsequently be conserved across long evolutionary time, in an attempt to explain why SDI might be conserved in jawed and jawless vertebrates but absent in invertebrates. We show that SDI can disrupt selection on the innate immune system and stabilise the evolution of a higher diversity of beneficial microbes, both of which may contribute to its long-term conservation.

Terminological considerations: mutualists, commensals, parasites and pathogens

Scientific works investigating interactions between hosts and microbiota and their significance for adaptive immunity use a variety of terms to characterise microbes with beneficial ("partners" [21]; "symbionts" [51, 52, 53, 32]; "commensals" [31]) and harmful ("pathogen" [32, 43, 22, 31]; "pathobiont" [54, 55]; "parasites" [56]) effects on their hosts. But this categorical distinction is often artificial, with microbe ecology and evolution blurring the line between neutral, harmful and beneficial microbes [57, 58]. Generally, microbe "virulence" (i.e., direct damage to the host) is highly context-dependent [59] and can for example be plastically induced via quorum sensing [60] and other microbe-microbe interactions [61, 55]. Indeed, well-known human "pathogens", such as *N. meningitidis*, *H. influenzae* and *E. Coli* are most commonly found as harmless commensals in hosts. Yet they can cause severe infections under opportune circumstances. Conversely, microbes classically viewed as beneficial, such as *Lactobacillus* and *Bacteroides*, can be harmful when multiplying in sensitive compartments of the host [62, 63]. Besides, 'harm' and 'benefit' are not restricted to direct effects on host tissues. When considering microbe-host interactions holistically, microbes consuming resources ('public goods') in the gut without producing nutrients or services that benefit the host, as well as those aiding the proliferation of virulent microbes [64, 55] also generate indirect costs to their host [65]. In summary, we acknowledge that identifying some microbes as unconditionally beneficial or harmful may seem unrealistic.

However, this dichotomy is not wholly meaningless. Some microbes do have higher predispositions than others to detriment their host [66], because their evolvability is generally constrained. For example, constraints in the availability of metabolic pathways result in convergent patterns of microbe adaptation across hosts producing similar habitats [67]. In addition, as shown by their pervasive use even in recent literature —a search in *The Web of Science* produces 1,782 results for "gut pathogen" for publications from 2024 alone— these labels still carry meaning in the scientific community. Accordingly, the opposition between stereotypically beneficial and harmful microbes is central in hypotheses linking the microbiota with the evolution of adaptive immunity [21] that were proposed by contrast with defence-centric metaphors of immunity as a response to "pathogens". Consequently, in the following we operate with the assumption that from a coarse viewpoint, there exists distinguishable groups of microbes that tend to detriment and benefit their hosts —respectively '**parasites**' and '**mutualists**', following similar existing computational analyses of microbes-host evolution [68, 69, 70].

---

2 Model summary

An overview of the model is presented in figure 1. We assumed that host fitness was entirely determined by the cumulative effect of beneficial and harmful microbes in their gut. Hosts influenced the microbial population in their gut by using either exclusively innate immunity or a combination of innate immunity and SDI. Innate immune reactions only depended on host genotypes and on

the perceived harm caused by the microbiota, the latter mimicking inflammatory responses. Innate immunity was environmentally responsive but deterministic, such that two host clones with the same microbiota would produce the same innate immune response. By contrast, SDI was based on the stochastic population dynamics of multiple lymphocyte lineages within each host, which were selectively reinforced by antigens sampled in the gut. This antigen-mediated feedback was itself biased by the innate immune system. Instead of the common assumption that the principal constraint on immunity is the cost of auto-immune reactions [24, 71], we considered that implicit costs of immunity would arise from immune reactions targeting 'mutualists' and from insufficient immune responses to 'parasites' (see box above for terminology). Thus, we did not assume intrinsic costs of either form of immunity. Hosts reproduced asexually in partially overlapping generations with fitness-proportionate selection. Fitness was updated cumulatively for up to 100 time-steps per host, including an initial developmental period of 25 time-steps where hosts could neither reproduce nor die. We tested multiple dose-response functions associating the density of each microbe species within the gut to its contribution to the host's fitness. In the following, unless specified otherwise beneficial effects are modelled with a logarithmic response function whereas harmful effects are modelled with an exponential response. This aimed to reflect the potential saturating effect of nutritional intake in mutualistic associations, by contrast with common density-threshold effects associated with parasites that often only produce harmful effects at sufficiently high densities [60]. Each reproductive event, host genotypes could mutate to modify their innate immune response to a subset of microbes. This represented an implicit set of innate receptors and associated responses pathways, which we assumed to evolve hand-in-hand. After co-evolutionary dynamics between hosts and microbes stabilised (usually $3e^4$ time-steps; see figure 4) we let SDI invade populations with a low mutation probability. For simplicity, we only considered antagonistic effects of host immunity towards microbes in the gut, assuming positive effects —such as the provision of nutrients like mucin [72]— to be implicit and shared across all microbes.

Figure 1. Overview of the model. We consider a population of hosts representing vertebrate ancestors, whose fitness ω only depends on microbe-host effects β weighted by microbe densities X and a dose-response function f. Each microbe species is subjected to sampling from the environment following a hurdle model with a Bernoulli trial of probability ϵ and normally-distributed amplitude. **E**: Microbe dynamics depend on their growth rate r_i, linear interaction terms α_{ij} with other microbes and host immune effects H_i (null outside of hosts). **A**: Hosts influence the population of microbes in their gut using an immune response H determined by an innate component and a facultative SDI component. Innate immune responses are proportional to the perceived weighted harm inflicted by microbes $\sum \beta_i X_i$, $\beta_i < 0$. **B**: SDI immune responses are formed by a variation-selection process in a population of lymphocytes L_i targeting each microbe species i. Variation is produced by stochastic and antigen-independent influx of lymphocytes $\gamma \cdot \sigma \cdot L$, following a Gaussian hurdle model of probability γ and amplitude σ. Lymphocyte lineages are selectively reinforced based on antigens sampled by the innate immune system in proportion to the gut content $R \cdot X$. **C**: Host can evolve by modifying their innate repertoire R, by modifying the overall intensity of their immune responses λ and by adopting or losing SDI, which is controlled by a discrete gene μ taking values 0 and 0.5. **D**: Microbes evolve by modifying β_i towards parasitism ($\beta_i < 0$) or mutualism ($\beta_i > 0$), which is driven by a positive relationship between parasitism and growth rate within hosts and a negative or neutral relationship between parasitism and growth rate in the external environment.

We let microbes evolve within hosts and in the external environment, such that parasites contributing negatively to host fitness have a higher growth rate within hosts. Microbes that are targeted by the immune system of their host do not benefit from higher growth rates, since their net growth becomes negative. Therefore, the host's immune systems can subject parasites to random drift, displacing them on the mutualism-parasitism spectrum and countering selection for parasitism. However, to maximise the benefits received from exploiting mutualists the hosts must not suppress

them with immune responses, which lets mutualists evolve towards parasitism. This produces a difficult evolutionary challenge for hosts because previously beneficial microbes can gradually become threats, which may prevent the maintenance of long-term mutualism [69]. To constrain and stabilise microbe evolution, we also let microbes evolve in the external environment where microbe-microbe interactions are the principal driver of selection, which constrains and stabilises host-microbe co-evolution. In short, host innate immune systems must keep up with microbes such that current threats are suppressed while beneficial microbes are left alone. In these conditions, the inherently stochastic nature of SDI produces a large direct cost by potentially targeting mutualists —while also delaying immune responses to parasites. In addition, the developmental noise generated by SDI produces a second indirect cost by disrupting the inheritance of fit innate immune systems, hindering their evolution. Conversely, when the innate immune system is error-prone the stochastic (albeit delayed) density-dependent responses of SDI may become beneficial by limiting the risk posed by abundant parasites that are not sufficiently suppressed by host innate repertoires.

We investigated the influence of four principal variables. 1) The number of microbe species in the environment, n, with each 'species' represented by a) a marker i serving as a target for the hosts' immune system; b) a microbe-host interaction term β_i characterising whether this species is beneficial or detrimental to the hosts; and c) a matrix of microbe-microbe interactions $\alpha_{i,j}$ representing competition and cooperation between microbe species. For simplicity, we assumed that microbe-microbe interactions did not change during our simulations because these were not the primary focus of this work. We let microbes evolve by changing the sign and amplitude of their effect β_i on hosts, thus varying on the mutualism-parasitism spectrum. We made the typical assumption that parasites multiplied faster than mutualists within hosts [73, 51, 74, 69], which created a selective pressure for parasitism. Conversely, parasitism did not necessarily increase growth rate in the external environment. 2) Additionally, we varied the relative advantage of parasites and mutualists in the external environment to indirectly control the proportion of beneficial and harmful microbes in our simulations. 3) We also varied the relative intensity of microbe-microbe competition to explore the potential role of competitive exclusion. Finally, 4) we investigated how the dose-response function of the host to the abundance of beneficial microbes influenced the evolution of SDI. A detailed account of the model and parameter choices is given in section Materials and Methods.

3 Results

3.1 A high diversity of beneficial microbes favours the evolution of SDI

As described in the model summary, SDI is costly because it produces stochastic variation in immune responses, whereas innate immunity deterministically targets parasites but only if selection on host genomes can 'keep up' with microbe evolution. However, when the evolution of innate immune systems is outpaced by microbial evolution SDI might be beneficial by limiting the threat posed by parasites that would escape the host's innate immune responses. As a consequence of this trade-off, we find that SDI only evolves in restricted conditions (figure 2). First, SDI only evolves when the diversity of microbe species in the environment and within hosts is high (¿60 species). This supports the general intuition behind the 'care for the microbiota' hypothesis: when the number of microbe species commonly interacting with hosts is low, selection on the innate immune system can rapidly follow changes in microbe populations. Therefore, the costs potentially posed by SDI are high, making innate immunity preferable. Conversely, when the number of microbe species

—each potentially capable of evolutionary switches between mutualism and parasitism— is high, the defence-focused paradigm of adaptive immunity may predict that innate immunity should be overwhelmed, making SDI a better alternative. Our results partially support this view, but with a significant caveat. Indeed, we find that even when microbe diversity is high SDI is rarely favoured when parasites are dominant (figure 2), panel A). In such conditions, hosts with innate immunity tended to tolerate very few microbes while suppressing the vast majority of others, effectively simplifying the evolutionary challenge posed by the larger diversity of microbes (see figure 4, panel D: microbe population evenness is markedly lower without SDI, indicating that few microbe species are dominant while most others are virtually absent). By contrast, we find that SDI is favoured when the proportion of mutualists is high, particularly when microbe-microbe competition is intense. This suggests that the benefits of SDI are more pronounced when the density of any given microbe within host is kept low by competitive interference, such that the risk of harm from any given parasites is lower. In short, when parasites are more dangerous they generate strong selection on the host's innate immune system. This forces hosts to 'keep up', at the cost of collateral immune responses targeting mutualists. In these conditions, SDI is strictly detrimental. Conversely, when parasites are comparatively less threatening they produce weaker selection than mutualists on the immune system of hosts. This leads to an imbalanced trade-off between defence and tolerance favouring SDI as a risk-mitigating strategy in hosts exploiting diverse mutualists.

3.2 SDI only evolves when hosts benefit from exploiting multiple microbe species simultaneously

Hosts generally do not linearly benefit from an increase in the abundance of a given mutualist species, nor do they proportionately benefit from an increase in microbe species richness. Depending on the number of microbiome ecological services that the host relies on, the loss of a single microbe species may be dramatic or fully inconsequential [75]. Borrowing the metaphor of "rivet-redundancy" *vs* "immediate catastrophe" [76] in ecological resilience studies, we analysed the effect of the sensitivity of hosts to microbiome diversity (e.g., from a nutritional perspective) on the evolution of SDI (figure 2, panel B and C). Note, there was always a saturation threshold in the dose-response function. But this saturation could either be reached rapidly, favouring hosts with an evenly distributed microbiota; or slowly and with few benefits for hosts at low microbe abundances, making microbiota with one or few dominant mutualists preferable. We find that SDI only evolved when host fitness was maximised by more even microbiota. Specifically, SDI was favoured by convex benefit functions (either logarithmic or logistic with a low growth rate $s < 1 =$) but never evolved when weighted microbe-host effects followed a concave (either exponential, logistic with $s > 1$ or step-like) shape with low benefits at low densities and higher benefits at higher densities.

This can be interpreted as follows. On the one hand, SDI produces stochastic and density-dependent responses that target abundant microbes, which hinders the ability of hosts to foster any particular mutualist species in high abundances. On the other hand, these same characteristics are ideal to maintain a balanced population of microbes within the host. Consequently, SDI evolved when hosts benefited more from an evenly distributed community of diverse mutualists.

Figure 2. Somatically diversified immunity (SDI) is only favoured by diverse and predominantly beneficial microbial communities. A: We indirectly control the proportion of beneficial microbes by modifying the sign and amplitude of the trade-off incurred by microbes in the external environment: and observe the frequency of SDI after evolutionary dynamics have stabilised. We find that SDI is only likely to evolve when the majority of microbes (at least 55%) are beneficial. In addition, the requirement of a predominantly beneficial microbiota was alleviated by increased microbe-microbe competition (full lines), indicating an effect of competitive exclusion. **B**: We examine the influence of the benefits function of hosts by modifying the parameter s ('Sigmoid factor') controlling the curvature of a dose-response function defined as a Hill function. When s is low, the weighted beneficial effects of microbes increase very rapidly with microbe density but plateau rapidly, such that hosts benefit more from exploiting a high diversity of microbes. When s is high, weighted positive effects from microbes follow an exponential shape such that hosts only receive large benefits from microbes present at high densities. This relationship is shown in **C** by the interaction between microbe community evenness and host fitness with respect to the benefit function. Hosts with a concave (s_i=1 and Logarithmic) response functions benefit from even (high Shannon index) microbiota, whereas hosts with a convex (s_i1 and Exponential) benefit function are favoured by uneven microbiota. We find that SDI only evolves when microbe diversity is high and hosts benefit more from exploiting many mutualists rather than one or few dominant microbes in an uneven community (top right corner of the heatmap).

3.3 Disrupted selection on innate immunity favours SDI and is self-sustaining across long evolutionary time

Results from the previous section show that the conditions favouring SDI are limited to environments with a high diversity of predominantly beneficial microbe species, where microbe-microbe competition

is high (fig 2, panel A) and where hosts benefit more from an evenly distributed community of diverse microbes than from a few dominant mutualists (panel B). These conditions are unlikely to apply across all vertebrate lifestyles and across their entire evolution, even if we assumed that such an environment arose at some point in vertebrate ancestry; unless the fixation of SDI in vertebrate ancestors produced and maintained at least some of the conditions that favour its own maintenance.

3.3.1   SDI disrupts selection on the innate immune system

In figure 4 (panel B), we show that the presence of SDI disturbs selection on the innate immune system, preventing the evolution of receptors that suppress parasites while ignoring mutualists. This is particularly marked when the evolutionary task is complicated by a high proportion of beneficial microbes, such that mutations to the innate immune system that improve defence against a parasite are likely to also increase the suppression of multiple beneficial microbes (fig. 3, right-most panels). In other words, we find that with time the innate immune system becomes increasingly poorer at suppressing parasites while ignoring mutualists in populations of model hosts where SDI has been fixated. Consequently, a loss of SDI and reversion to relying exclusively on a deprecated innate immune system becomes increasingly unlikely as the associated innate immune system becomes gradually incapable of functioning on its own. This is shown in figure (3, panel A) by a rapid plateau in the probability of a decrease in the frequency of SDI as its cumulative frequency (i.e., the time spent by a population of hosts while SDI was fixated) increases. When the number of microbe species is high (> 50) we even find a negative probability of decrease after 500 generations. This shows that mutational noise on the trait 'presence/absence of SDI' eventually becomes incapable of removing SDI from populations of hosts where SDI has been present for sufficiently long.

Figure 3. The persistence of SDI is self-stabilising. **A**: When microbe diversity is
sufficiently high in the host gut microbiota, the probability of loss of SDI plateaus and
eventually decreases in populations where SDI was fixated. **B**: The presence of SDI hinders
the evolution of innate immune receptor repertoires. We assess the performance of innate
immune receptors repertoires with a simple criterion shown in the top row ('Receptor loss':
innate receptors must suppress parasites while ignoring mutualists) in shorter simulations
where microbes did not evolve. Innate repertoires incrementally mutated using this criterion
are indistinguishable from the innate repertoires that would naturally be selected using a
fitness-based evaluation (grey and pink lines, respectively). Conversely, the presence of SDI
disturbs selection on innate repertoires, delaying and lowering the plateau of performance
both expressed as fitness and as 'receptor loss'.

3.3.2 SDI promotes the evolution of a stable and beneficial microbiota

In addition to affecting the evolution of the innate immune system, the presence of SDI markedly
modifies the evolutionary trajectories of the associated microbiota in our simulations (figure 4,

panels D-G). Even when SDI does not strongly affect the average fitness of hosts (e.g., when $n_{microbe\,species} = 40$; panel B) the associated microbiota markedly differs in hosts with SDI, as shown by sharp changes after SDI is allowed to invade by mutation (green line at $t = 3e4$, panels D, E and F). Specifically, microbe populations interacting with hosts relying on SDI become markedly more even (panel D) and more evolutionary stable , as shown by a significant decrease in their mutational variability (panel G). Note, these populations of microbes are still undergoing the same rate of per-generation mutation. However, mutations that provoke a transition from mutualism to parasitism and vice-versa become significantly less likely to be selected in hosts with SDI. In particular, the transition from mutualism to parasitism is markedly suppressed in hosts with SDI when microbe species diversity is high ($n >= 80$). This rapidly increases and stabilises the proportion of beneficial microbes relative to the total quantity of microbes (panel F). In addition to these changes, the evolution of SDI is associated with a rapid decrease in the strength of innate immune defences (fig. 4, panel C). This indicates that, irrespective of which microbes are targeted, suppressive responses from the innate immune system are on average weaker in the presence of SDI. Given that immune suppression from SDI depends on bias from the innate immune system sampling target microbe species from the gut and on the abundance of these targets, this shows a systemic shift from a focus on immune defence to a focus on tolerance in hosts with SDI. In hosts populations with a highly diverse microbiota ($n >= 80$) the decrease in the use of innate immunity to suppress microbes —following the evolution of SDI— is most apparent (panel C when $t > 4e4$). Together with the associated high proportion of beneficial microbes (fig. 4, panel F) and the requirement of high microbe-microbe competition for SDI to evolve (fig. 2), this indicates that competitive exclusion favours the evolution of SDI by gradually decreasing the need for a highly performant and defence-focused immune system, in favour of an immune system capable of tolerating many microbe species while infrequently reacting to potential threats.

Figure 4. Co-evolutionary interactions between SDI and the microbiota. Microbes were subjected to a minimal trade-off in their growth rate between the external environment. Within-host environment and microbe-microbe interactions were competitive on average (interaction term $\alpha_{i,j} \sim \mathcal{N}(0.7, 0.7)$). After evolutionary dynamics between hosts using exclusively innate immunity and their microbiota have stabilized, SDI was allowed to invade populations by mutation (green vertical line). **A**: SDI only evolves when the diversity of microbes (i.e., number of species that can be encountered) is high. **B**: After SDI evolves, there is a marked increase in the fitness of hosts interacting with a diverse microbiota. **C**: The average strength of innate immune defence rapidly decreases in populations where SDI evolves (number of microbe species¿50). **D** and **E**: SDI markedly increases the evenness of microbe populations boss within hosts and to a lesser extent in the external environment. **F** and **G**: SDI markedly increases the proportion of beneficial microbe species in the microbiota and significantly decreases the rates of variance on the mutualism-parasitism scale, particularly when the number of microbe species is high.

4 Discussion

We considered a minimal model of adaptive immunity with limited affinity maturation and recognition specificity (such that adaptive immune cells and innate receptors could in principle be equally specific), without long-term memory and restricted to mucosal and humoral interactions at the gut interface. Our results suggest that even with such restrictions, the selective pressure generated by interactions with the microbiota could induce the evolution of an ancestral form of adaptive immunity, under the following conditions. First, hosts must interact closely with their microbiota, and in such a way that they benefit from a diverse microbial community rather than a single or few symbionts as observed in contemporary invertebrates [77]. Second, the microbiota must be highly diverse and prevalently beneficial. Third, interactions between microbes must be prevalently

competitive, allowing beneficial microbes to protect hosts from parasites in a form of "co-immunity" [27]. Under these conditions, disrupted selection on innate immunity —caused in part by competitive exclusion from beneficial microbes— weakens the defensive function of the innate system. This opens the way for the evolution of a noisy (and therefore costly) form of immunity relying on somatic variation and feedback-mediated reinforcement to suppress dominant microbes, reducing the risk of parasite infection while preserving diversity within the gut community. These initial conditions may seem unlikely, but we find that they could be self-preserving. Hosts interacting closely with their microbiota may favour the preservation of a high proportion of mutualists, and a high proportion of mutualists favours adaptive immunity over innate immunity. Further, our results suggest that adaptive immunity may both reinforce the stability of microbe evolution [78] and disrupt positive selection on innate immunity, making a return to an innate-only immune system improbable.

4.1 A scenario for the evolutionary exception of adaptive immunity

4.1.1 Constraints on invertebrate genomes

To explain the absence of adaptive immunity in invertebrates, multiple authors suggest that evolutionary constraints, rather than selective pressures, facilitated the evolution adaptive immunity in vertebrates while preventing its apparition in invertebrates. Two notable events in vertebrate evolution are often mentioned in such explanations. First, the appearance of RAG-based recombination along with the cellular machinery necessary to generate and selectively reinforce somatically diversified receptor cell lineages has long been considered as an evolutionary "big bang" [19, 16]; that is, an extraordinarily rare event that conditioned the evolution of adaptive immunity. But this view has been challenged by recent evidence of other (convergent) mechanisms of somatic diversification of immune receptors in jawless vertebrates [79, 40] and even in some invertebrates [80, 81]. Similarly, the principles of cell-cell competition required for clonal selection may be more common than previously thought [82, 83]. Conversely, non-vertebrate chordates such as Cephalochordates possess RAG and genes analogues of the vertebrate Major Histocompatibility Complex (MHC) commonly associated with adaptive immunity, but lack functional equivalents of adaptive immunity [84]. Kasahara and colleagues (2010) suggest that a more compelling origin for the emergence of adaptive immunity may be the event of whole-genome duplication (followed by a second duplication in Gnathostomes) separating vertebrates from invertebrate chordates, where adaptive immunity is absent. The authors argued that the resulting increase in genome sizes may have facilitated the emergence of the genomic complexity required to regulate an adaptive immune system [85, 16]. However, this argument suffers from a flaw: in principle, genome-wide duplications could have accelerated the evolution of new innate receptors and response pathways and increased the immune genetic diversity of vertebrates instead of giving rise to adaptive immunity [86, 80]. Unless the ancestral trait of adaptive immunity —for instance, RAG-based somatic variation— produced a very marked fitness benefit, there is no reason to believe that an increase in genome size and genetic diversity would favour adaptive immunity over improved innate immunity. Our results may suggest a solution to this conundrum. We find that interactions with a prevalently mutualistic microbiota could result in a mismatch between innate immunity and microbe populations (figure 3, panel B; the pink and grey curves differ most when mutualists are dominant and microbe diversity is high). This mismatch primarily results from asymmetric selection for tolerance of mutualists over defence from parasites, rather than from constraints on genomes that would hinder the evolution of immune systems. Consequently, a sudden increase in the availability of genetic diversity and novelty though whole-genome duplication may have not improved their innate immune system of hosts, but could have favoured the appearance of

SDI instead.

4.1.2 Vertebrate and invertebrate microbial environment: same but different?

Arguably, ancestral vertebrates and invertebrates likely shared a similar environment; at the very least, there must have been overlaps between the ecological niches of some ancestral invertebrates and some vertebrates [87]. Despite this, differences in the way vertebrate and invertebrate ancestors experience external stimuli and mutational novelty could have induced distinct selective pressures, even in a shared environment. Accordingly, multiple specificities of the vertebrate body plan [88, 89, 12] and metabolism [56, 90] have been proposed as explanations for the evolution of adaptive immunity in vertebrates. For instance, Nakashima and colleagues (2018) remark that unlike invertebrates, for which the gut epithelia is reinforced by chitin and respiratory and digestive tracts are separated, the vertebrate digestive tract is formed of a single layer of cells and joint with the respiratory tract at the pharynx. This may have greatly increased the sensitivity of vertebrate ancestors to the composition of their gut microbiota —and to specific types of parasites such as 'helminths' [91]— while also facilitating the evolution of an immune system relying on the indirect monitoring and control of microbial population by exchanges through a thin epithelium at the pharyngeal and gut interface [41]. Our results offer potential targets to study how such differences in biological systems might influence the response to selection in immunity. For example, the relative permeability of the gut epithelia in vertebrates may constrain the dose-response curve linking mutualist abundance and host benefits, biasing hosts to favour a diverse and even community of microbes, or vice-versa.

Members of the Cephalochordates sub-phylum may be promising animal model candidates for such research. These basal Chordates possess many potential precursors to the emergence of a minimal adaptive immune system in vertebrate ancestors, including a single-layer epithelium, shared nutritional and respiratory function in the pharyngeal cavity, evidence of active phagocytosis, immune effector secretion in gut epithelial cells, as well as analogues of the MHC and RAG genes considered as hallmarks of vertebrate adaptive immunity (see Yuan and colleagues (2015) for a detailed account in Amphioxus). Yet, no evidence of an immune system based on recombination and clonal selection has been found. Whether Cephalochordate ancestors lost a primitive form of adaptive immunity or were subjected to different selective pressures from vertebrates has yet to be determined, to our knowledge.

4.1.3 Explaining the long-term conservation of adaptive immunity

Proposing that adaptive immunity may not have evolved as a superior form of defence [81] raises a secondary question. If adaptive immunity is not widely 'adaptive', how can we explain its conservation across vertebrate phylogeny? Ward and Rosenthal (2014) attribute the impoverished state of innate immune repertoires in vertebrates [93] relative to invertebrates to disrupted selection caused by adaptive immunity, with a proposed shift of innate receptor function from recognition to regulation. We propose a similar selectively neutral explanation, borrowing from the framework of "constructive neutral evolution" [94] and "evolutionary ratchet" [95]. Our results suggest that the exploitation of a diversity of beneficial microbes could have canalised the appearance of adaptive immunity by disrupting selection on innate immune defence, paving the way for a functionally redundant and genetically under-determined system in the form of SDI. Subsequently, both the presence of adaptive immunity and of protective symbionts may have further weakened selection

on the innate immune system and increased the reliance of hosts on adaptive immunity. In our simulations, this is shown by the relative stability of fitness both before and after the appearance of SDI (figure 4, panel B for $n = 60$) despite significant changes in the frequency of SDI (panel A) and in the average strength of innate immune responses (panel C). Crucially, such selectively neutral explanations are potentially robust to temporal and geographical environmental changes that would modify selection on a given trait. In a recent review, Hammer (2024) suggested that some hosts may be "addicted" to their microbiota across evolutionary time-frames. Such an effect has been experimentally demonstrated in *D. melanogaster*, where the presence of protective symbionts has been linked with weakened selection on immune defence [97]. We suggest that such interdependencies may have occurred broadly in vertebrate evolution and could explain the conservation of both adaptive immunity and host-microbiota interactions, even in cases where the evolutionary benefits are apparently unclear.

4.2   Immune systems as constructors of their own selection

The idea that hosts and parasites co-evolve is not novel [98]. However, here we considered microbes not as abstract antagonists in an evolutionary arms race with their hosts, but rather as dynamic components of an ecological community. Recent evidence indeed suggests that eco-evolutionary feedbacks may play a crucial role in co-evolution [99], for instance by imposing both ecological and evolutionary sweeps that produce marked phenotypic change in members of the microbiota [100] and limit genetic diversity, increasing the influence of drift [101]. As such, hosts can influence the evolution of microbes not only by imposing specific selective pressures within their gut but also by controlling the demography of microbial populations. The vertical [102, 69] and horizontal [103] transmission of microbes have been shown to respectively facilitate and disturb these demographic-level forms of control. Accordingly, our results suggest that when hosts exploit microbes well-adapted to the external environment, the bilateral exchange of microbes between hosts and the environment may itself be utilised to control microbe evolution, similarly to the coordinated recruitment and dismissal of microbes in the bobtail squid [104]. Thus, even with small-scale microbe migrations from hosts to the environment, the effect of each host's immunity may alter the distribution of microbes not only within hosts but also in the external environment, even when it contains far more microbes than each host (figure 4, panel E). Accordingly, microbes originating from the vertebrate gut have been isolated in the environment surrounding hosts, including on artificial surfaces [105] and in soils contaminated with faeces [106]. Together, this forms a picture of host-microbiota interactions as a form of reciprocal niche construction, where hosts and microbes may modify their respective environments by disrupting and stabilising the network of interactions composing the microbial community [107, 108].

Our understanding of the interactions between residents of the gut is generally limited to studies in humans and mice relying on temporal and geographical -omics data to infer interactions from non-random patterns of distributions [109]; and the building of complex knowledge-driven models to complete these largely correlation-based methods [110, 111]. However, the integration of various external environments and life-cycles into our view of microbial ecology is still lacking [112] despite recent progress in our understanding of microbiota in non-classic model animals [113] and free-living microbiomes [114]. Recent experiments have demonstrated the potential interest of controlling the demography of the microbiota, showing that k-selected microbiota promoted better health than r-selected microbiota in fish [115]. However, the ecological dynamics of microbes outside of hosts may be just as important for microbe-host co-evolution as within-host dynamics, as suggested by

the importance of the evolutionary trade-off affecting selection of parasitism within and outside of hosts in our model.

4.3   Model limits and future perspectives

To keep the model tractable, it only features a limited representation of adaptive immunity. 'SDI' lacks both long-term memory and the capacity for auto-immune reactions, which are both significant determinants of the impact of adaptive immunity on host fitness [8]. In addition, many functional properties of modern adaptive systems —such as local regulatory control, hypermutation and poly-reactivity— are absent from our model. We also confined our analysis to interactions between the host and the gut microbiota within the gut (or the pharyngeal lumen; see below). This focus on a minimal representation has multiple advantages. First, as stated in the introduction we isolated characteristics of the adaptive immune system that are conserved across most vertebrates —including modern fish [11, 116, 36, 117]— in order to keep our results generalisable and applicable to hypothetical vertebrate ancestors where primitive forms of adaptive immunity emerged. This approach offers a second benefit. Assuming that adaptive immunity emerged as a form of feedback-mediated mucosal immunity vastly reduces the requirements for the regulatory control necessary to prevent auto-immune reactions towards diversified cell types in systemic immunity. In our opinion, this simplified and highly conserved functional view of adaptive immunity therefore provides a parsimonious initial trait upon which subsequent adaptations [16] could have gradually introduced additional complexity, leading to systemic regulatory control, complex somatic diversification and long-term memory as observed in contemporary vertebrates. Coincidentally, while the VLRA/VLRB system of jawless vertebrates is functionally similar to the B and T lymphocyte system in jawed vertebrates, no functional equivalent to the MHC (regulatory self/non-self system) has been found in jawless vertebrates [40]. This may support the hypothesis that mucosal adaptive immunity was an ancestral trait shared by both groups, pre-dating distinct transitions to systemic adaptive immunity. In addition, gut-associated lymphoid tissues (GALT) are highly conserved in vertebrates, which further supports the hypothesis of an ancestral adaptive immune system located at the gut interface [88, 118]. Note, no GALT have been reported in jawless fish where lymphoid tissues are located at the tip of the gills [119]. Whether this invalidates the role of the microbiota in the evolution of jawless fish immunity is debatable, since immune control of the microbiota could have originated in pharyngeal tissues at the interface between what later became gut and respiratory tissues in contemporary vertebrates [120, 84].

A crucial assumption of the model is that beneficial microbes were readily available in the external environment, for instance in soils, water and food; and in greater abundance than potential parasites. Yet, many common beneficial residents of the gut of contemporary vertebrate —such as bacteria from the Bifidobacteria and Lactobacillus groups— do not thrive in the external environment [121, 122]. This might arise from long co-evolution between hosts and select microbial partners, as evidenced by patterns of "phylosymbiosis" [123]; albeit with a strong effect of diet [67]. Indeed, adaptation to living within hosts is usually associated with a marked decrease in genome sizes, preference for anaerobic and warmer environments and specific nutritional requirements such as metabolic pathways specialised in degrading starchy polymers [121, 124]. Long-time microbial partners may be "stuck at the end of the line" [125], such that mechanisms of host control, including diet choice, vertical transmission and metabolic control, might have become sufficient to prevent transitions to parasitism [70]. However, in our model it is not necessary for 'mutualists' to be adapted to benefit their hosts yet. Rather, beneficial microbes were less able to multiply rapidly in

the gut of their hosts, relative to parasites. Mutualists may simply replicate metabolic pathways that they use in the external environments and happen to benefit hosts, for instance by fermenting complex sugars or degrading fibres. Arguably, the real challenge presented by host-microbiota interactions might be to prevent "nomadic" [121] and free-living microbes from taking advantage of their host by evolutionary and plastic changes while ensuring the transmission of long-term mutualist partners across host generations and environments. In this context, the model proposed here may be viewed as a representation of the role of immunity in these early associations that pre-date mutual co-evolutionary adaptations between hosts and microbes, highlighting the need for a better understanding of the evolutionary pressures affecting transient microbial partners that have not yet committed to either a fully mutualistic or parasitic lifestyle. Mechanisms of host control such as the targeted sampling of specific food items [126], active recruitment of target microbe species [127] and direct transmission of mutualists from parent to offspring were absent in our minimal model, but may play a key role in the maintenance of host-microbiota cooperation [72]. For simplicity, our model focused a manageably small number of microbe species. However, such mechanisms of recognition-independent host control might simplify the task of immune control down to a comparably small (i.e., $50 - 100$ OTUs) number in natural systems, and should make the object of further empirical and theoretical work. Likewise, the extent to which the internal environment of ancestral chordates may have differed from the external environment is unknown, but might be inferred from further research in animals with simple body plans [104].

4.4 Conclusion

The present work investigates the potential role of beneficial and antagonistic microbe-host interactions in the ancient evolution of adaptive immunity. Focusing on a minimal model of adaptive immunity, we find the exploitation of a diverse and potentially protective microbiota to be a plausible selective explanation for the appearance and conservation of adaptive immunity. Following this framework, we propose considering stochastic variation in immune receptors and genes —even in the absence of well-identified mechanisms of feedback-mediated clonal reinforcement, for instance alternative splicing of immune proteins in arthropods [128, 16]— as potential candidates for precursory forms of adaptive immunity in non-vertebrate chordates, rather than other traits traditionally viewed as hallmarks of adaptive immunity such as long-term memory and the MHC. In particular, modern comparative approaches to the evolution of adaptive immunity are limited by the lack of species diversity and of ecological variation in jawless fish [129]. Continued research on basal chordates, in which some of these traits may have been overlooked, might inform future theoretical approaches and provide insights into whether our immune system truly evolved to moderate interactions with a diverse and beneficial microbiota.

5 Acknowledgements

The authors thank colleagues and collaborators for their feedback in the design and writing process of this work. This work was supported by the University of St Andrews and funded by the DSTL French-UK PhD scheme.

References

[1] Charles A Janeway et al. "Principles of innate and adaptive immunity". en. In: *Immunobiology: The Immune System in Health and Disease. 5th edition*. Garland Science, 2001. URL: https://www.ncbi.nlm.nih.gov/books/NBK27090/ (visited on 24/05/2025).

[2] Klaus Rajewsky. "Clonal selection and learning in the antibody system". en. In: *Nature* 381.6585 (June 1996), pp. 751–758. ISSN: 0028-0836, 1476-4687. DOI: 10.1038/381751a0. URL: http://www.nature.com/articles/381751a0 (visited on 25/11/2021).

[3] Victor Chardès et al. "Affinity maturation for an optimal balance between long-term immune coverage and short-term resource constraints". In: *Proceedings of the National Academy of Sciences* 119.8 (Feb. 2022), e2113512119. DOI: 10.1073/pnas.2113512119. URL: https://www.pnas.org/doi/10.1073/pnas.2113512119 (visited on 13/11/2024).

[4] Max D. Cooper and Matthew N. Alder. "The Evolution of Adaptive Immune Systems". English. In: *Cell* 124.4 (Feb. 2006), pp. 815–822. ISSN: 0092-8674, 1097-4172. DOI: 10.1016/j.cell.2006.02.001. URL: https://www.cell.com/cell/abstract/S0092-8674(06)00152-8 (visited on 18/03/2025).

[5] Masayuki Hirano et al. "Chapter 4 - The Evolution of Adaptive Immunity in Vertebrates". In: *Advances in Immunology*. Ed. by Frederick W. Alt. Vol. 109. Academic Press, Jan. 2011, pp. 125–157. DOI: 10.1016/B978-0-12-387664-5.00004-2. URL: https://www.sciencedirect.com/science/article/pii/B9780123876645000042 (visited on 17/03/2025).

[6] By Florienne Loder et al. "B Cell Development in the Spleen Takes Place in Discrete Steps and Is Determined by the Quality of B Cell Receptor–Derived Signals". In: *Journal of Experimental Medicine* 190.1 (July 1999), pp. 75–90. ISSN: 0022-1007. DOI: 10.1084/jem.190.1.75. URL: https://doi.org/10.1084/jem.190.1.75 (visited on 28/01/2026).

[7] Jared Feldman et al. "Naive human B cells engage the receptor binding domain of SARS-CoV-2, variants of concern, and related sarbecoviruses". In: *Science Immunology* 6.66 (Oct. 2021), eabl5842. DOI: 10.1126/sciimmunol.abl5842. URL: https://www.science.org/doi/10.1126/sciimmunol.abl5842 (visited on 01/10/2025).

[8] Thomas Pradeu and Louis Du Pasquier. "Immunological memory: What's in a name?" en. In: *Immunological Reviews* 283.1 (2018). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/imr.12652, pp. 7–20. ISSN: 1600-065X. DOI: 10.1111/imr.12652. URL: https://onlinelibrary.wiley.com/doi/abs/10.1111/imr.12652 (visited on 29/09/2022).

[9] Jacob Rimer, Irun R. Cohen and Nir Friedman. "Do all creatures possess an acquired immune system of some sort?" en. In: *BioEssays* 36.3 (2014). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/bies.20 pp. 273–281. ISSN: 1521-1878. DOI: 10.1002/bies.201300124. URL: https://onlinelibrary.wiley.com/doi/abs/10.1002/bies.201300124 (visited on 29/09/2022).

[10] Benjamin Gourbal et al. "Innate immune memory: An evolutionary perspective". en. In: *Immunological Reviews* 283.1 (2018). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/imr.12647, pp. 21–40. ISSN: 1600-065X. DOI: 10.1111/imr.12647. URL: https://onlinelibrary.wiley.com/doi/abs/10.1111/imr.12647 (visited on 09/05/2024).

[11] Susana Magadan, J. Oriol Sunyer and Pierre Boudinot. "Unique Features of Fish Immune Repertoires: Particularities of Adaptive Immunity Within the Largest Group of Vertebrates". en. In: *Pathogen-Host Interactions: Antigenic Variation v. Somatic Adaptations*. Ed. by Ellen Hsu and Louis Du Pasquier. Cham: Springer International Publishing, 2015, pp. 235–264. ISBN: 978-3-319-20819-0. DOI: 10.1007/978-3-319-20819-0_10. URL: https://doi.org/10.1007/978-3-319-20819-0_10 (visited on 21/03/2024).

[12] Gustav van Niekerk, Tanja Davis and Anna-Mart Engelbrecht. "Was the evolutionary road towards adaptive immunity paved with endothelium?" en. In: *Biology Direct* 10.1 (Sept. 2015), p. 47. ISSN: 1745-6150. DOI: 10.1186/s13062-015-0079-0. URL: https://doi.org/10.1186/s13062-015-0079-0 (visited on 08/08/2024).

[13] Akiko Iwasaki and Ruslan Medzhitov. "Control of adaptive immunity by the innate immune system". en. In: *Nature Immunology* 16.4 (Apr. 2015). Number: 4, pp. 343–353. ISSN: 1529-2916. DOI: 10.1038/ni.3123. URL: https://www.nature.com/articles/ni.3123 (visited on 04/04/2022).

[14] Thomas Boehm. "Evolution of Vertebrate Immunity". en. In: *Current Biology* 22.17 (Sept. 2012), R722–R732. ISSN: 0960-9822. DOI: 10.1016/j.cub.2012.07.003. URL: https://www.sciencedirect.com/science/article/pii/S0960982212007944 (visited on 03/04/2023).

[15] Masanori Kasahara. "Genome Duplication and T Cell Immunity". In: *Progress in Molecular Biology and Translational Science*. Ed. by Adrian Liston. Vol. 92. Development of T Cell Immunity. Academic Press, Jan. 2010, pp. 7–36. DOI: 10.1016/S1877-1173(10)92002-4. URL: https://www.sciencedirect.com/science/article/pii/S1877117310920024 (visited on 07/03/2024).

[16] Viktor Müller et al. "An evolutionary perspective on the systems of adaptive immunity". en. In: *Biological Reviews* 93.1 (2018). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/brv.12355, pp. 505–528. ISSN: 1469-185X. DOI: 10.1111/brv.12355. URL: https://onlinelibrary.wiley.com/doi/abs/10.1111/brv.12355 (visited on 15/11/2022).

[17] Mingming Zhao et al. "The mechanisms and factors that induce trained immunity in arthropods and mollusks". In: *Frontiers in Immunology* 14 (2023). ISSN: 1664-3224. URL: https://www.frontiersin.org/journals/immunology/articles/10.3389/fimmu.2023.1241934 (visited on 22/02/2024).

[18] Craig B. Thompson. "New insights into V(D)J recombination and its role in the evolution of the immune system". en. In: *Immunity* 3.5 (Nov. 1995), pp. 531–539. ISSN: 10747613. DOI: 10.1016/1074-7613(95)90124-8. URL: https://linkinghub.elsevier.com/retrieve/pii/1074761395901248 (visited on 02/11/2025).

[19] R M Bernstein et al. "Primordial emergence of the recombination activating gene 1 (RAG1): sequence of the complete shark gene indicates homology to microbial integrases." In: *Proceedings of the National Academy of Sciences* 93.18 (Sept. 1996), pp. 9454–9459. DOI: 10.1073/pnas.93.18.9454. URL: https://www.pnas.org/doi/10.1073/pnas.93.18.9454 (visited on 14/08/2024).

[20] Ian Miller and C. Jessica E. Metcalf. "Evolving resistance to pathogens". EN. In: *Science* (Mar. 2019). DOI: 10.1126/science.aaw8710. URL: https://www.science.org/doi/abs/10.1126/science.aaw8710 (visited on 20/01/2022).

[21] Margaret McFall-Ngai. "Care for the community". en. In: *Nature* 445.7124 (Jan. 2007). Number: 7124, pp. 153–153. ISSN: 1476-4687. DOI: 10.1038/445153a. URL: https://www.nature.com/articles/445153a (visited on 27/09/2022).

[22] Yun Kyung Lee and Sarkis K. Mazmanian. "Has the Microbiota Played a Critical Role in the Evolution of the Adaptive Immune System?" In: *Science* 330.6012 (Dec. 2010), pp. 1768–1773. DOI: 10.1126/science.1195568. URL: https://www.science.org/doi/10.1126/science.1195568 (visited on 27/06/2023).

[23] Daniela Gomez, J. Oriol Sunyer and Irene Salinas. "The mucosal immune system of fish: The evolution of tolerating commensals while fighting pathogens". In: *Fish & Shellfish Immunology* 35.6 (Dec. 2013), pp. 1729–1739. ISSN: 1050-4648. DOI: 10.1016/j.fsi.2013.09.032. URL: https://www.sciencedirect.com/science/article/pii/S1050464813007778 (visited on 18/03/2025).

[24] Craig L. Maynard et al. "Reciprocal interactions of the intestinal microbiota and immune system". en. In: *Nature* 489.7415 (Sept. 2012), pp. 231–241. ISSN: 1476-4687. DOI: 10.1038/nature11551. URL: https://www.nature.com/articles/nature11551 (visited on 18/03/2025).

[25] Katie L. Alexander, Stephan R. Targan and Charles O. Elson III. "Microbiota activation and regulation of innate and adaptive immunity". en. In: *Immunological Reviews* 260.1 (2014). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/imr.12180, pp. 206–220. ISSN: 1600-065X. DOI: 10.1111/imr.12180. URL: https://onlinelibrary.wiley.com/doi/abs/10.1111/imr.12180 (visited on 23/02/2024).

[26] Eduard Ansaldo, Taylor K. Farley and Yasmine Belkaid. "Control of Immunity by the Microbiota". In: *Annual Review of Immunology* 39.1 (2021). _eprint: https://doi.org/10.1146/annurev-immunol-0930 pp. 449–479. DOI: 10.1146/annurev-immunol-093019-112348. URL: https://doi.org/10.1146/annurev-immunol-093019-112348 (visited on 23/02/2024).

[27] Thomas Bazin, Lynn Chiu and Thomas Pradeu. "Host-Microbiota Co-immunity: An Intimate Relationship That Goes Beyond Protection". In: *Philosophy, Theory, and Practice in Biology* 14 (Jan. 2022). DOI: 10.3998/ptpbio.2097. URL: https://hal.science/hal-03883529 (visited on 25/06/2024).

[28] Lora V. Hooper, Dan R. Littman and Andrew J. Macpherson. "Interactions Between the Microbiota and the Immune System". In: *Science* 336.6086 (June 2012), pp. 1268–1273. DOI: 10.1126/science.1223490. URL: https://www.science.org/doi/full/10.1126/science.1223490 (visited on 20/03/2023).

[29] John Chaston and Heidi Goodrich-Blair. "Common trends in mutualism revealed by model associations between invertebrates and bacteria". In: *FEMS Microbiology Reviews* 34.1 (Jan. 2010), pp. 41–58. ISSN: 0168-6445. DOI: 10.1111/j.1574-6976.2009.00193.x. URL: https://doi.org/10.1111/j.1574-6976.2009.00193.x (visited on 17/04/2023).

[30] Zhanshan (Sam) Ma, Wendy Li and Peng Shi. "Microbiome–host-phylogeny relationships in animal gastrointestinal tract microbiomes". In: *FEMS Microbiology Ecology* 98.2 (Feb. 2022), fiac021. ISSN: 0168-6496. DOI: 10.1093/femsec/fiac021. URL: https://doi.org/10.1093/femsec/fiac021 (visited on 20/02/2023).

[31] Jesse C. Nussbaum and Richard M. Locksley. "Infectious (Non)tolerance—Frustrated Commensalism Gone Awry?" en. In: *Cold Spring Harbor Perspectives in Biology* 4.5 (Jan. 2012). Company: Cold Spring Harbor Laboratory Press Distributor: Cold Spring Harbor Laboratory Press Institution: Cold Spring Harbor Laboratory Press Label: Cold Spring Harbor Laboratory Press, a007328. ISSN: , 1943-0264. DOI: 10.1101/cshperspect.a007328. URL: http://cshperspectives.cshlp.org/content/4/5/a007328 (visited on 16/05/2024).

[32] Hiutung Chu and Sarkis K. Mazmanian. "Innate immune recognition of the microbiota promotes host-microbial symbiosis". en. In: *Nature Immunology* 14.7 (July 2013). Number: 7, pp. 668–675. ISSN: 1529-2916. DOI: 10.1038/ni.2635. URL: https://www.nature.com/articles/ni.2635 (visited on 22/02/2024).

[33] Kyoko Takahashi et al. "Epigenetic Control of the Host Gene by Commensal Bacteria in Large Intestinal Epithelial Cells *". English. In: *Journal of Biological Chemistry* 286.41 (Oct. 2011), pp. 35755–35762. ISSN: 0021-9258, 1083-351X. DOI: 10.1074/jbc.M111.271007. URL: https://www.jbc.org/article/S0021-9258(20)73782-8/abstract (visited on 17/03/2025).

[34] Nicole M. Gerardo, Kim L. Hoang and Kayla S. Stoy. "Evolution of animal immunity in the light of beneficial symbioses". In: *Philosophical Transactions of the Royal Society B: Biological Sciences* 375.1808 (Aug. 2020), p. 20190601. DOI: 10.1098/rstb.2019.0601. URL: https://royalsocietypublishing.org/doi/10.1098/rstb.2019.0601 (visited on 25/07/2024).

[35] Kayla S Stoy et al. "Host-associated transmission favors transition of a commensal toward antagonism". In: *Evolution* 77.11 (Nov. 2023), pp. 2512–2521. ISSN: 0014-3820. DOI: 10.1093/evolut/qpad173. URL: https://doi.org/10.1093/evolut/qpad173 (visited on 16/08/2024).

[36] Thomas Boehm and Jeremy B. Swann. "Origin and Evolution of Adaptive Immunity". In: *Annual Review of Animal Biosciences* 2.1 (2014). _eprint: https://doi.org/10.1146/annurev-animal-022513-114201, pp. 259–283. DOI: 10.1146/annurev-animal-022513-114201. URL: https://doi.org/10.1146/annurev-animal-022513-114201 (visited on 18/05/2023).

[37] Matt Pennell et al. "The evolutionary and functional significance of germline immunoglobulin gene variation". English. In: *Trends in Immunology* 44.1 (Jan. 2023), pp. 7–21. ISSN: 1471-4906, 1471-4981. DOI: 10.1016/j.it.2022.11.001. URL: https://www.cell.com/trends/immunology/abstract/S1471-4906(22)00236-8 (visited on 24/01/2024).

[38] Huan Chen et al. "BCR selection and affinity maturation in Peyer's patch germinal centres". en. In: *Nature* 582.7812 (June 2020). Number: 7812, pp. 421–425. ISSN: 1476-4687. DOI: 10.1038/s41586-020-2262-4. URL: https://www.nature.com/articles/s41586-020-2262-4 (visited on 22/01/2024).

[39] Jean-Laurent Casanova. "Adaptive immunity by convergent evolution". en. In: *Nature Reviews Immunology* 18.5 (May 2018). Number: 5, pp. 294–294. ISSN: 1474-1741. DOI: 10.1038/nri.2018.28. URL: https://www.nature.com/articles/nri.2018.28 (visited on 19/05/2023).

[40] Jose Ricardo Morales Poole, Julien Paganini and Pierre Pontarotti. "Convergent evolution of the adaptive immune response in jawed vertebrates and cyclostomes: An evolutionary biology approach based study". en. In: *Developmental & Comparative Immunology*. Impact of high throughput sequencing on comparative immunogenomics 75 (Oct. 2017), pp. 120–126. ISSN: 0145-305X. DOI: 10.1016/j.dci.2017.02.011. URL: https://www.sciencedirect.com/science/article/pii/S0145305X17300885 (visited on 18/05/2023).

[41] Wei-guang Kong et al. "Pharyngeal Immunity in Early Vertebrates Provides Functional and Evolutionary Insight into Mucosal Homeostasis". In: *The Journal of Immunology Author Choice* 203.11 (Dec. 2019), pp. 3054–3067. ISSN: 0022-1767. DOI: 10.4049/jimmunol.1900863. URL: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6859377/ (visited on 18/03/2025).

[42] Andrew J. Macpherson et al. "A Primitive T Cell-Independent Mechanism of Intestinal Mucosal IgA Responses to Commensal Bacteria". en. In: *Science* 288.5474 (June 2000), pp. 2222–2226. ISSN: 0036-8075, 1095-9203. DOI: 10.1126/science.288.5474.2222. URL: https://www.science.org/doi/10.1126/science.288.5474.2222 (visited on 17/03/2025).

[43] Emma Slack, Maria L. Balmer and Andrew J. Macpherson. "B cells as a critical node in the microbiota–host immune system network". en. In: *Immunological Reviews* 260.1 (2014). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/imr.12179, pp. 50–66. ISSN: 1600-065X. DOI: 10.1111/imr.12179. URL: https://onlinelibrary.wiley.com/doi/abs/10.1111/imr.12179 (visited on 22/02/2024).

[44] Andreas Mayer et al. "Regulation of T cell expansion by antigen presentation dynamics". In: *Proceedings of the National Academy of Sciences* 116.13 (Mar. 2019), pp. 5914–5919. DOI: 10.1073/pnas.1812800116. URL: https://www.pnas.org/doi/full/10.1073/pnas.1812800116 (visited on 16/07/2024).

[45] Rebecca A. Clemons, Chase H. Smith and Kelly R. Zamudio. "Primary regulatory T cell activator FOXP3 is present across Amphibia". eng. In: *Immunogenetics* 77.1 (Feb. 2025), p. 15. ISSN: 1432-1211. DOI: 10.1007/s00251-025-01372-0.

[46] Timothy E. O'Sullivan, Joseph C. Sun and Lewis L. Lanier. "Natural Killer Cell Memory". In: *Immunity* 43.4 (Oct. 2015), pp. 634–645. ISSN: 1074-7613. DOI: 10.1016/j.immuni.2015.09.013. URL: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4621966/ (visited on 20/02/2024).

[47] Andrew G. Soerens et al. "Functional T cells are capable of supernumerary cell division and longevity". en. In: *Nature* (Jan. 2023), pp. 1–5. ISSN: 1476-4687. DOI: 10.1038/s41586-022-05626-9. URL: https://www.nature.com/articles/s41586-022-05626-9 (visited on 07/02/2023).

[48] Aakanksha Jain and Chandrashekhar Pasare. "Innate Control of Adaptive Immunity: Beyond the Three-Signal Paradigm". en. In: *The Journal of Immunology* 198.10 (May 2017), pp. 3791–3800. ISSN: 0022-1767, 1550-6606. DOI: 10.4049/jimmunol.1602000. URL: https://www.jimmunol.org/content/198/10/3791 (visited on 25/11/2021).

[49] Rita Pettinello and Helen Dooley. "The Immunoglobulins of Cold-Blooded Vertebrates". en. In: *Biomolecules* 4.4 (Dec. 2014). Number: 4, pp. 1045–1069. ISSN: 2218-273X. DOI: 10.3390/biom4041045. URL: https://www.mdpi.com/2218-273X/4/4/1045 (visited on 18/03/2025).

[50] Guillaume Voisinne et al. "T Cells Integrate Local and Global Cues to Discriminate between Structurally Similar Antigens". In: *Cell Reports* 11.8 (May 2015), pp. 1208–1219. ISSN: 2211-1247. DOI: 10.1016/j.celrep.2015.04.051. URL: https://www.sciencedirect.com/science/article/pii/S2211124715004684 (visited on 16/07/2024).

[51] Kevin R. Foster et al. "The evolution of the host microbiome as an ecosystem on a leash". en. In: *Nature* 548.7665 (Aug. 2017). Number: 7665, pp. 43–51. ISSN: 1476-4687. DOI: 10.1038/nature23292. URL: https://www.nature.com/articles/nature23292 (visited on 11/04/2023).

[52] Yuezhou Chen et al. "Microbial symbionts regulate the primary Ig repertoire". In: *Journal of Experimental Medicine* 215.5 (Mar. 2018), pp. 1397–1415. ISSN: 0022-1007. DOI: 10.1084/jem.20171761. URL: https://doi.org/10.1084/jem.20171761 (visited on 24/01/2024).

[53] Spencer V. Nyholm and Joerg Graf. "Knowing your friends: invertebrate innate immunity fosters beneficial bacterial symbioses". en. In: *Nature Reviews Microbiology* 10.12 (Dec. 2012). Number: 12, pp. 815–827. ISSN: 1740-1534. DOI: 10.1038/nrmicro2894. URL: https://www.nature.com/articles/nrmicro2894 (visited on 17/04/2023).

[54] Nobuhiko Kamada et al. "Control of pathogens and pathobionts by the gut microbiota". en. In: *Nature Immunology* 14.7 (July 2013). Number: 7, pp. 685–690. ISSN: 1529-2916. DOI: 10.1038/ni.2608. URL: https://www.nature.com/articles/ni.2608 (visited on 17/04/2023).

[55] George Hajishengallis and Richard J. Lamont. "Dancing with the Stars: How Choreographed Bacterial Interactions Dictate Nososymbiocity and Give Rise to Keystone Pathogens, Accessory Pathogens, and Pathobionts". English. In: *Trends in Microbiology* 24.6 (June 2016), pp. 477–489. ISSN: 0966-842X, 1878-4380. DOI: 10.1016/j.tim.2016.02.010. URL: https://www.cell.com/trends/microbiology/abstract/S0966-842X(16)00047-0 (visited on 20/04/2025).

[56] Jens Rolff. "Why did the acquired immune system of vertebrates evolve?" eng. In: *Developmental and Comparative Immunology* 31.5 (2007), pp. 476–482. ISSN: 0145-305X. DOI: 10.1016/j.dci.2006.08.009.

[57] Pierre-Olivier Méthot and Samuel Alizon. "What is a pathogen? Toward a process view of host-parasite interactions". In: *Virulence* 5.8 (Oct. 2014), pp. 775–785. ISSN: 2150-5594. DOI: 10.4161/21505594.2014.960726. URL: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4601502/ (visited on 22/11/2022).

[58] Lajos Rózsa, Péter Apari and Viktor Müller. "The microbiome mutiny hypothesis: can our microbiome turn against us when we are old or seriously ill?" en. In: *Biology Direct* 10.1 (Jan. 2015), p. 3. ISSN: 1745-6150. DOI: 10.1186/s13062-014-0034-5. URL: https://doi.org/10.1186/s13062-014-0034-5 (visited on 19/03/2025).

[59] Sho Kitamoto et al. "Regulation of virulence: the rise and fall of gastrointestinal pathogens". en. In: *Journal of Gastroenterology* 51.3 (Mar. 2016), pp. 195–205. ISSN: 1435-5922. DOI: 10.1007/s00535-015-1141-5. URL: https://doi.org/10.1007/s00535-015-1141-5 (visited on 17/04/2023).

[60] Steven T. Rutherford and Bonnie L. Bassler. "Bacterial Quorum Sensing: Its Role in Virulence and Possibilities for Its Control". en. In: *Cold Spring Harbor Perspectives in Medicine* 2.11 (Jan. 2012), a012427. ISSN: , 2157-1422. DOI: 10.1101/cshperspect.a012427. URL: http://perspectivesinmedicine.cshlp.org/content/2/11/a012427 (visited on 19/03/2025).

[61] Bernadette C Young et al. "Severe infections emerge from commensal bacteria by adaptive evolution". In: *eLife* 6 (Dec. 2017). Ed. by Matthew TG Holden, e30637. ISSN: 2050-084X. DOI: 10.7554/eLife.30637. URL: https://doi.org/10.7554/eLife.30637 (visited on 19/03/2025).

[62] D. W. Harty et al. "Pathogenic potential of lactobacilli". eng. In: *International Journal of Food Microbiology* 24.1-2 (Dec. 1994), pp. 179–189. ISSN: 0168-1605. DOI: 10.1016/0168-1605(94)90117-1.

[63] Hassan Zafar and Milton H. Saier. "Gut Bacteroides species in health and disease". In: *Gut Microbes* 13.1 (), p. 1848158. ISSN: 1949-0976. DOI: 10.1080/19490976.2020.1848158. URL: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7872030/ (visited on 20/03/2025).

[64] Emily J. Stevens, Kieran A. Bates and Kayla C. King. "Host microbiota can facilitate pathogen infection". en. In: *PLOS Pathogens* 17.5 (May 2021), e1009514. ISSN: 1553-7374. DOI: 10.1371/journal.ppat.1009514. URL: https://journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.1009514 (visited on 19/03/2025).

[65] Elizabeth J. Culp and Andrew L. Goodman. "Cross-feeding in the gut microbiome: Ecology and mechanisms". English. In: *Cell Host & Microbe* 31.4 (Apr. 2023), pp. 485–499. ISSN: 1931-3128. DOI: 10.1016/j.chom.2023.03.016. URL: https://www.cell.com/cell-host-microbe/abstract/S1931-3128(23)00121-X (visited on 19/03/2025).

[66] Ashley Stengel et al. "Rethinking the Roles of Pathogens and Mutualists: Exploring the Continuum of Symbiosis in the Context of Microbial Ecology and Evolution". In: *Phytobiomes Journal* 6.2 (Jan. 2022), pp. 108–117. DOI: 10.1094/PBIOMES-05-21-0031-P. URL: https://apsjournals.apsnet.org/doi/full/10.1094/PBIOMES-05-21-0031-P (visited on 19/03/2025).

[67] Florent Mazel et al. "Is Host Filtering the Main Driver of Phylosymbiosis across the Tree of Life?" In: *mSystems* 3.5 (Oct. 2018), 10.1128/msystems.00097–18. DOI: 10.1128/msystems.00097-18. URL: https://journals.asm.org/doi/full/10.1128/msystems.00097-18 (visited on 18/04/2024).

[68] Léo Ledru et al. "Mutualists construct the ecological conditions that trigger the transition from parasitism". fr. In: *Peer Community Journal* 2 (2022). ISSN: 2804-3871. DOI: 10.24072/pcjournal.139. URL: https://peercommunityjournal.org/articles/10.24072/pcjournal.139/ (visited on 16/08/2024).

[69] Simon Van Vliet and Michael Doebeli. "The role of multilevel selection in host microbiome evolution". en. In: *Proceedings of the National Academy of Sciences* (2019), pp. 20591–20597. DOI: 10.1073/pnas.1909790116. URL: https://www.pnas.org/doi/10.1073/pnas.1909790116 (visited on 24/05/2024).

[70] Connor Sharp and Kevin R. Foster. "Host control and the evolution of cooperation in host microbiomes". en. In: *Nature Communications* 13.1 (June 2022), p. 3567. ISSN: 2041-1723. DOI: 10.1038/s41467-022-30971-8. URL: https://www.nature.com/articles/s41467-022-30971-8 (visited on 24/05/2024).

[71] Carl T. Bergstrom and Rustom Antia. "How do adaptive immune systems control pathogens while avoiding autoimmunity?" English. In: *Trends in Ecology & Evolution* 21.1 (Jan. 2006), pp. 22–28. ISSN: 0169-5347. DOI: 10.1016/j.tree.2005.11.008. URL: https://www.cell.com/trends/ecology-evolution/abstract/S0169-5347(05)00372-1 (visited on 11/09/2023).

[72] Jacob Wilde, Emma Slack and Kevin R. Foster. "Host control of the microbiome: Mechanisms, evolution, and disease". In: *Science* 385.6706 (July 2024), eadi3338. DOI: `10.1126/science.adi3338`. URL: `https://www.science.org/doi/full/10.1126/science.adi3338` (visited on 04/08/2024).

[73] Marco Archetti et al. "Economic game theory for mutualism and cooperation: Cooperation amongst non-kin and mutualism". en. In: *Ecology Letters* 14.12 (Dec. 2011), pp. 1300–1312. ISSN: 1461023X. DOI: `10.1111/j.1461-0248.2011.01697.x`. URL: `https://onlinelibrary.wiley.com/doi/10.1111/j.1461-0248.2011.01697.x` (visited on 29/07/2024).

[74] Jonas Schluter and Kevin R. Foster. "The Evolution of Mutualism in Gut Microbiota Via Host Epithelial Selection". en. In: *PLOS Biology* 10.11 (Nov. 2012), e1001424. ISSN: 1545-7885. DOI: `10.1371/journal.pbio.1001424`. URL: `https://journals.plos.org/plosbiology/article?id=10.1371/journal.pbio.1001424` (visited on 21/08/2024).

[75] Kayla Cross et al. "Microbiome metabolic capacity is buffered against phylotype losses by functional redundancy". EN. In: *Applied and Environmental Microbiology* (Feb. 2025). DOI: `10.1128/aem.02368-24`. URL: `https://journals.asm.org/doi/10.1128/aem.02368-24` (visited on 10/11/2025).

[76] Matthew R. Orr. "The biodiversity dose-response curve translates theory and practice from ecological restoration into research and clinical priorities for fecal microbiota transplantation". English. In: *Frontiers in Medicine* 9 (Nov. 2022). ISSN: 2296-858X. DOI: `10.3389/fmed.2022.1059148`. URL: `https://www.frontiersin.org/journals/medicine/articles/10.3389/fmed.2022.1059148/full` (visited on 05/11/2025).

[77] Piotr Łukasik and Michał R. Kolasa. "With a little help from my friends: the roles of microbial symbionts in insect populations and communities". In: *Philosophical Transactions of the Royal Society B: Biological Sciences* 379.1904 (May 2024), p. 20230122. DOI: `10.1098/rstb.2023.0122`. URL: `https://royalsocietypublishing.org/doi/full/10.1098/rstb.2023.0122` (visited on 20/08/2024).

[78] João Barroso-Batista, Jocelyne Demengeot and Isabel Gordo. "Adaptive immunity increases the pace and predictability of evolutionary change in commensal gut bacteria". en. In: *Nature Communications* 6.1 (Nov. 2015), p. 8945. ISSN: 2041-1723. DOI: `10.1038/ncomms9945`. URL: `https://www.nature.com/articles/ncomms9945` (visited on 11/03/2024).

[79] Martin F. Flajnik. "Re-evaluation of the Immunological Big Bang". English. In: *Current Biology* 24.21 (Nov. 2014), R1060–R1065. ISSN: 0960-9822. DOI: `10.1016/j.cub.2014.09.070`. URL: `https://www.cell.com/current-biology/abstract/S0960-9822(14)01266-4` (visited on 18/03/2025).

[80] Julie Ghosh et al. "Invertebrate immune diversity". In: *Developmental & Comparative Immunology*. Special issue: In Honour of Gregory Warr 35.9 (Sept. 2011), pp. 959–974. ISSN: 0145-305X. DOI: `10.1016/j.dci.2010.12.009`. URL: `https://www.sciencedirect.com/science/article/pii/S0145305X10003058` (visited on 02/10/2023).

[81] Thomas Pradeu. "Immune System: "Big Bang" in Question". In: *Science* 325.5939 (July 2009), pp. 393–393. DOI: `10.1126/science.325_393a`. URL: `https://www.science.org/doi/10.1126/science.325_393a` (visited on 18/03/2025).

[82] Nicholas E. Baker. "Emerging mechanisms of cell competition". en. In: *Nature Reviews Genetics* 21.11 (Nov. 2020), pp. 683–697. ISSN: 1471-0064. DOI: `10.1038/s41576-020-0262-8`. URL: `https://www.nature.com/articles/s41576-020-0262-8` (visited on 08/03/2025).

[83] Romain Levayer, Barbara Hauert and Eduardo Moreno. "Cell mixing induced by myc is required for competitive tissue invasion and destruction". en. In: *Nature* 524.7566 (Aug. 2015), pp. 476–480. ISSN: 1476-4687. DOI: 10.1038/nature14684. URL: https://www.nature.com/articles/nature14684 (visited on 30/04/2024).

[84] Shaochun Yuan et al. "Amphioxus as a model for investigating evolution of the vertebrate immune system". In: *Developmental & Comparative Immunology*. Specific immunity in invertebrates 48.2 (Feb. 2015), pp. 297–305. ISSN: 0145-305X. DOI: 10.1016/j.dci.2014.05.004. URL: https://www.sciencedirect.com/science/article/pii/S0145305X14001402 (visited on 26/02/2024).

[85] Martin F. Flajnik and Masanori Kasahara. "Origin and evolution of the adaptive immune system: genetic events and selective pressures". en. In: *Nature Reviews Genetics* 11.1 (Jan. 2010). Number: 1, pp. 47–59. ISSN: 1471-0064. DOI: 10.1038/nrg2703. URL: https://www.nature.com/articles/nrg2703 (visited on 18/05/2023).

[86] Allan Force et al. "Preservation of Duplicate Genes by Complementary, Degenerative Mutations". In: *Genetics* 151.4 (Apr. 1999), pp. 1531–1545. ISSN: 1943-2631. DOI: 10.1093/genetics/151.4.1531. URL: https://doi.org/10.1093/genetics/151.4.1531 (visited on 20/03/2023).

[87] Philip C. J. Donoghue and Mark A. Purnell. "The Evolutionary Emergence of Vertebrates From Among Their Spineless Relatives". en. In: *Evolution: Education and Outreach* 2.2 (June 2009), pp. 204–212. ISSN: 1936-6434. DOI: 10.1007/s12052-009-0134-3. URL: https://doi.org/10.1007/s12052-009-0134-3 (visited on 25/06/2024).

[88] T. Matsunaga. "Did the first adaptive immunity evolve in the gut of ancient jawed fish?" In: *Cytogenetics and Cell Genetics* 80.1-4 (July 1998), pp. 138–141. ISSN: 0301-0171. DOI: 10.1159/000014970. URL: https://doi.org/10.1159/000014970 (visited on 21/03/2024).

[89] Keisuke Nakashima et al. "Chitin-based barrier immunity and its loss predated mucus-colonization by indigenous gut microbiota". en. In: *Nature Communications* 9.1 (Aug. 2018), p. 3402. ISSN: 2041-1723. DOI: 10.1038/s41467-018-05884-0. URL: https://www.nature.com/articles/s41467-018-05884-0 (visited on 10/04/2024).

[90] Franziska C. Sandmeier and Richard C. Tracy. "The metabolic pace-of-life model: incorporating ectothermic organisms into the theory of vertebrate ecoimmunology". eng. In: *Integrative and Comparative Biology* 54.3 (Sept. 2014), pp. 387–395. ISSN: 1557-7023. DOI: 10.1093/icb/icu021.

[91] Joseph A. Jackson et al. "Review series on helminths, immune modulation and the hygiene hypothesis: Immunity against helminths and immunological phenomena in modern human populations: coevolutionary legacies?" en. In: *Immunology* 126.1 (Jan. 2009), pp. 18–27. ISSN: 1365-2567. DOI: 10.1111/j.1365-2567.2008.03010.x. URL: https://onlinelibrary.wiley.com/doi/10.1111/j.1365-2567.2008.03010.x (visited on 29/10/2025).

[92] Amanda E. Ward and Benjamin M. Rosenthal. "Evolutionary responses of innate immunity to adaptive immunity". In: *Infection, Genetics and Evolution* 21 (Jan. 2014), pp. 492–496. ISSN: 1567-1348. DOI: 10.1016/j.meegid.2013.12.021. URL: https://www.sciencedirect.com/science/article/pii/S1567134813004644 (visited on 24/01/2024).

[93] Nadia Danilova and Chris T. Amemiya. "Going Adaptive". en. In: *Annals of the New York Academy of Sciences* 1168.1 (2009). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1749-6632.2009.04881 pp. 130–155. ISSN: 1749-6632. DOI: 10.1111/j.1749-6632.2009.04881.x. URL: https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1749-6632.2009.04881.x (visited on 24/01/2022).

[94] T. D. P. Brunet. "Higher level constructive neutral evolution". en. In: *Biology & Philosophy* 37.4 (June 2022), p. 23. ISSN: 1572-8404. DOI: 10.1007/s10539-022-09858-x. URL: https://doi.org/10.1007/s10539-022-09858-x (visited on 05/04/2023).

[95] Julius Lukeš et al. "How a neutral evolutionary ratchet can build cellular complexity". en. In: *IUBMB Life* 63.7 (2011). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/iub.489, pp. 528–537. ISSN: 1521-6551. DOI: 10.1002/iub.489. URL: https://onlinelibrary.wiley.com/doi/abs/10.1002/iub.489 (visited on 26/06/2024).

[96] Tobin J. Hammer. "Why do hosts malfunction without microbes? Missing benefits versus evolutionary addiction". English. In: *Trends in Microbiology* 32.2 (Feb. 2024), pp. 132–141. ISSN: 0966-842X, 1878-4380. DOI: 10.1016/j.tim.2023.07.012. URL: https://www.cell.com/trends/microbiology/abstract/S0966-842X(23)00224-X (visited on 29/10/2025).

[97] Julien Martinez et al. "Addicted? Reduced host resistance in populations with defensive symbionts". In: *Proceedings of the Royal Society B: Biological Sciences* 283.1833 (June 2016), p. 20160778. DOI: 10.1098/rspb.2016.0778. URL: https://royalsocietypublishing.org/doi/10.1098/rspb.2016.0778 (visited on 25/07/2024).

[98] W D Hamilton, R Axelrod and R Tanese. "Sexual reproduction as an adaptation to resist parasites (a review)." In: *Proceedings of the National Academy of Sciences* 87.9 (May 1990), pp. 3566–3573. DOI: 10.1073/pnas.87.9.3566. URL: https://www.pnas.org/doi/abs/10.1073/pnas.87.9.3566 (visited on 18/09/2024).

[99] Guénolé Le Pennec et al. "Demographic fluctuations and selection during host–parasite co-evolution interactively increase genetic diversity". en. In: *Molecular Ecology* 33.10 (2024). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/mec.16939, e16939. ISSN: 1365-294X. DOI: 10.1111/mec.16939. URL: https://onlinelibrary.wiley.com/doi/abs/10.1111/mec.16939 (visited on 18/02/2025).

[100] Benjamin H. Good and Layton B. Rosenfeld. "Eco-evolutionary feedbacks in the human gut microbiome". en. In: *Nature Communications* 14.1 (Nov. 2023), p. 7146. ISSN: 2041-1723. DOI: 10.1038/s41467-023-42769-3. URL: https://www.nature.com/articles/s41467-023-42769-3 (visited on 12/08/2024).

[101] Andrei Papkou et al. "Host–parasite coevolution: why changing population size matters". In: *Zoology*. SI: Host-Parasite Coevolution 119.4 (Aug. 2016), pp. 330–338. ISSN: 0944-2006. DOI: 10.1016/j.zool.2016.02.001. URL: https://www.sciencedirect.com/science/article/pii/S0944200616300071 (visited on 18/02/2025).

[102] Cheng-Yu Chen et al. "Maternal gut microbes shape the early-life assembly of gut microbiota in passerine chicks via nests". In: *Microbiome* 8.1 (Sept. 2020), p. 129. ISSN: 2049-2618. DOI: 10.1186/s40168-020-00896-9. URL: https://doi.org/10.1186/s40168-020-00896-9 (visited on 15/04/2025).

[103] Alexandre R. T. Figueiredo and Jos Kramer. "Cooperation and Conflict Within the Microbiota and Their Effects On Animal Hosts". English. In: *Frontiers in Ecology and Evolution* 8 (May 2020). ISSN: 2296-701X. DOI: 10.3389/fevo.2020.00132. URL: https://www.frontiersin.org/journals/ecology-and-evolution/articles/10.3389/fevo.2020.00132/full (visited on 26/07/2024).

[104] Spencer V. Nyholm and Margaret McFall-Ngai. "A lasting symbiosis: how the Hawaiian bobtail squid finds and keeps its bioluminescent bacterial partner". en. In: *Nature Reviews Microbiology* 19.10 (Oct. 2021), pp. 666–679. ISSN: 1740-1534. DOI: 10.1038/s41579-021-00567-y. URL: https://www.nature.com/articles/s41579-021-00567-y (visited on 23/08/2024).

[105] Hilary P. Browne et al. "Transmission of the gut microbiota: spreading of health". en. In: *Nature Reviews Microbiology* 15.9 (Sept. 2017), pp. 531–543. ISSN: 1740-1534. DOI: 10.1038/nrmicro.2017.50. URL: https://www.nature.com/articles/nrmicro.2017.50 (visited on 16/09/2024).

[106] Joseph S. Palmer et al. "A review of the abundance, behaviour and detection of clostridial pathogens in agricultural soils". en. In: *European Journal of Soil Science* 70.4 (2019). _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/ejss.12847, pp. 911–929. ISSN: 1365-2389. DOI: 10.1111/ejss.12847. URL: https://onlinelibrary.wiley.com/doi/abs/10.1111/ejss.12847 (visited on 16/09/2024).

[107] Renee M. Borges. "Co-niche construction between hosts and symbionts: ideas and evidence". en. In: *Journal of Genetics* 96.3 (July 2017), pp. 483–489. ISSN: 0973-7731. DOI: 10.1007/s12041-017-0792-9. URL: https://doi.org/10.1007/s12041-017-0792-9 (visited on 19/09/2024).

[108] Lynn Chiu and Scott F. Gilbert. "17 - Niche construction and the transition to herbivory: Phenotype switching and the organization of new nutritional modes". In: *Phenotypic Switching*. Ed. by Herbert Levine et al. Academic Press, Jan. 2020, pp. 459–482. ISBN: 978-0-12-817996-3. DOI: 10.1016/B978-0-12-817996-3.00015-3. URL: https://www.sciencedirect.com/science/article/pii/B9780128179963000153 (visited on 19/09/2024).

[109] Meng Luo et al. "Progress on network modeling and analysis of gut microecology: a review". EN. In: *Applied and Environmental Microbiology* (Mar. 2024). DOI: 10.1128/aem.00092-24. URL: https://journals.asm.org/doi/10.1128/aem.00092-24 (visited on 04/11/2025).

[110] Chenhao Li et al. "Predicting microbial interactions through computational approaches". In: *Methods*. Pan-omics analysis of biological data 102 (June 2016), pp. 12–19. ISSN: 1046-2023. DOI: 10.1016/j.ymeth.2016.02.019. URL: https://www.sciencedirect.com/science/article/pii/S1046202315300943 (visited on 26/10/2025).

[111] Mollie Virgo, Serge Mostowy and Brian T. Ho. "Emerging models to study competitive interactions within bacterial communities". In: *Trends in Microbiology* 33.6 (June 2025), pp. 688–700. ISSN: 0966-842X. DOI: 10.1016/j.tim.2024.12.009. URL: https://www.sciencedirect.com/science/article/pii/S0966842X24003251 (visited on 05/11/2025).

[112] Nancy Obeng et al. "Evolution of Microbiota–Host Associations: The Microbe's Perspective". English. In: *Trends in Microbiology* 29.9 (Sept. 2021), pp. 779–787. ISSN: 0966-842X, 1878-4380. DOI: 10.1016/j.tim.2021.02.005. URL: https://www.cell.com/trends/microbiology/abstract/S0966-842X(21)00041-X (visited on 11/04/2024).

[113] Kelsey E. Huus and Ruth E. Ley. "Blowing Hot and Cold: Body Temperature and the Microbiome". EN. In: *mSystems* (Sept. 2021). DOI: 10.1128/msystems.00707-21. URL: https://journals.asm.org/doi/10.1128/msystems.00707-21 (visited on 11/11/2025).

[114] Jack A. Gilbert, Janet K. Jansson and Rob Knight. "Earth Microbiome Project and Global Systems Biology". In: *mSystems* 3.3 (Apr. 2018), 10.1128/msystems.00217–17. DOI: 10.1128/msystems.00217-17. URL: https://journals.asm.org/doi/full/10.1128/msystems.00217-17 (visited on 19/09/2024).

[115] Olav Vadstein et al. "K-Selection as Microbial Community Management Strategy: A Method for Improved Viability of Larvae in Aquaculture". English. In: *Frontiers in Microbiology* 9 (Nov. 2018). ISSN: 1664-302X. DOI: 10.3389/fmicb.2018.02730. URL: https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2018.02730/full (visited on 16/07/2024).

[116] Martin F. Flajnik. "A cold-blooded view of adaptive immunity". en. In: *Nature Reviews Immunology* 18.7 (July 2018). Number: 7, pp. 438–453. ISSN: 1474-1741. DOI: 10.1038/s41577-018-0003-9. URL: https://www.nature.com/articles/s41577-018-0003-9 (visited on 02/10/2023).

[117] J. Oriol Sunyer and Pierre Boudinot. "B-Cell Responses and Antibody Repertoires in Teleost Fish: From Ag Receptor Diversity to Immune Memory and Vaccine Development". en. In: *Principles of Fish Immunology : From Cells and Molecules to Host Protection*. Ed. by Kurt Buchmann and Christopher J. Secombes. Cham: Springer International Publishing, 2022, pp. 253–278. ISBN: 978-3-030-85420-1. DOI: 10.1007/978-3-030-85420-1_8. URL: https://doi.org/10.1007/978-3-030-85420-1_8 (visited on 19/02/2024).

[118] Irene Salinas and Robert D. Miller. "Chapter 10 - Comparative Phylogeny of the Mucosa-Associated Lymphoid Tissue". en. In: *Mucosal Immunology (Fourth Edition)*. Ed. by Jiri Mestecky et al. Boston: Academic Press, Jan. 2015, pp. 145–159. ISBN: 978-0-12-415847-4. DOI: 10.1016/B978-0-12-415847-4.00010-0. URL: https://www.sciencedirect.com/science/article/pii/B9780124158474000100 (visited on 20/03/2023).

[119] Peng Guo et al. "Dual nature of the adaptive immune system in lampreys". en. In: *Nature* 459.7248 (June 2009), pp. 796–801. ISSN: 1476-4687. DOI: 10.1038/nature08068. URL: https://www.nature.com/articles/nature08068 (visited on 07/05/2024).

[120] Christopher J. Lowe et al. "The deuterostome context of chordate origins". en. In: *Nature* 520.7548 (Apr. 2015), pp. 456–465. ISSN: 1476-4687. DOI: 10.1038/nature14434. URL: https://www.nature.com/articles/nature14434 (visited on 08/08/2024).

[121] Rebbeca M. Duar et al. "Lifestyles in transition: evolution and natural history of the genus Lactobacillus". In: *FEMS Microbiology Reviews* 41.Supp_1 (Aug. 2017), S27–S48. ISSN: 0168-6445. DOI: 10.1093/femsre/fux030. URL: https://doi.org/10.1093/femsre/fux030 (visited on 17/04/2025).

[122] Sabrina Duranti et al. "Exploring the Ecology of Bifidobacteria and Their Genetic Adaptation to the Mammalian Gut". en. In: *Microorganisms* 9.1 (Jan. 2021). Number: 1, p. 8. ISSN: 2076-2607. DOI: 10.3390/microorganisms9010008. URL: https://www.mdpi.com/2076-2607/9/1/8 (visited on 17/04/2025).

[123] Ruth E. Ley et al. "Worlds within worlds: evolution of the vertebrate gut microbiota". en. In: *Nature Reviews Microbiology* 6.10 (Oct. 2008). Number: 10, pp. 776–788. ISSN: 1740-1534. DOI: 10.1038/nrmicro1978. URL: https://www.nature.com/articles/nrmicro1978 (visited on 30/01/2023).

[124] Daniela Karasova et al. "Host Species Adaptation of Obligate Gut Anaerobes Is Dependent on Their Environmental Survival". en. In: *Microorganisms* 10.6 (June 2022). Number: 6, p. 1085. ISSN: 2076-2607. DOI: 10.3390/microorganisms10061085. URL: https://www.mdpi.com/2076-2607/10/6/1085 (visited on 29/08/2024).

[125] Georgia C. Drew, Emily J. Stevens and Kayla C. King. "Microbial evolution and transitions along the parasite–mutualist continuum". en. In: *Nature Reviews Microbiology* 19.10 (Oct. 2021), pp. 623–638. ISSN: 1740-1534. DOI: 10.1038/s41579-021-00550-7. URL: https://www.nature.com/articles/s41579-021-00550-7 (visited on 26/07/2024).

[126] Scott M. Villa et al. "Specialized acquisition behaviors maintain reliable environmental transmission in an insect-microbial mutualism". English. In: *Current Biology* 33.13 (July 2023), 2830–2838.e4. ISSN: 0960-9822. DOI: 10.1016/j.cub.2023.05.062. URL: https://www.cell.com/current-biology/abstract/S0960-9822(23)00724-8 (visited on 16/08/2024).

[127] Ramya Ganesan et al. "How It All Begins: Bacterial Factors Mediating the Colonization of Invertebrate Hosts by Beneficial Symbionts". In: *Microbiology and Molecular Biology Reviews* 86.4 (Oct. 2022), e00126–21. DOI: 10.1128/mmbr.00126-21. URL: https://journals.asm.org/doi/full/10.1128/mmbr.00126-21 (visited on 16/08/2024).

[128] Daniela Brites and Louis Du Pasquier. "Somatic and Germline Diversification of a Putative Immunoreceptor within One Phylum: Dscam in Arthropods". en. In: *Pathogen-Host Interactions: Antigenic Variation v. Somatic Adaptations*. Ed. by Ellen Hsu and Louis Du Pasquier. Results and Problems in Cell Differentiation. Cham: Springer International Publishing, 2015, pp. 131–158. ISBN: 978-3-319-20819-0. DOI: 10.1007/978-3-319-20819-0_6. URL: https://doi.org/10.1007/978-3-319-20819-0_6 (visited on 24/01/2024).

[129] Yoichi Sutoh and Masanori Kasahara. "The immune system of jawless vertebrates: insights into the prototype of the adaptive immune system". en. In: *Immunogenetics* 73.1 (Feb. 2021), pp. 5–16. ISSN: 1432-1211. DOI: 10.1007/s00251-020-01182-6. URL: https://doi.org/10.1007/s00251-020-01182-6 (visited on 02/05/2024).

[130] Philip A Waller et al. "Dose-response effect of Bifidobacterium lactis HN019 on whole gut transit time and functional gastrointestinal symptoms in adults". In: *Scandinavian Journal of Gastroenterology* 46.9 (Sept. 2011). _eprint: https://doi.org/10.3109/00365521.2011.584895, pp. 1057–1064. ISSN: 0036-5521. DOI: 10.3109/00365521.2011.584895. URL: https://doi.org/10.3109/00365521.2011.584895 (visited on 10/11/2025).

[131] Ruth Hamilton, Mike Siva-Jothy and Mike Boots. "Two arms are better than one: parasite variation leads to combined inducible and constitutive innate immune responses". In: *Proceedings of the Royal Society B: Biological Sciences* 275.1637 (Apr. 2008), pp. 937–945. DOI: 10.1098/rspb.2007.1574. URL: https://royalsocietypublishing.org/doi/full/10.1098/rspb.2007.1574 (visited on 24/11/2022).

[132] Polly Matzinger et al. "Tolerance, danger, and the extended family". In: *Annual review of immunology* 12.1 (1994), pp. 991–1045.

[133] Thomas Pradeu et al. "The conceptual foundations of innate immunity: Taking stock 30 years later". English. In: *Immunity* 57.4 (Apr. 2024), pp. 613–631. ISSN: 1074-7613. DOI: 10.1016/j.immuni.2024.03.007. URL: https://www.cell.com/immunity/abstract/S1074-7613(24)00128-6 (visited on 16/07/2024).