1 **Proteomic stress response by a novel methanogen enriched from the Great Salt Lake**
2

3 William C Christian[1,2,3], Zackary J Jay[1,2,3], Nikola Tolic[4], Carrie D Nicora[5], Reece Livingstone[1],
4 Stavros Trimmer[1,2,3], Timothy R McDermott[6], Roland Hatzenpichler[1,2,3,7,*]
5

Affiliations
[1]Department of Chemistry and Biochemistry, Montana State University, Bozeman, MT, USA
[2]Thermal Biology Institute, Montana State University, Bozeman, MT, USA
[3]Center for Biofilm Engineering, Montana State University, Bozeman, MT, USA
[4]Environmental Molecular Sciences Directorate, Pacific Northwest National Laboratory,
Richland, WA, USA.
[5]Biological Sciences Directorate, Pacific Northwest National Laboratory, Richland, WA, USA.
[6]Department of Land Resources and Environmental Sciences, Montana State University,
Bozeman, MT, USA
[7]Department of Microbiology and Cell Biology, Montana State University, Bozeman, MT, USA.
[*], corresponding author: R.H., rolandhatzenpichler@gmail.com
17

18 **ORCID numbers**
19 WC, 0000-0003-1682-8083. ZJ, 0000-0003-3062-4933. CDN, 0000-0003-2461-9548. NT, 0000-
20 0003-3950-9130. ST, 0009-0003-3873-4456. TM, 0000-0001-8293-5106. RH, 0000-0002-5489-
21 3444.

23 **Running title:** Proteomic response by a methanogen to stress
24

Abstract
Methanogenic archaea affect the climate through their production of the greenhouse gas,
methane. However, it is unclear how a changing climate and other anthropogenic influences
impact methanogen physiology and consequent methane flux. The Great Salt Lake (GSL) is an
environment that has been heavily impacted by human activity; more than doubling its salt
concentration since the last methanogen was cultured from it in 1985. In this study, we enriched
a novel methanogen, for which we propose the name *Candidatus* Methanohalophilus hillemani,
from the GSL at a time when its salinity reached a historical high. Interestingly, *Ca.* M. hillemani
does not increase expression of energy-conservation or osmo-tolerance proteins when challenged
with salinity or oxygen. In contrast, *Ca.* M. hillemani prioritizes trace metal uptake and immune
functions in response to the presence of the sulfate-reducing bacterium *Desulfovermiculus*. 16S
rRNA gene amplicon data from GSL shore soils with extremely high and variable methane flux
indicated the presence of *Ca.* M. hillemani. Our results show that *Ca.* M. hillemani is active
when challenged with environmental stressors and contributes to the methane flux emanating
from the GSL.

Importance

Methanogens are microbes that affect the climate through their production of the greenhouse gas, methane. Changes in climate and land-use patterns are drying up saline lakes, damaging their unique economic and ecological value. As lake levels across the globe fall, it is unclear how methanogens and the amount of methane they produce will concurrently shift. In this study, we measured high methane output from the Great Salt Lake (GSL) across seasons and identified a novel methanogen as part of a larger methanogenic community that is responsible for these methane emissions. We cultured this novel methanogen from GSL sediments and determined that its methane production was largely unaffected by stress conditions. Our findings indicate that methanogens in saline environments, including a novel cultivated species, may be important sources of methane and will continue to produce methane as salinity increases.

Introduction

Methane is a potent greenhouse gas that has contributed 0.3-0.8 °C to global warming since the start of the industrial revolution [1, 2]. Approximately 60% of methane emitted to the atmosphere is produced by methanogens, archaea that couple the production of methane to energy conservation [2, 3]. Due to the oxygen sensitivity of key enzymes in their metabolism, as well as the comparatively low free-energy yield of methanogenesis, it was long assumed that methanogens are restricted to anoxic environments with large stores of organic carbon. Specifically, wetlands, melting permafrosts, rice paddies, and animal digestive tracts are known for their high methane fluxes [2] and have been extensively studied, together with methanogens therein, due to their influence on climate change. However, many additional ecosystems contribute to the methane cycle. Environments experiencing fast changes due to anthropogenic influence are of particular interest because the magnitude and direction (i.e. emission vs. uptake) of methane flux can change on annual to decadal scales [4-7].

The Great Salt Lake (GSL) and other major terminal lakes around the globe are being altered by human activity [8-10]. Increased upstream water consumption and evaporation due to higher temperatures are causing lake levels to plummet [11-13]. In terminal lakes, a decrease in water volume is accompanied by an increase in salinity. It is well known that halophilic methanogens produce methane under hypersaline conditions [14], however, how they respond to increasing salinity, and how this impacts methane flux, is not well known. Recent studies suggest that increasing salinity in freshwater systems decreases methane emissions [15], but also that droughts will increase methane emissions from once-inundated sediments [16]. Studies specifically examining salinity gradients have shown mixed results, with some reporting no correlation between salinity and methane emissions [17] and others finding a negative relationship [18, 19]. These results have contradictory implications for methane emissions from drying terminal lakes, including the GSL. Unfortunately, data examining emissions from these unique environments is rare and to the best of our knowledge, only one study has quantified methane flux from a terminal lake [20], which measured low and spatially homogenous methane emissions from the GSL. Methane production has been observed from GSL sediments [21], and

a halophilic methanogen, *Methanohalophilus mahii*, was isolated from the GSL in 1985 [22-24].
Despite these observations, many microbial community analyses of the GSL did not find or
mention methanogens [25-27].

M. mahii is a methyl-dismutating methanogen that thrives at 7-12% NaCl [23]. The
Methanohalophilus genus includes six described species and several strains [28, 29]. Besides the
GSL, these species were isolated from deep sea salt brines, deep shale reservoirs, hypersaline
bays, and salt pans [29-33]. Known members of the *Methanohalophilus* genus all share methyl-
dismutating methanogenesis from MMA, DMA, TMA (with one exception [34]) and methanol,
but are incapable of growing via acetoclastic or CO_2-reducing hydrogenotrophic
methanogenesis. The salinity range varies by organism, but they have been shown to grow with
NaCl concentrations from 1% to 20% [28]. They tolerate osmotic stress through both 'salt-in'
strategies with many ion symporters and antiporters, and via the biosynthesis of the compatible
solute, glycine betaine [34]. All members of this genus also universally encode genes for the
detoxification of O_2, including thioredoxins, peroxiredoxins, alkylhydroperoxidases, and
superoxide reductase [28].

In this study, we cultured a novel species of the genus *Methanohalophilus* from the GSL
for which we propose the name *Candidatus* Methanohalophilus hillemani. Using metagenomics
and metaproteomics, we describe its unique features and explore how this organism responds to
environmental stressors, including salinity. Finally, we quantify methane flux from the GSL and
identify the microbes responsible. We combine environmental, microbiological, and meta-omics
techniques to examine how this methanogen responds to changes experienced in the GSL.
101

Materials and methods
Collection of samples and cultivation
Microbialite material was collected in July of 2022 from Bridger Bay in the South arm of
the GSL (Supplementary Figures 1, 2). Methanogens were enriched by adding 10 mM MMA to
0.2 µm filter-sterilized GSL water. Lake water was eventually replaced by media designed by
Paterek and Smith [22], originally used to isolate *Methanohalophilus mahii*, but adjusted to the
salinity of the current GSL (16%).
Methanohalophilus enrichment cultures were grown in Paterek and Smith 1985 media
[22], with differing concentrations of salts and TMA, and without the addition of volatile fatty
acid solution and sodium sulfide. The headspace consisted of a 50/45/5 ratio of $H_2/N_2/CO_2$ at
ambient pressure (~90 kPa in Bozeman, MT). The anoxic media included the following salts to
reflect the salinity of GSL water from April 13[th] 2023: 100 g/L NaCl, 45 g/L $MgCl_2$, 5.5 g/L
KCl, 1.5 g/L $CaCl_2$, 0.13 g/L LiCl, 19 g/L Na_2SO_4, 0.5 g/L NH_4Cl. Additionally, the media
contained (per L): 500 µL of 1 g/L sodium resazurin solution, 10 mL of DSMZ 141 trace
element solution, 23.81 mL of 1 M sodium bicarbonate, 18.87 mL 1 M sodium carbonate, 5 mL
of 40 g/L 2-mercaptoethane sulfonic acid sodium salt (Coenzyme M), and a small spatula tip (1-
2 mg) of sodium dithionite as reductant. The pH was adjusted to ~7.5 and cultures were grown in
9.5 mL of medium in 70 mL serum bottles. Vancomycin and streptomycin (10 µL of 50 mg/mL),

200 µL of 1 M TMA, and 300 µL of carbon source mix [22] were added to each bottle. Bottles
were inoculated with 100 µL from the previous culture and grown at 30 ℃ for approximately
one month prior to transfer.
123
124   **Methane measurements**
125   Methane was measured via GC-FID as previously described [35] with the following
126   modifications: (i) 1 mL of sample was injected into autosampler vials instead of 250 µL, (ii) a 3
127   mL plastic BD syringe equipped with a sample lock was used in place of a Hamilton syringe, and
128   (iii) an Rtx-1 GC Capillary column was used instead of a GS-CarbonPLOT column.
129
Gene amplicon analysis
16S rRNA gene analysis was performed as described in Krukenberg et al. [36] using
Earth Microbiome Project primers 515F and 806R [37, 38]. The mlas-mod-F and *mcrA*-rev-R
primers and the touchdown amplification program from Angel et al. 2012 were used to amplify
the alpha-subunit of methyl-coenzyme M reductase (*mcrA*) [39]. Further details can be found in
the supplementary methods section.
136
Metagenomics
A combination of Illumina short-read and Nanopore long read sequencing, following an
established workflow [35] with some modifications, was used to obtain the circular genome of
Ca. M. hillemani str. BB. Fragmented but complete (>99.5 %) genomes were also generated for
Halanaerobium and *Desulfovermiculus* spp.. The genomes can be found with the following
accession numbers for *Ca.* M. hillemani closed genome, *Desulfovermiculus* sp.*,* and
Halanaerobium sp.*,* respectively: SAMN54119603, SAMN54119604, SAMN54119605. Details
on sequencing, assembly, annotation, and genome comparisons are described in the
supplementary methods section.
146
Salt stress experiment
For methane production and metaproteomics experiments, *Methanohalophilus*
enrichment cultures were grown (i) as the control (176 g/L total salts; 50/45/5 of $H_2/N_2/CO_2$
headspace), (ii) with a 100% N_2 headspace, (iii) in low salt media (114 g/L salts; 50/45/5 of
$H_2/N_2/CO_2$ headspace), (iv) high salt media (227 g/L salts; 50/45/5 of $H_2/N_2/CO_2$ headspace),
and (v) micro-oxic conditions (176 g/L total salts; 50/45/5 of $H_2/N_2/CO_2$ headspace plus 1 mL of
filter-sterilized room air injected into the headspace every four days, achieving ca. 0.35% O_2
upon first injection). The ratio of the different salts for all conditions was the same as described
for the media above, with each salt being decreased or increased proportionally to achieve the
final concentrations. Cultures grown under control conditions were transferred (1:100) into
bottles with these conditions. To acclimate the test cultures to stress conditions, they were grown
under these test conditions for three transfers before methane data was recorded during the fourth
generation. That culture was then used to inoculate a fifth generation, which was harvested in

mid-log phase (based on methane production) for the metaproteomics experiment and the 16S
rRNA gene community analysis (n=3 replicates for each condition).
162

Metaproteomics sample preparation and data analysis

The harvested cultures were sent to the Pacific Northwest National Laboratory (Richland,
WA, USA) where 200 µg of protein was used for digestion. MS analysis of the peptides was
performed using an Orbitrap Exploris 480 mass spectrometer (Thermo Scientific, Waltham, MA,
USA). Peptide matching from the acquired datasets was performed using MSGF+ software [40]
against target meta-proteome composed from sequenced genomes (see Supplementary Methods).
Peptide abundances, integrated as area under the peptide elution peaks, were produced from raw
spectra using in-house developed software MASIC [41]. MASIC peptide abundances were log2
transformed to remove skewness in the distribution of measured abundances. Peptide abundances
were corrected for the relative abundance of organisms using the summed area under the peptide
curves mapping to each organism. Transformed abundance values of peptides, and later proteins,
were normalized using mean central tendency method implemented in InfernoR software [42].
Unless otherwise indicated, all proteomic data presented in this study represents normalized
protein abundances. Raw proteomics data has been deposited at massIVE under accession
number: MSV000100495. Further details can be found in the supplementary methods section.
178

Methane flux measurements

Methane flux data was collected in transects, perpendicular to the shoreline, at Bridger
Bay in the south arm of the GSL using A Li-COR 7810 trace gas analyzer (Li-COR, Lincoln,
NE, USA). Terrestrial flux measurements were made as previously described [43], but with a
one-minute measurement period and a 10 second deadband time. To generate depth-resolved 16S
rRNA gene amplicon data, soil cores were collected from within the boundary of the terrestrial
collar which registered the highest methane flux. Aquatic flux measurements were done using a
custom-made floating flux chamber to take duplicate, one minute, flux measurements. Aquatic
fluxes were calculated from the linear change in methane concentration over time, as previously
described [44].
189

Results and discussion
Enrichment of a novel methanogen

Microbialite and water samples from Bridger Bay in the South arm of GSL were
collected in July 2022, when the water level was at its lowest in recorded history (Supplementary
Figure 3). Methanogens were enriched using the growth medium used to isolate
Methanohalophilus mahii [22]. We increased the salinity of the medium to match GSL water
from April 13th 2023, provided a headspace consisting of 50% H_2, 45% N_2, and 5% CO_2, and
used trimethylamine (TMA) as the methanogenic substrate. The culture also produced methane
from MMA, DMA and methanol, but not from glycine betaine or hydrogen and carbon dioxide
(Supplementary Figure 4).

After 21 months of enrichment, 16S rRNA gene amplicon sequencing indicated that the culture consisted of 63% *Methanohalophilus*, 20% *Desulfovermiculus* (a sulfate reducing bacterium - SRB), and 16% *Halanaerobium* (an anaerobic fermenter). An additional 25 taxa contributed to the remaining <1% of the 16S rRNA gene reads. Metagenomic sequencing of this sample with both Illumina and Oxford Nanopore enabled assembly of a circular *Methanohalophilus* genome (2.09 Mb). Sequence coverage indicated that the culture consisted of 80% *Methanohalophilus*, 7% *Desulfovermiculus* sp., 4% *Halanaerobium* sp., with 9% of sequences remaining unbinned. In addition to these three assembled genomes, the metagenome contained 16S rRNA gene sequences belonging to *Vibrio cholerae*, a *Desulfohalobium* sp., and an *Izimaplasma* sp.. Attempts to isolate the methanogen using dilution to extinction, colony-picking from solid media, and additional antibiotics were not successful.

Characterization of the novel methanogen as *Ca.* Methanohalophilus hillemani

This newly assembled *Methanohalophilus* genome shares less than 95% ANI with all other genomes within its genus, indicating a novel species [45] and only 91.2% with *M. mahii* (Figure 1A). The 16S rRNA gene is less than 99.3% identical to other species, noticeably lower than other *Methanohalophilus* species to each other (Supplementary Figure 5). We propose the name *Candidatus* Methanohalophilus hillemani strain BB for this archaeon (see etymology section in the supplementary online materials).

Comparing protein-coding sequences encoded by *Ca.* M. hillemani to all other *Methanohalophilus* spp. showed it shares the major characteristics of the genus, including methyl-dismutating methanogenesis from MMA, DMA, TMA, and methanol [23, 28, 34 (SI Table 2)]. Like the other species, *Ca.* M. hillemani lacks hydrogenases that would be required for hydrogenotrophic methyl-reducing methanogenesis. Other conserved functions across the genus include glycine betaine transport and synthesis, synthesis and usage of the 22^{nd} amino acid, pyrrolysine (Pyl), and O_2 tolerance via thioredoxins, peroxidases, superoxide reductases, and peroxiredoxin. Unique genetic features of *Ca.* M. hillemani as compared to other members of the genus *Methanohalophilus* are discussed in the Supplementary Results section.

Increasing salinity in the Great Salt Lake

The biomass that led to culturing *Ca.* M. hillemani was collected from the GSL in July 2022, when the lake was at its lowest recorded level (Supplementary Figure 3). Additionally the sample came from Bridger Bay, which was cut off from the main lake by low water levels, leading to a salinity much higher (~24%) than the rest of the south arm of the GSL (~17%). Na^+ and Cl^- concentrations at this time were 3.6 and 4.2 M, respectively, more than twice the concentrations of these ions measured by Post et al. in 1977 (1.6 and 1.8 M) [47]. In contrast, *Methanohalophilus mahii* was isolated in 1985 [22], right before GSL reached recorded high water levels (Supplementary Figure 5). This led us to the hypothesis that *Ca.* M. hillemani may be adapted to higher osmotic stress levels as compared to *M. mahii*.

240
Methane production under salt stress

To understand how stressors in the GSL ecosystem might impact *Ca.* M. hillemani,
control cultures were compared to those incubated under low salt, high salt, 100% N_2, and micro-
oxic conditions (see methods, above). Proportions of major salt ions in all conditions matched
the GSL, which contrasts previous work on the genus that had only modulated NaCl
concentration [22, 29, 48]. The lag time for methane production increased slightly with
increasing salinity (Figure 2A). Nevertheless, the overall methane yield for *Ca.* M. hillemani
cultures was consistent across all conditions tested (Figure 2A). Mass balance of the methyl
groups provided via TMA shows that after 16 days 68 - 77% was converted to methane,
consistent with the 75% value for methyl-dismutating methanogenesis that has been reported for
some euryarchaeotal methanogens [49, 50].
16S rRNA gene amplicon data showed that *Ca.* M. hillemani was the most abundant
member in all conditions (Figure 2B); however, the community composition of these cultures
differed significantly from the 3-species enrichment indicated by the metagenome. The control,
low salt, and high salt conditions contained primarily (>68%) *Ca.* M. hillemani, while
D*esulfovermiculus* was the only other organism with a relative abundance over 0.5%. Over 96%
of amplicons in the O_2 condition belonged to *Ca.* M. hillemani. By contrast, in the N_2 condition
40% of amplicons belonged to *Ca.* M. hillemani and less than 0.1% to *Desulfovermiculus*. An
Izimaplasma species that was absent from all other samples represented 24% of the community,
and the *Halanaerobium* species that made up <0.1% of amplicons in all other conditions,
constituted 30% in the N_2 condition. Despite differences in culturing conditions and community
composition, especially in the N_2 cultures, all conditions yielded the same amount of methane.

263
Metaproteomics

265 To understand the biological basis for *Ca.* M. hillemani's consistent methane production
266 across stressful conditions, we performed metaproteomics. Triplicate cultures were harvested in
267 mid-log phase and peptides from cultures of each condition were mapped to predicted proteins
268 from the assembled genomes. Because no bins could be retrieved from *Izimaplasma* and
269 *Desulfohalobium* from the N_2 condition, we were not able to map peptides to these organisms.

270
Stress conditions do not impact methanogenesis protein expression

For all conditions, the proteins involved in the methyl-dismutating methanogenesis
pathway were among the top 10% of *Methanohalophilus* proteins expressed (Supplementary
Figure 6). The proteins responsible for the transfer of a methyl group to coenzyme M and its
subsequent reduction to methane were all among the top 1%, with one exception
(methylcobamide:CoM methyltransferase – in the top 1.1% of the N_2 condition). The only
methanogenesis-related protein that was differentially expressed was an MMA-methyltransferase
protein found in the control condition but not in the O_2 condition (Figure 3). However, *Ca.* M.

hillemani encodes two copies of this protein, and the other copy was in the top 1% of proteins
expressed in the O_2 condition, suggesting that the methanogenic pathway was not impacted.
It was previously reported that in response to long-term high and low-salt stress,
Methanohalophilus portucalensis transcripts for methylamine methyltransferases and corrinoid
proteins were upregulated [48]. Conversely, we show that the *Ca.* M. hillemani methanogenic
proteome is highly and uniformly expressed, regardless of high or low salt concentrations. One
possible explanation for the lack of observed methanogenic protein expression response, is the
high basal expression of energy-conservation proteins (Supplementary Figure 6). This may be
advantageous in an environment with constant stressors that require large amounts of energy to
tolerate (*i.e.* osmotic and oxidative stress). This strategy could be especially feasible in the GSL,
where methylated amines are likely always in high abundance due to the production of a
metabolic precursor, glycine betaine [51], as a compatible solute.
Another possibility is that methanogenic protein activity, as opposed to abundance, is
post-translationally regulated in response to stress. An investigation of protein phosphorylation
in *Methanohalophilus portucalensis* found that 26% of the phosphorylated proteins were
involved in methanogenesis, energy conservation, and osmolyte biosynthesis [52]. That study
found that a mutated glycine sarcosine N-methyltransferase (responsible for glycine betaine
synthesis), modified to act like a phosphorylated version, had reduced activity that resulted in
lower salinity tolerance when expressed in *E. coli*. While they did not have analogous data for
phosphorylated methanogenic proteins, they demonstrate that *Methanohalophilus* proteins can be
regulated beyond the level of protein expression. The transcriptomic and proteomic findings
discussed here, together with the lack of differential expression of methanogenic proteins,
suggests that future work on the *Methanohalophilus* genus should examine forms of regulation
downstream of protein expression.
303
304 Limited number of proteins differentially expressed in response to changing salinity
Contrary to our expectations, low and high salt conditions resulted in relatively few
proteins that were both highly and differentially expressed, compared to the control (Figure 3).
Both high and low salt conditions showed two upregulated proteins from an iron transport
system, FeoA and FeoB, of which FeoA was not detected in the control. Of the many DNA
repair proteins encoded by *Ca.* M. hillemani, none were both upregulated and expressed above
median protein expression in the high salt condition. The lack of DNA-repair protein expression
in response to high salinity was unexpected [53], though basal expression of several of these
proteins was observed in the control. Alternatively, the need for DNA repair may be mitigated by
osmo-tolerance via high expression of compatible solute. pathways. Indeed, glycine betaine
synthesis and import genes are in the top 10% of expressed proteins in the low-salt, high-salt,
and control conditions. As discussed above, post-translational regulation could also play a role in
osmo-tolerance without requiring differential expression of proteins [52].
Protein expression was more affected by the O_2 growth condition. Two proteins
associated with protein modification and two DNA-repair proteins were up-regulated and

expressed above the median (Figure 3). An additional three DNA-repair proteins were up regulated but not expressed above the median. Thiamine is known to confer antioxidative stress properties; two proteins involved in thiamine biosynthesis, hydroxyethylthiazole kinase and thiamine phosphate synthase, were upregulated in the O_2 condition. Surprisingly, none of the oxygen detoxification genes were upregulated. Constitutive expression could also explain this lack of response to oxygen—superoxide reductase, peroxiredoxin, and catalase-peroxidase are all in the top 10% of expressed proteins in the control.

Unique *Ca.* M. hillemani proteins are differentially expressed in the N_2 condition

When comparing the number of differentially expressed proteins between the control and test conditions, the N_2 condition resulted in twice as many differentially expressed proteins (Supplementary Table 8). This was consistent when filtering for only the proteins that were expressed above the median for each condition or control (Supplementary Table 9). Notably, all 26 differentially and highly expressed *Ca.* M. hillemani proteins in the N_2 condition were less abundant than in the control (Figure 3). Additionally, eight of these 26 differentially and highly expressed proteins are unique proteins to *Ca.* M. hillemani, among the *Methanohalophilus* genus (see Supplementary Results). However, none of these unique and highly expressed proteins were detected in the N☐ condition; in contrast, all eight were found in every other condition, with seven expressed above the median. We hypothesize that these distinct protein expression patterns in the N_2 condition are due to the differences in community composition (Figure 2B).

Metal uptake, DNA repair, and immunity down-regulated in the N_2 condition

Down-regulated functions in the N_2 condition included immune response and DNA repair, cell-surface features, and metal metabolism (Figure 3). A cobaltochelatase and a cobalt/magnesium transporter were not detected and down-regulated 19-fold, respectively, in the N_2 condition (Figure 3). Additionally, one iron transporter was down regulated 29-fold and two other iron transporters, detected in all other conditions, were absent from the N_2 condition.

Highly expressed immune-response and DNA-repair proteins were noticeably absent from the N_2 condition. Seven proteins in this category (Figure 3) were undetectable in the N_2 condition yet were expressed above the median in all other conditions. Eleven additional proteins involved in immune response or DNA repair were also detected in every condition except for N_2. These included two CRISPR-related proteins, two proteins involved in the sulfur-modification of DNA [54], four DNA helicases, two DNA repair proteins, and an additional copy of the type 1 restriction-modification system protein, HsdS.

Ca.* M. hillemani is adapted to sulfide stress and competition with *Desulfovermiculus

The genomic and proteomic data do not support the hypothesis that osmotic tolerance distinguishes *Ca.* M. hillemani and *M. mahii*. Both species are genetically capable of synthesizing and taking up the compatible solute, glycine betaine, and both encode ion transporters to maintain osmotic balance. While *Ca.* M. hillemani encodes unique proteins

involved in DNA repair and immune response, it encodes no unique proteins predicted to
specifically promote osmotolerance (Supplementary Table 10). In fact, differing salt
concentrations resulted in a relatively small proteomic change in this halophilic methanogen.
Instead, the main proteomic differences were observed from the N_2 condition. Immune-
response and DNA-repair proteins, including proteins unique to *Ca.* M. hillemani, were highly
expressed in all conditions except for the N_2 condition. It was previously shown that DNA-repair
transcripts were upregulated by methanogens in response to the presence of sulfide [55].
Additionally, high levels of sulfide can be detrimental to methanogens [56, 57] and known to
damage DNA [58]. Sulfide is produced by SRBs using H_2 as an electron donor, and notably, all
conditions in our study, except for the N_2 condition, contained a headspace with 50% H_2. While
we did not take sulfide measurements during the proteomics experiment, conditions with H_2 in
the headspace yielded cultures with a 15-500 times higher relative abundance of the SRB
Desulfovermiculus, compared to the N_2 condition (Figure 2B). A different SRB,
Desulfohalobium, was present in the N_2 condition at lower relative abundances. However, since
no bin for this genus was retrieved from our metagenome, we were unable to map proteins to this
organism. Together, these results indicate that either high levels of sulfide or the presence of the
specific SRB *Desulfovermiculus* induced a distinct proteomic response by *Ca.* M. hillemani.
Unique immune-response and DNA-repair proteins expressed by *Ca.* M. hillemani in the
presence of *Desulfovermiculus* potentially represent novel adaptations to protect or repair DNA
from SRB-related damage.
Additionally, reduced expression of metal uptake and chelation by *Ca.* M. hillemani in
the N_2 condition, compared to all other conditions, suggests competition for metals with
Desulfovermiculus. Cobalt is crucial for methyl-dismutating methanogenesis as it is the cofactor
in corrinoid proteins used to transfer methyl groups to MCR (Supplementary Figure 6) [59].
However, the conversion of sulfate to sulfide by SRB can directly diminish the availability of
cobalt through Co^{2+} - sulfide co-precipitation [60]. Additionally, SRB utilize cobalt as an
enzymatic cofactor for various metabolisms, including dissimilatory sulfate reduction [61, 62],
which could lead to competition. Indeed, *Desulfovermiculus* expresses two cobaltochelatases in
the control condition, one just below median normalized expression and the other well above.
Tolerating sulfide stress and competing against *Desulfovermiculus* for cobalt may be important
factors impacting *Ca.* M. hillemani physiology when H_2 is available.
390

Ca. M. hillemani uses pyrrolysine sparingly

Methylated amine-utilizing methanogens are known to encode the 22^{nd} amino acid,
pyrrolysine (Pyl), which plays an important role in methylamine activation in tri, di, and
monomethylamine methyltransferases (MttB, MtbB, MtmB, respectively) [63-65]. Indeed, *Ca.*
M. hillemani encodes the genes required for Pyl synthesis (PylBC) and uses (Pyl-tRNA ligase).
An mRNA secondary structure element, PYLIS (analogous to SECIS for selenocysteine), has
been suggested to enable Pyl insertion into proteins [66, 67] and PYLIS elements were annotated
within *Ca.* M. hillemani's methylated-amine methyltransferase genes. Accordingly, Pyl residues

were identified in peptidesmapping to all three types of methylated-amine methyltransferases
encoded by *Ca.* M. hillemani.

Kivenson *et al.* have recently proposed that some methanogens, including
Methanohalophilus mahii, *M. euhalobius*, and *M. levihalophilus,* have reprogrammed the TAG
codon to universally incorporate Pyl, instead of being read as a Stop-codon, across the proteome
[68]. While the authors confirmed their hypothesis for a marine and a human-gut-associated
methanogen, experimental data on halophilic methanogens was not yet available. Moreover, how
and if microbes regulate the incorporation of Pyl at TAG codons is a topic of current debate [68,
69]. Having already identified Pyl residues in peptides mapping to methyltransferase proteins,
we used Kivenson's modification to Prodigal [68], as well as our own modifications to Prokka,
to re-predict and annotate genes for *Ca.* M. hillemani. This resulted in 95 predicted proteins
containing an in-frame TAG codon. In this reanalysis, we found peptide hits from predicted-Pyl-
containing proteins downstream of the predicted-Pyl residue. This could suggest that the TAG
codon is resulting in a Pyl residue, not being read as a Stop-codon. However, for non-
methyltransferase proteins predicted to contain pyrrolysine, only 7 of the 3,463 (0.2%) detected
peptides associated with those proteins contained Pyl. In contrast, 2,920 out of the 27,114
(10.7%) peptides associate with methyltransferase proteins contained Pyl. The low percentage of
non-methyltransferase peptides identified as containing Pyl could indicate that TAG codons, in
contrast to genomic predictions for the genus [68], are primarily read as Stop in *Ca.* M.
hillemani.

High methane flux from the Great Salt Lake
To assess methanogen activity *in situ*, we measured methane flux from the GSL. We
focused on Bridger Bay, where the *Ca.* M. hillemani inoculum was obtained (Supplementary
Figure 2). Methane fluxes were very high but also variable, with fluxes reaching 3 mmol
CH_4/m^2/day observed across sampling dates (Figure 4A, Supplementary Figure 7). These fluxes
are comparable to high-methane flux environments such as thawing permafrost [43, 70] and
freshwater wetlands [71]. The highest flux measured was 377 mmol CH_4/m^2/day (Supplementary
Figure 8), which greatly exceeds values reported from either permafrost or wetlands.

Fluxes were measured in transects perpendicular to the water's edge, including both
terrestrial and aquatic measurements. We repeatedly observed a zone of elevated terrestrial
methane flux between 5 and 15 meters from the shoreline (Figure 4A). This was counterintuitive,
as we expected that the most anoxic and methanogen-conducive conditions to be right at the
shoreline. However, this trend was observed across sampling dates and transects (Supplementary
Figure 7).

Our results contrast the only other study that measured methane flux from the GSL [20],
which observed similar aquatic fluxes (0.42 +/- 0.08 mmol CH_4/m^2/day), but much lower
terrestrial fluxes (0.47 +/- 0.1 mmol CH_4/m^2/day). Their findings of similar terrestrial and aquatic
fluxes and minimal flux variability between sites did not corroborate our results. We hypothesize
that this disparity is due to the difference in spatial scales for our two studies. Cobo *et al.* [20]

examined differences between basins, over kilometer scales. Their transects included steep
vertical gradients over a kilometer from the contemporary GSL shoreline. Our measurements
were on the meter scale and centered on the contemporary lake shore. It is only with our higher
spatial resolution (and smaller scope), that we observed extremely high methane fluxes and flux
variability. Together, Cobo's and our results show that the GSL does have extremely high
methane flux across seasons; however, the zone of highest flux is located just meters from the
current shoreline.
446
Novel methanogens likely contribute to Great Salt Lake methane flux
To examine the ecological relevance of the novel methanogen in the GSL ecosystem, we
performed 16S rRNA and *mcrA* gene amplicon sequencing on samples from a soil core taken
from the transect A 10-meter sample, where extremely high and variable methane flux was
observed (Supplementary Figure 8). The 16S rRNA gene amplicon data revealed that across all
depths, *Ca.* M. hillemani was the second most abundant methanogen, while other
Methanohalophilus sequences were the fourth most abundant (Figure 4C). *Desulfovermiculus*
was the most abundant genus of SRB (Supplementary Figure 9). *Ca.* M. hillemani was similarly
abundant across all depths, and it represented the largest fraction of the methanogenic
community near the surface. This suggests some level of O_2 tolerance, consistent with the higher
relative abundance of *Ca.* M. hillemani in the O_2 condition of our culture experiments (Figure
2B). The most abundant methanogens, by far, were represented by a group of 11 poorly
characterized sequences. A classifier trained on the ARB-silva 16S rRNA gene database could
only identify these sequences as members of the order *Methanofastidiosales* [72], while BLAST
only retrieved "uncultured archaeon clone" sequences. The relative abundance of all
methanogens increased with depth, ranging from ~0.25% at the surface to over 2% of the total
community at the 15-18 cm depth.
The most abundant *mcrA* gene amplicon sequences were also poorly characterized – their
top BLAST hits were to sequences recovered from deep sea anoxic brines (Figure 4C).
Methanohalophilus sequences that could not be assigned to a specific species were the next most
abundant. Of the *Methanohalophilus* sequences that were identified to the species level, *M. mahii*
was the most abundant, followed by *M. portucalensis* and *Ca.* M. hillemani. Like the 16S rRNA
gene amplicon data, the poorly characterized sequences increased in relative abundance with
increasing depth. In both the 16S rRNA and *mcrA* gene amplicon datasets, the most abundant
methanogenic sequences were poorly characterized, indicating that novel methanogens may be
the dominant community in the high-methane-flux GSL soil.
473
Conclusion
Ca. Methanohalophilus hillemani is a novel methanogen, collected from the south arm of
the GSL when its salinity was at an all-time high. Metaproteomic analysis of this organism
showed that methanogenesis and osmo-tolerance proteins had high basal expression. The
expression of these proteins was not impacted by salinity changes, suggesting that future
proteomic studies of the *Methanohalophilus* genus should focus on post-translational

modification and other forms of regulation, for example on protein activity or metabolome level.
Large differences in the expressed proteome were instead observed in the N_2 condition, which
lacked a large population of the SRB, *Desulfovermiculus*. In conditions under which *Ca.* M.
hillemani and *Desulfovermiculus* were both present, the methanogen highly expressed metal
transporters and chelatases. We hypothesize that, with sufficient H_2, SRB directly compete for
and reduce the availability of trace metals, forcing *Ca.* M. hillemani to invest in uptake of metals
vital to its energy metabolism. Coculturing of the two also led to the upregulation of unique
immune-response and DNA-repair proteins by *Ca.* M. hillemani, indicating the adaptation of
novel strategies to tolerate stress and/or *Desulfovermiculus*. High methane flux was measured
from the shore of the GSL, and amplicon data of a soil core identified *Ca.* M. hillemani to be
present at significant levels, confirming the ecological relevance of this novel methanogen. In
summary, this study cultured a novel methanogen from the GSL, demonstrated that its
physiology may not be controlled as much by changes in salinity as by community interactions,
and identified it in GSL soils exhibiting high methane flux.
494

<u>Acknowledgements</u>
This work was supported by the NASA Exobiology award 80NSSC21K0487 (to T.R.M.) and
Department of Energy, Office of Science, Biological and Environmental Research award DE-
SC0025661 (to R.H.). A portion of this research was performed under FICUS award 509790
(DOI:10.46936/fics.proj.2023.60847/60008905) and used resources at the Environmental
Molecular Sciences Laboratory, a DOE Office of Science User Facility sponsored by the
Biological and Environmental Research program under contract no. DE-AC05-76RL01830.
Battelle operates PNNL for the DOE under contract DE-AC0576RLO01830. We thank Bonnie
Baxter (Westminster University) for her guidance on field sampling, Angel Nieto and Ciara Pike
for assistance during field work, Veronika Kivenson (UC Berkeley) for helpful discussions on
pyrrolysine and alternative codon use, and the laboratory of Samantha Joye (University of
Georgia) for help in quantifying major ions in GSL water. Finally, we thank the family of the
late Dr. Maurice Hilleman for allowing us to name the newly cultured methanogen after him in
recognition of his life-saving contributions to microbiology, vaccine development, and public
health (see Supplementary Etymology).

Contributions
Conceptualization: W.C.C., T.R.M. and R.H.
Methodology: W.C.C., T.R.M. and R.H.
Validation: W.C.C., Z.J.J., N.T., T.R.M., and R.H.
Formal analysis: W.C.C., Z.J.J. and N.T.
Investigation: W.C.C., C.D.N., S.T. and R.L.
Resources: T.R.M. and R.H.
Data curation: W.C.C., Z.J.J., and N.T.
Writing–original draft: W.C.C.

Writing–review and editing: W.C.C., Z.J.J., N.T., T.R.M., and R.H.

Visualization: W.C.C. and Z.J.J.

Supervision: T.R.M. and R.H.

Project administration: T.R.M. and R.H.

Funding acquisition: T.R.M. and R.H.

References

1. *Summary for Policymakers*, in *Climate Change 2021 – The Physical Science Basis:*
Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental
Panel on Climate Change, C. Intergovernmental Panel on Climate, Editor. 2023,
Cambridge University Press: Cambridge. p. 3-32.

2. Saunois, M., Martinez, A., Poulter, B., Zhang, Z., Raymond, P.A., Regnier, P., Canadell,
J.G., Jackson, R.B., Patra, P.K., Bousquet, P., Ciais, P., Dlugokencky, E.J., Lan, X.,
Allen, G.H., Bastviken, D., Beerling, D.J., Belikov, D.A., Blake, D.R., Castaldi, S.,
Crippa, M., Deemer, B.R., Dennison, F., Etiope, G., Gedney, N., Höglund-Isaksson, L.,
Holgerson, M.A., Hopcroft, P.O., Hugelius, G., Ito, A., Jain, A.K., Janardanan, R.,
Johnson, M.S., Kleinen, T., Krummel, P.B., Lauerwald, R., Li, T., Liu, X., McDonald,
K.C., Melton, J.R., Mühle, J., Müller, J., Murguia-Flores, F., Niwa, Y., Noce, S., Pan, S.,
Parker, R.J., Peng, C., Ramonet, M., Riley, W.J., Rocher-Ros, G., Rosentreter, J.A.,
Sasakawa, M., Segers, A., Smith, S.J., Stanley, E.H., Thanwerdas, J., Tian, H., Tsuruta,
A., Tubiello, F.N., Weber, T.S., van der Werf, G.R., Worthy, D.E.J., Xi, Y., Yoshida, Y.,
Zhang, W., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q., *Global Methane Budget 2000–*
2020. Earth Syst. Sci. Data, 2025. **17**(5): p. 1873-1958.

3. Conrad, R., *The global methane cycle: recent advances in understanding the microbial*
processes involved. Environmental Microbiology Reports, 2009. **1**(5): p. 285-292.

4. Keuschnig, C., Larose, C., Rudner, M., Pesqueda, A., Doleac, S., Elberling, B., Björk,
R.G., Klemedtsson, L., and Björkman, M.P., *Reduced methane emissions in former*
permafrost soils driven by vegetation and microbial changes following drainage. Global
Change Biology, 2022. **28**(10): p. 3411-3425.

5. Wang, N., Zhu, X., Zuo, Y., Liu, J., Yuan, F., Guo, Z., Zhang, L., Sun, Y., Gong, C.,
Guo, D., Song, C., and Xu, X., *Microbial mechanisms for methane source-to-sink*
transition after wetland conversion to cropland. Geoderma, 2023. **429**: p. 116229.

6. Shoemaker, J.K., Keenan, T.F., Hollinger, D.Y., and Richardson, A.D., *Forest ecosystem*
changes from annual methane source to sink depending on late summer water balance.
Geophysical Research Letters, 2014. **41**(2): p. 673-679.

7. He, T., Ding, W., Cheng, X., Cai, Y., Zhang, Y., Xia, H., Wang, X., Zhang, J., Zhang, K.,
and Zhang, Q., *Meta-analysis shows the impacts of ecological restoration on greenhouse*
gas emissions. Nature Communications, 2024. **15**(1): p. 2668.

8. Stone, R., *Saving Iran's great salt lake.* Science, 2015. **349**(6252): p. 1044-1047.

9.   Aladin, N., Plotnikov, I., and T, B., *The Aral Sea: water level, salinity and long-term changes in biological communities of an endangered ecosystem - past, present and future*. 2008.

10.   Patel, K. *Getting Saltier*. 2019.

11.   Baxter, B.K. and Butler, J.K., *Climate Change and Great Salt Lake*, in *Great Salt Lake Biology: A Terminal Lake in a Time of Change*, B.K. Baxter and J.K. Butler, Editors. 2020, Springer International Publishing: Cham. p. 23-52.

12.   Wurtsbaugh, W.A., Miller, C., Null, S.E., DeRose, R.J., Wilcock, P., Hahnenberger, M., Howe, F., and Moore, J., *Decline of the world's saline lakes*. Nature Geoscience, 2017. **10**(11): p. 816-821.

13.   Wang, J., Song, C., Reager, J.T., Yao, F., Famiglietti, J.S., Sheng, Y., MacDonald, G.M., Brun, F., Schmied, H.M., Marston, R.A., and Wada, Y., *Recent global decline in endorheic basin water storages*. Nature Geoscience, 2018. **11**(12): p. 926-932.

14.   McGenity, T.J. and Sorokin, D.Y., *Methanogens and Methanogenesis in Hypersaline Environments*, in *Biogenesis of Hydrocarbons*, A.J.M. Stams and D.Z. Sousa, Editors. 2019, Springer International Publishing: Cham. p. 283-309.

15.   Soued, C., Bogard, M.J., Finlay, K., Bortolotti, L.E., Leavitt, P.R., Badiou, P., Knox, S.H., Jensen, S., Mueller, P., Lee, S.C., Ng, D., Wissel, B., Chan, C.N., Page, B., and Kowal, P., *Salinity causes widespread restriction of methane emissions from small inland waters*. Nature Communications, 2024. **15**(1): p. 717.

16.   Paranaíba, J.R., Aben, R., Barros, N., Quadra, G., Linkhorst, A., Amado, A.M., Brothers, S., Catalán, N., Condon, J., Finlayson, C.M., Grossart, H.P., Howitt, J., Oliveira Junior, E.S., Keller, P.S., Koschorreck, M., Laas, A., Leigh, C., Marcé, R., Mendonça, R., Muniz, C.C., Obrador, B., Onandia, G., Raymundo, D., Reverey, F., Roland, F., Rõõm, E.I., Sobek, S., von Schiller, D., Wang, H., and Kosten, S., *Cross-continental importance of CH(4) emissions from dry inland-waters*. Sci Total Environ, 2022. **814**: p. 151925.

585   17.   Welti, N., Hayes, M., and Lockington, D., *Seasonal nitrous oxide and methane emissions across a subtropical estuarine salinity gradient*. Biogeochemistry, 2017. **132**(1): p. 55-69.

588   18.   Patel, L., Gowd, S.C., Singh, R., and Thottathil, S.D., *Salinity and phosphorus as key regulators of methane emissions in tropical aquaculture ponds*. Aquaculture International, 2025. **33**(4): p. 302.

591   19.   Hartman Wyatt, H., Bueno de Mesquita Clifton, P., Theroux Susanna, M., Morgan-Lang, C., Baldocchi Dennis, D., and Tringe Susannah, G., *Multiple microbial guilds mediate soil methane cycling along a wetland salinity gradient*. mSystems, 2024. **9**(1): p. e00936-23.

595   20.   Cobo, M., Goldhammer, T., and Brothers, S., *A desiccating saline lake bed is a significant source of anthropogenic greenhouse gas emissions*. One Earth, 2024. **7**(8): p. 1414-1423.

598   21.   Baxter, B.K. and Zalar, P., *Chapter 4 - The extremophiles of Great Salt Lake: Complex*
599   *microbiology in a dynamic hypersaline ecosystem*, in *Model Ecosystems in Extreme*
600   *Environments*, J. Seckbach and P. Rampelotto, Editors. 2019, Academic Press. p. 57-99.
601   22.   Paterek, J.R. and Smith, P.H., *Isolation and characterization of a halophilic methanogen*
602   *from great salt lake.* Appl Environ Microbiol, 1985. **50**(4): p. 877-81.
603   23.   Spring, S., Scheuner, C., Lapidus, A., Lucas, S., Glavina Del Rio, T., Tice, H., Copeland,
604   A., Cheng, J.F., Chen, F., Nolan, M., Saunders, E., Pitluck, S., Liolios, K., Ivanova, N.,
605   Mavromatis, K., Lykidis, A., Pati, A., Chen, A., Palaniappan, K., Land, M., Hauser, L.,
606   Chang, Y.J., Jeffries, C.D., Goodwin, L., Detter, J.C., Brettin, T., Rohde, M., Göker, M.,
607   Woyke, T., Bristow, J., Eisen, J.A., Markowitz, V., Hugenholtz, P., Kyrpides, N.C., and
608   Klenk, H.P., *The genome sequence of Methanohalophilus mahii SLP(T) reveals*
609   *differences in the energy metabolism among members of the Methanosarcinaceae*
610   *inhabiting freshwater and saline environments.* Archaea, 2010. **2010**: p. 690737.
611   24.   PATEREK, J.R. and SMITH, P.H., *Methanohalophilus mahii gen. nov., sp. nov., a*
612   *Methylotrophic Halophilic Methanogen†*. International Journal of Systematic and
613   Evolutionary Microbiology, 1988. **38**(1): p. 122-123.
614   25.   Tazi, L., Breakwell, D.P., Harker, A.R., and Crandall, K.A., *Life in extreme*
615   *environments: microbial diversity in Great Salt Lake, Utah.* Extremophiles, 2014. **18**(3):
616   p. 525-35.
617   26.   Shoemaker, A., Maritan, A., Cosar, S., Nupp, S., Menchaca, A., Jackson, T., Dang, A.,
618   Baxter, B.K., Colman, D.R., Dunham, E.C., and Boyd, E.S., *Wood–Ljungdahl pathway*
619   *encoding anaerobes facilitate low-cost primary production in hypersaline sediments at*
620   *Great Salt Lake, Utah.* FEMS Microbiology Ecology, 2024. **100**(8).
621   27.   Kanik, M., Munro-Ehrlich, M., Fernandes-Martins Maria, C., Payne, D., Gianoulias, K.,
622   Keller, L., Kubacki, A., Lindsay Melody, R., Baxter Bonnie, K., Vanden Berg Michael,
623   D., Colman Daniel, R., and Boyd Eric, S., *Unexpected Abundance and Diversity of*
624   *Phototrophs in Mats from Morphologically Variable Microbialites in Great Salt Lake,*
625   *Utah.* Applied and Environmental Microbiology, 2020. **86**(10): p. e00165-20.
626   28.   Guan, Y., Ngugi, D.K., Vinu, M., Blom, J., Alam, I., Guillot, S., Ferry, J.G., and Stingl,
627   U., *Comparative Genomics of the Genus Methanohalophilus, Including a Newly Isolated*
628   *Strain From Kebrit Deep in the Red Sea.* Frontiers in Microbiology, 2019. **10**.
629   29.   L'Haridon, S., Haroun, H., Corre, E., Roussel, E., Chalopin, M., Pignet, P., Balière, C., la
630   Cono, V., Jebbar, M., Yakimov, M., and Toffin, L., *Methanohalophilus profundi sp. nov.,*
631   *a methylotrophic halophilic piezophilic methanogen isolated from a deep hypersaline*
632   *anoxic basin.* Systematic and Applied Microbiology, 2020. **43**(5): p. 126107.
633   30.   Davidova, I.A., Harmsen, H.J.M., Stams, A.J.M., Belyaev, S.S., and Zehnder, A.J.B.,
634   *Taxonomic description of Methanococcoides euhalobius and its transfer to the*
635   *Methanohalophilus genus.* Antonie van Leeuwenhoek, 1997. **71**(4): p. 313-318.
636   31.   Katayama, T., Yoshioka, H., Mochimaru, H., Meng, X.Y., Muramoto, Y., Usami, J.,
637   Ikeda, H., Kamagata, Y., and Sakata, S., *Methanohalophilus levihalophilus sp. nov., a*

slightly halophilic, methylotrophic methanogen isolated from natural gas-bearing deep aquifers, and emended description of the genus Methanohalophilus. Int J Syst Evol Microbiol, 2014. **64**(Pt 6): p. 2089-2093.

32. Wilharm, T., Zhilina, T.N., and Hummel, P., *DNA-DNA Hybridization of Methylotrophic Halophilic Methanogenic Bacteria and Transfer of Methanococcus halophilusvp to the Genus Methanohalophilus as Methanohalophilus halophilus comb. nov.* International Journal of Systematic and Evolutionary Microbiology, 1991. **41**(4): p. 558-562.

33. Boone, D.R., Mathrani, I.M., Liu, Y., Menaia, J.A.G.F., Mah, R.A., and Boone, J.E., *Isolation and Characterization of Methanohalophilus portucalensis sp. nov. and DNA Reassociation Study of the Genus Methanohalophilus.* International Journal of Systematic and Evolutionary Microbiology, 1993. **43**(3): p. 430-437.

34. Borton, M.A., Daly, R.A., O'Banion, B., Hoyt, D.W., Marcus, D.N., Welch, S., Hastings, S.S., Meulia, T., Wolfe, R.A., Booker, A.E., Sharma, S., Cole, D.R., Wunch, K., Moore, J.D., Darrah, T.H., Wilkins, M.J., and Wrighton, K.C., *Comparative genomics and physiology of the genus Methanohalophilus, a prevalent methanogen in hydraulically fractured shale.* Environmental Microbiology, 2018. **20**(12): p. 4596-4611.

35. Kohtz, A.J., Petrosian, N., Krukenberg, V., Jay, Z.J., Pilhofer, M., and Hatzenpichler, R., *Cultivation and visualization of a methanogen of the phylum Thermoproteota.* Nature, 2024. **632**(8027): p. 1118-1123.

36. Krukenberg, V., Reichart, N.J., Spietz, R.L., and Hatzenpichler, R., *Microbial Community Response to Polysaccharide Amendment in Anoxic Hydrothermal Sediments of the Guaymas Basin.* Frontiers in Microbiology, 2021. **Volume 12 - 2021**.

37. Parada, A.E., Needham, D.M., and Fuhrman, J.A., *Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples.* Environ Microbiol, 2016. **18**(5): p. 1403-14.

38. Thompson, L.R., Sanders, J.G., McDonald, D., Amir, A., Ladau, J., Locey, K.J., Prill, R.J., Tripathi, A., Gibbons, S.M., Ackermann, G., Navas-Molina, J.A., Janssen, S., Kopylova, E., Vázquez-Baeza, Y., González, A., Morton, J.T., Mirarab, S., Zech Xu, Z., Jiang, L., Haroon, M.F., Kanbar, J., Zhu, Q., Jin Song, S., Kosciolek, T., Bokulich, N.A., Lefler, J., Brislawn, C.J., Humphrey, G., Owens, S.M., Hampton-Marcell, J., Berg-Lyons, D., McKenzie, V., Fierer, N., Fuhrman, J.A., Clauset, A., Stevens, R.L., Shade, A., Pollard, K.S., Goodwin, K.D., Jansson, J.K., Gilbert, J.A., Knight, R., Rivera, J.L.A., Al-Moosawi, L., Alverdy, J., Amato, K.R., Andras, J., Angenent, L.T., Antonopoulos, D.A., Apprill, A., Armitage, D., Ballantine, K., Bárta, J., Baum, J.K., Berry, A., Bhatnagar, A., Bhatnagar, M., Biddle, J.F., Bittner, L., Boldgiv, B., Bottos, E., Boyer, D.M., Braun, J., Brazelton, W., Brearley, F.Q., Campbell, A.H., Caporaso, J.G., Cardona, C., Carroll, J., Cary, S.C., Casper, B.B., Charles, T.C., Chu, H., Claar, D.C., Clark, R.G., Clayton, J.B., Clemente, J.C., Cochran, A., Coleman, M.L., Collins, G., Colwell, R.R., Contreras, M., Crary, B.B., Creer, S., Cristol, D.A., Crump, B.C., Cui, D., Daly, S.E., Davalos, L., Dawson, R.D., Defazio, J., Delsuc, F., Dionisi, H.M., Dominguez-Bello,

M.G., Dowell, R., Dubinsky, E.A., Dunn, P.O., Ercolini, D., Espinoza, R.E., Ezenwa, V.,
Fenner, N., Findlay, H.S., Fleming, I.D., Fogliano, V., Forsman, A., Freeman, C.,
Friedman, E.S., Galindo, G., Garcia, L., Garcia-Amado, M.A., Garshelis, D., Gasser,
R.B., Gerdts, G., Gibson, M.K., Gifford, I., Gill, R.T., Giray, T., Gittel, A., Golyshin, P.,
Gong, D., Grossart, H.-P., Guyton, K., Haig, S.-J., Hale, V., Hall, R.S., Hallam, S.J.,
Handley, K.M., Hasan, N.A., Haydon, S.R., Hickman, J.E., Hidalgo, G., Hofmockel,
K.S., Hooker, J., Hulth, S., Hultman, J., Hyde, E., Ibáñez-Álamo, J.D., Jastrow, J.D., Jex,
A.R., Johnson, L.S., Johnston, E.R., Joseph, S., Jurburg, S.D., Jurelevicius, D., Karlsson,
A., Karlsson, R., Kauppinen, S., Kellogg, C.T.E., Kennedy, S.J., Kerkhof, L.J., King,
G.M., Kling, G.W., Koehler, A.V., Krezalek, M., Kueneman, J., Lamendella, R., Landon,
E.M., Lane-deGraaf, K., LaRoche, J., Larsen, P., Laverock, B., Lax, S., Lentino, M.,
Levin, I.I., Liancourt, P., Liang, W., Linz, A.M., Lipson, D.A., Liu, Y., Lladser, M.E.,
Lozada, M., Spirito, C.M., MacCormack, W.P., MacRae-Crerar, A., Magris, M., Martín-
Platero, A.M., Martín-Vivaldi, M., Martínez, L.M., Martínez-Bueno, M., Marzinelli,
E.M., Mason, O.U., Mayer, G.D., McDevitt-Irwin, J.M., McDonald, J.E., McGuire, K.L.,
McMahon, K.D., McMinds, R., Medina, M., Mendelson, J.R., Metcalf, J.L., Meyer, F.,
Michelangeli, F., Miller, K., Mills, D.A., Minich, J., Mocali, S., Moitinho-Silva, L.,
Moore, A., Morgan-Kiss, R.M., Munroe, P., Myrold, D., Neufeld, J.D., Ni, Y., Nicol,
G.W., Nielsen, S., Nissimov, J.I., Niu, K., Nolan, M.J., Noyce, K., O'Brien, S.L.,
Okamoto, N., Orlando, L., Castellano, Y.O., Osuolale, O., Oswald, W., Parnell, J.,
Peralta-Sánchez, J.M., Petraitis, P., Pfister, C., Pilon-Smits, E., Piombino, P., Pointing,
S.B., Pollock, F.J., Potter, C., Prithiviraj, B., Quince, C., Rani, A., Ranjan, R., Rao, S.,
Rees, A.P., Richardson, M., Riebesell, U., Robinson, C., Rockne, K.J., Rodriguezl, S.M.,
Rohwer, F., Roundstone, W., Safran, R.J., Sangwan, N., Sanz, V., Schrenk, M.,
Schrenzel, M.D., Scott, N.M., Seger, R.L., Seguin-Orlando, A., Seldin, L., Seyler, L.M.,
Shakhsheer, B., Sheets, G.M., Shen, C., Shi, Y., Shin, H., Shogan, B.D., Shutler, D.,
Siegel, J., Simmons, S., Sjöling, S., Smith, D.P., Soler, J.J., Sperling, M., Steinberg, P.D.,
Stephens, B., Stevens, M.A., Taghavi, S., Tai, V., Tait, K., Tan, C.L., Tas, N., Taylor,
D.L., Thomas, T., Timling, I., Turner, B.L., Urich, T., Ursell, L.K., van der Lelie, D.,
Van Treuren, W., van Zwieten, L., Vargas-Robles, D., Thurber, R.V., Vitaglione, P.,
Walker, D.A., Walters, W.A., Wang, S., Wang, T., Weaver, T., Webster, N.S., Wehrle,
B., Weisenhorn, P., Weiss, S., Werner, J.J., West, K., Whitehead, A., Whitehead, S.R.,
Whittingham, L.A., Willerslev, E., Williams, A.E., Wood, S.A., Woodhams, D.C., Yang,
Y. and The Earth Microbiome Project, C., *A communal catalogue reveals Earth's*
multiscale microbial diversity. Nature, 2017. **551**(7681): p. 457-463.
39.   Angel, R., Claus, P., and Conrad, R., *Methanogenic archaea are globally ubiquitous in*
aerated soils and become active under wet anoxic conditions. The ISME Journal, 2012.
6(4): p. 847-862.

40.   Kim, S., Gupta, N., and Pevzner, P.A., *Spectral Probabilities and Generating Functions*
of Tandem Mass Spectra: A Strike against Decoy Databases. Journal of Proteome
Research, 2008. **7**(8): p. 3354-3363.
41.   Monroe, M.E., Shaw, J.L., Daly, D.S., Adkins, J.N., and Smith, R.D., *Brief*
Communication: MASIC: A software program for fast quantitation and flexible
visualization of chromatographic profiles from detected LC-MS(/MS) features. Comput.
Biol. Chem., 2008. **32**(3): p. 215–217.
42.   Polpitiya, A.D., Qian, W.J., Jaitly, N., Petyuk, V.A., Adkins, J.N., Camp, D.G., 2nd,
Anderson, G.A., and Smith, R.D., *DAnTE: a statistical tool for quantitative analysis of -*
omics data. Bioinformatics, 2008. **24**(13): p. 1556-8.
43.   Farina, M., Christian, W., Hasson, N., McDermott, T., Powell, S., Hatzenpichler, R.,
Webb, H., LaRue, G., Okano, K., Sproles, E.A., and Watts, J.D., *Methane emission*
hotspots in a boreal forest-fen mosaic potentially linked to deep taliks. Environmental
Research Letters, 2025. **20**(10): p. 104029.
44.   Alowaifeer, A.M., Wang, Q., Bothner, B., Sibert, R.J., Joye, S.B., and McDermott, T.R.,
Aerobic methane synthesis and dynamics in a river water environment. Limnology and
Oceanography, 2023. **68**(8): p. 1762-1774.
45.   Jain, C., Rodriguez-R, L.M., Phillippy, A.M., Konstantinidis, K.T., and Aluru, S., *High*
throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries.
Nature Communications, 2018. **9**(1): p. 5114.
46.   Price, M.N., Dehal, P.S., and Arkin, A.P., *FastTree 2 – Approximately Maximum-*
Likelihood Trees for Large Alignments. PLoS ONE, 2010. **5**(3): p. e9490.
47.   Post, F.J., *The microbial ecology of the Great Salt Lake.* Microb Ecol, 1977. **3**(2): p. 143-
65.
48.   Shih, C.J. and Lai, M.C., *Differentially expressed genes after hyper- and hypo-salt stress*
in the halophilic archaeon Methanohalophilus portucalensis. Can J Microbiol, 2010.
56(4): p. 295-307.
49.   Keltjens, J.T. and Vogels, G.D., *Conversion of Methanol and Methylamines to Methane*
and Carbon Dioxide, in *Methanogenesis: Ecology, Physiology, Biochemistry & Genetics*,
J.G. Ferry, Editor. 1993, Springer US: Boston, MA. p. 253-303.
50.   Hippe, H., Caspari, D., Fiebig, K., and Gottschalk, G., *Utilization of trimethylamine and*
other N-methyl compounds for growth and methane formation by Methanosarcina
barkeri. Proceedings of the National Academy of Sciences, 1979. **76**(1): p. 494-498.
51.   Daly, R.A., Borton, M.A., Wilkins, M.J., Hoyt, D.W., Kountz, D.J., Wolfe, R.A., Welch,
S.A., Marcus, D.N., Trexler, R.V., MacRae, J.D., Krzycki, J.A., Cole, D.R., Mouser, P.J.,
and Wrighton, K.C., *Microbial metabolisms in a 2.5-km-deep ecosystem created by*
hydraulic fracturing in shales. Nature Microbiology, 2016. **1**(10): p. 16146.
52.   Wu, W.-L., Lai, S.-J., Yang, J.-T., Chern, J., Liang, S.-Y., Chou, C.-C., Kuo, C.-H., Lai,
M.-C., and Wu, S.-H., *Phosphoproteomic analysis of Methanohalophilus portucalensis*

755   *FDF1T identified the role of protein phosphorylation in methanogenesis and*
756   *osmoregulation.* Scientific Reports, 2016. **6**(1): p. 29013.

53.   Sepúlveda-Correa, A., Monsalve, L., Polania, J., Mestanza, O., and Vanegas, J., *Effect of*
salinity on genes involved in the stress response in mangrove soils. Antonie van
Leeuwenhoek, 2023. **116**(11): p. 1171-1184.

54.   Wang, L., Jiang, S., Deng, Z., Dedon, P.C., and Chen, S., *DNA phosphorothioate*
modification—a new multi-functional epigenetic system in bacteria. FEMS Microbiology
Reviews, 2019. **43**(2): p. 109-122.

55.   Bechtold, E.K., Ellenbogen, J.B., Xin, D., Pacheco, M., Toner, B.M., Chin, Y.-P.,
Arnold, W.A., Bansal, S., and Wilkins, M.J., *Sulfide stress tolerance as a controller of*
methane production in temperate wetlands. The ISME Journal, 2025. **19**(1): p. wraf196.

56.   Khanal, S.K. and Huang, J.C., *Effect of high influent sulfate on anaerobic wastewater*
treatment. Water Environ Res, 2005. **77**(7): p. 3037-46.

57.   Gupta, A., Flora, J.R.V., Sayles, G.D., and Suidan, M.T., *Methanogenesis and sulfate*
reduction in chemostats—II. Model development and verification. Water Research, 1994.
28(4): p. 795-803.

58.   Hoffman, M., Rajapakse, A., Shen, X., and Gates, K.S., *Generation of DNA-Damaging*
Reactive Oxygen Species via the Autoxidation of Hydrogen Sulfide under Physiologically
Relevant Conditions: Chemistry Relevant to Both the Genotoxic and Cell Signaling
Properties of H2S. Chemical Research in Toxicology, 2012. **25**(8): p. 1609-1615.

59.   Martens, J.H., Barg, H., Warren, M.J., and Jahn, D., *Microbial production of vitamin*
B12. Appl Microbiol Biotechnol, 2002. **58**(3): p. 275-85.

60.   Krumholz, L., Elias, D., and Suflita, J., *Immobilization of cobalt by sulfate-reducing*
bacteria in subsurface sediments. Geomicrobiology Journal, 2003. **20**(1): p. 61-72.

61.   Ekstrom, E.B. and Morel, F.M.M., *Cobalt Limitation of Growth and Mercury*
Methylation in Sulfate-Reducing Bacteria. Environmental Science & Technology, 2008.
42(1): p. 93-99.

62.   Gavel, O.Y., Bursakov, S.A., Calvete, J.J., George, G.N., Moura, J.J.G., and Moura, I.,
ATP Sulfurylases from Sulfate-Reducing Bacteria of the Genus Desulfovibrio. A Novel
Metalloprotein Containing Cobalt and Zinc. Biochemistry, 1998. **37**(46): p. 16225-
16232.

63.   Mahapatra, A., Patel, A., Soares, J.A., Larue, R.C., Zhang, J.K., Metcalf, W.W., and
Krzycki, J.A., *Characterization of a Methanosarcina acetivorans mutant unable to*
translate UAG as pyrrolysine. Mol Microbiol, 2006. **59**(1): p. 56-66.

64.   Hao, B., Gong, W., Ferguson, T.K., James, C.M., Krzycki, J.A., and Chan, M.K., *A new*
UAG-encoded residue in the structure of a methanogen methyltransferase. Science, 2002.
296(5572): p. 1462-6.

65.   O'Donoghue, P., Prat, L., Kucklick, M., Schäfer, J.G., Riedel, K., Rinehart, J., Söll, D.,
and Heinemann, I.U., *Reducing the genetic code induces massive rearrangement of the*

proteome. Proceedings of the National Academy of Sciences, 2014. **111**(48): p. 17206-
17211.

66.   Zhang, Y., Baranov, P.V., Atkins, J.F., and Gladyshev, V.N., *Pyrrolysine and*
selenocysteine use dissimilar decoding strategies. J Biol Chem, 2005. **280**(21): p. 20740-
51.

67.   Théobald-Dietrich, A., Giegé, R., and Rudinger-Thirion, J., *Evidence for the existence in*
mRNAs of a hairpin element responsible for ribosome dependent pyrrolysine insertion
into proteins. Biochimie, 2005. **87**(9-10): p. 813-7.

68.   Kivenson, V., Peters, S.L., Borrel, G., Kivenson, A., Roe, L.T., Hamlish, N.X.,
Fadhlaoui, K., Schepartz, A., Gribaldo, S., Hettich, R.L., and Banfield, J.F., *Sporadic*
distribution of a new archaeal genetic code with all TAG codons as pyrrolysine. bioRxiv,
2024: p. 2024.09.30.615893.

69.   Shalvarjian, K.E., Chadwick, G.L., Pérez, P.I., Woods, P.H., Orphan, V.J., and Nayak,
D.D., *The Ambiguous Genetic Code of Methanogenic Archaea that Grow on*
Methylamines. bioRxiv, 2025: p. 2025.06.11.659114.

70.   Walter Anthony, K.M., Anthony, P., Hasson, N., Edgar, C., Sivan, O., Eliani-Russak, E.,
Bergman, O., Minsley, B.J., James, S.R., Pastick, N.J., Kholodov, A., Zimov, S.,
Euskirchen, E., Bret-Harte, M.S., Grosse, G., Langer, M., and Nitzbon, J., *Upland*
Yedoma taliks are an unpredicted source of atmospheric methane. Nature
Communications, 2024. **15**(1): p. 6056.

71.   Cui, S., Liu, P., Guo, H., Nielsen, C.K., Pullens, J.W.M., Chen, Q., Pugliese, L., and Wu,
S., *Wetland hydrological dynamics and methane emissions.* Communications Earth &
Environment, 2024. **5**(1): p. 470.

72.   Nobu, M.K., Narihiro, T., Kuroda, K., Mei, R., and Liu, W.-T., *Chasing the elusive*
Euryarchaeota class WSA2: genomes reveal a uniquely fastidious methyl-reducing
methanogen. The ISME Journal, 2016. **10**(10): p. 2478-2487.

820
821

Figure legends

Figure 1: *Ca.* Methanohalophilus hillemani represents a novel species within the genus
Methanohalophilus. (A) Pairwise average nucleotide identity (ANI) and average amino acid
identity (AAI) on bottom left and top right, respectively, demonstrating *Ca.* M. hillemani
represents a new species within the *Methanohalophilus* genus. (B) Maximum likelihood tree of
65 aligned and concatenated single-copy *Methanohalophilus* proteins constructed with FastTree
v2.1.11 [46] and 1,000 bootstraps. Circles represent bootstrap values. Nodes with <50 support
were collapsed. Accession numbers for the other *Methanohalophilus* species can be found in SI
Table 1.

Figure 2: *Ca.* Methanohalophilus hillemani produces similar amounts of methane under a
variety of conditions. (A) Methane production of the *Methanohalophilus* enrichment culture
grown under various stress conditions. Error bars (where visible) represent the standard deviation
of triplicate cultures. (B) Community composition of same samples that were processed for
metaproteomics, as assessed by 16S rRNA gene amplicon sequencing. Amplicon sequence
variants that constituted less than 0.5% of the total community were clustered into the "Other"
category.
839
Figure 3: *Ca.* Methanohalophilus hillemani differential protein expression between control
and test conditions. Maximum differentially expressed proteins occurred between the control
and N_2 conditions, all of which were more highly expressed in the control. Down-regulated
proteins in the N_2 condition included proteins involved in immune response, metal metabolism,
and the cell-surface. Proteins shown in this figure meet three criteria: (i) above median
normalized protein expression for that condition; (ii) statistically significantly differentially
expressed ($p < 0.05$); and (iii) function could be predicted (*i.e.* hypotheticals not included). A
negative log fold change indicates the protein was more abundant in the control, positive
indicates more abundant in the test condition. Asterisks in the boxes indicate that the protein was
not detected in either that test condition or the control and a log2 fold change of 10 or -10 was
imputed. Bolded proteins indicate that the coding sequence for this protein was unique to *Ca.* M.
hillemani and no homologs were identified in other *Methanohalophilus* genomes. Proteins with
grey boxes do not indicate absence in a specific condition, but that the protein did not meet all
three criteria listed above. See Supplementary Tables 3-7 for raw values.
854
Figure 4: Methane flux and methanogenic community from Bridger Bay on the Great Salt
Lake. (A) Methane fluxes measured from soil within Transect A at Bridger Bay in May and
July of 2025 showing a zone of high methane flux 10-15 m from the shoreline. (B) Relative
abundance of different methanogens in a sediment core taken from the Transect A 10 m flux site
from July 15, 2025 based on 16S rRNA gene amplicon sequencing. For most taxa, taxonomic
classification was done at the genus level. *Ca.* M. hillemani was manually categorized based on a
one-nucleotide difference from other *Methanohalophilus* 16S rRNA genes in their amplified
region. Sequences that could not be assigned a genus are labeled "Family" or "Order" prior to
the name. "Uncultured Methanogens" (possibly members of the order *Methanofastidiosales*, see
main text) were the most abundant sequences. Only methanogen sequences are shown. (C) *mcrA*
gene amplicon relative abundance in the same core samples as (B), corroborating the presence of
Ca. M. hillemani. Other *Methanohalophilus mcrA* gene sequences were also present, including
some that could not be classified to the species level. For most taxa, taxonomic classification was
done at the species level. For taxa that could not be assigned to species, "Family" or "Genus" is
included in the name.
870

A

B 16S rRNA Gene Relative Abundance – July 15th, 2025 Transect A 10m Core

Genus
- Family Methanobacteriaceae
- Methanoregula
- Methanosphaera
- Methanonatronarchaeum
- Candidatus_Methanomethylicus
- Family Methanomassiliicoccaceae
- Methanohalophilus
- Methanohalobium
- Ca. Methanohalophilus hillemani
- Uncultured Methanogen

C *mcrA* Gene Relative Abundance – July 15th, 2025 Transect A 10m Core

Species
- Methanohalophilus profundi
- Methanosaeta concilii GP-6
- Family Methanomicrobiaceae
- Genus Methanolobus
- Family Methanosarcinaceae
- Ca. Methanohalophilus hillemani
- Methanohalophilus portucalensis
- Uncultured Archaeal Clone
- Methanohalobium evestigatum Z-7303
- Methanohalophilus mahii
- Genus Methanohalophilus
- Deep Sea Brine Archaea

Supplementary figures

Proteomic stress response by a novel methanogen enriched from the Great Salt Lake

William C Christian[1,2,3], Zackary J Jay[1,2,3], Nikola Tolic[4], Carrie D Nicora[5], Reece Livingstone[1], Stavros Trimmer[1,2,3], Timothy R McDermott[6], Roland Hatzenpichler[1,2,3,7,*]

Supplementary Figure 1

Microbialite material from which the enrichment culture was started. Material shown in the spoon was the exact material from which *Ca*. Methanohalophilus hillemani was enriched.

Supplementary Figure 2: Methane flux transects from July 15th 2025

Locations of the transects A, B and C as sampled on July 15th 2025. The pinned location represents the shoreline (1m) flux measurement. Measurements extended further into the shore and out into the water, perpendicularly, from this point.

17

Supplementary Figure 3: Change in the Great Salt Lake during past half century

Google Earth satellite imagery from the Great Salt Lake near its recorded high lake level in 1986 (A) and near its recorded low in 2022 (B). (C) Lake level and the reciprocal salinity varied substantially at the times when *M. mahii* and *Ca.* M. hillemani were cultured. Lake level data was retrieved for Saltair Boat Harbor from the USGS website. "USGS Salinity" data was only available on the website starting in August of 1995. Salinity data prior to this was retrieved from Arnow and Stephens 1990. Salinity data from Bridger Bay was collected in this study and measured using a refractometer.

Supplementary Figure 4: Methane production from various substrates

The GSL methanogenic enrichment culture produced methane from MMA, DMA, TMA, and methanol, but not from glycine betaine, or H_2 and CO_2.

16S rRNA Gene % Identity Heatmap

99.93																
99.93	100															
99.19	99.12	99.12														
98.63	98.63	98.63	99.73													
99.12	99.05	99.05	99.46	99.45												
99.25	99.25	99.25	99.8	99.86	99.53											
99.25	99.25	99.25	99.8	99.86	99.53	100										
99.12	99.05	99.05	99.46	99.45	100	99.53	99.53									
99.12	99.12	99.12	99.66	99.59	99.53	99.73	99.73	99.53								
99.12	99.12	99.12	99.66	99.59	99.53	99.73	99.73	99.53	100							
99.12	99.12	99.12	99.66	99.59	99.53	99.73	99.73	99.53	99.86	99.86						
99.19	99.12	99.12	99.46	99.73	99.59	99.53	99.53	99.59	99.39	99.39	99.39					
99.19	99.19	99.19	99.73	99.73	99.46	99.8	99.8	99.46	99.8	99.8	99.8	99.46				
99.19	99.19	99.19	99.73	99.73	99.46	99.8	99.8	99.46	99.8	99.8	99.8	99.46	100			
99.19	99.19	99.19	99.73	99.73	99.46	99.8	99.8	99.46	99.8	99.8	99.8	99.46	100	100		
99.19	99.19	99.19	99.73	99.73	99.46	99.8	99.8	99.46	99.8	99.8	99.8	99.46	100	100	100	
95.74	95.74	95.74	96.21	96.17	96.01	96.28	96.28	96.01	96.28	96.28	96.34	95.88	96.21	96.21	96.21	96.21

Row labels (top to bottom):
Ca. M. hillemani 01
Ca. M. hillemani 02
Ca. M. hillemani 03
M. profundi 01
M. profundi 02
M. sp. WG1 01
M. portucalensis 01
M. portucalensis 02
M. euhalobius 01
M. mahii 01
M. mahii 02
M. mahii_03
M. sp. RSK 01
M. halophilus 01
M. halophilus 02
M. halophilus 03
M. halophilus 04
M. levihalophilus 01

Column labels (left to right):
Ca. M. hillemani 01
Ca. M. hillemani 02
Ca. M. hillemani 03
M. profundi 01
M. profundi 02
M. sp. WG1 01
M. portucalensis 01
M. portucalensis 02
M. euhalobius 01
M. mahii 01
M. mahii 02
M. mahii_03
M. sp. RSK 01
M. halophilus 01
M. halophilus 02
M. halophilus 03
M. halophilus 04
M. levihalophilus 01

Legend: 100, 99, 98, 97, 96

Supplementary Figure 5: Heatmap of 16S rRNA gene similarity

Heatmap showing the percent identity of the 16S rRNA genes in the *Methanohalophilus* genus. All copies from each genus are shown, should they have multiple copies.

Supplementary Figure 6: Expression of proteins involved in methyl-dismutating methanogenesis

For each condition the normalized protein values were compared to others within that condition to assess the percentile of expression. In the case of duplicate proteins, the most highly expressed copy was used for this figure. Abbreviations: MttB, Trimethylamine Methyltransferase; MttC, Trimethylamine Corrinoid Protein; MtbB, Dimethylamine Methyltransferase; MtbC, Dimethylamine Corrinoid Protein; MtmB, Monomethylamine Methyltransferase; MtmC, Monomethylamine Corrinoid Protein; MtbA, Methylcobamide:CoM methyltransferase; McrA, Methyl Coenzyme M Reductase Subunit α; MtrA, Tetrahydromethanopterin S-methyltransferase subunit A; Mer, 5,10-methylenetetrahydromethanopterin reductase; Mtd, F420-dependent methylenetetrahydromethanopterin dehydrogenase; Mch, Methenyltetrahydromethanopterin cyclohydrolase; Ftr, Formylmethanofuran--tetrahydromethanopterin formyltransferase; FwdA, formylmethanofuran dehydrogenase subunit A.

Supplementary Figure 7: Methane flux across all dates and all transects from Bridger Bay in the GSL

The time points were selected to represent the lowest seasonal lake level (Sept. 15, 2024), the highest seasonal lake level (May 21, 2025), and the time of year which *Ca.* Methanohalophilus hillemani was collected from the lake (July 15, 2025) (see Supplementary Figure 10).

Methane Flux – Transect A 10m Collar Jul. 15 2025

Supplementary Figure 8: Methane flux dynamics from a single location

Terrestrial flux measurements made over the course of 6 hours from the exact same location, highlighting the variability of methane flux. This was the exact location from which the sediment core was collected that was used to generate the amplicon data presented in main text Figure 4. This was the most extreme example of flux variability observed at a single location. The atmospheric pressure was dropping that afternoon and the wind was increasing, so it is possible that a low-pressure system evacuated gas from the sediment, resulting in high methane fluxes. However, the pressure continued dropping, as did methane flux, from 11:56 to 13:45. Unless all the methane that had been "stored up" in the sediment was evacuated, meteorology cannot entirely explain the extreme variability in methane flux.

Supplementary Figure 9: *Desulfovermiculus* **is the most abundant SRB present in GSL soils**

Relative abundance of the different SRB genera from a sediment core taken from the Transect A 10m flux site from July 15, 2025 based on 16s rRNA gene amplicon sequencing. Amplicons plotted belong to the taxonomic Classes *Desulfovibrionia*, *Desulfobacteria*, *Desulfobulbia*, *Desulfuromonadia*, *Desulfotomaculia*, and *Desulfitobacteriiain*, and excluded orders shown in the literature not to perform sulfate reduction. Amplicon sequence variants that constituted less than 0.1% of the total community were clustered into the "Other" category. Sequence variants that could not be assigned a genus are labeled "Family" prior to the name.

Supplementary Figure 10: Methane flux sampling dates compared to historical lake levels

Red triangles indicate the date that methane flux measurements were taken from the GSL. Lake level data was retrieved for Saltair Boat Harbor from the USGS website.

Supplementary Figure 11: Phylogenetic tree of *Methanohalophilus* McrA sequences

Maximum likelihood tree of 573 amino acid positions constructed with IQtree2 using 1,000 ultrafast bootstrap replicates.

1 **Supplementary text**
2
3 **Proteomic stress response by a novel methanogen enriched from the Great Salt Lake**
4
5 William C Christian[1,2,3], Zackary J Jay[1,2,3], Nikola Tolic[4], Carrie D Nicora[5], Reece Livingstone[1],
6 Stavros Trimmer[1,2,3], Timothy R McDermott[6], Roland Hatzenpichler[1,2,3,7,*]

8 <u>**Supplementary methods**</u>
9 **Enrichment**
10  Microbialite material was incubated in gas-sealed bottles with filter-sterilized GSL water
collected on July 12, 2022 (salinity of 23.7%), and spiked with 10 mM MMA to enrich for
methanogens [1]. From July 2022 to December 2022, enrichments were maintained by
transferring inoculum, in a 1:10 ratio, into fresh lake water + MMA. When the GSL water
collected in July 2022 was depleted, enrichments then used filter-sterilized GSL water collected
on April 13th 2023 (salinity of 16%) as the liquid media until October 2023. Lake water was then
replaced by media designed by Paterek and Smith [2], originally used to isolate
Methanohalophilus mahii, but adjusted to salinity of GSL water from April 2023. In February
2024, TMA replaced MMA as the methanogenic substrate, as it yielded more methane
production. In December of 2024, we stopped adding sodium sulfide and the volatile fatty acid
solution from the Paterek and Smith 1985 [2] recipe – no impact was observed on methane
production. Enrichment cultures have been maintained in this manner since December 2024.
More details on exact conditions and media preparation can be found in the 'Culturing
Conditions' section.
24

25 **DNA Extractions and 16S rRNA gene amplicon sequencing**
26  In April of 2024, after ~660 days of incubation and 23 transfers, phenol-chloroform-
isoamyl alcohol DNA extractions were done on the methanogenic enrichments being grown in
the Paterek and Smith media [2]. 1.5 mL of culture was harvested and pelleted at 20,000 x G for
10 minutes. The supernatant was discarded and the pellet was resuspended in 100 µL of 20 mM
Tris-HCl (pH 8). 2 µL of 1% SDS was added to the tubes, which were then vortexed for 2
minutes. 100 µL of 25:24:1 phenol:chloroform:isoamyl alcohol was added to the tubes which
were scraped against the tube rack to mix. The tubes were then centrifuged again, as before, and
the top layer (~100 µL) was pipetted into fresh tubes containing 75 µL of ice-cold isopropanol.
These tubes were then vortexed, and spun at 20,000 x G for 5 minutes, the supernatant was
decanted, and 500 µL of 75% ethanol was added to the (frequently invisible) pellet. The tubes
were then stored at -20 ℃ overnight. The next day, the tubes were spun at 20,000 x G for 5
minutes and the supernatant was decanted. Tubes were then dried in a 37 ℃ incubator until all
ethanol had evaporated off (~20 minutes) and the (frequently invisible) pellet was resuspended in
32 µL of 20 mM Tris-HCl (pH 8).

16S rRNA gene amplification, barcoding, sequencing, and subsequent bioinformatics
data analysis was performed as described in Krukenberg et al. [3]. The V4 hypervariable region
of the 16S rRNA gene was amplified using the 515F and 806R Earth Microbiome Project
primers, allowing approximately genus-level resolution of most previously characterized
bacterial and archaeal taxa [4, 5]. The amplicon sequencing was performed via Illumina MiSeq
by SeqCenter (Pittsburgh, PA). QIIME2 [6] was used to process the data and DADA2 was used
to merge forward and reverse reads [7].
47
Metagenome sequencing
Metagenome libraries were constructed from the remaining DNA of a 16S rRNA
amplicon sample that contained 63% relative abundance of *Methanohalophilus*. Long-read
sequencing libraries were prepared using the rapid barcoding kit (SQK-RBK114-24; Oxford
Nanopore Technologies) according to the manufacturer's protocol. Sequencing was performed
on a MinION device equipped with an R10.4.1 flow cell (Oxford Nanopore Technologies).
Basecalling was conducted with Dorado v0.7.2 (Oxford Nanopore Technologies). Short-read
sequencing libraries were generated on an Illumina NovaSeq X Plus platform (SeqCenter,
Pittsburgh, PA, USA) generating 2 x 151 bp paired-end reads.
57
Metagenome assembly and annotation
Nanopore reads were assembled with MetaFlye 2.9.4-b1799 [8]. Manual evaluation of
the resulting 85 contigs (≥2 kb) with SendSketch (BBTools Suite v38.94; [9]) and BLASTn
(v2.14.0+) identified 10 *Methanohalophilus*-like contigs (2.21 Mb), including a 1.78 Mb
fragment. These contigs were used as references for mapping Illumina short-reads with BBsplit
(default parameters) and Nanopore reads with Minimap2 v2.26-r1175 [10] using default
parameters prior to hybrid assembly with Unicycler v0.5.0 (--mode bold) [11]. The curated
assembly produced 4 scaffolds (2.08 Mb), which were subsequently used for a second round of
Illumina and Nanopore read mapping and assembly with Unicycler. Curation of this second
assembly resulted in 2 *Methanohalophilus*-like contigs (2.10 Mb), which were further closed
with LongStitch v1.0.5 [12] and Nanopore read mapping.
The closed genome was polished with long-reads using four rounds of Racon v1.5.0 [13]
followed by one round of Medaka v2.1.0 (Oxford Nanopore Technologies) with the
r104_e82_400bps_hac_v5.0.0 model. Short-read polishing was performed with two rounds of
Polypolish v0.6.1 [14] and one round of POLCA v4.1.1 [15], both utilizing BWA v0.7.17-r1188
[16] alignments. The final circular genome was 2,093,478 bp in length with a GC content of 41.5
%. Genome completeness (99.99 %) was estimated using CheckM2 v1.0.1 [17].
Illumina reads were also assembled with MetaSpades v3.15.5 [18]. Coverage was
calculated with BBmap (ambiguous=random), and sequences ≥2 kb were binned using MetaBAT
v2.12.1 (with and without coverage; [19]), MaxBin v2.2.7 [20], CONCOCT v1.0.0 [21],
Autometa v1 (archaeal and bacterial modes; including the machine learning option; [22]), and
MetaDecoder v1.0.19 [23]. Final bin evaluation was performed with DAS Tool v1.1.7 [24],

resulting in fragmented but complete (>99.5 %) genomes of *Halanaerobium* and
Desulfovermiculus spp. at ~17X and ~25X lower coverage, respectively, compared to
Methanohalophilus (Supplementary Table 11). Genomes were annotated with Prokka v1.14.6
[25] using both default and custom in-house annotation databases. These genomes can be found
with the following accession numbers for *Ca.* M. hillemani str. BB.*, Desulfovermiculus* sp.*,* and
Halanaerobium sp.*,* respectively: SAMN54119603, SAMN54119604, SAMN54119605
86

Marker gene analyses and comparative genomics

Publicly available *Methanohalophilus* spp. 16S rRNA sequences were aligned and
masked using SSU-ALIGN v0.1.1 [26], resulting in a final alignment of 1,376 positions.
Maximum likelihood analysis was performed with FastTree v2.1.11 [27] using default
parameters. A single nucleotide polymorphism in one of the three *Ca.* M. hillemani 16S rRNA
gene copies did not affect phylogenetic placement. Average 16S rRNA gene sequence identities
were determined with pairwise BLASTn comparisons.

Sequence alignments of *Methanohalophilus* spp. McrA were generated with MAFFT-linsi
v7.522 [28] and trimmed with trimAL v1.4.rev22 [29] using a 50 % gap threshold, producing a
final alignment of 573 positions. Maximum likelihood phylogenetic analysis was conducted with
IQtree2 v2.0.6 [30] using the LB+C60+F+G model and 1,000 ultrafast bootstrap replicates
(Supplementary Figure 11).

A set of 65 single-copy marker proteins (Supplementary Table 12) were collected from
the genomes of isolated *Methanohalophilus* spp., aligned with MUSCLE v3.8.1551 [31], and
concatenated to generate a final alignment of 14,606 positions. Phylogenetic inference was
performed with FastTree (option -wag) using 1,000 ultrafast bootstrap replicates.

Whole-genome comparisons were conducted to calculate average nucleotide identity
(ANI) with fastANI (-fragLen 2,000; [32]) and average amino acid identity (AAI) with
CompareM v0.1.2 (default parameters; https://github.com/dparks1134/CompareM), respectively.
Additionally, Mauve v20150226 [33] was used to identify orthologous protein-coding sequences
(≥70 % amino acid ID and ≥70 % aligned length) (Supplementary Tables 13 - 15). Orthologs
were further evaluated with BLASTp alignments, and proteins were considered unique if they
exhibited <50 % amino acid identity or aligned with <25 % of any other *Methanohalophilus*
proteins (Supplementary Table 16 , 17).
111

Culturing conditions

Since December 2024, *Ca.* M. hillemani cultures were grown in Paterek and Smith 1985
media [2], with differing concentrations of salts and TMA, and without the addition of volatile
fatty acid solution and sodium sulfide. The headspace consisted of a 50/45/5 ratio of $H_2/N_2/CO_2$
at ambient pressure (~90kPa in Bozeman, MT). These conditions served as the control for the
methane production and proteomics experiments. Cultures were grown with 10 mL of liquid
medium in a 70 mL borosilicate glass serum bottle. The salt levels in our media were increased
to match GSL lake water collected on Apr. 13[th] 2023, as measured by the Joye Lab at The

University of Georgia. The media was made by adding the following salts: 100 g/L NaCl, 45 g/L
MgCl$_2$, 5.5 g/L KCl, 1.5 g/L CaCl$_2$, 0.13 g/L LiCl, 19 g/L Na$_2$SO$_4$, 0.5 g/L NH$_4$Cl. This results in
ion concentrations for Na$^+$, Cl$^-$, Ca^{2+}, Mg^{2+}, K$^+$, and SO$_4$$^{2-}$ of 2,040, 2,770, 14, 221, 74, and 134
mM, respectively. This salt solution was then autoclaved using a 30-minute sterilization liquid
cycle. After autoclaving, the media was immediately sparged for 10 minutes with 90/10 N$_2$/CO$_2$
gas to purge O$_2$. The media bottle was then sealed and left to cool. Once cooled, the following
components were added to make 1 L final volume of media: 500 µL of 1 g/L sodium resazurin
solution, 10 mL of DSMZ 141 trace element solution (Modified Wolin's mineral solution), 23.81
mL of 1 M sodium bicarbonate, 18.87 mL 1 M sodium carbonate, 5 mL of 40 g/L 2-
mercaptoethane sulfonic acid sodium salt (Coenzyme M), and a small spatula tip (1-2 mg) of
sodium dithionite as a reducing agent. Finally, pH was adjusted to ~7.5. When new bottles were
to be inoculated, 9.5 mL of media was added to the 70 mL serum bottles. Then, 10 µL of 50
mg/mL vancomycin and streptomycin were added to each bottle along with 200 µL of 1 M TMA
and 300 µL of carbon source mix. This mix consisted of 33.33 g/L of yeast extract, 33.33 g/L of
trypticase peptone, and 16.67 g/L of cysteine HCl [2]. Cultures were inoculated with 100 µL
from the previous culture and then grown to stationary at 30 ℃. Cultures were grown for
approximately a month before being transferred to fresh media.
137
Isolation efforts
Dilution to extinction and plating were attempted to isolate *Ca.* Methanohalophilus
hillemani. Solid media were prepared by adding 16 g/L agar to the medium before autoclaving.
Plates were poured and streaked from active liquid cultures in a Coy chamber (Coy Lab
Products, Grass Lake, MI, USA). The plates were then placed into a gas-sealed metal container
(Liquid Impact Water Jet & Fabrication Fort Collins, CO, USA) and incubated at 30 ℃. We tried
100% N$_2$ and 50/45/5 H$_2$/N$_2$/CO$_2$ headspaces in the gas-sealed container. Adding additional
antibiotics, chloramphenicol and kanamycin to solid and liquid media, was also attempted.
Dilution to extinction and plating efforts resulted in methane-producing cultures, but, based on
microscopy, neither resulted in an isolate.
148
Metaproteomics sample preparation
The harvested cultures were sent to the Pacific Northwest National Laboratory (Richland,
WA, USA) where protein digestion was performed using S-Trap microfilters (Protifi, Fairport
NY) following an optimized procedure adapted from the manufacturer's guidelines with reagents
purchased from Sigma (St. Louis, MO, USA). Cell pellets were resuspended in 50 µL of 5 %
(w/v) sodium dodecyl sulfate (SDS) prepared in 50 mM triethylammonium bicarbonate (TEAB;
pH 7.5, LC-MS–grade water) and lysed by alternating brief sonication and vortex agitation until
complete solubilization was achieved. Total protein concentration was quantified by
bicinchoninic acid (BCA) assay (Thermo Scientific, Waltham, MA, USA), and 200 µg of protein
from each sample was used for digestion. Disulfide bonds were reduced by addition of
dithiothreitol (DTT) to a final concentration of 20 mM, followed by incubation at 95 °C for 10

min. After cooling to ambient temperature, cysteine residues were alkylated with 40 mM
iodoacetamide in the dark for 30 min.
Samples were acidified with phosphoric acid to a final concentration of 1.2% (v/v) and diluted
six-fold with S-Trap binding buffer composed of 90% methanol and 100 mM TEAB (pH 7.1).
The mixtures were gently vortexed and applied to S-Trap filter cartridges. Proteins were captured
on the filter by centrifugation at approximately 2,000 × g for 30–60 s, and the flow-through was
reapplied twice to ensure complete binding. The filters were then washed three times with 200
µL of binding buffer to remove residual detergents and contaminants.
168   Bound proteins were digested on-filter with sequencing-grade trypsin (Promega,
Madison, WI) and Lys-C (Fujifilm Wako, Richmond, VA) in 150 µL of 50 mM TEAB, pH 8.0
(digestion buffer) at 37 °C for 16 h. To prevent membrane desiccation during the overnight
incubation, a digestion volume of at least 100 µL was maintained. Peptides were sequentially
eluted using three stepwise buffers (each 80 µL): 50 mM TEAB, 0.2% formic acid in water, and
50% acetonitrile containing 0.2% formic acid. The combined eluates were dried by vacuum
centrifugation and desalted using C18 macro-spin columns (Harvard Apparatus, Holliston, MA,
USA). Cleaned peptide samples were subsequently dried and stored at –80 °C until LC-MS/MS
analysis.
177   Raw proteomics data has been deposited at massIVE under accession number:
178   MSV000100495
179
180   **Metaproteomic MS measurements and data analysis**
181   MS analysis was performed using an Orbitrap Exploris 480 mass spectrometer (Thermo
Scientific, Waltham, MA, USA) outfitted with a homemade nano electrospray ionization
interface using data-dependent acquisition (DDA) mode (QC plots in Supplementary Table 18).
The ion transfer tube temperature and spray voltage were 300°C and 2.2 kV, respectively. Data
were collected for 120 min following a 10 min delay after completion of sample trapping and
start of gradient. FT MS spectra were acquired from 300 to 1800 m/z at a resolution of 60 K
(AGC target 3e6) and up to the top 20 FT HCD MS/MS spectra were acquired in DDA mode
with an isolation window of 0.7 m/z at a resolution of 30 K (AGC target 1e5) using a normalized
collision energy of 30, dynamic exclusion time of 45 s , and detected charge state of an ion 2 to
6.
191   Peptide matching from the acquired datasets was performed using MSGF+ software [34]
against proposed target meta-proteome composed from sequenced genomes (see 'Metagenome
Assembly and Annotation' section) of novel members of *Desulfovermiculus*, *Halanaerobium*,
and *Methanohalophilus* genera totaling 7,403 protein sequences. *Ca.* M. hillemani protein
collection incorporated Prokka [25] predicted protein sequences with non-standard amino acid
pyrrolysine (symbol O in FASTA file) and common contaminant sequences were added to the
protein collection used. MSGF+ was used in target/decoy mode with 20 ppm parent ion
tolerance, partial tryptic rule with static carbamidomethylation of cysteine (+57.0215) and
variable modifications for the oxidation of methionine (+15.9949), deamidation of (+0.984) of

asparagine and glutamine, and a methyl loss (-14.0157) for pyrrolysine (O) residues. Best
matches from the MSGF+ searches were filtered at 1% FDR based on target/decoy model; this
set of peptides was used in consequent quantitation analysis (Supplementary Table 19). Spectral
counts (SC) and normalized spectral abundance factors (NSAF) were derived from counts of 1%
false detection rate (FDR) detected peptides without consideration of peptide-protein specificity.
For MASIC reports, to optimize peptide-protein specificity and protein coverage, the best protein
for each peptide was guessed based on the full set of protein-peptide mappings.
Peptide abundances, integrated as area under the peptide elution peaks, were produced
from raw spectra using in-house developed software MASIC [35]. MASIC peptide abundances
were log2 transformed to remove skewness in the distribution of measured abundances
(Supplementary Table 20). Transformed abundance values were normalized using mean central
tendency method implemented in InfernoR software [36]. For each sample, *Ca.*
Methanohalophilus hillemani peptide abundances were normalized to the relative abundance of
Ca. M. hillemani by dividing the summed area under the curve (AUC) for *Ca.* M. hillemani
peptides by the summed AUC for all peptides. The AUC for each *Ca.* M. hillemani peptide in
each sample was then divided by this sample-specific relative abundance value (Supplementary
Table 21). Peptides were then aggregated to the protein level by summing all peptide AUCs
mapping to specific proteins (Supplementary Table 22). For protein level comparison, pairwise
z-score analysis (standard deviation from the mean) between samples was used to extract
significant differences in protein expression with significance attributed to a z-score of 2 or
higher (Supplementary Tables 3 - 7). Unless otherwise indicated, all proteomic data presented in
this study represents normalized protein abundances.
222
Methane flux data collection
Methane flux data was collected at Bridger Bay in the south arm of the GSL on three
different dates: Sept, 15, 2024; May 21, 2025, and July 15, 2025. This is the same location where
the original sample for cultivation was obtained. A Li-COR 7810 trace gas analyzer (Li-COR,
Lincoln, NE, USA) was used to measure methane flux. For terrestrial flux, we used a Li-COR
8200–01S Smart Chamber placed on cylindrical, PVC soil collars (10 cm diameter). Three
replicates were collected per sample. Each replicate consisted of a one-minute measurement with
a 10 second deadband time, followed by a one-minute period to flush prior to the next replicate.
The SoilFluxPro Software (Li-COR, Lincoln, NE, USA) was used to analyze the data, and the
exponential fit was chosen to estimate methane flux.
Aquatic flux measurements were done using a custom-made floating flux chamber to take
duplicate, one minute, flux measurements. The chamber consisted of a gas-tight coffee can fitted
with a foam floatation collar. The internal volume of the flux chamber was 2,584 mL. The
chamber was removed from the water's surface between duplicates to allow the chamber gas
composition to return to atmospheric conditions. Aquatic fluxes were calculated from the linear
change in methane concentration over time, as previously described [37].

On each sampling date, flux was collected in a transect, perpendicular to the GSL Bridger
Bay shoreline. Measurements extended toward to shore and out into the water, perpendicularly,
from the shore.
242

Soil core collection and *mcrA* gene analysis

Soil cores were collected from within the boundary of the terrestrial collar with the
highest methane flux (transect A - 15m for May 2025 and transect A – 10m for July 2025) and
transported back to Montana State University on ice. In the lab, the core was separated into
sections of three or six cm each. DNA from each soil core section was extracted with MP Bio
Soil Extraction kits, and 16S rRNA gene amplicon sequencing was performed, as described
above. For the July 15, 2025 core, the DNA extract was also used for amplifying and sequencing
an amplicon for the gene encoding the alpha-subunit of methyl-coenzyme M reductase (*mcrA*).
The mlas-mod-F and *mcrA*-rev-R primers and the touchdown amplification program from Angel
et al. 2012 were utilized [38]. Sequencing and analysis of data was performed as described
above. However, due to variable lengths of the *mcrA* gene amplicon, the merging of forward and
reverse reads via DADA2 was not uniformly successful. Therefore, the *mcrA* gene amplicon data
was analyzed using only the ~227 bases from the forward read.
256

<u>Supplementary results</u>

Unique genetic features of *Ca.* Methanohalophilus hillemani

We identified 64 coding sequences in *Ca.* M. hillemani that have no homologs in any of
the other 13 *Methanohalophilus* genomes (Supplementary Tables 16, 17). When analyzing these
unique (among *Methanohalophilus*) sequences, there were many proteins that appeared to be part
of potential operons. Three operons of unique *Ca.* M. hillemani genes involved in immune
response were found. One consists of three genes encoding type 1 restriction modification
system proteins. Type 1 restriction-modification systems are responsible for recognizing self-
DNA (hsdS), modifying self-DNA via methylation (hsdM), and destroying foreign DNA via
restriction endonuclease activity (hsdR) [39, 40]. *Ca.* M. hillemani encodes a second type 1
restriction-modification system operon, however, these proteins have significant homology to
sequences from other *Methanohalophilus* species. A second unique operon encodes a toxin
production protein, a CRISPR associated protein, and two antiviral radical SAM proteins. A third
consists of three antibiotic resistance ABC transports and a protease. Borton et al. in 2018
suggested that viral predation may drive strain differentiation in *Methanohalophilus* species [41].
Our data corroborates that immune response is a differentiating factor between *Ca.* M. hillemani
and other *Methanohalophilus* genomes.
A large operon of unique coding sequences for DNA maintenance and modification
proteins was also found in the *Ca.* M. hillemani genome. This putative operon includes genes
coding for a chromosome segregation protein, a DNA helicase, a DNA sulfur modification
protein, and a RecB family nuclease. In addition to this operon, *Ca.* M. hillemani encodes three
unique DNA helicases and a unique double strand break repair protein. DNA repair mechanisms

are shared across the genus, including the DNA repair proteins RadA, B, and C. However, the
unique *Ca.* M. hillemani sequences in this category indicate that DNA maintenance may be of
particular importance.
Cobalamin (vitamin B12) is a cobalt-containing corrinoid cofactor that is essential for
methyl-dismutating methanogenesis [42], and all *Methanohalophilus* species share most genes
required for its biosynthesis. While *Ca.* M. hillemani encodes cobalamin biosynthesis, its
genome also contains two operons with unique coding sequences involved in cobalamin import.
The first operon encodes the Cobalamin import permease, BtuC, the Cobalamin import ATP-
binding protein, BtuD, and an ABC transporter. The other operon contains a second copy of
BtuD and two ABC transporters. Other *Methanohalophilus* species encode genes annotated as
BtuC and BtuD; however, *Ca.* M. hillemani appears to have acquired/evolved divergent uptake
pathways, as its BtuC and D proteins have low ($<50\%$) amino acid identity with any proteins in
the other genomes.
292
Etymology
The newly cultured archaeon is a methanogen affiliated with the genus Methanohalophilus,
family Methanosarcinaceae, order Methanosarcinales, class Methanomicrobia, phylum
Methanobacteriota, for which we propose the name *Candidatus* Methanohalophilus hillemani sp.
nov. strain BB.
Me. tha. no. ha. lo. phi'lus. M. L. n. *methanum*, methane; Gr. n. *halo*, salt; Gr. no. *philos*,
lover; M. L. n. *Methanohalophilus*. methane-producing salt lover.
hil.le.ma'ni N.L. gen. masc. n. *hillemani,* of Hilleman; named in honor of Maurice
Hilleman (1919-2005), a microbiologist and vaccinologist born in Miles City, Montana.
Hilleman graduated from Montana State University and, over the course of his career, developed
over 40 vaccines including eight that are currently recommended for children and adults in the
United States. His vaccines significantly reduced morbidity and mortality during two 20[th]
century influenza pandemics, and his cumulative contributions are credited with saving 10s of
millions of lives worldwide. See https://hillemanfilm.com/dr-hilleman for more information.
Appropriately, the results of this study show that *Ca.* M. hillemani invests heavily in immune
response.
The strain name BB refers to Bridger Bay on the Great Salt Lake from which *Candidatus*
Methanohalophilus hillmani was cultured from.
311
Citations

1.   Wang, Q., *Aerobic Bacterial Methane Synthesis Encoded by a Single Gene*. 2021: Proc.
Natl. Acad. Sci. U. S. A.
2.   Paterek, J.R. and P.H. Smith, *Isolation and characterization of a halophilic methanogen*
from great salt lake. Appl Environ Microbiol, 1985. 50(4): p. 877-81.

3.   Krukenberg, V., et al., *Microbial Community Response to Polysaccharide Amendment in*
Anoxic Hydrothermal Sediments of the Guaymas Basin. Frontiers in Microbiology, 2021.
Volume 12 - 2021.

4.   Parada, A.E., D.M. Needham, and J.A. Fuhrman, *Every base matters: assessing small*
subunit rRNA primers for marine microbiomes with mock communities, time series and
global field samples. Environ Microbiol, 2016. 18(5): p. 1403-14.

5.   Thompson, L.R., et al., *A communal catalogue reveals Earth's multiscale microbial*
diversity. Nature, 2017. 551(7681): p. 457-463.

6.   Bolyen, E., et al., *Reproducible, interactive, scalable and extensible microbiome data*
science using QIIME 2. Nature Biotechnology, 2019. 37(8): p. 852-857.

7.   Callahan, B.J., et al., *DADA2: High-resolution sample inference from Illumina amplicon*
data. Nature Methods, 2016. 13(7): p. 581-583.

8.   Kolmogorov, M., et al., *metaFlye: scalable long-read metagenome assembly using repeat*
graphs. Nature Methods, 2020. 17(11): p. 1103-1110.

9.   Bushnell, B. *BBMap: A Fast, Accurate, Splice-Aware Aligner*. 2014.

10.   Li, H., *New strategies to improve minimap2 alignment accuracy.* Bioinformatics, 2021.
37(23): p. 4572-4574.

11.   Wick, R.R., et al., *Unicycler: Resolving bacterial genome assemblies from short and long*
sequencing reads. PLoS computational biology, 2017. 13(6): p. e1005595.

12.   Coombe, L., et al., *LongStitch: high-quality genome assembly correction and scaffolding*
using long reads. BMC Bioinformatics, 2021. 22(1): p. 534.

13.   Vaser, R., et al., *Fast and accurate de novo genome assembly from long uncorrected*
reads. Genome Research, 2017: p. gr.214270.116.

14.   Wick, R.R. and K.E. Holt, *Polypolish: Short-read polishing of long-read bacterial*
genome assemblies. PLoS Computational Biology, 2022. 18(1): p. e1009802.

15.   Zimin, A.V. and S.L. Salzberg, *The genome polishing tool POLCA makes fast and*
accurate corrections in genome assemblies. PLOS Computational Biology, 2020. 16(6):
p. e1007981.

16.   Li, H. and R. Durbin, *Fast and accurate short read alignment with Burrows-Wheeler*
transform. Bioinformatics, 2009. 25(14): p. 1754-60.

17.   Chklovski, A., et al., *CheckM2: a rapid, scalable and accurate tool for assessing*
microbial genome quality using machine learning. Nature Methods, 2023. 20(8): p. 1203-
1212.

18.   Nurk, S., et al., *metaSPAdes: a new versatile metagenomic assembler.* Genome Research,
2017. 27(5): p. 824-834.

19.   Kang, D.D., et al., *MetaBAT, an efficient tool for accurately reconstructing single*
genomes from complex microbial communities. PeerJ, 2015. 3: p. e1165-e1165.

20.   Wu, Y.-W., et al., *MaxBin: an automated binning method to recover individual genomes*
from metagenomes using an expectation-maximization algorithm. Microbiome, 2014.
2(1): p. 26.

21. Alneberg, J., et al., *Binning metagenomic contigs by coverage and composition.* Nature
Methods, 2014. 11(11): p. 1144-1146.
22. Miller, I.J., et al., *Autometa: automated extraction of microbial genomes from individual*
shotgun metagenomes. Nucleic Acids Research, 2019.
23. Liu, C.-C., et al., *MetaDecoder: a novel method for clustering metagenomic contigs.*
Microbiome, 2022. 10(1): p. 46.
24. Sieber, C.M.K., et al., *Recovery of genomes from metagenomes via a dereplication,*
aggregation and scoring strategy. Nature Microbiology, 2018. 3(7): p. 836-843.
25. Seemann, T., *Prokka: rapid prokaryotic genome annotation.* Bioinformatics, 2014.
30(14): p. 2068-2069.
26. Nawrocki, E., *Structural RNA homology search and alignment using covariance models.*
2009, Washington University: St. Louis, MO, USA. p. 281.
27. Price, M.N., P.S. Dehal, and A.P. Arkin, *FastTree 2 – Approximately Maximum-*
Likelihood Trees for Large Alignments. PLoS ONE, 2010. 5(3): p. e9490.
28. Katoh, K. and D.M. Standley, *MAFFT multiple sequence alignment software version 7:*
Improvements in performance and usability. Molecular Biology and Evolution, 2013.
30(4): p. 772-780.
29. Capella-Gutiérrez, S., J.M. Silla-Martínez, and T. Gabaldón, *trimAl: a tool for automated*
alignment trimming in large-scale phylogenetic analyses. Bioinformatics, 2009. 25(15):
p. 1972-1973.
30. Nguyen, L.T., et al., *IQ-TREE: a fast and effective stochastic algorithm for estimating*
maximum-likelihood phylogenies. Mol. Biol. Evol., 2015. 32(1): p. 268-74.
31. Edgar, R.C., *MUSCLE: multiple sequence alignment with high accuracy and high*
throughput. Nucleic Acids Research, 2004. 32(5): p. 1792-1797.
32. Jain, C., et al., *High throughput ANI analysis of 90K prokaryotic genomes reveals clear*
species boundaries. Nature Communications, 2018. 9(1): p. 5114.
33. Darling, A.E., B. Mau, and N.T. Perna, *progressiveMauve: Multiple genome alignment*
with gene gain, loss and rearrangement. PLoS ONE, 2010. 5(6): p. e11147.
34. Kim, S., N. Gupta, and P.A. Pevzner, *Spectral Probabilities and Generating Functions of*
Tandem Mass Spectra: A Strike against Decoy Databases. Journal of Proteome Research,
2008. 7(8): p. 3354-3363.
35. Monroe, M.E., et al., *Brief Communication: MASIC: A software program for fast*
quantitation and flexible visualization of chromatographic profiles from detected LC-
MS(/MS) features. Comput. Biol. Chem., 2008. 32(3): p. 215–217.
36. Polpitiya, A.D., et al., *DAnTE: a statistical tool for quantitative analysis of -omics data.*
Bioinformatics, 2008. 24(13): p. 1556-8.
37. Alowaifeer, A.M., et al., *Aerobic methane synthesis and dynamics in a river water*
environment. Limnology and Oceanography, 2023. 68(8): p. 1762-1774.

38.   Angel, R., P. Claus, and R. Conrad, *Methanogenic archaea are globally ubiquitous in*
396   *aerated soils and become active under wet anoxic conditions.* The ISME Journal, 2012.
397   6(4): p. 847-862.
398   39.   Murray, N.E., *Type I restriction systems: sophisticated molecular machines (a legacy of*
399   *Bertani and Weigle).* Microbiol Mol Biol Rev, 2000. 64(2): p. 412-34.
400   40.   Bickle, T.A. and D.H. Krüger, *Biology of DNA restriction.* Microbiol Rev, 1993. 57(2): p.
401   434-50.
402   41.   Borton, M.A., et al., *Comparative genomics and physiology of the genus*
403   *Methanohalophilus, a prevalent methanogen in hydraulically fractured shale.*
404   Environmental Microbiology, 2018. 20(12): p. 4596-4611.
405   42.   Martens, J.H., et al., *Microbial production of vitamin B12.* Appl Microbiol Biotechnol,
406   2002. 58(3): p. 275-85.
407
408