Bottom trawling impacts on faunal and microbial communities increase seafloor carbon loss

Guido Bonthond[1,2,3,4]*, Jan Beermann[3], Alexa Wrede[3], Vera Sidorenko[3], Peter J. Schupp[1,4], Lars Gutow[3]

[1]   Institute for Chemistry and Biology of the Marine Environment (ICBM), School of Mathematics
and Science, Carl von Ossietzky Universität Oldenburg, Ammerländer Heerstraße 114-118, 26129
Oldenburg, Germany
[2]   Thünen Institute of Sea Fisheries, Herwigstraße 31, 27572 Bremerhaven, Germany
[3]   Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Am Handelshafen 12,
27570 Bremerhaven, Germany
[4]   Max Planck Institute for Marine Microbiology, Bremen, Germany
[5]   Helmholtz Institute for Functional Marine Biodiversity at the University of Oldenburg
(HIFMB), Ammerländer Heerstrasse 231, D-26129 Oldenburg, Germany
*   Correspondence: guido.bonthond@awi.de

ABSTRACT

The seafloor is the largest organic carbon sink on Earth. Its benthic fauna and microbiota jointly
regulate long-term carbon storage by assimilating and degrading sedimentary organic matter (SOM).
Bottom trawling threatens the long-term storage of organic matter by physical disturbance of the
seafloor and remobilization of SOM. Trawling also impacts faunal and microbial communities and their
diversity, but the implications of these ecological consequences for long-term carbon storage
dynamics are poorly understood.

Here, we combined datasets of paired faunal and microbial communities from the North Sea and
used structural equation modeling to disentangle direct and biotically mediated effects of bottom
trawling on SOM. One third of trawling-induced SOM loss is indirect and mediated by shifts in benthic
faunal and microbial communities. Microbial community structure directly affected SOM and
represented a key link in this pathway. Microbial effects on SOM were strongly tied to changes in
metabolic traits. SOM loss coincides with increased microbial aerobic respiration potential, whereas
SOM increases correlated with microbial assimilation processes, including dark carbon fixation.
Benthic fauna impact SOM indirectly by shaping benthic microbial communities. Faunal diversity and
species' traits related to bioturbation promoted shifts in benthic microbiota that favor OM storage,
corroborating their indirect but essential role in seafloor carbon dynamics.

This study reveals that bottom trawling alters seafloor carbon storage not only through physical
disturbance but also by disrupting faunal-microbial interactions that regulate sediment carbon
dynamics. Taxonomic and functional shifts in benthic microbiota are a key, but underestimated,
component of the processes regulating the seafloor carbon sink and warrant better integration of
benthic biodiversity and ecological processes into assessments of seafloor carbon storage under
anthropogenic pressures.

Keywords: Carbon sequestration, bottom trawling, structural equation models, sediment microbiota,
benthic fauna

INTRODUCTION

Marine sediments cover nearly 70% of Earth's surface and critically contribute to the long-term storage of organic matter (OM), of which organic carbon (OC) is the major constituent (Snelgrove *et al.* 2014; Atwood *et al.* 2020). Sedimentary organic matter (SOM) originates from marine primary production (phytoplankton and macrophytes) and terrestrial runoff that is deposited as detritus on the seafloor. This detritus is degraded by benthic invertebrates and microbes, but a fraction escapes heterotrophic consumption and may be buried and stored in the seafloor for thousands to millions of years (Estes *et al.* 2019; Atwood *et al.* 2020). As the ultimate decomposers of OM, benthic invertebrates and microbial heterotrophs jointly regulate the storage of OM in the seafloor and thereby play a key role in the carbon cycle.

Carbon sequestration is especially pronounced in productive coastal shelf seas, globally accounting for more than 85% of all OC that is annually buried in marine sediments (Atwood *et al.* 2020; Epstein *et al.* 2022). At the same time, shelf seas are most exposed to multiple human impacts, such as pollution, sand extraction, mining, infrastructure construction, and bottom trawling by demersal fisheries (Amoroso *et al.* 2018; Coates *et al.* 2014; Harris 2020; Hiddink *et al.* 2017; Porz *et al.* 2026), which may affect the capacity of sediments for OC sequestration (Epstein *et al.* 2022). In particular, bottom trawling impacts benthic ecosystems by altering seabed morphology (Puig *et al.* 2012) and resuspending large quantities of sediment (Breimann *et al.* 2022), thereby reducing OC storage in exchange for carbon dioxide emissions (Bradshaw *et al.* 2024; Tiano *et al.* 2024; Zhang *et al.* 2024). Furthermore, trawling causes high mortality in benthic macrofauna (Tillin *et al.* 2006; Hiddink *et al.* 2017; Sciberras *et al.* 2018) and alters benthic microbial communities and metabolism (Bruce *et al.* 2022; Bonthond *et al.* 2023), which may even be detectable in microbial communities of seafloor-inhabiting holobionts, such as flatfish (Gwinner *et al.* 2026).

The impact of bottom trawling on seafloor carbon storage is a matter of growing concern. Trawling promotes the degradation of the more reactive fraction of SOC, but its effect on total SOC is context-dependent (Porz *et al.* 2024; Tiano *et al.* 2024; Zhang *et al.* 2024). While mechanical impacts (i.e., heavy gear disturbing the sediment) directly remobilize OC, the depletion of benthic fauna can also have consequences for SOC (Epstein *et al.* 2022), as these organisms degrade and assimilate detritus and actively rework the sediment matrix through bioturbation. Bioturbation broadly refers to all faunal transport processes within the sediment, including particle reworking and burrow ventilation (Kristensen *et al.* 2012). Bioturbating macrofauna also influence benthic microbial communities by transporting oxygen and OC to deeper sediment layers, and by homogenizing benthic microbiota organized along redox gradients (Deng et al. 2020; Middelburg 2018). Accordingly, trawling-induced shifts in microbial diversity and metabolism (Bruce *et al.* 2022; Bonthond *et al.* 2023) may be partially mediated by changes in macrofaunal communities.

Benthic microbiota in turn degrade and assimilate detritus and therewith impact the storage and cycling of OC (Jiao *et al.* 2010; LaRowe *et al.* 2020). However, the roles of microbes in SOM cycling, and how their interactions with benthic macrofauna influence SOM, are not well understood (Jiao *et al.* 2010; Middelburg 2018; LaRowe *et al.* 2020). In terrestrial environments, microbiota are known to critically impact the accumulation and degradation of OC in soils. Due to their capacity to decompose recalcitrant OC, microbial respiration can enhance OC remineralization and reduce soil OC (De Graaff *et al.* 2015). At the same time, microbial biomass and byproducts promote OC stabilization and sequestration (Derrien *et al.* 2023; Tao *et al.* 2023). If microbial processes are of similar importance for OC storage in the seafloor, the interplay among macrofauna and microbiota could importantly mediate anthropogenic impacts, such as bottom trawling, on seafloor carbon storage.

Here, we used an extensive paired dataset of benthic macrofaunal and microbial communities to disentangle direct and indirect effects of bottom trawling on SOM. One hundred and fifty sites were sampled across a regional scale in the southeastern North Sea, a region subject to variable fishing intensities and characterized by a broad range of SOM contents. We used structural equation models (SEMs) to analyze how SOM is jointly shaped by abiotic factors (sediment characteristics, temperature, shear stress, trawling) and by macrofaunal and microbial communities. We hypothesized that benthic macrofauna and microbiota mediate the effects of bottom trawling on SOM. To test this hypothesis,

98 we developed a conceptual framework including different potential abiotic and biotic pathways
99 through which trawling may influence SOM. Further, we used multivariate and penalized generalized
100 linear models to identify key faunal and microbial traits and taxa, responding to trawling, and with
101 impact on SOM.
102
103 RESULTS
104 **Sampling and data description.** Macrofaunal and microbial communities were characterized for
105 150 sites in the southeastern North Sea (Figure 1). Sediment organic matter contents varied from
106 0.063 to 5.02% weight and trawling intensities (in swept area ratio per year, SAR y^{-1}) from close to zero
107 (0.016 SAR y^{-1}) to heavy trawling rates of up to 1.262 SAR y^{-1}.
108
Benthic microbiota. The microbial dataset counted 69,556 OTUs with 1,293 different genus
classifications. Non-metric multidimensional scaling (NMDS) revealed various trends associated with
microbial community structure (Figure 2A). Apart from shear stress and microbial diversity, all
variables were significantly ordinated on the NMDS space (p < 0.05, Table S1). SOM increased along
the second NMDS axis, while trawling intensity decreased (Figure 2B-C). Microbial community
structure also varied with faunal community properties. The biomass, the community bioturbation
potential (BPc, Queirós *et al.* 2013), and the community bioirrigation potential (IPc, Wrede *et al.* 2018)
were ordinated in similar directions, increasing along NMDS1 and NMDS2. Faunal diversity decreased
along both axes.
Pseudomonadota was the most abundant phylum, followed by Bacteroidetes and Planctomycetes.
Compositional changes along NMDS1 included a shift within the Pseudomonadota from Delta- to
Gammaproteobacteria. Bacteroidetes decreased along NMDS1 while Nitrospirae increased.
Actinobacteria transitioned from Actinomarinales to Microtrichales (Figure 2B). Compositional
changes along NMDS2 were characterized by a strong decrease in Planctomycetes and an increase in
Nitrospirae, but were otherwise less pronounced at higher taxonomic ranks (Figure 2C).
124
Benthic macrofauna. In total, 301 faunal taxa were identified, of which 218 taxa were classified to
species and 39 to genus level. Besides bottom shear stress, all variables were significantly ordinated
onto the NMDS space (p < 0.05, Figure 2D). Compositional changes along NMDS1 coincided with a
decrease in mud content and an increase in temperature and median grain size. Along NMDS2, SOM
increased and trawling intensity decreased (Figure 2F). Faunal biomass, BPc, and IPc also increased
along NMDS2. Faunal diversity decreased along both NMDS axes, and microbial diversity increased
along NMDS1 and decreased along NMDS2.
Annelida were the largest and most diverse phylum (36.7% relative abundance, 115 taxa, Figure
2E-F). Phoronida and Chordata were also among the most abundant phyla (24.0% and 15.2% relative
abundance), but counted only two taxa each. NMDS1 showed a strong shift within the Annelida from
Spionida to Phyllodocida and Polygordiidae, and to communities dominated by *Branchiostoma*
lanceolatum (Chordata, Figure 2E). Along NMDS2, faunal community composition gradually shifted
from communities dominated by Annelida, along high abundances of both Annelida and Arthropoda,
toward communities dominated by *Phoronis muelleri* (Phoronidae, Figure 2F).
139
Structural equation modeling of abiotic and biotic effects on microbial composition and SOM. The
initial SEM did not support faunal biomass and the IPc as informative predictors for microbial
community composition or SOM. Accordingly, they were excluded from the final model, resulting in a
better fit (Fisher's C = 51.893, df = 42, p = 0.141, AICc = 4433.95, Table S2-4). Median grain size, mud
content, and trawling intensity had direct effects on faunal community composition. Faunal diversity
was directly dependent on faunal community composition, increased with temperature, and
decreased with trawling intensity (Figure 3A). Direct abiotic effects on microbial community structure
were in descending order of strength: median grain size, mud content, and trawling intensity.
Microbial community structure was also dependent on faunal composition, the BPc, and faunal
diversity. Consequently, fauna mediated the effects of temperature, mud content, median grain size,

and trawling intensity on microbial composition. These indirect effects were substantial and equaled
about half of the direct effect in case of mud content and median grain size. For trawling intensity, the
indirect effect on benthic microbial composition was even larger than the direct effect (Figure 3B).
Microbial diversity varied with microbial and faunal composition, but not significantly with faunal
diversity. Shear stress was the only predictor for which no effects were statistically supported.
Bottom trawling effort directly reduced SOM. Additionally, the microbial latent variable NMDS2
was significant, indicating a direct effect of microbial community structure on SOM (Figure 3A). The
effect of microbial diversity on SOM was not significant, and neither were faunal composition, BPc and
faunal diversity. Several pathways were supported through which faunal composition, BPc and faunal
diversity influenced SOM indirectly. The indirect effects of faunal diversity and BPc on SOM where
both positive, whereas the indirect effect of trawling was negative (Figure 3C).
161
Bioturbation related traits and taxon level effects. Trawling impacted in total 74 predicted
pathways (Figure 4A, Figure S3A), which were mostly related to biosynthesis (13 decreasing, 64
increasing). In addition, especially pathways related to energy metabolism (belonging to the categories
'generation of precursor metabolites' and 'degradation/utilization/assimilation)' increased in response
to trawling (6 and 5, respectively), including the TCA cycle I (*TCA*) and VI (*PWY-5913*), and aerobic
respiration I (*PWY-3781*). Various OTUs varied with trawling effort (Figure 4B). The relative abundance
of OTUs within the Actinobacteria and Planctomycetes, particularly increased with trawling effort,
whereas Alphaproteobacteria and Deltaprotebacteria tended to decrease. Fifteen faunal traits
declined with trawling effort and none increased (Figure 4C), suggesting an overall decrease in faunal
functional diversity. This was also reflected at the species level, where the abundance of 27 species
decreased and only 12 species increased in response to trawling (Figure 4D). In particular, species
belonging to Annelida, Mollusca and Nemertea decreased with trawling intensity, whereas in other
phyla increasing and decreasing taxa were more balanced.
In total, eight trait categories were identified with pGLMs to explain changes in microbial
community structure (Figure 4A). Most of these traits were related to mobility (limited movement,
free movement through sediment matrix, free movement through burrows) and reworking (bio-
diffusors, up- and downward conveyors, surficial modifiers), but also included the feeding type
category 'subsurface filter feeders', and the injection pocket depth category 'below 10 cm'. The model
retained all environmental covariates (i.e., mud content, median grain size, trawling intensity), the
faunal latent variables, and faunal diversity. When analyzing taxa instead of traits, 30 faunal taxa were
identified to shape microbial community composition (Figure 4B). More than half of these taxa
belonged to the Annelida, with the two species *Eteone longa* and *Aonides paucibranchiata* having the
highest impact. But also *B. lanceolatum* (Chordata) and *Echinocardium pennatifidum* (Echinodermata)
were among the species with highest effect scores.
Twenty-seven predicted microbial pathways were resolved as informative predictors of SOM
(Figure 4C). Eighteen pathways were negatively associated with SOM, of which the strongest effect
was observed for aerobic respiration (*PWY-3781*). Out of the nine pathways associated with an
increase in SOM, the two strongest regularized coefficients were found for gluconeogenesis
(*GLUCONEO-PWY*) and the Calvin cycle (*CALVIN-PWY*). Twenty-five OTUs were associated with SOM
decrease, and four OTUs with SOM increase (Figure 4D). Planctomycetes were prominent among the
OTUs related to SOM decrease (nine in total), but the strongest negative association was found for an
Actinobacterial OTU, classified to Sva0996 marine group. From the covariates, trawling intensity,
microbial NMDS2, and predicted functional microbial diversity were retained in both pGLMs
evaluating trait and species effects on SOM.
196
DISCUSSION
Our study confirms that bottom trawling reduces OC storage in marine sediments. While the
trawling impact is partially direct, likely resulting from physical seabed disturbance, approximately one
third of the overall bottom trawling impact on SOM is indirect, mediated by changes in benthic
macrofaunal and microbial communities. Our analysis provides insight into these indirect trawling

effects, and reveals several pathways through which coexisting fauna and microbiota shape the
sequestration of carbon. The direct link between benthic microbial community structure and SOM is
central for these pathways. Consequently, benthic microbiota serve as a key mediator of various biotic
and abiotic effects on OC storage. While macrofaunal communities could not be directly linked with
SOM, they strongly impact microbial composition, marking an indirect, but pivotal, effect of
macrofauna on SOM. The net effects of faunal diversity and bioturbation on SOM are positive. These
results indicate that the indirect impacts of bottom trawling on SOM, cascading through macrofaunal
and microbial communities, have net negative consequences for the seafloor carbon stocks. Together,
our findings emphasize the central role of benthic fauna and microbiota, and the interaction among
them, in OC sequestration and underscore the need to consider both physical and ecological
processes for the assessment of long-term consequences of bottom trawling on seafloor carbon
dynamics.
214
Trawling reduces SOM by restructuring benthic fauna and microbiota. The impact of bottom
trawling on OM storage in sediments entails a combination of mechanisms (Epstein *et al.* 2022; Zhang
et al. 2024). The mechanical resuspension of OM is the most direct effect, as it locally mobilizes OM
from the sediment, which can then become subject to export by currents, and is partially
remineralized by heterotrophs before being deposited, and buried again. In addition, physical remixing
of the upper sediment layers enhances benthic remineralization of OM in the sediment through
aerobic microbial respiration. Previous studies have demonstrated that the extended exposure to
oxygen, caused by resuspension and remixing, decreases SOM, providing estimates of trawling
induced loss of OC stored in the seafloor (Sala *et al.* 2021; Hiddink *et al.* 2023; Porz *et al.* 2024; Zhang
et al. 2024). Our results confirm the negative direct impact of trawling on SOM. Additionally, our study
reveals that trawling-induced shifts in benthic macrofaunal and microbial communities also have
important consequences for carbon storage, supporting our hypothesis that macrofauna and
microbiota mediate the effects of bottom trawling on SOM. Trawling alters the communities of
organisms involved in the processing and degradation of SOM. The contribution of benthic biota to
SOM dynamics has been acknowledged in previous studies (Epstein *et al.* 2022; Zhang *et al.* 2024).
However, the biotic pathways involved have been poorly characterized and are not explicitly
considered in predictive models of sedimentary carbon storage. Instead, these models typically
estimate SOM remineralization as a function of extended oxygen exposure associated with trawling-
induced sediment disturbance, treating the benthic biota rather as a 'black-box' without resolving the
contribution of the fauna and microbiota mechanistically (Sala *et al.* 2021; Zhang *et al.* 2024). Our
analysis, however, demonstrates that indirect, biota-mediated effects are crucial and complex,
accounting for roughly one third of the total trawling impact on SOM (Figure 3C).
Benthic microbiota were identified as a key mediator of trawling impacts on SOM. Our model
showed that compositional shifts in benthic microbiota are directly linked to changes in SOM,
highlighting the prominent role of microbial processes in shaping SOM. Bottom trawling induces
changes in the composition and energy metabolism of benthic microbial communities promoting
specifically aerobic metabolic pathways (Bonthond *et al.* 2023). The responses to trawling varied
between microbial groups with some groups apparently benefiting from trawling while other groups
clearly declined in response to trawling. The study confirmed that the predicted aerobic respiration
pathway increased with trawling effort, indicating that microbial aerobic heterotrophic metabolism is
more prevalent in sediments exposed to intense trawling. These taxonomic and functional shifts in the
microbial communities explain a net loss in SOM, highlighting microbial aerobic respiration as the
critical process by which SOM is removed from the food web (Middelburg 2018).
The impacts of trawling on benthic fauna also have negative consequences for seafloor carbon
storage. In line with previous research (see Sciberras *et al.* 2018 and references therein), various traits
related to bioturbation also benthic faunal diversity reduced with trawling intensity. This trawling-
induced decline in structural and functional diversity of the benthic macrofauna coincided with, but
did not directly cause, a decrease in SOM. Instead, trawling induces a cascade of responses involving
both benthic macrofauna and microbiota. Losses in response to trawling were particularly pronounced

in annelids and mollusks and especially these groups impacted microbiota. As macrofaunal diversity
and bioturbation regulate benthic microbiota, there is an indirect negative net effect of trawling on
SOM, mediated by the benthic fauna. Accordingly, marine protection efforts to enhance benthic
faunal diversity have the potential to also promote seafloor carbon storage.
258
Carbon storage is directly affected by benthic microbiota. Aerobic respiration is the terminal
process that catalyzes the conversion of OC to carbon dioxide (Middelburg 2018; Jørgensen *et al.*
2022). In the pelagic environment, OC is primarily recycled through the microbial loop, where OC
assimilated by microbes is partially transferred to higher trophic levels through predation and grazing
(Azam *et al.* 1983). The transfer of microbially processed OC to higher trophic levels in sediments is,
however, limited and most OC consumed by microbes is eventually lost through microbial respiration,
a concept described as the 'inverted microbial loop' (Middelburg 2018). In accordance with this,
microbial aerobic respiration emerged in our analysis as the most important pathway associated with
SOM loss. In Bonthond *et al.* (2023) and in this study, we found that the predicted aerobic respiration
pathway increased with trawling effort, indicating that microbial aerobic heterotrophic metabolism is
more prevalent in sediments exposed to more trawling. Both macrofauna and microbes carry out
aerobic respiration. However, they partially utilize different OM substrates from different sources. This
likely determines whether SOM is being decomposed and cycled back into the food web through
assimilation, or whether it escapes degradation and becomes sequestered. Living tightly attached to
sand grains, microbes utilize sediment-bound particulate OM (Arnosti 2011). This OM is characterized
by its low reactivity with the labile OM fraction typically accounting for <10% (Pusceddu *et al.* 2009;
Arndt *et al.* 2013). Among prokaryotes, sophisticated metabolic pathways have evolved, enabling the
degradation of even the most recalcitrant molecules (Jiao *et al.* 2010; LaRowe *et al.* 2020), which are
stable and most likely to be sequestered. Essentially, under adequate conditions, there are no
biomolecules known that cannot be degraded by microorganisms (LaRowe *et al.* 2020; Dittmar *et al.*
2021). One of the phylum-level microbial shifts associated with decreasing SOM was a relative
increase in Planctomycetes, which include microbial groups specialized in decomposing high molecular
weight and diverse recalcitrant OM substrates (Probandt *et al.* 2017; Wiegand *et al.* 2018).
Additionally, Planctomycetes were prevalent among the OTUs that increased with trawling effort,
indicating that bottom trawling not only promotes OM remineralization by enhancing oxygen
exposure, but also by favoring microbial metabolic degradation of less reactive OM, which is more
probable to be sequestered. Thus, trawling not only enhances the exposure of SOM to oxygen, but
also to microbial decomposers capable of remineralizing low-reactive OM.
Our analysis identified the Calvin cycle as a primary predicted microbial function associated with
SOM increase. As light availability at the sampled sites is negligible to absent, this likely reflects dark
carbon fixation by chemoautotrophic microbes, a common process in marine sediments, largely driven
by Gammaproteobacteria (Hügler & Sievert 2011; Middelburg 2011; Dyksma *et al.* 2016). The strong
increase of SOM with the gluconeogenesis pathway is further in line with this. Gluconeogenesis is an
assimilatory pathway, which synthesizes glucose from smaller precursors, such as fatty acids and
amino acids, and indirectly from carbon dioxide (Sharkey & Weise 2012). Dark carbon fixation in
sediments is recognized as an important contribution to the input of new OC into the oceans and its
burial in sediments (Middelburg 2011), but has not been considered in the context of bottom trawling.
Accordingly, trawling may not only enhance the remineralization of OC but also reduce the benthic OC
production by chemoautotrophs.
The weak positive effect of the predicted functional microbial diversity on SOM suggests that
degradation and accumulation of SOM involves 'diversity effects'. Diversity effects define
contributions of a community to processes that cannot be explained by the sum of the contribution of
its single members, and are known to play an important role in OC storage in terrestrial ecosystems
(Gessner *et al.* 2010). The decomposition of organic substrates involves complex trophic interactions,
as well as competition and cooperation among primary degraders of recalcitrant molecules (e.g.,
cellulose, lignin, laminarin) and secondary consumers of more labile OM (e.g., acetate, lactate, amino
acids; Gralka *et al.* 2020). Functional diversity and redundancy influence these interactions and

increase carbon use efficiency, potentially promoting OC respiration and decomposition with negative impact on OM storage (De Graaff *et al.* 2015). At the same time, the increased generation of microbial biomass and more stable by-products can promote OC accumulation (Tao *et al.* 2023). Our results indicate that microbial functional diversity influences OC cycling in marine sediments in a similar manner to terrestrial soils. This is of particular importance in the context of bottom trawling impacts on seafloor carbon dynamics, which can decrease microbial taxonomic and functional diversity (Bonthond *et al.* 2023).

Faunal communities influence SOM by regulating benthic microbiota. Faunal composition, bioturbation, and diversity had no direct effect on SOM but significantly influenced microbial community structure, highlighting that benthic microbial diversity and community composition are shaped not only by abiotic factors (Bonthond *et al.* 2023)**,** but also by the composition of macrofaunal communities. Consequently, macrofauna mediate the effects of different environmental processes on benthic microbiota, constituting a key link between bottom trawling and carbon sequestration.

A direct effect of bioturbation on SOM was not supported, suggesting only minor effects of sediment reworking on SOM. Alternatively, antagonistic processes inherent to bioturbation, such as up- and downward OC transport, may minimize the net effect of bioturbation. Our results indicate that bioturbation essentially modulates benthic microbiota. This is in agreement with previous work, that demonstrated that faunal bioturbation activity shapes benthic microbiota by vertically mixing OC substrates and homogenizing microbial communities (Deng *et al.* 2020). The BPc, which is a commonly used index to estimate the community bioturbation potential (Queirós *et al.* 2013), explained shifts in benthic microbiota. Specifically, faunal traits related to mobility and sediment reworking activity, which are used to calculate the BPc, were identified as important predictors of SOM (Figure 5A) and also declined with trawling intensity (Figure 4C). Bioturbators have been shown to promote aerobic microbial OC degradation by enhancing oxygen supply to the sediment and its microbiota (Baranov *et al.* 2016). At the same time, macrofauna compete with the microbiota for oxygen (Middelburg 2018; Jørgensen *et al.* 2022), which may locally limit microbial aerobic respiration. The strong impact of the faunal trait 'limited movement' on benthic microbiota may indicate a suppression of microbial aerobic respiration by stationary animals that consume oxygen within the sediment but enhance its resupply only minimally. Additionally, benthic fauna may contribute to the stabilization of SOM as their feces and remains contain more recalcitrant OC compared to freshly deposited detritus and are therefore degraded less efficiently by microbes (Arndt *et al.* 2013; LaRowe *et al.* 2020).

Both the SEM and pGLMs consistently showed that changes in the microbial community were explained by macrofaunal diversity, suggesting that benthic microbiota are not only shaped by specific animal traits, but may also depend on 'diversity effects', such as resource partitioning, cooperation and facilitation (Fridley 2001; Gessner *et al.* 2010). Moreover, the positive indirect effect of macrofaunal diversity on SOM implies that diverse faunal communities mitigate microbial remineralization of SOM. In other words, faunal diversity favors OC storage. In marine sediments, studies on diversity effects are scarce, particularly in the context of interacting faunal and microbial communities. In terrestrial soils, complementarity processes (i.e., niche partitioning and functional differentiation among taxa) can increase OC assimilation into the food web and enhance OC decomposition (Gessner *et al.* 2010). Our results indicate similar processes in marine sediments. While this study focused on macrofauna and microbiota, meiofauna, an important component of benthic diversity and biomass (Gerlach 1971), were not considered. Meiofauna constitute a critical link between macrofauna and microbiota in the regulation of marine biogeochemical processes (Bonaglia *et al.* 2014). Accordingly, involving meiofauna in future studies will improve the mechanistic understanding of how the different components of the benthic biota interact and control carbon sequestration in marine sediments.

CONCLUSIONS

The impacts of bottom trawling on SOM are not solely the result of direct physical disturbance, but
also emerge from cascading effects involving benthic fauna and microbiota. Our results underscore
the importance of ecological processes in trawling impacts on carbon sequestration, and demonstrate
that benthic microbiota play a central role in seafloor carbon dynamics. Approximately one third of
the trawling-associated decline in SOM is mediated by shifts in faunal and microbial communities,
highlighting the need to consider biotic pathways in assessments of the impacts of bottom trawling on
carbon sequestration. Considering these ecological processes in predictive models will be critical for
more accurately forecasting the development of seafloor carbon stocks in the face of increasing
anthropogenic disturbance. Future research should further disentangle how specific microbial and
faunal functions interact with labile and recalcitrant OM to ultimately improve our understanding of
seafloor carbon sequestration.
368
369
370 **MATERIALS AND METHODS**
371
Sampling and sediment properties. Macrofauna and microbiota were sampled with van Veen grabs
(0.1 m^2, minimum penetration depth: 10 cm) from 150 sites in the southeastern North Sea (Figure 1)
with the German research vessel 'Heincke' in August 2019 (cruise HE538, doi:10.2312/cr_he538) and
September 2020 (cruise HE562, doi:10.48433/cr_he562). At each site, one or three grabs were taken
to characterize macrofaunal communities, and sediment samples were obtained from an additional
grab taken to characterize microbial communities, grain size metrics, and SOM. The sediment
temperature was measured on board using a standard thermometer (~3 cm depth) from the first grab
immediately after recovery. GPS coordinates were recorded from the ship-logs as the ship position at
the time of bottom contact.
Samples for DNA extraction were taken from the top centimeter of the sediment directly in the
grab (HE538) or with small vertical mini-cores (HE562), from which the upper centimeter was later
sampled in the lab (details in Bonthond *et al.* 2023). A separate core (ø = 4.5 cm, penetration depth: 6
cm) was taken to characterize the sediment grain size distribution (Wentworth scale, Wentworth
1922) and total SOM content. Samples were stored at -20 °C until further processing. Subsamples of
40 g wet weight were dried and combusted for 5 hours at 500 °C to calculate SOM via loss on ignition
(LOI) and characterize grain size distributions using a sieve cascade (details in Bonthond *et al.* 2023).
388
Characterization of benthic microbiota. Benthic microbiota were characterized previously in
Bonthond *et al.* (2023), where DNA extractions, library preparation, bioinformatic processing, OTU
clustering (based on the 97% similarity criterion), and taxonomic classification are described in detail.
For the present study, the final OTU table was rarefied by subsampling the count table a 100 times to
a 1000 reads. Metagenomes were predicted with PICRUST2 v2.5.3 (Douglas *et al.* 2020), and MetaCyc
pathway annotations were used for analyses.
395
Characterization of benthic macrofauna and community properties. The van Veen grab contents
were sieved (1 mm mesh size) onboard. Retrieved individuals of the benthic macrofauna were
preserved in buffered 4 % formaldehyde-seawater solution and identified to the lowest taxonomic
level possible in the laboratory and counted. The taxonomy was matched against the World Register
of Marine Species (WoRMS Editorial Board 2024, https://www.marinespecies.org). The biomass was
measured as fresh weight at the species level.
402
Data preparation. To combine microbial and macrofaunal community data, datasets were reduced
to a single observation per site. For sites with multiple mini-cores per sediment grab, the microbial
community from the core taken from the most central position inside the grab was used. For sites
with multiple macrofauna grabs, the grab nearest to the microbial grab was selected. As potential
informative abiotic predictors for microbial and macrofaunal community composition and SOM, we
considered the variables median grain size (µm), mud content (% dry weight), sediment temperature

(°C), bottom shear stress (N/m^2), and fishing intensity in terms of swept area ratio per year (SAR y^{-1}).
Fishing intensity was obtained from the OSPAR data & information management system (OSPAR 2017)
and bottom shear stress was derived from the barotropic FESOM-C model (Androsov *et al.* 2019).
Equivalent numbers of species, OTUs, and predicted MetaCyc pathways were used as a measure of
faunal diversity, microbial diversity, and predicted functional microbial diversity, respectively (Jost
2006). The community bioturbation potential (BPc) was calculated according to Queirós *et al.* (2013),
based on the biomass and abundance of macrofauna species, and species level scores for the traits
'Mobility' and 'Sediment reworking'. For the community bioirrigation potential (IPc) we used the
equation of Wrede *et al.* (2018), which uses the biomass and abundance of macrofauna species, and
the traits 'Burrowing', 'Feeding type' and 'Injection pocket depth' (see Table S5 for more details on
traits).
As general biotic predictors, we considered faunal biomass, faunal diversity, the BPc, IPc, and
microbial diversity. Correlations among abiotic and biotic variables were explored using Spearman's
rank coefficients (Figure S1).
423
NMDS to characterize macrofaunal and microbial community composition. Non-metric Multi-
Dimensional Scaling (NMDS) was conducted with the R package *vegan* (v2.6-6.1, Oksanen 2010), using
Aitchison distances for microbial communities (accounting for compositionality) and Bray-Curtis
dissimilarities for macrofaunal communities. Abiotic variables and community properties were
ordinated in NMDS spaces with the *enfvit()* function, evaluating significance with permutation tests
with 9,999 permutations restricted within expeditions (Figure 2A, B).
430
Linear mixed effect models (SEM pieces). LMMs were fitted with restricted maximum Likelihood
(REML) using the R-package *lme4* (v1.1-35.5, Bates *et al.* 2015), including the factor 'Expedition' as
random intercept to account for the non-independence associated with expeditions. Moran's I was
computed to infer whether models accounted reasonably for spatial autocorrelation, considering
autocorrelation negligible when |Moran's I|< 0.1. To meet the model assumptions, we log-
transformed the faunal biomass, BPc, IPc, and SOM. Variance inflation factors (VIFs) were computed
to assess multicollinearity, and predictors with VIF values > 5 were excluded from LMMs.
Macrofaunal community structure was modeled by regressing both faunal latent variables against
the abiotic predictors. Faunal biomass, faunal diversity, BPc, and IPc were fitted in response to the
faunal latent variables. For the BPc and IPc, faunal biomass was also included as predictor. As we
hypothesized that the impacts of abiotic variables on benthic microbiota are partially mediated by
changes in macrofaunal communities, LMMs fitted on the microbial latent variables included the
abiotic predictors, faunal biomass, BPc, IPc, faunal diversity, and the faunal latent variables. Based on
variance inflation factor (VIF) scores, the faunal NMDS2 was excluded from both LMMs. Microbial
diversity was regressed against the microbial latent variables, faunal diversity, and faunal latent
variables. The final LMM, with SOM as response, included the microbial latent variables, representing
microbial community composition, and faunal NMDS1 to represent microbial community composition.
In addition, we included faunal and microbial diversity as predictors as they are known to influence
OM cycling in soils (De Graaff *et al.* 2015; Gessner *et al.* 2010; Nielsen *et al.* 2011; Tao *et al.* 2023).
Further, faunal biomass, the BPc and the IPc were included as community level traits with potential
impact on SOM as they may influence SOM positively (e.g., enhancing downward transport of OM)
and negatively (e.g., faunal decomposition, upward transport and resuspension of SOM, Epstein *et al.*
2022). Finally, the model included the abiotic variables bottom shear stress and fishing intensity,
which may reduce SOM through resuspension, potentially resulting in subsequent degradation of OM
in the water column or transport to other areas (Epstein *et al.* 2022). Hence, temperature and
sediment properties were only included as indirect effects on SOM in our SEMs, as they influence
macrofaunal and microbial processes by regulating (heterotrophic) metabolic rates (Clarke & Fraser
2004) and the oxygen penetration depth (Neumann *et al.* 2021) but do not influence SOM directly.
459

Structural Equation Models (SEMs). We used structural equation modeling to disentangle the direct and indirect effects of abiotic and biotic variables on benthic microbiota and SOM. Faunal and microbial NMDS axes were used as latent variables, representing community structure. First, a conceptual causal network was constructed (Figure S2A) based on findings from previous work (Bonthond *et al.* 2023) and from literature (Gessner *et al.* 2010; Sciberras *et al.* 2018; Deng *et al.* 2020; Tiano *et al.* 2024). Using the framework of the *piecewiseSEM* R package (v2.3.0.1, Lefcheck 2016), LMMs were compiled into a SEM. After inspecting correlations, we included the following environmental variables: temperature, median grain size, mud content, bottom shear stress, and fishing effort. As it was found in Bonthond *et al.* (2023) that the median grain size and mud content can have opposite effects on microbial diversity and certain functions, it was decided to retain both variables in the analysis, even though they both represent sediment properties. In the first model, we included the faunal biomass, BPc, and IPc together, although these were strongly correlated ($\rho > 0.75$, Figure S1). Shipley's test of d-separation revealed the presence of direct effects of temperature and trawling intensity on faunal diversity, which were added to the model (Figure S2). In addition, correlation terms were added to the global SEM between the IPc and BPc, microbial NMDS1 and NMDS2, and faunal NMDS2 and microbial NMDS2. As the faunal biomass and IPc were not informative as predictors, a reduced SEM was fitted without both responses, and both models were refitted with Maximum Likelihood and compared based on the corrected Akaike information criterion (AICc). The overall SEM fit was evaluated with Shipley's test of d-separation obtained through Fisher's C statistic and the significance of direct effects was assessed by bootstrapping the overall SEM with 9,999 iterations.

Multivariate GLMs to identify trawling impacts on traits and taxa. To identify key traits and taxa responding to trawling effort, multivariate GLMs were fitted using the R package mvabund (v4.2.1), assuming a negative binomial distribution. First a model was fitted with the 100 relatively most abundant predicted pathways in response to the variables also considered in SEMs (temperature, median grain size, mud content, bottom shear stress and bottom trawling effort). P-values, indicating significant changes in relative OTU abundances in response to the predictors were obtained by resampling the model with 499 iterations with the *anova.manyglm* function, using 'Expedition' as blocking variable. The same model was fitted on the faunal traits and on OTUs and faunal taxa with > 0.1% relative abundance.

Penalized GLMs to identify key traits and taxa. The effects of faunal traits and species on microbial community composition and microbial traits and OTUs on SOM were evaluated with separate pGLMs with trait category and species abundances as predictors, using the R package glmmLasso (v1.6.3). First, pGLMs were fitted on both microbial latent variables, using faunal trait categories related to bioturbation ('Mobility', 'Reworking', Queirós *et al.* 2013) and bioirrigation ('Burrow type', 'Feeding type', 'Injection pocket depth' – Wrede *et al.* 2018) as predictors, along with the environmental and community property covariates that were used in the SEM analysis. In a subsequent step, similar models were fitted where faunal traits were replaced by faunal taxon abundances. Second, a pGLM was fitted on SOM, using the microbial traits as predictors in combination with environmental variables and faunal community properties (faunal latent variables, BPc, faunal diversity) as covariates and microbial diversity and predicted functional microbial diversity. This step was repeated using the 100 most abundant OTUs as predictors instead of microbial traits.
In all pGLMs, faunal and microbial traits and species abundances were standardized by subtracting the mean and dividing by the standard deviation. A default lasso penalty of 10 was used to shrink coefficients toward zero in all models. The significance of selected predictors was inferred from 95% confidence intervals obtained by bootstrapping the models with 999 repeated iterations. Finally, for microbial community structure, we pooled the coefficients from both models (fitted on microbial NMDS1 and NMDS2) to estimate a single overall effect size for each predictor.

Data Availability. Fastq files with raw de-multiplexed V3-V4-16S gene amplicon reads were generated in Bonthond *et al.* (2023) and are archived in the sequence read archive (SRA, Bioproject accession number PRJNA988469). The macrofauna data was deposited in CRITTERBASE (https://critterbase.awi.de; Teschke *et al.* 2022). Rendered R-scripts and respective input data for all analyses is available on GitHub at https://github.com/gbonthond/macrofauna-microbiota.

Acknowledgements

The authors are grateful to the crew of the Research Vessel Heincke for support during both expeditions (HE538 in 2019, HE562 in 2020). Dr. Sahar Khodami and Dr. Pedro Martinez-Arbizu are thanked for amplicon sequencing at the Senckenberg am Meer Metabarcoding and SNG laboratory (this is publication number 112).

Funding

This research was part of the project MGF North Sea within the German Marine Research Alliance and was funded by the German Federal Ministry of Education and Research (Grant numbers: 03F0847B, 03F0936C) and the German Federal Agency for Nature Conservation (Grant numbers: 3519532201, 3522521401, LABEL project).

Author contributions

GB, JB and LG conceptualized the study. Field collections were conducted by GB, JB, and LG. GB conducted laboratory work. GB, VS and JB processed data. GB conducted the formal analysis and drafted the manuscript. All authors contributed to writing and revising the manuscript.

Conflict of interest

All authors declare they have no conflicts of interest.

REFERENCES

Amoroso, R.O., Pitcher, C.R., Rijnsdorp, A.D., McConnaughey, R.A., Parma, A.M., Suuronen, P., *et al.* (2018). Bottom trawl fishing footprints on the world's continental shelves. *Proceedings of the National Academy of Sciences*, 115, E10275–E10282.

Androsov, A., Fofonova, V., Kuznetsov, I., Danilov, S., Rakowsky, N., Harig, S., *et al.* (2019). FESOM-C v.2: coastal dynamics on hybrid unstructured meshes. *Geosci. Model Dev.*, 12, 1009–1028.

Arndt, S., Jørgensen, B.B., LaRowe, D.E., Middelburg, J.J., Pancost, R.D. & Regnier, P. (2013). Quantifying the degradation of organic matter in marine sediments: A review and synthesis. *Earth-Science Reviews*, 123, 53–86.

Arnosti, C. (2011). Microbial Extracellular Enzymes and the Marine Carbon Cycle. *Annu. Rev. Mar. Sci.*, 3, 401–425.

Atwood, T.B., Witt, A., Mayorga, J., Hammill, E. & Sala, E. (2020). Global Patterns in Marine Sediment Carbon Stocks. *Front. Mar. Sci.*, 7, 165.

Azam, F., Fenchel, T., Field, J.G., Gray, J.S., Meyer-Reil, L.-A. & Thingstad, F. (1983). The ecological role of water-column microbes in the sea. *Marine ecology progress series. Oldendorf*, 10, 257–263.

Baranov, V., Lewandowski, J. & Krause, S. (2016). Bioturbation enhances the aerobic respiration of lake sediments in warming lakes. *Biol. Lett.*, 12, 20160448.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *J. Stat. Soft.*, 67.

Bonaglia, S., Nascimento, F.J.A., Bartoli, M., Klawonn, I. & Brüchert, V. (2014). Meiofauna increases bacterial denitrification in marine sediments. *Nat Commun*, 5, 5133.

Bonthond, G., Beermann, J., Gutow, L., Neumann, A., Barboza, F.R., Desiderato, A., *et al.* (2023).
Benthic microbial biogeographic trends in the North Sea are shaped by an interplay of
environmental drivers and bottom trawling effort. *ISME commun*, 3.
Bradshaw, C., Iburg, S., Morys, C., Sköld, M., Pusceddu, A., Ennas, C., *et al.* (2024). Effects of bottom
trawling and environmental factors on benthic bacteria, meiofauna and macrofauna communities
and benthic ecosystem processes. *Science of The Total Environment*, 921, 171076.
Breimann, S.A., O'Neill, F.G., Summerbell, K. & Mayor, D.J. (2022). Quantifying the resuspension of
nutrients and sediment by demersal trawling. *Continental Shelf Research*, 233, 104628.
Bruce, S.A., Aytur, S.A., Andam, C.P. & Bucci, J.P. (2022). Metagenomics to characterize sediment
microbial biodiversity associated with fishing exposure within the Stellwagen Bank National Marine
Sanctuary. *Sci Rep*, 12, 9499.
Clarke, A. & Fraser, K.P.P. (2004). Why Does Metabolism Scale with Temperature? *Functional Ecology*,
18, 243–251.
Coates, D.A., Deschutter, Y., Vincx, M. & Vanaverbeke, J. (2014). Enrichment and shifts in
macrobenthic assemblages in an offshore wind farm area in the Belgian part of the North Sea.
Marine Environmental Research, 95, 1–12.
De Graaff, M.-A., Adkins, J., Kardol, P. & Throop, H.L. (2015). A meta-analysis of soil biodiversity
impacts on the carbon cycle. *SOIL*, 1, 257–271.
Deng, L., Bölsterli, D., Kristensen, E., Meile, C., Su, C.-C., Bernasconi, S.M., *et al.* (2020). Macrofaunal
control of microbial community structure in continental margin sediments. *Proc. Natl. Acad. Sci.
U.S.A.*, 117, 15911–15922.
Derrien, D., Barré, P., Basile-Doelsch, I., Cécillon, L., Chabbi, A., Crème, A., *et al.* (2023). Current
controversies on mechanisms controlling soil carbon storage: implications for interactions with
practitioners and policy-makers. A review. *Agron. Sustain. Dev.*, 43, 21.
Dittmar, T., Lennartz, S.T., Buck-Wiese, H., Hansell, D.A., Santinelli, C., Vanni, C., *et al.* (2021).
Enigmatic persistence of dissolved organic matter in the ocean. *Nat Rev Earth Environ*, 2, 570–583.
Douglas, G.M., Maffei, V.J., Zaneveld, J.R., Yurgel, S.N., Brown, J.R., Taylor, C.M., *et al.* (2020). PICRUSt2
for prediction of metagenome functions. *Nat Biotechnol*, 38, 685–688.
Dyksma, S., Bischof, K., Fuchs, B.M., Hoffmann, K., Meier, D., Meyerdierks, A., *et al.* (2016). Ubiquitous
Gammaproteobacteria dominate dark carbon fixation in coastal sediments. *The ISME Journal*, 10,
1939–1953.
Epstein, G., Middelburg, J.J., Hawkins, J.P., Norris, C.R. & Roberts, C.M. (2022). The impact of mobile
demersal fishing on carbon storage in seabed sediments. *Global Change Biology*, 28, 2875–2894.
Estes, E.R., Pockalny, R., D'Hondt, S., Inagaki, F., Morono, Y., Murray, R.W., *et al.* (2019). Persistent
organic matter in oxic subseafloor sediment. *Nature Geosci*, 12, 126–131.
Fridley, J.D. (2001). The influence of species diversity on ecosystem productivity: how, where, and
why? *Oikos*, 93, 514–526.
Gerlach, S.A. (1971). On the importance of marine meiofauna for benthos communities. *Oecologia*,
176–190.
Gessner, M.O., Swan, C.M., Dang, C.K., McKie, B.G., Bardgett, R.D., Wall, D.H., *et al.* (2010). Diversity
meets decomposition. *Trends in Ecology & Evolution*, 25, 372–380.
Gralka, M., Szabo, R., Stocker, R. & Cordero, O.X. (2020). Trophic Interactions and the Drivers of
Microbial Community Assembly. *Current Biology*, 30, R1176–R1188.
Gwinner, M., Haslob, H., Neumann, H., Khodami, S., Schupp, P.J. & Bonthond, G. (2026). Flatfish
intestinal microbiota depend on various host traits, and vary with sediment type and bottom
trawling effort. *Sci Rep*.
Harris, P.T. (2020). Anthropogenic threats to benthic habitats. In: *Seafloor Geomorphology as Benthic
Habitat*. Elsevier, pp. 35–61.
Hiddink, J.G., Jennings, S., Sciberras, M., Szostek, C.L., Hughes, K.M., Ellis, N., *et al.* (2017). Global
analysis of depletion and recovery of seabed biota after bottom trawling disturbance. *Proc. Natl.
Acad. Sci. U.S.A.*, 114, 8301–8306.

Hiddink, J.G., Van De Velde, S.J., McConnaughey, R.A., De Borger, E., Tiano, J., Kaiser, M.J., *et al.*
(2023). Quantifying the carbon benefits of ending bottom trawling. *Nature*, 617, E1–E2.

Hügler, M. & Sievert, S.M. (2011). Beyond the Calvin Cycle: Autotrophic Carbon Fixation in the Ocean.
Annu. Rev. Mar. Sci., 3, 261–289.

Jiao, N., Herndl, G.J., Hansell, D.A., Benner, R., Kattner, G., Wilhelm, S.W., *et al.* (2010). Microbial
production of recalcitrant dissolved organic matter: long-term carbon storage in the global ocean.
Nat Rev Microbiol, 8, 593–599.

Jørgensen, B.B., Wenzhöfer, F., Egger, M. & Glud, R.N. (2022). Sediment oxygen consumption: Role in
the global marine carbon cycle. *Earth-Science Reviews*, 228, 103987.

Jost, L. (2006). Entropy and diversity. *Oikos*, 113, 363–375.

Kristensen, E., Penha-Lopes, G., Delefosse, M., Valdemarsen, T., Quintana, C. & Banta, G. (2012). What
is bioturbation? The need for a precise definition for fauna in aquatic sciences. *Mar. Ecol. Prog.*
Ser., 446, 285–302.

LaRowe, D.E., Arndt, S., Bradley, J.A., Estes, E.R., Hoarfrost, A., Lang, S.Q., *et al.* (2020). The fate of
organic carbon in marine sediments - New insights from recent data and analysis. *Earth-Science*
Reviews, 204, 103146.

Lefcheck, J.S. (2016). PIECEWISESEM : Piecewise structural equation modelling in R for ecology,
evolution, and systematics. *Methods Ecol Evol*, 7, 573–579.

Middelburg, J.J. (2011). Chemoautotrophy in the ocean. *Geophys. Res. Lett.*, 38, n/a-n/a.

Middelburg, J.J. (2018). Reviews and syntheses: to the bottom of carbon processing at the seafloor.
Biogeosciences, 15, 413–427.

Neumann, A., van Beusekom, J.E.E., Eisele, A., Emeis, K., Friedrich, J., Kröncke, I., *et al.* (2021).
Macrofauna as a major driver of bentho-pelagic exchange in the southern North Sea. *Limnology &*
Oceanography, 66, 2203–2217.

Nielsen, U.N., Ayres, E., Wall, D.H. & Bardgett, R.D. (2011). Soil biodiversity and carbon cycling: a
review and synthesis of studies examining diversity–function relationships. *European J Soil Science*,
62, 105–116.

Oksanen, J. (2010). Vegan: community ecology package. *http://vegan. r-forge. r-project. org/*.

OSPAR. (2017). OSPAR Bottom Fishing Intensity - Surface & Subsurface.

Porz, L., Chen, J., Yilmaz, R., Kuhlmann, J., Zhang, W. & Schrum, C. (2026). Dredging and dumping
impact coastal fluxes of sediment and organic carbon. *Nat Commun*, 17, 216.

Porz, L., Zhang, W., Christiansen, N., Kossack, J., Daewel, U. & Schrum, C. (2024). Quantification and
mitigation of bottom-trawling impacts on sedimentary organic carbon stocks in the North Sea.
Biogeosciences, 21, 2547–2570.

Probandt, D., Knittel, K., Tegetmeyer, H.E., Ahmerkamp, S., Holtappels, M. & Amann, R. (2017).
Permeability shapes bacterial communities in sublittoral surface sediments: Permeability shapes
benthic bacterial communities. *Environ Microbiol*, 19, 1584–1599.

Puig, P., Canals, M., Company, J.B., Martín, J., Amblas, D., Lastras, G., *et al.* (2012). Ploughing the deep
sea floor. *Nature*, 489, 286–289.

Pusceddu, A., Dell'Anno, A., Fabiano, M. & Danovaro, R. (2009). Quantity and bioavailability of
sediment organic matter as signatures of benthic trophic status. *Mar. Ecol. Prog. Ser.*, 375, 41–52.

Queirós, A.M., Birchenough, S.N.R., Bremner, J., Godbold, J.A., Parker, R.E., Romero-Ramirez, A., *et al.*
(2013). A bioturbation classification of E uropean marine infaunal invertebrates. *Ecology and*
Evolution, 3, 3958–3985.

Sala, E., Mayorga, J., Bradley, D., Cabral, R.B., Atwood, T.B., Auber, A., *et al.* (2021). Protecting the
global ocean for biodiversity, food and climate. *Nature*, 592, 397–402.

Sciberras, M., Hiddink, J.G., Jennings, S., Szostek, C.L., Hughes, K.M., Kneafsey, B., *et al.* (2018).
Response of benthic fauna to experimental bottom fishing: A global meta-analysis. *Fish and*
Fisheries, 19, 698–715.

Sharkey, T.D. & Weise, S.E. (2012). Autotrophic Carbon Dioxide Fixation. In: *Photosynthesis*, Advances
in Photosynthesis and Respiration (eds. Eaton-Rye, J.J., Tripathy, B.C. & Sharkey, T.D.). Springer
Netherlands, Dordrecht, pp. 651–674.

Snelgrove, P.V.R., Thrush, S.F., Wall, D.H. & Norkko, A. (2014). Real world biodiversity–ecosystem
functioning: a seafloor perspective. *Trends in Ecology & Evolution*, 29, 398–405.
Tao, F., Huang, Y., Hungate, B.A., Manzoni, S., Frey, S.D., Schmidt, M.W.I., *et al.* (2023). Microbial
carbon use efficiency promotes global soil carbon storage. *Nature*, 618, 981–985.
Teschke, K., Kraan, C., Kloss, P., Andresen, H., Beermann, J., Fiorentino, D., *et al.* (2022). CRITTERBASE,
a science-driven data warehouse for marine biota. *Sci Data*, 9, 483.
Tiano, J., De Borger, E., Paradis, S., Bradshaw, C., Morys, C., Pusceddu, A., *et al.* (2024). Global meta-
analysis of demersal fishing impacts on organic carbon and associated biogeochemistry. *Fish and*
Fisheries, faf.12855.
Tillin, H., Hiddink, J., Jennings, S. & Kaiser, M. (2006). Chronic bottom trawling alters the functional
composition of benthic invertebrate communities on a sea-basin scale. *Mar. Ecol. Prog. Ser.*, 318,
31–45.
Wentworth, C.K. (1922). A scale of grade and class terms for clastic sediments. *The journal of geology*,
30, 377–392.
Wiegand, S., Jogler, M. & Jogler, C. (2018). On the maverick Planctomycetes. *FEMS Microbiology*
Reviews, 42, 739–760.
Wrede, A., Beermann, J., Dannheim, J., Gutow, L. & Brey, T. (2018). Organism functional traits and
ecosystem supporting services – A novel approach to predict bioirrigation. *Ecological Indicators*, 91,
737–743.
Zhang, W., Porz, L., Yilmaz, R., Wallmann, K., Spiegel, T., Neumann, A., *et al.* (2024). Long-term carbon
storage in shelf sea sediments reduced by intensive bottom trawling. *Nat. Geosci.*
Zhu, D., Delgado-Baquerizo, M., Ding, J., Gillings, M.R. & Zhu, Y.-G. (2021). Trophic level drives the host
microbiome of soil invertebrates at a continental scale. *Microbiome*, 9, 189.
689
690
691

694
695
696   **Figure 1.** Geographic overview of all study sites in the southeastern North Sea where macrofaunal and
697   microbial samples were obtained in parallel. The background heatmap displays the bottom trawling
698   intensity calculated as swept area ratio (SAR) per year.
699

Figure 2. Community composition of benthic microbiota **(A-C)** and macrofauna **(D-F)** across 150 stations. **(A, D)** NMDS ordination plots based on **(A)** benthic microbial community composition using Aitchison distances (stress = 0.097), and **(D)** benthic faunal community composition using Bray-Curtis distances (stress = 0.128). Predictor and response variables fitted on the ordination are indicated by arrows (showing only variables with $p < 0.05$). **(B-C, E-F)** Stacked barplots showing the 25 relative most abundant microbial **(B-C)** and faunal **(E-F)** orders of all 150 sites, ordered by their scores on the respective NMDS axes. Border plots on top of each barplot show the log transformed SOM% (green), and trawling intensity in SAR y^{-1} (red).

711
Figure 3. (A) Structural equation model of SOM in response to abiotic and biotic variables. Red and
green arrows indicate significant ($p < 0.05$) negative and positive effects. Black arrows indicate
significant effects associated with changes in composition and dashed gray arrows show hypothesized
relationships which were statistically not supported. Arrows widths are proportional to their
standardized effect sizes. Observed variables are shown in squared boxes and latent variables in
rounded boxes. The marginal and conditional R^2 values of the LMMs are shown in boxes of the
respective response variables, indicating the model variation explained by fixed effects (R^2m), and by
both fixed and random effects (R^2c). **(B)** Absolute direct (blue) and indirect (yellow) standardized
effects on microbial community composition. Displayed values are the sum of the significant ($p < 0.05$)
standardized effects of the same variable on microbial NMDS1 and NMDS2. **(C)** Net direct (blue) and
indirect (yellow) standardized effects of log transformed SOM%.
723

724
Figure 4. Trawling impacts on microbial **(A, B)** and faunal **(C, D)** traits and species identified through
mGLMs. **(A)** Microbial traits (predicted pathways) with significant responses to trawling effort and
effect sizes > 0.1. **(B)** Microbial taxa (OTU numbers in brackets) with significant responses to trawling
effort and effect sizes > 0.1. **(C)** Faunal traits with significant responses to trawling effort. **(D)** Faunal
taxa with significant responses to trawling effort. Colors correspond to MetaCyc functional categories
(A), faunal trats **(C)** phylum-level taxonomy **(B, D)**. Se supplementary 1 for the extended figure
including micorbial traits and taxa with smaller effect sizes.
732

Figure 5. The most important faunal **(A, B)** and microbial **(C, D)** traits and species identified through pGLMs as key predictors of microbial community composition **(A, B)** and SOM% **(C, D)**. **(A)** Pooled effect sizes of faunal traits and covariates retained from penalized regressions fitted on microbial NMDS1 and NMDS2. **(B)** Pooled effect sizes of faunal species and covariates retained from penalized regressions fitted on microbial NMDS1 and NMDS2. Pooled effects were calculated as the square root of the sum of the squared regularized effects of each predictor on NMDS1 and NMDS2. Regularized effect sizes from individual regressions are visualized in heatmaps adjacent to the plots. **(C)** Regularized effect sizes of microbial traits (e.g., pathways) and covariates retained from penalized regressions fitted on log-transformed SOM%. Pathway numbers are labeled within the plot area. **(D)** Regularized effect sizes of microbial OTUs and covariates retained from penalized regressions fitted on log-transformed SOM%. Colors correspond to phylum-level taxonomy.