Morph bias in inflorescences and individual plants reduces opportunities for geitonogamy

in a monomorphic enantiostylous species

Authors: S. B. Rhuthuparna[1] and Vinita Gowda[1*]

[1]Tropical Ecology and Evolution Lab, Department of Biological Sciences, Indian Institute of

Science Education and Research, Bhopal, Madhya Pradesh – 462066, India

[*] Corresponding author: Vinita Gowda

E-mail: gowdav@iiserb.ac.in

ORCiD:

S. B. Rhuthuparna: 0009-0004-4235-5587

Vinita Gowda: 0000-0001-8533-0014

Summary

● Evolution of enantiostyly, a stylar polymorphism characterised by left- or right-
handed flowers, is predicted to counter selfing through disassortative pollen
movement. However, in monomorphic enantiostyly, both morphs are present on the
same individual, and geitonogamy has been proposed to depend on the morph ratio
within an individual.

● Using a monomorphic enantiostylous *Didymocarpus podocarpus* (Gesneriaceae)
endemic to the Eastern Himalayas, we hypothesised that geitonogamous events would
be reduced when plants have a biased morph ratio, due to disassortative pollen
movement. We first established natural morph ratios and reproductive compatibilities
of the morphs. We next examined the role of morph ratios in pollen movement by
recording pollinator visitation patterns using visual observations and quantum dots.

● Our results show that inflorescences and individual plants exhibit biased morph
ratios, while the population maintains an isoplethic ratio. Pollen transfer was higher
between morphs than within morphs, and pollination success in inter-morph
treatments was higher than in intra-morph treatments. Finally, inter-morph switches
by the pollinator within plants decreased with an increase in morph bias.

● We show that biased morph ratios within plants, combined with disassortative pollen
movement, can limit the occurrence of geitonogamous events, thus acting as a
strategy to increase pollination success.

Keywords: Bee pollination, Eastern Himalayas, Gesneriaceae, Reproductive strategies, Self-
pollination, Stylar polymorphism, Quantum dots

Introduction

In hermaphroditic flowers, the spatial arrangement of male and female reproductive structures can affect their reproductive success (Webb & Lloyd, 1986; Richards, 1997; Barrett *et al*., 2000; Barrett, 2002; Li *et al*., 2009; Barrett, 2010). Hermaphroditic plants with inter-mating floral morphs, such as plants with stylar polymorphisms, are predicted to have evolved to counter selfing and to promote outcrossing (Barrett et al., 2000; Barrett & Harder, 2005; Pauw, 2005; Barrett, 2010; De Almeida & de Castro, 2019). In species with stylar polymorphisms, on a population scale, pollination and its outcome depend on the number of flowers of each floral morph that are displayed simultaneously within and between plants (Waites & Ågren, 2004; Brys & Jacquemyn, 2010). For instance, in stylar polymorphism like heterostyly, individual plants display only one type of floral morph. This facilitates pollen movement among individuals carrying dissimilar morphs, that is, inter-morph cross-pollination, thus ensuring higher outcrossing rates within a population (Keller *et al*., 2014). However, in enantiostyly, which represents a distinct type of stylar polymorphism characterised by left (L-morph) or right-handed (R-morph) flowers, the morphs can be present on the same individual or segregated between individuals (Table 1). Enantiostylous flowers are identified by left or right-handed orientation of the style (non-reciprocal enantiostyly) or both style and stamen (reciprocal enantiostyly). The two morphs may be present in two separate individuals, resulting in dimorphic enantiostyly (DE), or within the same individual, as observed in monomorphic enantiostyly (ME). A majority of the ME species are reported to be self-compatible (Table 1; Wang et al., 1995; Gao et al., 2006; Tang & Huang, 2005; Ren et al., 2013; Richman & Venable, 2018; Mora-Carrera et al., 2019; Paudel *et al.,* 2024; Johnson *et al*., 2025) and thus, the presence of both the morphs within an individual can pose a risk of geitonogamous pollen transfer.

Floral handedness in enantiostylous species has been shown to enhance the movement of pollen between flowers of different morphs, resulting in a disassortative pollen movement (i.e., L to R or R to L), thus reducing self-pollen transfer (geitonogamy) when compared with plants that lack enantiostyly (Table 1; Richards, 1997; Barrett *et al.*, 2000; Jesson *et al*., 2003; Jesson & Barrett, 2005). Between DE and ME types of enantiostyly, geitonogamy is lowest in DE species, whereas ME species may exhibit intermediate levels of geitonogamy (Jesson & Barrett, 2005;

Saltini *et al*., 2025). Thus, rates of geitonogamy may vary with the type of enantiostyly, where
the main differentiating factor is the distribution of L and R morphs within an individual.

Among ME species, a population may consist of individual plants with an equal
distribution of L and R flowers (Wang *et al*., 1995; Gao *et al*., 2006; Martins, 2008; Ren *et al*.,
2013; Richman & Venable, 2018; Mora-Carrera *et al*., 2019; Braga *et al*., 2022) or a skewed
proportion of morphs, where individual plants exhibit partial or complete bias towards either one
of the morphs (Dulberger & Ornduff, 1980; Tang & Huang, 2005; De Almeida *et al*., 2018;
Robertson *et al*., 2025). Since ME species bear both L and R morphs on the same individual
plant, both the visitation sequence of pollinators within an individual as well as the relative
abundance of L and R morphs, are predicted to contribute towards geitonogamous pollen transfer
(Table 1; Jesson & Barrett, 2005; Tang & Huang, 2005; Mora-Carrera *et al*., 2019; Barrett &
Fairnie, 2024). However, in ME individuals with an equal number of L and R morphs, that is, an
isoplethic morph ratio, geitonogamous pollen transfer may be inevitable since both active
morph-switches and random switches by pollinators will result in high geitonogamous pollen
transfer. Thus, a reduced opportunity for geitonogamous pollen transfer may be achieved in an
ME species if morph ratios are highly skewed within an individual. A highly skewed morph ratio
within a plant can ensure that even under random pollinator movement within an individual, the
disassortative pollen movement (L to R and R to L) is mostly between individuals.

Enantiostylous forms and their morph ratios	Reproductive compatibility	Expected rate of successful self- and cross-pollination events given the type of enantiostyly and reproductive compatibility.		
		Autogamy	**Geitonogamy**	**Allogamy**
Dimorphic enantiostyly

(Plants have only one type of morph per individual) | Self-compatible (Johnson *et al*., 2023; Jesson & Barrett, 2002) | Exiguous | Exiguous | High |
| | Self-incompatible (Minnaar & Anderson, 2021) | Absent | Absent | High |

		Autogamy	Geitonogamy	Allogamy
Monomorphic enantiostyly (Plants have isoplethic morph ratio)	Self-compatible (Wang *et al*., 1995; Gao *et al*., 2006; Ren *et al*., 2013; Richman & Venable., 2018; Mora-Carrera *et al.,* 2019)	Exiguous	Intermediate[a] to High[a, b]	Intermediate[a]
	Self-incompatible (Morais *et al*., 2020)	Absent	Absent	Intermediate[a]
Monomorphic enantiostyly (Plants have a biased proportion of morphs)	Self-compatible (Tang & Huang, 2005)	Exiguous	Exiguous[b] to low[b]	Intermediate[a] to High
	Self-incompatible (Dulberger & Ornduff, 1980; Robertson et al., 2025)	Absent	Absent	Intermediate[a] to High
Non-polymorphic hermaphroditic species (Included for comparison with enantiostylous floral forms)	Self-compatible (Brys *et al*., 2013)	Intermediate to High	Intermediate to High	Low[a]
	Self-incompatible (Li *et al*., 2013)	Absent	Absent	Intermediate[a] to High

Table 1 The potential for successful self-pollination (autogamy and geitonogamy) and cross-pollination (allogamy) events in different types of enantiostylous plants. Floral arrangement of L morph (L), R morph (R), and hermaphroditic flowers (F) within two inflorescences (1 and 2) and within a plant are illustrated. The pollination

evaluation is based on the morph ratios characteristically known for the specific type of enantiostyly and their

known reproductive compatibility (references included). The superscripts refer to pollination outcomes under two

specific conditions: a—when within-plant pollen loss is expected, b—when floral arrangement can affect pollen

movement within an inflorescence on a plant.

To the best of our knowledge, the effect of the relative abundance of morphs within the

plant on opportunities for geitonogamy in natural populations of ME plants remains to be tested.

Our present research aims to fill this void by investigating the ME species *Didymocarpus*

podocarpus (Gesneriaceae), an understorey, perennial, mostly lithophytic herb restricted to the

eastern Himalayas. We carried out *in situ* studies on two populations of *D. podocarpus* and

tested the hypothesis that geitonogamous events will be reduced when plants have a biased

morph ratio, especially due to disassortative pollen movement between morphs. Though ME has

been reported from multiple genera within Gesneriaceae (Harrison *et al*., 1999; Gao *et al*., 2006;

Ling *et al*., 2020; Prasanna, 2023), enantiostyly remains an understudied topic. The study was

conducted across two populations of *D. podocarpus* to ensure that the morph ratio is not a

population-specific character.

The aims of our investigation were as follows: (1) to quantify monomorphic enantiostyly

by documenting morph ratios in natural populations across multiple years; (2) to perform hand-

pollination experiments to establish autonomous self-pollination, and to check intra- and inter-

morph compatibility; (3) to evaluate pollen transfer patterns by the pollinators in the two morphs,

and (4) to finally integrate morph ratios, reproductive compatibilities of the two morphs, and

pollinator visitation patterns to infer the potential of geitonogamy within a plant. Based on these

objectives, we address the following key questions: (i) Do the morph ratios within a plant and a

population deviate from an isoplethic ratio? (ii) Do self and cross treatments yield similar

reproductive success (fruit set and seed set) in the two morphs? (iii) Do morph ratio and

pollinator visitation patterns within an individual affect opportunities for geitonogamous pollen

transfer?

Materials and methods

Study species and study sites

Didymocarpus podocarpus (Gesneriaceae) is a perennial herb distributed in the Himalayan foothills of Northeast India, Nepal, Bhutan, and China. It is mostly lithophytic (occasionally terrestrial), found on moss-covered rocks. The species flowers between late July and early September, and the flowering within an individual plant lasts for a maximum of 15–20 days. The plants produce 1–9 cymose inflorescences, each bearing ≤ 10 open flowers (mean ± SE number of open flowers is 6.28 ± 0.41). The species exhibits monomorphic, reciprocal enantiostyly with distinct L and R morphs (Fig. 1; Fig. S1). The flowers are tubular, last for five days, and have a nectary at the base of the tube. Stigma receptivity did not differ between the two morphs across their floral longevity, based on the hydrogen peroxide test (Kearns & Inouye, 1993). All experiments were conducted in two natural populations of *D. podocarpus* within India, the Takdah Reserve Forest, West Bengal (27°01'58" N, 88°20'23" E, elevation 1812m), and Gangtok, Sikkim (27°20'30" N, 88°37'16" E, elevation 2001m), during August in the years 2022, 2024, and 2025. We studied a total of 153 plants for the estimation of morph ratio (section 2.2), out of which 72 were used in quantifying pollinator visitation sequence within individual plants (section 2.6). Out of 153, only 109 plants were available to measure fruit set. Since experimentation on this species in shade house conditions is challenging due to both collection restrictions from the wild and survivability rates in the shade house, all studies were carried out in the wild populations. The primary pollinator (98.8% visits) of *D. podocarpus* was identified as *Bombus breviceps* (Apidae; Williams, 2022), and throughout this study, the term '*Bombus* bee' refers only to this species.

Fig. 1 Floral and fruit morphology of enantiostylous *D. podocarpus*. (a) From left to right of the image - L morph

and R morph. Arrows point to the reciprocal placement of anther (A) and stigma (S) on the L and R morphs. (b)

Floral dissection (from left to right) - adaxial surface of calyx, abaxial surface of calyx, abaxial surface of corolla

tube, abaxial surface of corolla tube showing fusion of the anther to the corolla tube, and pistil. (c) mature fruit, and

microscopic image of seeds.

Estimation of morph ratios (MR) to quantify L bias and R bias

In ME, since both morphs can be present on the same inflorescence, an inflorescence can show L

bias, R bias, or an isoplethic ratio. To estimate morph ratio within an inflorescence, within an

individual, and within a population, we sampled 30–33 individual plants at random, across three

years in Takdah, in 2022 (T22), 2024 (T24), and 2025 (T25), and two years in Gangtok in 2024

(G24) and 2025 (G25). For each individual, we counted the number of open L and R morphs on

all the inflorescences. Next, to quantify the relative abundance of the morphs within

inflorescences and individual plants, we calculated the 'morph ratio' (MR) of inflorescences and

individual plants as: MR = the total number of flowers of L morph ÷ the total number of flowers.

Thus, MR represents the relative bias of inflorescences or plants with reference to only the L

morphs and is a metric that retains morph identity (L bias or R bias). That is, when all flowers in

an inflorescence or plant are L morphs, the MR value = 1, and can be identified as L bias. Later,

we introduce another metric, morph bias (MB), which designates the overall morph bias of an

inflorescence or plant, without retaining morph identity. To investigate yearly variation in the

MR of plants, we fitted a Generalised Linear Mixed Model (GLMM) with a binomial

distribution using the lme4 package (Bates et al., 2015) in R (version 4.4.0; R Core Team, 2024).

The year of sampling was treated as a fixed effect, and the population as a random factor. A chi-

square goodness-of-fit test was used to evaluate whether the total number of L and R morph

flowers within populations across years deviated from a 1:1 expectation.

To investigate whether individual plants exhibit L or R biased inflorescences, based on

the MR values, we categorised each inflorescence within a plant into three groups: inflorescences

with (i) complete bias (MR values of 0 for R bias and 1 for L bias); (ii) partial bias (0.01–0.44

for partial R bias and 0.56–0.99 for partial L bias), and (iii) isoplethic ratio (0.45–0.55). The

number of inflorescences in each category of morph ratio within an individual plant was

compared using a GLMM with a negative binomial distribution, where morph ratio was treated

as a fixed effect, and plant identity and population were treated as random effects. The

significance of the individual fixed effects was tested using Wald's χ^2 test (type III) using the car

package (Fox & Weisberg, 2018), and multiple contrast analysis was carried out using the

emmeans package (Lenth, 2025) with Sequential Bonferroni Correction. All statistical analyses

were carried out in R (version 4.4.0; R core team, 2024).

Reproductive compatibility between morphs

To quantify the self and cross-compatibility between the two morphs, we carried out the

following hand-pollination treatments: (i) inter-morph outcrossing (n = 40 flowers from 29

plants), (ii) intra-morph outcrossing (n = 41 flowers from 30 plants), (iii) inter-morph

geitonogamy (n = 42 flowers from 25 plants), (iv) intra-morph geitonogamy (n = 40 flowers

from 25 plants), and (v) unmanipulated, bagged flowers to check for autonomous self-pollination

(n = 52 flowers from 34 plants). Flowers were bagged pre- and post-treatment to avoid pollinator

visitation, emasculated before anther dehiscence, and hand-pollination was carried out within the

first 18 hours post-anthesis. Each treatment was carried out on flowers of different plants, with

no plant receiving more than one treatment. A maximum of two flowers per plant was used for

any given treatment. For outcrossing, mixed pollen from multiple plants at least 3 m apart was

used. Mature fruits, except the ones herbivored, were collected after three weeks.

The seeds of *D. podocarpus* are tiny - *c.* 0.3 × 0.1 mm and are reported to be 'numerous'.

Therefore, seed counting for mature fruits was carried out using ImageJ software (Rueden *et al.*,

2017). All seeds were carefully removed from the capsule and placed in a white weighing boat.

High-resolution images of these weighing boats were taken using a digital camera (Nikon

D3500), and the total number of seeds per capsule was quantified using the 'analyse particle'

function in the ImageJ software following the method mentioned in Ochogavía (2022). This was

carried out for a total of 84 fruits: (i) inter-morph outcrossing (n = 25), (ii) intra-morph

outcrossing (n = 21), (iii) inter-morph geitonogamy (n = 23), (iv) intra-morph geitonogamy (n =

15).

We fitted separate GLMM models to check whether the fruiting success rate and seed

count differed between (i) different hand pollination treatments, (ii) between L and R morphs,

(iii) between outcrossing and selfing treatments, and (iv) between inter-morph and intra-morph

treatments. Fruiting success was analysed using a binomial distribution, whereas seed count was

analysed using a Poisson distribution. Seed count between L and R morphs was compared using

a negative binomial distribution due to overdispersion and poor fit of the Poisson distribution.

Plant identity was used as a random effect in all models. The significance of the individual fixed

effect and multiple contrast analysis was carried out using the method mentioned under the

section 'estimation of morph ratio, and to quantify L and R bias'.

Furthermore, to compare the pollen production between the two morphs, we counted the

pollen grains from mature buds of both morphs. Paired mature buds (one L and one R morph)

were collected from ten individual plants (five each from Takdah and Gangtok). Undehisced

anthers were stored in microcentrifuge tubes containing 1 ml of 70% ethanol. The anthers were

macerated, and pollen grains were counted using a Neubauer chamber following the method

mentioned in Dafni (1992) and Kearns & Inouye (1993). The difference in pollen count was

compared using a paired t-test.

Pollen placement on the pollinator

To examine if the pollen from the two morphs is differentially placed on the *Bombus* bees, we

carried out controlled manipulative experiments using stained pollen grains and euthanised bees.

The pollen grains of the L and R flowers were labelled using heavy-metal-free CuInSexS2-x/ZnS

(core/shell) quantum dots with Zinc Oleate ligands dissolved in hexane (5 mg/mL), following the

method mentioned in Minnaar and Anderson (2019). To distinguish the pollen grains of L and R

morphs, we labelled them using 2 μL of yellow (580 nm) and red (650 nm) quantum dots,

respectively. Experiments were carried out on 10 freshly opened flowers of each morph, and the

individual euthanised *Bombus* bees ($n = 10$) were manually inserted into a labelled L morph

flower, followed by an R morph using sterile forceps, mimicking their natural probing behaviour

(Fig. 5a). To quantify the number of labelled pollen grains deposited on the pollinator, we

categorised the body of the bees into four parts (Fig. 5a): (i) left proximal region (left side of

head and thorax), (ii) left distal region (left side of the abdomen), (iii) right proximal region

(right side of the head and thorax), and (iv) right distal region (right side of the abdomen). The

number of fluorescent pollen grains was counted in these four regions using a 60X hand

microscope with an inbuilt UV light source. We tested for statistical differences in pollen

placement on the pollinator using GLMM with a negative binomial distribution, where the
interaction of morph type and body parts of the bee was the fixed effect, and bee identity was the
random effect. The significance of the fixed effects and multiple contrast analysis was carried
out as mentioned in the section 'Estimation of morph ratios (MR) to quantify L bias and R bias'.

Pollen movement between morphs using quantum dots

To confirm if pollen transfer to stigmas is highest between the two morphs (L to R and R to L) in
natural conditions, we tracked the pollen using quantum dots. We selected 12 plots (eight from
Takdah and four from Gangtok) of 3×2 m with a minimum distance of 1.5 km between plots.
To mimic the natural ratio of morphs within a population in our experimental plots, we
maintained the morph ratio at an isoplethic ratio by manually removing flowers when necessary.
Since *D. podocarpus* has a restricted distribution in its native habitat and the available flowering
density allowed only a single experimental setup, in the interest of the species and its
conservation, we did not set up alternate experiments, such as with variable morph ratios. All the
inflorescences within the plot were bagged a day before the quantum dot application to avoid
pollen loss. To ensure recording of pollen movement in natural conditions, pollen grains of
approximately one-fourth of the total number of L and R flowers were labelled between 06:00
and 07:00 h using yellow and red colored quantum dots. After 8 hours of natural pollination, all
flowers within the plot were collected between 15:00 and 16:00 h and immediately screened for
the number of fluorescent pollen grains deposited on the stigmas using a 60X hand microscope
with an inbuilt UV light source. We tested if inter-morph pollen transfers were higher than intra-
morph pollen transfer by fitting a negative binomial GLMM, with the interaction between morph
type of the donor and recipient as fixed effects, and plant and plot identity as random effects. The
significance of the individual fixed effect was tested, and pairwise comparisons were carried out
using the method mentioned in the section 'Estimation of morph ratios (MR) to quantify L bias
and R bias'.

Quantifying pollinator constancy towards a floral morph

Pollinator preference for a morph and its subsequent foraging behaviour can influence pollen
movement between morphs. To explore if individual pollinators show a preference for one
morph while foraging across multiple plants, which may result in biased morph switches, we

quantified the floral constancy of *Bombus* bees towards a specific morph in a single foraging

bout. We tracked individual *Bombus* bees (n = 50) for up to 10–30 consecutive floral visits, and

recorded their total intra-morph switches (indicating high pollinator constancy towards a morph)

and inter-morph switches (indicating low pollinator constancy towards a morph) during a single

foraging bout. We used a paired-sample t-test to assess whether the frequency of intra-morph

switches significantly differed from that of inter-morph switches.

Effect of morph bias on pollinator movement within the plant

Complementary to the above experiment, we next tested whether pollinator switches within a

plant may be governed by the plant's morph bias. For this, we first measured pollinator switches

within a plant in 72 of the 153 experimental plants (40 in Takdah and 32 in Gangtok). Each plant

was observed for two 25-minute periods from 07:00 to 15:00 h during the peak pollinator

activity time, resulting in a total of 60 hours of observation (2 slots × 25 min × 72 plants). We

recorded 76 *Bombus* bees and their within-plant inter-morph switches (R to L and L to R) and

intra-morph switches (R to R and L to L) during one visitation bout. We next assigned a morph

bias value for each experimental plant, where 'morph bias' (MB) was computed as: MB = 1-

(number of flowers of the rare morph ÷ number of flowers of the abundant morph). Here, MB =

0 would represent individuals with equal proportions of both morphs, while MB =1 would

indicate a complete skewness towards one morph. This metric identifies only if a plant deviates

from an isoplethic morph ratio and does not identify the type of bias, that is, L bias or R bias. In

ME, since both morphs are present on a plant, and all inter-morph switches (L to R and R to L)

will result in self-pollen transfer (geitonogamy), the direction of the switch is not critical. To

examine the effect of morph bias on the frequency of inter-morph switches, we fitted a GLMM

with a binomial distribution, with MB as a fixed effect and plant identity and population as

random variables.

Finally, we quantified the effect of MB on the pollination success of plants by counting

the total number of fruits, normalised by the total number of flowers, per plant, in a total of 109

(out of 153) naturally pollinated plants. To test the effect of morph bias on the fruit set, we used

a binomial GLMM with morph bias as a fixed effect and population as a random variable.

Results

Morph ratio (MR) within populations

We calculated the MR of a total of 226 inflorescences from 153 plants, from two populations
(Table S1). Of the total number of inflorescences, 85.39% and 79.46% exhibited a biased MR
value (MR value outside the isoplethic range of 0.45–0.55) in the Takdah and Gangtok
population, respectively (Fig. 2a). Similarly, of the 153 individuals, 74.19% and 75% of
individual plants exhibited a biased MR value in the Takdah and Gangtok populations,
respectively (Fig. 2b). GLMM analysis revealed that the MR of plants across the years and for
the two populations was not significantly different (χ^2 = 5.39, df = 4, p = 0.24; Fig. S2), and
therefore, in all subsequent analyses, population identity has not been retained.

Among all the inflorescences examined from the two populations (226 inflorescences),
49.11% exhibited a complete bias towards a single morph, whereas 34.32% showed a partial
morph bias, and only 16.57% of inflorescences were noted to display isoplethic morph ratio. The
number of inflorescences across the three categories of morph ratios within individual plants
differed significantly (χ^2 = 46.6, df = 2, p < 0.001). Individual plants exhibited a significantly
higher number of inflorescences with complete bias compared to inflorescences with partial bias
(Estimate = 0.36, z = 2.81, p < 0.005) and isoplethic ratio (Estimate = 1.09, z = 6.8, p < 0.001).
However, within a population, the total number of L and R morph flowers across the years did
not significantly deviate from an isoplethic ratio (T22: χ^2 = 3.47, p = 0.06; T24: χ^2 = 1.38, p =
0.24; T25: χ^2 = 0.04, p = 0.85; G24: χ^2 = 0, p = 1; G25: χ^2 = 0.59, p = 0.44; Table. 2).

Fig. 2 Observed morph ratio distributions in (a) inflorescences and (b) individual plants within two populations of
D. podocarpus. Zero value on the y-axis represents R-bias, and one represents L-bias, while 0.5 represents an
absolute isoplethic ratio. The isoplethic region is shown in grey shade (0.45−0.55). The red open circle represents
the average of all the morph ratios distributed within the population. When the average value (red open circle) is at
0.5, it suggests that the population displays an equal number of L and R morphs (Gangtok population), while a slight
shift towards the lower part of the isoplethic region suggests a slightly higher number of R morphs in the population
(Takdah population).

Reproductive compatibilities of morphs

We carried out a total of 215 hand pollinations, of which 103 sired fruits. The per cent fruiting
success of each pollination treatment was recorded as follows: (i) inter-morph outcrossing: 80%,
(ii) intra-morph outcrossing: 60.9%, (iii) inter-morph geitonogamy: 66.6%, (iv) intra-morph
geitonogamy: 45%, and (v) autonomous self-pollination with zero success rate (Fig. 3a). We
observed a significant effect of the type of pollination treatment (excluding the autonomous self-
pollination) on the fruiting success ($\chi^2 = 8.33$, df = 3, $p < 0.05$), but the pairwise comparisons
with sequential Bonferroni correction showed significant differences in fruit set only between
inter-morph outcrossing and intra-morph geitonogamy (Table S2). The two morphs did not differ
in their fruiting success (Estimate = -0.29, $z = -0.8$, $p = 0.42$). A marginally significant difference
in fruit set was observed when all outcrossing treatments were compared with all selfing
treatments, which included both intra and inter-morph crosses (Estimate = -0.75, $z = -1.8$, $p =$

0.07). However, the fruit set of all inter-morph pollination treatments was significantly higher

than all intra-morph pollination treatments (Estimate = -0.99, *z* = -2.44, *p* = 0.01; Fig. 3a),

irrespective of whether these crosses were carried out within a plant (selfing) or between plants

(outcrossing).

We did not observe a significant effect of the type of pollination treatment on seed count

(χ^2 = 3.15, df = 3, *p* = 0.37; Fig. 3b), and it did not differ between the two morphs (Estimate = -

0.03, *z* = -0.19, *p* = 0.85). No significant difference was observed between all types of

outcrossing and selfing treatments (Estimate = 0.09, *z* = 0.37, *p* = 0.71). However, consistent

with the fruiting success, all inter-morph treatments produced significantly higher seeds

compared to intra-morph treatments (Estimate = -0.5, *z* = -2.11, *p* < 0.05; Fig. 3b). Intra-morph

treatments (both within and between plants) exhibited a 28.33% reduction in fruiting success and

23.3% reduction in mean seed count per fruit compared to that of inter-morph treatments (both

within and between plants). Finally, pollen count was also noted to be different between the

morphs, where flowers of the R morph show significantly higher pollen count when compared to

the L morph (t = -3.18, df = 9, p-value = 0.01; Fig. 4a).

Fig. 3 Results of fruiting success and seed count from four pollination treatments: inter-morph outcrossing (inter-

out), inter-morph geitonogamy (inter-geit), intra-morph outcrossing (intra-out), and intra-morph geitonogamy (intra-

geit). (a) Fruiting success rate showing significant effect between inter-morph treatments and intra-morph

treatments, and (b) mean number of seeds per fruit (± SE) showing significant effect between inter-morph

treatments and intra-morph treatments. * indicates a statistical significance at $p < 0.05$.

Fig. 4 Comparative pollen count in the two morphs and on the pollinator. (a) Comparison of pollen count between L

and R morph flowers (mean ± SE). * indicates a statistical significance at $p < 0.05$. (b) Asymmetrical pollen

deposition was noted on a *B. breviceps* after natural visits to approximately 30 flowers on different *D. podocarpus*

plants. Arrows point to the larger pollen patch on the bee's right side of the thorax, the contact area of the R morph

(R), while the smaller pollen patch on the bee's left side represents the contact area of the L morph (L).

Pollen placement on the pollinator

All *B. breviceps* individuals (n=50) observed during their natural visitation to *D. podocarpus*

probed the flowers using their proboscis by positioning their proximal part (head and thorax) at

the rim of the flower. Since the anther and stigma are situated at the rim of the floral opening, the

anthers of the L and R morphs were observed to come in contact with the left and right proximal

regions of the Bombus bees, respectively (Fig. 4b). This observation was further corroborated by

results from the pollen placement experiment using euthanized bees and pollen stained using

quantum dots which show that the left and right proximal regions of the bees received pollen

from the two morphs differentially (χ^2 = 75.65, df = 3, p < 0.001; Fig. 5b). The following pair-

wise comparison shows that the left proximal region of the *Bombus* bee received a significantly

higher number of L morph pollen grains than the R morph (Estimate = 2.6, z = 4.89, *p* < 0.001),

and the right proximal region received more R morph pollen grains than the L morph (Estimate =

-3.54, z = -6.64, *p* < 0.0001; Fig. 5b). We further found that the R morph deposited significantly

higher labelled pollen grains on the pollinator compared to the L morph (χ^2 = 6.68, df = 1, *p* <

0.01).

Fig. 5 Pollen deposition across different parts of *B. breviceps*. (a) The natural probing position exhibited by *B*

breviceps and the four regions in which pollen deposition was quantified: left proximal (L-Pro), left distal (L-Dist),

right proximal (R-Pro), and right distal (R-Dist). (b) Mean (± SE) number of pollen grains labelled with quantum

dots recovered from the euthanised *B. breviceps* after manual probing. Asterisks indicate statistical significance:

***, $p \leq 0.001$; *, $p \leq 0.05$; ns, non-significant.

Pollen movement using quantum dots

All 12 plots in which pollen tracking experiments were carried out consisted of a total of 582

flowers, in which we found movement of labelled pollen grains in stigmas of 217 flowers

(37.28%; Table S3). Inter-morph pollen transfer identified on the stigmas was significantly

higher than the intra-morph pollen transfer ($\chi^2 = 126.3$, df $= 1$, $p < 0.0001$; Fig. 6a). The

difference in inter-morph pollen transfer was true when the L morph was the recipient flower

(that is, R × L > L × L, donor × recipient; Estimate = -2.35, z = -10.86, $p < 0.0001$) as well as

when the R morph was the recipient flower (that is, L × R > R × R; Estimate = 1.01, z = 4.98, p

< 0.0001). However, irrespective of intra- or inter-morph comparisons, stigmas of the L morph

received significantly higher labelled pollen than the R morph ($\chi^2 = 59.12$, df $= 1$, $p < 0.0001$;

Fig. 6).

Fig. 6 Pollen deposition on the stigmas of the two morphs in Takdah (T) and Gangtok (G) population. (a) Mean ($\pm$

SE) number of inter-morph (black) and intra-morph (grey) quantum dot-labelled pollen grains deposited on the

stigma of L and R morph recipients. *** indicate statistical significance at $p \leq 0{\cdot}001$. (b) A schematic

representation of the percentage of pollen movement recorded within morphs and between morphs of *D.*

podocarpus. Illustrations by Abhishek Thakur.

Pollinator constancy and pollinator movement within a plant

Paired t-test results did not show any significant difference between the number of intra-morph

switches (pollinator constancy on a morph) and inter-morph switches (absence of pollinator

constancy) by *Bombus* bees ($n = 50$) during their single foraging bout ($t = 0.53$, df $= 49$, $p =$

0.6). We noted a significant negative correlation between the frequency of inter-morph switches

and MB (Estimate $= -2.54$, z $= -3.69$, $p < 0.001$; Fig. 7a). Specifically, pollinators exhibited

higher inter-morph switches when the morphs were in equal proportion (isoplethic), and the

frequency of these switches decreased as the morph bias approached one (complete bias, Fig.

7a). Finally, GLMM analysis also identified a significant positive association between the MB

and the fruit set (Estimate $= 0.35$, z $= 3.44$, $p < 0.001$; Fig. 7b). That is, individual plants with a

biased morph ratio exhibited higher fruit sets.

Fig. 7 Relationship between morph bias (MB) and pollinator switches, and MB and fruiting success. **(a)** Correlation

between the morph bias of individual plants and the frequency of inter-morph switches by the pollinator. (b)

Correlation between the morph bias of individual plants and the natural fruit set. The values range from 0

(isoplethic), which represents an equal number of both morphs, to 1, which represents complete skewness towards

either one of the morphs. The grey shaded area along the x-axis represents plants with isoplethic morph ratio.

Statistical significance is indicated in the top right corner.

Discussion

The relative abundance of morphs within an individual plant and its effect on opportunities for selfing (geitonogamy) remain one of the least explored aspects of monomorphic enantiostylous plants. The key findings from our study are that while individuals of the monomorphic enantiostylous *D. podocarpus* produced inflorescences with a biased morph ratio, the overall number of morphs within the population met the isoplethic ratio. Further, we noted inter-morph pollen transfer to be higher than intra-morph pollen transfer and hand-pollination treatments between morphs yielded higher fruit set and seed set than treatments carried out within morphs. Therefore, we propose that the predominant disassortative pollen movement and reduced inter-morph switches by the pollinators within individuals with higher morph bias may play a critical role in reducing self-pollination (geitonogamy) and increasing cross-pollination.

Individuals with biased morph ratios show reduced geitonogamous events

The frequent morph bias observed within inflorescences and the majority of individuals (Table S1) in the two populations of *D. podocarpus* across multiple years suggests ecological factors that favour a biased morph ratio and a strong selective advantage for the underlying developmental process. Similar morph bias has also been reported from other ME species (Dulberger & Ornduff, 1980; Tang & Huang, 2005; De Almeida *et al*., 2018; Robertson *et al*., 2025). Additionally, in *D. podocarpus,* we noted that the morph ratio of individual plants mirrored the most common morph ratio noted among all of its inflorescences, suggesting that individual plants and their inflorescences tend to exhibit a bias towards the same morph (Fig. S3). In ME species, a morph bias within inflorescences and within individual plants can reduce the frequency of geitonogamous selfing events. This is because when there are multiple flowers on an individual plant, interactions specifically between flowers within an inflorescence can affect their reproductive outcome since pollinators tend to visit flowers within the same inflorescence, causing geitonogamy (Harder & Barrett, 1995; Harder *et al*., 2000; Harder *et al*., 2004). Therefore, any inflorescence trait that might influence mating outcomes, such as a biased morph ratio, is likely under a strong selective pressure (Harder *et al*., 2000), as noted in the majority of individuals of the two *D. podocarpus* populations.

Enantiostyly in *D. podocarpus* is unique because it differs from expected floral traits in

enantiostylous flowers, such as the presence of heteranthery, the lack of nectary, and an open

floral design (outward facing; Jesson & Barrett, 2003). Instead, *D. podocarpus* lacks

heteranthery, has nectaries, and exhibits tubular flowers where the stigma and anther are exerted

at the rim of the floral tube (Fig. 1). As expected from the floral morphology, and consistent with

other studies, we found that inter-morph pollen transfers (L to R and R to L) were higher than

intra-morph (L to L and R to R) pollen transfers, confirming disassortative pollen movement

within *D. podocarpus* (Fig. 6; Jesson & Barrett, 2005; De Almeida *et al*., 2013; Minnaar &

Anderson, 2021). Although the inter-morph pollen transfer in ME species can result in

geitonogamous pollination, the extent of geitonogamy will be low in ME species compared to

non-enantiostylous species, where all flowers within a plant are potential mates (Jesson &

Barrett, 2005; Saltini *et al*., 2025). Studies on heterostylous species have shown that, apart from

the arrangement of sex organs within flowers, the floral display of individual plants and the

morph composition of neighbouring plants affect the degree of disassortative pollination and

quality of pollen deposited on the stigma (Waites & Ågren, 2004; Brys & Jacquemyn, 2010).

While these factors are also true for ME species, the morph ratio within individual plants adds an

additional layer of complexity.

Results from our study provide empirical evidence of the theoretical prediction that the

extent of geitonogamy that may occur in ME species depends on a) the relative abundance of L

and R morph flowers within an individual plant, and b) the visitation sequence by the pollinator

(Jesson & Barrett, 2005; Saltini *et al*., 2025). We found that a biased morph ratio within

individual plants limited the occurrence of geitonogamous events, and thus may influence both

the male and female fitnesses by limiting the effect of both pollen and ovule discounting. The

potential outcome of reduced selfing was also reflected in the fruit set, where the fruit set

increased with an increase in the morph bias.

We expected that the predominant pollen movement between flowers of different morphs

might mandate an equal proportion of both morphs in a population. Our results validate this

because, despite the biased morph ratio in the inflorescence and in the individual plants, the total

number of L and R morph flowers within the two populations across multiple years did not

significantly vary from the isoplethic ratio. This result is also consistent with the majority of

previous studies on ME species (Fenster, 1995; Ren *et al*., 2013; Morais *et al*., 2020; Contreras-

Varela *et al*., 2023). This suggests that, while individuals exhibit a biased morph ratio, the

isoplethic ratio at a population level is likely driven by negative frequency-dependent selection

since any deviation from this ratio can lead to lower pollination success (Jesson & Barrett, 2002).

467 Flowers of the same morph show reduced compatibility

Studies examining reproductive compatibilities among morphs have reported that stylar

polymorphisms, like distyly and tristyly, are linked to heteromorphic incompatibility associated

with variations in pollen size and number, size of the stigmatic papillae, and stigma shape

between morphs, potentially leading to intra-morph pollen recognition at the stigma, style, or

ovary, thus reducing fruiting success and seed count (Richards, 1977; Barrett & Cruzan, 1994;

Massinga *et al*., 2005; Costa *et al*., 2017). We noted moderate intra-morph incompatibility in the

form of lower fruit set and lower seed count per fruit in *D. podocarpus,* irrespective of whether

the intra-morph treatment was within or between plants. These observations strongly suggest that

'morph type' is being recognised regardless of whether the pollen is from within the individual

or a different individual. Intra-morph incompatibility has been previously reported in the

dimorphic enantiostylous species *Wachendorfia paniculata* (Ornduff & Dulberger, 1978),

although a subsequent study failed to corroborate this (Jesson & Barrett, 2002). We propose that

the moderate incompatibility between flowers of the same morphs in *D. podocarpus* can act as

an additional mechanism to ensure disassortative pollen movement (Costa *et al*., 2017), where

the fruiting success of pollen transfer between flowers of the same morph is significantly

reduced.

Finally, in our experiments to track pollen movement, despite an equal number of L and

R morphs in our experimental plots, the stigmas of L morphs received significantly higher

numbers of labelled inter-morph pollen compared to stigmas of R morphs (Fig. 6). Similarly, R

morphs deposited significantly higher pollen on the pollinator compared to the L morphs (Fig.

4b; Fig. 5b). In concordance with this, the comparison of pollen count between the two morphs

show that the R morph indeed produces significantly higher pollen compared to the L morph in

D. podocarpus (Fig. 4a). This difference was observed only when the morph was a pollen donor,

and the fruiting success of the two morphs as a female parent was not different in our hand

pollination experiments. These results suggest that the two morphs may have dissimilar sexual

roles due to their differential contribution to the population's overall pollen pool. The R morph in

D. podocarpus appears to display higher male function through higher pollen donation, whereas

the L morph may display higher female function by receiving a higher number of pollen grains.

These patterns loosely share similarities with a few andromonoecious systems, where male

flowers are reported to produce more pollen compared to that of hermaphrodite flowers

(Narbona *et al*., 2005; Dai & Galloway, 2012). While male flowers of andromonoecious plants

are solely males, in the monomorphic enantiostylous *D. podocarpus* R morphs may emphasise

more on male function by producing and exporting more pollen without being exclusively male.

Conclusion

We demonstrate that the morph ratio of inflorescences and in individual plants has a significant

influence on the pollination success in monomorphic enantiostylous species. A biased morph

ratio within inflorescences and individuals, in combination with disassortative pollen movement

and intra-morph incompatibility, acts as a multi-pronged strategy to reduce opportunities for

selfing (geitonogamy). Further, the presence of intra-morph incompatibility and morph-specific

pollen count differences together suggests that there may be differences in sexual roles between

the two morphs that remain to be explored. Monomorphic enantiostylous plants represent one of

the unique innovations in labile reproductive strategies among all organisms, where they do not

represent true and fixed genetic polymorphisms as known from other stylar polymorphisms.

Instead, they represent morphs whose ratios are probably maintained within a population,

although developmentally regulated within a plant. This opens unique insights into how morphs

are distributed within a plant and population in order to optimise high outcrossing rates,

reproductive success, and genetic admixtures.

Acknowledgements

We express our gratitude to the forest departments of Sikkim and West Bengal for the necessary

permits. We would also like to thank the local community, especially Mr Provesh Gurung and

his family, for their hospitality and support. We would like to thank Sanika Goray, Nevil S,

Manila Chingtham, Rohan Dandavate, and Jyotil Dave for their assistance during the fieldwork

and Kirti and Arti for helping with the seed count. We thank Dr Paul Williams, Natural History

Museum, London, and Dr A Rameshkumar, Zoological Survey of India, for helping us identify

the *Bombus* bee specimens. We acknowledge Dr N. S. Prasanna for giving us the details of the

study sites. We thank Abhishek Thakur for the illustrations. We would like to acknowledge the

Science and Engineering Research Board (SERB ECR/2017/001073; POWER Grant

SPG/2021/000793) and DBT-NER (BT/PR24525/NER/95/754/2017) and Ministry of Human

Resource Development (Ministry of Education), for the research fund awarded to VG, IISER

Bhopal for infrastructural support, the University Grant Commission (UGC; 191620076099) for

a fellowship to RSB and the American Society of Plant Taxonomists (ASPT) Graduate Student

Research Grant, International Association for Plant Taxonomist (IAPT) Graduate Student

Research Grant and Society for Tropical Ecology Student Research Grant awarded to RSB for

fieldwork.

Competing interests

533   The authors have no conflict of interest to declare.

Author contributions

535   RSB and VG conceived and designed the study. RSB carried out the field experiments and

536   collected the data. RSB carried out the statistical analyses with inputs from VG. RSB drafted the

537   manuscript, and VG provided conceptual advice and edited the manuscript. Both authors gave

538   final approval for publication.

Data availability

540   The dataset and R-scripts used for this study will be made available through a public repository

541   once the study is published.

Literature cited

Barrett, S. C., & Cruzan, M. B. (1994). Incompatibility in heterostylous plants. In *Genetic

control of self-incompatibility and reproductive development in flowering plants* (pp. 189–219).

Dordrecht: Springer Netherlands.

Barrett, S. C., & Fairnie, A. L. (2024). The neglected floral polymorphism: Mirror-image flowers
emerge from the shadow of heterostyly. *Evolutionary Journal of the Linnean Society, 3*(1),
kzae004. https://doi.org/10.1093/evolinnean/kzae004

Barrett, S. C., & Harder, L. D. (2005). The evolution of polymorphic sexual systems in daffodils
(*Narcissus*). *New Phytologist, 165*(1), 45–53. https://doi.org/10.1111/j.1469-8137.2004.01183.x

Barrett, S. C., Jesson, L. K., & Baker, A. M. (2000). The evolution and function of stylar
polymorphisms in flowering plants. *Annals of Botany, 85*(suppl 1), 253–265.
https://doi.org/10.1006/anbo.1999.1067

Barrett, S. C. (2002). The evolution of plant sexual diversity. *Nature Reviews Genetics, 3*(4),
274–284. https://doi.org/10.1038/nrg776

Barrett, S. C. (2010). Darwin's legacy: The forms, function, and sexual diversity of flowers.
Philosophical Transactions of the Royal Society B: Biological Sciences, 365(1539), 351–368.
https://doi.org/10.1098/rstb.2009.0212

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R. H. B., Singmann, H., … &
Bolker, M. B. (2015). Package *lme4* (Version 12 [1], 2).

Braga, B. L. P., Matias, R., Consolaro, H., Souza, J. T., & Almeida, N. M. (2022). Inaccuracy of
sexual organ position and spatial variation of styles in monomorphic enantiostylous flowers of
Senna rugosa (Fabaceae-Caesalpinioideae). *Flora, 293*, 152112.
https://doi.org/10.1016/j.flora.2022.152112

Brys, R., & Jacquemyn, H. (2010). Floral display size and spatial distribution of potential mates
affect pollen deposition and female reproductive success in distylous *Pulmonaria officinalis*
(Boraginaceae). *Plant Biology, 12*(4), 597–603. https://doi.org/10.1111/j.1438-
8677.2009.00259.x

Brys, R., Geens, B., Beeckman, T., & Jacquemyn, H. (2013). Differences in dichogamy and
herkogamy contribute to higher selfing in contrasting environments in the annual *Blackstonia*
perfoliata (Gentianaceae). *Annals of Botany, 111*(4), 651–661.
https://doi.org/10.1093/aob/mct031

Contreras-Varela, X., García-Franco, J. G., & Aguirre-Jaimes, A. (2023). Floral and reproductive

biology of *Commelina erecta* L. (Commelinaceae) in a coastal dune system on the Gulf of

Mexico. *Flora, 302*, 152259. https://doi.org/10.1016/j.flora.2023.152259

Costa, J., Castro, S., Loureiro, J., & Barrett, S. C. (2017). Experimental insights on the function

of ancillary pollen and stigma polymorphisms in plants with heteromorphic incompatibility.

Evolution, 71(1), 121–134. https://doi.org/10.1111/evo.13082

Dafni, A. (1992). *Pollination ecology: A practical approach*. Oxford University Press.

Dai, C., & Galloway, L. F. (2012). Male flowers are better fathers than hermaphroditic flowers in

andromonoecious *Passiflora incarnata*. *New Phytologist*, *193*(3), 787-796.

https://doi.org/10.1111/j.1469-8137.2011.03966.x

De Almeida, N. M., & de Castro, C. C. (2019). Enantiostyly in angiosperms. In *Asymmetry in*

plants (pp. 319–328). CRC Press.

De Almeida, N. M., De Castro, C. C., de Lima Leite, A. V., Novo, R. R., & Machado, I. C.

(2013). Enantiostyly in *Chamaecrista ramosa* (Fabaceae-Caesalpinioideae): Floral morphology,

pollen transfer dynamics, and breeding system. *Plant Biology, 15*(2), 369–375.

https://doi.org/10.1111/j.1438-8677.2012.00651.x

De Almeida, N. M., Souza, J. T., Oliveira, C. R. S., Bezerra, T. T., Novo, R. R., Siqueira Filho,

J. A., … & Castro, C. C. (2018). Functional dimorphic enantiostyly in monomorphic

enantiostylous species of the subtribe Cassiinae (Fabaceae-Caesalpinioideae). *Plant Biology,*

20(4), 797–801. https://doi.org/10.1111/plb.12718

Dulberger, R., & Ornduff, R. (1980). Floral morphology and reproductive biology of four

species of *Cyanella* (Tecophilaeaceae). *New Phytologist, 86*(1), 45–56.

https://doi.org/10.1111/j.1469-8137.1980.tb00778.x

Fenster, C. B. (1995). Mirror-image flowers and their effect on outcrossing rate in *Chamaecrista*

fasciculata (Leguminosae). *American Journal of Botany, 82*(1), 46–50.

https://doi.org/10.1002/j.1537-2197.1995.tb15647.x

Fox, J., & Weisberg, S. (2018). *An R companion to applied regression.* Sage Publications.

Gao, J. Y., Ren, P. Y., Yang, Z. H., & Li, Q. J. (2006). The pollination ecology of *Paraboea*

rufescens (Gesneriaceae): A buzz-pollinated tropical herb with mirror-image flowers. *Annals of*

Botany, 97(3), 371–376. https://doi.org/10.1093/aob/mcj044

Harder, L. D., & Barrett, S. C. (1995). Mating cost of large floral displays in hermaphrodite

plants. *Nature, 373*(6514), 512–515. https://doi.org/10.1038/373512a0

Harder, L. D., Barrett, S. C., & Cole, W. W. (2000). The mating consequences of sexual

segregation within inflorescences of flowering plants. *Proceedings of the Royal Society B:*

Biological Sciences, 267(1441), 315–320. https://doi.org/10.1098/rspb.2000.1002

Harder, L. D., Jordan, C. Y., Gross, W. E., & Routley, M. B. (2004). Beyond floricentrism: The

pollination function of inflorescences. *Plant Species Biology, 19*(3), 137–148.

https://doi.org/10.1111/j.1442-1984.2004.00110.x

Harrison, C. J., Möller, M., & Cronk, Q. C. B. (1999). Evolution and development of floral

diversity in *Streptocarpus* and *Saintpaulia*. *Annals of Botany, 84*(1), 49–60.

https://doi.org/10.1006/anbo.1999.0887

Jesson, L. K., & Barrett, S. C. (2002). Enantiostyly in *Wachendorfia* (Haemodoraceae): The

influence of reproductive systems on the maintenance of the polymorphism. *American Journal of*

Botany, 89(2), 253–262. https://doi.org/10.3732/ajb.89.2.253

Jesson, L. K., & Barrett, S. C. (2003). The comparative biology of mirror-image flowers.

International Journal of Plant Sciences, *164*(S5), S237-S249.

Jesson, L. K., & Barrett, S. C. (2005). Experimental tests of the function of mirror-image

flowers. *Biological Journal of the Linnean Society, 85*(2), 167–179.

https://doi.org/10.1111/j.1095-8312.2005.00480.x

Jesson, L. K., Barrett, S. C., & Day, T. (2003). A theoretical investigation of the evolution and

maintenance of mirror-image flowers. *The American Naturalist, 161*(6), 916–930.

https://doi.org/10.1086/375176

Johnson, S. D., Midgley, J. J., & Illing, N. (2023). The enantiostylous floral polymorphism of

Barberetta aurea (Haemodoraceae) facilitates wing pollination by syrphid flies. *Annals of*

Botany, *132*(6), 1107-1118. https://doi.org/10.1093/aob/mcad118

Johnson, S. D., Midgley, J. J., & Illing, N. (2025). Functions of heteranthery and enantiostyly for

wing pollination by pollen-collecting bees in *Dilatris ixioides* (Haemodoraceae). *Annals of*

Botany, mcaf189. https://doi.org/10.1093/aob/mcaf189

Kearns, C. A., & Inouye, D. W. (1993). *Techniques for pollination biologists*. University Press

of Colorado.

Keller, B., Thomson, J. D., & Conti, E. (2014). Heterostyly promotes disassortative pollination

and reduces sexual interference in Darwin's primroses: Evidence from experimental studies.

Functional Ecology, 28(6), 1413–1425. https://doi.org/10.1111/1365-2435.12274

Lenth, R. (2025). *emmeans: Estimated marginal means, aka least-squares means* (Version

1.11.1-00001) [R package]. https://rvlenth.github.io/emmeans/

Li, X. X., Zou, Y., Xiao, C. L., Gituru, R. W., Guo, Y. H., & Yang, C. F. (2013). The differential

contributions of herkogamy and dichogamy as mechanisms of avoiding self-interference in four

self-incompatible *Epimedium* species. *Journal of Evolutionary Biology, 26*(9), 1949–1958.

https://doi.org/10.1111/jeb.12193

Li, Y. X., Quan, Q. M., & Sun, G. L. (2009). Effect of floral morphology on fruit set in

Epimedium sagittatum (Berberidaceae). *Plant Systematics and Evolution, 279*, 51–58.

https://doi.org/10.1007/s00606-009-0145-0

Ling, S. J., Lu, T., & Ren, M. X. (2020). A novel type of mirror-image flowers caused by lateral

bending of the floral tube in a bumblebee-pollinated plant. *Flora, 272*, 151694.

https://doi.org/10.1016/j.flora.2020.151694

Martins, D. J. (2008). Pollination observations of the African violet in the Taita Hills, Kenya.

Journal of East African Natural History, 97(1), 33–42. https://doi.org/10.2982/0012-

8317(2008)97[33:POOTAV]2.0.CO;2

Massinga, P. H., Johnson, S. D., & Harder, L. D. (2005). Heteromorphic incompatibility and

efficiency of pollination in two distylous *Pentanisia* species (Rubiaceae). *Annals of Botany,*

95(3), 389–399. https://doi.org/10.1093/aob/mci040

Minnaar, C., & Anderson, B. (2019). Using quantum dots as pollen labels to track the fates of

individual pollen grains. *Methods in Ecology and Evolution, 10*(5), 604–614.

https://doi.org/10.1111/2041-210X.13155

Minnaar, C., & Anderson, B. (2021). A combination of pollen mosaics on pollinators and floral

handedness facilitates the increase of outcross pollen movement. *Current Biology, 31*(14), 3180–

3184. https://doi.org/10.1016/j.cub.2021.04.074

Mora-Carrera, E., Castañeda-Zárate, M., Fornoni, J., Boege, K., & Domínguez, C. A. (2019). On

the adaptive value of monomorphic versus dimorphic enantiostyly in *Solanum rostratum*. *Annals*

of Botany, 123(1), 205–212. https://doi.org/10.1093/aob/mcy162

Morais, J. M., Consolaro, H. N., Bergamini, L. L., & Ferrero, V. (2020). Reproductive biology

and pollinators in two enantiostylous *Qualea* species (Vochysiaceae) in the Brazilian Cerrado.

Plant Biology, 22(5), 872–880. https://doi.org/10.1111/plb.13091

Narbona, E., Ortiz, P. L., & Arista, M. (2005). Dichogamy and sexual dimorphism in floral traits

in the andromonoecious Euphorbia boetica. *Annals of Botany, 95*(5), 779-787.

https://doi.org/10.1093/aob/mci077

Ochogavía, A. C. (2022). Quantifying the reproductive progression of sunflower using FIJI

(Image J). *MethodsX, 9*, 101879. https://doi.org/10.1016/j.mex.2022.101879

Ornduff, R., & Dulberger, R. (1978). Floral enantiomorphy and the reproductive system of

Wachendorfia paniculata (Haemodoraceae). *New Phytologist, 80*(2), 427-434.

https://doi.org/10.1111/j.1469-8137.1978.tb01577.x

Paudel, B. R., Fairnie, A. L., McCarren, S., Anderson, B., Illing, N., & Barrett, S. C. (2024).

Loss of buzz pollination results in chronic pollen limitation in an enantiostylous plant. South

African Journal of Botany, 171, 592-601. https://doi.org/10.1016/j.sajb.2024.06.019

Pauw, A. (2005). Inversostyly: A new stylar polymorphism in an oil-secreting plant, *Hemimeris*

racemosa (Scrophulariaceae). *American Journal of Botany, 92*(11), 1878–1886.

https://doi.org/10.3732/ajb.92.11.1878

Prasanna, N. S. (2023). *Phylogenetic and biogeographic approaches to understand the*

diversification of the Asian genus Didymocarpus Wall. (Gesneriaceae) (Doctoral dissertation).

Indian Institute of Science Education and Research, Bhopal. http://hdl.handle.net/10603/525010

R Core Team. (2025). *R: A language and environment for statistical computing* [Software]. R

Foundation for Statistical Computing. https://www.R-project.org/

Ren, M. X., Zhong, Y. F., & Song, X. Q. (2013). Mirror-image flowers without buzz pollination

in the Asian endemic *Hiptage benghalensis* (Malpighiaceae). *Botanical Journal of the Linnean*

Society, 173(4), 764–774. https://doi.org/10.1111/boj.12101

Richards, A. J. (1997). *Plant breeding systems.* Garland Science.

Richman, S. K., & Venable, D. L. (2018). Aggregate enantiostyly: Floral visitor interactions with

a previously unreported form of floral display. *Journal of Pollination Ecology, 22*, 49–54.

https://doi.org/10.26786/1920-7603(2018)seven

Robertson, C., Xue, H., Saltini, M., Fairnie, A. L., Lang, D., Kerstens, M. H., … & Lenhard, M.

(2025). Spiral phyllotaxis predicts left–right asymmetric growth and style deflection in mirror-

image flowers of *Cyanella alba*. *Nature Communications, 16*(1), 3695.

https://doi.org/10.1038/s41467-025-58803-5

Rueden, C. T., Schindelin, J., Hiner, M. C., DeZonia, B. E., Walter, A. E., Arena, E. T., &

Eliceiri, K. W. (2017). ImageJ2: ImageJ for the next generation of scientific image data. *BMC*

bioinformatics, 18(1), 529. https://doi.org/10.1186/s12859-017-1934-z

Saltini, M., McCarren, S., Illing, N., Barrett, S. C., & Anderson, B. (2025). Selective contexts

favouring asymmetry: A theoretical and empirical study of pollen transfer in mirror-image

flowers. bioRxiv, 2025-09. https://doi.org/10.1101/2025.09.23.677969

Tang, L. L., & Huang, S. Q. (2005). Variation in daily floral display and the potential for

geitonogamous pollination in two monomorphic enantiostylous *Monochoria* species. *Plant*

Systematics and Evolution, 253, 201–207. https://doi.org/10.1007/s00606-005-0309-5

Waites, A. R., & Ågren, J. O. N. (2004). Pollinator visitation, stigmatic pollen loads and among-

population variation in seed set in *Lythrum salicaria*. *Journal of Ecology*, *92*(3), 512-526.

https://doi.org/10.1111/j.0022-0477.2004.00893.x

Wang, G., Miura, R., & Kusanagi, T. (1995). The enantiostyly and the pollination biology of

Monochoria korsakowii (Pontederiaceae). *Acta Phytotaxonomica et Geobotanica*, *46*(1), 55-65.

https://doi.org/10.18942/bunruichiri.KJ00001079078

Williams, P. H. (2022). *The Bumblebees of the Himalaya: An Identification Guide*. Royal

Belgian Institute of Natural Sciences.

Supporting information

Fig. S1 Image of the inflorescence and individual plant of *D. podocarpus*.

Fig. S2 Yearly variation in the morph ratio of inflorescences and individual plants in the two

populations Takdah (T) and Gangtok (G).

Fig. S3 Correlation between the morph ratio of individual plants and their inflorescences.

Table S1 Number of individuals showing complete morph bias, partial morph bias, and

isoplethic ratio across years in all the study population.

Table S2 Results of the pairwise comparison of fruiting success between hand-pollination

treatments.

Table S3 Number of flowers of L and R morphs in experimental plots for pollen tracking on the

stigma.