Molecular architecture of the tumor microenvironment caused by *BRCA1* and *BRCA2* somatic mutations in lung adenocarcinoma

Gaoming Liao[1, 2, †, *], Xinbin Yang[3, †], Qi Liu[2, †], Shufeng Nan[2], Yan Liu[2], Jinwei Li[2], Si Huang[2], Wang Ning[3], Xionghai Qin[4, *], Gang Xu[3, *]

[1]State Key Laboratory of Respiratory Disease, National Clinical Research Center for Respiratory Disease, National Center for Respiratory Medicine, The First Hospital of Guangzhou Medical University, Guangzhou 510000, Guangdong, China.

[2]Guangzhou Medical University, Guangzhou 510000, Guangdong, China.

[3]Department of Thoracic Surgical Oncology, Affiliated Cancer Hospital & Institute of Guangzhou Medical University, Guangzhou 510095, Guangdong, China.

[4]Department of Thoracic Surgery, Harbin Medical University Cancer Hospital, Harbin 150040, Heilongjiang, China.

†Gaoming Liao, Xinbin Yang, and Qi Liu contributed to this work as first author.

*To whom correspondence should be addressed: lgm179496478@163.com (Gaoming Liao), qxh9069@163.com (Xionghai Qin), and xugang1973@126.com (Gang Xu).

ABSTRACT

Objectives: Homologous recombination repair (HRR) deficiency is associated with improved immunotherapy responses in non-small cell lung cancer (NSCLC) patients. The HRR genes *BRCA1/2* are key regulators of DNA repair, yet their impact on the tumor microenvironment (TME) in lung adenocarcinoma (LUAD) remains unclear.

Methods: Using single-cell sequencing and multi-omics data, we characterized *BRCA1/2* mutation-associated transcriptional programs, immune cell composition, and functional alterations in T cells, investigating the molecular and immune architecture of BRCA-mutant LUAD patients.

Results: BRCA1/2 mutations were associated with increased genomic instability and poor prognosis in LUAD patients, but predicted better clinical outcomes following immune checkpoint blockade (ICB) treatment. *BRCA1* mutations correlated with an upregulated type I IFN/IFN-γ signature and CD8+ T cell activation. *BRCA2* mutations were associated with alveolar/stress/inflammatory responses and enhanced MHC-II antigen presentation, linked to CD4+ T cell differentiation. Both alterations coincided with reduced CD28 co-stimulation and CTL activity, hinting of immune evasion. We identified two tissue-resident memory T cell (Trm) subsets as predictors of clinical outcomes and ICB response. *BRCA1* mutations were associated with CD8+ Trm expansion, whereas *BRCA2* mutations linked to tumor CD4+ Trm expansion and peripheral T/NK cell cytotoxicity. Furthermore, a cancer-promoting program activated by *BRCA1* mutation was vulnerable to histone deacetylase inhibitors, which inhibited LUAD tumor growth.

Conclusions: This study provides a preliminary characterization of the BRCA-mutant TME in LUAD patients, revealing distinct transcriptional and immune patterns that highlight differences in *BRCA1/2*-associated molecular architecture and offer a framework for improving therapy efficacy in LUAD.

Keywords: Lung adenocarcinoma, *BRCA1/2* mutations, Tumor microenvironment, Single-cell sequencing, Tissue-resident memory T cells.

1. Introduction

Lung adenocarcinoma (LUAD), the most common histological subtype of non-small cell lung cancer (NSCLC), remains a leading cause of cancer-related mortality worldwide [1]. The genetic landscape of this disease is complex, involving multiple driver mutations such as *EGFR*, *KRAS*, and genomic instability alterations, which collectively contribute to tumor progression and metastasis [2-4]. The tumor microenvironment (TME) plays a critical role in shaping tumor progression, immune evasion, and therapeutic response, particularly in the context of genetic mutations affecting DNA repair mechanisms [5, 6]. Homologous recombination repair (HRR) deficiency, characterized by mutations in key DNA repair genes, has been increasingly recognized as a determinant of tumor genomic instability in breast cancer, ovarian cancers, and lung cancer [7-9].

BReast CAncer gene 1 (*BRCA1*) and gene 2 (*BRCA2*) are crucial mediators of genomic stability, and their mutations often lead to increased DNA damage, heightened immune surveillance, and altered tumor-immune interactions [10, 11]. Previous studies revealed that mutations in *Brca1* and *Brca2* mediate distinct immune microenvironment effects in murine models of breast cancer, and showed that *Brca2* deficiency improved response to immune checkpoint inhibitors [10, 12]. Among the key genetic mutations that drive LUAD, *BRCA1* and *BRCA2* mutations have been increasingly implicated in the pathogenesis and progression of lung cancer [9, 13]. Recent studies suggest that HRR-deficient tumors exhibit distinct immunogenic profiles, influencing immune cell infiltration, antigen presentation, and responses to immune checkpoint blockade (ICB) therapy in NSCLC [14, 15]. *BRAC1* mRNA can predict the efficacy of second-line cisplatin chemotherapy in patients with metastatic NSCLC and has a good predictive value for patients with advanced NSCLC receiving platinum-based chemotherapy [16, 17]. One report showed a case of *BRCA2*-positive lung adenocarcinoma, in which the disease was stable for about 2 years after treatment with olaparib [18]. Despite this, the molecular architecture of the TME in LUAD with *BRCA1* and *BRCA2* mutations remains incompletely understood.

Understanding the molecular mechanisms through which *BRCA1* and *BRCA2*
mutations alter the TME in LUAD is crucial for identifying potential therapeutic
targets and improving treatment strategies. To address these challenges, we leveraged
single-cell RNA sequencing (scRNA-seq) and single-cell T cell receptor sequencing
(scTCR-seq) to comprehensively dissect the *BRCA1/2*-driven TME in LUAD.
scRNA-seq enables transcriptomic profiling at the single-cell level, providing
unparalleled insights into cellular heterogeneity, lineage differentiation, and gene
expression dynamics within tumors [19, 20]. Unlike bulk RNA sequencing, which
averages gene expression across heterogeneous cell populations, scRNA-seq allows
for the identification of distinct malignant and immune cell subsets, as well as their
functional states [21]. Furthermore, scTCR-seq offers a powerful approach for
understanding T cell diversity, clonal expansion, and antigen-specific immune
responses [22].

In this study, we leveraged large-scale genomic and transcriptomic datasets to
investigate the prognostic implications of *BRCA1/2* mutations in LUAD patients
(Nearly 2,000 samples). By integrating scRNA-seq and scTCR-seq, this study
provides a high-resolution characterization of the immune microenvironment and
antigen-specific T cell responses in *BRCA1/2*-mutant LUAD. Our findings reveal how
BRCA1/2 mutations shape the immune landscape, influence CD8+ and CD4+ T cell
populations, and modulate immunotherapy outcomes. These insights offer
comprehensive view of the molecular mechanisms underlying *BRCA1* and *BRCA2*
mutations in LUAD and highlights new opportunities for personalized therapeutic
strategies.

2. Materials and methods

2. Materials and methods

2.1. Human specimens

Patient-derived samples for this study were collected at Affiliated Cancer Hospital &
Institute of Guangzhou Medical University, Guangdong, China. All patients were
treatment-naïve and had invasive lung adenocarcinoma and PD-L1 negative
confirmed based on immunohistochemistry. Fresh tumor and adjacent normal tissues

of patients were collected through surgery. This study was approved by the Ethics

Committee of Affiliated Cancer Hospital and Institute of Guangzhou Medical

University (Approval No. KY-2025003) in compliance with the international

guidelines. All patients have signed informed consent. The samples were divided into

two parts, one for single-cell sequencing and the other for whole exome sequencing.

In addition, we collected and obtained public datasets of multiple cohorts. The

genomic variation, gene expression, and clinical data of LUAD patients (n=564) of

the cancer genome atlas (TCGA) program were downloaded from cBioPortal data

resource (https://www.cbioportal.org/). The genomic variation and survival data of

LUAD in individuals of East Asian ancestry (OncoSG cohort, n=305) [23] also from

cBioPortal. The mutation and survival data of LUAD patients receiving ICB treatment

from SU2C MARK cohort (n=309) [24]. The gene expression, survival, and response

data of NSCLC patients receiving ICB treatment from OAK cohort (n=699) [25].

Tumor mutation burden (TMB), homologous recombination deficiency (HRD) score,

and neoantigen load data for TCGA-LUAD patients were obtained from a previous

study [26]. The HRD score was determined bysumming specific genomic alterations,

including loss of heterozygosity (LOH), large-scale state transitions (LST), and

telomeric allelic imbalances (ntAI).

2.2. DNA extraction and library construction of WES

Genomic DNA were extracted from lung adenocarcinoma tissues and blood using

CretMag[TM] Multi Sample DNA Kit. Genomic DNA were captured using Agilent

SureSelect Human All Exon v6 library following the manufacturer's protocol (Agilent

Technologies, USA). DNA libraries were constructed following the protocols

provided by Illumina. Then these libraries were sequenced on the Illumina sequencing

platform (Illumina NovaSeq X plus) and 150 bp paired-end reads were generated. The

whole exome sequencing was conducted by OE Biotech Co., Ltd. (Shanghai, China).

2.3. Somatic variant identification

The raw reads (fastq) were pre-processed and trimmed with fastp (Version: 0.23.4)

based on default parameters. Clean reads were aligned to the reference human genome (GRCh38) utilizing the BWA (version 0.7.17). The mapped reads were sorted and indexed by using SAMtools (Version 1.5). The GATK (Version 4.0.5.1) was used for recalibration of the base quality score and for single nucleotide polymorphism (SNP) and insertion/deletion (INDEL) realignment, and used for marking duplicate reads, to obtain analysis-ready BAM files. The final BAM files were used as input files for variant calling. Identification of somatic mutations using the GATK Mutect2 method based on matched blood sample and population germline resource from gnomAD (https://gnomad.broadinstitute.org/). ANNOVAR was used to annotated variations during SNP&INDEL calling.

2.4. Single-cell suspension preparation

The solid tissues (including tumor and adjacent tissues) were processed to prepare single-cell suspensions. The tissue samples were cut into small pieces, and the cells were washed twice with precooled RPMI 1640 + 0.04% BSA medium under aseptic conditions. The tissues were cut into small pieces of approximately 0.5 mm^3 using surgical scissors, placed in freshly prepared hydrolase and digested at a constant temperature of 37° C for 60 minutes, mixing every 10 minutes. The tissue was filtered 2 times using a cell sieve and centrifuged (300g) for 5 minutes at 4° C. After resuspending the sediment with an appropriate amount of medium, an equal volume of erythrocyte lysis buffer (MACS, catalog number: 130-094-183) was added. After mixing, the cells were incubated at 4° C for 10 minutes and centrifuged for 5 minutes (300g), and the number of viable cells, as determined by Trypan blue staining, was determined with a hemocytometer. Human fresh peripheral blood was collected in an anticoagulant tube and diluted with PBS at 1:1. Then, the sample was added to the Ficoll separation medium (Merck) and centrifugated 20 minutes (75g). After centrifugation, a white-grayish layer consisting of mononuclear cells will be found on top of the Ficoll-Paque. We carefully collected the mononuclear cells with a Pasteur pipette, transferred them to a new centrifuge tube, and washed them by centrifuging for 10 minutes (300g). The supernatant was discarded and resuspended in

RPMI-1640 supplemented with 0.04% BSA. The number of viable cells, as

determined by Trypan blue staining, was determined with a hemocytometer.

2.5. Single-cell RNA and V(D)J sequencing

The cell concentration of the freshly prepared single cell suspension was adjusted to

700-1200 cell/μl. The cDNA library amplification was performed according to the

operating instructions of 10× Genomics Chromium Next GEM Single Cell 5′ Reagent

Kits v2.0 (PN-1000263). DNA library construction was performed using the 10x

Genomics Chromium Single Cell 5' Kit (PN-1000190) following the manufacturer's

guidelines. Purified cDNA libraries were sequenced using the paired-end 150 mode

on an Illumina platform (Illumina NovaSeq X plus). For T-cell receptor (TCR), T cell

V(D)J enrichment was performed using the 10× Genomics single cell V(D)J

Enrichment Kit (TCR, PN-1000005) according to the human T cell operating

instructions. The TCR library was amplified using the Chromium Single Cell TCR

amplification Kit (TCR, PN-1000252), and the experimental operation was performed

according to the product instructions. The constructed libraries were performed

high-throughput sequencing (Illumina NovaSeq X plus) using the paired-end 150

mode at OE Biotech Co. Ltd (Shanghai, China).

2.6. scRNA-seq and scTCR-seq data processing

Raw read files were processed with Cell Ranger 7.1.0 and mapped to the GRCh38

genome assembly and counted with GRCh38 annotations. Unique molecular identifier

(UMIs) was calculated by using the "cellranger count" function. As a result, a cellular

gene expression matrix containing the number of gene UMIs detected in each cell was

generated. In addition, we took several steps to obtain high-quality data. First, cells

with high mitochondrial gene expression were removed, as dead cells often exhibit

mitochondrial contamination [27]. Specifically, we followed the previous method and

used the median-centered median absolute deviation (MAD)-variance normal

distribution to fit the expression levels of mitochondrial genes, and removed cells

with expression levels significantly higher than expected

(Benjamini-Hochberg-corrected FDR < 0.01) [28]. Second, we removed cells with

less than 500 or more than 6000 detected genes. Third, we used DoubletFinder [29]

and scDblFinder [30] methods to identify and remove potential doublets. The 90th

percentile of the doublet score of DoubletFinder was used as the cutoff value. And the

expected doublet rate is 10% in scDblFinder method. Raw TCR reads were aligned to

the GRCh38 reference genome and consensus TCR annotation was performed using

"cellranger vdj" function (Cell Ranger 7.1.0). TCR annotation was performed using

the default parameters of the 10X cellranger vdj pipeline. The R package scRepertoire

was employed for clonotype assignment and dynamics analysis.

2.7. Normalization, clustering and visualization of scRNA-seq data

The gene expression matrix of the selected high-quality cells was subjected to

downstream analysis using the R package Seurat (v5.0.3) [31]. The main steps include

the following. First, only genes expressed in more than 5 cells were retained. Second,

the raw UMI count matrix was log-normalized and scaled to 10,000 using the

"LogNormalize" function. Third, 2,000 highly variable genes were identified using

the "FindVariableFeatures" function based on the normalized matrix. Fourth, the gene

expression matrix was scaled and centered using the "ScaleData" function. Fifth,

unsupervised clustering was performed by constructing a shared nearest neighbor

(SNN) graph using the "FindNeighbors" function and the Louvain algorithm. The first

30 principal components were considered and the clustering resolution was 1.6. For

uniform manifold approximation and projection (UMAP) visualization, the first two

dimensions of UMAP was calculated using the "RunUMAP" function. Sixth,

differential expression analysis between clusters was performed using the

"FindAllMarkers" function from Seurat (using default parameters but setting min.pct

to 0.5). Use the "FindMarkers" function to find differential expression genes for

specific clusters. Finally, two complementary approaches are used to annotate cell

clusters: (1) canonical markers; (2) markers were in the top rank of differential

expressed genes.

2.8. Signature score calculation at single-cell and bulk level

Signature (module) score at single-cell level was calculated for selected gene-sets

(including pathways and signatures) using "AddModuleScore_UCell" function from

UCell R package. The pathways or signatures involved in this study include: immune

pathways and receptors from ImmPort database (https://immport.org/shared/home),

tissue-specific T cell signatures [32] and known cancer meta-programs [33] from

previous studies, as were as several specific signatures from the molecular signatures

database (MSigDB) human collections. In addition, tumor tissue upregulated and

downregulated gene signatures are the top 60 up-regulated genes and the bottom 60

down-regulated genes, respectively, calculated using the TCGA-LUAD cohort. CD8+

Trm and CD4+ Trm signatures were identified based on "FindMarkers" function at

single-cell level. Signature score at bulk level was calculated using single sample

gene set enrichment analysis (ssGSEA) algorithm in R package GSVA.

2.9. Functional enrichment analysis

Differential expression analysis between malignant cells and normal cells was

performed, and the up-regulated genes of malignant cells were screened according to

the thresholds avg_log2FC≥0.5 and Padj≤0.01. According to the descending sorting

of avg_log2FC, the top 200 genes of malignant cells were selected. The GO BP

functional enrichment analysis was performed using the R package clusterProfiler (the

threshold used were P≤0.001 and Padj≤0.01). Padj calculations were based on

Benjamini-Hochberg correction. The same approach was used for the comparison

between the two types of malignant cells and the comparison between malignant cells

in *BRCA1* and *BRCA2* mutation samples. In addition, we obtained immune signatures

and gene modules of immune cell type from MSigDB hallmarks, ImmPort database,

and previous studies [34, 35]. Gene set enrichment analysis (GESA) was employed to

compare the immune response profile between *BRCA1* and *BRCA2* mutations using

the R package enrichplot.

For all malignant cells, we performed pathway enrichment analysis using

ActivePathways (v.2.0.3) R package for the terms from MSigDB (human collection

C5) based on the gene sets of two meta-programs. The terms were removed once the

number of genes was less than 10 or more than 500. Terms with BH-corrected

significance adjusted P≤0.05 are considered to be significantly enriched by the gene

sets of two meta-programs and will be retained. The significantly enrichment results

were used as input to the EnrichmentMap plugin in cytoscape (v.3.9.1) software to

draw a network diagram.

2.10. Single-cell copy number variation inference

The inferCNV (v.1.19.1) R package was used to distinguish malignant cells by

inferring chromosomal CNVs based on the single-cell expression data. The cells from

CD8+ T cells as normal reference were used to estimate CNVs for the population of

malignant cells. We prepared gene order files containing the chromosomal start and

end positions of each gene from the GRCh38 assembly as input to the

"gene_order_file" parameter in the "CreateInfercnvObject" function. To perform the

infercnv operation, the count matrix and annotation file are input to create the

infercnv object, and then inferCNV is run with cutoff=0.1 in the "run" function.

2.11. Identification of potential drugs targeting BRCA1-mutant genes

This study identified potential drug small molecules based on *BRCA1*

mutation-related cancer-promoting factors. Gene expression profiles of lung

adenocarcinoma cell lines before and after small molecule drug treatment were

obtained from the NIH library of integrated network-based cellular signatures (LINCS)

data resource. The connectivity Map (cMap) analysis was employed to identify

potential drugs associated with molecular perturbations of four *BRCA1* mutation

prognostic risk factors (4-R genes) [36]. To demonstrate the inhibitory effect of

HDAC inhibitors on the growth of lung adenocarcinoma cell lines, the replication

levels at different HDAC inhibitor concentrations were obtained from the cancer cell

line encyclopedia (CCLE) data resource. However, due to limited data, we only

obtained data on two HDAC inhibitors, entinostat and vorinostat, in 43 lung

adenocarcinoma cell lines. Since drug concentrations (C) were divergent, we

categorized them into 4 groups: 0 μM<C≤2 μM ([0-2]), 2 μM <C≤6 μM ([2-6]), 6

μM <C≤10 μM ([6-10]) and vehicle control (C=0 μM). The mean of each group is the

average replication level of the cell line.

2.12. Cell culture and reagents

The human lung cancer cell line A549 (#SCSP-503) and HEK293T (#SCSP-502)

were purchased from the Type Culture Collection of the Chinese Academy of

Sciences, China. Short Tandem Repeat (STR) analyses were performed to

authenticate the identity of each Cell line used in this article. The A549 cells were

cultured in RPMI-1640 medium (#C11875500BT; Gibco). The HEK293T cells were

cultured in Dulbecco's Modified Eagle's medium (#C11995500BT; Gibco). All two

cell lines were supplemented with 10% fetal bovine serum (#S712-012S; Lonsera),

and grown in a humidified 5% CO_2 atmosphere at 37 °C.

2.13. Vectors and lentiviral transfection

All the short hairpin RNAs (shRNA) were cloned into pLKO.1 vector. Target

sequences are as follows:

LDHA#1: GCCACAGATTTACCCGTGGAT,

LDHA#2: GCCAACAACTTGTGTCTCAAT,

S100A10#1: TAAGGAGCCAAATACCTTGCG,

S100A10#2: CATGAAACACAAACGGCAAAT,

GAPDH#1: GTGCGGAGTGTAATCAGTATT,

GAPDH#2: TCAGGTTGTACGGGATCAAAT.

Lentiviral particles were produced by transfecting the shRNA plasmid, packaging

vectors psPAX2 and pMD2.G in HEK293T cells at a ratio of 4:3:1 by JetPRIME In

Vitro DNA Transfection Reagent (#101000046 (114-15); Polyplus). The media was

changed after 6 h. Then 48 h after media change, the media containing the virus were

collected and concentrated. Viral supernatants were centrifuged at 1500 x g for 45 min

and viral pellets were resuspended with DMEM medium. The lentivirus was frozen at

–80 °C for further use. The efficiency of lentiviral shRNA clones was determined by

western blotting.

2.14. Lentiviral infection

302   1×10^5 A549 cells were seeded into each well of 6well plates, at the same time, added

303   with 400 μL of lentivirus and 8 μg/mL polybrene (#28728-55-4; Selleck). Plates were

304   incubated at 37 °C for 12 h. The cells were then passaged and seeded into new culture

305   dishes. 24 hours after cell passage, infected lentiviral cells were selected with

306   Puromycin (1 μg/mL, #P8833-10mg; Sigma-Aldirch) for 3 days.

2.15. Cell proliferation assay

Cell proliferation was measured over a 4-day time course using the CellTiter-Glo®

Luminescent Cell Viability Assay kit (#G9242; Promega). For this assay, 1000 cells

were seeded into each well of a 96-well plate containing 100 μL of medium per well

and cultured in normal conditions. At the time point of detection, plates were cooled

down to room temperature, then 100 μL of the CellTiter-Glo reagent was added to

each well. Mix the contents for 2 minutes on an orbital shaker to induce cell lysis.

Allow the plate to incubate at room temperature for 10 minutes to stabilize the

luminescent signal. Finally, plates were read using a microplate reader (#SLXFA-SN;

Gene Company). All data were normalized to day 1 and presented as mean ± SD.

2.16. Drug treatment

For Cell viability assay, 5000 cells of A549 were seeded into 96-well plates

containing 100 μL of medium per well. After 24 h of culture at 5% CO2 at 37 °C,

cells were treated with a gradient concentration of corresponding agents (Vorinostat,

S1047; Belinostat, S1085; Entinostat, S1053; Pracinostat, S1515; Selleckchem) for 48

h. Cell viability was determined using the CellTiter-Glo® Luminescent Cell Viability

Assay kit (#G9242; Promega) as described above. For detecting the protein level of

the targeted gene induced by drug treatment, 4×10^5 A549 cells were seeded into each

60 mm cell culture dish. After 24 h of culture at 5% CO2 at 37 °C, cells were treated

with 1 and 3 μM Vorinostat or Belinostat for 2 days. Then cells were harvested for

detecting the protein level of the targeted genes.

2.17. Western blotting

In the case of the efficiency of lentiviral shRNA clones and drug treatment studies. Cells were washed in PBS twice and collected by trypsinization and centrifugation (1500 x g for 5 min). Cell pellets were washed in PBS twice, resuspended in RIPA (#P0013B; Beyotime), supplemented with protease inhibitor cocktail (#78440, Invitrogen), incubated on ice and boiled for 10 min at 100°C. Protein lysates were resolved on a 4-12% YoungPAGE™ Precast Gels (#PK0931, GenScript) and transferred to 0.45 μm Immobilon-P PVDF membrane (#IPVH00010, Merck millipore). Membranes were incubated for 1 hr in TBST (#BL602A, Biosharp) containing 5% non-fat dry milk and then incubated for 1 hr at RT or overnight at 4°C with primary antibodies. Detection was achieved using appropriate horseradish peroxidase (HRP)-conjugated secondary antibodies. Finally, the membrane was incubated with an enhanced chemiluminescence ECL (#CW0049M; CWBIO), and the images were obtained by using MiniChemi®580 + Imaging System (MiniChemi®580; SINSAGE). The following antibodies were used: GAPDH Rabbit mAb (#A19056, 1:100,000; Abclonal), S100A10 Rabbit mAb (#A13634, 1:1000; Abclonal), LDHA Rabbit mAb (#A0861, 1:1000; Abclonal), β-Tubulin Rabbit mAb (#A12289, 1:10000; Abclonal) and HRP Goat anti-Rabbit IgG(H+L) (#111-035-003, 1:10000; Jackson).

2.18. RNA isolation and RT-qPCR

Total cellular RNA was isolated using TRIzol Reagent (#15596018; Thermo Fisher Scientific). The RNA concentration was measured using Nanodrop 2000 (Thermo Fisher Scientific, USA). Total RNA was reverse transcribed into cDNA using HiScript III 1st Strand cDNA Synthesis Kit (+gDNA wiper) (#R312-02; Vazyme) according to the manufacturer's instructions. Quantitative reverse transcription PCR was performed per the manufacturer's protocol on the SLAN-96P Real-time PCR system using ChamQ SYBR qPCR Master Mix (#Q311-02; Vazyme) and gene-specific primers. The qPCR primers sequences are as follows:

LDHA-fwd: 5′-ACCGTGTTATTGGAAGCGGT -3′,

LDHA-rev: 5′-CTCCATGTTCCCCAAGGACC -3′,

GAPDH-fwd: 5′-GTCAAGGCTGAGAACGGGAA -3′,

GAPDH-rev: 5′-AAATGAGCCCCAGCCTTCTC -3′,

β-actin-fwd: 5′-CACCATTGGCAATGAGCGGTTC -3′,

β-actin-rev: 5′-AGGTCTTTGCGGATGTCCACGT -3′.

2.19. Statistics and survival analysis

All statistical analysis were conducted using R software (version 4.2.3,

http://www.r-project.org). The two-sided Wilcoxon rank-sum test was employed to

count and compare the differences between the two groups. Kruskal-Wallis test was

used to compare the differences among the three groups. The significance level of

discrete variables was calculated by Chi-squared test. The identification of prognostic

factors associated with *BRCA1* and *BRCA2* mutations was based on the up-regulated

genes of the corresponding mutations using univariate Cox regression analysis.

Survival analysis was performed by the log-rank test and compared using

Kaplan-Meier curves. Statistical significance for pairwise comparisons across the

multiple groups was assessed using the Benjamini-Hochberg false discovery rate

(FDR) correction. For all statistical tests, $*P \leq 0.05$; $**P \leq 0.01$; $***P \leq 0.001$;

$****P \leq 0.0001$.

3. Results

3.1. BRCA mutations are associated with genomic instability and poor prognosis

in lung adenocarcinoma

Using the TCGA-LUAD cohort data, we examined the effect of somatic mutations of

BRCA1 and *BRCA2* on genomic instability. The results showed that the mutations in

both genes could significantly increase the homologous recombination repair

deficiency (HRD) score and present a higher tumor mutation burden (TMB) and

fraction of genome altered (FGA; Fig. 1A and Fig. S1A), indicating elevated genomic

instability in LUAD tumors with *BRCA1/2* mutations. This phenomenon was also

observed in *BRCA1/2* mutated patients with NSCLC (Fig. S1B-C). Furthermore,

BRCA1 and *BRCA2* mutations increased the neoantigen load in LUADs (Fig. 1A),

suggesting that *BRCA1/2*-mutated tumors might be more immunogenic. By analyzing

the proportion of immune subtypes [26], we found that patients with *BRCA1*

mutations had a higher proportion of C1 subtype (wound healing; P=5.80e-04,

Chi-squared test), while patients with *BRCA2* mutations and wild-type had a higher

proportion of C3 subtype (inflammatory; P=6.61e-03, Fig. 1B). *BRCA1*-mutated

patients also showed activated DNA damage repair pathways and exhibited higher

DNA replication and cell cycle activity compared to wild-type patients (Fig. 1C-D),

suggesting that *BRCA1* mutations may promote tumor growth through compensatory

DNA repair activation—a pattern that may reflect partial HRD rather than complete

loss of function.

Next, we explored the impact of *BRCA1/2* mutations on clinical outcomes of

LUAD patients and found that these mutations were associated with worse prognosis

for disease-free survival (DFS, P=8.46e-03, Log-rank test; Fig. 1E) and overall

survival (OS, P=0.03; Fig. 1F) in multiple LUAD cohorts. In particular, *BRCA1* and

BRCA2 mutations as risk factors shortened the survival of LUAD patients (P=0.028;

Fig. 1E). Interestingly, LUAD patients with BRCA mutations had significantly longer

progression-free survival after ICB treatment (Fig. 1G). Through transcriptional

analysis, we found that upregulation of both *BRCA1* and *BRCA2* was associated with

worse prognosis after ICB treatment (P<0.05, Fig. 1H), consistent with *BRCA1/2*

expression as a risk factor for lung cancers [9]. The apparent paradox between

favorable ICB outcomes linked to *BRCA1/2* mutations and poorer outcomes

associated with high *BRCA1/2* expression may be explained by distinct biological

roles—where mutations induce genomic instability and immunogenicity, while high

expression reflects proficient DNA repair and tumor cell fitness. These results suggest

that BRCA abnormalities may have potential to predict ICB treatment outcomes. In

addition, we found the inconsistencies in lymphocyte infiltration between *BRCA1* and

BRCA2 mutations, suggesting heterogeneity in the immune microenvironment (Fig.

1I).

3.2. Single-cell landscape of BRCA mutations in LUAD patients

Exome sequencing of this study showed that the two types of tumor tissues carried non-synonymous mutations of driver genes *TP53*, *EGFR*, and *FAT1* (Table S1), and had similar mutation frequencies in oncogenic signaling pathways, including p53 and Notch pathways, as well as significant mutation gene set (MutSig) scrutinized by MutSigCV [37] (Fig. 1J). To preliminarily explore the impact of *BRCA1/2* mutations on the immune microenvironment, scRNA-seq was performed on LUAD patients carrying *BRCA1* or *BRCA12* somatic mutations (a total of 6 samples, including 2 tumors, 2 adjacent normal tissues, and 2 peripheral blood samples from one *BRCA1*-mutant and one *BRCA2*-mutant patient). A total of 69,180 high-quality cells were obtained after QC, of which 27,851 (40.3%) were from tumor tissues, 25,035 (36.2%) were from adjacent normal tissues, and 16,294 (23.5%) were from blood (Fig. 2A and Fig. S2A-B). Of these cells, 35,450 cells (51.2%) and 33,730 cells (48.8%) were from the *BRCA1*- and *BRCA2*-mutant samples, respectively. After batch correction (Harmony), we performed cell clustering and cataloged all cells into 19 main cell types annotated with canonical marker genes (Fig. 2A-C and Fig. S2A-C), identifying 4 epithelial compartments (alveolar type 1 [AT1], alveolar type 2 [AT2], ciliated, and club cells), 2 stromal compartments (fibroblasts and endothelial cells), 5 lymphocyte compartments (CD4+ T, CD8+ T, natural killer-NK/NKT, B cells, and plasma) and 6 myeloid compartments (macrophages, dendritic cells-DCs, classical/nonclassical monocytes, neutrophil, and mast cells). Among immune cells, the most abundant cell types were T lymphocytes and macrophages (Fig. 2B).

To distinguish malignant cells, we first inferred copy number variation (CNV) of all epithelial cell types in solid tissues using the inferCNV pipeline [38] (Fig. S2D-E). Using immune cells as a control and comparing the CNV scores of each epithelial cell subset, we identified two malignant cell subsets that were significantly clustered together using the K-means clustering algorithm and exhibited higher scores of oncogenic CNV in LUAD (Fig. 2D-E). For example, both chromosome 1q21 region and chromosome 7—frequently amplified in LUAD tumors [39, 40]—showed

significantly higher CNV scores in the malignant cells of this study (Fig. 2D-E). The previously reported loss of chromosome 3q [41] was also observed in malignant cells (Fig. 2D-E). To demonstrate the properties of malignant cells, we used bulk RNA-seq data from the TCGA-LUAD cohort to identify tumor upregulated and downregulated gene signatures, and calculated module scores in different epithelial cell types. The malignant cell subsets had significantly higher tumor upregulated signature scores and lower downregulated signature scores (Fig. 2F-G and Fig. S2F). In addition, tumor marker genes, such as *CEACAM6*, *CLDN4*, and *WFDC2*, showed higher expression in malignant cells (Fig. 2H and Fig. S2C).

3.3. Distinct malignant transcriptional programs associated with *BRCA1* and *BRCA2* mutations in LUAD

We have identified two subsets of malignant cells (13,694 cells in total) from the *BRCA1/2*-mutant samples. We then asked which key molecules were specifically activated in malignant cell subsets and whether there were differences in the oncogenic programs between the *BRCA1*- and *BRCA2*-mutant cases. To address this, we first performed differential expression analysis comparing malignant cell subsets versus other epithelial cells. The results showed that tumor markers *WFDC2* and *CLDN4* were significantly upregulated in the malignant1 subset, while *CEACAM6* was significantly upregulated in both malignant cell types (Fig. S3A and Table S2). Previous studies show that *DUSP1* could promote angiogenesis, invasion, and metastasis in NSCLC, and *FTH1* expression protects cells from cell death induced by the GPX4 inhibitor RSL3 [42, 43]. These genes upregulated in malignant cells are often also highly expressed in tumor tissues relative to paracancerous tissue (Fig. S3B). The upregulated genes in malignant1 subset were mainly enriched in ribonucleotide biosynthetic and nucleoside-related metabolic process (Fig. 3A and Table S3), suggesting enhanced DNA replication and growth abilities. Differently, the upregulated genes of malignant2 were mainly enriched in cell adhesion, antigen processing and presentation of peptide antigen, and MHC protein complex assembly (Fig. 3A and Table S3), suggesting involvement in lymphocyte recruitment. These

patterns were also observed in direct comparison between the two malignant subsets

(Fig. S4A and Table S4-5).

We then applied nonnegative matrix factorization (NMF) algorithm to identify

transcriptional programs consisting of co-expressed genes across all malignant cells.

Using this method, we extracted a total of 4 programs and identified two

meta-programs (MP1, MP2) based on program similarities (Fig. 3B). The MP1

module contains 156 genes (Table S6), whose activity was significantly higher in

BRCA2-mutant tumor cells (Fig. 3C) and was associated with better prognosis in

LUAD patients (P=0.031, Fig. 3D). The favorable prognosis linked to MP1 was

related to its involvement in lymphocyte migration, T-cell chemotaxis, and myeloid

leukocyte differentiation (Fig. 3E and Table S7). The MP2 module contains 200 genes

involved in functions such as ATP synthesis/metabolic process and mitochondrial

translation, and its activity was higher in *BRCA1*-mutant tumors (Fig. 3C, 3E, Table

S6-7). To further contextualize MP1 and MP2, we obtained 41 known cancer MPs

defined by Gavish et al.[33]. Among these, 15 MPs irrelevant to LUAD were

excluded and 26 MPs were included. Comparing our MPs with known MPs, we found

that MP1 was mainly enriched in alveolar, stress and secretion-related MPs, while

MP2 overlapped with DNA damage cell cycle (HMG-rich) and MYC-related MPs

(Fig. 3F). Furthermore, *BRCA1* mutations were associated with hypoxia and

oncogenic MYC-related MPs, while *BRCA2* mutations were linked to alveolar, stress

and secretion-related MPs (Fig. 3G and Fig. S4B). Comparing key molecules, *BRCA1*

mutation was associated with upregulation of type I IFN genes (e.g., *IFI27, IFITM3*),

angiogenesis genes (*S100A4*, *S100A10*) and *LDHA* (Fig. 3H, Fig. S4C, and Table S8).

BRCA2 mutation was associated with upregulation of stress-related genes

(e.g., *GADD45B*, *DUSP1*, and *SOD2*), immune cell-related genes (*NFKBIA*, *ICAM1*),

inflammatory-related genes (*TNFAIP3*, *CEBPB*, and *CXCL2*), and apoptosis gene

CFLAR (Fig. 3H, Fig. S4C, and Table S8).

3.4. Inferred transcriptional trajectories from epithelial to malignant cells in

***BRCA1*- and *BRCA2*-mutant tissues**

Tumor evolution is a process of gradual accumulation of somatic mutations from

normal cells [44]. To infer potential transcriptional transitions during malignant

transformation, we analyzed the transcriptional trajectories using Monocle 2 [45],

revealing a dynamic transitional spectrum from AT1/2 cells to malignant cells (Fig.

4A and Fig. S5A). The early pseudotime was dominated by AT1 and AT2 cells,

primarily from tumor-adjacent and *BRCA2*-mutant tumor tissues (Fig. 4A). The

trajectory then extended into two branches: one dominated by malignant1 subset (87.6%

from *BRCA1*-mutant tumor), and the other almost all malignant2 subset, allowing us

to explore expression gradients associated with *BRCA1* and *BRCA2* mutations (Fig.

4A). Branch fate determination gene analysis showed that the malignant1 branch

upregulated genes related to oxidative phosphorylation, ATP synthesis and

metabolism, and ROS response (Fig. S5B). The malignant2 branch upregulated

immune-related genes such as *IL1B*, *TNFAIP3*, and *CD74*, enriched in

cytokine-mediated signaling, antigen processing and presentation, and T cell

activation and differentiation (Fig. S5B). This suggests heterogeneous transcriptional

fates during the transition from normal lung epithelial cells to malignant cells in these

two cases. We further investigated the expression differences along the inferred

trajectories. The *BRCA1* mutation branch showed activation of innate immune

response-related pathways such as type I IFN production, response to IFN-α/β, and

upregulated genes related to T cell homeostasis/migration and T cell receptor

signaling (Fig. 4B), suggesting an association with innate immune response and T cell

activation. The *BRCA2* mutation branch showed activation of cell growth and

development, response to wounding, and wound healing-related functions, along with

promotion of MHC protein complex assembly and lymphocyte-mediated immunity

(Fig. 4B), suggesting that *BRCA2* mutation may be associated with immune

engagement alongside tumor growth.

Based on pseudotime order, divided trajectories into 10 bins and analyze the

activity changes of related features. The module scores of tumor downregulated genes

gradually decreased along pseudotime (Fig. 4C), while tumor upregulated genes

increased (Fig. S5C). Cancer MPs, such as cell cycle G2/M (MP1) and stress (MP5) increased along pseudotime in both *BRCA1*- and *BRCA2*-mutant tissues (Fig. 4D-E). Density analysis suggested that stress activation was more prominent in *BRCA2*-mutant tumor cells, while cell cycle HMG-rich program was mainly activated in the *BRCA1*-mutant group (Fig. 4F). Differences in expression programs between *BRCA1* and *BRCA2* mutations were mainly apparent in mid-to-late pseudotime. For example, *BRCA1* mutation-associated fate-determination functions such as response to IFN-α and type I IFN showed higher activity in mid-to-late pseudotime bins (Fig. 4G-H and Fig. S5D). This pattern was also observed in cGAS-STING innate immune sensing pathways, such as cGAS and IFN-γ response (Fig. 4H and Fig. S5E). In contrast, TGF-β signaling and stress modules were more active in late pseudotime in *BRCA2*-mutant cells (Fig. 4I). MHC class I and II molecules were more active in late pseudotime in *BRCA1*- and *BRCA2*-mutant cells, respectively (Fig. 4G-I), a pattern also reflected in cell density analysis (Fig. 4J). CD8+ Tcm and Th1 signatures showed higher activity in late pseudotime in *BRCA1*- and *BRCA2*-mutant cells, respectively (Fig. S5F-G), suggesting differential association with CD8+ vs. CD4+ T cell engagement.

3.5. *BRCA1* and *BRCA2* mutations reshape distinct immune microenvironment in LUAD tumors

We next explored how *BRCA1/2* mutations reshape tumor immune microenvironment of. *BRCA2* mutation-upregulated genes were enriched in MHC class II-related antigen presentation and lymphocyte-mediated immunity functions (Fig. 5A). GSEA showed that MHC class II molecules were upregulated in *BRCA2* mutation group, while MHC class I molecules were upregulated in *BRCA1* mutation group (Fig. 5B-C), consistent with cell density distribution (Fig. 4J). Similarly, response to IFN-α/γ and type I IFN-related genes, as well as TNF-α family members and receptors, were enriched in *BRCA1*-mutant tumor cells (Fig. 5B-C and Fig. S6A-B). Similar patterns were observed for DNA repair, fatty acid metabolism, and CD8+ T cell signatures (Fig. 5B, 5D, Fig. S6A). DNA double-strand breaks (DSBs) can lead to release of cytoplasmic

dsDNA, which can be sensed by cGAS-STING signaling to trigger innate immune

responses and type I IFN production, promoting anti-tumor immunity [46-48]. Based

on these observations, we hypothesized that *BRCA1* mutation triggers the activation

of cGAS-STING signaling and CD8+ T cell infiltration. Indeed, *BRCA1* mutations

were upregulated DNA damage-related checkpoints, significantly activated

DSBs-related DNA repair pathways (e.g., DNA double-strand break repair,

homology-directed repair, and HRR), and had higher activities of cGAS-STING

signaling and STING mediated induction of host immune responses compared to

BRCA2 mutations (Fig. 5E-G and Fig. S6C-F). *cGAS*, *STING1*, and downstream

factors *STAT1* and *CCL5* were upregulated in *BRCA1*-mutant tumor cells (Fig. 5H).

Immune activity analysis suggested that *BRCA1* mutation associated with enhanced

activity of CD8+ T cells and myeloid cells (Fig. 5I), consistent with GSEA results.

Notably, *BRCA1* mutation increased infiltration of CD8+ T cells was also confirmed

in lymphocyte subsets (Fig. 5J).

Unlike *BRCA1* mutation, upregulated genes in *BRCA2*-mutant tumor cells were

enriched in inflammatory response, interleukins receptor, and chemokines (Fig. 5B

and Fig. S6A), which was confirmed by cell activity analysis. For example, the

IL2/IL17 pathway, 4-1BB pathway, and cytokine pathway were more active in

BRCA2-mutant tumor cells (Fig. 5E and Fig. S6C), consistent with the higher

proportion of inflammatory subtypes in *BRCA2*-mutant patients (Fig. 1B). In addition,

BRCA2 mutation was also associated with activation and differentiation of CD4+ T

cells (Fig. 5B and 5J). For example, *BRCA2*-upregulated genes were enriched in

CD4+ Tem signature and showed higher activity of Th1/Th2 pathways, Th1 toxicity

and MHC class II modules (Fig. 5B-E and Fig. S6C). Indeed, immune activity

analysis also suggested that *BRCA2*-mutant tumor cells were more associated with

activation of Th1, Th2, and Treg cells compared to *BRCA1* mutations (Fig. 5I).

Together, these observations suggested that *BRCA1* mutations are associated with

cGAS-STING-mediated host immune responses and CD8+ T cell infiltration, while

BRCA2 mutations are linked to MHC II molecule-mediated antigen presentation,

inflammatory responses and CD4+ T cell differentiation.

3.6. *BRCA1* and *BRCA2* mutations facilitate the clonal expansion of tissue-resident memory T cells

To examine the effect of *BRCA1/2* mutations on T lymphocytes, we obtained T cell data from BRCA wild-type in LUAD as controls from a previous study (LME) [49] and combined with our T cell data for integrated analysis. Nine T cell subsets (67,823 cells) were identified (Fig. 6A and Fig. S7A-C). Naïve T cells were mainly distributed in blood, while Teff cells were more abundant in blood and paracancerous tissues than in tumors (Fig. 6B). Treg cells were decreased in *BRCA1/2* mutant tumors compared to wild-type (Fig. S7D). We identified two tissue-resident memory T cell (Trm) subsets, CD8+ Trm and CD4+ Trm, both predominantly derived from tumor tissues (Fig. 6B). CD8+ Trm cells were more abundant in *BRCA1*-mutant sample, whereas CD4+ Trm cells were higher in *BRCA2*-mutant sample (Fig. 6B and Fig. S7D). CD8+ Trm cells highly expressed TNF family members and showed high activity for interferons and chemokines, similar to Teff and NK/NKT cells, whereas CD4+ Trm cells upregulated chemokine receptors and cytokines receptors (Fig. S7E). Trm score of both ubsets could predict the overall survival of NSCLC patients after ICB treatment ($P<0.01$, Fig. S7G-H) and were associated with ICB response ($P<0.05$, Fig. S7I), suggesting Trm cells as potential biomarkers for ICB treatment in lung cancer.

Consistent with malignant cell results, *BRCA1* mutation was associated with upregulation of TNF family members in T lymphocytes compared to *BRCA2* mutation and wild type (Fig. S7E-F and Fig. S8). Chemokine receptor and TGF-β were more active in *BRCA2*-mutant tumor tissues (Fig. 6C-D, Fig. S7J, and Fig. S8A). *BRCA1/2* mutations were associated with enhanced antigen presentation and cytokine signaling compared to wild type (Fig. S7K and Fig. S8A). Using tissue-specific T-cell signatures [32], we calculated activity at the single-cell level (Fig. 6E and Fig. S9A), and found that *BRCA1/2* mutations were associated with reduced CD28 family co-stimulation and decreased cytotoxic T lymphocytes (CTLs) and NKT cell infiltration compared to wild type (Fig. 6E-F), which explains why patients with

BRCA1/2 mutations showed worse prognosis (Fig. 1E-F). *BRCA1* mutations were associated with reduced TIL dysfunction and exhaustion dysfunction in tumors (Fig. 6E-F), consistent with lower expression of T cell exhaustion markers (Fig. 6G). In blood samples, *BRCA1* mutations were associated with enhanced TCR and its downstream signaling, while *BRCA2* mutations were linked to enhanced T and NK-mediated cytotoxicity and NK cell activation receptors (Fig. 6E).

Furthermore, we performed scTCR-seq to analyze the TCR clonal expansion. Teff, CD8+ Trm, and CD4+Trm showed the highest hyperexpanded clonal types (Fig. 6H-I and Fig. S9B-C), suggesting antigen-driven activation. *BRCA1*-mutant samples had a higher proportion of expanded TCR clonotypes compared to *BRCA2*, across sample types (Fig. 6J and Fig. S9D). In addition, *BRCA1* mutations were associated with higher TCR diversity in tumor and blood, while *BRCA2* mutations increased TCR richness in blood (Fig. S9E). In *BRCA1*-mutant patient, shared clonotypes tended to expand in paracancerous tissue and were more shared between blood and tumor tissues (Fig. 6K and Fig. S9F-G). However, in *BRCA2*-mutant patient, shared clonotypes tended to expand in blood and paracancerous tissue (Fig. 6K and Fig. S9F). *BRCA1* mutations were associated expansion of Teff, CD8+ Trm, and Tem in solid tissues, while CD4+ Trm expansion was more prominent in *BRCA2* mutations (Fig. 6L and Fig. S9H). This was especially evident among the top 50 expanded TCR clonotypes (Fig. 6M). Teff was the most expansion subset in blood, regardless of mutation type (Fig. 6M and Fig. S9I). Together, these results suggest that *BRCA1* mutations are associated with increase antigen exposure and CD8+ Trm expansion, while *BRCA2* mutations are linked to CD4+ Trm expansion in tumor tissue and enhanced T/NK toxicity in blood.

3.7. Targeting a *BRCA1* mutation-associated transcriptional program inhibits LUAD tumor growth in vitro

Our analyses suggested that *BRCA1/2* mutations are associated with distinct transcriptional programs. Then, we identified BRCA mutation-related prognostic factors using Cox regression analysis. The factors associated with *BRCA2* mutation

were linked to longer patient survival (HR<1, Table S9), and a *BRCA2* signature score

predicted better survival (P=0.0015, Fig. S10A), consistent with upregulation of MHC

complex assembly genes. This score can also predicted better survival in

TCGA-LUAD (P<0.0001; Fig. S10B and Fig. S11A) and OAK (P<0.0001; Fig.

S10C-D) cohorts. In contrast, a *BRCA1* mutation prognostic factor was associated

with shorter survival (HR>1, Table S9), and its signature score predicted worse

survival in TCGA-LUAD (P=0.0012; Fig. 7A and Fig. S11B) and OAK (P=0.0013;

Fig. 7B and Fig. S10D, suggesting that *BRCA1* mutation activates a tumor-promoting

program. Based on this, we identified four *BRCA1* mutation-associated risk genes,

S100A10, *LDHA*, *MYL12A,* and *GAPDH* (4-R genes) for subsequent analysis.

Although the expression of 4-R genes was heterogeneous in LUAD patients, it was

associated with worse overall survival (Fig. 7C and Fig. S10E-F). Previous studies

have shown that *S100A10* can promote cancer metastasis by recruiting MDSC cells,

and increased *LDHA* activity contributes to tumor immune escape [50, 51].

Knockdown of *S100A10*, *LDHA*, and *GAPDH* reduced LUAD cell proliferation *in*

vitro (Fig. 7D-E).

We next asked whether small molecules could reduce the expression of these risk

genes and inhibit LUAD growth. Using gene expression profiles of A549 cells before

and after drug treatment from LINCS data resource, we performed cMap analysis to

identify drugs that perturb 4-R gene expression. Interestingly, histone deacetylase

(HDAC) inhibitors consistently downregulated 4-R genes in A549 cells (Fig. 7F and

Table S10). Cell growth decreased significantly with increasing HDAC inhibitor

concentrations (Fig. 7 G-H, Fig. S10G-H, and Table S11). Vorinostat (FDA-approved

for cutaneous T cell lymphoma) significantly inhibited 4-R gene expression (NCS<-1,

P<0.01; Fig. 7F and Table S10). Furthermore, vorinostat and belinostat reduced

expression of *LDHA*, *S100A10*, and *GAPDH* in A549 cells in a dose-dependent

manner (Fig. 7I-J and Fig. S10I). Other HDAC inhibitors (mocetinostat, entinostat,

and SB-939) also showed similar effects on the perturbation of 4-R genes (Fig. 7F-H

and Table S10). These findings suggest that HDAC inhibitors may target *BRCA1*

mutation-associated risk genes and inhibit LUAD tumor growth *in vitro*, offering a
preliminary therapeutic hypothesis for further testing.

4. Discussion

Our study provides a comprehensive molecular and cellular characterization of the
TME in LUAD with *BRCA1* and *BRCA2* mutations, using large-scale multi-omics
data and single-cell data. We describe transcriptional and immune patterns associated
with these mutations, which may contribute to genomic instability, immune cell
infiltration, and immune modulation within the TME. These observations offer
hypotheses for understanding HRR deficiency in LUAD and for developing potential
therapeutic strategies, including ICB and targeted therapies. HRR deficiency is a key
contributor to tumorigenesis and treatment response in multiple cancers [7]. Although
HRR deficiency has been correlated with enhanced sensitivity to DNA-damaging
agents and ICB in cancers such as breast, ovarian, and NSCLC, our study suggests
that in LUAD, *BRCA1/2* mutations may be associated with distinct TME features that
could influence immunotherapy effectiveness. This underscores the complexity of
BRCA-driven biology in LUAD.

A key aspect of this study is the single-cell transcriptomic profiling of
BRCA1/2-mutant LUAD tumors, paracancerous tissues, and peripheral blood patients.
This enabled high-resolution examination of immune signatures, though the sample
size limits generalizability. *BRCA1* mutations were associated with the upregulation
of cGAS-STING and type I IFN response gene, which can enhance innate immune
responses, antigen presentation, and cytotoxic T-cell activity [52]. *BRCA1*-mutant
tumors showed higher CD8+ T cell activation and MHC-I expression, possibly due to
increased neoantigen load. This suggests that *BRCA1*-mutant tumors may be more
immunogenic and potentially responsive to immunotherapy. In contrast, *BRCA2*
mutation was associated with upregulated MHC-II expression, which plays a role in
antigen presentation and CD4+ T cell recruitment [53], suggesting a shift toward
adaptive immunity and antigen-presenting cell involvement. *BRCA2*-mutant tumors
showed higher CD4+ T cell activation, particularly inflammatory subsets, and were

associated with increased NK cell activity and T cell cytotoxicity in blood, suggesting systemic immune engagement.

Despite these immune signatures, *BRCA1/2* mutations were associated with downregulation of CD28 family co-stimulatory molecules, essential for effective T cell activation [54]. This may indicate adaptive resistance mechanisms. ICB may overcome such resistance (e.g., blocking PD-L1), unleashing the pre-existing, infiltration-potentiated immune response (Fig. 1G). We identified two Trm subsets (CD8+ Trm and CD4+ Trm) as potential biomarkers for ICB response in NSCLC. *BRCA1* mutation was associated with CD8+ Trm TCR expansion, while *BRCA2* mutation was linked to CD4+ Trm expansion, which may contribute to maintain a chronic inflammatory TME [55, 56]. Given the observed T cell infiltration and IFN-driven immune activation, *BRCA1*-mutant tumors might initially respond well to anti-PD-1/PD-L1 therapies. However, *BRCA1* mutations also activated risk genes (*S100A10*, *LDHA*, *GAPDH*, and *MYL12A*). We identified HDAC inhibitors as potential agents targeting this program. HDAC inhibitors can modulate chromatin accessibility, reduce immune suppression, and enhance tumor immunogenicity, suggesting that combining them with ICB might enhance immune response and overcome resistance mechanisms in *BRCA1*-driven tumors.

Despite the comprehensive multi-omics and single-cell analyses conducted in this study, several limitations should be acknowledged. Our study primarily relied on retrospective data from public cohorts and a limited number of in-house samples. Although we included both tumor tissues and matched paracancerous and blood samples, the sample size remains modest, particularly for patients with *BRCA1* or *BRCA2* mutations. While our single-cell RNA-seq and TCR-seq analyses provide high-resolution insights into the cellular and clonal dynamics of the TME, the functional validation of key mechanisms—such as the causal role of *BRCA1*-driven cGAS-STING activation in CD8+ T cell recruitment—remains largely correlative. Experimental models, such as *BRCA1/2*-knockout cell lines or patient-derived organoids, coupled with in vivo studies, are required to establish causal relationships

and elucidate the underlying molecular pathways. Finally, the therapeutic implications

of HDAC inhibitors in *BRCA1*-mutant LUAD, although supported by *in vitro* data,

warrant further preclinical and clinical investigation. The efficacy and safety of

combining HDAC inhibitors with ICB in *BRCA1*-mutant LUAD patients remain

unexplored and represent an important avenue for future research.

5. Conclusions

Overall, this study provides a high-resolution molecular map of the TME in

BRCA1/2-mutant LUAD, highlighting distinct immune and transcriptional signatures

associated with each mutation in the profiled cases. By uncovering these molecular

distinctions, our study offers a hypothesis-generating framework for understanding

HRR deficiency in LUAD and lays groundwork for future validation and precision

immunotherapy strategies tailored to *BRCA1/2* mutation status.

Abbreviations

NSCLC    Non-small cell lung cancer

LUAD    Lung adenocarcinoma

HRR    Homologous recombination repair

HRD    Homologous recombination repair deficiency

TMB    Tumor mutation burden

TME    Tumor microenvironment

ICB    Immune checkpoint blockade

Trm    Tissue-resident memory T cell

HDAC    Histone deacetylase

BRCA1/2    BReast CAncer gene 1/2

TCGA    The cancer genome atlas

UMI    Unique molecular identifier

UMAP    Uniform manifold approximation and projection

TCR    T-cell receptor

MSigDB    Molecular signatures database

761	ssGSEA	Single sample gene set enrichment analysis
762	NMF	Nonnegative matrix factorization
763	cMap	Connectivity Map
764	LINCS	Library of integrated network-based cellular signatures
765	CCLE	Cancer cell line encyclopedia

CRediT authorship contribution statement

Gaoming Liao: Writing - original draft, Visualization, Formal analysis, Data curation,
Conceptualization, Software, Methodology, Funding acquisition. **Xinbin Yang**:
Writing - review & editing, Investigation, Methodology. **Qi Liu**: Writing - review &
editing, Investigation, Validation. **Shufeng Nan**: Validation, Software. **Yan Liu**:
Software. **Jinwei Li**: Validation. **Si Huang**: Validation. **Wang Ning**: Validation.
Xionghai Qin: Writing - review & editing, Supervision, Conceptualization. **Gang Xu**:
Writing - review & editing, Project administration, Supervision, Conceptualization.

Funding

This study was supported by the National Natural Science Foundation of China (Grant
No. 82403816), the Postdoctoral Fellowship Program of CPSF (Grant No.
GZB20230180), the Guangdong Basic and Applied Basic Research Foundation (Grant
No. 2023A1515110297), the China Postdoctoral Foundation (Grant No.
2023M740846), and the Fundamental Research Funds for the Provincial Universities
(2023-KYYWF-0212).

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal
relationships that could have appeared to influence the work reported in this paper.

Data availability

The raw data of single-cell sequencing (including scRNA-seq and scTCR-seq) and the
processed data generated in this study have been deposited at the Gene Expression
Omnibus (GEO) repository, with the accession code GSE292700. Open-source R

packages and software, as well as standard workflows were used in this study. No previously unreported custom code was developed for the analyses presented. All codes and other processed data are available from the corresponding author upon reasonable request.

Acknowledgements

The authors wish to thank the staff members from Affiliated Cancer Hospital and Institute of Guangzhou Medical University. High-throughput sequencing was performed by the OE Biotech Co., Ltd. (Shanghai, China). The contribution of the bioinformatics core facility at Guangzhou medical university is gratefully acknowledged.

References

[1] D.J. Myers, J.M. Wallen, Lung Adenocarcinoma, StatPearls, Treasure Island (FL) ineligible companies. Disclosure: Jason Wallen declares no relevant financial relationships with ineligible companies., 2025.

[2] F. Skoulidis, J.V. Heymach, Co-occurring genomic alterations in non-small-cell lung cancer biology and therapy, Nat Rev Cancer 19(9) (2019) 495-509.

[3] B. Nguyen, C. Fong, A. Luthra, S.A. Smith, R.G. DiNatale, S. Nandakumar, H. Walch, W.K. Chatila, R. Madupuri, R. Kundra, C.M. Bielski, B. Mastrogiacomo, M.T.A. Donoghue, A. Boire, S. Chandarlapaty, K. Ganesh, J.J. Harding, C.A. Iacobuzio-Donahue, P. Razavi, E. Reznik, C.M. Rudin, D. Zamarin, W. Abida, G.K. Abou-Alfa, C. Aghajanian, A. Cercek, P. Chi, D. Feldman, A.L. Ho, G. Iyer, Y.Y. Janjigian, M. Morris, R.J. Motzer, E.M. O'Reilly, M.A. Postow, N.P. Raj, G.J. Riely, M.E. Robson, J.E. Rosenberg, A. Safonov, A.N. Shoushtari, W. Tap, M.Y. Teo, A.M. Varghese, M. Voss, R. Yaeger, M.G. Zauderer, N. Abu-Rustum, J. Garcia-Aguilar, B. Bochner, A. Hakimi, W.R. Jarnagin, D.R. Jones, D. Molena, L. Morris, E. Rios-Doria, P. Russo, S. Singer, V.E. Strong, D. Chakravarty, L.H. Ellenson, A. Gopalan, J.S. Reis-Filho, B. Weigelt, M. Ladanyi, M. Gonen, S.P. Shah, J. Massague, J. Gao, A. Zehir, M.F. Berger, D.B. Solit, S.F. Bakhoum, F. Sanchez-Vega, N. Schultz, Genomic characterization of metastatic patterns from prospective clinical sequencing of 25,000 patients, Cell 185(3) (2022) 563-575 e11.

[4] N.M. Novikov, S.Y. Zolotaryova, A.M. Gautreau, E.V. Denisov, Mutational drivers of cancer cell migration and invasion, Br J Cancer 124(1) (2021) 102-114.

[5] K.E. de Visser, J.A. Joyce, The evolving tumor microenvironment: From cancer initiation to metastatic outgrowth, Cancer Cell 41(3) (2023) 374-403.

[6] N.A. Giraldo, R. Sanchez-Salas, J.D. Peske, Y. Vano, E. Becht, F. Petitprez, P. Validire, A. Ingels, X. Cathelineau, W.H. Fridman, C. Sautes-Fridman, The clinical role of the TME in solid cancer, Br J Cancer 120(1) (2019) 45-53.

[7] M.M. Hoppe, R. Sundar, D.S.P. Tan, A.D. Jeyasekharan, Biomarkers for Homologous Recombination Deficiency in Cancer, J Natl Cancer Inst 110(7) (2018) 704-713.

[8] G. Liao, Z. Jiang, Y. Yang, C. Zhang, M. Jiang, J. Zhu, L. Xu, A. Xie, M. Yan, Y. Zhang, Y. Xiao, X. Li, Combined homologous recombination repair deficiency and immune activation analysis for predicting intensified responses of anthracycline, cyclophosphamide and taxane chemotherapy in triple-negative breast cancer, BMC Med 19(1) (2021) 190.

[9] B. Yan, B. Xie, M. Huang, J. Guo, J. Sun, J. Chen, Y. Tao, D. Xiao, Mutations and expressions of breast cancer 1/2 in lung cancer, Thorac Cancer 14(18) (2023) 1753-1763.

[10] R.M. Samstein, C. Krishna, X. Ma, X. Pei, K.W. Lee, V. Makarov, F. Kuo, J. Chung, R.M. Srivastava, T.A. Purohit, D.R. Hoen, R. Mandal, J. Setton, W. Wu, R. Shah, B. Qeriqi, Q. Chang, S. Kendall, L. Braunstein, B. Weigelt, P. Blecua Carrillo Albornoz, L.G.T. Morris, D.L. Mandelker, J.S. Reis-Filho, E. de Stanchina, S.N. Powell, T.A. Chan, N. Riaz, Mutations in BRCA1 and BRCA2 differentially affect the tumor microenvironment and response to checkpoint blockade immunotherapy, Nat Cancer 1(12) (2021) 1188-1203.

[11] M. Bruand, D. Barras, M. Mina, E. Ghisoni, M. Morotti, E. Lanitis, N. Fahr, M. Desbuisson, A. Grimm, H. Zhang, C. Chong, J. Dagher, S. Chee, T. Tsianou, J. Dorier, B.J. Stevenson, C. Iseli, C. Ronet, S. Bobisse, R. Genolet, J. Walton, M. Bassani-Sternberg, L.E. Kandalaft, B. Ren, I. McNeish, E. Swisher, A. Harari, M. Delorenzi, G. Ciriello, M. Irving, S. Rusakiewicz, P.G. Foukas, F. Martinon, D. Dangaj Laniti, G. Coukos, Cell-autonomous inflammation of BRCA1-deficient ovarian cancers drives both tumor-intrinsic immunoreactivity and immune resistance via STING, Cell Rep 36(3) (2021) 109412.

[12] J.C. Francis, L. Melchor, J. Campbell, H. Kendrick, W. Wei, J. Armisen-Garrido, I. Assiotis, L. Chen, I. Kozarewa, K. Fenwick, A. Swain, M.J. Smalley, C.J. Lord, A. Ashworth, Whole-exome DNA sequence analysis of Brca2- and Trp53-deficient mouse mammary gland tumours, J Pathol 236(2) (2015) 186-200.

[13] Y.C. Lee, Y.C. Lee, C.Y. Li, Y.L. Lee, B.L. Chen, BRCA1 and BRCA2 Gene Mutations and Lung Cancer Sisk: A Meta-Analysis, Medicina (Kaunas) 56(5) (2020).

[14] R.M. Chabanon, M. Pedrero, C. Lefebvre, A. Marabelle, J.C. Soria, S. Postel-Vinay, Mutational Landscape and Sensitivity to Immune Checkpoint Blockers, Clin Cancer Res 22(17) (2016) 4309-21.

[15] Z. Zhou, Z. Ding, J. Yuan, S. Shen, H. Jian, Q. Tan, Y. Yang, Z. Chen, Q. Luo, X. Cheng, Y. Yu, X. Niu, L. Qian, X. Chen, L. Gu, R. Liu, S. Ma, J. Huang, T. Chen, Z. Li, W. Ji, L. Song, L. Shen, L. Jiang, Z. Yu, C. Zhang, Z. Tai, C. Wang, R. Chen, D.P. Carbone, X. Xia, S. Lu, Homologous recombination deficiency (HRD) can predict the therapeutic outcomes of immuno-neoadjuvant therapy in NSCLC patients, J Hematol Oncol 15(1) (2022) 62.

[16] C. Papadaki, M. Sfakianaki, G. Ioannidis, E. Lagoudaki, M. Trypaki, K. Tryfonidis, D. Mavroudis, E. Stathopoulos, V. Georgoulias, J. Souglakos, ERCC1 and BRAC1 mRNA expression levels in the primary tumor could predict the effectiveness of the second-line cisplatin-based chemotherapy in pretreated patients with metastatic non-small cell lung cancer, J Thorac Oncol 7(4) (2012) 663-71.

[17] L. Bonanno, C. Costa, M. Majem, A. Favaretto, M. Rugge, R. Rosell, The predictive value of BRCA1 and RAP80 mRNA expression in advanced non-small-cell lung cancer patients treated with platinum-based chemotherapy, Ann Oncol 24(4) (2013) 1130-2.

[18] T. Motohashi, K. Isobe, T. Yoshizawa, Y. Usui, H. Shimizu, M. Sekiya, S. Miyoshi, Y. Nakamura, N. Urabe, S. Sakamoto, S. Homma, S. Sadamoto, N. Tochigi, K. Kishi, BRCA2-positive lung adenocarcinoma treated with olaparib: A case report, Respirol Case Rep 12(3) (2024) e01317.

[19] D. Jovic, X. Liang, H. Zeng, L. Lin, F. Xu, Y. Luo, Single-cell RNA sequencing technologies and
applications: A brief overview, Clin Transl Med 12(3) (2022) e694.

[20] K.J. Travaglini, A.N. Nabhan, L. Penland, R. Sinha, A. Gillich, R.V. Sit, S. Chang, S.D. Conley, Y.
Mori, J. Seita, G.J. Berry, J.B. Shrager, R.J. Metzger, C.S. Kuo, N. Neff, I.L. Weissman, S.R.
Quake, M.A. Krasnow, A molecular cell atlas of the human lung from single-cell RNA sequencing,
Nature 587(7835) (2020) 619-625.

[21] A. Haque, J. Engel, S.A. Teichmann, T. Lonnberg, A practical guide to single-cell
RNA-sequencing for biomedical research and clinical applications, Genome Med 9(1) (2017) 75.

[22] M.L. Frank, K. Lu, C. Erdogan, Y. Han, J. Hu, T. Wang, J.V. Heymach, J. Zhang, A. Reuben,
T-Cell Receptor Repertoire Sequencing in the Era of Cancer Immunotherapy, Clin Cancer Res
29(6) (2023) 994-1008.

[23] J. Chen, H. Yang, A.S.M. Teo, L.B. Amer, F.G. Sherbaf, C.Q. Tan, J.J.S. Alvarez, B. Lu, J.Q. Lim,
A. Takano, R. Nahar, Y.Y. Lee, C.Z.J. Phua, K.P. Chua, L. Suteja, P.J. Chen, M.M. Chang, T.P.T.
Koh, B.H. Ong, D. Anantham, A.A.L. Hsu, A. Gogna, C.W. Too, Z.W. Aung, Y.F. Lee, L. Wang,
T.K.H. Lim, A. Wilm, P.S. Choi, P.Y. Ng, C.K. Toh, W.T. Lim, S. Ma, B. Lim, J. Liu, W.L. Tam,
A.J. Skanderup, J.P.S. Yeong, E.H. Tan, C.L. Creasy, D.S.W. Tan, A.M. Hillmer, W. Zhai,
Genomic landscape of lung adenocarcinoma in East Asians, Nat Genet 52(2) (2020) 177-186.

[24] A. Ravi, M.D. Hellmann, M.B. Arniella, M. Holton, S.S. Freeman, V. Naranbhai, C. Stewart, I.
Leshchiner, J. Kim, Y. Akiyama, A.T. Griffin, N.I. Vokes, M. Sakhi, V. Kamesan, H. Rizvi, B.
Ricciuti, P.M. Forde, V. Anagnostou, J.W. Riess, D.L. Gibbons, N.A. Pennell, V. Velcheti, S.R.
Digumarthy, M. Mino-Kenudson, A. Califano, J.V. Heymach, R.S. Herbst, J.R. Brahmer, K.A.
Schalper, V.E. Velculescu, B.S. Henick, N. Rizvi, P.A. Janne, M.M. Awad, A. Chow, B.D.
Greenbaum, M. Luksza, A.T. Shaw, J. Wolchok, N. Hacohen, G. Getz, J.F. Gainor, Genomic and
transcriptomic analysis of checkpoint blockade response in advanced non-small cell lung cancer,
Nat Genet 55(5) (2023) 807-819.

[25] N.S. Patil, B.Y. Nabet, S. Muller, H. Koeppen, W. Zou, J. Giltnane, A. Au-Yeung, S. Srivats, J.H.
Cheng, C. Takahashi, P.E. de Almeida, A.S. Chitre, J.L. Grogan, L. Rangell, S. Jayakar, M.
Peterson, A.W. Hsia, W.E. O'Gorman, M. Ballinger, R. Banchereau, D.S. Shames, Intratumoral
plasma cells predict outcomes to PD-L1 blockade in non-small cell lung cancer, Cancer Cell 40(3)
(2022) 289-300 e4.

[26] V. Thorsson, D.L. Gibbs, S.D. Brown, D. Wolf, D.S. Bortone, T.H. Ou Yang, E. Porta-Pardo, G.F.
Gao, C.L. Plaisier, J.A. Eddy, E. Ziv, A.C. Culhane, E.O. Paull, I.K.A. Sivakumar, A.J. Gentles, R.
Malhotra, F. Farshidfar, A. Colaprico, J.S. Parker, L.E. Mose, N.S. Vo, J. Liu, Y. Liu, J. Rader, V.
Dhankani, S.M. Reynolds, R. Bowlby, A. Califano, A.D. Cherniack, D. Anastassiou, D.
Bedognetti, Y. Mokrab, A.M. Newman, A. Rao, K. Chen, A. Krasnitz, H. Hu, T.M. Malta, H.
Noushmehr, C.S. Pedamallu, S. Bullman, A.I. Ojesina, A. Lamb, W. Zhou, H. Shen, T.K. Choueiri,
J.N. Weinstein, J. Guinney, J. Saltz, R.A. Holt, C.S. Rabkin, N. Cancer Genome Atlas Research,
A.J. Lazar, J.S. Serody, E.G. Demicco, M.L. Disis, B.G. Vincent, I. Shmulevich, The Immune
Landscape of Cancer, Immunity 48(4) (2018) 812-830 e14.

[27] T. Ilicic, J.K. Kim, A.A. Kolodziejczyk, F.O. Bagger, D.J. McCarthy, J.C. Marioni, S.A.
Teichmann, Classification of low quality cells from single-cell RNA-seq data, Genome Biol 17
(2016) 29.

[28] B. Pijuan-Sala, J.A. Griffiths, C. Guibentif, T.W. Hiscock, W. Jawaid, F.J. Calero-Nieto, C. Mulas,
X. Ibarra-Soria, R.C.V. Tyser, D.L.L. Ho, W. Reik, S. Srinivas, B.D. Simons, J. Nichols, J.C.

Marioni, B. Gottgens, A single-cell molecular map of mouse gastrulation and early organogenesis,
Nature 566(7745) (2019) 490-495.

[29] C.S. McGinnis, L.M. Murrow, Z.J. Gartner, DoubletFinder: Doublet Detection in Single-Cell RNA
Sequencing Data Using Artificial Nearest Neighbors, Cell Syst 8(4) (2019) 329-337 e4.

[30] P.L. Germain, A. Lun, C. Garcia Meixide, W. Macnair, M.D. Robinson, Doublet identification in
single-cell sequencing data using scDblFinder, F1000Res 10 (2021) 979.

[31] Y. Hao, T. Stuart, M.H. Kowalski, S. Choudhary, P. Hoffman, A. Hartman, A. Srivastava, G. Molla,
S. Madad, C. Fernandez-Granda, R. Satija, Dictionary learning for integrative, multimodal and
scalable single-cell analysis, Nat Biotechnol 42(2) (2024) 293-304.

[32] J. Wienke, L.L. Visser, W.M. Kholosy, K.M. Keller, M. Barisa, E. Poon, S. Munnings-Tomes, C.
Himsworth, E. Calton, A. Rodriguez, R. Bernardi, F. van den Ham, S.R. van Hooff, Y.A.H. Matser,
M.L. Tas, K.P.S. Langenberg, P. Lijnzaad, A.L. Borst, E. Zappa, F.J. Bergsma, J.G.M. Strijker,
B.M. Verhoeven, S. Mei, A. Kramdi, R. Restuadi, A. Sanchez-Bernabeu, A.M. Cornel, F.C.P.
Holstege, J.C. Gray, G.A.M. Tytgat, M.A. Scheijde-Vermeulen, M. Wijnen, M.P. Dierselhuis, K.
Straathof, S. Behjati, W. Wu, A.J.R. Heck, J. Koster, S. Nierkens, I. Janoueix-Lerosey, R.R. de
Krijger, N. Baryawno, L. Chesler, J. Anderson, H.N. Caron, T. Margaritis, M.M. van Noesel, J.J.
Molenaar, Integrative analysis of neuroblastoma by single-cell RNA sequencing identifies the
NECTIN2-TIGIT axis as a target for immunotherapy, Cancer Cell 42(2) (2024) 283-300 e8.

[33] A. Gavish, M. Tyler, A.C. Greenwald, R. Hoefflin, D. Simkin, R. Tschernichovsky, N. Galili
Darnell, E. Somech, C. Barbolin, T. Antman, D. Kovarsky, T. Barrett, L.N. Gonzalez Castro, D.
Halder, R. Chanoch-Myers, J. Laffy, M. Mints, A. Wider, R. Tal, A. Spitzer, T. Hara, M.
Raitses-Gurevich, C. Stossel, T. Golan, A. Tirosh, M.L. Suva, S.V. Puram, I. Tirosh, Hallmarks of
transcriptional intratumour heterogeneity across a thousand tumours, Nature 618(7965) (2023)
598-606.

[34] S. Mariathasan, S.J. Turley, D. Nickles, A. Castiglioni, K. Yuen, Y. Wang, E.E. Kadel, III, H.
Koeppen, J.L. Astarita, R. Cubas, S. Jhunjhunwala, R. Banchereau, Y. Yang, Y. Guan, C. Chalouni,
J. Ziai, Y. Senbabaoglu, S. Santoro, D. Sheinson, J. Hung, J.M. Giltnane, A.A. Pierce, K. Mesh, S.
Lianoglou, J. Riegler, R.A.D. Carano, P. Eriksson, M. Hoglund, L. Somarriba, D.L. Halligan, M.S.
van der Heijden, Y. Loriot, J.E. Rosenberg, L. Fong, I. Mellman, D.S. Chen, M. Green, C. Derleth,
G.D. Fine, P.S. Hegde, R. Bourgon, T. Powles, TGFbeta attenuates tumour response to PD-L1
blockade by contributing to exclusion of T cells, Nature 554(7693) (2018) 544-548.

[35] G. Bindea, B. Mlecnik, M. Tosolini, A. Kirilovsky, M. Waldner, A.C. Obenauf, H. Angell, T.
Fredriksen, L. Lafontaine, A. Berger, P. Bruneval, W.H. Fridman, C. Becker, F. Pages, M.R.
Speicher, Z. Trajanoski, J. Galon, Spatiotemporal dynamics of intratumoral immune cells reveal
the immune landscape in human cancer, Immunity 39(4) (2013) 782-95.

[36] G. Liao, Y. Yang, A. Xie, Z. Jiang, J. Liao, M. Yan, Y. Zhou, J. Zhu, J. Hu, Y. Zhang, Y. Xiao, X. Li,
Applicability of Anticancer Drugs for the Triple-Negative Breast Cancer Based on Homologous
Recombination Repair Deficiency, Front Cell Dev Biol 10 (2022) 845950.

[37] F. Sanchez-Vega, M. Mina, J. Armenia, W.K. Chatila, A. Luna, K.C. La, S. Dimitriadoy, D.L. Liu,
H.S. Kantheti, S. Saghafinia, D. Chakravarty, F. Daian, Q. Gao, M.H. Bailey, W.W. Liang, S.M.
Foltz, I. Shmulevich, L. Ding, Z. Heins, A. Ochoa, B. Gross, J. Gao, H. Zhang, R. Kundra, C.
Kandoth, I. Bahceci, L. Dervishi, U. Dogrusoz, W. Zhou, H. Shen, P.W. Laird, G.P. Way, C.S.
Greene, H. Liang, Y. Xiao, C. Wang, A. Iavarone, A.H. Berger, T.G. Bivona, A.J. Lazar, G.D.
Hammer, T. Giordano, L.N. Kwong, G. McArthur, C. Huang, A.D. Tward, M.J. Frederick, F.

McCormick, M. Meyerson, N. Cancer Genome Atlas Research, E.M. Van Allen, A.D. Cherniack,

G. Ciriello, C. Sander, N. Schultz, Oncogenic Signaling Pathways in The Cancer Genome Atlas,

Cell 173(2) (2018) 321-337 e10.

[38] I. Tirosh, A.S. Venteicher, C. Hebert, L.E. Escalante, A.P. Patel, K. Yizhak, J.M. Fisher, C.

Rodman, C. Mount, M.G. Filbin, C. Neftel, N. Desai, J. Nyman, B. Izar, C.C. Luo, J.M. Francis,

A.A. Patel, M.L. Onozato, N. Riggi, K.J. Livak, D. Gennert, R. Satija, B.V. Nahed, W.T. Curry,

R.L. Martuza, R. Mylvaganam, A.J. Iafrate, M.P. Frosch, T.R. Golub, M.N. Rivera, G. Getz, O.

Rozenblatt-Rosen, D.P. Cahill, M. Monje, B.E. Bernstein, D.N. Louis, A. Regev, M.L. Suva,

Single-cell RNA-seq supports a developmental hierarchy in human oligodendroglioma, Nature

539(7628) (2016) 309-313.

[39] E. Munkhbaatar, M. Dietzen, D. Agrawal, M. Anton, M. Jesinghaus, M. Boxberg, N. Pfarr, P.

Bidola, S. Uhrig, U. Hockendorf, A.L. Meinhardt, A. Wahida, I. Heid, R. Braren, R. Mishra, A.

Warth, T. Muley, P.S.P. Poh, X. Wang, S. Frohling, K. Steiger, J. Slotta-Huspenina, M. van

Griensven, F. Pfeiffer, S. Lange, R. Rad, M. Spella, G.T. Stathopoulos, J. Ruland, F. Bassermann,

W. Weichert, A. Strasser, C. Branca, M. Heikenwalder, C. Swanton, N. McGranahan, P.J. Jost,

MCL-1 gains occur with high frequency in lung adenocarcinoma and can be targeted

therapeutically, Nat Commun 11(1) (2020) 4527.

[40] Z. Wang, Y. Wang, M. Chang, Y. Wang, P. Liu, J. Wu, G. Wang, X. Tang, X. Hui, P. Liu, X. Guo, B.

Xing, Y. Wang, Z. Han, W. Ma, Single-cell transcriptomic analyses provide insights into the

cellular origins and drivers of brain metastasis from lung adenocarcinoma, Neuro Oncol 25(7)

(2023) 1262-1274.

[41] T. Karasaki, D.A. Moore, S. Veeriah, C. Naceur-Lombardelli, A. Toncheva, N. Magno, S. Ward,

M.A. Bakir, T.B.K. Watkins, K. Grigoriadis, A. Huebner, M.S. Hill, A.M. Frankell, C. Abbosh, C.

Puttick, H. Zhai, F. Gimeno-Valiente, S. Saghafinia, N. Kanu, M. Dietzen, O. Pich, E.L. Lim, C.

Martinez-Ruiz, J.R.M. Black, D. Biswas, B.B. Campbell, C. Lee, E. Colliver, K.S.S. Enfield, S.

Hessey, C.T. Hiley, S. Zaccaria, K. Litchfield, N.J. Birkbak, E.L. Cadieux, J. Demeulemeester, P.

Van Loo, P.S. Adusumilli, K.S. Tan, W. Cheema, F. Sanchez-Vega, D.R. Jones, N. Rekhtman, W.D.

Travis, A. Hackshaw, T. Marafioti, R. Salgado, J. Le Quesne, A.G. Nicholson, T.R. Consortium, N.

McGranahan, C. Swanton, M. Jamal-Hanjani, Evolutionary characterization of lung

adenocarcinoma morphology in TRACERx, Nat Med 29(4) (2023) 833-845.

[42] V. Moncho-Amor, I. Ibanez de Caceres, E. Bandres, B. Martinez-Poveda, J.L. Orgaz, I.

Sanchez-Perez, S. Zazo, A. Rovira, J. Albanell, B. Jimenez, F. Rojo, C. Belda-Iniesta, J.

Garcia-Foncillas, R. Perona, DUSP1/MKP1 promotes angiogenesis, invasion and metastasis in

non-small-cell lung cancer, Oncogene 30(6) (2011) 668-78.

[43] H. Bayir, S.J. Dixon, Y.Y. Tyurina, J.A. Kellum, V.E. Kagan, Ferroptotic mechanisms and

therapeutic targeting of iron metabolism and lipid peroxidation in the kidney, Nat Rev Nephrol

19(5) (2023) 315-336.

[44] A. Salcedo, M. Tarabichi, A. Buchanan, S.M.G. Espiritu, H. Zhang, K. Zhu, T.H. Ou Yang, I.

Leshchiner, D. Anastassiou, Y. Guan, G.H. Jang, M.F.E. Mootor, K. Haase, A.G. Deshwar, W. Zou,

I. Umar, S. Dentro, J.A. Wintersinger, K. Chiotti, J. Demeulemeester, C. Jolly, L. Sycza, M. Ko, P.

Evolution, G. Heterogeneity Working, S.M.-H. Participants, D.C. Wedge, Q.D. Morris, K. Ellrott,

P. Van Loo, P.C. Boutros, Crowd-sourced benchmarking of single-sample tumor subclonal

reconstruction, Nat Biotechnol   (2024).

[45] J. Cao, M. Spielmann, X. Qiu, X. Huang, D.M. Ibrahim, A.J. Hill, F. Zhang, S. Mundlos, L.

Christiansen, F.J. Steemers, C. Trapnell, J. Shendure, The single-cell transcriptional landscape of mammalian organogenesis, Nature 566(7745) (2019) 496-502.

[46] S.M. Harding, J.L. Benci, J. Irianto, D.E. Discher, A.J. Minn, R.A. Greenberg, Mitotic progression following DNA damage enables pattern recognition within micronuclei, Nature 548(7668) (2017) 466-470.

[47] J. Kwon, S.F. Bakhoum, The Cytosolic DNA-Sensing cGAS-STING Pathway in Cancer, Cancer Discov 10(1) (2020) 26-39.

[48] Q. Chen, L. Sun, Z.J. Chen, Regulation and function of the cGAS-STING pathway of cytosolic DNA sensing, Nat Immunol 17(10) (2016) 1142-9.

[49] Y. Deng, L. Xia, J. Zhang, S. Deng, M. Wang, S. Wei, K. Li, H. Lai, Y. Yang, Y. Bai, Y. Liu, L. Luo, Z. Yang, Y. Chen, R. Kang, F. Gan, Q. Pu, J. Mei, L. Ma, F. Lin, C. Guo, H. Liao, Y. Zhu, Z. Liu, C. Liu, Y. Hu, Y. Yuan, Z. Zha, G. Yuan, G. Zhang, L. Chen, Q. Cheng, S. Shen, L. Liu, Multicellular ecotypes shape progression of lung adenocarcinoma from ground-glass opacity toward advanced stages, Cell Rep Med 5(4) (2024) 101489.

[50] J. Li, C. Zhou, X. Gao, T. Tan, M. Zhang, Y. Li, H. Chen, R. Wang, B. Wang, J. Liu, P. Liu, S100A10 promotes cancer metastasis via recruitment of MDSCs within the lungs, Oncoimmunology 13(1) (2024) 2381803.

[51] M. Certo, C.H. Tsai, V. Pucino, P.C. Ho, C. Mauro, Lactate modulation of immune responses in inflammatory versus tumour microenvironments, Nat Rev Immunol 21(3) (2021) 151-161.

[52] A. Hu, L. Sun, H. Lin, Y. Liao, H. Yang, Y. Mao, Harnessing innate immune pathways for therapeutic advancement in cancer, Signal Transduct Target Ther 9(1) (2024) 68.

[53] M.L. Axelrod, R.S. Cook, D.B. Johnson, J.M. Balko, Biological Consequences of MHC-II Expression by Tumor Cells in Cancer, Clin Cancer Res 25(8) (2019) 2392-2402.

[54] K.P. Burke, A. Chaudhri, G.J. Freeman, A.H. Sharpe, The B7:CD28 family and friends: Unraveling coinhibitory interactions, Immunity 57(2) (2024) 223-244.

[55] H. Chi, M. Pepper, P.G. Thomas, Principles and therapeutic applications of adaptive immunity, Cell 187(9) (2024) 2052-2078.

[56] B. Kruse, A.C. Buzzai, N. Shridhar, A.D. Braun, S. Gellert, K. Knauth, J. Pozniak, J. Peters, P. Dittmann, M. Mengoni, T.C. van der Sluis, S. Hohn, A. Antoranz, A. Krone, Y. Fu, D. Yu, M. Essand, R. Geffers, D. Mougiakakos, S. Kahlfuss, H. Kashkar, E. Gaffal, F.M. Bosisio, O. Bechter, F. Rambow, J.C. Marine, W. Kastenmuller, A.J. Muller, T. Tuting, CD4(+) T cell-induced inflammatory cell death controls immune-evasive tumours, Nature 618(7967) (2023) 1033-1040.

Figure legend

1034

Fig. 1. *BRCA1* and *BRCA2* mutations are associated with genomic instability and poor prognosis of LUAD. A. Distribution of HRD score, TMB and neoantigen load in patients with wild-type, *BRCA1*- and *BRCA2*- mutated lung adenocarcinoma from the TCGA-LUAD cohort (n=566). B. Proportion of immune subtypes. The significance level was calculated by Chi-squared test. C1: wound healing, C2: IFN-g dominant, C3: inflammatory, C4: lymphocyte depleted, C6: TGF-b dominant. C-D. Activity of DNA damage repair and cell cycle-related pathways in wild-type, *BRCA1*- and *BRCA2*- mutated patients (TCGA-LUAD cohort). E-F. Survival analysis of BRCA mutations in TCGA-LUAD (E) and OncoSG-LUAD (F) cohorts. G. Survival analysis of BRCA mutations in LUAD patients after ICB treatment. H. Survival analysis based on *BRCA1* (left) and *BRCA2* (right) expression in patients with ICB treatment. I. Lymphocyte infiltrates in patients with wild-type, *BRCA1* and *BRCA2* mutations (TCGA-LUAD cohort). J. Mutation frequencies of the oncogenic signaling pathway genes (in-house data).

1049

Fig. 2. Single-cell transcriptome analysis and LUAD malignant cells identification. A. UMAP visualization of cell types in patients with *BRCA1/2* mutations (in-house data). B. The number of each cell type and its proportion in different samples, including tumor tissue (T), adjacent normal tissue (N), and peripheral blood (B). C. The bubble plot shows the expression of the markers used for cell type identification. D. The heat map inferred CNVs in epithelial cells, malignant cells, and reference cells. E. Distribution of CNV scores across different cell types on specific chromosomes. F. UMAP visualization of CNV score in epithelial cells and malignant cells. G. UMAP visualization (left) and box plot (right) of tumor upregulated signature score. H. Expression of *CEACAM6* in epithelial and malignant cells.

Fig. 3. Identification of key programs in *BRCA1*/2 mutated tumors. A. Functional enrichment analysis of genes that upregulated in malignant cells relative to normal epithelial cells. B. The heat map of the similarity between pairs of programs is identified based on the NMF algorithm. C. MP score between *BRCA1* and *BRCA2* mutated malignant cells. D. K-M curves for MP1 score survival analysis. E. Enrichment map of the functions enriched by MP1 and MP2 genes. F. The overlap of in-house MPs and known cancer MPs. G. The heat map shows the activity of known cancer MPs in *BRCA1* and *BRCA2* mutated malignant cells. H. Volcano diagram of differentially expressed genes between *BRCA1* and *BRCA2* mutated malignant cells.

1070

Fig. 4. Tumor evolutionary analysis of *BRCA1* and *BRCA2* mutations. A. Pseudotime analysis of epithelial and malignant cells. The branched trajectory was colored by pseudotime (left) and cell types (right). The pie chart shows the proportion of sample groups in each branch. B. The top 50 cell fate genes and their enriched GO BP terms of *BRCA1/2* mutations in malignant2 cell type. C-E. Pseudotime was broken down into 10 bins to smooth gene expression patterns. Average gene module score between

BRCA1 and *BRCA2* mutation groups for tumor down-regulation (C), MP1 cell cycle
G2/M (D), and MP5 stress (E) gene modules. F, J. Cell density distribution based on
cell cycle HMG rich versus stress gene modules (F) and MHC class I versus MHC
class II gene modules (J) in the same cells between *BRCA1* and *BRCA2* mutations.
Color represents the density of cells. The dotted line represents the median value of
the corresponding module score. G. Average gene module score for IFN-α response
(top) and MHC class II (bottom) gene signatures. H-I. The activities of representative
signatures between *BRCA1* and *BRCA2* mutations in bin 10 of malignant cells
according to pseudotime.

Fig. 5. *BRCA1/2* mutations are associated with tumor lymphocyte activation. A. Functional enrichment of differentially expressed genes in malignant cells between *BRCA1* and *BRCA2* mutation groups. B. GESA of immune response profile between *BRCA1* and *BRCA2* mutations. Red (blue) indicates the gene set enriched in *BRCA2* (*BRCA1*)-mutated malignant cells. C-D. GESA plot of immune signature (C) and gene

module of immune cell type (D) in malignant2 subset. NES: normalized enrichment score. E. Density ridge plot of representative pathways in two types of malignant subset with *BRCA1* and *BRCA2* mutations. HRR: homologous recombination repair. F. UMAP visualization of cGAS pathway score. G. Mean of pathway activities related to DNA damage repair across two types of malignant cells with *BRCA1* and *BRCA2* mutations. H. The expression of the representative markers. I. Immune cell activities between *BRCA1* and *BRCA2* mutations in bin 10 of malignant cells according to pseudotime. J. Lymphocyte activity between *BRCA1* and *BRCA2* mutations in lymphocyte cells.

Fig. 6. T lymphocyte infiltration and TCR clonal expansion analysis. A-B. UMAP visualization (A) and proportion (B) of T lymphocyte subsets in combined dataset. C-D. Box plot (C) and UMAP plot (D) of representative immune pathways from ImmPort database. E-F. The activity of tissue-specific T cell signatures from Judith Wienke et al. G. Terminal exhaustion marker expression levels. H. UMAP of lymphocyte subsets in in-house dataset. I. Quantification of clonal size across sample types. Clonotypes are ranked by expansion level, including: single (1 cell), small (>1 and <5 cells), medium (>5 and <20 cells), large (>20 and <100 cells), and

hyperexpanded (>100 and <500 cells). J. Expanded clonotypes distribution in different samples. K. Proportional and dynamic changes in shared clonotypes (top5 per group) between different samples within the same patient. The colors represent different clonotypes. L. The proportion of clonally expanded cells (≥5 clone size) in *BRCA1/2* mutated tumor tissues for specific cell types. M. The sample and cell proportions of the 50 most abundant TCR motifs.

Fig. 7. Targeting *BRCA1* mutation-related risk genes inhibits tumor growth. A-B. Survival analysis of *BRCA1*-mutant signature score in TCGA-LUAD (A) and OAK (B) cohorts. C. Survival analysis of *S100A10* (left) and *LDHA* (right) in TCGA-LUAD cohort according to the expression median. D. The effects of sh-*LDHA*, *S100A10*, *GAPDH*, and vector on cell proliferation were determined by a cell proliferation assay in A549 LUAD cell line. E. The effects of sh-*LDHA*, *S100A10*, and

GAPDH on protein levels in A549 cell line (The original image is shown in Fig. S12A). F. The molecular perturbations of 4-R genes in A549 cell line at 10 μM concentration (24 h) from the LINCS data resource. NCS: Normalized connectivity score. G. The effects of HDAC inhibitors concentration on cell proliferation in A549 cell line. H. Cell replication levels after treatment with different concentrations of entinostat (left) and vorinostat (right) in LUAD cell lines. I. The protein levels of *LDHA*, *S100A10*, and *GAPDH* after treatment with vorinostat (left) and belinostat (right) in A549 cell line were measured using western blotting (The original image is shown in Fig. S12B). J. The mRNA levels of *LDHA* and *GAPDH* after treatment (3 μM) with vorinostat (left) and belinostat (right) in A549 cell line were measured using qPCR. Significance levels were determined by Student's t-test. For all statistical tests, *P≤0.05; **P≤0.01; ***P≤0.001.