Title: Are Synaptic Clefts Directionally Oriented?

Dexuan Tang[1*], Zhi-De Deng[2], Bethanny Danskin[3], Daniel Berger[4], Mark Ingersoll[1,5], Hanbing Lu[6], Bruce Rosen[7], Marom Bikson[8], Gregory Noetscher[1], Sergey Makaroff[1,7]

Affiliations

(1) Department of Electrical and Computer Eng., Worcester Polytechnic Inst., Worcester MA, USA

(2) Computational Neurostimulation Research Program, Experimental Therapeutics and Pathophysiology Branch, NIMH, National Institute of Health, Bethesda MD, USA

(3) Allen Institute for Brain Science, Seattle WA, USA

(4) Dept. of Molecular and Cellular Biology, Harvard Univ., Boston MA, USA

(5) University of Massachusetts Amherst, Amherst MA, USA

(6) Neuroimaging Research Branch, NIDA, Intramural Research Program, National Institute of Health, Baltimore MD, USA

(7) Athinoula A. Martinos Ctr. for Biomedical Imaging, Massachusetts General Hospital, Charlestown MA, USA

(8) Department of Biomedical Engineering, The City College of New York, New York NY, USA

*Corresponding author: Dexuan Tang

Email: dtang2@wpi.edu

ORCID: 0009-0002-8928-9041

Competing Interest Statement: The authors declare that they have no competing interests.

Data Availability Statement: The data that supports the findings of this study are available in the supplementary material of this article.

Keywords:

Synaptic clefts

Synaptic orientation

Synaptic architecture

MICrONS 1 mm^3 cortical volume

H01 1 mm^3 cortical volume

Abstract

Synapses are fundamental building blocks of cortical circuits, yet their geometry is typically regarded as a local property, independent of mesoscale architecture. The prevailing assumption is that synaptic clefts are isotropically oriented in space. Here, we test this assumption by analyzing approximately 117 million synaptic clefts from two independent 1 mm³ electron microscopy datasets: the human H01 middle temporal gyrus and the mouse MICrONS primary visual cortex, using three independent cleft-extraction methods. Across both volumes, we observe that synaptic cleft orientations are not randomly distributed, but instead show statistically significant and spatially coherent directional biases across cortical layers. This mesoscale anisotropy is conserved across species, yet is stronger and more consistent in human association cortex than in mouse sensory cortex, a difference that may reflect the expanded dendritic arbors and greater integrative demands of human pyramidal neurons. We propose that cleft orientation bias is a geometric consequence of the axonal and dendritic architecture that shapes synapse formation, representing a new candidate organizational feature of cortical microarchitecture with potential implications for circuit computation and neuromodulation. These findings motivate targeted physiological studies to determine whether synaptic orientation contributes causally to cortical function.

Main Text

1. Introduction

The cerebral cortex exhibits architectural regularities that recur across species and scales, from the laminar alignment of cell bodies to the columnar organization of dendrites and axons [1],[2],[3]. These structural motifs constrain the flow of information within local circuits and shape the way cortical populations integrate activity across regions [4],[5]. Despite extensive ultrastructural characterization of synapses over several decades [6], one fundamental geometric property has remained largely unexplored: the orientation of synaptic clefts relative to cortical axes.

At the ultrastructural level, a synaptic cleft is a thin disk-shaped gap between pre- and postsynaptic membranes. To make orientation analysis tractable across large-scale electron microscopy reconstructions, we define cleft orientation along the disk's normal axis, typically pointing from the presynaptic to the postsynaptic side. Even when clefts deviate from an ideal disk, their orientation can be quantified as the dominant averaged normal vector of the apposed membranes. The prevailing view, derived from analyses of small datasets, has been that cleft orientations are isotropically distributed at the mesoscale, a local geometric property determined by pre- and postsynaptic specializations but otherwise independent of cortical architecture [7],[8].

This isotropy assumption has never been tested at scale. The emergence of large-volume electron
microscopy reconstructions now makes it possible to do so directly.

If cleft orientations are systematically biased, such anisotropy would constitute a previously
unmeasured structural property of cortical microarchitecture, analogous to the preferential
alignment of axons or the radial orientation of apical dendrites. Mechanistically, orientation bias is
expected as a geometric consequence of the dendritic and axonal architecture that governs where
and how synapses form: pyramidal neurons extend apical dendrites radially toward the pial surface,
basal dendrites tangentially, and en passant axonal boutons contact these compartments at angles
constrained by the local geometry. Because trans-synaptic adhesion molecules such as neurexins
and neuroligins are differentially expressed across cell types, layers, and dendritic compartments
[9], the distribution of synapse types and cleft orientations across cortical space need not be
isotropic. Detecting such anisotropy at the mesoscale would suggest that synaptic geometry
contributes to the spatial patterning of cortical circuits.

In this work, we test the isotropy assumption directly by analyzing approximately 117 million
synaptic clefts from two mesoscale cortical volumes – human H01 [10],[11],[12] and mouse
MICrONS [13],[14],[15],[6] – using three independent cleft-extraction methods. We observe that
synaptic clefts of excitatory neurons in mammalian neocortex are not randomly distributed. Instead,
they exhibit statistically significant, spatially coherent anisotropy across cortical layers. This
mesoscale orientation is evident in both mouse and human cortex, consistent with an evolutionarily
conserved organizational principle. The magnitude of anisotropy is significantly greater in human
association cortex compared to in mouse primary visual cortex, consistent with the expanded
dendritic arbors and greater integrative demands of human pyramidal neurons [16].

We also evaluate methodological artifacts arising from sample preparation and processing
that can bias apparent cleft geometry and therefore require careful control. Subject to these
constraints, the observed mesoscale anisotropy identifies synaptic cleft orientation as an additional,
previously unmeasured axis of cortical microarchitecture. Functionally, a consistent orientation bias
could shape synaptic integration across minicolumns, and thereby support lateral information flow
in association areas [17],[18]. Biophysically, because synaptic clefts form nanoscale dipoles within
the extracellular medium, their collective alignment could influence the generation of endogenous
field potentials, as well as alter how tissue couples to exogenous stimulation [19],[20]. Together,

these results can link synaptic ultrastructure to mesoscale circuit organization and to the electrical
signals used to measure and perturb cortical activity.
68
69 **2. Results**
70
We performed a directional analysis on two mesoscale cortical volumes obtained using scanning
electron microscopy (H01) and transmission electron microscopy (MICrONS), applying three
independent synaptic-cleft extraction approaches and testing methods described in the *Methods*
section.
- **Mesh-touch method (H01, human):** This method was applied to neuronal membrane
meshes generated by Google researchers [11],[12] for excitatory neurons whose somas
were located within the 1 mm³ H01 cortical volume. We processed approximately 940
pyramidal neurons from the repository (a representative subset is shown in Fig.1b,c),
yielding ~1.6 million proofread and post-processed synaptic-cleft geometries (cf. Fig.1l,m).
The clefts were distributed across cortical layers as follows: L1–~0.06 million; L2–~0.5
million; L3–~0.6 million; L4–~0.2 million; L5–~0.2 million; L6–~0.1 million; with the
remainder located in white matter.
- **Voxel-based method (H01, human):** This analysis utilized the excitatory-synapse dataset
provided by the H01 team [10] in voxel format, comprising all putative excitatory synapses
(~111 million) within the H01 sample, downloaded from repository. Unlike the mesh-touch
method, the voxel representation encompasses synapses originating from all neuronal
processes, not solely those of pyramidal neurons.
- **Synapse-table method (MICrONS, mouse):** The publicly available MICrONS cortical
column dataset [15] includes pre- and post-synaptic coordinates of all column neurons
spanning all cortical layers (Figs.6,7). For each synapse, the orientation was defined by
the unit normal vector pointing from the pre- to the post-synaptic side. The total number of
processed clefts was ~3.2 million (~1.4 million in L2/L3, ~0.6 million in L4, ~0.4 million in
L5, and ~0.8 million in L6).
94
2.1. Synaptic clefts of H01 human brain dataset
Fig.1a shows the topology of the H01 human brain sample [10]. The x-axis is (approximately)
directed from L6 to L1. Fig.1b is a typical postsynaptic excitatory pyramidal neuron of the sample
(cyan). Red "hairs" are the associated presynaptic processes. Fig.1c is a close-up view of the apical
dendrite branch located in the top right of Fig.1b. Figs.1d-k show a collection of synaptic geometries
and types studied. Most typical are the axodendritic excitatory synapses of the excitatory neurons
in Figs.1d-g (estimated 81% based on the mesh touch method). Among them, *en passant* boutons

and synapses (Fig.1e-g) dominate terminal synapses (Fig.1d) as approximately 10:1. Figs.1h-j show less typical axosomatic connections. In Fig.1h, a part of the soma membrane is shown and axo-dendritic shaft (in Figs.1i,j) synapses of the excitatory neurons. These synapses are putatively considered inhibitory; their relative number is estimated as ~15% again based on the mesh touch method. Fig.1k shows putative axoaxonic synapses; their relative number is estimated as approximately the remaining 4%. Those synapses are very rare and may be subject to false detection.

2.2. Directional synaptic vectors of H01 human brain datasets

Using the mesh touch method (*Methods* section), normal vectors of the facets of a refined and healed presynaptic membrane (yellow in Fig.1l,m) are found first. They are shown by small red arrows in Fig.1 m. These normal vectors are further averaged (weighted by facet areas), which results in the effective directional vector of the synapse (magenta in Fig.1m). Fig.1n shows an alternative independent approach – synaptic cleft directions extracted using direct voxel data from labeling presynaptic terminals with clustered synaptic vesicles [21]. There, a boundary between pre- and postsynaptic voxels is extracted and triangulated via Delaunay triangulation, in full or partially. Normal vectors of the boundary facets are found and then averaged again weighted by facet areas. Fig.1o demonstrates overlap of voxel and mesh touch synaptic representations (wherever possible). It indicates that the voxel and mesh touch methods may generate similar, but not identical results. Additional details and direct evidence substantiating this assumption are presented in the *Discussion* and *Materials and Methods* sections.

2.3. "Synaptic pattern"

By analogy with the antenna radiation pattern [22], a 3D "synaptic pattern" (a 3D polar plot) has been constructed for each cortical layer of both brain samples. First, all synaptic clefts physically located within a given cortical layer are moved (only translation, no rotation) to the origin. After that, for every discrete direction in 3D characterized by a pair of polar and azimuthal angles, we introduce an elementary solid angle, $d\Omega$, enclosing this direction; the sum of all such non-overlapping $d\Omega$ is equal to 4π. Next, we find the total number of synaptic clefts whose primary axis or direction from Figs.1m,n – which is always oriented from pre- to postsynaptic process – lies within $d\Omega$. The synaptic pattern of the layer is constructed as a set of such absolute numbers corresponding to

133 elementary solid angles for all different pairs of polar and azimuthal angles. The synaptic pattern is
134 finally normalized with respect to its maximum value and is encoded via a color palette.

135   This definition a priori assumes that the dominant signal in the resulting synaptic pattern is
136 coming from true orientation structure, not from a spatial sampling bias. More details and direct
137 proof of this assumption are available in the *Discussion* section and *Materials and Methods* section.

2.4 Layer-by layer synaptic patterns of the 1 mm^3 H01 human brain dataset (from association
cortex) in three principal planes

Figs.2a–d, 3a–d, and 4a–d show the synaptic patterns of the H01 sample in three principal planes:
xz-plane (Fig.2), yz-plane (Fig.3), and xy-plane (Fig.4), along with the corresponding sample
topology and composition. For each plane we show in Figs.2a,3a,4a side views of the H01 sample,
with cortical layers labeled from L1 (left, dark blue) to L6 (right, yellow), and representative cross-
sectional planar rectangles used for plotting the Neuroglancer cell maps for each layer. We also
show in Figs.2b,3b,4b the corresponding Neuroglancer slice. In all cases, all synaptic clefts located
within a given cortical layer were included in constructing that layer's synaptic pattern. We present
two independent sets of patterns. The first set in (Figs.2c,3c,4c), corresponds to synaptic clefts
extracted from membrane meshes using the precise mesh-touch method (Fig.1l,m and *Methods*).
These patterns included both excitatory and inhibitory synapses (Fig.1d–k) of such excitatory
neurons whose somas are located within the sample – approximately 1.6 million processed
synaptic clefts in total. The second set in (Fig.2d,3d,4d), was obtained by processing the H01
synaptic voxel data (Fig.1n and *Methods*), which comprise 111 million voxel-based synaptic cleft
geometries – all putative excitatory synapses within the sample. The same dimensionless
normalized scale (0.3–1.0) is used for all human synaptic patterns.

2.5 Layer-by layer synaptic patterns of the 1 mm^3 MICrONS mouse cortical column (from
primary visual cortex) in three principal planes

The same three-plane layout is shown for the MICrONS column in Figs.2e–g, 3 e–g, and 4e–g, in
the order of yz-plane, xz-plane, and xy-plane. For each plane we show in Fig.2e,3e,4e side views
of the MICrONS sample, labeled from L2/3 (left, red) to L6 (right, yellow). Representative cross-
sectional planar rectangles, used for plotting the Neuroglancer cell maps, are also indicated for
each layer. We also show in Figs.2f,3f,4f the corresponding Neuroglancer slice. The synaptic cleft
orientations (Figs.2g,3g,4g) were acquired directly from the MICrONS database as described in
the *Methods* section. The mouse brain patterns include only excitatory-neuron synapses and are
based on approximately 3.2 million processed synaptic clefts of the MICrONS cortical column.

Although different Cartesian coordinate systems were originally used for human and
mouse samples, we maintained consistent anatomical orientations across all three figures. Again,

all synaptic clefts residing within a cortical layer were included in the construction of that layer's

synaptic pattern. The dimensionless normalized scale (0.5-1.0) is used for mouse synaptic patterns

in Figs.2,3,4. It differs from that used for the human data (0.3-1.0), providing finer contrast and

allowing smaller variations in the synaptic pattern to be resolved.

3. Discussion

3.1 Findings

Across all three cleft-extraction approaches, and across both species, synaptic clefts in

mammalian cortex were not randomly oriented. Instead, they exhibited a conserved, statistically

significant anisotropy (Figs.2–4). This is confirmed by randomization controls showing that the

result is not an artifact of finite sampling (Fig.7d). The principal orientation axis did not

consistently align with cortical columns but instead appeared to reflect location-specific cortical

anisotropy, as visible in Figs. 2b, 3b, 4b. Orientation patterns showed moderate layer-wise

variability and were bidirectional, consistent with the geometry of symmetric pre- and postsynaptic

membranes. Anisotropy was present in both species, but its magnitude and layer-wise

consistency were substantially greater in human association cortex than in mouse primary visual

cortex. The one exception was mouse L6, where artifactual distortions associated with column

curvature and increased myelination complicate interpretation.

3.2 Methodological robustness and limitations

The voxel-derived patterns of the H01 dataset (Figs.2d,3d,4d) exhibited the weakest variability

across cortical layers and a consistent strong minimum along the anterior–posterior (z) direction.

This minimum is not biological in origin but reflects a major methodological artifact. As explained

in Fig.8, the synapse classifiers used to construct the H01 dataset were trained on EM images

with high xy-resolution (4 nm) but substantially lower z-resolution (33 nm). Postsynaptic densities

oriented within xy-planes were therefore detected more readily than those aligned along z, and

this anisotropic detectability was inherited by the machine-learning classifiers. In contrast, the

mesh-touch method is less affected by this bias (Figs.2c,3c,4c). A separate, sample-compression

correction described in the *Materials and Methods* section was also applied.

In the MICrONS cortical column, which exhibits comparable imaging anisotropy, the

synaptic orientation patterns display analogous minima (Figs.2g,3g,4g). These patterns consist

primarily of a directional component superimposed on an artificial rectangular "dumbbell" feature

that is most pronounced in Layer 6. Imperfections in the manually trained synapse classifiers may

contribute to this artifact. Bending of the cortical column in L6 (Fig.6f) likely amplifies this

distortion, while additional segmentation challenges associated with the increased myelination
toward L6 may further accentuate it.
In both datasets, irregular sample geometries can potentially distort unweighted
directional patterns. To mitigate this, we applied a weighted directional analysis analogous to
those used in materials science and crystallography [23],[24] and, implicitly, in neuroscience [25],
where local density normalization is employed to reduce anatomical bias in orientation fields. The
Methods section details this methodology and provides a side-by-side comparison for the
MICrONS dataset, yielding results that closely match those shown in Figs.2–4. A similar
validation was performed for the H01 dataset, confirming that spatial sampling density does not
vary systematically with direction – that is, sampling is sufficiently isotropic.
Robustness was further tested by extracting multiple spherical subvolumes from different
cortical layers in both datasets and repeated the analyses. The resulting patterns were nearly
identical, indicating that the dominant features in our directional histograms – except for the z-
direction artifact in the H01 dataset – reflect genuine synaptic-cleft orientation structure rather
than geometry-specific sampling bias. Unavoidable residual effects may also include annotation
errors, meshing inaccuracies, and tissue deformations introduced during sample preparation that
could not be fully accounted for. Notably, across both datasets and all three extraction methods,
the reconstructed synaptic patterns converge toward a stable form even when relatively few clefts
(~25,000–50,000) are analyzed. As the number of included clefts increases, the overall
anisotropic patterns remain robust and change only modestly.
A further interpretive limitation is that the two datasets originate from different cortical
regions: human association cortex (middle temporal gyrus) and mouse primary visual cortex (V1).
The greater anisotropy in the human sample could therefore reflect regional specialization rather
than a species difference per se. Distinguishing these possibilities will require datasets from
matched cortical regions across species, which are not currently available at comparable scale
and resolution.
230
3.3 Possible mechanistic basis for cleft orientation
Cortical pyramidal neurons extend apical dendrites radially toward the pial surface and basal
dendrites tangentially within their home layer. En passant axonal boutons, which dominate over
terminal synapses in both datasets, contact these dendritic compartments at angles that are
constrained by local geometry. Because the normal to the synaptic cleft is determined by the
contact angle between the axonal process and its dendritic target, the distribution of cleft
orientations across a cortical volume will directly reflect the distribution of dendritic orientations,
which is far from isotropic.

At the molecular level, this geometric argument is supported by the known cell-type and

compartment specificity of trans-synaptic adhesion molecules. Neurexin splice isoforms are

differentially expressed across neuron types and brain regions in patterns that are temporally and

spatially regulated [9],[26]. Neuroligin isoforms are similarly segregated: Nlgn1 localizes

predominantly to excitatory synapses, Nlgn2 to inhibitory synapses, and Nlgn3 to subsets of both

[27]. Because different neuroligin isoforms are concentrated on different dendritic compartments,

and because these compartments have preferred orientations relative to cortical axes, the

aggregate distribution of cleft orientations will inherit the geometric bias of the dendritic arbor. No

field-sensitivity or orientation-specific molecular signaling is required; the anisotropy emerges

from the spatial distribution of where synapses form.

One supporting line of evidence is recent work on the mouse brain synaptome [28] which

demonstrated that synapses are not uniform but exhibit region- and layer-specific patterns of

molecular composition and morphology. Zhu and colleagues [28] showed that these spatial

gradients in synapse diversity shape network topology and functional connectivity, providing

evidence that synaptic properties contribute to mesoscale circuit organization. Our study extends

this principle by identifying orientation as an additional dimension of synaptic architecture.

Whereas Zhu et al. highlighted diversity in molecular identity and spine morphology, our results

indicate that the geometry of the synaptic cleft itself is patterned relative to cortical axes.

Together, these findings suggest that synaptic microstructure may contribute to cortical

computation along multiple orthogonal dimensions: molecular diversity, morphological

specialization, and spatial orientation. This framework positions synaptic architecture as a

multiscale substrate for evolutionary and functional differences in cortical processing.

The anisotropy was present in both mouse and human cortex, establishing it as a

conserved rule across species. However, the magnitude and layer consistency of the bias were

greater in human than in mouse cortex. This difference aligns with broader comparative

neuroanatomy: human cortex is thicker, and pyramidal neurons possess more extensive dendritic

arbors, with apical and basal branches orders of magnitude longer than in mouse, while human

neurons also receive substantially more synaptic inputs per dendritic segment [29],[30],[31].

We hypothesize that stronger anisotropy in human association cortex may reflect (i)

evolutionary scaling of synaptic geometry to support greater integration across cortical columns

and/or; (ii) specialization of the brain region in which it occurs – as pointed out, for example, in

Barbas et al 2022 [32].

Recent cross-species imaging work reinforces the first interpretation. Liu et al.

demonstrated that cortical architecture exhibits species- and layer-specific trajectories [33]: in

both humans and mice, deep layers thin with age while layer IV becomes relatively thicker and

more myelinated, but the magnitude and functional implications of these changes differ between

species. Their findings underscore that laminar microstructure is not fixed but evolves in ways

that support distinct modes of integration. In parallel, our observation that synaptic clefts are more

strongly and consistently anisotropic across layers in human cortex suggests that orientation bias

is another measurable property along which cortical microarchitecture scales with integrative

demands. Taken together, these results argue that both laminar architecture and synaptic

orientation anisotropy represent species-sensitive structural adaptations that enhance cross-

columnar and cortico–cortical communication.

On the other hand, there could be significant differences in the columnar and laminar

organizations between different brain regions. Barbas et al. noted that feedforward connections,

where the signal travels from earlier to later processing cortices, have a vertical organization. This

is in contrast with feedback connections, where the signal travels from later to earlier processing

areas have a more laminar organization. This could also be a possible explanation for the

difference in anisotropy observed in the human and mouse data, as the two regions (association

cortex vs primary sensory cortex) may have a preference in the direction which the signal travels

[32].

Several factors may contribute to the observed anisotropy. At the cellular level, the

expanded dendritic arbors of human pyramidal neurons increase the likelihood of tangentially

aligned synaptic contacts. Developmental guidance signals and structural scaffolds, including

radial glia and molecular gradients [34],[35], may impose constraints that orient presynaptic and

postsynaptic membranes relative to cortical axes. At the tissue level, interactions with

extracellular matrix could reinforce preferred orientations. Together, these cellular and

developmental mechanisms may embed orientation bias as a byproduct of cortical architecture.

Orientation bias may enhance lateral signal integration across minicolumns and modules

and vice versa. In association cortex, such integration is essential for combining multimodal

inputs and supporting higher-order cognition. By aligning synapses preferentially in tangential

planes, cortical microcircuits could potentially facilitate cross-columnar communication while

reducing redundancy along the radial axis. This aligns with evidence that human cortex favors

cortico-cortical connectivity emerging from supragranular layers [17],[18]. At a broader scale,

synaptic anisotropy could shape the propagation of activity across cortical networks, influencing

oscillatory dynamics and inter-areal coordination.

Bodor et al. (2025) [36], analyzing the same MICrONS dataset, showed that layer 5 thick-

tufted pyramidal neurons (extratelencephalic cells) in mouse visual cortex predominantly form

local synapses onto inhibitory interneurons, which reciprocally inhibit them, while their longer-

range projections target excitatory populations such as L6 pyramidal neurons and L5-IT cells.

This architecture implies local competition and long-range coordination, consistent with predictive

coding and the generation of gamma oscillations. Notably, Bodor et al. highlight that such local

inhibitory targeting differs from cat and monkey cortex, where excitatory neurons more commonly

synapse onto other local excitatory cells. These findings are in line with our observation that

synaptic cleft orientation anisotropy is weaker and less consistent in mouse cortex than in human

cortex: both results point to a circuit-level strategy in rodents that emphasizes inhibitory balance

locally, in contrast to the more integrative excitatory connectivity characteristic of higher order

animals.

In conclusion, our analyses suggest that synaptic clefts in mammalian cortex are not

randomly oriented but may exhibit a conserved anisotropy at the mesoscale. This orientation bias

was evident in both human and mouse datasets, yet it appeared stronger and more consistent in

human association cortex. We propose that anisotropy in cleft orientation may reflect the

geometry of axonal and dendritic arbors and the developmental scaffolds that shape them, and

that its amplification in humans could relate to the expanded integrative capacity of association

cortex. Furthermore, the collective alignment of synapses may influence how cortical tissue

interacts with endogenous field potentials and with externally applied brain stimulation.

At the same time, we emphasize that the present findings must be interpreted with

caution. Both datasets are subject to methodological artifacts, resolution anisotropies, and

potential biases in segmentation and classification. The observed anisotropy may therefore, at

least in part, reflect technical or sampling effects rather than an intrinsic biological property.

Definitive validation will require independent datasets acquired at truly isotropic voxel resolution

(~10×10×10 nm), improved AI-based reconstructions with minimized geometric distortion, and

targeted physiological and computational studies to link structure with function.

Despite these limitations, the possibility that a conserved, orientation-dependent

organization exists across mammalian synapses is intriguing. By revealing a candidate

mesoscale feature of synaptic architecture and by framing it explicitly within its methodological

constraints this work aims to motivate deeper experimental and theoretical investigations into how

microstructural geometry contributes to cortical computation and neuromodulation.

4. Materials and Methods

4.1 Synaptic cleft extraction and processing for H01 dataset

Chemical synaptic clefts were extracted from a micrometer thick slab of human cortex (H01) from

the anterior part of the middle temporal gyrus, with a total volume of just over one cubic

millimeter. The sample was removed to gain access to an epileptic focus in the underlying

hippocampus [10]. In its longest dimension, the sample spans all cortical layers (6 through 1) in

the x-direction (from deep to superficial) but is relatively narrow (~175 µm) in the anterior-

posterior (z) direction. By light microscopy-based neuropathological examination, the sample was

deemed normal [10]. To align the cortical direction with the x-axis, the entire sample was rotated

by 35° about the z-axis, resulting in the orientation shown in Fig.1a of the main text.

Mesh touch method for extracting cleft directions

The starting point of our analysis is the synaptic repository [11] of the H01 sample publicly

available via Google Cloud storage [12]. Two parts of the repository were used: (i) surface

membrane meshes created by a Google meshing team in the STL format for individual pyramidal

excitatory postsynaptic neurons with adjacent presynaptic neuronal (predominantly axonal)

processes (as shown in Fig.1b,c) and; (ii) the corresponding CSV synaptic tables (one such table

per postsynaptic neuron) that list pre- and postsynaptic surface mesh IDs, as well as the

coordinates for each side (side center) of the synaptic cleft. In this analysis, we did not

differentiate between excitatory and inhibitory synapses for the membrane surface meshes

although the dominant axodendritic synapses are mostly excitatory.

Based on the presynaptic and postsynaptic center positions, and using manipulations

with the available triangular surface meshes, we have extracted the membrane topology near the

cleft within approximately 1 µm from its center as shown in Fig.1l). We have further extracted

both pre- (yellow in Fig.1l) and postsynaptic (magenta in Fig.1l) membrane interfaces. Those are

defined here as separated from each other by at most 100 nm. A test was done when reducing

this value to 75 nm and very similar results were obtained. The corresponding meshes were

further refined by 1:4 manifold triangular subdivision followed by surface-preserving Taubin

smoothing [37].

In this way, we processed 944 pyramidal neurons of the repository which resulted in a

total of 1,566,358 proofread and postprocessed synaptic clefts. The synaptic clefts themselves

are mostly located between layers L2 to L6 (L1: 56,649; L2: 482,282; L3: 559,968; L4: 165,621;

L5: 172,241; L6: 99,995; the rest is in white matter).

As shown in Fig.1m, for every yellow presynaptic membrane interface, we found the total

average outer normal vector by averaging the normal vectors of the respective triangular facets

(weighted by facet areas) of the presynaptic interface. This average normal vector (magenta

arrow in Fig.1m) has been chosen as the (pre) synaptic orientation of the cleft or its primary

directional axis. This axis denotes the approximate direction in which the neurotransmitters travel

from pre- to postsynaptic processes.

Voxel method for extracting cleft directions

One potential limitation of the previous method is that it includes only neurons whose somas are

located within the sample volume, thereby excluding orphaned fragments of the neuropil

contained inside the dataset. To address this issue, we additionally analyzed the excitatory

synaptic data provided in voxel format, which encompass all putative excitatory synapses

(111,272,315 in total) within the sample. The same Google Cloud repository also contains the

voxel representation of synaptic clefts spanning the entire sample volume [12]. Unlike the mesh-

based approach, the voxel representation includes synapses originating from all neuronal

processes, not only from pyramidal neurons.

For our analysis, we used two available volumetric renderings decompressed from the

Neuroglancer format [38]: (i) *Volumetric E/I Synapses* that contains a voxel representation of the

entire synapse (both pre- and postsynaptic sites, but the voxels are not discriminated between

the two types, shown in red in Fig.1n) and; (ii) *Incoming excitatory synapses* that contains voxel

representation of the excitatory postsynaptic density or PSD (shown in blue in Fig.1n). Both

volumetric renderings have identical dimensions (515,892×356,400×5,249 voxels) and

resolutions (8 nm per voxel in the x and y directions and 33 nm per voxel in the z-direction).

The entire volume was originally divided into ~6.4 million regions of interests, each with a

size of 1,000×1,000×150 voxels. For each region, voxels belonging to the individual synapses

were separately extracted using the *ind2sub* and *bwconncomp*-based clustering method in

MATLAB. After that, we analyzed every individual synaptic assembly separately. Namely, for

every presynaptic voxel, we found all postsynaptic voxels within a 40 nm distance. In total,

111,272,315 excitatory synapses (all synapses available) were processed.

To ensure that no synaptic neighborhood was artificially truncated at the borders of our

1,000×1,000×150-voxel blocks, each chunk was extended by a volumetric "margin" whose

thickness in X, Y, and Z matched the maximum synapse-to-synapse search radius (40 nm

converted to voxels). In practice, for every region we first loaded an additional pad of voxels on all

six faces, built connected-component labels using *ind2sub* and *bwconncomp*-based clustering

and distance searches over the full padded region, and only after computing each synapse's

orientation did we discard any result whose centroid lay within that padding. This approach

guaranteed that every potential partner voxel within one search radius (40 nm) of a presynaptic

site was present in the analysis – even at chunk boundaries – while at the same time preventing

any synapse from being counted more than once.

By filtering out synapses whose centroids fell in the padded margins, we retained exactly

one fully correct orientation vector per synapse and avoided artifacts that would otherwise arise

when a search sphere is clipped by an artificial cube face. The result is a seamless tiling of

orientation vectors across the entire volume, with no edge-effects and no overlap between

adjacent blocks. Crucially, although we discard every synapse whose centroid lies in the padded

"margin for each processed region," no synapse is ever lost. The reasons being:

(i) Every synapse appears fully inside at least one padded block. By extending each

1,000×1,000 × 150-voxel chunk by a "margin" equal to the 40 nm search radius

on all six faces, even a synapse whose true centroid lies exactly on the boundary

of one central block will lie well inside the padded volume of its neighbor. In that

neighboring block the synapse never touches the padded-minimum faces and so

is processed with its full 40 nm neighborhood intact.

(ii)   We record only synapses whose centroids lie in the original (unpadded) block.

After computing every orientation vector in the padded region, we tested each synapse's

global centroid against the global coordinate ranges of each region and saved only those that fall

inside. Since a given centroid can lie in exactly one block's central region, every synapse is

committed exactly once – and only once.

Delaunay triangulation-based method for extracting cleft directions from voxel data

Our method is illustrated in Fig.5a–c. Extraction of synaptic cleft orientation from the original

voxelized synaptic data (Fig.5a) begins by identifying the boundary between the two voxel sets

shown in Fig.5b. This boundary is then triangulated (Fig.5c), after which the normal vectors of the

resulting triangular facets are computed and subsequently averaged to determine the cleft

direction. In cases where Delaunay triangulation performs poorly due to complex synapse

geometries, partial triangulation is applied to multiple subsets of the synaptic boundary. This

procedure was used to extract the orientation of all synaptic clefts analyzed in this study.

The mesh-touch method may be more accurate than the voxel-based method

In most cases, the two methods produce reasonably consistent results, as illustrated in Fig.5d,e.

However, there are instances in which the voxel-based method fails to capture portions of a

synapse, as shown in Fig.5f. Consequently, the estimated cleft direction (indicated by magenta

and green arrows in Fig.5f) deviates between the two methods. In other cases, the synaptic cleft

is only partially identified or misclassified altogether (Fig.5g). The exact frequency of such

occurrences remains undetermined.

4.2 Sample compression correction for the H01 dataset

It is important to consider that artifacts introduced during H01 sample preparation may influence

the measured synaptic orientations. The geometry of the H01 dataset originates from multibeam

scanning electron microscopy (ATUM–mSEM) rather than transmission electron microscopy

(TEM). In the ATUM workflow, each 30–33 nm section is cut, mounted on Kapton tape, and

scanned using 61 electron beams at 4 × 4 nm pixel resolution over an area of approximately 4.5

mm². Mechanical pickup during this process anisotropically flattens each section: the authors of

[10] reported nearly 28% compression along the axis perpendicular to the knife edge (x-axis) and

only ~0.3 % along the axis parallel to it (y-axis) (see Supplementary Information of Shapson-Coe

et al., 2024 [10]).

By contrast, the serial-section TEM used for the MICrONS dataset images entire sections

in transmission. In this technique, dimensional distortions are typically isotropic—on the order of

2–5 % shrinkage—arising from resin polymerization and drying, rather than the directional

flattening inherent to tape-collect SEM [39],[40].

To correct for this effect (not accounted for in the publicly available datasets), we

expanded the H01 sample by 28% along the x-axis and correspondingly reduced the x-

components of all synaptic normal vectors by the same proportion.

4.3 Synaptic cleft extraction and processing for MICrONS dataset

Using synaptic tables

We analyzed the *Minnie65* subset of the MICrONS dataset [6], representing the primary visual

cortex of a mouse. This subset contains a cortical column comprising 1,356 neurons and

measuring approximately 200 μm in diameter. The column spans all cortical layers and is shown

in Fig.6 (axons – cyan; apical dendrites – magenta; basal dendrites – blue; somas – orange). It

occupies a substantial portion of the overall sample volume and exhibits a slight curvature in layer

6. The *Minnie65* dataset was accessed via CAVEclient (Python 3.11.9), using materialization

v1181, which was the most recent public release at the time of analysis. For the present study,

we selected all excitatory neurons (1,097 in total) within this cortical column.

The synapse table (synapses_pni_2) includes pre- and post-synaptic side point positions.

This information was used for determining effective synaptic cleft orientations (and thicknesses if

necessary). Additionally, the manual cell types (V1 column) table

(allen_v1_column_types_slanted_ref) was referred to for determining detailed cell type

information necessary for filtering and categorizing downloaded cell data. The root IDs were

extracted and organized by layer into the four categories (L2/3, L4, L5, and L6). These extracted

root IDs were used to filter the synapse table query by specifying them as post_ids. Cell data

from the synapse query were downloaded as csv files and organized by layer. The total number

of processed clefts is ~3.2 million (1.4 million in L2/3, 0.6 million in L4, 0.4 million in L5, and 0.8

million in L6). To determine the orientation of each cleft, unit normal vectors were constructed

pointing from the pre- to the post-synaptic side. No differentiation between excitatory and

inhibitory synapses was made.

Layer-by-layer separation of synaptic clefts

Approximate layer boundaries (Fig.6a) were determined based on local nuclear features [40]. In

this region of the visual cortex, layers 2 and 3 are not well demarcated and are therefore

collectively referred to as Layers 2/3. The generated meshes delineate the borders between

Layer 1, Layers 2/3, Layer 4, Layer 5, Layer 6, and the white matter. Manually drawing layer

boundaries in approximately the correct anatomical locations was considered sufficient for the

purposes of this analysis. All synapses were then assigned to their respective layers according to

spatial position, and three-dimensional synaptic orientation patterns were subsequently

constructed for each layer as described in the Results section.

4.4 Weighted versus unweighted synaptic patterns

To minimize the effect of the sample shape and other inhomogeneities, we applied the weighted

directional analysis used in materials science and crystallography [23],[24] and, implicitly, in

neuroscience [25]) where the authors apply local density normalization in histological mapping to

avoid anatomical bias in orientation fields. In essence, the previous definition of the synaptic

pattern was modified by

$$Weighted\ Directional\ Contribution = \sum_{i=1}^{N} \frac{n_i}{\rho(x_i)}, \quad \rho(x_i) = \sum_{j=1}^{N} exp\left(-\frac{\|x_i - x_j\|^2}{2\sigma^2}\right) \tag{1}$$

503   Here, x_i are positions of N synapses, n_i are the cleft orientation vectors, $\rho(x_i)$ is the local spatial

504   density of the synaptic clefts which is found as a probability density estimate. This method in

505   particular corrects for a non-uniform sample geometry.

506   The results presented in Figs. 2, 3, and 4 of the main text were re-computed using the

new definition (Eq. 1) with a moderate standard deviation of 0.3, and the resulting patterns were

compared across datasets. The outcomes were broadly similar for both samples. Figs. 7b,c

provide a side-by-side comparison of the MICrONS patterns obtained using the unweighted

definition of the synaptic pattern (from Figs. 2–4) and the weighted definition based on Eq. (1).

The weighted formulation is intended to compensate for potential non-uniformities in sample

geometry. Considering the top view of the MICrONS cortical column (Fig.7a) and the

corresponding two-dimensional projections, only minor differences are observed between the two

approaches—except perhaps in Layer 5, where the discrepancy is somewhat more noticeable,

though the overall pattern shape remains consistent. Both methods thus yield comparable results,

indicating that the spatial sampling density does not vary systematically with direction; in other

words, sampling is sufficiently isotropic that a computationally intensive weighting correction is

not required.

4.5 Randomized controls

As an additional control, we assigned random orientations to all synaptic clefts, drawing each direction from a three-dimensional standard normal distribution (mean = 0, standard deviation = 1). This procedure was applied to both the mesh-touch human synaptic patterns and the mouse synaptic patterns. The resulting distribution for the mouse cortical column is shown in Fig.7d. In every such randomized dataset, the resulting patterns were nearly omnidirectional, as illustrated in Fig.7d. The corresponding human results were similar.

4.6 Known (major) artifact(s)

Fig.8 explains why the apparent strong minimum in the anterior–posterior (z) direction observed in the voxel-based patterns of the H01 sample (Figs. 2d, 3d, and 4d of the main text) arises from methodological artifacts rather than a genuine biological feature. Fig.8a shows three consecutive H01 slices containing a synapse oriented primarily within the xy-plane (highlighted by yellow circles). This synapse is readily detected during manual labeling and classifier training on electron microscopy cross-sections, which have high in-plane ($x - y$) resolution of 4 nm but lower axial (z) resolution of 33 nm. In contrast, Fig.8b shows three consecutive H01 slices containing a synapse oriented predominantly along the z-axis. Such a synapse is often visible in only a single z-slice, resulting in poor detection and contributing to their apparent underrepresentation.

A comparable artifact may also occur in the MICRONS cortical column, which was imaged at a similarly anisotropic resolution. The synaptic patterns in Figs.2g,3g,4g of the main text and in Fig.7b,c display corresponding minima that align closely with the Cartesian axes.

Acknowledgments

SNM has received support from NIH grant R01MH130490, R01EB035484; BD has received support from NIH grant U24NS120053; ZDD has received support from NIMH grant ZIAMH002955; HL has received support NIDA grant ZIADA000638; MB has received support from NIDA grant UG3DA048502, NIGMS grant T34GM137858, T32GM136499, NINDS grant R01NS112996, R01NS101362; DB has received support from NIMH grant UG3MH123386, NINDS grant U19NS104653, U24NS109102, NIBIB grant U01EB026996; BR has received support from NIBIB grant P41EB030006.

References

[1]   Mountcastle VB. The columnar organization of the neocortex. Brain. 1997 Apr;120 (Pt 4):701-22. doi: 10.1093/brain/120.4.701.

[2]   Douglas RJ, Martin KA. Neuronal circuits of the neocortex. Annu Rev Neurosci. 2004;27:419-51. doi: 10.1146/annurev.neuro.27.070203.144152.

[3]   Rakic P. Confusing cortical columns. Proc Natl Acad Sci U S A. 2008 Aug 26;105(34):12099-100. doi: 10.1073/pnas.0807271105. Epub 2008 Aug 20. Erratum in: Proc Natl Acad Sci U S A. 2008 Sep 30;105(39):15220.

[4]   Harris KD, Shepherd GM. The neocortical circuit: themes and variations. Nat Neurosci. 2015 Feb;18(2):170-81. doi: 10.1038/nn.3917.

[5]   Yuste R. From the neuron doctrine to neural networks. Nat Rev Neurosci. 2015 Aug;16(8):487-97. doi: 10.1038/nrn3962.

[6]   Turner NL, Macrina T, Bae JA, Yang R, Wilson AM, Schneider-Mizell C, Lee K, Lu R, Wu J, Bodor AL et al. Reconstruction of neocortex: Organelles, compartments, cells, circuits, and activity. Cell. 2022 Mar 17;185(6):1082-1100.e24. doi: 10.1016/j.cell.2022.01.023.

[7]   GRAY EG. Axo-somatic and axo-dendritic synapses of the cerebral cortex: an electron microscope study. J Anat. 1959 Oct;93(Pt 4):420-33. PMID: 13829103; PMCID: PMC1244535.

[8]   Peters, A., Palay, S. L., Webster, H. The Fine Structure of the Nervous System. Neurons and their Supporting Cells. 3rd edition. Oxford University Press: New York, 1991.

[9]   Gomez AM, Traunmüller L, Scheiffele P. Neurexins: molecular codes for shaping neuronal synapses. Nat Rev Neurosci. 2021 Mar;22(3):137-151. doi: 10.1038/s41583-020-00415-7.

[10] Shapson-Coe A, Januszewski M, Berger DR, Pope A, Wu Y, Blakely T, Schalek RL, Li PH, Wang S, Maitin-Shepard J, et al. A petavoxel fragment of human cerebral cortex reconstructed at nanoscale resolution. Science. 2024 May 10;384(6696):eadk4858. doi: 10.1126/science.adk4858.

[11] H01 Human brain sample repository. Accessed 10/2024. Online: https://h01-release.storage.googleapis.com/data.html.

[12] H01 Google Cloud Storage. Accessed 10/2024. Online: gs://h01-release/data/.

[13] MICrONS Consortium. Functional connectomics spanning multiple areas of mouse visual cortex. Nature. 2025 Apr;640(8058):435-447. doi: 10.1038/s41586-025-08790-w.

[14] Ding Z, Fahey PG, Papadopoulos S, Wang EY, Celii B, Papadopoulos C, Chang A, Kunin AB, Tran D, Fu J, et al. Functional connectomics reveals general wiring rule in

mouse visual cortex. Nature. 2025 Apr;640(8058):459-469. doi: 10.1038/s41586-025-
08840-3.

[15] MICrONS Consortium. A functional connectome containing 200,000 cells, 75,000
neurons with physiology, and 523 million synapses. Online: https://www.microns-
explorer.org/cortical-mm3.

[16] Wang XJ, Kennedy H. Brain structure and dynamics across scales: in search of rules.
Curr Opin Neurobiol. 2016 Apr;37:92-98. doi: 10.1016/j.conb.2015.12.010. Epub 2016
Feb 8.

[17] DeFelipe, J. and Jones, E. G. Neocortical microcircuits. Handbook of Brain Microcircuits,
Shepherd, G. and Grillner, S. Eds. Oxford University Press. 2010.

[18] Markov NT, Vezoli J, Chameau P, Falchier A, Quilodran R, Huissoud C, Lamy C, Misery
P, Giroud P, Ullman S, et al. Anatomy of hierarchy: feedforward and feedback pathways
in macaque visual cortex. J Comp Neurol. 2014 Jan 1;522(1):225-59. doi:
10.1002/cne.23458.

[19] Fröhlich F, McCormick DA. Endogenous electric fields may guide neocortical network
activity. Neuron. 2010 Jul 15;67(1):129-43. doi: 10.1016/j.neuron.2010.06.005.

[20] Reato D, Rahman A, Bikson M, Parra LC. Low-intensity electrical stimulation affects
network dynamics by modulating population rate and spike timing. J Neurosci. 2010 Nov
10;30(45):15067-79. doi: 10.1523/JNEUROSCI.2059-10.2010.

[21] Radman T, Su Y, An JH, Parra LC, Bikson M. Spike timing amplifies the effect of electric
fields on neurons: implications for endogenous field effects. J Neurosci. 2007 Mar
14;27(11):3030-6. doi: 10.1523/JNEUROSCI.0095-07.2007.

[22] Balanis CA. Antenna Theory. Analysis and Design. 4th Ed. New York: Wiley, 2016, 1104
p., ISBN: 1118642066.

[23] Wenk, HR, Van Houtte P. Texture and anisotropy. Reports on Progress in Physics,
2004, 67(8), 1367–1428. https://doi.org/10.1088/0034-4885/67/8/R03.

[24] Kocks UF, Tomé CN, Wenk HR. Texture and anisotropy: Preferred orientations in
polycrystals and their effect on materials properties. Cambridge University Press. 1998.
https://doi.org/10.1017/CBO9780511755523.

[25] Zhou Z, Liu Z, Chen S, Zhang Z, Liu H, Zhang D, Lin L, Yuan J, Ren M, Qiu L, Xu T,
Zeng H. Functional connectome and microarchitecture of the mouse brain. Nature. 2020
Oct;588(7837):103–107. doi:10.1038/s41586-020-2969-6.

[26] Südhof TC. Synaptic Neurexin Complexes: A Molecular Code for the Logic of Neural
Circuits. Cell. 2017 Nov 2;171(4):745-769. doi: 10.1016/j.cell.2017.10.024.

[27] Bemben MA, Shipman SL, Nicoll RA, Roche KW. The cellular and molecular landscape
623   of neuroligins. Trends Neurosci. 2015 Aug;38(8):496-505. doi:
624   10.1016/j.tins.2015.06.004.

625   [28] Zhu F, Cizeron M, Qiu Z, Benavides-Piccione R, Kopanitsa MV, Skene NG, Koniaris B,
626   DeFelipe J, Fransén E, Komiyama NH, Grant SGN. Architecture of the Mouse Brain
627   Synaptome. Neuron. 2018 Aug 22;99(4):781-799.e10. doi:
628   10.1016/j.neuron.2018.07.007.

629   [29] Elston GN. Cortex, cognition and the cell: new insights into the pyramidal neuron and
630   prefrontal function. Cereb Cortex. 2003 Nov;13(11):1124-38. doi:
631   10.1093/cercor/bhg093.

632   [30] Galakhova AA, Hunt S, Wilbers R, Heyer DB, de Kock CPJ, Mansvelder HD,
633   Goriounova NA. Evolution of cortical neurons supporting human cognition. Trends Cogn
634   Sci. 2022 Nov;26(11):909-922. doi: 10.1016/j.tics.2022.08.012.

635   [31] Loomba S, Straehle J, Gangadharan V, Heike N, Khalifa A, Motta A, Ju N, Sievers M,
636   Gempt J, Meyer HS, Helmstaedter M. Connectomic comparison of mouse and human
637   cortex. Science. 2022 Jul 8;377(6602):eabo0924. doi: 10.1126/science.abo0924.

638   [32] Barbas H, Zikopoulos B, John YJ. The inevitable inequality of cortical columns. Front
639   Syst Neurosci. 2022 Sep 20;16:921468. doi: 10.3389/fnsys.2022.921468. PMID:
640   36203745; PMCID: PMC9532056.

641   [33] Liu P, Doehler J, Henschke JU, Northall A, Knaf-Serian A, Loaiza-Carvajal LC, Budinger
642   E, Schwarzkopf DS, Speck O, Pakan JMP, Kuehn E. Layer-specific changes in sensory
643   cortex across the lifespan in mice and humans. Nat Neurosci. 2025 Sep;28(9):1978-
644   1989. doi: 10.1038/s41593-025-02013-1.

645   [34] Casingal CR, Descant KD, Anton ES. Coordinating cerebral cortical construction and
646   connectivity: Unifying influence of radial progenitors. Neuron. 2022 Apr 6;110(7):1100-
647   1115. doi: 10.1016/j.neuron.2022.01.034.

648   [35] Whitford KL, Dijkhuizen P, Polleux F, Ghosh A. Molecular control of cortical dendrite
649   development. Annu Rev Neurosci. 2002;25:127-49. doi:
650   10.1146/annurev.neuro.25.112701.142932.

651   [36] Bodor AL, Schneider-Mizell CM, Zhang C, Elabbady L, Mallen A, Bergeson A, Brittain D,
652   Buchanan J, Bumbarger DJ, Dalley R, et al. The synaptic architecture of layer 5 thick

tufted excitatory neurons in mouse visual cortex. Nat Neurosci. 2025 Aug;28(8):1704-1715. doi: 10.1038/s41593-025-02004-2.

[37] Täubin G. Estimating the tensor of curvature of a surface from a polyhedral approximation. Technical Report. RC-19860, IBM Research, December 1994.

[38] Maitin-Shepard J. Precomputed sharded format – Neuroglancer documentation. GitHub repository, Google; 2022. github.com/google/neuroglancer/blob/master/src/datasource/precomputed/sharded.md

[39] Mollenhauer HH. Artifacts caused by dehydration and epoxy embedding in transmission electron microscopy. Microsc Res Tech. 1993 Dec 15;26(6):496-512. doi: 10.1002/jemt.1070260604. PMID: 8305727.

[40] Ströh S, Hammerschmith EW, Tank DW, Seung HS, Wanner AA. In situ X-ray-assisted electron microscopy staining for large biological samples. Elife. 2022 Oct 20;11:e72147. doi: 10.7554/eLife.72147. PMID: 36263931; PMCID: PMC9665849.

[41] Neuroglancer. Online: https://spelunker.cave-explorer.org/#!middleauth+https://global.daf-apis.com/nglstate/api/v1/5810339900817408

     Figures and Tables

Figure 1. a) Topology of the H01 sample. The x-axis is directed from L6 to L1. b) Postsynaptic
excitatory pyramidal neuron of the sample (blue); associated presynaptic processes (red). c)
Close up view of the apical dendrite branch located top right in b). d,e,f,g) Axodendritic excitatory
synapses of the excitatory neurons: en passant boutons and synapses e,f,g) and terminal
synapse d). h) Axosomatic synapse with part of the soma membrane. i,j) Axo-dendritic shaft
synapses. k) Rare putative axoaxonic synapses, possible false detection. l,m) Estimating synaptic
cleft direction using the mesh touch method. Red normal vectors of the presynaptic membrane
(yellow facets) are averaged and result in the effective directional vector (magenta). n) Estimating
synaptic cleft direction using voxel data from labeling presynaptic terminals with clustered
synaptic vesicles. A boundary between pre and postsynaptic voxels is extracted and subjected to
Delaunay triangulation. o) Overlap of voxel and mesh touch synaptic representations (wherever
possible).

a) L1 L2 L3 L4 L5 L6

Anterior-posterior z

Superficial x Deep

b) L1 L2 L3 L4 L5 L6

c) XZ plane view of H01 synaptic pattern (perpendicular to layers) obtained with the mesh touch method

d) XZ plane view of H01 synaptic pattern (perpendicular to layers) obtained from the voxel data

e) z Anterior

L2/3 L4 L5 L6

Superficial Deep y

f) L2/3 L4 L5 L6

g)

Figure 2. a) xz-View (side view) of the H01 sample with cortical layers labeled from L1 (left, dark
blue) to L6 (right, orange) and cross-sectional planar rectangles for each layer. b) Neuroglancer
slices taken along the cross-sectional planar rectangles shown in panel a). c) Layer-by-layer
mesh-based synaptic patterns in the same xz-plane. d) Layer-by-layer voxel-based synaptic
patterns in xz-plane. e) yz-View (side view) of the MICrONS sample with cortical layers labeled
from L2/L3 (left, red) to L6 (right, yellow), and representative cross-sectional planar rectangles for
each layer. f) Neuroglancer slices taken along the cross-sectional planar rectangles shown in
panel e). g) Layer-by-layer synaptic patterns of the MICrONS sample in the same yz -plane. For
both samples, different Cartesian coordinate systems were originally used, but the anatomical
directions in the panels are the same.

Top view of H01 synaptic pattern (into layers) obtained with the mesh touch method

Top view of H01 synaptic pattern (into layers) obtained from the voxel data

Figure 3. a) yz-View of the H01 sample with labeled cortical layers from L1 (front, dark blue) to

L6 (back, orange) and cross-sectional planar rectangles for each layer. b) Neuroglancer slices in

the cross-sectional planar rectangles from panel a). c) Layer-by-layer mesh-based synaptic

patterns in the same yz-plane. d) Layer-by-layer voxel-based synaptic patterns in the yz-plane. e)

xz -View of the MICrONS sample with labeled cortical layers from L2/L3 (left, red) to L6 (right,

yellow) and cross-sectional planar rectangles. f) Neuroglancer slices in the cross-sectional planar

rectangles from panel e). g) Layer-by-layer synaptic patterns of the MICrONS sample in the same

xz-plane

Figure 4. a) xy -View of the H01 sample with labeled cortical layers from L1 (top, dark blue) to L6

(bottom, orange) and cross-sectional planar rectangles for each layer. b) Neuroglancer slices in

the cross-sectional planar rectangles from panel a). c) Layer-by-layer mesh-based synaptic

patterns in the same xy-plane. d) Layer-by-layer voxel-based synaptic patterns in the same xy-

plane. e) xy-View of the MICrONS sample with labeled cortical layers from L2/L3 (left, red) to L6

(right, yellow) and cross-sectional planar rectangles where the Neuroglancer cell maps will be

plotted) for each layer. f) Neuroglancer slices in the cross-sectional planar rectangles from panel

e). g) Layer-by-layer synaptic patterns of the MICrONS sample in the same xy-plane.

Figure 5. a,b,c) Illustration of Delaunay-triangulation method used for extracting cleft directions from original voxel data in a). In b), a boundary between two voxel sets is extracted, which is further triangulated in c). Normal vectors of the triangular facets are then found and finally averaged. d,e,f,g) Correlations between voxel and mesh touch synapse representations. Frequently, both methods agree reasonably well as in d,e). However, there are many cases where the voxel-based method is missing a part of the synapse as in f). As a result, the cleft direction shown by the respective arrow becomes different. In certain cases, the cleft is identified by the voxel method totally incorrectly as in g).

Figure 6. a) Side view of the IARPA Phase III MICrONS V1 cortical column showing 1,097 excitatory neurons overlaid on a cross-section of the *Minnie65* cortical volume at $z = 937\ \mu m$. Cell membranes are shown in black, the vasculature in red, and somas in white. Approximate layer boundaries are indicated by orange lines. Axons are shown in cyan, apical dendrites in magenta, and basal dendrites in blue. The table on the left quantifies the number of cells with somas located within each layer and the number of synaptic clefts physically contained in that layer. b,c,d) Separate visualizations of apical dendrites (b), basal dendrites (c), and somas (d). e,f) Illustration of cortical column curvature in the yz-plane, with somas shown in orange.

Figure 7. a) Top view (into layers) of IARPA Phase III MICrONS V1 cortical column with 1,097

excitatory neurons. Axons – cyan, apical dendrites – magenta, basal dendrites – blue. The

average length of the axonal arbor per neuron is 4.9 mm. The column overlaps with the original

Minnie 65 sample's cross-section at y=600 µm where cell membranes are shown in black, the

blood network – in red, and the somas are white. b). Top view (xz -plane into the cortical column)

of the same sample. c, d) Side-by side comparison between the patterns for the MICrONS

sample obtained with the unweighted definition of the synaptic pattern (from Figs. 2,3,4) and

using the weighted definition from Eq. (1), respectively. The same xz-view is displayed. e) The

741   same xz-view of synaptic patterns obtained when all synaptic cleft directions are assigned

742   random values chosen from the 3D standard normal distribution with mean zero and standard

743   deviation one.

Case 1: Synapse with the normal vector primarily in the xy-plane

Case 2: Synapse with the normal vector perpendicular to the xy-plane

746   **Figure 8.** H01 synapses oriented primarily within the xy-plane (highlighted by yellow circles) are

747   readily detected during manual labeling and classifier training on electron microscopy cross-

748   sections, which have high in-plane $(x - y)$ resolution of 4 nm but lower axial (z) resolution of 33

749   nm. (b) In contrast, H01 synapses oriented predominantly along the z-axis are often visible in only

750   a single z-slice, resulting in poor detection and contributing to their apparent underrepresentation.