Polyploid cardiomyocytes define disease-specific transcriptional states in the mammalian heart

Authors: Paul Kiessling[1#], Mehdi Joodaki[2,3#], Daryna Pikulska[1#], Emilia Scheidereit[1#], Giulia Cesaro[4#], Mayra Ruiz[2,3], Damin Kühn[3,5], Kai Peng[2,3], Osman Goni[1], Sebastian Foecking[1], Samaneh Samiei[1], Xian Liao[1], Kai Li[1], Zihao Feng[1], Delin Wang[1], Lampros Mavrommatis[1], Anna-Maria Vllaho[1], Merwan Rombach[1], Giulia Cucinella[1], Mingbo Cheng[2], Till Lautenschläger[6,7,8], David Rodriguez Morales[6,7,8], Rogier J. A. Veltrop[9,10,11,12,13], Leon J. Schurgers[1,9], Moritz von Scheidt[14,15], Barbara Mara Klinkhammer[16], Florian Kahles[17], Jennifer Kranz[18,19], Aitor Aguirre[20], Anne Loft[21,22], Niklas Klümper[23,24], Markus Eckstein[25], Thomas Seidel[26], Giancarlo Forte[27,28,29], Rik Westland[30], Man Zhang[31], Heng Zhao[31], Feng Ren[31,32], Sikander Hayat[1,33,34], Junedh Amrute[35], Benjamin Kopecky[36], Rebekka K. Schneider[37,38,39], Hind Medyouf[40], Pierre-Louis Tharaux[41], Peter Boor[16], Rafael Kramann[1,42], Linda W. van Laake[43], Annelotte Vos[44], Bernadette S. de Bakker[45,46], Jermo Hanemaaijer-van der Veer[45,46], Kory Lavine[35], Nikolaus Marx[17], Karin Klingel[47], Michael T. Schaub[3,5, 48], Stefanie Dimmeler[6,7,8], Alex Zhavoronkov[31,32, 49], Hendrik Milting[50*], Ivan G. Costa[2,3*] and Christoph Kuppe[1,3,9*]

Affiliations:
[1]Division of Nephrology and Clinical Immunology, RWTH Aachen University, Medical Faculty, Aachen, Germany.
[2]Institute of Computational Genomics, RWTH Aachen University, Medical Faculty, Aachen, Germany.
[3]Center for Computational Life Sciences, RWTH Aachen University, Aachen, Germany.
[4]Department of Information Engineering, University of Padova, Padova, Italy.
[5]Department of Computer Science, RWTH Aachen University, Aachen, Germany.
[6]Institute of Cardiovascular Regeneration, Goethe University Frankfurt, Frankfurt am Main, Germany.
[7]German Center for Cardiovascular Research (DZHK), Partner Site Rhine-Main, Germany.
[8]Cardio-Pulmonary Institute (CPI), Frankfurt, Germany.
[9]Department of Biochemistry, Cardiovascular Research Institute Maastricht (CARIM), Maastricht University, Maastricht, The Netherlands.
[10]Department of Cardiology, Division of Heart and Lungs, University Medical Center Utrecht, Utrecht University, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands.
[11]Regenerative Medicine Center Utrecht and Circulatory Health Research Center, University Medical Center Utrecht, Utrecht University, Heidelberglaan 8, 3584 CS Utrecht, The Netherlands.
[12]LMNAcardiac.org, LMNA Patient Organization, Vredehofstraat 13761, HA Soest, The Netherlands.
[13]Netherlands Heart Institute, Holland Heart House, Moreelsepark 1, 3511 EP Utrecht, The Netherlands.
[14]Department of Cardiology, TUM Klinikum Deutsches Herzzentrum, Technical University Munich, Munich, Germany.
[15]Deutsches Zentrum für Herz- und Kreislaufforschung (DZHK), Partner Site Munich Heart Alliance, Munich, Germany.
[16]Institute of Pathology, RWTH Aachen University Hospital, Aachen, Germany.
[17]Department of Internal Medicine I-Cardiology, University Hospital Aachen, RWTH Aachen University, Aachen, Germany.
[18]Department of Urology and Pediatric Urology, University Hospital RWTH Aachen, Aachen, Germany.
[19]Department of Urology and Kidney Transplantation, Martin-Luther-University, Halle, Germany.
[20]Institute for Quantitative Health Science and Engineering, Michigan State University, East Lansing, MI, USA; Department of Biomedical Engineering, Michigan State University, East Lansing, MI, USA.

[21]Center for Functional Genomics and Tissue Plasticity (ATLAS), Department of Biochemistry and Molecular Biology, University of Southern Denmark (SDU), Odense, Denmark.
[22]The Novo Nordisk Foundation Center for Genomic Mechanisms of Disease, Broad Institute of MIT and Harvard, Cambridge, MA, USA.
[23]Department of Urology and Pediatric Urology, University Hospital Bonn, Bonn, Germany.
[24]Institute of Experimental Oncology, University Hospital Bonn, Bonn, Germany.
[25]Institute of Pathology, Universitätsklinikum Erlangen, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Erlangen, Germany; Comprehensive Cancer Center Erlangen-EMN (CCC ER-EMN), Erlangen, Germany; Comprehensive Cancer Center Alliance WERA (CCC WERA), Erlangen, Germany; Bavarian Cancer Research Center (BZKF), Erlangen, Germany.
[26]Institute of Cellular and Molecular Physiology, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany.
[27]International Clinical Research Center (ICRC), St Anne's University Hospital, Brno, Czech Republic.
[28]Faculty of Medicine, Masaryk University, Brno, Czech Republic.
[29]School of Cardiovascular and Metabolic Medicine and Sciences, King's College London, London, UK.
[30]Department of Pediatric Nephrology, Amsterdam UMC-Emma Children's Hospital, Location University of Amsterdam, Meibergdreef 9, 1105 AZ, Amsterdam, The Netherlands.
[31]Insilico Medicine Shanghai Ltd., Shanghai, China.
[32]Insilico Medicine Hong Kong Ltd., Hong Kong SAR, China.
[33]Cardiovascular Research Institute and Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA.
[34]Windreich Department of Artificial Intelligence and Human Health, Icahn School of Medicine at Mount Sinai, New York, NY, USA.
[35]Center for Cardiovascular Research, Division of Cardiology, Department of Medicine, Washington University in St. Louis School of Medicine, St. Louis, MO, USA.
[36]Department of Medicine, Division of Cardiology, University of Colorado Anschutz Medical Campus, Aurora, CO, USA.
[37]Department of Developmental Biology Erasmus Medical Center Rotterdam, The Netherlands.
[38]Oncode Institute Erasmus Medical Center Cancer Institute Rotterdam, The Netherlands.
[39]Department of Cell and Tumor Biology, Faculty of Medicine University Hospital RWTH Aachen, Aachen, Germany.
[40]Department of Hematology, Oncology, Hemostaseology, Stem Cell Transplantation, Medical Faculty, Center for Integrated Oncology (CIO), RWTH Aachen University, Aachen, Germany.
[41]Université Paris Cité, INSERM, Paris Cardiovascular research Center (PARCC), Paris, France.
[42]Department of Internal Medicine, Nephrology and Transplantation, Erasmus Medical Center, Rotterdam, The Netherlands.
[43]Department of Cardiology, Division of Heart & Lungs, University Medical Centre Utrecht and Utrecht University, Utrecht, The Netherlands.
[44]Department of Pathology, University Medical Center Utrecht, Utrecht, The Netherlands.
[45]Amsterdam UMC location University of Amsterdam, Dept. of Obstetrics and Gynecology, Amsterdam, The Netherlands.
[46]Amsterdam Reproduction and Development Research Institute, Amsterdam, The Netherlands.
[47]Cardiopathology, Institute for Pathology and Neuropathology, University Hospital Tübingen, Tübingen, Germany.
[48]Department of Biology, RWTH Aachen University, Aachen, Germany.
[49]Insilico Medicine US Inc., Cambridge, MA, USA.
[50]Heart- and Diabetescenter NRW, Erich and Hanna Klessmann-Institute for Cardiovascular Research and Development, University Hospital of the Ruhr University Bochum, Bad Oeynhausen, Germany.

#co-first authors, *co-senior authors
Correspondence to:
Ivan Costa, PhD
Institute of Computational Genomics
Medical Faculty RWTH Aachen University,
Pauwelsstrasse 30, 52074 Aachen, Germany

104   Phone: +49-241-80-89760
105   E-mail: igesteiracostafilho@ukaachen.de
106
Christoph Kuppe, MD, PhD
Division of Nephrology and Clinical Immunology,
Medical Faculty RWTH Aachen University,
Pauwelsstrasse 30, 52074 Aachen, Germany
Phone: +49-241-80-37308
E-mail: christoph.kuppe@googlemail.com

Abstract

The adult mammalian heart has a limited regenerative capacity. Following injury, cardiomyocytes undergo a hypertrophic response accompanied by polyploidization, which has been described as a barrier to proliferation and regeneration of the heart[1,2]. However, the unique molecular programs of polyploidy, or genome multiplied cardiomyocytes, and their influence on the disease-related myocardial remodelling process remains unclear. Here, we integrate single-nuclei and high-resolution spatial multi-omics across human, rat, and mouse hearts to define novel cardiac cell states and their tissue niches in ischemic and non-ischemic heart disease. Computational analysis across scales allowed us to generate detailed networks of the cardiac tissue remodelling process as well as tissue and sub-cellular environments uniquely enriched in polyploid cardiomyocytes or their diploid origins. We identify a conserved, dichotomous transcriptional program distinguishing diploid from polyploid cardiomyocytes. Polyploid cardiomyocytes demonstrated rewired metabolic and chromatin-remodeling transcriptional programs and recapitulate the gene signature of immature human fetal cardiomyocytes. Notably, we observe that polyploid cardiomyocytes—rather than the general myocyte population—are the primary sites of enrichment for major heart-failure drug targets, including the mineralocorticoid, β1-adrenergic, and glucagon-like peptide-1 receptors. Based on our cross-species dataset we further identified TNIK, a Wnt-pathway regulator expressed in polyploid cardiomyocytes across species, as a potential therapeutic target and demonstrate that pharmacological TNIK inhibition improves cardiac function after myocardial infarction in rats. Together, this species-spanning, disease-resolved study redefines cardiomyocyte heterogeneity in heart disease and suggests a therapeutic path to heart failure treatment by targeting polyploid cardiomyocytes.

Results

Heart failure represents a major global health burden, affecting nearly half a billion individuals worldwide and causing more deaths than most forms of cancer[3]. Despite significant treatment advancements, current therapies typically address the symptoms and systemic consequences of heart failure, but do not directly target the pathways

driving maladaptive remodeling or those that restrict cardiac regeneration[4]. The human
heart possesses a minimal intrinsic regenerative capacity[5] compared to that of other
species such as zebrafish[6] and neonatal mice[7]. Human cardiomyocytes (CMs) originate
from proliferative mesodermal precursors during embryogenesis but rapidly lose their
regenerative capacity shortly after birth, coinciding with a metabolic shift from glycolysis
to fatty acid oxidation and the initiation of polyploidization[1,8,9]. The adult human heart
predominantly responds to injury with maladaptive remodeling, stress-associated gene
activation, concomitant fibrosis, and progressive functional loss[10,11]. This stress response
is typically associated with upregulated expression of the genes NPPB and NPPA, which
are co-regulated by an evolutionarily conserved enhancer cluster and induced by
mechanical stress[12,13]. Recent studies suggest that metabolic perturbations, specifically
inhibition of fatty acid oxidation, induce a fetal cell state shift in CMs and can induce
cardiomyocyte proliferation and regeneration highlighting the influence of metabolic
states on cardiomyocyte cell state plasticity[14]. It remains unclear whether additional CM
cell states, beyond the canonical NPPB/NPPA stress-response, drive cardiac tissue
remodelling.
160
Polyploidization, characterized by increased genome copy numbers or multinucleation in
CMs, emerges postnatally as the heart matures[15]. It represents a fundamental cell growth
principle in biology, utilized by different organisms and is common in plants and salivary
glands of insects[16–18]. Polyploid cell types can be found in various tissues, including
heart[19], muscle[20], liver[21] and pancreas[22] and have been suggested to play a role in tissue
repair and regeneration[21]. While several mechanisms like cell fusion, mitotic slippage,
and endocycling have been identified as polyploid mechanisms, endocycling, i.e. rounds
of DNA replication without cell division, has been identified as the major mechanism in
the heart, which is particularly linked to mitochondrial function[23]. Polyploidization has
been hypothesized to serve as a barrier to CM proliferation, thereby limiting cardiac
regeneration[2]. This appears paradoxical given that polyploidy is beneficial in other
organs, such as the liver[24], where it supports tissue regeneration[24] and protects against
genotoxic stress[25,26]. Despite extensive research[27], the functional and molecular
distinctions between diploid and polyploid CMs in the mammalian heart, especially
regarding their disease-modifying relevance in the human heart, have remained elusive.
176
Here, we present a single-cell and spatial multi-omics analysis of over 4.5 million cardiac
nuclei from healthy and diseased hearts across 105 human, 16 rat, and 16 mouse donors.
This includes a temporal map of mouse myocardial infarction (MI) and the evaluation of
a novel therapeutic approach targeting polyploid CMs in a rat model. Selection of samples
was performed based on pathologist assessment and refined by UNI, a foundation model
for pathology image analysis[28]. Samples were measured using probe-based single-
nucleus RNA sequencing (snRNA, Chromium Fixed RNA Profiling) and spatial

transcriptomics (MERFISH, Xenium) techniques using fixed-frozen or formalin-fixed paraffin-embedded (FFPE) cardiac tissue. An advanced computational framework exploring histology and sample level spatial information identified several novel disease-associated cell states and profibrotic tissue niches in ischemic and non-ischemic heart disease. To take full advantage of the subcellular resolution of our spatial transcriptomic measurements, we studied patterns in the position of individual transcripts across CMs. We identify multiple cell state-specific subcellular patterns, among others a specific signature enriched in the intercalated disk (ICD) region, the cardiomyocyte-cardiomyocyte interface. In summary, our study provides a high-resolution, cross-species dataset that distinguishes polyploid from diploid CM states, resolves their interactions with other cardiac cell types, and delineates the regulatory networks underlying myocardial remodeling in ischemic and non-ischemic failing human myocardium. Specifically, we identify polyploid CMs as key regulators of maladaptive cardiac remodelling and TNIK (TRAF2 and NCK interacting kinase), selectively expressed in polyploid CMs, as a novel therapeutic target limiting cardiac maladaptive remodelling.

Spatial multi-omics analysis of human heart disease

To characterize cellular heterogeneity and tissue remodelling in ischemic and non-ischemic human heart disease, we analysed heart tissue of 105 adult and fetal human donors and 16 rat and 16 mouse samples (**Fig. 1a-c**), using probe-based snRNA-seq (Flex) and high-resolution spatial transcriptomics (ST; MERFISH/Xenium) (**Fig. 1d**, **Extended Data Fig. 1a-b; Supplementary Table 1**). In total, the snRNA-seq dataset includes 997,362 nuclei and the ST dataset 3,593,941 nuclei (**Fig. 1d**). Human samples were obtained from left ventricular and septal walls, encompassing controls from non-failing donor hearts and patients with ischemic cardiomyopathy (ICM), non-ischemic dilated cardiomyopathy (DCM) of idiopathic etiology, and acute myocardial infarction (ICM_AMI) to capture a wide range of tissue remodelling dynamics (**Supplementary Table 1**). The tissue specimens were collected at heart transplantation or during implantation of a ventricular assist device. The five fetal heart tissue samples were obtained from human fetuses from terminated pregnancies between 12 and 21 post-conception weeks (p.c.w.) (**Fig. 1a**) from the Dutch Fetal Biobank. In addition, we analyzed heart tissues following drug treatment during the cardiac tissue remodelling phase (**Fig. 1b**) in rats (total n=52, n=16 for snRNA-seq) and following myocardial infarction (MI) at five different time points (**Fig. 1c, Extended Data Fig. 1a; Supplementary Table 1**) in mice (n=16, snRNA-seq + Xenium). For each cardiac sample, we obtained a 25 µm tissue section from either fixed-frozen or FFPE tissue specimens for nuclei isolation. We optimized our probe-based snRNA-seq isolation protocol to achieve high data quality and the lowest possible background from ambient RNA. 5 µm tissue sections were collected for high-resolution spatial transcriptomics (**Fig.**

222 **1d**, **Methods**). From each sample, an H&E staining was performed and evaluated by an
223 expert cardiac pathologist.
224
For snRNA dataset annotation we followed a two-step approach. An initial graph-based
clustering[29] and annotation based on marker genes resulted in the annotation of 20 major
cell types across the three species (**Supplementary Table 2**). Each major cell type was
then subclustered into detailed dataset-specific cell states with a focus on cross-species
cardiomyocyte diversity (**Fig. 1e-g**, **Supplementary Fig. 1-3)**. Our snRNA-seq data
contained a median of 2,358 UMIs and 1,403 detected genes per nucleus for the adult
human, 1,190 UMIs and 612 detected genes per nucleus for rat, and 1,512 UMIs and
1,051 detected genes per nucleus for mouse (**Supplementary Table 2**).
233
Altogether, our cross-species snRNA data identified several novel CM cell states, as well
as cell types not captured in previous studies, such as neutrophils (**Fig. 1e-g**). The
enhanced recovery originates from a nuclei-isolation protocol optimized for fixed heart
tissue that reduces ambient-RNA bias and minimizes loss of nuclei with atypical
morphology (e.g. neutrophils)[30,31]. While the transcriptomic complexity (genes detected
per UMI) was comparable to our previous polyA-3'-enriched snRNA dataset[11], our new
data achieves substantially higher gene detection rate (up to 14,452 genes vs 6,202
genes) and deeper sequencing coverage (2,534 vs 2,117 median UMIs per cell),
indicating improved capture efficiency and transcriptome sampling (**Extended Data Fig.**
1c).
244
Marker gene analysis identified two primary CM clusters conserved across three
mammalian species, each containing distinct subclusters (**Fig. 1e-h**). The first major
group was characterized by *DES* expression and included two specific CM
subpopulations: one defined by *NPPA* and *NPPB* expression, and another human-
specific subcluster defined by *DES*⁺/*SYNPO*⁺ expression. These *DES*⁺/*SYNPO*⁺ CMs
displayed a transcriptional program enriched for cytoskeletal remodeling genes—
including *SYNPO*, *SYNPO2L*, and *SYNPO2*—alongside increased *TAZ* (*WWTR1*)
expression, consistent with a mechano-adaptive, disease-associated state (**Extended**
Data Fig. 1d). The second major CM group was defined by the expression of *CFLAR* and
MYH7B (or *Myh7b* in rodents). *MYH7B* represents an ancient member of the myosin
heavy chain motor protein family, and its expression has not previously been described
in a unique adult human CM subpopulation[32]. Unlike the alpha and beta myosin heavy
chains (encoded by *MYH6* and *MYH7*), *MYH7B* does not produce a full-length protein
due to post-transcriptional exon-skipping[33]; instead, it functions as a long non-coding RNA
that regulates the ratio of alpha to beta myosin heavy chains and modulates the
expression of other sarcomeric genes[32]. Within this second group, we also identified a
small, human-specific subfraction of CMs expressing *HAND2*, a cardiac-enriched

transcription factor essential for lineage specification during development whose
reactivation in adult cells is linked to stress- and remodeling-associated gene programs.
CFLAR (also known as c-FLIP), an apoptotic regulator with a reported role in beneficial
cardiac remodeling[34], serves as an additional key marker for this novel CM group in
humans (**Fig. 1h**). Notably, these novel CM states were not detectable in two previously
published datasets[11,35] utilizing polyA-3' snRNA technology (**Extended Data Fig. 1e-f**),
highlighting the technical sensitivity required to resolve these populations.
269
In human hearts, CMs were approximately evenly distributed between the CFLAR$^+$ and
DES$^+$ groups at the cohort level, however, substantial inter-individual variability was
observed. Across patients, the percentage of CFLAR$^+$ CMs ranged from 20% to 80%.
(**Extended Data Fig. 1g**). The proportion of CFLAR$^+$ CM was higher in ICM and DCM
than in ICM_AMI (**Extended Data Fig. 1g**). Notably, variation in the CFLAR$^+$/DES$^+$
distribution was not associated with clinical measures of cardiac function, including left
ventricular ejection fraction (Spearman's ρ = 0.23, p = 0.17, n = 36) or left-ventricular end-
diastolic diameter (LVEDD) (Spearman's ρ = −0.27, p = 0.10, n = 36), in samples with
available metadata.
279
To further resolve disease-associated compositional changes, we analyzed cell
abundance shifts in our human snRNA data (**Extended Data Fig. 1h**). ICM_AMI samples
exhibited a pronounced increase in immune cell populations, including SPP1$^+$
macrophages, monocytes, plasma cells, and mast cells, alongside expansion of
myofibroblasts and CTHRC1$^+$ fibroblasts (**Extended Data Fig. 1h**). CM sub-states also
showed disease-specific remodeling patterns. CFLAR$^+$ CMs were significantly expanded
in ICM and DCM, whereas ICM_AMI was characterized by a reduction of multiple CM
states, including DES$^+$, SLC25A4$^+$, and specific CFLAR$^+$ CM substates (0, 3, and 4)
(**Extended Data Fig. 1h**). Additionally, perivascular and ELN$^+$ fibroblasts increased
significantly only in ICM and DCM, while in ICM_AMI myofibroblast and CTHCR1$^+$
fibroblasts were significantly increased, highlighting divergent stromal remodeling
programs between chronic and acute cardiac injuries (**Extended Data Fig. 1h**).
292
We used consecutive FFPE sections to perform spatial transcriptomic experiments with
probe sets covering 500 (MERFISH) or up to 5,100 (Xenium) genes (**Extended Data
Fig.1b**). Panel design was based on multiple publicly available datasets (**Methods**). In
larger cells like CMs, this allowed us to recover up to 10,000 transcripts per cell, while
across all cell-types 175 to 306 median genes per segmented cell/nuclei were detected
(**Fig. 1i-j**, **Extended Data Fig. 2a**). Out of 150 biosamples (human, mouse and rat), about
50% were measured in both snRNA and spatial modalities (**Extended Data Fig. 1a**).
300

Figure 1 Cross-species single-nucleus and spatial multimodal atlas identifies conserved and human-specific cardiomyocyte states in heart disease. a, Study design and human cardiac tissue cohort. Adult human heart samples (n = 100) were obtained from non-failing controls, ischemic cardiomyopathy (ICM), dilated cardiomyopathy (DCM), and acute myocardial infarction (AMI), complemented by fetal heart samples (n = 5). **b,** Rat MI model. Hearts were harvested at day 7 post-MI and after treatment at day 28. **c,** Mouse myocardial infarction (MI) model. Hearts were collected at baseline and at multiple time points following MI (days 1, 3, 7, 14, and 28). **d,** Overview of spatial multimodal profiling. FFPE and frozen cardiac tissues were processed for histology (H&E), spatial transcriptomics (Xenium and MERFISH), and single-nucleus RNA sequencing (snRNA-seq). In total, 4,591,303 nuclei were profiled across snRNA-seq and spatial transcriptomics. **e,** UMAP embedding of human snRNA-seq data (n = 533,866 nuclei), colored by major cardiac cell types. Cardiomyocyte states are highlighted, including DES⁺, NPPA/NPPB⁺, SLC25A4⁺, DES_SYNPO⁺, HAND2⁺, and CFLAR⁺ cardiomyocytes. **f,** UMAP embedding of rat snRNA-seq data (n = 120,413 nuclei), revealing conserved cardiomyocyte states, including DES⁺,

NPPA/NPPB+, MYH7B+, and mixed cardiomyocyte populations. **g,** UMAP embedding of mouse snRNA-
seq data (n = 75,825 nuclei), identifying DES+, NPPA/NPPB+, and MYH7B+ cardiomyocyte states alongside
non-myocyte populations. **h,** Heatmaps showing expression of selected cardiomyocyte marker genes
across cardiomyocyte states in human (top), rat (middle), and mouse (bottom), highlighting conserved and
species-specific transcriptional programs. Expression in logarithmic counts. **i,** Spatial mapping of
cardiomyocyte patch clusters in human cardiac tissue using MERFISH. Representative H&E-stained
sections and corresponding patch cluster annotations are shown for control and ICM samples. Bottom
panels show high-resolution MERFISH images with DAPI and selected marker genes (MYH7B). Scale bar
indicates 1 mm. **j,** Spatial mapping of myocardial patch clusters in mouse hearts using Xenium across
control, MI day 0, day 7, and day 14. H&E staining, patch cluster annotations, and spatial transcriptomic
overlays are shown. Marker expression highlights cardiomyocyte and fibroblast-rich regions. Scale bar
indicates 1 mm. **k,** Patch-level cardiomyocyte (path-CM) composition analysis in human Xenium tissue
microarrays (TMAs), showing enrichment of cardiomyocyte states across myocardial, damaged
myocardium, fibrotic/scar, and granulation tissue regions. Asterisks indicate statistically significant
enrichments. Significance was determined by Fisher's exact test with adjusted p-values < 0.05. **l,** Patch-
condition analysis comparing AMI, DCM, and ICM to controls in human Xenium TMAs, revealing condition-
specific enrichment of myocardial damage and fibrotic tissue states. Significance was determined by t-test
with adjusted p-values < 0.05. Asterisks denote statistically significant differences.

Segmentation of CMs is an unsolved challenge, impeded by their elongated and irregular
shape. We trained "heartbreaker", a Cellpose[36] model finetuned on expert annotated
sections of our MERFISH data. This approach demonstrated superior performance
relative to other state-of-the-art algorithms (**Extended Data Fig. 2b-c; Methods**). Due to
the lower signal-to-noise ratio of cell membrane staining in Xenium, we opted for nuclei
segmentation using CellPose for the processing of those samples (**Extended Data Fig.
2d**). For our subcellular analysis of one tissue microarray (TMA) with 26 samples, we
performed a post-Xenium IF staining with wheatgerm agglutinin (WGA)[37] and DAPI, which
allowed us to confidently segment cells and nuclei (**Extended Data Fig. 2e**).

We annotated each spatial dataset by transferring cell state labels from the corresponding
species-matched snRNA-seq dataset. To ensure robust annotations, we used five state-
of-the-art label transfer tools[38–42] and assigned each cell the label predicted by the
majority of tools. The agreement between tools served as a measure of annotation
confidence, indicating which cell states could be reliably distinguished in the spatial data
despite capturing only a fraction of the genes measured in the whole-transcriptome
snRNA-seq datasets. When tools disagreed, cell types were merged into broader
categories until at least three out of five tools agreed. This resulted in comparable cell
type proportions between the spatial and snRNA-seq datasets from the same donor
(Spearman's ρ > 0.8).

The inclusion of histology (e.g., H&E staining) provided an additional modality that
enabled an integrative, cross-species analysis of the spatial data (**Fig. 1i+j**). To analyze
the H&E stainings, we employed a deep learning model specialized in heart tissue[28]. We
divided the images into small patches and used the model to classify them based on their
histological features (**Extended Data Fig. 3a**). This method successfully distinguished

several specific tissue compartments, effectively bridging the analysis between human
and mouse samples (**Extended Data Fig. 3b**). The identified tissue compartments
effectively captured the transition from the healthy heart (myocardium, adipose tissue)
through stress (damaged myocardium 1, 2 and 3), major damage (fresh infarct with
demarcation, granulation tissue, necrotic tissue) and chronic damage (fibrotic tissue, scar
tissue) (**Fig. 1i, Extended Data Fig. 3b**).

While the majority of tissue compartments were consistent with the pathologist's
annotation, we observed patch sub-clusters (damaged myocardium 1, 2 and 3), that were
not readily identifiable by the experts (**Extended Data Fig. 3c**). To uncover compositional
differences we related the cell types identified in our spatial transcriptomic measurement
(Xenium measurement of human TMAs) to the H&E-derived annotation. We detected
differences in the CM composition of different areas (**Fig. 1k**). While CFLAR$^+$ CMs were
enriched in myocardium, damaged myocardium 1 and granulation tissue, DES$^+$ CMs
enriched in damaged myocardium 2 together with DES-SYNPO$^+$ CMs, which were also
found in fibrotic and scar tissue regions. NPPA-NPPB$^+$ CMs (border-zone stressed
cardiomyocyte signature) were found in high levels in the patches annotated as
myocardium damaged 3 in both human and mouse. We could identify an enrichment of
immune cells (macrophages, monocytes) in the fibrotic and granulation tissue (**Extended
Data Fig. 3d**). Myofibroblasts were localized especially in damaged myocardium 2 and 3
and fibrotic scar tissue regions. Comparing the different disease groups (ICM_AMI, ICM,
DCM) we were able to identify an enrichment of myocardium and damaged myocardium
1 in controls whereas damaged myocardium 3 was enriched in DCM as well as
fibrotic/scar tissue (Fig. 1l). Quantitative analysis of nearest-neighbor counts revealed that
DES+ CMs form more cohesive spatial clusters (based on nearest-neighbor distances),
while CFLAR+ CMs are more spread out (**Extended Data Fig. 3e**), supporting the model
of distinct spatial patterning of CM states in the diseased human myocardium.

Together, these analyses establish a spatially resolved, cross-species dataset integrating
single-nucleus transcriptomics, spatial gene expression, and histology-based tissue
annotation. We identify novel CM (CFLAR$^+$, DES$^+$, DES-SYNPO$^+$, SLC25A4$^+$) beyond
the established stressed NPPA-NPPB$^+$ CM state. By anchoring transcriptomic states to
specific morphological zones—ranging from healthy myocardium to chronic scar—we
demonstrate that novel CM states (CFLAR$^+$, DES$^+$, etc.) are not transient signatures but
are related to specific stages of pathological remodeling.

Spatial niches of human heart disease

While the histology-based compartments provided a macro-scale framework of tissue
injury, they represent broad morphological zones rather than precise functional units.
Cellular colocalization serves as a proxy for the complex communication networks that

operate at different length scales to maintain homeostasis or drive pathology. To bridge
the gap between these large-scale regions and cellular behavior, we moved from tissue
compartments to spatial niches—repeating tissue structures with co-localized cells—and
characterized how they differed across patients and disease conditions (ICM_AMI, ICM,
or DCM).
405
We performed a sample level cell–cell colocalization analysis using a framework
combining NicheSphere and PILOT[43] (**Extended Data Fig. 4a, Methods**). Using
NicheSphere, we identified six distinct niches in the cell interaction networks in the
contrast control vs. ICM samples (**Fig. 2a**). Similar to the H&E spatial data analysis, our
spatial transcriptomic analysis revealed that CM states were localized in separate niches
e.g. CFLAR$^+$ and SLC25A4$^+$ CM together with macrophages, pericytes and lymphatic
endothelial cells in niche 4. Cells comprising this niche, characteristic of healthy
myocardium, showed diminished spatial colocalization in ICM. Regarding niches with
increased colocalization in ICM, niche 3 includes CM cell states (DES$^+$, NPPA_NPPB$^+$
and DES_Synpo$^+$). Niche 1 includes myofibroblasts clustered together with
proinflammatory macrophages, monocytes and fibroblasts forming a unique profibrotic
and proinflammatory tissue niche (**Fig. 2a**). Focusing on colocalization statistics
associated with the different novel CM subtypes, only myofibroblasts increase their
colocalization with all CM subtypes (**Fig. 2b**). However we also noted distinct differences
between CFLAR$^+$ and SLC25A4$^+$ CMs compared to the other CMs. These CM states lost
their colocalization to macrophages, pericytes and mast cells, whereas DES_SYNPO$^+$
CMs uniquely increased their localization to tissue resident macrophages (**Fig. 2b**). We
next compared the spatial niches (based on spatial transcriptomics) with our identified
patch clusters (based on H&E staining) and observed strong concordance between the
two approaches. (**Extended Data Fig. 4b**). The niche 4 (CFLAR+, SLC25A4+ CMs) was
related to healthy myocardium, whereas niche 3 (DES, NPPA-NPPB-CMs) was
significantly enriched in damaged myocardium 1-3 (**Extended Data Fig. 4b**). By resolving
the cellular neighborhoods of each CM state, we demonstrate that cardiac remodeling
involves the formation and dissolution of specific multicellular niches: loss of homeostatic
CFLAR$^+$/SLC25A4$^+$-macrophage-pericyte networks in ICM parallels emergence of
profibrotic DES$^+$/myofibroblast assemblies, revealing niche-level reorganization as a
hallmark of disease progression.
433
To study the dynamical changes of spatial niches, we performed a sample level trajectory
analysis using PILOT (**Fig. 2c**). A statistical analysis on niche changes over the trajectory
indicated a decline of niche 4 (CFLAR myocardium) and expansion of niches 1, 2 and 5
(macrophage, myofibroblast, lymphoid cell, and resident immune niches), as well as a
transient increase of niche 3 with DES-CMs (**Fig. 2d; Extended Data Fig. 4c**). Selected
cores display changes in niche, patches, cells and genes among the disease trajectory

(**Fig. 2e**). For example, CCDC80 was identified as a gene with a significant increase in
expression in PILOT's trajectory for fibroblasts and myofibroblast cells (**Fig. 2e**). We
further found this gene was highly expressed in fibroblasts (not shown) particularly in the
CTHRC1[+] fibroblast subpopulation in the snRNA-seq data (**Extended Data Fig. 4d**).
Other trajectory-related genes for CFLAR-CMs include CM contractile genes (*TNNT2*,
MYBPC3) showing a reduction, whereas *TTN*, *PDE4DIP* and *DST* have an increased
expression during disease progression (**Fig. 2e; Extended Data Fig. 4e**). Thus, PILOT
trajectory analysis suggests that CFLAR[+] CMs in ICM undergo a transition from a
sarcomere-rich contractile state toward a stress-adaptive, cytoskeletal, and ECM-
anchored (extracellular matrix) phenotype (based on the DST and PDE4DIP gene
changes)[44,45] which is accompanied by increasing activation and presence of ECM-
secreting fibroblast subsets.
452
Comparing controls vs. DCM we identified five cardiac tissue niches (**Extended Data Fig.
4f-h**). Two niches had similar cell types and changes to ICM niches, i.e. decrease in
colocalization of cells in the CM-rich CFLAR myocardium (niche 4) and increase of
colocalization of cells in myofibroblast and macrophages (niche 2) (**Extended Data Fig.
4f**). However, our analysis revealed that the DES_Synpo[+] CM state constituted their own
cardiac niche in DCM with endothelial and dendritic cells (niche 3) (**Extended Data Fig.
4f-g**). Similar to ICM, all CMs increased their spatial colocalization with myofibroblasts
(**Extended Data Fig. 4g**). Interestingly, the DES_Synpo[+] CMs demonstrated more spatial
interactions with other cell-types compared to ICM, indicating a potential unique role in
DCM. Moreover, in DCM samples, niche 3 with DES-SYNPO[+] CM was highly enriched in
fibrotic tissue patches (**Extended Data Fig. 4i**). A comparison of niche distribution in ICM
vs. DCM samples (**Fig. 2e** vs. **Extended Data Fig. 4j**) revealed distinct spatial
architectures. ICM samples exhibited significantly higher spatial clustering compared to
the diffuse, fine-grained distribution observed in DCM (Wilcoxon rank-sum statistic = 4.09,
p < 0.0001) (**Extended Data Fig. 4k**). This focal organization was especially pronounced
for inflammatory ICM niches 5 and 6, which formed contiguous domains.
469
In CFLAR[+] CMs of contrast control vs. DCM samples, contractile genes (*TNNT2*,
MYBPC3) decline early as in ICM (**Extended Data Fig. 4l**), while *TTN*, *OBSCN* and the
NMD kinase *SMG1* increase, indicating a CM-intrinsic program of sarcomere scaffold
realignment. Comparative niche analysis (ICM vs. DCM) demonstrates conserved
localization of CFLAR[+] and NPPA_NPPB[+] CMs in distinct cardiac tissue niches across
cardiomyopathy etiologies, contrasted by disease-specific redistribution of DES-SYNPO[+]
CM states (**Extended Data Fig. 5a**). For the controls vs. ICM_AMI contrast, we detected
six tissue niches using NicheSphere (**Extended Data Fig. 5b**). Niche 1 contained all CM
substates and represented the niche losing colocalization interactions across other cell
types (**Extended Data Fig. 5b**). Interestingly, an increased colocalization is observed

between myofibroblasts and CFLAR+ CM cells as in other contrasts (**Extended Data Fig.**
5b). Among the niches with increased colocalization in ICM_AMI condition, we also
observe a fibrotic niche (niche 4) and a tissue resident (niche 6) with increase in ICM_AMI
(**Extended Data Fig. 5d**). The latter niche included SPP1+ tissue-resident macrophages,
which represent the cell type with the highest amount of ICM_AMI specific interactions,
recapitulating our previous findings[11].

Figure 2 Spatial tissue niches and subcellular transcript organization reveal cardiomyocyte state–specific remodeling in cardiomyopathy. a, NicheSphere interaction network control vs. ICM. Network representation of spatial tissue niches inferred from Xenium-TMA sections comparing ischemic cardiomyopathy (ICM) with controls. Nodes represent cell-type–specific niches and edges indicate significantly altered spatial interactions, with red and blue edges denoting increased or decreased interactions in ICM, respectively. **b,** Spatial colocalization analysis. Heatmap of differential colocalization (diffColoc score) between cardiomyocyte (CM) states and other cardiac cell types across Xenium-TMA samples. For **a** and **b**, only statistically significant interactions are shown (adjusted Wilcoxon's rank sums test; p-value < 0.05). **c,** PILOT pseudotime trajectory (control as blue dots vs. ICM as red dots)**.** Trajectory of TMA samples illustrating progressive niche remodeling from control to ICM tissue states. **d,** Niche proportion changes along pseudotime. Changes in relative abundance (y-axis) of myocardium-associated CFLAR+ CMs (top) and myofibroblast–macrophage niches (bottom) along pseudotime (x-axis), indicating reciprocal dynamics during disease progression (p-value - goodness of fit test on the regression of the cell proportion and pseudo-time). **e,** Tissue niche composition across six selected Xenium-TMA samples along the Trajectory from early to middle and late stages. Spatial maps showing niche assignments (top), histopathology-based tissue annotation (middle; myocardium, damaged myocardium, fibrotic and granulation tissue), and CM state distributions (bottom) across selected control and ICM samples. Representative gene expression patterns of genes with trajectory expressific changes are shown: DST (cardiomyocytes) and CCDC80 (fibroblasts) . **f,** Subcellular clustering. Spatial clustering of transcripts within cardiomyocytes identifies four recurrent subcellular domains across Xenium-TMA sections. Scale bar, 25 µm. **g,** Reference-based domain annotation**.** Mapping of Xenium-derived subcellular domains to APEX-seq reference signatures, assigning nuclear, intercalated disc (ICD), peri-nuclear, and cytoplasmic transcript compartments. **h,** Imaging validation of subcellular domains. Post-Xenium wheat germ agglutinin (WGA) staining and DAPI highlighting cardiomyocyte structure and nuclei, with arrows indicating enrichment of domain-specific transcripts. Bottom panel shows spatial localization of nuclear and ICD-associated transcripts. **i,** ICD-associated gene enrichment. Bar plot of genes enriched in intercalated disc–associated transcripts identified in Xenium-TMA data. **j,** Cytoplasmic gene enrichment. Bar plot of genes enriched in cytoplasmic transcript domains in Xenium-TMA data. **k,** snRNA-seq validation of subcellular gene programs. Heatmap showing mean expression of domain-specific genes across CM states in snRNA-seq data, confirming nuclear, peri-nuclear, cytoplasmic, and ICD-associated transcriptional programs. Particular enrichment is found of ICD domain 2 associated transcripts in DES_SYNPO+ CMs.

Subcellular features of human cardiomyocytes

Next we analyzed the subcellular localization patterns of RNA molecules across individual human CMs using BENTO[46], a toolkit for the analysis of subcellular RNA distribution. Unsupervised clustering revealed four distinct subcellular domains characterized by recurring micrometer-scale expression patterns **(Fig. 2f-j).** Enrichment analysis of these domains based on known subcellular organelle mRNA signatures from APEXseq[47] confirmed a nuclear (domain 1), perinuclear (domain 3), and two cytoplasmic domains (domain 2, 4) **(Fig. 2f-h, Extended Data Fig. 5e).** The two cytoplasmic domains were enriched for different mRNAs (**Fig. 2i-j, Extended Data Fig. 5e**). We hypothesized that the cytoplasmic domain 2 could correspond to CM-CM cell contacts, the area where the ICD supports the synchronized contraction of the heart also based on previous reports[48]. This was confirmed by co-staining with WGA, where the ICD is distinguishable by a higher signal intensity compared to lateral membranes. (**Fig. 2h-h', Extended Data Fig. 5e-h**). The ICD domain was enriched with genes related to membrane-cytoskeleton linkage like *CMYA5*, an important regulator of the cardiac dyad architecture[49] (**Extended Data Fig. 5h**). Notably, the RNA-level presence of the RNA-binding protein *SAMD4A*[50] and of the translation initiation factor *EIF3A*[51] is consistent with localized post-transcriptional

regulation and translational activity, raising the possibility of spatially organized protein
synthesis in proximity to the ICD (**Fig. 2i**). After identifying genes with specific subcellular
expression patterns, we explored their expression levels in our human snRNA data (**Fig.**
2k). Our analysis shows an enrichment of ICD-related (domain 2) transcripts in
DES_SYNPO[+] CMs and of nuclear and perinuclear genes (domain 1 and 3) in CFLAR[+]
CM clusters (**Fig. 2k**). In summary, we define distinct subcellular mRNA domains in
human CMs and demonstrate that their utilization differs systematically across our novel
CM cell states, with ICD-associated transcripts preferentially enriched in DES_SYNPO[+]
CMs and nuclear transcripts in CFLAR[+] CMs, consistent with divergent adaptive
subcellular RNA localization programs in human heart disease.

Novel human cardiomyocyte cell states
Next, we analyzed the molecular differences of the CM cell-states (**Fig. 3a-b**). We based
our annotation of the CMs clusters on unique marker gene expression (**Fig. 3b**). Desmin
(*DES* gene), as intermediate filament in CMs, has important structural roles; it connects
several CMs specific subcellular structures like the ICD complexes and with organelles
like mitochondria and the nucleus[52,53]. HAND2[+] CMs represented the least represented
CM cell state with 1.77% (**Supplementary table 2**). The re-expression of *HAND2*, a fetal
CM transcription factor[54], combined with *MYH7B* points towards a fetal gene expression
shift. *CFLAR*, or *c-FLIP*, a direct Casp8 inhibitor and master anti-apoptotic regulator[55], is
uniquely and highly expressed in human CMs when compared to mouse and rat snRNA-
seq (**Fig. 3b, Fig. 1h**). It has been demonstrated that the gene *CFLAR* protects CMs from
cell death in MI[56]. Based on the apparent specificity of CFLAR[+] marker genes to enlarged
nuclei in our spatial transcriptomics data we hypothesized that the novel HAND2[+] and
CFLAR[+] CMs states could be related to polyploid, or genome duplicated CM cell states.
Nuclear DAPI stainings have been used to identify polyploid nuclei in tissues previously[57].
To confirm our hypothesis we extracted morphometric properties (using CellProfiler[58]) of
CM nuclei from our spatial transcriptomics Xenium dataset and checked for association
with our transcriptomically defined CM cell state identities. This analysis showed a large
degree of heterogeneity in nuclear appearance for the different CM states (**Fig. 3c-d,**
Extended Data Fig. 6a-d), and especially between CFLAR[+] and DES[+] CMs. Differences
in DAPI texture and nuclei size were also obviously distinct upon visual inspection (**Fig.**
3d, Extended Data Fig. 6a). Among the most distinct nuclear features in CFLAR[+] CMs
were the area of the nuclei followed by features related to DAPI intensity, edge intensity,
entropy and granularity, consistent with differences in polyploidy between the analyzed
CM states and potentially linked to chromatin changes (**Extended Data Fig. 6b-c**). A
machine learning model trained on these nuclei images was able to differentiate CFLAR[+]
and DES[+] CM based on both CellProfiler morphometrics as well as vision model (Dinov3)
features (**Extended Data Fig. 6d, Methods**).
574

To validate this result we next performed FANS (fluorescent activated nuclei sorting)
based on DAPI staining and resorted the nuclei to confirm the successful enrichment of
diploid (2N), tetraploid (4N) and octaploid (8N) nuclei (**Fig. 3e, Extended Data Fig. 6e-i**).
We performed this experiment on the tissue specimen of four patients where enough
material was available and sorted for 2N, 4N and 8N DAPI content, respectively. This
yielded eight enriched snRNAseq datasets (see schematic in **Extended Data Fig. 6e**). A
median UMI count per nuclei of 3,233 and median of 1,640 genes per nuclei indicated
that the FANS data had similar quality as our non-FANS isolation approach
(**Supplementary Table 1**). Non-CM cells were mainly found in the diploid (2N) nuclei
fraction, while 95% of nuclei in the polyploid fraction were derived from CM (**Extended**
Data Fig. 6f). Clustering of all CMs across fractions revealed two distinct CM populations
(**Extended Data Fig. 6g**). To connect this data to our main non-FANS snRNA dataset we
performed enrichment analysis. The 8N fraction was strongly enriched for the CFLAR$^+$
CM signature, while the 2N fraction scored high for the DES$^+$ signature (**Fig. 3e,**
Extended Data Fig. 6i). Nuclei in the 4N fraction were split into high DES+ and high
CFLAR cells, which may represent a transitional cell state (**Fig. 3e, Extended Data Fig.**
6h-i). Taken together the DAPI and FANS experiments confirmed that the CFLAR$^+$ CM
gene signature is linked to the ploidy of the CMs, while the DES$^+$ CM gene signature was
more indicative of diploid CM.
594

While we could not identify the novel CM signatures in two previously published 3' based
snRNA datasets (**Extended Data Fig. 1e+f**), we searched for scRNA-seq or snRNA-seq
studies of human heart tissue identifying similar heterogeneous CM states, independent
of atrial vs. ventricular phenotypes. The only exception being a dataset from Wehrens et
al.[59], where such CM clusters are described derived from human HCM patients after
septal myectomy and whole CM sequencing using SORT-seq (**Extended Data Fig. 6j**).
When we mapped the CFLAR$^+$ CM and DES$^+$ CM gene scores in the Wehrens scRNA-
seq dataset, we could identify a clear enrichment and separation of the two scores, fitting
to our observation of novel CM states in ICM and DCM patient samples (**Extended Data**
Fig. 6k). To increase our confidence in this mapping we employed a dual approach. Using
ClusterFoldSimilarity[60] we calculated similarity scores between the two datasets, based
on differential genes expressed for each cluster. In addition, we performed a hyper
geometric test (500 genes per cluster), which confirmed an overlap of expression patterns
between Wehrens cluster 5 and our polyploid HAND2$^+$ and CFLAR$^+$ CMs (**Extended**
Data Fig. 6l).
610

We next mapped structural, proliferative, metabolic and chromatin programs across CM
states (**Fig. 3f+g, Extended Data Fig. 7a-j**). Contractile sarcomere genes diverged in a
CM state-resolved manner: thin-filament and titin/Z-disc gene sets separated clearly
across subtypes, indicating remodeling of the sarcomere. In particular *OBSCN*, *LDB3* and

TTN were upregulated in HAND2$^+$ and CFLAR$^+$ CMs (**Fig. 3f**). Metabolic pathway
analysis revealed a coordinated metabolic re-wiring (**Fig. 3g, Extended Data Fig. 7a-d**).
Diploid DES$^+$ CMs demonstrated upregulated β-oxidation genes, whereas polyploid
HAND2$^+$/CFLAR$^+$ states showed increase in glycolytic and glutaminolysis gene
expression (**Fig. 3g, Extended Data Fig. 7a-c**). Over representation analysis comparing
DES$^+$ and CFLAR$^+$ CMs revealed significant enrichment of mitochondrial, oxidative
phosphorylation, and aerobic respiration–related gene sets in DES$^+$ CM (**Extended Data**
Fig. 7d).
623
We next compared the expression of cell-cycle regulatory genes (**Extended Data Fig.**
7e-f). In adult CMs three reactivation modes emerged. HAND2$^+$ CMs exhibited G1/S-
associated gene expression (*MYC, CDK4, CDK6, CCNE1, CCNE2*), with relative
enrichment for S-phase signatures compared to other states. Polyploid CFLAR$^+$ CMs re-
expressed late G2/M genes (*CDK1, CCNB1, AURKA, PRC1, HMMR*) in the absence of
G1/S induction, a pattern compatible with endoreplication-like genome duplication without
productive cytokinesis (**Extended Data Fig. 7e-f**). DES_SYNPO$^+$ CMs also upregulated
cell-cycle associated genes, including *CCND1/CCND2* (G1), *MKI67* (pan-proliferation),
TOP2A (S/G2), and *CDK1* (mitotic entry)-consistent with partial engagement of cell-cycle
transcriptional programs (**Extended Data Fig. 7e-f**).
634
Ion-channel programs segregated sharply by CM state (**Extended Data Fig. 7g**).
Polyploid CFLAR$^+$ subsets showed the strongest pacemaker/diastolic-depolarization
signature (*HCN1*) together with an L-type Ca^{2+} module (*CACNA1C* with *CACNB2*) and
broad TRP upregulation (*TRPC1, TRPC3, TRPM4, TRPM7*) (**Extended Data Fig. 7g**). In
contrast, diploid subtypes (NPPA_NPPB$^+$, SLC25A4$^+$, DES$^+$) preferentially expressed the
noncanonical connexin GJA3 (C*X46*), whereas polyploid CM states expressed GJC1
(CX45), indicating a potential junctional coupling shift at the ICD. Together, these patterns
point to a polyploid-biased increase in depolarizing drive (*HCN1*), Ca^{2+} influx capacity
(*CACNA1C/CACNB2*), mechanosensitive entry (*TRPM4/7*; *TRPC1/3*), and an ICD
connexin switch (*GJA3→GJC1*) (**Extended Data Fig. 7g**). Chromatin and nuclear stability
gene programs were likewise CM state-specific (**Extended Data Fig. 7h**). Polyploid
CFLAR$^+$ CMs, particular the HAND2$^+$ state, were enriched for histone
acetylation/methylation genes, chromatin-remodeling factors, and nuclear-stability
components, consistent with broad epigenomic reorganization accompanying metabolic
shift (**Extended Data Fig. 7h**). This was further supported by the unique upregulation of
nuclear receptor gene expression involved in metabolism in CMs, like *PPARα and PPARγ*
(**Extended Data Fig. 7i).**
652
653   Gene set analysis revealed a clear enrichment of cardiac disease terms in polyploid CMs
654   (CFLAR$^+$ and HAND2$^+$) including hypertrophic cardiomyopathy (HCM) and dilatative

cardiomyopathy terms (DCM) as well as the associated maladaptive cell signalling pathways in cardiac remodelling including WNT signalling pathway (**Fig. 3h, Extended Data Fig. 7j**). NPPA-NPPB⁺ CMs demonstrated enrichment in systolic heart failure gene signature and cardiac hypertrophic response. DES⁺ diploid CMs enriched in metabolic terms oxidative phosphorylation and fatty acid metabolism as well as myogenesis and cardiac muscle contraction (**Fig. 3h, Extended Data Fig. 7d and 7j**).

Based on single-nucleus-resolved RNA expression and open chromatin information from our previous work[11] we constructed an enhancer based gene regulatory network (eGRN) with scMEGA[61]. The estimated eGRN detected two main regulatory modules (**Supplementary Table 4**). One module was associated with cells at the beginning of the trajectory (DES⁺ CMs) and one connected to cells at the end of the trajectory (CFLAR⁺, DES-SYNPO⁺ and NPPA-NPPB⁺ CMs). *CTCF*, which is crucial in chromatin organization of the developing heart[62], was the most well connected transcription factor (TF) (as determined by PageRank centrality) in the GRN module related to DES⁺ CMs (**Supplementary Table 4**). The integrated dataset was then used to construct a cell differentiation tree with PHLOWER[63]. By using DES+ CM cells as root, PHLOWER indicated branches leading to DES-SYNPO⁺ CMs and another branching leading to CFLAR⁺ or NPPA-NPPB⁺ CMs (**Fig. 3i**). We observe lower TF activity of *CTCF* in disease related CM sub-states such as CFLAR⁺, NPPA_NPPB+ and DES_SYNPO⁺ CMs (**Extended Data Fig. 7k**). Along this trajectory, CFLAR⁺ CMs displayed dynamic regulation of WNT pathway modulators, including a decrease expression of the WNT antagonist *DKK3* (**Fig. 3j**) and *SFRP1* (**Fig. 3k**), alongside induction of canonical WNT target genes such as *AXIN2* and the kinase *TNIK* (**Fig. 3k**). The main transcription factor regulated to DES-SYNPO⁺ branch was *WT1* (**Supplementary Table 4**), which is a known epicardial regulator[64] and is known to directly regulate synaptopodin gene expression e.g. in podocytes in the kidney[65]. Its knock-out is known in zebrafish to lead to a loss of CM expression programs and upregulation of epicardial gene expression[66]. For CFLAR⁺ CMs, predicted regulators include the sonic hedgehog pathway TF *GLI3*, which has a known muscle repair–associated angiogenesis role in myocytes[67] and *NR2C2* (or *TR4*), a reporter repressor of PPARα nuclear receptor signaling[68] (**Supplementary Table 4**).

We next analyzed the expression of ligands and receptors of known canonical fetal cell-cell communication and fibrosis related pathways in CMs (**Extended Fig. Data 8a-c**). CFLAR⁺ and HAND2⁺ CMs showed the highest expression among CMs of receptor and ligands associated with maladaptive remodelling including the receptor *NOTCH2* and ligands *JAG2, DLL1* together with non-canonical WNT ligands (*WNT5A, WNT11*). Among these, HAND2⁺ uniquely presented upregulated *SHH* expression, whereas CFLAR⁺ CM states demonstrated high *PTCH1/SMO* levels. In contrast, DES⁺, NPPA_NPPB⁺, and SLC25A4⁺ CMs preferentially expressed WNT antagonists (*DKK3, SFRP1/2*) and showed generally attenuated Notch/HH signatures, consistent with a contractile/stress-

adapted phenotype that restrains canonical Wnt activity (**Extended Fig. data 8a**). We
next analyzed fibrogenic cell-cell communication pathways (TGFβ, PDGF, endothelin,
VEGF/ANGPT, FGF/NRG axes) across CM states (**Extended Fig. data 8b**). DES$^+$,
NPPA_NPPB$^+$, SLC25A4$^+$, and especially DES_SYNPO$^+$ were enriched for distinct
profibrotic and proinflammatory ligands, including *TGFB* isoforms, *PDGFR-A/B*, *VEGFB*
and *VEGFC*, *IL6* and *CXCL12* consistent with a CM-derived "injury signal" that can
activate fibroblasts and vascular/perivascular cells (**Extended Fig. data 8b-c**). In
contrast, CFLAR$^+$ and HAND2$^+$ CMs preferentially expressed growth-factor receptors
(*TGFBR1/2*) together with *ERBB2/ERBB4* and the profibrotic ligands EDN1 and FGF2,
indicating a divergent send-receiver profibrotic phenotype between CM states (**Extended**
Fig. Data 8b-c).
705
In order to consider both cell-cell colocalization and changes in ligand-receptor
expression across disease conditions, we performed a cell-cell communication analysis
with scSeqComm[69] and CrossTalkeR[70] by focusing on colocalized cell pairs predicted by
NicheSphere (**Fig. 2a; Methods**). We first used CrossTalkeR pagerank to identify highly
influential cell types in ICM disease. This indicates the importance (higher communication
in ICM) of diploid CM clusters (SLC25A4$^+$, DES_SYNPO$^+$, DES+$^+$) with a highest score
for the SLC25A4$^+$ CMs and immune cells like neutrophils, macrophages (**Extended Data**
Fig. 8d). We next analyzed differences in intracellular signalling in CM from disease
samples. scSeqComm calculates differential cellular communication scores, which reflect
how cell-cell communication affects down-stream pathways and transcriptional
regulation. This indicates an upregulated activity of WNT, NOTCH and Hedgehog
signalling in CFLAR$^+$ CMs in the ICM vs. control contrast (**Fig. 3l**).
718
If we consider the signals related to WNT activation towards CFLAR (**Fig. 3m-n**), we
observe increased signalling from myofibroblasts including *CTHRC1[71], CCN2 and*
WNT11 (**Fig. 3m**). In ICM myofibroblasts secreted *CTHRC1*[71] can bind *FZD6* on CFLAR$^+$
CMs to potentiate non-canonical Wnt signaling. These predictions are consistent with our
prior spatial analysis showing interaction of CFLAR$^+$ CM with myofibroblast in all disease
conditions and indicates the activation of WNT signalling in CFLAR$^+$ CM cells.
Reciprocally, CFLAR$^+$ CM show an increased secretion of VEGFA towards
myofibroblasts (**Fig. 3n**). Other signals that drive the fibrotic niche created by CFLAR+
CMs are *AREG*, *DLL1* and *FAT4* (**Extended Data Fig. 8e-f**).
728
729 Our colocalization analysis (**Fig. 2g**) indicated an unique role of DES-SYNPO$^+$ CMs,
730 which formed a DCM specific niche not detected in ICM samples. Moreover, DES-
731 SYNPO$^+$ CMs had the strongest proinflammatory gene signatures including IL6, CCL2
732 and CXCL8 (**Extended Data Fig. 8c**). We identified a reciprocal signalling axis of
733 neutrophils and NK cells to DES-SYNPO$^+$ CMs involving *GNAI2* and *CCL5* highlighting a

distinct immune-CM axis in DCM (**Extended Data Fig. 8h-i**). When evaluating the intracellular pathway activity across contrasts, we observe an increase in WNT signalling in CFLAR+ CM in all disease conditions in particular in DCM samples (**Fig. 3o**).

Figure 3 Molecular, structural, metabolic, and signaling features of novel human cardiomyocyte cell states. a, UMAP embedding of snRNA-seq highlighting distinct CM populations, including CFLAR+, DES+, DES_SYNPO+, HAND2+, NPPA_NPPB+, and SLC25A4+ CMs. Arrows indicate cluster identities. **b,** Cell state marker genes. Heatmap showing scaled mean expression of representative marker genes across CM

states, confirming transcriptional distinctions between CMs. **c,** Nuclei morphology analysis. PCA of
CellProfiler-derived nuclear morphology features from DAPI staining separates CM states, indicating
systematic differences in nuclear shape and size across transcriptional identities. **d,** Representative nuclei
images. Immunofluorescence images of DES⁺ and CFLAR⁺ cardiomyocyte nuclei illustrating distinct
nuclear morphologies (arrows). Scale bar, 20 μm. **e,** Polyploidy enrichment using FANS. Violin plots of cell-
state signature scores across nuclei with different ploidy levels (2N, 4N, 8N) obtained by FANS. CFLAR
and DES signature enrichment differs by ploidy state (p-value < 0.0001). **f,** Sarcomere gene expression.
Schematic of sarcomeric organization (thin filaments, thick filaments, titin and Z-disc) and heatmap of
corresponding gene expression across CM states, highlighting differential regulation of structural
components. **g,** Metabolic gene programs. Heatmaps showing expression of genes involved in β-oxidation
(left) and glycolysis (right) across CM states, indicating metabolic specialization. **h,** Gene set enrichment
analysis (GSEA) of hallmark, KEGG, and WikiPathways terms across CM states, revealing differences in
hypertrophic signaling, metabolism, WNT signaling, and cardiac contraction programs. **i,** Differentiation
trajectory of CM states. PHLOWER-based trajectory inference depicting transitions between CM states
along pseudotime, with relative contributions of CFLAR⁺, DES⁺, DES_SYNPO⁺, HAND2⁺, and
NPPA_NPPB⁺ CMs. **j,** DKK3 expression dynamics. Smoothed expression of DKK3 along the CM
pseudotime trajectory, highlighting state-specific regulation. **k,** WNT pathway–related genes along
cardiomyocyte trajectories. Expression dynamics of SFRP1, AXIN2, and TNIK across CM states along
pseudotime, illustrating coordinated regulation of WNT signaling components. **l,** Intracellular pathway
activity (ICM vs. control). Heatmap summarizing inferred intracellular activity scores of key signaling
pathways (e.g., Toll-like receptor, JAK–STAT, WNT, TGF-β, NF-κB, Hedgehog) across CM states,
highlighting differential transcriptional response between ICM versus control hearts. **m,** WNT upstream
signaling (ICM vs. control). Sankey plot display ligand-receptor interactions upstream of WNT signalling in
CFLAR+ CM cells, comparing ICM with control hearts. Positive values indicate increased signaling strength
in ICM, whereas negative values indicate reduced interaction strength. **n,** Cell-cell communication
downstream of WNT signaling (ICM vs. control). Sankey plot displaying ligand–receptor interactions
downstream of WNT pathway in CFLAR+ CM cells comparing ischemic cardiomyopathy (ICM) compared
with controls; ligands correspond to transcriptional targets of WNT signaling. **o,** Disease-specific pathway
activity. Heatmap of inferred intracellular pathway activity across CM states comparing ischemic
cardiomyopathy (ICM), dilated cardiomyopathy (DCM), and acute myocardial infarction (AMI) with controls,
focusing on Hippo, TGF-β, and WNT signaling pathways.

Collectively, our results suggest the existence of distinct CM states characterized by
polyploidy linked remodeling of contractile, electrophysiological, metabolic, and chromatin
programs. These CM state-specific programs appear to converge further on specific
trajectories, raising the hypothesis that cardiomyocyte genome multiplications may serve
as a central organizing principle of CM-linked remodeling in human heart disease.

Fetal vs. adult human cardiomyocytes
A central observation in cardiac transcriptomic analysis is that postnatal hearts adopt a
fetal-like transcriptional state, also referred to as the "fetal gene hypothesis" in response
to stress or chronic injury[72]. Given the expression of cardiac fetal genes like *MYH7B* and
HAND2 in polyploid CMs, we next addressed how conserved this gene expression
signature is. To this end, we generated snRNA data using fresh frozen human fetal heart
tissues from five timepoints preserved in RNAlater solution (p.c.w. 12 - 22) (**Fig. 4a**). In

total we recovered 146,025 nuclei (**Extended Data Fig. 9a**). The nuclei contained a
median of 3,531 UMIs and expressed a median of 1,886 genes. Our analyses focused
on CMs of the left ventricle and identified six major cell states, which recapitulated the
divergent expression profiles of CFLAR⁺ and DES⁺ populations observed in the adult
dataset (**Fig. 4b-c**, all cells shown in **Extended Data Fig. 9a-b** and **Supplementary table**
2).
793
Compared to other fetal heart datasets[73,74], our data for the first time revealed all stages
of fetal CM proliferation (**Fig. 4d**). The subclusters cycling and proliferating CMs mapped
to the cell-cycle S-phase and G2/M-phase respectively (**Fig. 4d**, **Extended Data Fig. 9b**).
We next compared our novel human adult CM states with the cell-cycle scores (S and
G2/M) of our human fetal snRNA data (**Extended Data Fig. 9c-d**). While we found the
vast majority of adult human CMs in G1-phase and 1% in G2M, none of them were
enriched in cell cycle genes to the same level as the proliferating cells in the fetal heart
(**Extended Data Fig. 9d**).
802
In addition to these cell-cycle CMs, which we grouped for downstream analysis, we were
able to identify CMs in multiple stages of maturation based on the canonical maturation
markers *HOPX*[75] and *TCAP*[76] (**Fig. 4b**, **Extended Data Fig. 9e**). Gene-set enrichment
analysis revealed an increase of cell-proliferation terms in cycling CMs, WNT pathway
terms and glycolysis for immature CMs and myofibril-assembly and cardiac contraction
for mature CMs (**Extended Data Fig. 9f**), which corroborated our annotation.
Interestingly, we could detect a similar shift in sarcomere gene expression of fetal
immature vs. mature CMs (**Fig. 4e**) and changes in β-oxidation and glycolysis regulating
genes compared to adult human CFLAR⁺DES⁺ and DES⁺CFLAR⁺ CM (**Fig. 4f**). In order
to study the spatial arrangement of these fetal CM states we mapped their location in the
tissue using a publicly available spatial transcriptomics dataset[74]. We saw the expected
enrichment of our atrial and trabecular cells in their respective compartments (**Extended**
Data Fig. 9g-h). Immature and mature CM were more evenly spread throughout the
heart, although there was a higher abundance of immature cells around the right ventricle
and a higher abundance of mature cells in atrioventricular plane papillary muscle, which
might relate to differing mechanical strain in these distinct tissue regions (**Fig. 4g,**
Extended Data Fig. 9g).
820
Next we investigated gene expression changes during CM maturation. We partitioned our
data into immature, differentiating and mature CMs and calculated a diffusion map, to
visualize the continuous trajectory of the maturation process (**Fig. 4h**). By assigning cells
a pseudo-temporal ordering we were then able to study expression trends along this
trajectory. We found that an increase of the DES⁺ and a decrease of CFLAR⁺ signatures,
derived from the adult snRNA dataset, were among the most significant results (**Fig. 4i**).

Maturation was also accompanied by increased cellular respiration and oxidative
phosphorylation (**Fig. 4i**).
829

830
831
Figure 4 Single-nucleus and spatial dissection of human embryonic cardiomyocyte differentiation.
a, Overview of human fetal heart samples profiled by snRNA-seq across developmental stages (12–22+
post-conception weeks, pcw). **b,** UMAP embedding of snRNA-seq data from human fetal hearts, revealing
distinct cardiomyocyte (CM) populations including proliferating, cycling, differentiating, immature, stress-
responsive, trabecular, and terminally differentiated left-ventricular (LV) CMs. **c,** Heatmap of selected
marker genes defining major embryonic LV-CM states (scaled mean expression). **d,** Cell-cycle phase
annotation of embryonic CMs projected onto the UMAP, highlighting G1, S-phase, and G2/M populations.
e, Expression of sarcomeric gene modules (thin filament, titin/Z-disc, thick filament) across LV-CM
maturation states. **f,** Metabolic gene expression programs associated with β-oxidation and glycolysis across
LV-CM states.   **g,** Reference spatial domains from the human embryonic heart dataset (Lazar et al.),
annotated by anatomical and functional regions. Spatial mapping of embryonic CM states identified in this

study onto the reference human embryonic heart sections, showing regional enrichment of immature,
mature, and trabecular LV-CMs, alongside major non-CM lineages (estimated counts per Visium spot). **h,**
Diffusion map embedding depicting the continuous differentiation trajectory from immature through
differentiating to mature LV-CMs. **i,** Gene set enrichment analysis (GSEA) along the human embryonic CM
trajectory, highlighting pathways decreasing (blue) or increasing (red) during maturation. **j,** Cluster-to-
cluster similarity analysis comparing embryonic LV-CM states (y-axis) to adult human LV-CM populations
(x-axis), revealing selective correspondence between embryonic differentiation states and adult CM
phenotypes. Stars indicate significance (p-value <0.05) with an hypergeometric test.

In the next step we used ClusterFoldSimilarity[60] and a hypergeometric test to compare
the embryo subclusters to the adult human CMs. We found that adult HAND2$^+$ and
CFLAR$^+$ were significantly enriched in proliferating fetal CMs (**Fig. 4j**). Mapping of fetal
cell types into the PHLOWER cell differentiation tree based on adult CMs confirmed that
immature LV-CMs are related to the CFLAR+ CM branch **(Extended Data Fig. 9i)**.

Overall, our data uncover a conserved fetal differentiation trajectory of CMs that provides
a reference framework for adult CM states in human heart disease. Polyploid HAND2$^+$
and CFLAR$^+$ CMs transcriptionally revert toward immature fetal programs, whereas
diploid DES$^+$ and DES–SYNPO$^+$ CMs represent terminally differentiated, contractile end
states. Notably, polyploid CMs do not reacquire fetal proliferative signatures, supporting
a model of incomplete fetal reprogramming rather than regenerative proliferation.

Temporal disease association and HF targets in CMs

To study the novel CM heterogeneity and their dynamics with spatiotemporal resolution,
we performed snRNA-seq and Xenium in mice after LAD ligation across four post-MI time
points from day 3 to day 28 (**Fig. 5a**). In total, we recovered 75,825 nuclei from snRNA-
seq and 543,586 nuclei from spatial transcriptomics (**Fig. 1g; Supplementary Table 2**),
which allowed the detection of six transcriptionally distinct cardiomyocyte states (**Fig. 5b**).
Marker-gene expression of the novel human CM clusters demonstrated conservation in
the mouse dataset (**Fig. 5c, Extended Data Fig. 10a**), with a major cluster high in *Des*
and a second one expressing fetal genes like *Myh7b*, although *Cflar* was expressed at
much lower levels than in the human.

We next quantified shifts in CM state abundances across the post-MI time course[77]. Both
Des$^+$ and Myh7b$^+$ CMs were reduced at day 3 following the acute MI, whereas
NPPA_NPPB$^+$ CMs represented the only CM population that increased at this early injury
time point (**Fig. 5d, Extended Data Fig. 10b**). By day 7, the trajectories of Des$^+$ and
Myh7b$^+$ CMs diverged: Des$^+$ CMs remained low, while Myh7b$^+$ CMs expanded
progressively and became the dominant CM population at later stages (**Fig. 5d**). This
delayed but pronounced enrichment of Myh7b$^+$ CMs after injury highlights them as an
adaptive CM state that emerges during post-ischemic remodelling.

To link mouse and human CM states, we applied ClusterFoldSimilarity[60] and enrichment
test, which showed that the Des$^+$ CM nuclei from mouse overlapped in their expression
with diploid CMs of the adult human heart (DES$^+$, SLC25A4$^+$, DES_SYNPO$^+$,
NPPA_NPPB$^+$), whereas the Myh7b$^+$ CM mouse nuclei were more similar to human
polyploid HAND2$^+$ and CFLAR$^+$ CM nuclei (**Fig. 5e**). Nppa$^+$ mouse CMs overlapped
significantly with NPPA$^+$ human CMs but also shared markers with human HAND2$^+$ and
CFLAR$^+$ CMs (**Fig. 5e**). We then compared the mouse clusters to our fetal human data.
Des+ CMs were most similar to terminally differentiated CM, while Myh7b$^+$ CM showed
correspondence to fetal immature MYH7B+ cells, a similar trend to what we observed in
our adult to fetal comparison (**Extended Data Fig. 10c**).

Next we compared transcriptional programs between mouse CM subtypes (**Fig. 5f-g and**
Extended Data Fig. 10d–i). Sarcomere-associated genes, including *Obscn*, *Ldb3*, *Ttn*
and *Mybpc3*, were significantly upregulated in Myh7b$^+$ CMs, whereas genes involved in
β-oxidation regulation were upregulated in Des$^+$ CMs mirroring the human polyploid
pattern (**Fig. 3f-g, 5f-g**). Analyses of glycolytic and glutaminolysis genes (**Extended Data**
Fig. 10d-e) showed that Myh7b$^+$ CMs demonstrated reduced gene expression consistent
with a metabolically rewired, stress-adapted state, whereas Des$^+$ CMs maintained a more
oxidative, metabolic increased profile (**Extended Data 10d-f**). Because our human data
implicated polyploid CMs in fibrotic and developmental signalling, we next systematically
assessed these pathways in the mouse model (**Extended Data Fig. 10g-i**). Our analysis
shows that canonical profibrotic ligands and receptors including Tgfb isoforms, *Tgfbr1/2,*
Pdgfa/b, Pdgfra/b, Edn1/Ednra/Ednrb, *Vegfa* and *Il6/11*,were strongly enriched in Nppa$^+$
and Myh7b$^+$ CMs compared to Des$^+$ CMs, while Des+ CMs preferentially expressed
Vegfb and *Vegfc* (**Extended Data Fig. 10g-i**). This pattern is only partially concordant
with the human heart, where polyploid CFLAR+/HAND2+ CMs demonstrated higher
upregulated gene expression of fibrosis signalling ligands and receptors (**Extended Data**
Fig. 8b, 10g-i). However a marked bias of WNT (particularly WNT5/11) and Hedgehog
signaling components towards Myh7b$^+$ CMs remained: Myh7b$^+$ cells expressed multiple
WNT ligands and receptors, NOTCH ligands and receptors, and Sonic hedgehog
pathway members, whereas Des$^+$ CMs predominantly expressed inhibitory modulators
such as *Sfrp1* and *Dkk3* (**Extended Data Fig. 8g+i).**

Intracellular pathway analysis of mouse CM states revealed notable differences
compared to human CMs (**Extended Data Fig. 10j**). Specifically, WNT, TGFβ, and Notch
signaling were predominantly upregulated in murine Nppa+ CMs rather than Myh7b+
CMs, highlighting a species- or model-dependent divergence in signaling. Despite this,
the general functional stratification was conserved: the adaptive Nppa+ state in mice
concentrated profibrotic and developmental signaling capacities, whereas Des+ CMs
retained a restrained, homeostatic profile similar to that of humans (**Extended Data Fig.**
10j).

Figure 5 Spatiotemporal organization of cardiomyocyte states after myocardial infarction a,
Experimental design and sampling scheme of murine hearts subjected to myocardial infarction (MI), with
snRNA-seq performed at days 0, 3, 7, 14, and 28 post-MI. **b,** UMAP embedding of snRNA-seq profiles from
murine cardiomyocytes, identifying Myh7b+, Nppa+, and Des+ cardiomyocyte (CM) subclusters, as well as
atrial CMs. **c,** Heatmap of marker gene expression defining CM subclusters, highlighting differential
expression of contractile, stress-response, and cytoskeletal genes. **d,** Temporal dynamics of CM subcluster
abundances following MI, showing early expansion of Nppa+ CMs and progressive enrichment of Des+ CM
states during remodeling. **e,** Cross-species similarity analysis mapping murine CM subclusters to human
CM states, including HAND2+, CFLAR+, NPPA/ NPPB+, DES+, SLC25A4+, and DES_SYNPO+

cardiomyocytes. **f,** Scaled mean expression of sarcomeric gene modules (thin filament, titin/Z-disc, thick
filament) across CM subclusters, indicating progressive structural remodeling. **g,** Scaled mean expression
of β-oxidation–related genes, revealing reduced oxidative metabolic signatures in Myh7b$^+$ and Nppa$^+$ CMs
relative to Des$^+$ CM states. **h,** Nichesphere interaction network derived from Xenium spatial transcriptomics
comparing ischemic cardiomyopathy (ICM) and control myocardium, highlighting CM-centered tissue
niches enriched for immune, stromal, and vascular cell interactions. **i,** Differential spatial colocalization
scores for CM subclusters with non-myocyte cell types in ICM versus control hearts, indicating selective
niche rewiring of Nppa$^+$ and Des$^+$ CMs. For **h** and **i**, only significant interactions are shown(adjusted
Wilcoxon's rank sums test; p-value < 0.05). **j,** Enrichment of CM-associated tissue patches across
anatomical and pathological regions, including fibrotic scar, infarct border zones, necrotic tissue, and
damaged myocardium. Significance was determined by Fisher's exact test with adjusted p-values < 0.05.
k, Temporal dynamics of heart-failure–relevant drug targets across murine CM states following MI. **l,**
Expression landscape of clinically established heart-failure drug targets across human cardiac cell types,
with the "fantastic four" therapeutic classes highlighted, demonstrating preferential target enrichment in
distinct CM and non-CM compartments.

Next, we performed spatial colocalization analysis, using NicheSphere of Xenium mouse
data, contrasting controls and ICM conditions (days 7,14 and 28), which detected six
major tissue niches (**Fig. 5h**). In the mouse heart, Des$^+$ and Myh7b$^+$ CMs largely shared
a common myocardial niche (niche 6 - myocardium) indicating that diploid and polyploid
CMs coexist within the same anatomical compartments (**Fig. 5h-i**). Cells in this niche
tended to lose their colocalization to other cell-types upon diseases. By contrast, NPPA$^+$
CMs dominated a distinct niche that showed a gain of colocalization with myofibroblasts
in ICM, consistent with their transcriptional injury response program (**Fig. 5i**). This
organisation differs from the human heart, where Des$^+$ and polyploid CFLAR$^+$/HAND2$^+$
CMs segregate into two spatially distinct niches suggesting that niche partitioning of
diploid versus polyploid CMs is, at least in part, time- and species-dependent. NPPA$^+$
mouse CMs in niche 2 mapped predominantly to the "damaged myocardium 3" region in
the heart patch atlas, in which also the human NPPA_NPPB CMs enriched (**Fig. 5j**). This
underscores conserved features of the injury-associated NPPA+ states across
myocardial tissue from different mammalian species. Next we performed a PILOT
trajectory analysis on spatial niches, which indicates the decrease of myocardium (niche
1) and increase of Nppa+ (niche 2) and the myofibroblast_enriched niche (niche 3) among
others and revealing an overall increase in periostin expression fitting to our previous data
(**Fig. 5i, Extended Data Fig. 11a-b**).

Finally, we analyzed the expression of heart failure drug targets in our mouse snRNA
data (**Extended Data Fig. 11c**). This revealed a conserved, dichotomous expression
pattern: three out of four major heart-failure targets, β1-adrenergic receptor (*Adrb1*),
mineralocorticoid receptor (*Nr3c2*) and angiotensin II receptor (*Agtr1*) were predominantly
expressed in Myh7b$^+$ CMs, while *Pde5a* and *Gucy1a1* were enriched in pericytes and
vascular smooth muscle cells (**Extended Data Fig. 11c**). *Glp1r* expression was mainly
restricted to T cells. Adrenoceptor (β1) expression overall was higher in Myh7b$^+$ than in

Des[+] CMs. Comparing the dynamics of mineralocorticoid receptor (*Nr3c2*) expression,
we noted a marked expression dynamics only in Myh7b+ CMs (**Fig. 5k).** We next asked
if this pattern of HF drug target expression is recapitulated in our human snRNA data and
curated a panel of established drug targets and well-supported HF mediators and profiled
their expression across all cardiac cell subtypes (**Fig. 5l**). Of the four standard of care
drugs - β1-blockers, ACEi/ARB, mineralocorticoid-receptor (MR) antagonists, and SGLT2
inhibitors, we observed a selective enrichment of transcripts encoding two of these targets
(MR and adrenergic receptor β1) in polyploid CMs (HAND2+/CFLAR+) (**Fig. 5l, Extended**
Data Fig, 11d). Additionally, we found a significant enrichment of *GLP1R* in polyploid
CMs, which in mice was restricted to Treg and CD8-T-cells (**Fig. 5l, Extended Data Fig.**
11c). *AGTR1* was primarily expressed in cardiac pericytes and fibroblasts (**Fig. 5l**) as well
as transcripts for *PDE5A* and the soluble guanylyl cyclase α1 subunit (*GUCY1A1*).

Taken together, our HF target gene analysis indicates that polyploid CMs constitute a
major —previously unrecognized— target cell for heart failure therapies, while
pharmacologic control points for the cGMP pathway are predominantly enriched in the
vascular/perivascular compartment, and not the endothelium. In sum, our analysis
suggests a compartment-specific division of therapeutic action in human heart failure
treatment, with polyploid CMs as a major myocardial drug-addressable population.

TNIK inhibition ameliorates maladaptive remodelling by targeting polyploid CMs

Based on prior evidence implicating TNIK in WNT-dependent fibrotic programmes[78], we
examined TNIK expression across our multispecies snRNA-seq atlas and observed
consistent enrichment in CFLAR+ or Myh7b+ CMs across human, mouse, and rat
(**Extended Data Fig. 12a-d**). While TNIK expression was not strictly CM-restricted, being
detectable in human mast cells and select murine T-cell populations, the rat heart
displayed the highest CM specificity (**Extended Data Fig. 12d**). This enrichment
suggested a potential role for TNIK in regulating polyploid CM states.

To investigate the functional consequences of TNIK inhibition in CMs, we assessed its
effects both *in vitro* using human iPSC-derived CMs (**Fig. 6a-g)** and *in vivo* in a rat MI
model (**Fig. 6h-p, Extended Data Fig. 12e-j**). Long-term differentiated beating CMs from
two independent iPSC lines were treated for seven days with the TNIK inhibitor NCB-
0846[79] or vehicle and profiled by snRNA-seq. Unsupervised clustering identified two
major CM populations (**Fig. 6b**), which we annotated as mature and immature based on
enrichment analysis (**Fig. 6c**). The more immature CM showed the CFLAR signature,
while the mature cluster closely resembled DES[+] CM. ClusterFoldSimilarity[60] and a
hypergeometric test verified the similarity to DES[+] and CFLAR+ clusters from the adult
dataset (**Fig. 6d)**.

1008

Fig. 6 TNIK inhibition promotes cardiomyocyte maturation and improves post-MI cardiac function.
a, Experimental workflow for iPSC-derived cardiomyocyte (CM) differentiation and TNIK inhibition with NCB-0846. **b,** UMAP of iPSC-CMs showing segregation into immature and mature CM states. **c,** Gene set enrichment analysis highlighting immature (CFLAR-associated) versus mature (DES-associated) CM programs. **d,** Transcriptomic similarity between iPSC-CMs and adult clusters, color indicates ClusterFoldSimilarity, *p-value < 0.05 hypergeometric test. **e,** Increased proportion of mature CMs following TNIK inhibition in iPSC-CM, * indicates FDR < 0.1 for the scCODA abundance test. **f,** Pathways decreased or increased upon TNIK inhibition in iPSC-CMs. **g,** Expression of WNT pathway genes across control and TNIK-inhibited iPSC-CMs. **h,** Rat myocardial infarction (MI) study design and treatment groups. **i,** snRNA-seq of rat hearts identifying CM subpopulations after MI and treatment. **j,** Sirius Red staining and

quantitative analysis demonstrating reduced fibrosis with TNIK inhibition, scale bars indicate 4 mm.
k) Changes in cell abundance between treatment groups,* indicates FDR < 0.1 for the scCODA abundance
test. **l)** Increased proportion of mature CMs following TNIK inhibition in rats , * indicates FDR < 0.1 for the
scCODA abundance test. **m)** Compatarative enrichment analysis of selected pathways. ***p-value < 0.001.
n-p), Improved cardiac function following TNIK inhibition, including left ventricular ejection fraction,
and fractional shortening. Data are shown as mean ± s.e.m.; *p-value <
0.05, **p-value < 0.01, ***p-value < 0.001.

Cell-type abundance analysis revealed a significant increase in mature CM states
following TNIK inhibition (**Fig. 6e**). Gene set enrichment analysis demonstrated a marked
reduction in hypoxia-associated transcriptional programmes, metabolic changes and
changes in RNA transport and splicing after TNIK inhibition (**Fig. 6f**). Although global
WNT pathway activity was not significantly altered (**Extended Data Fig. 12g**), TNIK
inhibition resulted in coordinated expression changes of non-canonical WNT
components, including decreased *WNT11* and *NFATC4* expression and induction of the
WNT antagonists *SFRP1* and *SFRP5* (**Fig. 6g**). Together, these findings indicate that
TNIK inhibition promotes maturation-associated CM states alongside reduced hypoxia
signalling and selective remodelling of WNT-related transcriptional programmes.

We next investigated whether TNIK inhibition confers therapeutic benefit in the injured
heart in vivo. Rats underwent permanent LAD ligation or sham surgery and, following a
one-week recovery period, received vehicle, the clinical-stage TNIK inhibitor INS018_055
(rentosertib, NCT05938920, NCT05975983) at two doses, or the antifibrotic agent
pirfenidone as a reference therapy for 28 days (**Fig. 6h**). Rentosertib is a novel AI-
designed TNIK inhibitor whose efficacy was recently demonstrated in a phase IIa trial for
idiopathic pulmonary fibrosis[81]. Single-nucleus RNA-seq was performed on FFPE rat
heart tissue at endpoint and identified similar CM populations to the human and mouse
infarct datasets (**Fig. 6i**). TNIK inhibition resulted in a reduction of fibrosis, as shown by
sirius red staining (**Fig. 6j, Extended Data Fig. 12h**). Abundance analysis revealed a
significant, dose-dependent increase in DES+ cardiomyocytes following TNIK inhibition,
accompanied by a reduction in fibroblast and macrophage populations (**Fig. 6k-l**).
Furthermore, treatment with both doses of rentosertib resulted in a marked decrease of
hypoxia and TGFβ signalling pathway in CMs (**Fig. 6m**), while WNT signatures did not
significantly change between any of the groups. iPSC-derived CMs and rat CM shared a
large overlap in genes that were down-regulated upon treatment. More than 400 hypoxia-
associated genes were decreased in both cohorts, including transcripts implicated in
cross-talk between HIF-1α signalling and WNT/β-catenin or TGF-β pathways, such as
KCTD11 (REN), SEMA4B, PDK1, TGFBI and ARRDC3. Rats treated with rentosertib
additionally demonstrated improved cardiac function as detected by echocardiography,
which reached comparable levels to the group treated with antifibrotic control pirfenidone
(**Fig. 6n-p, Extended Data Fig. 12i-j**). Collectively, the concordant human iPSC-CM and
rat MI data demonstrate that pharmacological TNIK inhibition mitigates maladaptive post-
infarction remodelling, in part by shifting (*in vitro*) or leading (*in vivo*) to increased mature

CM transcriptional states and suppressing hypoxia- and fibrosis-associated signalling
programmes. Together, these results support the concept that pharmacological
modulation of disease-associated polyploid CM states can favorably modulate post-
infarction remodelling of the heart.

Discussion

In this study, we provide a multi-scale, multimodal view of how CM heterogeneity, ploidy
state and microenvironmental context shape cardiac remodeling in human heart failure.
We integrate single-nucleus transcriptomics, chromatin profiling, high-resolution spatial
transcriptomics and histology across ischemic and non-ischemic cardiomyopathies,
human development, and in three mammalian species. Our analysis revealed that
disease progression is not only encoded in changes of cell-type composition and fibrosis,
but in the emergence of distinct diploid and polyploid cardiomyocyte states that shape
spatial niches and drive cardiac remodeling.

We systematically dissect diploid versus polyploid CMs in the human heart. Using DAPI-
based nuclear profiling, FANS-enriched ploidy fractions and integrative snRNA-seq, we
show that polyploid CMs form distinct transcriptional states, marked by CFLAR and
HAND2, which are enriched in 4N/8N nuclei, engage specific metabolic and chromatin-
remodeling programs, and display a disease-biased remodeling signature. In contrast,
predominantly diploid DES$^+$ CMs preserve oxidative phosphorylation–dominated
metabolic profiles and mature sarcomere programs. Direct comparison to human fetal
hearts indicates that HAND2$^+$ and CFLAR$^+$ polyploid CMs align with proliferating and
immature fetal states, whereas DES$^+$ and DES_SYNPO$^+$ CMs resemble terminally
differentiated, contractile end states. Together, these findings provide a cellular and
ploidy-based resolution to the fetal gene hypothesis. We demonstrate that developmental
reactivation in human heart disease is partial, selective, and strongly shaped by CM
ploidy, rather than representing a generic fetal-like state.

At the tissue level, CM states organize into distinct spatial niches that differ between
ischemic and non-ischemic cardiomyopathies. Polyploid CFLAR$^+$/HAND2$^+$ CMs
preferentially localize to healthy-appearing myocardium and damaged myocardium,
whereas DES$^+$ and DES_SYNPO$^+$ states colocalize with myofibroblasts and tissue-
resident macrophages in fibrotic and chronically remodeled regions. The use of deep
learning models for histology analysis revealed distinct damaged cardiomyocyte regions
conserved in both mouse and human, that could not be identified even by expert
pathologists. These distinct areas have differences in both their abundance of damaged
NPPA_NPPB CMs and lymphocytes.

Subcellular spatial transcriptomics further reveals a discrete ICD transcriptome
compartment, enriched for mRNAs encoding junctional regulators, cytoskeletal linkers

and local mRNA-regulatory components. Differential analysis of this ICD program across
cardiomyocyte states and ploidy suggests that altered mRNA localization and junctional
remodeling may contribute to conduction and mechanical dysfunction in heart disease.
All these technical advances are released as community resources, i.e. machine learning
models that empower researchers to segment CMs in microscopy and spatial
transcriptomic datasets, identify regions of interest in H&E sections, and detect
cardiomyocyte polyploidy from DAPI images.

Ligand–receptor and drug-target mapping position these CMs states within a structured
signaling and therapeutic landscape. Polyploid CFLAR$^+$ and HAND2$^+$ CMs act as major
receivers and senders of distinct developmental and fibrotic cues, with enriched Notch,
non-canonical Wnt and Hedgehog receptors, as well as TGFβ and ERBB signaling
components. More differentiated DES$^+$, NPPA_NPPB$^+$ and SLC25A4$^+$ CMs preferentially
express Wnt antagonists.

Analysis of heart failure drug targets demonstrated that molecular targets of standard-of-
care therapies, such as β-blockers and mineralocorticoid receptor antagonists, show
marked enrichment in polyploid CMs. These data suggest a compartmentalized division
of therapeutic action and highlight polyploid CMs as a major drug-addressable myocardial
population in patients.

This study is limited by the observational nature of human end-stage cardiac tissues,
targeted spatial gene panels, and reliance on transcript-based inference for signaling and
drug-target activity. In addition, ploidy assignment is based on DAPI content and
enrichment rather than direct lineage tracing. Furthermore, our pharmacological
perturbation of TNIK in an *in vivo* model is not CM-subtype specific. Future work
combining CM-subtype-specific genetic perturbations, higher-order multiplexing spatial
profiling, and functional assays of electrophysiology and force generation will be required
to test causal roles of individual CM states and to define how genome multiplication
mechanistically constrains -or enables- cardiomyocyte plasticity.

Finally, through cross-species analyses we nominated TNIK as a central regulator in
polyploid cardiomyocyte remodeling. TNIK inhibition in a rodent model of myocardial
infarction ameliorated remodelling, reduced fibrosis and improved myocardial function.
We anticipate that these insights will serve as a reference to dissect when and how diploid
CMs transition into maladaptive polyploid states, and to guide the design of interventions
enabling the development of novel cardiac therapeutic strategies, such as TNIK inhibition,
that selectively reprogramming pathological CM states to restore myocardial function.

Methods

Experimental procedures

All human tissue protocols were approved by the local ethics committees of Ruhr University Bochum (Bad Oeynhausen; Reg.-No. 21/2023), RWTH Aachen University (EK 24-337), University of Chicago (STU00216865), University Medical Center Utrecht (UCC-UNRAVEL #12-387; TCBio 25-242), Amsterdam University Medical Center (METC 2016_285; Dutch Fetal Biobank[81]), and Washington University in St. Louis (No. 201104172). Human myocardial tissue was obtained from non-transplanted donor hearts, from patients with myocardial infarction undergoing heart transplantation, and from patients receiving a total artificial heart or left ventricular assist device (LVAD) implantation. The study complied with the Declaration of Helsinki and with the Dutch Code of Conduct for the Responsible Use of Human Tissue. All participants provided written informed consent. This study was conducted within the framework of the UCC-UNRAVEL biobank[82] (University Medical Center Utrecht; www.unravelrdp.nl). Tissue collection was approved by the scientific advisory boards of the UMC Utrecht Biobank (UNRAVEL) and the Amsterdam University Medical Center (Dutch Fetal Biobank).

Human tissue processing and screening

Mainly FFPE heart tissues were used. Most samples were obtained between 2021 and 2023. 5 μm tissue sections were performed and the H&E stainings were analyzed by a cardiac pathologist. For RNA quality control, two scrolls of 25 μm tissue sections were used as input for the Qiagen FFPE RNA isolation kit. DV200 measurements were subsequently performed using TapeStation (Agilent). For MERFISH spatial transcriptomics (Vizgen) all samples underwent QC and only samples with a DV200 value of 30% or higher were included. For Xenium and Flex (10XGenomics) no DV200 measurements were performed. For H&E and IF imaging a Zeiss Axioscan was used.

Animal studies

Mouse experiments were performed according to the animal welfare guidelines and German national laws and were authorized by the local authority (Regierungspräsidium Darmstadt, Hessen, Germany and Baden-Württemberg, Germany) under the animal experiment license FU/2084. C57BL/6J wildtype animals were obtained from Janvier Labs. Female and male mice were used at the age of 10 to 14 weeks. Myocardial infarction was induced by ligation of the left anterior descending coronary artery as described previously[83]. Experiments were performed in both females and males. Animals were held at 23°C ambient temperature and 60% humidity in 10 h/14 h light/dark cycle. At the end of each experiment, animals were euthanized by cervical dislocation after isoflurane inhalation (3–3.5 vol%).

1167

Rat studies were performed by WuXi AppTec (Nantong, China, IACUC protocol number
GP02-QD079-2023v1.2). Sprague–Dawley rats (male, 6-8 weeks, purchased from Vital
Rivers Animal Technology Co., Ltd., weighted around 200-300 g) were obtained and
housed at the WuXi AppTec (Nantong) facility under controlled conditions (22–25 °C, 40–
70% humidity). A timed lighting system maintained a 12 h light/12 h dark cycle (06:00–
18:00). Rats had ad libitum access to chow and water. Left anterior descending coronary
artery (LAD) ligation was performed on SD rats to induce heart failure. On the day of
model induction, rats were anesthetized with a combination of Zoletil (50 mpk, i.m.) and
xylazine (8 mpk, i.p.). The trachea was connected to a ventilator for assisted respiration.
A thoracotomy was performed between the third and fourth ribs; the pericardium was
opened, and the LAD was ligated with 6-0 silk suture. The ribs and skin were closed with
4-0 silk sutures, and rats were placed on a heating pad postoperatively until recovery
from anesthesia. In the sham group (G1), all procedures were identical except the ligation
of the LAD. Animals were administered meloxicam (1 mg/kg, i.m.) and gentamicin
hydrochloride (8 mg/kg, i.p.) postoperatively for analgesia and infection prophylaxis. One
week later, successfully modeled rats (30% or more reduction in LVEF) selected by
echocardiographic measurement, were grouped by LVEF into four groups G2 (Model),
G3 (INS018-055, 2.5 mg/kg, BID), G4 (INS018-055, 5 mg/kg, BID), and G5 (pirfenidone,
350 mg/kg, QD). Echocardiography was performed on Vevo®1100 (Fuji), readout
includes Left ventricular ejection fraction (LVEF), fractional shortening (FS), left
ventricular end-diastolic diameter (LVEDD), left ventricular end-systolic diameter
(LVESD), left ventricular end-diastolic volume (LVEDV), left ventricular end-systolic
volume (LVESV), stroke volume (SV), cardiac output (CO), left ventricular mass (LV
mass), left ventricular anterior wall thickness in systole/diastole (LVAWTs/d), and left
ventricular posterior wall thickness in systole/diastole (LVPWTs/d). After grouping, rats
received vehicle or test drugs by oral gavage for 4 weeks: G1-G4 were dosed twice daily
(12 h apart), and G5 once daily. INS018-055 was given in 50 mM citrate buffer,
pirfenidone in 0.5% CMC-Na. General condition was monitored and abnormalities
recorded throughout. After 4 weeks of treatment, endpoint echo measurements and
sample collection were performed. The hearts were fixed in 4% paraformaldehyde (PFA)
and subjected to Sirius Red (SR) staining to evaluate myocardial fibrosis. Sirius red
staining was analyzed by Halo AI 3.6. 4 samples each from groups 1-4 were then selected
for snRNA processing.

Human iPSC culture

WTC-11 (Male, Coriell, GM25256) and AICS-TTN (Male, Coriell, AICS-0048-039) human
iPSC lines were used in this study. iPSCs cultures were maintained in Matrigel coated
plates (Corning, 356231) in mTeSR™ Plus medium (STEMCELL, 100-0276) and
passaged upon reaching 70% confluency. Briefly, iPSCs, with no signs of differentiation,

were dissociated chemically using TryPLE Select (ThermoScientific, 12563029) and
seeded at 1:4 ratio onto Matrigel-coated plates with mTeSR™ Plus medium,
supplemented with 10 µM Y-27632 dihydrochloride (MedChemExpress, HY-10071).
iPSCs cultures were regularly tested for mycoplasma contamination.

Human iPSCs derived cardiomyocyte differentiation protocol

iPSCs cultures at 70-80% confluency and with no apparent signs of spontaneous
differentiation, were differentiated to cardiac lineage by following established
conditions[84,85]. Briefly, to promote gastrulation on the iPSC cultures at Day 0, the culture
media , switched to RPMI 1640 (Gibco, 21875034) basal media supplemented with 1%
penicillin/streptomycin, 2% B27 supplement minus insulin (Gibco, A1895601, hereafter
referred to as RPMI/B27−), and 6 µM CHIR99021 (Cayman Chemical, 13122). Culture
was directed on Day 2 towards cardiac mesodermal lineage, by supplementing to
RPMI/B27− basal media with 5 µM IWP2 (Tocris, 3533). On day 4, the medium was
changed to RPMI/B27−. On day 6 and 8, the medium was changed to RPMI 1640 basal
media supplemented with 1% P/S and 2% B27 supplement (Gibco, 17504044; hereafter
referred to as RPMI/B27+). Cardiomyocytes were enriched until day 10, by switching the
medium to glucose-free RPMI 1640 (Gibco, 11879020) supplemented with 0.1% (v/v)
sodium DL-lactate (Sigma, L4263-100 mL). On day 13, the medium was reverted to
RPMI/B27+, and thereafter, medium changes were performed every 2 days.

TNIK perturbations on hiPSCs derived cardiomyocytes

At day 50 of differentiation, beating cardiomyocyte cultures were treated with 1 µM NCB-
0846 (MedChemExpress, HY-100830), with DMSO-treated cultures serving as controls,
for 7 days.The medium was changed every two days[78,86]. On day 7 of treatment, iPSC-
derived cardiomyocytes (iPSC-CMs) were mechanically harvested for nuclei isolation.
Nuclei were isolated using the 10x Nuclei Isolation kit (10XGenomics) with bulk
cardiomyocyte cultures handled according to the tissue protocol.

Tissue dissociation and nuclei isolation

Two to three 50 µm sections were taken from FFPE samples and stored at 4°C until
processing. To isolate single nuclei from FFPE tissue, samples were deparaffinized with
Xylene (3x) for 10 min each to remove the paraffin (Sigma-Aldrich, 108298) and
rehydrated in an decreasing ethanol concentrations (100%, 70%, 50%, 30%) (prepared
from Fisher Scientific, BP2818500), nuclease-free water (invitrogen, 10977035) and 1X
PBS (Gibco, 11594516). To digest the tissue sections, they were processed with
digestion buffer (1.68 ml RPMI 1640 medium (Thermo Scientific, A1049101), 420 µl
Liberase TH (final concentration 1mg/ml, Roche, 5401135001), 21 µl Collagenase D (final
concentration 1mg/ml, Roche, 11088858001), 8.4 µl RNase inhibitor (final concentration
0.2 U/µl, New England Biolabs, M0314L)) on the gentleMACS™ Octo Dissociator

(Miltenyi Biotec, 130-096-427) with the 37C_FFPE_1 protocol. After digestion, the
samples were transferred to glass tubes, cell lysis mix was added (796 µl nuclei EZ lysis
buffer (Sigma Aldrich, N3408), 200 µl 10% BSA (Miltenyi Biotech, 130-091-376), 4 µl
RNase inhibitor), pestled with A and B (10-15x), and incubated for 5 min to dissociate the
tissue further. Afterwards, the nuclei/cell solution was filtered with a 70 µm filter (PN:
431007060, PluriSelect) and washed. To reduce debris, a density centrifugation step was
performed at 700 rcf for 10 min at 4°C in debris removal buffer (550 µl debris removal
reagent, PN-2000560, Chromium Nuclei Isolation Reagents (PN-1000447), 10X
Genomics), 0.55 µl reducing agent B (10X Genomics, Reducing Agent B (PN-1000450),
PN-2000087)). The nuclei suspension was then washed and resuspended in
resuspension buffer (2733.5 µl 0.5X PBS, 5.5 µl 10% BSA, 11 µl RNase inhibitor). Nuclei
concentration was determined by staining an aliquot of the nuclei suspension with
ethidium homodimer 1:100 (Invitrogen, E1169) and counting on the Invitrogen™
Countess™ 3 FL Automated Cell Counter (Fisher Scientific, 16832556). Only samples
with over 100,000 nuclei were input for probe hybridization.

Single-Nucleus RNA- sequencing (Flex)

1258

Single-nucleus RNA sequencing (snRNA-seq) was performed following the 10x
Genomics protocol CG000527 (Rev. E). Fixed nuclei or cells were hybridized and
incubated overnight (16-24 h) with mRNA transcript probe sets targeting approximately
18,000 genes per species, with probes carrying a sample-specific barcode in the left
handle. After hybridization, nuclei were counted and pooled in equal proportions, followed
by multiple washes with post-hybridization wash buffer (10X Genomics) to remove
unbound probes. The pooled nuclei were then counted again, and a volume
corresponding to a target recovery of 20,000 nuclei or cells per sample was combined
with a master mix containing enzymes for probe ligation and extension during GEM
incubation. The reaction mixture was loaded onto a NEXT GEM Chip Q together with gel
beads and partitioning oil, and GEMs were generated using the Chromium X system.
Following GEM recovery, cDNA was amplified and used for library preparation. A fraction
of the amplified material was subsequently re-amplified using library-specific P5 and P7
indices.

QC and Sequencing

1273

1274 All libraries were quantified with the Qubit fluorometer's high sensitivity dsDNA reagents
1275 (Thermo Fisher Scientific) and sized on the Agilent 2200 Tapestation with high-sensitivity
1276 tapes and reagents. All libraries were run on the NextSeq2000 and NovaSeq X (Illumina).
1277 Sample sequencing was conducted to achieve a depth of 20,000 reads per nuclei/cell as
1278 recommended by 10XGenomics.

Fluorescent-activated nuclei sorting (FANS)

Nuclei were stained with DAPI (1:10,000, 1 mg/dl stock, Sigma-Aldrich) and diluted in 1X PBS and analyzed on the FACS SH800 (SONY) using standard laser configuration and a 100 µm microfluidic chip (SONY) for sorting. Events were first gated on FSC and SSC profiles, followed by singlet gating using FSC-H and FSC-A and subsequent gating on DAPI as indicated. Sorts were completed with semi-purity mode, and a resort was always performed to check sorting efficiency.

MERFISH

FFPE samples were sectioned to 4.5 µm, placed on a functionalized slide (Vizgen, 20400100) and gently pressed onto the slide to enhance tissue adhesion. Subsequently, the slide was dried at room temperature for 60 minutes, then baked at 55°C for 30 minutes and placed at room temperature again for 3 hours before storage at 4°C until use. After equilibrating the slides to room temperature, deparaffinization was performed with deparaffinization buffer (Vizgen, 20300112) in two cycles of 55°C for 5 minutes, followed by a single cycle at room temperature for 5 minutes. The tissue was rehydrated by three washes with 100% ethanol (Fisher Scientific, BP2818500) and one consecutive wash with 90% and 70% ethanol. The slides were dried at room temperature for 30 minutes, then incubated with decrosslinking buffer (Vizgen, 20300115) at 90°C for 15 minutes. Before further processing, FFPE samples were split into MERFISH chemistry V1 and V2. For V1, conditioning buffer (Vizgen, 20300116) was added to the slide for 30 minutes at 37°C. For pre-RNA anchoring, 5 µl pre-anchoring activator (Vizgen, 20300113) and 5 µl RNase inhibitor (Roche Diagnostics, 09537643103) were diluted in 100 µl conditioning buffer and incubated on the tissue for 2 hours. Multiplex protein staining steps were performed before adding formamide wash buffer (Vizgen, 20300002) at 37°C for 30 min and an overnight incubation in 100 µl RNA anchoring buffer (Vizgen, 20300117). RNA anchoring buffer was removed from the tissue by adding formamide wash buffer at 47°C for 15 min and sample prep wash buffer (Vizgen, 20300001) afterwards. Polyacrylamide gel was prepared by mixing 5 ml gel embedding premix (Vizgen, 20300118) with 25 µl 10% ammonium peroxodisulfate (Karl Roth, 9592.2) and 5 µl N,N,N′,N′-Tetramethylethylendiamin (Sigma-Aldrich, T7024-25ML). The slides were coated in gel solution for 5 min, then the solution was removed and 50 µl gel solution added to the center of the tissue. The gel was evenly spread over the tissue with a coverslip (Vizgen, 30200004) coated with gel-slick solution (Lonza, 50640) and incubated until fully polymerized. Afterwards, the coverslip was carefully removed and the tissue digested overnight at 47°C by adding 5 ml clearing premix (Vizgen, 20300114) with 50 µl Proteinase K (New England Biolabs, P8107S). To reduce tissue autofluorescence, the tissue was bleached with blue LED light for 3 hours (Vizgen, 10100003). After three washes with sample prep wash buffer and one with formamide wash buffer at 37°C for 30 min, 100 µl custom gene probes were added onto the gel and spread evenly by placing

a small piece of parafilm on top. Probe hybridization with V1 probes was performed for
48 h at 37°C. To remove unbound gene probes, two washes with formamide wash buffer
were performed at 47°C for 30 min each, followed by two washes in sample prep wash
buffer. MERFISH chemistry V2 provided an improved RNA detection rate and overall
increased sensitivity by optimizing anchoring mechanisms, gene probes and augmented
readout-probes. For this version, cell staining was performed immediately after
decrosslinking, followed by an anchoring pretreatment with a conditioning buffer (Vizgen,
20300116) incubation for 15 min and a pre-anchoring buffer incubation (100 μl
conditioning buffer, 5 μl pre-anchoring activator (Vizgen, 20300113), 5 μl RNase inhibitor)
overnight. RNA anchoring was performed the following day for 2 hours at 37°C. The
samples were then embedded into gel and cleared with tissue clearing solution at 47°C
overnight. After autofluorescence quenching and probe hybridization with V2 probes, an
additional enhancer hybridization step was added after the post-hybridization wash with
formamide buffer. For this, pre-warmed enhancer (Vizgen, 30300491) was added onto
the embedded sample and incubated at 37°C overnight, followed by two additional
washing steps with enhancer wash buffer (Vizgen 20300192). All slides were
consecutively stained with DAPI and PolyT staining reagent (Vizgen, 20300021) in the
dark for 15 min, then washed again with formamide wash buffer and sample prep wash
buffer before proceeding to imaging.
1337
Fresh frozen samples were cut to 10 μm sections, placed on functionalized slides (Vizgen,
20400001) and equilibrated to -20°C for 5 min to improve tissue adherence. Then, slides
were dried at 47°C for 3 min and immediately fixed with pre-warmed 4% PFA (Electron
Microscopy Sciences, 15714-S) at 47°C for 30 min. PFA was removed by washing in 1X
PBS (Gibco, 11594516) three times, then the tissue was permeabilized in 70% ethanol
at 4°C overnight. After a wash with 1X PBS, multiplex protein staining was performed.
The slide was then incubated in formamide wash buffer at 37°C for 30 min before adding
a 50 μl custom gene panel and placed at 37°C for 48 h for probe hybridization. Excess
probes were removed by two incubations in formamide wash buffer at 47°C for 30 min
each, before the tissue was embedded in a polyacrylamide gel prepared from gel
embedding premix (Vizgen, 20300004) and cleared in 5 ml clearing premix (Vizgen,
20300003) with 50 μl Proteinase K at 47°C overnight. Photobleaching for 3 hours was
initiated the next day. Analogous to the FFPE sample preparation, the slides were washed
with sample prep wash buffer, before staining with DAPI and PolyT in the dark for 15 min
and a final wash in formamide wash buffer and sample prep wash buffer before imaging.
1353
For multiplex protein staining, the slides were blocked with blocking buffer C premix
(Vizgen, 20300100) for 60 min before adding the primary antibody mix (Anti-Desmin 1:50
(Novus Biologicals, AF3844), Anti-Myeloperoxidase 1:100 (Abcam, ab300651), Anti-
Vimentin 1:100 (Abcam, ab24525), cell boundary primary stain mix 1:100 (Vizgen,

20300010) diluted in blocking buffer C premix) for 60 min. Unbound primary antibodies
were washed away with 1X PBS before secondary antibody mix (Anti-goat Aux 6 protein
stain 1:100 (Vizgen, 20300103), Anti-rat Aux 7 protein stain 1:100 (Vizgen, 20300104),
Anti-chicken Aux 9 protein stain 1:100 (Vizgen, 20300106), cell boundary secondary stain
mix 1:33 (Vizgen, 20300011) diluted in blocking buffer C premix) was added onto the slide
for 60 min. After further washing with 1X PBS, a second blocking step with diluted rabbit
IgG (Thermo Fisher, #02-6102) was performed for 30 min. During this time, a conjugate
of Anti-Connexin 43 (Proteintech, 26980-1-AP) and a custom 5'acrydite-Aux 5-
conjugated Anti-rabbit nanobody (Massive Photonics) was prepared and diluted in
blocking buffer C premix to a final antibody concentration of 1:100. After washing away
excess rabbit antibody with 1X PBS, the antibody-nanobody mix was incubated on the
tissue for 60 minutes. Rnase inhibitor was added to all steps to ensure RNA preservation.
Afterwards, more washes with 1X PBS and a light fixation with 4% PFA for 15 min were
performed. The slide was then washed again with 1X PBS and sample prep wash buffer.
Imaging Imaging cartridges (Vizgen, 20300019) were activated by adding a mix of 250 µl
imaging buffer activator (Vizgen, 20300022) and RNase inhibitor (New England Biolabs,
M0314L) to the cartridge port and layering mineral oil (Sigma-Aldrich, M5904-500ML) on
top to avoid oxidation. The slides were placed into the flow cell, both imaging cartridge
and assembled flow cell were connected to the instrument. After ROI selection and focus
acquisition, the measurement was initiated and processing of the raw data performed
according to Vizgen's proprietary processing pipeline.

Xenium spatial transcriptomics

Spatial in situ sequencing data of human and mouse heart tissues were generated using
the Xenium Prime In Situ Gene Expression workflow for FFPE tissue (10X Genomics)
with the Xenium Prime 5K Human or Mouse Pan Tissue & Pathways Panel (10X
Genomics, 1000724/1000725, RevE). Samples were sectioned to 5 µm and carefully
placed in the sample area of Xenium slides following the Tissue Preparation Guide (10X
Genomics, CG000578). Slides were dried for 3 h at 42°C, then placed in a desiccator at
room temperature overnight. Xenium slides were fixed and permeabilized according to
the Xenium In Situ Gene Expression Fixation and Permeabilization protocol (10X
Genomics, CG000580, RevE) prior to probe hybridization, ligation, rolling circle
amplification and cell segmentation staining according to the manufacturer's instructions
in Xenium In Situ Gene Expression with optional Cell Segmentation Staining (10X
Genomics, CG000760, RevA). For human heart samples, a custom add-on probe panel
was added to the priming hybridization and probe hybridization step. Slides were imaged
on the Xenium Analyzer (10X Genomics, CG000584, RevF) and an H&E staining was
performed afterwards following the manufacturer's recommendations (10X Genomics,
CG000613, RevB). A post-Xenium WGA-FITC staining was performed by incubating
1:100 WGA-FITC on the Xenium-slide in 1X PBS for 30 min, followed by washing in 1X

PBS for 3x 5min. Imaging was performed using an AxioScan (Zeiss) at a magnification of
20x.

Bioinformatics Analysis

Data Acquisition

Spatial transcriptomics data was acquired with onboard software versions 234c
(MERSCOPE) and 4.0.1 (Xenium). Microscopy images were acquired with Zeiss Zen
3.1.9. snRNA data was aligned with Cellranger 7.1.0. Reads were aligned to the human
reference genome GRCh38 using the pre-built reference package version 2024-A
provided by 10x Genomics. This reference incorporates the GENCODE v44 annotation
and the GRCh38 primary assembly sequence. Reads were aligned to the mouse
reference genome GRCm39 using the pre-built reference package version 2024-A
provided by 10x Genomics. This reference incorporates the GENCODE vM33 annotation
and the GRCm39 primary assembly. Single-nucleus RNA-seq data were aligned to the
Rat reference genome mRatBN7.2 with annotations from Ensembl Release 110.

snRNA processing

To preprocess and analyse the snRNA-seq dataset we created a standardised
Snakemake[87] pipeline that includes the following steps: ambient RNA detection and
removal with SoupX[88], quality control and normalisation with Scanpy[89], doublet detection
with scrublet[90] and doubletdetection[91], integration with Harmonypy[92] and differentially
expressed marker genes analysis with Presto[93]. Each dataset underwent identical
preprocessing, followed by a semi-manual cell type annotation with support from
CellTypist[94]. Human, mouse and rat data annotations were harmonized on the cell type
level, while cell state annotation remained dataset specific.

1420

Cells were filtered using a multi-step quality control procedure. First, cells expressing
fewer than 2 genes were removed. Subsequently, outlier cells were identified using a
median absolute deviation (MAD)-based approach across three quality metrics: total
counts (log-transformed), number of genes detected (log-transformed), and percentage
of counts in the top 20 genes. For each metric, cells deviating more than 5 MADs from
the median were flagged as outliers (similar to Germain et al.[95]). The SoupX ambient RNA
detection was then applied with default parameters with the exceptions of the iPSC
snRNA-seq data due to the extremely homogeneous nature of the cell line data, as
suggested by the package authors. For the integration with Harmony, we selected 4000
highly variable genes across samples using a batch-aware approach with the unique
sample identifier provided as the batch key. Expression counts were normalised to the
total library size, log-transformed, and scaled without zero-centring. Then we performed
principal component analysis (PCA) using the default number of 50 principal components

(PCs) on the scaled data and used Harmony for the batch correction with the sample
label as the integration variable and a maximum of 100 iterations. Nearest-neighbour
graphs were constructed using the Harmony-corrected principal components. Cell
clustering was performed with the Leiden algorithm at two resolutions 1.5 and 3 for further
manual annotation and cluster assignments were stored separately. Then we generated
two-dimensional embeddings using PaCMAP and UMAP applied to the Harmony-
corrected PCA space. We performed automatic cell type annotation using CellTypist on
the integrated dataset, following the authors instructions. Additionally, we calculated cell
cycle phase scores for S-phase and G2/M-phase gene expression using predefined gene
sets[96].

We identified marker genes of the integrated datasets for both clustering resolutions using
a two-sided Wilcoxon rank-sum test implemented in the Presto framework. Genes were
ranked based on the Wilcoxon AUC statistic. Marker genes were filtered to retain the top
100 genes per cluster with an AUC of at least 0.5, expression in all cells of the cluster,
and an adjusted p-value below 0.05. For all datasets during the initial manual cell type
annotation process we used overclustered resolution (leiden res = 3) to annotate major
cell types and removed all small low quality, high in doublets subclusters that had a mixed
marker gene signatures. Additionally, during the preprocessing of the Embryo human
heart snRNA-seq dataset we first annotated all cell types then subsetted, reclustered and
reannotated only CMs to achieve higher annotation accuracy. During this step some low-
quality, high in doublets mixed clusters were removed.

Enrichment analysis and gene expression comparison

All gene expression comparisons (e.g. expression heatmaps) were created from
normalized, log1p transformed UMI matrices (see processing details above). Differential
gene expression and enrichment analysis was performed on the pseudobulk level with
edgeR[97]. Pseudobulks were formed at the donor and cell type level as required by the
individual analysis and were based on the sum of the count matrix after ambient RNA
removal. Enrichment analysis was performed using the competitive gene set test
CAMERA[98] implemented in edgeR[97] with default settings. The enrichment (t-statistic) of
selected gene sets was calculated with a univariate linear model as implemented in the
python version of the decoupler library[99] based on differential genes calculated by edgeR
(quasi-likelihood negative binomial generalized log-linear model). Gene sets were
retrieved from msigdb[100] and PROGENy[101]. Complete enrichment scores are given in
Supplementary Table 2.

Gene signature scoring per nucleus

1470 We computed per-nucleus signature scores using Scanpy's gene scoring function for
1471 DES[+] and CFLAR[+] CMs. Then we analyse the signature scores in relation to cellular

1472　ploidy using the polyploid annotated data. Pairwise comparisons between ploidy groups
1473　2N, 4N, and 8N were performed separately for each signature using the two-sided
1474　Wilcoxon rank-sum test.

Abundance testing

Cell-type abundance differences were analyzed using scCODA, a Bayesian framework
for compositional single-cell data[102]. In all analyses, control samples were specified as
the reference condition, resulting in direct comparisons between control and disease
samples, as well as between untreated and treated iPSC samples. For human and mouse
snRNA-seq data, iPSC derived samples and rat samples, credible changes in cell-type
abundances were identified using an estimated false discovery rate (FDR) of 0.1.
Changes in cell-type abundances were quantified as scCODA-estimated log2-fold
changes relative to control samples and visualized using heatmaps with markers
indicating credibly changing cell types.

Comparison of cardiomyocyte substates across data sets

We use two complementary approaches to contrast cardiomyocyte sub-states/clusters
across novel (human adult, human embryo, mouse, rat) and public data sets[59] The first
approach, ClusterFoldSimilarity[60], quantifies the similarity of cell clusters across datasets
by their gene fold changes measured against the remaining clusters within the respective
dataset and comparing these to the fold change in clusters from a reference dataset[60].
As the resulting similarity score is dependent on the specific context, we refrain from
comparing the magnitude of specific values across pairs of considered datasets and only
interpret these as indicators of matching or distinct cell types.

As a second approach, we perform a one-sided hypergeometric test to measure the
statistical significance of the overlap between cell markers for each cluster[103]. Based on
the hypergeometric distribution, the test calculates the probability of observing at least as
many overlapping genes by chance when drawing two random sets from the shared
background of genes expressed in at least 5% of CMs in the compared datasets. To
perform this test, the differentially expressed genes of the subtypes are ranked within
each considered dataset (tie-corrected two-sided Wilcoxon test). The resulting top 500
differentially expressed genes of each CM subtype are then compared pairwise across
all datasets to determine the number of matching genes. The match of top ranked genes
between two subtypes is considered as significant, if the random chance of obtaining at
least as many matching genes under the hypergeometric distribution is less than
alpha=0.01. As we are only interested in contrasting CM populations, all marker genes
and fold similarities described above are based on contrasting only CM cells.

Spatial transcriptomics panel design

1507

An initial MERFISH V1 500 gene panel was designed based on publicly available scRNA
datasets of the human heart and human heart disease[11,104–108] in addition to canonical
marker genes. Cell type annotation was harmonized between public datasets. We then
selected the most informative genes using the Spapros[1] probe selection pipeline. The
selection procedure was carried out in three steps. Spapros was first run jointly on all
datasets, selecting informative genes based differential expressed (DE) gene statistics.
Afterwards, the tool was run on the individual datasets to select informative genes for cell
states based on PCA loadings. The performance of the panel was confirmed by using
Spapros' evaluation mode which trains a random forest classifier to distinguish cell types
in scRNA data based on subsets of genes. In cases where cell types were commonly
confused by the classifier (i.e. the correct cell type was selected less than 60% of the
time) additional genes were selected with NSForest[2], a marker gene selection tool based
on a random forest classifier. To reduce the chance of optical crowding the panel adhered
to an overall FPKM (Fragments Per Kilobase Million) limit of 9000 in the manufacturers
reference RNAseq dataset. Genes with very high FPKM were replaced with genes that
showed similar expression patterns but lower FPKM values. This replacement was based
on Pearson correlation values.

1525

We generated a second panel for MERFISH (v2) by including genes related to newly
discovered CM states, which were not previously defined in the previous analysis. For the
MERFISH v2 dataset the procedure was as follows: The snRNA dataset was subset to
only cardiomyocytes. Cells were then assigned to the major clustered CFLAR+ and DES+
and marker genes were selected with the tie-corrected two-sided Wilcoxon rank sum test,
as implemented in Scanpy[89]. The bottom 15 genes in the initial panel, as ranked by
importance according to Spapros, were replaced with CM state marker genes. For the
MERFISH v2 panel we added an additional step to further decrease the chance of optical
crowding occurring. The final MERFISH v2 codebook, which assigns binary barcodes to
transcripts[109], was shuffled in a way where highly expressed genes in the same cell type
were not hybridized with fluorescent probes during the same detection round. This
optimization procedure[110] relied on simulated annealing and used our human snRNA
dataset as input. All MERFISH panels were purchased from Vizgen Inc.

1539

For the Xenium experiments (10X Genomics) we used the preconfigured 5000 pan tissue
panel in the mouse and human variant (PO 1000725 and 1000724). A custom 100 gene
human addon panel optimized for cardiomyocyte state differentiation. The procedure was
the same as for MERFISH, but only considered 100 top genes. This panel was added to
all measurements of human samples. A detailed description of all panels can be found in
Supplementary Table 3.

Segmentation of spatial transcriptomic datasets

For MERFISH, the segmentation of heart muscle cells was extensively optimized. We fine-tuned a Cellpose 3 cell membrane model, "heartbreaker", on 216 (1000 x 1000 px) expert annotated images from our MERFISH measurements, specifically Vizgen's "Cell Boundary Stain 3" and the DAPI channel. Images were split into 200 training and 16 test images. The model was initialized with "cyto3" weights and then trained with a learning rate of 0.1 and a weight decay of 1e-05 for 200 epochs. In a benchmark we compared the performance of our model to several image and transcript-based segmentation algorithms. For image-based we considered the default MERSCOPE segmentation based on Cellpose 2, Cellpose 3 "Nuclei" model and our model. We additionally tested a configuration where the result of heartbreaker was combined with the output of the "Nuclei" Cellpose 3 model.

All image-based tools were run on boundary stain 3 from the Vizgen MERSCOPE cell boundary stain kit. Transcript-based segmentation methods tested included Baysor[111], BIDCell[112] and Proseg[113]. Performance was judged by comparison to a total of 10 hand-annotated image sections (1500 x 1500 px) from two MERFISH datasets. Tools were scored based on how similar their output was to the expert annotation. For this purpose the aggregated jaccard index (AJI)[114] and mean average precision at intersection over union of at least 50% (mAP IoU 50%) were used. We further ranked tools on the percentage of all transcripts assigned to cells and based on the number of segmented cells in the two tested MERFISH samples. As the total true number of cells in a sample is unknown, we compared the number of output cells to the number retrieved from nuclei segmentation. The absolute value of the difference between the estimated total number and number output by a tool was min-max scaled between the tools.

Heartbreaker achieved the overall best performance, combination with a second round of nuclei segmentation did not improve results (**Extended Data Fig. 2b**). Cellulose segmentation of the cell boundary stain closely adhered to human annotation, while Baysor, BIDCell and Proseg tended to oversegment large heart muscle cells.The default MERSCOPE segmentation mostly segmented nuclei and split cardiomyocytes into a large amount of small round cells. Nuclei segmentation with Cellpose reliably segmented nuclei (**Extended Data Fig. 2c**), but did not assign a large number of transcripts (**Extended Data Fig. 2b**). Based on these results all MERFISH datasets were segmented with heartbreaker.

For Xenium, datasets were run with the "Multimodal Cell Segmentation" kit, based on ATP1A1, E-Cadherin and CD45 antibodies for membrane staining and recommendation in CG000750. Compared to the MERSCOPE, the membrane stain was less well-suited for cardiomyocyte segmentation and suffered from lower contrast between foreground

1586 and background (**Extended Data Fig. 2d**). To minimize segmentation errors we therefore
1587 opted for nuclei segmentation with the Cellpose "Nuclei" model for all Xenium data except
1588 the TMA used in the subcellular analysis experiment (**Supplementary Table 3**).

1589

Subcellular analysis (**Fig. 2f-k**) required us to precisely segment cells and their nuclei in
our Xenium data. Here, we employed a new approach. A single TMA with 26 samples
was treated with a post-run WGA and DAPI stain, as described above. This resulted in
sharper boundaries with less background compared to the 10x segmentation kit
(**Extended Data Fig. 2e**). The post-run stain was aligned with the images acquired during
the Xenium measurements by affine transformation. Matching landmarks from both DAPI
channels were picked in the Xenium explorer software. The WGA staining was then run
through a white top hat filter[115] and segmented by morphological watershed in Napari[116].

Spatial transcriptomic processing

All datasets were processed in a standardized fashion with Sopa[8] which performed
segmentation, assignment of transcripts, calculations of protein intensity and conversion
to the Spatialdata[9] format. Cells that contained less than 20 transcripts (50 for human
Xenium samples) or expressed less than 3 unique genes were not considered in further
analysis tasks. Samples where a majority of cells were filtered this way were not
considered for further analysis. The data was then processed in a similar fashion as
described above for the snRNA data. For the Xenium data 2000 variable genes were
calculated, while this step was skipped for the MERFISH data due to the limited panel
size.

1608

Cell type annotation was performed by label transfer from the snRNA dataset of the
corresponding species. Detailed annotation on the cell state level was used for this
transfer (**Supplementary Table 2**). Annotation was performed with Insitutype, TACCO[40],
SingleR, Phi-Space[42] and RCTD[39]. For every cell majority voting was performed and the
number of tools in agreement was calculated. These values were used to calculate the
average number of method agreements per cell state. For cell states where this number
was below three, the cell state's annotation was merged with the alternative states
proposed by the dissenting tools. This resulted in datasets with annotation that was as
detailed as possible, while still being reproducible by the different annotation tools. As an
additional quality control step we calculated Spearman's correlation and Jensen–
Shannon divergence between spatial transcriptomic samples and snRNA samples from
the same patients. This showed strong correspondence between cell type proportions
(Spearman's ρ > 0.8). The cell type annotation per dataset is further described in
Supplementary Table 3.

HeartNet patch based histology analysis

We jointly analyzed the histology of our human and mouse samples. This included WSIs from 53 human and 15 mouse Xenium samples, as well as 23 MERFISH samples. We obtained a patch level embedding (1536 dimensions) by extracting 1,135,625 non-overlapping 224 × 224 pixel (38 μm × 38 μm) patches and running them through the UNI foundation model[28]. Next we assigned expert annotations to the patches. We applied PCA, (50 components) to the embedding followed by Harmony integration[92]. We then performed Leiden clustering[29] with the resolution 0.3 on the integrated embeddings, which revealed nine distinct patch clusters. These clusters were subsequently annotated by expert assessment of representative examples for every cluster.

To enable scalable annotation of new incoming samples, we developed HeartNet, a deep learning classifier trained on 1,135,625 patch embeddings with expert-annotated clusters from our integrated multi-platform dataset. HeartNet is implemented as a fully connected deep neural network operating on patch-level embeddings. The architecture consists of three hidden layers with 256, 128, and 64 neurons, respectively, each followed by batch normalization, ReLU activation, and dropout (rate = 0.3) to improve training stability and reduce overfitting. The network is trained using the Adam optimizer with weight decay (1e-4) and cross-entropy loss. Early stopping based on validation accuracy and adaptive learning-rate scheduling are employed to ensure robust generalization. Using a train/validation/test split (70%/10%/20%) on balanced classes, the model achieved 93.4% accuracy on independent test data, outperforming traditional machine learning approaches (Logistic Regression: 86.9%, Random Forest: 90.4%). We then applied the trained model to annotate 526,242 patches across 9 MERFISH samples using consistent feature extraction pipelines. In total, our comprehensive analysis encompassed 100 samples and 1,661,867 patches across all platforms.

To relate patches with cellular and nuclear information of spatial transcriptomic experiments, we performed a spatial alignment between transcriptomic data and H&E histology. Two different strategies were deployed, based on the difficulty of the task. For the human Xenium samples we used affine transformation as implemented in the Xenium explorer. Mouse Xenium samples showed higher levels of distortion after H&E staining, while MERFISH alignment was difficult because the H&E staining was performed on a subsequent tissue slice. Here we applied a more advanced alignment strategy. We picked matching landmarks in the BigWarp[117] Fiji[118] plugin. We fit thin plate splines[119] to the landmarks and transformed the centers of cells or nuclei from spatial transcriptomic coordinates to the H&E image space. For MERFISH this was only possible for 9 samples. The overall average number of nuclei per patch for mouse, human Xenium data and Merfish was 5.16.

NicheSphere based Differential Colocalization

We used NicheSphere to detect condition specific niches in Xenium experiments (https://github.com/CostaLab/Nichesphere). For this, NicheSphere first calculates pair-wise colocalization probabilities for different cell type pairs by considering the k-nearest neighbors in space (k=5) for each sample. Afterwards, we compared the distribution of colocalization probabilities for each cell type pair in two contrasts (Controls vs. ICM,Controls vs. DCM, Controls vs. AMI) via two-sided Wilcoxon tests[120]. Only significant edges are kept in the colocalization graph and the signed weights are set to the W statistic. Then we computed an adjacency matrix of the graph by measuring an Euclidean distance on the weights and used this as input for Louvain graph clustering[121]. Clusters indicate spatial niches. Finally, we use the niche information to perform a graph embedding for visualization purposes using the community layout (https://github.com/alexodavies/CommunityLayout).

Patient-level analysis

Next, we used NicheSphere niches for sample level trajectory analysis using PILOT[43]. Due to the use of the disease contrasts to define the niches, PILOT provides a semi-supervised trajectory sorting samples according to how well they have niches across the two contrasts, i.e. healthier samples have a lower pseudo-time and more diseased samples have higher pseudo-time and milder cases are in between. PILOT uses optimal transport to calculate distances between pairs of samples based on cell type proportions per sample. Then, a sample trajectory is inferred based on those distances. We modified PILOT, which originally works on transporting cell clusters distributions, using cells similarly as transport cost. Here, PILOT uses the niche sizes as distributions and the cosine distance (PILOT's default option) on the proportion matrix is used as cost function.

Next, we used linear and non-linear regression models in PILOT to characterize niche proportion changes and gene expression changes associated with the disease progression. In other words, for each feature in gene space and niche abundances, we fitted regression models of increasing complexity, including linear, quadratic, and combined linear–quadratic forms. Model significance was evaluated using the F-statistic, with corresponding p-values calculated to test the null hypothesis that all regression coefficients are simultaneously zero. Features whose best-fitting model produced a p-value exceeding 0.05 were deemed statistically non-significant and subsequently excluded from further analysis. Gene selection was performed in a cell type specific way. Moreover, for gene features, we use a Wald Test to contrast the cell type specific patterns with the pattern from other cells. Due to the panel characteristic of the Xenium data, we used a lenient threshold (Wald statistics > 1.8) and a fold change larger than 0.5.

Subcellular pattern identification

After nuclei and cells were segmented we applied a strict preprocessing to the Xenium TMA measured for subcellular analysis. All nuclei with less than 10 transcript counts, an isoperimetric quotient (measure of circularity) of less than 4 and an average DAPI intensity of less than 200 were removed. After label transfer from snRNA, as described above, we additionally removed all cells and nuclei which were not labeled as cardiomyocytes. All cells which were smaller than twice the 95th percentile of nuclei area were removed. In addition all cells with an area larger than 300,000 square pixels were removed. We then removed all nuclei not intersecting cells and all cells intersecting more than 2 nuclei and cells with less than 50 transcript counts.

We then performed the subcellular analysis with the Bento[46] package. A raster grid with a square length of 10 pixels was laid over the segmented cells. We then computed a square-wise vector defined as the difference between the local neighborhood gene composition and the total cell composition. These vectors were normalized and projected into a lower-dimensional space using truncated singular value decomposition as a measure of intracellular spatial variance. In the next step we clustered this embedding using Self-Organizing Maps. The optimal number of clusters was determined dynamically by evaluating the quantization error across a range of 2 to 14 clusters and selecting the inflection point (elbow method). Only squares with at least 100 transcripts were considered for the clustering. This resulted in 4 final domains being assigned to the data.

To interpret these clusters we relied on public APEXseq data[47] which assigned RNA molecules likelihoods for occurring in different cellular compartments. 500,000 squares were randomly selected. We then performed enrichment analysis using the weighted aggregate method implemented in decoupler[99]. We labeled the domains as nuclear, perinuclear, cytoplasm and intercalated disc based on the enrichment scores and overlay with nuclear (DAPI) and cell membrane (WGA) stainings.

Nuclear morphology analysis

We analyzed nuclear morphology based on DAPI staining on the strictly preprocessed TMA described above. 167 classical morphological features related to DAPI intensity and cell shape were calculated. Calculations were performed with squidpy[122] and cp_measure[123] which implement features offered by Cellprofiler[124] and scikit image[119]. The raw feature by nucleus matrix was then post-processed. We used Pycytominer[125] to perform robust median absolute deviation transformation and selected features based on variance and correlation thresholds (default values). This resulted in 154 features being retained. We then formed a pseudo-bulk dataset for every cardiomyocyte state based on the mean of every feature for that cell state and calculated PCA[89].

1739

Morphological features were compared between DES$^+$ CM and CFLAR$^+$ CM populations.
Numeric features were min-max scaled to the unit interval [0,1]. Differential analysis was
performed using Ordinary Least Squares regression with sample included as a fixed
effect. To ensure valid inference in the presence of intra-sample correlation,
heteroscedasticity-consistent covariance matrices were estimated using the sandwich
estimator clustered by sample (R packages sandwich[126] and lmtest[127]). The derived t-
statistics were used to compute p-values, which were subsequently adjusted using the
Benjamini-Hochberg procedure (FDR). Effect sizes were derived from the regression
coefficients and pooled standard deviations to generate Cohen's *d*. Shown are only
selected features based on variance and Cohen's d, all features can be found in
(Supplementary Table 3).

1751

In addition we trained classifiers to predict DES$^+$ CM /CFLAR$^+$ CM status from the
morphology of a nucleus. We compared classification performance based on the
morphological features described above to the embedding generated by the self-
supervised transformer Dinov3[128]. Binary classification of nuclei morphology was
performed using a Logistic Regression classifier with L1 regularization (Lasso). The
regularization parameter (*C*) was set to 0.1 to encourage model sparsity and
generalization. To address potential data imbalance, class weights were automatically
adjusted inversely proportional to class frequencies. Model performance was assessed
using stratified 5-fold cross-validation, and results were reported using mean, precision,
recall, and F1-scores. Classification of nuclei based on the Dinov3 embedding was more
reliable (higher recall, precision and F1 score).

Cell-cell communication

To investigate changes in cell-cell communication across conditions, we integrated
NicheSphere-defined niches with cell-cell communication inference tools. In short, we
used scSeqComm to estimate both ligand-receptor (LR) interactions and intracellular
activity downstream of these LR interactions. Finally, LR interactions between two
conditions were given as input for CrossTalker for a differential cell-cell communication
analysis.

1770

In detail, we first extracted gene expression matrices from the snRNA-seq data for each
experimental group (controls, ICM, ICM_AMI and DCM). We then applied scSeqComm
tool[69] using default parameters to infer LR interaction strengths independently for each
experimental group, applying the ConnectomeDB2020[129] human and CellTalkDB[130]
mouse LR database. Subsequently, we applied CrossTalkeR[70] to identify changes in cell–
cell communication by comparing the predicted LR interactions between each disease
condition and the control (i.e. ICM vs control, ICM_AMI vs control, DCM vs control). The

interactions considered by CrossTalkeR were filtered from the scSeqComm output using
a threshold of LR strength S_inter > 0.5 and restricted to pairs of cell types identified as
significantly colocalized based on NicheSphere analysis for the corresponding spatial
contrast.
1782
To estimate differences in intracellular signaling activity between each experimental
group and control samples for each signaling pathway, we applied the scSeqComm
differential intracellular scoring scheme[131], leveraging the KEGG pathway resources[132]
with pathway integrations from STRINGDB[133] (for WNT signaling) and TTRUSTv2[134],
RegNetwork[135], and HTRIdb[136] databases for transcriptional regulatory networks. The
method quantifies differential intracellular activity based on evidence of altered
transcriptional regulation of target genes, assessed using a two-sided Wilcoxon rank-sum
test, within the same cell-type in disease versus control conditions downstream of
receptor-mediated interactions.

Multi-ome Integration, Trajectory Analysis and Gene regulatory network
We employed an algorithm based on Fused Unbalanced Gromov-Wasserstein
(FUGW)[137] to integrate cardiomyocytes from the human single nucleus RNA-seq and
cardiomyocytes from a previous single nucleus ATAC-seq data[11], which builds upon
SCOT framework[138]. First,we applied dimensionality reduction for both modalities. For
RNA, Principal Component Analysis (PCA) was performed, retaining Principal
Components (PCs) with an explained variance ratio exceeding 10^{-3}. For ATAC, we
utilized scOpen[139] with the dimension set to 100. The intra-modal cost matrices were
calculated as Euclidean distance matrices derived from the RNA PCA embedding and
the ATAC scOpen embedding, respectively. To construct the inter-modal cost matrix, we
first identified common genes shared between the RNA gene expression matrix and the
ATAC gene activity matrix. After normalization, a PCA model was fitted on the RNA data
restricted to these common genes. The ATAC data (common genes) was then projected
into this shared space using the trained RNA PCA model to ensure alignment. We used
the pairwise Euclidean distances between the two modalities in this shared PCA space
as cost function. We assumed uniform marginal distributions for both modalities. The
FUGW hyperparameters were set as follows: the weight balancing Wasserstein and
Gromov-Wasserstein distances *alpha* was set to 0.1, the unbalanced constraint
parameter *rho* to 1.1, and the entropic regularization parameter *epsilon* to 10^{-5}. Finally,
the resulting optimal transport plan was used to project the ATAC embeddings from the
scOpen space into the RNA PCA space. Once aligned, each RNA cell was paired with its
nearest ATAC neighbor based on Euclidean distance.
1814
Subsequently, we employed the Phlower[63] framework to infer differentiation trajectories
of single-cell RNA-seq data. Initially, a k-nearest neighbor (k-NN) graph was constructed

in the PCA space with K=15 to capture the local manifold geometry. To capture the global
reachability and developmental timescales, we implemented a diffusion process on the
graph followed by spectral decomposition, yielding 14 diffusion components. Taking the
designated "DES+ CM" type cell group as the root, we applied a Hodge Laplacian (HL)
decomposition to compute a continuous pseudotime ordering; Subsequently, 10^4
trajectories were sampled by stochastic traversals of the kNN graph initiated from the root
cells. Finally, cells were ordered along the inferred pseudotime and trajectory paths to
reconstruct the dynamic expression kinetics of genes and transcription factors (TFs)
throughout the differentiation process.
1826
Based on the differentiation trajectories inferred by PHLOWER, we performed
transcription factor enrichment analysis for each identified branch. Specifically, cells were
first partitioned into branch-specific populations versus the remaining cell pool according
to the trajectory topology. Differential expression analysis was then conducted to identify
candidate regulatory factors. A TF was considered a potential regulator if it exhibited a
significant upregulation in the target branch compared to other branches, defined by a
log2 fold-change (log2FC) threshold of 0.13. The statistical significance (t-test) was
further evaluated by assessing the expression consistency among intra-branch cells,
yielding corresponding p-values. For all branches, TFs with a p-value < 0.05 were
aggregated as the final set of candidate regulators for downstream analysis. We used the
above method to select branch specific TFs (**Supplementary Table 4**).
1838
To construct the enhancer based gene Regulatory Network (GRN), we utilized scMEGA[61]
to calculate gene-TF correlations. In the resulting network, TF nodes were annotated with
pseudotime values derived from PHLOWER. We filtered for significant TFs by intersecting
those selected by PHLOWER (p-value < 0.05) and only considering genes with a
correlation > 0.3 when contrasting the gene expression and TF activity scores. Finally,
edges were filtered to retain only those with a gene-TF correlation > 0.5. To mitigate
network density for visualization, we further restricted the network to the top 500 edges
per TF based on correlation strength. The final GRN was produced with a Fruchterman-
Reingold force-directed network algorithm.

Pseudo-time analysis of embryonal cardiomyocytes

For the differentiation of embryonal cardiomyocytes, the dataset was subset to
"Differentiating LV-CMs", "Immature LV-CMs" and "Mature LV-CMs". A trajectory was
then calculated with scMEGA[61] using a diffusion map[140] based on harmony-corrected
PCA. The trajectory was defined in the order from "Immature LV-CMs" - "Differentiating
LV-CMs" - "Mature LV-CMs" with archR[141] and the first three dimensions of the diffusion
map were input into the algorithm. ArchR constructs a cellular trajectory using the
centroids of user-defined clusters in reduced dimension, aligning individual cells via the

nearest cell-to-trajectory distance. We then analyzed gene expression along the
trajectory. Pearson correlation was calculated between normalized log gene expression
and the pseudo-time assigned to a cell and genes were ranked based on this value.
Enrichment analysis was then performed with Camera[98] in pre-ranked mode as
implemented in edgeR[97]. The queried gene sets were gathered from the msigdb[100] and
encompassed the hallmark[100] collection, KEGG[132], Reactome[142], biocarta and GO[143].
Additionally, we added three custom gene sets composed of 100 marker genes each for
DES,CFLAR and NPPA cardiomyocytes to the test. The marker genes were calculated
using a two-sided tie-corrected Wilcoxon rank-sum test.

Spatial statistics

The spatial segregation index was calculated on the MERFISH v2 data (n=13) and was
calculated with the dixon function[144] as implemented in the dixon library[145]. Spatial
segregation tests every cell against its nearest neighbor and assesses whether a cell of
the same type is found more often than would be expected by chance. We specifically
analyzed "self-enrichment" (homotypic neighbor relationships) to determine the degree to
which niches form contiguous clusters versus scattered distributions. A comparison
between the Z-values for the cell types of interest output by the segregation test was then
calculated with the two-sided Wilcoxon rank-sum test. For the comparison of niches
between ICM and DCM the mean Z score for niches in every sample was compared using
a one-sided Wilcoxon rank-sum test (n=57).

Analysis of public fetal Visium data

The dataset was retrieved from the original publication[74]. We performed deconvolution
with our embryo data as reference with Cell2location[146]. We removed mitochondrial
genes and selected genes of interest as suggested by the Cell2location authors. The
deconvolution was performed with cells per location set to 15, detection alpha set to 20
and otherwise default parameters. We then calculated the mean number of cell types per
Visium spot for every heart compartment annotated by the original authors.

Analysis of rat in vivo measurements

Sirius red staining was quantified with Halo AI 3.6. Statistics on heart function were
calculated in GraphPad Prism. A one-way ANOVA followed by Dunnett's multiple
comparisons test was used for all measurements except bodyweight, where a two-way
ANOVA followed by Dunnett's multiple comparisons test was used.

Statistics and reproducibility

Sample sizes were determined based on sample availability and no statistical method
was used to predetermine sample size. Data generation was randomized with respect to

disease and control samples. The investigators were not blinded to allocation during experiments and outcome assessment. Criteria for data exclusion, including filtering of nuclei or cells based on size and gene counts, are detailed in the methods section. Microscopic images shown are representative examples and not exhaustive. Statistical tests and exact sample sizes are provided in the figure legends. Correction for multiple hypothesis testing was performed using the Benjamini–Hochberg procedure, unless specified otherwise. For heatmaps, the use of z-scoring or min-max scaling (0–1) is indicated in the figure legends. For boxplots, the median is indicated by the center line, with whiskers extending from the first and third quartiles to the farthest data point within 1.5× of the inter-quartile range. Violin plots represent a kernel density estimate of the underlying distribution; densities were normalized so that all violins within a panel have the same width.

Reporting summary

Further information on study design is available in the Nature Portfolio Reporting Summary attached to this article.

Data availability

Processed snRNA data and spatial transcriptomics are available from https://rwth-aachen.sciebo.de/s/c6oay6kCrAdXgM7. Data will be deposited in public repositories upon publication. Human Sequencing data will be deposited to the European Genome-phenome Archive (EGA), rat and mouse sequencing data will be deposited to the European Nucleotide Archive (ENA). Processed anndata files will be shared on Cell x Gene. Spatial transcriptomic datasets will be shared through the BioImage Archive. The data will additionally be presented in an interactive viewer at heartatlas.kuppelab.org.

Code availability

Analysis code is available from https://rwth-aachen.sciebo.de/s/LSCsL5j7cnrHFrY. and will be made available on Github and Zenodo upon publication. Code and tutorials for NicheSphere are available at https://github.com/Costalab/NicheSphere. A model checkpoint for HeartNet is available at https://github.com/Costalab/HeartNet.

Acknowledgements

This project has been funded by the German Research Foundation received by C.K. (CRU344-417911533, CRU5011-445703531, Emmy Noether EN-459969915); received by I.C. (DFG; 3888802535, CRU5011-445703531) by the the German Ministry of Science, Technology and Space (BMFTR) consortia Graphs4Patients for C.K., M. S. and I.C and E:MED Fibromap and CureFib for I.C. and R.K. This work was further supported

by grants to CK from the European Research Council (ERC-StG-101040726), the
Boehringer Ingelheim Foundation, Else Kroener Fresenius Foundation (EKFS, Physician
Scientist Professorship CK) and the Aventis Foundation. R.K. received support from the
German Research Foundation (DFG: SFBTRR219 322900939, CRU344
4288578857858, CRU5011 445703531, KR4073/18-1 (561848128) & INST 222/1582-1),
the European Research Council (ERC-CoG 101043403 & 101001791, ERC-PoC
101212974 and 10104303), the Leducq Foundation (ImmunoFibHF 20CVD02 and
COMET 24CVD02). G.C. was partially supported by Fondazione Ing. Aldo Gini. Views
and opinions expressed are those of the authors only and do not necessarily reflect those
of the European Union or the European Research Council Executive Agency. Neither the
European Union nor the granting authority can be held responsible for them. We thank
Dr. Igor Efimov from the Northwestern University, Chicago, IL, USA, for providing fixed
human donor heart samples. NM is supported by the Deutsche Forschungsgemeinschaft
(German Research Foundation; TRR 219; Project-ID 322900939. We thank the IZKF
Aachen Genomics Core facility for sequencing experiments. The Dutch Fetal Biobank
received funding from Amsterdam UMC (Innovation Impuls) and the Amsterdam
Reproduction and Development research institute.
1942
1943   We thank all supporting centers, physicians, midwives and nurses for the information they
1944   have provided to parents who considered donating their fetuses to science and in
1945   particular to the DFB. Most importantly, we express our gratitude to all mothers and
1946   couples who have donated their most precious gift to the scientific community.
1947
1948   This study is part of the UCC-UNRAVEL biobank (www.unravelrdp.nl). The study was
1949   approved by the local institutional ethics review board (University Medical Center Utrecht,
1950   protocol UCC-UNRAVEL #12-387 and TCBio 25-242) ERN UNRAVEL is Member of the
1951   European Reference Network for rare, low prevalence and complex diseases of the heart:
1952   ERN GUARD-Heart (ERN GUARD HEART; http://guardheart.ern-net.eu)
1953
1954   The data used in this publication was managed using the research data management
1955   platform Coscine with storage space granted by the Research Data Storage (RDS) of the
1956   DFG and Ministry of Culture and Science of the State of North Rhine-Westphalia (DFG:
1957   INST222/1261-1 and MKW: 214-4.06.05.08 - 139057).
1958
1959   The authors gratefully acknowledge the computing time provided to them at the NHR
1960   Center NHR4CES at RWTH Aachen University (project number p0020567). This is
1961   funded by the Federal Ministry of Education and Research, and the state governments
1962   participating on the basis of the resolutions of the GWK for national high performance
1963   computing at universities (www.nhr-verein.de/unsere-partner).
1964

1965 The authors wish to thank the many other people who assisted in the completion of the
1966 manuscript. Among them Louis Kümmerle for helping with spatial transcriptomic panel
1967 design, Clarence K. Mah for his advice on BENTO and Beth Cimini for input on cell
1968 segmentation and image alignment. In addition we thank Vizgen, Inc. for giving us early
1969 access to MERFISH v2 chemistry and technical support by Marc Reudelsterz and 10x
1970 Genomics, Inc. for giving us access to FLEX rat probes.

Author Contributions

C.K. and I.G.C. conceived and designed the study and supervised the project. CK
acquired the majority of funding for the study. D.P., E.S., O.G., S.F., S.S., X.L., M.v.S.,
M.E., T.S., B.K., J.A., K.L., B.S.d.B., J.H.-v.d.V., S.D., and H.M. organized patient tissue
collection, biobanking, and patient consent. C.K. generated tissue microarrays. E.S. and
O.G. performed spatial transcriptomics and single-nucleus RNA-sequencing
experiments. T.L., D.R.M., and S.D. provided mouse myocardial infarction time-course
tissues. M.Z., H.Z., F.R., and A.Z. generated and provided rat tissues and snRNA-
sequencing datasets. Human iPSC-derived experiments, including differentiation,
functional assays, and snRNA-seq, were performed by L.M., Z.F., and D.W., with
guidance from A.A. and C.K. P.K. designed the MERFISH and Xenium panels, optimized
the image segmentation approach, and performed subcellular spatial transcriptomic
analyses. K.K. performed histological analyses; H&E-stained cardiac tissues were
evaluated by K.K. and M.E. Computational analyses were structured as follows: M.R.
performed niche analyses; G.C. conducted cell–cell communication analyses; M.J.
performed PILOT and histology patch analysis. These analyses were advised by I.G.C.
and C.K. All other computational analyses—including data preprocessing, quality control,
cross-species integration, clustering, differential expression analysis, trajectory inference,
multi-omic integration, segmentation optimization, and subcellular transcriptomic
analyses—were conducted by P.K., M.J., D.P., G.C., M.R., D.K., K.P., and M.C., with
guidance from I.G.C., M.S. and C.K. Data interpretation was performed by P.K., I.G.C.,
and C.K., with input from all co-authors. P.K., I.G.C., and C.K. wrote the manuscript and
organized the figures. N.M., K.K., R.K., I.G.C., H.M., M.S., and C.K. critically revised the
manuscript and advised on data interpretation. All authors read and approved the final
manuscript.

Competing Interest

1997 C.K. received honoria from within the last 2 years from BAYER and received research
1998 funding from InsilicoMedicine for this study to perform rat snRNA-sequencing
1999 experiments. R.K. acknowledges the following outside of the submitted work, he is
2000 founder and board member of Sequantrix GmbH, received honoraria from Bayer, Chugai,
2001 Pfizer, Roche, Genentech, Eli Lilly, and GSK, AMGEN, Sobi, Hybridize Therapeutic,

Valerio Therapeutics, Exigent Therapeutics for advisory board meetings and received research funding from Travere Therapeutics, Galapagos, Chugai, Novo Nordisk, and Ask Bio.

F.R., H.Z., M.Z., and A.Z. are employees of Insilico Medicine, which develops generative artificial intelligence and related technologies for drug discovery, drug development, and aging research. Insilico Medicine has ongoing therapeutic programs, including rentosertib (INS018_055) in fibrotic and other disease areas.

The remaining authors declare no competing interests.

References

1. Koopmans, T. & van Rooij, E. Molecular gatekeepers of endogenous adult mammalian cardiomyocyte proliferation. *Nat. Rev. Cardiol.* (2025) doi:10.1038/s41569-025-01145-y.

2. González-Rosa, J. M. *et al.* Myocardial polyploidization creates a barrier to heart regeneration in zebrafish. *Dev. Cell* **44**, 433–446.e7 (2018).

3. Global Burden of Cardiovascular Diseases and Risks 2023 Collaborators. Global, regional, and national burden of cardiovascular diseases and risk factors in 204 countries and territories, 1990-2023. *J. Am. Coll. Cardiol.* (2025) doi:10.1016/j.jacc.2025.08.015.

4. Bozkurt, B. Contemporary pharmacological treatment and management of heart failure. *Nat. Rev. Cardiol.* **21**, 545–555 (2024).

5. Derks, W. *et al.* A latent cardiomyocyte regeneration potential in human heart disease. *Circulation* **151**, 245–256 (2025).

6. Poss, K. D., Wilson, L. G. & Keating, M. T. Heart regeneration in zebrafish. *Science* **298**, 2188–2190 (2002).

7. Wang, Z. *et al.* Mechanistic basis of neonatal heart regeneration revealed by transcriptome and histone modification profiling. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 18455–18465 (2019).

8. Swift, S. K. *et al.* Cardiomyocyte ploidy is dynamic during postnatal development and varies across genetic backgrounds. *Development* **150**, (2023).

9. Paredes, A. *et al.* γ-Linolenic acid in maternal milk drives cardiac metabolic maturation. *Nature* **618**, 365–373 (2023).

10. Kanemaru, K. *et al.* Spatially resolved multiomics of human cardiac niches. *Nature* **619**, 801–810 (2023).

11. Kuppe, C. *et al.* Spatial multi-omic map of human myocardial infarction. *Nature* **608**, 766–777 (2022).

12. Man, J. C. K. *et al.* Genetic dissection of a super enhancer controlling the Nppa-Nppb cluster in the heart. *Circ. Res.* **128**, 115–129 (2021).

13. Sergeeva, I. A. *et al.* Identification of a regulatory domain controlling the Nppa-Nppb gene cluster during heart development and stress. *Development* **143**, 2135–2146 (2016).

14. Li, X. *et al.* Inhibition of fatty acid oxidation enables heart regeneration in adult mice. *Nature* **622**, 619–626 (2023).

15. Derks, W. & Bergmann, O. Polyploidy in cardiomyocytes. *Circ. Res.* **126**, 552–565 (2020).

16. Schoenfelder, K. P. & Fox, D. T. The expanding implications of polyploidy. *J. Cell Biol.* **209**, 485–491 (2015).

17. Gjelsvik, K. J., Besen-McNally, R. & Losick, V. P. Solving the polyploid mystery in
health and disease. *Trends Genet.* **35**, 6–14 (2019).
18. Pandit, S. K., Westendorp, B. & de Bruin, A. Physiological significance of
polyploidization in mammalian cells. *Trends Cell Biol.* **23**, 556–566 (2013).
19. Gan, P., Patterson, M. & Sucov, H. M. Cardiomyocyte polyploidy and implications for
heart regeneration. *Annu. Rev. Physiol.* **82**, 45–61 (2020).
20. Peterson, N. G. & Fox, D. T. Communal living: the role of polyploidy and syncytia in
tissue biology. *Chromosome Res.* **29**, 245–260 (2021).
21. Sladky, V. C., Eichin, F., Reiberger, T. & Villunger, A. Polyploidy control in hepatic
health and disease. *J. Hepatol.* **75**, 1177–1191 (2021).
22. Matondo, R. B. *et al.* Atypical E2f functions are critical for pancreas polyploidization.
PLoS One **13**, e0190899 (2018).
23. Bischof, C. *et al.* Mitochondrial-cell cycle cross-talk drives endoreplication in heart
disease. *Sci. Transl. Med.* **13**, eabi7964 (2021).
24. Matsumoto, T., Wakefield, L., Tarlow, B. D. & Grompe, M. In vivo lineage tracing of
polyploid hepatocytes reveals extensive proliferation during liver regeneration. *Cell Stem*
Cell **26**, 34–47.e3 (2020).
25. Zhang, S. *et al.* The polyploid state plays a tumor suppressive role in the liver. *bioRxiv*
(2017) doi:10.1101/149799.
26. Hayashi, K. *et al.* Polyploidy mitigates the impact of DNA damage while simultaneously
bearing its burden. *Cell Death Discov.* **10**, 436 (2024).
27. Yekelchyk, M., Guenther, S., Preussner, J. & Braun, T. Mono- and multi-nucleated
ventricular cardiomyocytes constitute a transcriptionally homogenous cell population. *Basic*
Res. Cardiol. **114**, 36 (2019).
28. Chen, R. J. *et al.* Towards a general-purpose foundation model for computational
pathology. *Nat. Med.* **30**, 850–862 (2024).
29. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-
connected communities. *Sci. Rep.* **9**, 5233 (2019).
30. Elz, A. E. *et al.* Evaluating the practical aspects and performance of commercial single-
cell RNA sequencing technologies. *bioRxivorg* (2025) doi:10.1101/2025.05.19.654974.
31. De Simone, M. *et al.* A comprehensive analysis framework for evaluating commercial
single-cell RNA sequencing technologies. *Nucleic Acids Res.* **53**, (2025).
32. Broadwell, L. J. *et al.* Myosin 7b is a regulatory long noncoding RNA (lncMYH7b) in the
human heart. *J. Biol. Chem.* **296**, 100694 (2021).
33. Lee, L. A., Broadwell, L. J., Buvoli, M. & Leinwand, L. A. Nonproductive splicing
prevents expression of MYH7b protein in the mammalian heart. *J. Am. Heart Assoc.* **10**,
e020965 (2021).
34. Giampietri, C. *et al.* c-Flip overexpression reduces cardiac hypertrophy in response to
pressure overload. *J. Hypertens.* **26**, 1008–1016 (2008).
35. Reichart, D. *et al.* Pathogenic variants damage cell composition and single cell
transcription in cardiomyopathies. *Science* **377**, eabo1984 (2022).
36. Stringer, C. & Pachitariu, M. Cellpose3: one-click image restoration for improved
cellular segmentation. *Nat. Methods* **22**, 592–599 (2025).
37. Greiner, J. *et al.* A deep learning-enabled toolkit for the 3D segmentation of ventricular
cardiomyocytes. *J. Physiol.* (2025) doi:10.1113/JP288557.
38. Danaher, P. *et al.* Insitutype: likelihood-based cell typing for single cell spatial
transcriptomics. *Bioinformatics* (2022).
39. Cable, D. M. *et al.* Robust decomposition of cell type mixtures in spatial transcriptomics.
Nat. Biotechnol. **40**, 517–526 (2022).
40. Mages, S. *et al.* TACCO unifies annotation transfer and decomposition of cell identities
for single-cell and spatial omics. *Nat. Biotechnol.* **41**, 1465–1473 (2023).

41. Aran, D. *et al.* Reference-based analysis of lung single-cell sequencing reveals a
transitional profibrotic macrophage. *Nat. Immunol.* **20**, 163–172 (2019).
42. Mao, J., Deng, Y. & Lê Cao, K.-A. Φ-Space: Continuous phenotyping of single-cell
multi-omics data. *Bioinformatics* (2024).
43. Joodaki, M. *et al.* Detection of PatIent-Level distances from single cell genomics and
pathomics data with Optimal Transport (PILOT). *Mol. Syst. Biol.* **20**, 57–74 (2024).
44. Yoshioka, N. *et al.* Isoform-specific mutation in Dystonin-b gene causes late-onset
protein aggregate myopathy and cardiomyopathy. *Elife* **11**, (2022).
45. Granzier, H. L. & Labeit, S. Discovery of titin and its role in heart function and disease.
Circ. Res. **136**, 135–157 (2025).
46. Mah, C. K. *et al.* Bento: A toolkit for subcellular analysis of spatial transcriptomics data.
bioRxiv 2022.06.10.495510 (2023) doi:10.1101/2022.06.10.495510.
47. Fazal, F. M. *et al.* Atlas of Subcellular RNA Localization Revealed by APEX-Seq. *Cell*
178, 473–490.e26 (2019).
48. Scarborough, E., Uchida, K., Kehat, I. & Prosser, B. RNAs are localized and translated
at discrete cardiomyocyte microdomains. *J. Mol. Cell. Cardiol.* **173**, S163 (2022).
49. Lu, F. *et al.* CMYA5 establishes cardiac dyad architecture and positioning. *Nat.*
Commun. **13**, 2185 (2022).
50. Baez, M. V. *et al.* Smaug1 mRNA-silencing foci respond to NMDA and modulate
synapse formation. *J. Cell Biol.* **195**, 1141–1157 (2011).
51. Hinnebusch, A. G. eIF3: a versatile scaffold for translation initiation complexes. *Trends*
Biochem. Sci. **31**, 553–562 (2006).
52. West, G., Sedighi, S., Agnetti, G. & Taimen, P. Intermediate filaments in the heart: The
dynamic duo of desmin and lamins orchestrates mechanical force transmission. *Curr. Opin.*
Cell Biol. **85**, 102280 (2023).
53. Tsikitis, M. *et al.* Molecular insights into the role of desmin intermediate filament
network in chromatin landscape, cardiomyocyte differentiation, and maturation. *Cell Death*
Dis. **16**, 723 (2025).
54. George, R. M. *et al.* Single cell evaluation of endocardial Hand2 gene regulatory
networks reveals HAND2-dependent pathways that impact cardiac morphogenesis.
Development **150**, (2023).
55. Yang, C.-Y. *et al.* Reverse hierarchical DED assembly in the cFLIP-procaspase-8 and
cFLIP-procaspase-8-FADD complexes. *Nat. Commun.* **15**, 8974 (2024).
56. Liu, D. *et al.* Cellular FADD-like IL-1β-converting enzyme-inhibitory protein attenuates
myocardial ischemia/reperfusion injury via suppressing apoptosis and autophagy
simultaneously. *Nutr. Metab. Cardiovasc. Dis.* **31**, 1916–1928 (2021).
57. Russell, N. J. *et al.* Spatial ploidy inference using quantitative imaging. *Cell Rep.*
Methods **5**, 101249 (2025).
58. Stirling, D. R. *et al.* CellProfiler 4: improvements in speed, utility and usability. *BMC*
Bioinformatics **22**, 433 (2021).
59. Wehrens, M. *et al.* Single-cell transcriptomics provides insights into hypertrophic
cardiomyopathy. *Cell Rep.* **39**, 110809 (2022).
60. González-Velasco, O. *et al.* Identifying similar populations across independent single
cell studies without data integration. *NAR Genom. Bioinform.* **7**, lqaf042 (2025).
61. Li, Z., Nagai, J. S., Kuppe, C., Kramann, R. & Costa, I. G. scMEGA: single-cell multi-
omic enhancer-based gene regulatory network inference. *Bioinform Adv* **3**, vbad003 (2023).
62. Gomez-Velazquez, M. *et al.* CTCF counter-regulates cardiomyocyte development and
maturation programs in the embryonic heart. *PLoS Genet.* **13**, e1006985 (2017).
63. Cheng, M. *et al.* PHLOWER leverages single-cell multimodal data to infer complex,
multi-branching cell differentiation trajectories. *Nat. Methods* **22**, 2328–2336 (2025).
64. Velecela, V. *et al.* Epicardial cell shape and maturation are regulated by Wt1 via

transcriptional control of Bmp4. *Development* **146**, dev178723 (2019).

65. Ettou, S. *et al.* Epigenetic transcriptional reprogramming by WT1 mediates a repair
response during podocyte injury. *Sci. Adv.* **6**, eabb5460 (2020).

66. Marques, I. J. *et al.* Wt1 transcription factor impairs cardiomyocyte specification and
drives a phenotypic switch from myocardium to epicardium. *Development* **149**, (2022).

67. Renault, M.-A. *et al.* Gli3 regulation of myogenesis is necessary for ischemia-induced
angiogenesis. *Circ. Res.* **113**, 1148–1158 (2013).

68. Sakamoto, T. *et al.* A critical role for estrogen-related receptor signaling in cardiac
maturation. *Circ. Res.* **126**, 1685–1702 (2020).

69. Baruzzo, G., Cesaro, G. & Di Camillo, B. Identify, quantify and characterize cellular
communication from single-cell RNA sequencing data with scSeqComm. *Bioinformatics* **38**,
1920–1929 (2022).

70. Nagai, J. S., Leimkühler, N. B., Schaub, M. T., Schneider, R. K. & Costa, I. G.
CrossTalkeR: analysis and visualization of ligand-receptorne tworks. *Bioinformatics* **37**,
4263–4265 (2021).

71. Ruiz-Villalba, A. *et al.* Single-cell RNA sequencing analysis reveals a crucial role for
CTHRC1 (collagen triple helix repeat containing 1) cardiac fibroblasts after myocardial
infarction. *Circulation* **142**, 1831–1847 (2020).

72. Razeghi, P. *et al.* Metabolic gene expression in fetal and failing human heart.
Circulation **104**, 2923–2931 (2001).

73. Farah, E. N. *et al.* Spatially organized cellular communities form the developing human
heart. *Nature* **627**, 854–864 (2024).

74. Lázár, E. *et al.* Spatiotemporal gene expression and cellular dynamics of the developing
human heart. *Nat. Genet.* **57**, 2756–2771 (2025).

75. Friedman, C. E. *et al.* Single-cell transcriptomic analysis of cardiac differentiation from
human PSCs reveals HOPX-dependent cardiomyocyte maturation. *Cell Stem Cell* **23**, 586–
598.e8 (2018).

76. Knöll, R. *et al.* Telethonin deficiency is associated with maladaptation to biomechanical
stress in the mammalian heart. *Circ. Res.* **109**, 758–769 (2011).

77. Buettner, M., Ostner, J., Mueller, C. L., Theis, F. J. & Schubert, B. scCODA is a
Bayesian model for compositional single-cell data analysis. *Nat. Commun.* **12**, 1–10 (2021).

78. Ren, F. *et al.* A small-molecule TNIK inhibitor targets fibrosis in preclinical and clinical
models. *Nat. Biotechnol.* (2024) doi:10.1038/s41587-024-02143-0.

79. Masuda, M. *et al.* TNIK inhibition abrogates colorectal cancer stemness. *Nat. Commun.*
7, 12586 (2016).

80. Xu, Z. *et al.* A generative AI-discovered TNIK inhibitor for idiopathic pulmonary fibrosis:
a randomized phase 2a trial. *Nat. Med.* **31**, 2602–2610 (2025).

81. Dawood, Y. *et al.* Human embryonic and fetal biobanking: Establishing the Dutch Fetal
Biobank and a framework for standardization. *Dev. Cell* **58**, 2826–2835 (2023).

82. Sammani, A. *et al.* UNRAVEL: big data analytics research data platform to improve
care of patients with cardiomyopathies using routine electronic health records and
standardised biobanking. *Neth. Heart J.* **27**, 426–434 (2019).

83. Tombor, L. S. *et al.* Single cell sequencing reveals endothelial plasticity with transient
mesenchymal activation after myocardial infarction. *Nat. Commun.* **12**, 681 (2021).

84. Kornienko, J. *et al.* Mislocalization of pathogenic RBM20 variants in dilated
cardiomyopathy is caused by loss-of-interaction with Transportin-3. *Nat. Commun.* **14**, 4312
(2023).

85. Grancharova, T. *et al.* A comprehensive analysis of gene expression changes in a high
replicate and open-source dataset of differentiating hiPSC-derived cardiomyocytes. *Sci.*
Rep. **11**, 15845 (2021).

86. Sugano, T. *et al.* Pharmacological blockage of transforming growth factor-β signalling

by a Traf2- and Nck-interacting kinase inhibitor, NCB-0846. *Br. J. Cancer* **124**, 228–236
(2021).
87. Mölder, F. *et al.* Sustainable data analysis with Snakemake. *F1000Res.* **10**, 33 (2021).
88. Young, M. D. & Behjati, S. SoupX removes ambient RNA contamination from droplet-
based single-cell RNA sequencing data. *Gigascience* **9**, (2020).
89. Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression
data analysis. *Genome Biol.* **19**, 15 (2018).
90. Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational identification of cell
Doublets in Single-cell transcriptomic data. *Cell Syst.* **8**, 281–291.e9 (2019).
91. Gayoso, A. & Shor, J. *JonathanShor/DoubletDetection: Doubletdetection v4.3.0.post1*.
(Zenodo, 2025). doi:10.5281/ZENODO.2678041.
92. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat. Methods* **16**, 1289–1296 (2019).
93. *Presto: Fast Wilcoxon and auROC*. (Github).
94. Domínguez Conde, C. *et al.* Cross-tissue immune cell analysis reveals tissue-specific
features in humans. *Science* **376**, eabl5197 (2022).
95. Germain, P.-L., Sonrel, A. & Robinson, M. D. pipeComp, a general framework for the
evaluation of computational pipelines, reveals performant single cell RNA-seq
preprocessing tools. *Genome Biol.* **21**, 227 (2020).
96. Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by
single-cell RNA-seq. *Science* **352**, 189–196 (2016).
97. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140
(2010).
98. Wu, D. & Smyth, G. K. Camera: a competitive gene set test accounting for inter-gene
correlation. *Nucleic Acids Res.* **40**, e133 (2012).
99. Badia-I-Mompel, P. *et al.* decoupleR: ensemble of computational methods to infer
biological activities from omics data. *Bioinform. Adv.* **2**, vbac016 (2022).
100. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set
collection. *Cell Syst.* **1**, 417–425 (2015).
101. Schubert, M. *et al.* Perturbation-response genes reveal signaling footprints in cancer
gene expression. *Nat. Commun.* **9**, 20 (2018).
102. Büttner, M., Ostner, J., Müller, C. L., Theis, F. J. & Schubert, B. scCODA is a
Bayesian model for compositional single-cell data analysis. *Nat. Commun.* **12**, 6876 (2021).
103. Fury, W., Batliwalla, F., Gregersen, P. K. & Li, W. Overlapping probabilities of top
ranking gene lists, hypergeometric distribution, and stringency of gene selection criterion.
Conf. Proc. IEEE Eng. Med. Biol. Soc. **2006**, 5531–5534 (2006).
104. Tucker, N. R. *et al.* Transcriptional and cellular diversity of the human heart.
Circulation **142**, 466–482 (2020).
105. Koenig, A. L. *et al.* Single-cell transcriptomics reveals cell-type-specific diversification
in human heart failure. *Nat. Cardiovasc. Res.* **1**, 263–280 (2022).
106. Chaffin, M. *et al.* Single-nucleus profiling of human dilated and hypertrophic
cardiomyopathy. *Nature* **608**, 174–180 (2022).
107. Litviňuková, M. *et al.* Cells of the adult human heart. *Nature* **588**, 466–472 (2020).
108. Amrute, J. M. *et al.* Targeting RUNX1 in Macrophages Facilitates Cardiac Recovery.
Immunology (2025).
109. Chen, K. H., Boettiger, A. N., Moffitt, J. R., Wang, S. & Zhuang, X. RNA imaging.
Spatially resolved, highly multiplexed RNA profiling in single cells. *Science* **348**, aaa6090
(2015).
110. Pan, X. *MERFISH_probe_design*. (Github, 2024).
111. Petukhov, V. *et al.* Cell segmentation in imaging-based spatial transcriptomics. *Nat.*

Biotechnol. **40**, 345–354 (2022).

112. Fu, X. *et al.* BIDCell: Biologically-informed self-supervised learning for segmentation of subcellular spatial transcriptomics data. *Nat. Commun.* **15**, 509 (2024).

113. Jones, D. C. *et al.* Cell simulation as cell segmentation. *Nat. Methods* **22**, 1331–1342 (2025).

114. Kumar, N. *et al.* A dataset and a technique for generalized nuclear segmentation for computational pathology. *IEEE Trans. Med. Imaging* **36**, 1550–1560 (2017).

115. Serra, J. (jean). *Image Analysis and Mathematical Morphology: V. 1*. (Academic Press, San Diego, CA, 1982).

116. Sofroniew, N. *et al. Napari: A Multi-Dimensional Image Viewer for Python*. (Zenodo, 2025). doi:10.5281/ZENODO.3555620.

117. Bogovic, J. A., Hanslovsky, P., Wong, A. & Saalfeld, S. Robust registration of calcium images by learned contrast synthesis. in *2016 IEEE 13th International Symposium on Biomedical Imaging (ISBI)* 1123–1126 (IEEE, 2016).

118. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).

119. van der Walt, S. *et al.* scikit-image: image processing in Python. *PeerJ* **2**, e453 (2014).

120. Wilcoxon, F. Individual comparisons of grouped data by ranking methods. *J. Econ. Entomol.* **39**, 269 (1946).

121. Traag, V. A. Faster unfolding of communities: speeding up the Louvain algorithm. *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **92**, 032801 (2015).

122. Palla, G. *et al.* Squidpy: a scalable framework for spatial omics analysis. *Nat. Methods* **19**, 171–178 (2022).

123. Muñoz, A. F. *et al.* cp_measure: API-first feature extraction for image-based profiling workflows. *ArXiv* (2025).

124. Carpenter, A. E. *et al.* CellProfiler: image analysis software for identifying and quantifying cell phenotypes. *Genome Biol.* **7**, R100 (2006).

125. Serrano, E. *et al.* Reproducible image-based profiling with Pycytominer. *Nat. Methods* **22**, 677–680 (2025).

126. Zeileis, A. & Lumley, T. sandwich: Robust Covariance Matrix Estimators. *CRAN: Contributed Packages* The R Foundation https://doi.org/10.32614/cran.package.sandwich (2004).

127. Hothorn, T., Zeileis, A., Farebrother, R. W. & Cummins, C. lmtest: Testing Linear Regression Models. *CRAN: Contributed Packages* The R Foundation https://doi.org/10.32614/cran.package.lmtest (1999).

128. Siméoni, O. *et al.* DINOv3. *arXiv [cs.CV]* (2025).

129. Hou, R., Denisenko, E., Ong, H. T., Ramilowski, J. A. & Forrest, A. R. R. Predicting cell-to-cell communication networks using NATMI. *Nat. Commun.* **11**, 5011 (2020).

130. Shao, X. *et al.* CellTalkDB: a manually curated database of ligand-receptor interactions in humans and mice. *Brief. Bioinform.* **22**, (2021).

131. Cesaro, G., Baruzzo, G., Tussardi, G. & Di Camillo, B. Differential cellular communication inference framework for large-scale single-cell RNA-sequencing data. *NAR Genom. Bioinform.* **7**, lqaf084 (2025).

132. Kanehisa, M. & Goto, S. KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.* **28**, 27–30 (2000).

133. Szklarczyk, D. *et al.* The STRING database in 2025: protein networks with directionality of regulation. *Nucleic Acids Res.* **53**, D730–D737 (2025).

134. Han, H. *et al.* TRRUST v2: an expanded reference database of human and mouse transcriptional regulatory interactions. *Nucleic Acids Res.* **46**, D380–D386 (2018).

135. Liu, Z.-P., Wu, C., Miao, H. & Wu, H. RegNetwork: an integrated database of transcriptional and post-transcriptional regulatory networks in human and mouse. *Database*

(Oxford) **2015**, bav095 (2015).

136. Bovolenta, L. A., Acencio, M. L. & Lemke, N. HTRIdb: an open-access database for
experimentally verified human transcriptional regulation interactions. *BMC Genomics* **13**,
405 (2012).

137. Thual, A. *et al.* Aligning individual brains with Fused Unbalanced Gromov-
Wasserstein. *arXiv [q-bio.NC]* (2022) doi:10.48550/ARXIV.2206.09398.

138. Demetci, P., Santorella, R., Sandstede, B., Noble, W. S. & Singh, R. SCOT: Single-
cell multi-omics alignment with optimal transport. *J. Comput. Biol.* **29**, 3–18 (2022).

139. Li, Z. *et al.* Chromatin-accessibility estimation from single-cell ATAC-seq data with
scOpen. *Nat. Commun.* **12**, 6386 (2021).

140. Angerer, P. *et al.* destiny: diffusion maps for large-scale single-cell data in R.
Bioinformatics **32**, 1241–1243 (2016).

141. Granja, J. M. *et al.* ArchR is a scalable software package for integrative single-cell
chromatin accessibility analysis. *Nat. Genet.* **53**, 403–411 (2021).

142. Milacic, M. *et al.* The reactome pathway knowledgebase 2024. *Nucleic Acids Res.* **52**,
D672–D678 (2024).

143. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene
Ontology Consortium. *Nat. Genet.* **25**, 25–29 (2000).

144. Dixon, P. M. Nearest-neighbor contingency table analysis of spatial segregation for
several species. *Écoscience* **9**, 142–151 (2002).

145. de la Cruz Rot, M. Métodos para analizar datos puntuales. in *Introducción al análisis*
espacial de datos en ecología y ciencias ambientales: métodos y aplicaciones 75–127
(Servicio de Publicaciones, 2008).

146. Kleshchevnikov, V. *et al.* Cell2location maps fine-grained cell types in spatial
transcriptomics. *Nat. Biotechnol.* **40**, 661–671 (2022).

ED Data Figure 1 Cross-species identification and segregation analysis of CFLAR+ cardiomyocytes across spatial and single-nucleus datasets. a, Overview of human (left panel) and mouse (right panel) cardiac datasets used in this study. Bar plots indicate the number of samples per modality, including Xenium, snRNA-seq, MERFISH and MERFISH V2.0. UpSet-style plots summarize dataset overlap and availability across modalities in human and mouse hearts. **b,** Overview of the spatial transcriptomic panels used and their overlap. **c,** Comparison of library complexity (adult human snRNA) with Kuppe et al., 2022. Shown are unique genes per UMI (library complexity). The gene detected by counts per nucleus slope is comparable, while this dataset shows a higher top end. **d,** Heatmaps showing mean expression of SYNPO, SYNPO2, SYNPO2L, and WIF1 across major cardiomyocyte (CM) subtypes in human. DES_SYNPO+ CMs display selective enrichment of SYNPO family genes,

indicating a distinct cytoskeletal-associated CM program conserved across species. **e,** Projection of cardiomyocyte nuclei from Kuppe et al., 2022 onto a low-dimensional embedding, annotated by CM clusters. Feature plots illustrate expression of CFLAR, MYL2, DES, and MYH7B. The accompanying heatmap summarizes mean gene expression per CM cluster, highlighting segregation of CFLAR[+] CMs from classical contractile CM states. **f,** Independent validation in the Reichardt et al., 2022 human ventricular cardiomyocyte dataset. Left, UMAP embedding of ventricular CM subclusters, where clusters are name as in the original publication (vCM1–vCM5). Middle, feature plots showing expression of CFLAR, MYL2, DES, and MYH7B. Right, heatmap of scaled mean expression per vCM subtype, confirming selective enrichment of CFLAR and MYH7B in distinct CM populations. **g,** Quantification of the proportion of CFLAR[+] cardiomyocytes across disease conditions (control, ischemic cardiomyopathy with acute myocardial infarction (ICM_AMI), ischemic cardiomyopathy (ICM), and dilated cardiomyopathy (DCM)). Each dot represents one patient sample, illustrating disease-associated shifts in CM state composition. Stars indicate a FDR<0.1 based on scCODA abundance test. **h,** Differential cell-type proportion analysis across disease groups. Heatmap depicts log-fold changes in relative cell abundance between pairwise disease comparisons. Asterisks indicate credible changes by the scCODA test (False Discovery Rate=0.1). CFLAR[+] CM populations show selective expansion in cardiomyopathy compared to control hearts, while other CM and non-CM populations display distinct disease-specific remodeling patterns.

Extended Data Figure 2 Cardiac tissue segmentation optimization. a, Representative human cardiac tissue section profiled by high-resolution spatial transcriptomics Xenium 5k. Left, spatial distribution of detected transcripts (>10,000) within a manually outlined cardiomyocyte. Right, corresponding immunofluorescence image showing wheat germ agglutinin (WGA, cell boundaries; green) and DAPI (nuclei; blue). Scale 10 um. **b,** Quantitative comparison of segmentation methods using image-based and feature-based metrics. Image metrics include aggregated Jaccard index and mean average precision (mAP, IoU ≥50%) relative to reference annotations. Feature metrics include accuracy of cell number estimation (vs nucleus segmentation) and fraction of assigned transcripts. Our heartbreaker model

achieves the best overall balance between spatial accuracy and transcript assignment, whereas nucleus-only and default platform segmentations underperform in cardiomyocyte-dense tissue. **c,** Visual comparison of segmentation outputs overlaid on cardiac tissue images (segmented boundaries in white). Methods include Cellpose (membrane), Baysor, BIDCell, Proseg, Vizgen default segmentation, and Cellpose nuclei. Membrane-aware methods more accurately capture elongated cardiomyocyte morphology and reduce over-segmentation. **d,** Comparison of cell-boundary staining strategies for spatial transcriptomics platforms. MERFISH (Vizgen), left section, relies on intrinsic membrane signal, whereas Xenium (10X Genomics), right section, benefits from additional post-hoc membrane labeling to improve boundary definition in cardiomyocytes. **e,** Xenium cardiac tissue segmentation comparison. Top, default Xenium cell boundary staining and segmentation. Bottom, post-Xenium WGA staining followed by watershed segmentation, resulting in improved delineation of elongated cardiomyocytes and more accurate cell boundaries.

Extended Data Figure 3 Histopathology foundation model–based tissue state annotation across spatial transcriptomics platforms. a, Schematic overview of the histology-based tissue classification workflow using a deep learning foundation model. Whole-slide images (WSI) and tissue microarrays (TMA) stained with haematoxylin and eosin (H&E) were subdivided into non-overlapping image patches (224 × 224 pixels) and embedded using the UNI transformer foundation model (Chen *et al.*, 2024). Latent representations were clustered in an unsupervised manner to define recurrent myocardial tissue states, including intact myocardium, multiple damaged myocardium states, fresh infarct with demarcation, granulation tissue, fibrotic scar tissue, necrotic tissue, and adipose tissue. Representative image patches for each tissue state are shown. **b,** Joint embedding of spatial transcriptomics

data annotated by foundation model–derived tissue classes across platforms and species. UMAP projections of Xenium (human), MERFISH (human), and Xenium (mouse) datasets demonstrate consistent segregation of myocardial tissue states, indicating cross-platform and cross-species robustness of the histology-driven tissue classification. **c,** Comparison of expert tissue annotation and foundation model–based tissue analysis on whole heart sections. Left, tissue expert annotations highlighting infarcted cardiac tissue, vasculature, adipose tissue, and pericardium. Right, corresponding foundation model predictions resolving fine-grained myocardial damage states and infarct-associated tissue regions. Arrows indicate regions of concordance and refined subclassification by the model. **d,** Association between tissue states and cellular composition. Heatmap showing enrichment and depletion of cell types within foundation model–defined tissue regions using Fisher's exact test ($\log_2$ odds ratios). Red indicates enrichment, blue depletion; asterisks denote statistically significant associations after multiple-testing correction (adjusted p-values < 0.05). Fibrotic and granulation tissue regions are enriched for fibroblasts, macrophages, endothelial, and perivascular cells, whereas intact myocardium is selectively associated with cardiomyocyte populations. **e,** Segregation analysis of cardiomyocyte subtypes in MERFISH 2.0 data (n=13). Segregation index (SI) comparison shows significantly stronger spatial segregation (clumping) of DES$^+$ cardiomyocytes compared with CFLAR$^+$ cardiomyocytes, indicating differential spatial organization relative to tissue damage states. Box plots show median, interquartile range, and individual samples; *** p-value < 0.001 Wilcoxon rank-sum test.

Extended Data Figure 4 Spatial niche remodeling and disease trajectory in human cardiomyopathy. a, Overview of the spatial transcriptomics workflow. Xenium spatial transcriptomic data from control and ischemic cardiomyopathy (ICM) samples were analyzed using the NicheSphere framework to quantify cell–cell colocalization, graph-based interactions, and to define spatial tissue niches. TMA samples were subsequently ordered along a disease trajectory using PILOT pseudotime analysis in accordance to their niches. **b,** Comparison of tissue niche composition versus patch-based histological annotation. Heatmap shows enrichment of defined tissue niches (Xenium-TMA) across histopathological regions, including myocardium, damaged myocardium (subtypes 1–3), fibrotic/scar tissue, and granulation tissue. Asterisks indicate statistical significance (adjusted p-value < 0.05; Fisher's exact test). **c,** Changes in niche composition (y-axis) across disease progression (x-axis).

Scatter plots show proportions of selected niches along the inferred disease trajectory, including DES$^+$/DES_SYNPO$^+$/NPPA_NPPB$^+$ cardiomyocytes, resident immune niches, lymphoid endothelial niches, and neutrophil–venous endothelial niches. Lines indicate smoothed trends; p-values from F statistic test denote significance of association with disease progression. **d,** Violin plots of CCDC80 gene expression (normalized log counts) across major fibroblast-related cell states in snRNA, including C7$^+$ fibroblasts, ELN$^+$ fibroblasts, general fibroblasts, myofibroblasts, and CTHRC1$^+$ fibroblasts. **e,** Heatmap with genes selected (by PILOT) to be associated with the trajectory in CFLAR CM cells. They include cardiomyocyte structural and cytoskeletal genes across disease progression, illustrating coordinated downregulation of sarcomeric genes (e.g., *TNNT2*, *MYBPC3*) and upregulation of remodeling-associated genes (e.g., *DST*, *PDE4DIP*, *TTN*). **f,** NicheSphere interaction network comparing DCM and control samples (Xenium-TMA). Nodes represent cell types or niches, edges represent differential co-localization scores, and edge color indicates increased (red) or decreased (blue) interactions in disease. **g,** Differential co-localization heatmap showing changes in spatial associations between cardiomyocyte states and surrounding cell types in control versus DCM samples. Color scale indicates diffColoc scores. For **f** and **g**, only statistically significant interactions are shown(adjusted Wilcoxon's rank sums test; p-value < 0.05). **h,** Top, disease trajectory using PILOT pseudotime analysis, Bottom, Proportional changes of selected cardiomyocyte-centered niches along disease progression, including myocardium_CFLAR$^+$ cardiomyocytes, DES_SYNPO$^+$ cardiomyocytes, and myofibroblast–macrophage niches. Trend lines and p-values indicate significant associations with pseudotime. **i,** Association between spatial niches and histological patches. Heatmap displays the fraction of each niche within defined tissue patches, highlighting preferential localization of cardiomyocyte niches to myocardium and damaged myocardium, and fibroblast–immune niches to fibrotic or granulation tissue. Significance was determined by Fisher's exact test with adjusted p-values (p < 0.05). **j,** Representative Xenium-TMA sections (five selected samples along the Trajectory from early to middle and late stages) illustrating (top) spatial distribution of tissue niches, (middle) corresponding histopathological patches, and (bottom) cardiomyocyte subtype localization and marker gene expression. MYL6 expression highlights cardiomyocytes, whereas CCDC80 expression marks fibroblast-rich regions. Arrow indicates progression along the PILOT pseudotime trajectory. **k,** Comparison of spatial clustering of niches in ICM and DCM samples. Shown are mean self-enrichment scores (spatial segregation) per sample. (Wilcoxon rank-sum test: Z = 4.09, P < 0.0001) **l,** Heatmap of cardiomyocyte gene expression dynamics across disease progression, showing gradual loss of contractile gene programs (e.g., *TNNT2*, *MYL6*, *CASQ2*, *MYBPC3*) and induction of stress and remodeling-associated genes (e.g., *SMG1*, *OBSCN*).

Extended Data Figure 5 Differential spatial niches and subcellular RNA organization in cardiomyopathy. a, Heatmap summarizing differences in spatial niche composition between ischemic cardiomyopathy (ICM)/DCM and control samples. Columns represent annotated cell types, rows indicate identified spatial niches. **b,** NicheSphere interaction network comparing acute myocardial infarction (AMI) with control samples (Xenium-TMA). Nodes represent cell types, edges indicate differential spatial co-localization, with red edges denoting increased and blue edges decreased interactions in disease. Edge thickness corresponds to the magnitude of the differential co-localization score. **c,** Differential co-localization analysis of cardiomyocyte subtypes in Xenium-TMA data. Heatmap displays diffColoc scores for DES$^+$, DES_SYNPO$^+$, NPPA_NPPB$^+$, CFLAR$^+$, and SLC25A4$^+$ cardiomyocytes with surrounding cell types, highlighting disease-associated remodeling of cardiomyocyte neighborhoods. **d,** Proportional changes of niches along disease progression. **e,** Heatmap of subcellular spatial gene expression patterns in cardiomyocytes. Genes are ordered by preferential enrichment across four subcellular compartments—nucleus, perinuclear region, cytosol, and intercalated disc—revealing compartment-specific transcript localization programs. **f,** Enrichment of APEX-seq–derived subcellular reference signatures across the four identified RNA domains. Heatmap shows concordance of spatial transcript localization with known cytosolic, ER membrane (ERM), outer mitochondrial membrane (OMM), ER lumen, lamina, nuclear pore, nucleolus, and nucleus-associated RNA signatures. **g,** Top 15 genes enriched in each subcellular compartment (nucleus, perinuclear region, cytosol, and intercalated disc), ranked by enrichment score, highlighting distinct functional gene sets underlying compartment-specific RNA localization. **h,** Representative high-resolution microscopy image illustrating cardiomyocyte subcellular architecture. White arrows indicate intercalated disc regions. Nuclei are counterstained with DAPI (blue). **h′,** Same field of view as in **h**, overlaid with detected CMYA5 mRNA molecules (yellow dots), demonstrating preferential localization of CMYA5 transcripts at intercalated disc regions. Boundary markers (ATP1A1/CD45/E-cadherin), interior RNA (18S), and interior protein markers (alphaSMA/Vimentin) are shown as indicated in the legend.

Extended Data Figure 6 Nuclear morphology–based classification and polyploidy-associated cardiomyocyte programs. a, Representative nuclear morphologies. DAPI images of isolated cardiomyocyte nuclei from CFLAR[+] and DES[+] CMs illustrating distinct nuclear shapes and texture features (arrows). Scale bar, 20 μm. **b,** Quantitative nuclear morphology features. Comparison of selected CellProfiler-derived DAPI morphology features between DES[+] and CFLAR[+] CM nuclei. Bars indicate Cohen's d effect size; positive values indicate enrichment in DES[+], negative values enrichment in CFLAR[+] nuclei. p-values from two-sided tests (p-value< 0.05; p-value < 0.001). **c,** Morphology feature heatmap. Heatmap of z-scored DAPI morphology features across cardiomyocyte states (DES[+], DES_SYNPO[+], CFLAR[+], NPPA_NPPB[+], SLC25A4[+]), highlighting systematic differences in size, texture, shape, and Zernike moments. **d,** Performance of morphology-based classification (logistic regression). Precision, recall, and F1-score for distinguishing CFLAR[+] and DES[+] CMs using either DINOV3 feature vectors (top) or CellProfiler based nuclear morphology measurements (bottom). **e,** Polyploid nuclei isolation by FANS. Experimental overview and cell numbers of fluorescence-activated nuclei sorting (FANS) from FFPE samples into 2N, 4N, and 8N

fractions, including sample-specific contributions. Bottom panel shows representative DNA content distributions. **f,** Integrated single-nucleus landscape. UMAP embedding of all cardiac cell types (left), annotated cell types with cardiomyocytes highlighted (middle), and integrated samples colored by donor (right). **g,** Cardiomyocyte-only embedding. UMAP representation restricted to cardiomyocytes, capturing continuous transcriptional variation across CM states. **h,** Cell-state signature projection. UMAPs showing enrichment of CFLAR$^+$, DES$^+$, and NPPA_NPPB$^+$ gene signatures across cardiomyocytes. **i,** Polyploid gene signatures. Heatmap summarizing enrichment of CFLAR, DES, and NPPA_NPPB signatures across ploidy classes (2N, 4N, 8N). **j,** External CM classification reference Wehrens et al.. UMAP of cardiomyocytes annotated according to the classification by Wehrens *et al.*, showing correspondence to DES-like, NPPA_NPPB$^+$, CFLAR-like, and unclassified CM populations. **k,** Signature score validation. UMAPs displaying CFLAR$^+$ CM and DES$^+$ CM signature scores, confirming spatial separation of CM programs. **l,** Cluster fold-similarity analysis. Heatmap showing similarity between adult human CM states and reference DES$^+$ or CFLAR$^+$ programs. Asterisks denote significant enrichment ($P < 0.05$).

Extended Data Figure 7 Metabolic, cell-cycle, electrophysiological, and epigenetic programs distinguish diploid and polyploid cardiomyocyte states. a, Schematic overview of central metabolic pathways analyzed in cardiomyocytes, highlighting glycolysis, fatty acid uptake and β-oxidation, and mitochondrial acetyl-CoA entry into the tricarboxylic acid (TCA) cycle. Key transporters and enzymes are indicated. **b,** Heatmap showing expression of fatty acid uptake and β-oxidation–related genes across cardiomyocyte states, including CFLAR[+] subclusters, HAND2[+], DES[+], NPPA_NPPB[+], SLC25A4[+], and DES_SYNPO[+] cardiomyocytes. Values represent mean expression scaled per gene **c,** Heatmap of glutaminolysis and TCA cycle–associated genes across cardiomyocyte states, illustrating differential utilization of anaplerotic pathways. **d,** Enrichment analysis -log10 (pvalues). Enriched terms are dominated

by mitochondrial respiration, oxidative phosphorylation, and ATP synthesis pathways. **e,** Conceptual schematic of core cell-cycle regulators and their activity across G1, S, G2, and M phases, including MYC, cyclins, CDKs, and mitotic regulators. **f,** Heatmap of cell-cycle regulator expression across cardiomyocyte states, revealing selective enrichment of G1/S and G2/M regulators in specific CFLAR$^+$ subclusters relative to other cardiomyocyte populations. **g,** Heatmap of ion channels, transporters, connexins, and receptors relevant for cardiomyocyte electrophysiology and intercellular coupling. Channels are grouped by functional class (e.g., HCN, Ca^{2+}, K$^+$, Cl$^-$ channels, connexins). Mean expression is scaled per gene. **h,** Heatmap of genes involved in histone acetylation, histone/DNA methylation, chromatin remodeling, genome architecture, and nuclear stability across cardiomyocyte states, highlighting enhanced epigenetic and chromatin regulatory programs in CFLAR$^+$ cardiomyocytes. **i,** Heatmap summarizing higher-order functional programs across cardiomyocyte states, including metabolic maturation and mitochondrial programs, postnatal maturation and hypertrophy, lipid/cholesterol and inflammatory crosstalk, developmental/regenerative signaling axes, and xenobiotic/stress responses. **j,** Conceptual model contrasting diploid and polyploid cardiomyocytes. Diploid cardiomyocytes are characterized by preserved metabolic maturation, mitochondrial organization, and WNT antagonism (e.g., DKK3, SFRP1), whereas polyploid cardiomyocytes display activation of Hippo, WNT, TGFβ, and SHH signaling, altered genome architecture, and stress-adaptive transcriptional programs. **k**, TF Activity of CTCF within the PHLOWER estimated CM trajectory.

Extended Data Figure 8 Disease-specific intercellular signaling programs linking cardiomyocyte states to fibrosis and immune remodeling. a–c, Heatmaps showing mean scaled expression of pathway-associated genes across cardiomyocyte states. **a,** Developmental signaling pathways, including WNT, SHH, and NOTCH components, display selective activation in CFLAR+ and HAND2+ cardiomyocytes. **b,** Fibrosis-associated signaling genes (e.g. TGFβ, PDGF, VEGF, FGF pathways) highlight enhanced profibrotic communication potential of specific cardiomyocyte states. **c,** Immune signaling genes, including chemokines and cytokines, indicate differential immune-modulatory capacity across cardiomyocyte populations. **d,** PageRank analysis of cell–cell communication networks in ICM versus control. Positive values indicate higher cell-communication in ICM and negative values indicate lower cell-cell communication in ICM. **e–f,** Sankey plot with top ligand–receptor

interaction highlighting ongoing CFLAR$^+$-myofibroblast specific communication relative to other cardiomyocyte states in ICM vs control. **e,** Sankey plot with top ligand-receptor interactions between CFLAR$^+$ cardiomyocytes towards myofibroblasts, emphasizing fibrosis-related signaling axes. **f,** Reciprocal interactions from myofibroblasts, illustrating cardiomyocyte-driven profibrotic signaling. **g,** PageRank analysis of cell–cell communication networks in DCM versus control. Positive values indicate higher cell-communication in DCM and negative values indicate lower cell-cell communication in DCM. **h,** Sankey diagram with top ligand-receptor interactions with DES_SYNPO$^+$ as target cell in DCM versus control. **i,** Sankey diagram with top ligand-receptor interactions with DES_SYNPO$^+$ as source cell in DCM versus control.

Extended Data Fig. 9 Cell-cycle activity and maturation trajectories of embryonic cardiomyocytes. a, UMAP of embryonic human snRNA-seq data highlighting cardiomyocyte differentiation states. **b**, Min-max scaled expression of canonical cell-cycle genes across embryonic LV-CM subtypes. **c**, Density and joint distribution of S- and G2/M-phase enrichment scores in embryonic versus adult cardiomyocytes. **d**, Cell-cycle phases across matched cardiomyocyte states in adult and embryonic datasets. **e**, Expression of cardiomyocyte maturation markers across embryonic LV-CM subsets. **f,** Gene set enrichment analysis (GSEA) of embryonic cardiomyocyte states highlighting developmental, metabolic, and cell-

cycle programs. **g,** Quantitative association of embryonic CM states and non-CM cell types with anatomical spatial domains, illustrating coordinated patterning of CM maturation with cardiac tissue architecture, as measured by scaled cell counts per compartment estimated by deconvolution **h,** Embryonic cell states mapped to Visium data (estimated counts per Visium spot) on a representative sample. **i,** PHLOWER trajectory with selected embryonic cardiomyocytes cells mapped onto adult CM branches, indicating immature LV-CMs to be related to the CFLAR branch.

ED Figure 10 Cardiomyocyte state composition, metabolic rewiring, signaling programs, and spatial niches during disease progression in mice. a, Heatmap showing mean expression of selected cardiomyocyte (CM) marker genes across Myh7b[+], Nppa[+], and Des[+] CM subclusters, confirming transcriptional identity of polyploid (Myh7b[+], Nppa[+]) and

diploid (Des[+]) CM states. Expression is scaled per gene. **b,** Differential cell proportion analysis across experimental conditions using scCODA. Heatmap indicates credible increases (blue) or decreases (red) in cell type abundance relative to control (FDR = 0.1). Asterisks denote statistically credible changes. Comparisons include control vs d3, d7 vs d3, d14 vs d3, and d28 vs d3. **c,** Enrichment of CM differentiation and cell cycle–related gene signatures across CM states. Polyploid Myh7b[+] and Nppa[+] CMs show enrichment of immature, stress-responsive, and cycling programs, whereas Des[+] CMs align with terminal differentiation states. **d–f,** Heatmaps showing scaled mean expression of genes involved in **glycolysis** (d), **glutaminolysis** (e), and **fatty acid oxidation** (f) across CM subclusters. Polyploid CM states display increased glycolytic and glutaminolytic programs, while diploid Des[+] CM states retain higher fatty acid oxidation signatures. **g–i,** Expression of signaling pathway components across CM states. **g,** Developmental signaling pathways (e.g., WNT, NOTCH, Hedgehog). **h,** Fibrosis-associated signaling (e.g., TGF-β, PDGF, VEGF, EGFR). **i,** Immune-related signaling and chemokine expression. Polyploid CM states preferentially activate developmental and pro-fibrotic signaling programs. **j,** Heatmap of intracellular pathway activity scores in cardiomyocyte (CM) subtypes at day 28 post-injury compared with controls. Pathways include Toll-like receptor, Notch, TGF-β, chemokine, JAK–STAT, Wnt, NF-κB, and Hedgehog signaling. Values represent normalized pathway activity, with clustering revealing differential inflammatory and developmental signaling across CM states (Nppa[+], Des[+], and Myh7b[+] subsets).

Extended Data Fig. 11 Spatial and transcriptional remodeling of cardiomyocyte states and therapeutic target engagement in ischemic cardiomyopathy. a, Low-dimensional embedding illustrating the trajectory of disease progression, separating control samples (blue) from ischemic cardiomyopathy (ICM; red), indicating a coordinated shift in global transcriptional states with disease. **b,** Spatial tissue organization inferred from five samples selected along the trajectory of **a**. Top: tissue niches (1–6) highlighting myocardium, damaged

myocardium, fibrotic and scar tissue, and granulation tissue. Middle: spatial patches derived from unsupervised segmentation, reflecting recurrent micro-architectural patterns. Bottom: spatial expression of *Postn* in fibroblasts, illustrating fibrosis-associated gene expression across disease-affected regions. **c,** Cell type–resolved expression of major heart failure drug targets in mouse, including the "fantastic four" (β-blockers, MR antagonists, ARNI, and SGLT2 inhibitors; bold). Heatmap shows relative target gene expression across cardiomyocyte subtypes, vascular cells, adipocytes, and immune populations. Red dashed boxes highlight cardiomyocyte-enriched targets, while green dashed boxes indicate pericyte/VSMC-associated signaling nodes (e.g., PDE5A and sGC pathways), underscoring multicellular mechanisms of therapeutic action. **d**, Volcano plot between CFLAR+ and DES+ human CM. Highlighted are genes encoding heart failure targets. A line indicates the threshold of 0.05 FDR.

Extended Data Fig. 12 TNIK expression across species and models and functional effects of TNIK inhibition. a, Mean TNIK expression across human cardiac and non-cardiac cell types, highlighting enrichment in cardiomyocyte subpopulations, including CFLAR[+] and HAND2[+] cardiomyocytes, compared with stromal and immune compartments. **b,** Mean Tnik expression across mouse heart cell types, demonstrating conserved cardiomyocyte-biased expression relative to non-myocyte populations. **c,** TNIK expression across iPSC-derived cardiomyocyte differentiation stages and subtypes, with highest expression observed in immature and MYH7B[+] cardiomyocyte states. **d,** Mean TNIK expression across rat cardiac cell populations, confirming cross-species conservation of cardiomyocyte-enriched TNIK expression. **e,** UMAP projection of iPSC cardiomyocytes colored by sample and experimental condition, showing robust integration across datasets. **f,** The same UMAP colored by health status, demonstrating overlap between control and diseased cardiomyocytes without condition-driven batch effects. **g,** Pathway enrichment scores inferred from cardiomyocyte transcriptomes of WTC11 and AICS-TTN iPSC-derived cardiomyocytes under control

conditions or following TNIK inhibition (NCB-0846). *** p-value < 0.001 versus vehicle, two-sided Student's t-test of the linear model slope **h**, Quantification of sirius red positive cells for all treatment groups **i,** Weekly body weight trajectories following myocardial infarction (MI) and treatment with vehicle, INS018_055 (two doses), or pirfenidone. TNIK inhibition does not induce weight loss compared with vehicle or antifibrotic control. Data are shown as mean ± s.e.m.; * p-value < 0.05, ** p-value < 0.01 versus MI + vehicle. **j,** Left ventricular mass, LVAWTd and LVAWTs for all treatments groups, Representative M-mode echocardiography images from sham-operated animals, MI + vehicle, and MI treated with INS018_055 (2.5 mg/kg, BID), illustrating improved cardiac contractile performance following TNIK inhibition.