1

Temperature and intrinsic Ca^{2+} reshape TRPM4 pharmacology

Jinhong Hu[1], Sofia Ievleva[1], Sung Jin Park[1], Junuk Lee[1], Jennifer Cheng[1], Garrett O'Dea[1],
Jiangnan Sheng[1], Juan Du[#,1,2,3] & Wei Lü[#,1,2,3]

1. Department of Molecular Biosciences, Northwestern University, Evanston, IL, USA
2. Department of Pharmacology, Feinberg School of Medicine, Northwestern
University, Chicago, IL, USA
3. Chemistry of Life Processes Institute, Northwestern University, Evanston, IL, USA

21 **#CORRESPONDING AUTHOR**

22 Correspondence and requests for materials should be addressed to J. D. (email:

23 juan.du@northwestern.edu) TEL: (847) 467-6719, or W. L. (email: wei.lu@northwestern.edu).

24 TEL: (847) 491-5440

Abstract

Proteins operate in dynamic environments where ions, lipids, and temperature collectively define their properties, yet most studies rely on simplified conditions that overlook these intrinsic variables. Here, we show two such factors—temperature and Ca^{2+}—profoundly remodel the function and pharmacology of TRPM4, an ion channel implicated in cardiac conduction, immune regulation, and cancer. At physiological temperature and Ca^{2+}, TPPO—previously considered a selective TRPM5 inhibitor inactive toward TRPM4—potently activates TRPM4, revealing strong synergy among temperature, Ca^{2+}, and ligand binding. In contrast, Necrocide-1, a necroptotic activator targeting the same binding pocket, defies this logic: it opens TRPM4 without Ca^{2+} but is antagonized by Ca^{2+}. Meanwhile, the inhibitors NBA and CBA engage a nearby pocket, locking the channel in a non-conductive pre-open state. Our findings highlight that even rigid binding pockets can exhibit temperature-dependent ligand recognition, revealing hidden pharmacology and informing selective, environment-aware therapeutic strategies.

Introduction

Protein activity is inherently governed by the cellular environment in which it operates. A wide array of factors—including ions, lipids, temperature, pH, and protein–protein interactions—collectively determine how a protein functions in its native context. Yet, because of technical constraints, most biophysical and pharmacological studies have been conducted under simplified conditions, which have yielded foundational insights but often fail to capture the full physiological complexity.

Increasing efforts now aim to bridge this gap by incorporating more native variables—for example, using nanodiscs, liposomes or membrane-derived vesicles to preserve native lipidic environments for membrane proteins[1-4], and purifying proteins from native sources to retain endogenous protein compositions and modifications[5-8]. Despite these advances, one fundamental determinant, temperature, is largely overlooked in mechanistic and pharmacological studies, even though it influences every protein in the human body, where proteins operate at 37°C. Because most in vitro assays are performed at room temperature or lower, therefore, the broad impact of temperature on protein conformation, ligand recognition, efficacy and mechanism remains unexplored, with only a few recent exceptions[9,10].

Alongside temperature, intracellular Ca^{2+} represents another intrinsic physiological variable that exerts pervasive influence on protein behavior. Although not universal, Ca^{2+} acts as a selective yet broad regulator of cellular physiology. It is among the most tightly controlled cellular cofactors, with cytosolic concentrations fluctuating dynamically from tens of nanomolar at rest to low micromolar during signaling, and rising to abnormally high levels under pathological conditions such as ischemia, excitotoxicity, or cancer[11]. Many proteins are directly regulated by Ca^{2+} through dedicated binding sites, while others respond indirectly through Ca^{2+}-dependent signaling cascades. Although the structural and functional roles of Ca^{2+} have been extensively characterized, its impact on protein–ligand interactions—particularly in combination with other intrinsic factors like temperature—has rarely been considered. This represents an important gap in our understanding of how Ca^{2+}-regulated proteins engage ligands under physiological and pathological conditions.

The Ca^{2+}-activated, temperature-sensitive channel Transient Receptor Potential Melastatin 4 (TRPM4) provides an ideal model to investigate how intrinsic physiological factors remodel

pharmacology[12-15]. TRPM4 is a non-selective monovalent cation channel that contributes to

cardiac conduction, insulin secretion, neuronal excitability, immune regulation, and intestinal

fluid homeostasis[16-25]. Pathogenic variants have been implicated in Brugada syndrome, inherited

conduction disorders, and diverse cancers[26-34]. Despite its broad physiological and pathological

relevance, TRPM4 has remained largely untargeted by clinical pharmacology, with the single

exception of bisacodyl—a widely used drug for chronic constipation—and its active metabolite

deacetyl bisacodyl, which we recently identified and characterized as clinically relevant TRPM4

ligands[25].

Previous structural and functional studies established that TRPM4 adopts two major

conformations, a "cold" and a "warm" state, whose equilibrium depends on both temperature and

cytosolic Ca^{2+} concentration[10]. These findings suggest that biophysical and pharmacological

assays performed under room temperature and undefined Ca^{2+} conditions may not accurately

capture the channel's behavior under physiological settings.

Here, we systematically examine TRPM4 pharmacology under conditions that incorporate

both temperature and Ca^{2+}. Through combined cryo-EM and electrophysiology, we reveal that

these intrinsic factors profoundly shape ligand recognition, efficacy, and mechanism. Our results

identify the S1–S4 domain as a dynamic regulatory hub that integrates environmental and

chemical cues to govern activation and inhibition. More broadly, this work establishes an

environment-aware framework for pharmacology, in which incorporating intrinsic variables

exposes hidden drug activities and new mechanisms of regulation. For widely expressed proteins

like TRPM4, such principles open opportunities to design therapeutics that exploit pathological

local conditions—such as abnormally elevated Ca^{2+} or altered thermal states—for selective

action in disease, while preserving normal physiological function and minimizing side effects.

Results

Temperature reveals hidden pharmacology of TRPM4

During the cold-to-warm transition of TRPM4, Ca^{2+} and temperature drive major conformational

changes in the intracellular domain but only subtle shifts in the transmembrane domain[10]. From

these structures, we identified three categories of ligand binding sites that emerge elusively in the

warm conformation of the intracellular domain[10]. While this outcome may be anticipated from

the magnitude of ICD rearrangement, it nevertheless raises two central questions: whether drug

screening performed exclusively at room temperature may have overlooked active compounds,

and whether temperature-dependent ligand recognition is confined to regions undergoing large

conformational change (such as the intracellular domain) or extends to regions with more modest

movements (such as the transmembrane domain).

To address these questions, we examined several previously reported TRPM4 ligands, as well

as compounds considered inactive toward TRPM4. We focused on hydrophobic ligands predicted

to interact with the transmembrane domain, including triphenylphosphine oxide (TPPO),

necrocide-1 (NC1), and the anthranilic acid derivatives CBA (4-chloro-2-[2-(2-chloro-phenoxy)-

acetylamino]-benzoic acid) and NBA (4-chloro-2-(2-(naphthalene-1-yloxy) acetamido) benzoic

acid). TPPO, originally identified as a selective TRPM5 inhibitor with an IC_{50} of 12 μM, has

been found inactive on TRPM4 when tested at room temperature using fluorescence-based

membrane potential assays[35]. NC1 was recently characterized as a TRPM4 activator that

promotes Na^+ influx and triggers necrotic cell death[36], suggesting therapeutic potential in cancer.

NBA and CBA represent the best-characterized TRPM4 inhibitors to date with IC_{50} in the sub-

micromolar range[37]. Importantly, all prior functional and structural studies were conducted at

room temperature or lower.

Using patch-clamp electrophysiology, we tested these compounds at both room temperature

and 37°C. Under previously reported assay conditions—room temperature with either zero free

intracellular Ca^{2+} or basal free intracellular Ca^{2+} (100 nM), we reproduced established results: 50

μM TPPO showed negligible activity on TRPM4—comparable to the control group, where basal

Ca^{2+} alone is insufficient to drive channel activation (Fig. 1a,b; Extended Data Fig. 1a,b); NC1

activated the channel, eliciting largely linear, voltage-independent currents (Fig. 2a,b); and NBA

and CBA produced robust inhibition of Ca^{2+}-induced TRPM4 current (Extended Data Fig. 2a).

Strikingly, elevating the temperature to 37°C revealed a hidden pharmacological behavior. While

NC1, NBA, and CBA retained their room-temperature profiles (Fig. 2d,e; Extended Data Fig.

2a), TPPO displayed a completely different response—robustly activating TRPM4 currents (Fig.

1e; Extended Data Fig. 1e). These observations highlight a critical gap in conventional

pharmacological screening: assays performed under non-physiological conditions may both

overlook potential active compounds and misclassify ligand selectivity.

Ca^{2+} as a key determinant of agonist recognition and potency in TRPM4

Ca^{2+} is the only known endogenous agonist of TRPM4, and together with temperature, drives the transition between its cold to warm conformations[10]. Building on our findings that temperature reshapes TRPM4 pharmacology (Fig. 1e), we next explored whether Ca^{2+} modulates the binding and potency of the exogenous activators TPPO and NC1.

Under Ca^{2+}-free conditions achieved by EGTA buffering, 50 uM TPPO produced negligible activation of TRPM4 (Fig. 1d; Extended Data Fig. 1d), whereas adding 100 nM free intracellular Ca^{2+} enabled robust activation (Fig. 1e, red traces; Extended Data Fig. 1e). Thus, even basal Ca^{2+} concentrations—although insufficient to open the channel alone (Fig. 1e, black trace; Extended Data Fig. 1e)—markedly enhance TPPO-induced TRPM4 activation. To quantify this apparent synergy, we determined TPPO dose-response relationships. Physiological temperature increased TPPO potency by ~14-fold under basal Ca^{2+} (Fig. 1g,h; Extended Data Fig. 1g,h), while basal Ca^{2+} increased TPPO potency by ~6-fold at 37 °C (Fig. 1g,i; Extended Data Fig. 1g,i). In the absence of both Ca^{2+} and physiological temperature, TPPO activated TRPM4 only weakly at submillimolar concentrations and failed to reach saturation even at millimolar levels. When intracellular Ca^{2+} reach activating level (1 µM), the effect of TPPO is largely masked, consistent with the notion that TPPO and Ca^{2+} act though a shared activation pathway, with Ca^{2+} dominating at high concentration, as both produce the hallmark outwardly rectified currents (Fig. 1c,f; Extended Data Fig 1c,f).

Together, these data demonstrate that physiological context is essential for TPPO efficacy, revealing a three-way synergy between temperature, Ca^{2+}, and ligand binding. This context dependence overturns the view that TPPO is inactive toward TRPM4, revealing instead that it potently activates the channel—opposite to its inhibitory action on the closely related TRPM5[35]. This finding underscores how ligand behavior can be fundamentally mischaracterized when assays are performed under non-physiological conditions.

We next examined whether Ca^{2+} influences NC1, previously reported as the first Ca^{2+}-independent agonist of TRPM4[36]. Under Ca^{2+}-free and 100 nM Ca^{2+} conditions, NC1 evoked nearly identical currents with linear current-voltage relationships distinct from Ca^{2+}-activated currents, confirming that NC1 can activate TRPM4 independently with an EC$_{50}$ in the submicromolar range (Fig. 2a,b,d,e,g,h; Extended Data Fig. 3a). However, at 1 µM Ca^{2+}, a

physiologically relevant level during cellular stress that robustly activates TRPM4, the response

changed dramatically, becoming strongly outwardly rectified (Fig. 2c,f)—a hallmark of

canonical Ca^{2+}-mediated activation[14]. This striking reversal suggests that NC1 is not simply

Ca^{2+}-independent: once Ca^{2+} reaches activating levels, its action dominates, functionally

overriding NC1-mediated gating even at saturating NC1 concentrations (Fig. 2i). Thus, rather

than bypassing the Ca^{2+} requirement, NC1 displays a biphasic Ca^{2+} dependence: permissive at

basal Ca^{2+} yet antagonized at higher concentrations, likely either through binding competition or

allosteric coupling between NC1 and Ca^{2+} binding sites. This mechanism is discussed further in

the section "Mechanistic basis for Ca^{2+}-dependent gating reversal of NC1".

Taken together, these results reveal distinct modes of Ca^{2+}-agonist interplay in TRPM4.

TPPO requires basal Ca^{2+} (and temperature) to act effectively on the channel, whereas NC1

functions independent of Ca^{2+} at rest but loses efficacy once Ca^{2+} rises to stimulatory levels. This

contrast illustrates how the same intrinsic cofactor can differentially tune ligand action through

cooperative or antagonistic coupling. More broadly, they reinforce the importance of evaluating

ion channel pharmacology under physiological conditions, as environmental context can

fundamentally alter both efficacy and apparent selectivity.

Structural basis of the three-way synergy between TPPO, temperature and Ca^{2+}

To elucidate how temperature and Ca^{2+} shape TPPO activation of TRPM4, we determined cryo-

EM structures of the channel in the presence of TPPO at 18 °C and 37 °C, under both Ca^{2+}-free

and Ca^{2+}-saturated conditions. All the structures were resolved to high quality at ~2.5–2.9 Å

resolutions, allowing unambiguous identification of ligand binding sites (Extended Data Figs. 4–

6; Extended Data Table 1).

In the consensus map of the Ca^{2+}/TPPO–TRPM4–37°C dataset, we observed a prominent

tripod-shaped density in the S1–S4 domain of each protomer. This feature closely matches the

shape of TPPO, and sits immediately above the Ca_{TMD} site (Fig. 3a). We therefore designated this

pocket as the S1–S4$_{Upper}$ site. This site spatially overlaps with the deacetyl bisacodyl binding

pocket we identified previously[25]. Subunit-level classification revealed that where warm and cold

conformations coexist in this dataset, with the warm state predominating (Extended Data Fig. 4).

The ligand density is markedly stronger in the warm state, whereas a weaker yet still discernible

density exists in the cold state (Fig. 3d; this minor cold state is discussed at the end of this

section). Within this pocket, TPPO occupies a hydrophobic cavity and forms a critical polar

interaction between its phosphoryl moiety and R1072 on the TRP helix (Fig. 3d, left panel). To

validate this assignment, we mutated key coordinating residues to alanine and performed

electrophysiological recording. Most mutations impaired Ca^{2+}-dependent activation, likely by

disrupting the integrity of the adjacent Ca_{TMD} site, making it impossible to unambiguously isolate

their effects on TPPO, as TPPO requires Ca_{TMD} to manifest its activity (Extended Data Fig. 1j).

In contrast, R1072A preserved Ca^{2+}-evoked gating yet abolished TPPO-dependent activation,

thereby confirming the functional relevance of the TPPO binding site (Extended Data Fig. 1k).

In the Ca^{2+}-bound cold conformation from the Ca^{2+}/TPPO–TRPM4–18°C dataset, only a

faint and incomplete signal was discernible at the S1–S4$_{Upper}$ site at very low contour levels (Fig.

3b). The density formed a triangular outline reminiscent of TPPO but was markedly smaller than

the full ligand, consistent with partial or unstable occupancy. In the Ca^{2+}-free cold conformation

from the EGTA/TPPO–TRPM4–37°C dataset, this feature disappeared entirely, with no trace of

TPPO density detectable (Fig. 3c).

Thus, TPPO binding exhibits a clear dependence on the cold–warm conformational

equilibrium of TRPM4, a transition that is cooperatively governed by temperature and Ca^{2+}

(ref[10]). This structural observation coincides with our electrophysiological finding that TPPO,

temperature, and Ca^{2+} act in three-way synergy to promote channel activation (Fig. 1a,b,d,e).

Together, these results suggest that TPPO efficacy is tightly coupled to the conformational

landscape defined by these two intrinsic modulators.

To dissect the mechanism underlying this coupling, we compared the Ca^{2+}-free cold and

Ca^{2+}/TPPO-bound warm structures. In the Ca^{2+}-free cold conformation, R1072 is displaced from

the S1–S4$_{Upper}$ pocket and stabilized by an interaction with Q831 on S2—a key residue involved

in Ca_{TMD} binding—thereby precluding TPPO binding (Fig. 3e). Upon transition to the warm

state, upward movement of the intracellular domain tilts the TRP helix, moving R1072 into

position for TPPO coordination (Fig. 3e). Concurrently, Ca^{2+} binding at the Ca_{TMD} site recruits

Q831 for Ca^{2+} coordination, releasing R1072 from its locked pose for direct interaction with the

TPPO phosphoryl group (Fig. 3e). Together, these coupled rearrangements prime the S1–S4$_{Upper}$

pocket for effective ligand engagement.

Interestingly, a weak TPPO-shaped density was also detected in the minor Ca^{2+}-bound cold

conformation present in the Ca^{2+}/TPPO–TRPM4–37°C dataset, identified through single-subunit

classification (Extended Data Fig. 4). This density was weaker than in the warm conformation

but stronger than in the corresponding Ca^{2+}-bound cold conformation resolved at 18°C (Fig. 3b,

lower panel; Fig. 3d; Extended Data Fig. 7), indicating that TPPO occupancy increases with

temperature even within the same overall protein conformation. Because the backbone

architecture of the binding site remains nearly identical between the two cold structures, this

enhancement likely arises from subtle, temperature-dependent sidechain rearrangements—below

the resolution of our maps (~2.6 Å local)—that fine-tune ligand packing and render the

interaction thermodynamically more favorable.

Together, these results reveal that TPPO selectively engages the warm conformation of

TRPM4 by exploiting temperature- and Ca^{2+}-dependent changes in local binding energetics. This

provides a direct structural basis for the observed three-way synergy, illustrating how intrinsic

modulators reshape the binding pocket to control ligand efficacy at the atomic level.

Mechanistic basis for Ca^{2+}-dependent gating reversal of NC1

Building on our observation that NC1 activates TRPM4 in the absence of Ca^{2+} but loses efficacy

once Ca^{2+} reaches activating levels (Fig. 2a–f), we sought to define the structural basis of this

gating reversal. To this end, we determined cryo-EM structures of TRPM4 in complex with NC1

at 37 °C under Ca^{2+}-free and Ca^{2+}-bound conditions, resolved to 2.6 and 2.8 Å, respectively

(Extended Data Figs. 6,8; Extended Data Table 2).

In the Ca^{2+}-free dataset, a well-defined tripod-shaped density was observed in the S1–S4

domain (~2.6 Å local resolution), occupying the same pocket as the TPPO and deacetyl

bisacodyl binding sites[25], while no Ca^{2+} density appeared at the Ca_{TMD} site, as expected (Fig. 4a).

The shape and dimensions of this density unambiguously matched NC1, which, despite being

larger than TPPO, adopts a comparable three-ring scaffold with its indolinone carbonyl oxygen

positioned analogously to TPPO's phosphoryl oxygen (Figs. 3d, left panel; Fig. 4a).

Consequently, NC1 binds in a similar pose to TPPO, engaging an equivalent network of residues,

including a key polar contact between its indolinone carbonyl oxygen and R1072 on the TRP

helix (Fig. 4a). Consistent with this, alanine substitution of R1072 and other residues within the

binding site abolished or markedly reduced NC1-mediated activation (Fig. 4c; Extended Data

Fig. 3b), confirming that NC1, like TPPO, binds to the S1–S4$_{Upper}$ site.

Interestingly, in the Ca^{2+}-bound dataset, NC1 density at the same S1–S4$_{Upper}$ pocket was

markedly weaker (Fig. 4b). This diminished occupancy aligns with our electrophysiological

findings, where 1 uM Ca^{2+} strongly suppressed the linear, NC1-evoked currents characteristic of

Ca^{2+}-free conditions and restored the canonical Ca^{2+}-dependent gating pattern (Fig. 2a–f). The

weak residual density, likely arising from the oversaturating NC1 concentration used for cryo-

EM rather than genuine co-binding, supports that NC1 and Ca^{2+} binding are mutually

incompatible. Importantly, the antagonism does not result from direct competition for a shared

site but instead from allosteric interference between the NC1 pocket and the Ca$_{TMD}$ site (Fig. 4b).

To understand the basis of this interference, we compared the Ca^{2+}-free NC1–TRPM4 and

Ca^{2+}–TRPM4 complexes. Two local structural changes accompany Ca^{2+} coordination at the

Ca$_{TMD}$. First, Ca^{2+} binding repositions the TRP helix, shifting R1072 away from the indolinone

carbonyl oxygen that is central to NC1 recognition (Fig. 4d). Second, Ca^{2+} binding brings the

positively charged R905 into close proximity with NC1's hydrophobic seven-membered

cycloheptyl ring, creating an electrostatically unfavorable juxtaposition that further destabilizes

ligand binding (Fig. 4d). Together, these antagonistic rearrangements explain why NC1 affinity

drops at increased Ca^{2+} concentration.

In summary, NC1 represents the first example of a TRPM4 activator that is temperature-

independent yet negatively regulated by intrinsic Ca^{2+}. Structural and functional analyses

together uncover a Ca^{2+}-dependent gating reversal, whereby Ca^{2+} reshapes the allosteric network

to displace a previously bound activator. This mechanism contrasts sharply with TPPO, whose

efficacy requires Ca^{2+} synergy at physiological temperature, underscoring the opposing logic by

which intrinsic modulators shape TRPM4 pharmacology.

NC1 overrides Ca^{2+}-dependent ICD–TMD allosteric regulation

The defining feature of NC1 is that it activates TRPM4 without requiring Ca^{2+} at either of the

two regulatory sites (Fig. 2a,d): the agonist site Ca$_{TMD}$ and the temperature-dependent allosteric

site in the intracellular domain (Ca$_{ICD}$). In canonical gating, Ca^{2+} and temperature cooperatively

couple the ICD and TMD to drive activation: Ca^{2+} binding at the temperature-sensitive Ca$_{ICD}$

shifts the channel from the cold to the warm state, establishing the primed conformation in which

Ca^{2+} binding at the Ca_{TMD} opens the pore at physiologically relevant membrane potentials[10]. To

understand how NC1 circumvents this mechanism, we compared three structural states: Ca^{2+}-free

cold, Ca^{2+}-bound warm, and NC1-bound. The transition from Ca^{2+}-free cold to Ca^{2+}-bound warm

involves major ICD rearrangements coupled to a substantial shift in the TRP helix and TMD

(Fig. 4e). By contrast, in the NC1-bound structure the ICD remains in the cold-like, yet the TMD

closely resembles the warm state conformation even in the absence of Ca^{2+} at the Ca_{TMD} (Fig.

4f). These observations suggest that NC1 functionally substitutes for Ca^{2+} at the TMD while

operating independently of Ca^{2+}-dependent priming at the ICD.

Electrophysiology supported this model: NC1 elicited robust currents of comparable

amplitude at both room and physiological temperatures (Fig. 2b,e)—in contrast to Ca^{2+}-

dependent TRPM4 activation, which is strongly enhanced at 37 °C through Ca_{ICD}-mediated

ICD–TMD coupling[10,15]. Thus, NC1 activates the channel without engaging the Ca_{ICD}-driven

allosteric control. To directly test this mechanism, we examined the E396A mutant, which

abolishes Ca_{ICD} binding and locks TRPM4 in the cold-like conformation regardless of

temperature[10]. Remarkably, NC1 still produced strong, temperature-independent activation in

this mutant, confirming that NC1 does not require the ICD–TMD regulation (Extended Data Fig.

3c). This unprecedented mode of action reveals an alternative allosteric pathway within TRPM4

and highlights NC1 as a promising scaffold for therapeutic development, particularly under

conditions where intracellular Ca^{2+} fluctuates.

NBA and CBA inhibit TRPM4 through the S1–S4$_{Lower}$ site

Pharmacological inhibition of TRPM4 is of broad clinical importance, as channel overactivation

contributes to arrhythmogenic cardiac disorders, ischemic injury, and cancer progression[18,21,38].

However, our previous work revealed that the endogenous antagonist ATP markedly loses

potency at physiological temperature because its binding location shifts between the cold and

warm conformations of the channel[10]. This temperature sensitivity limits ATP's ability to

suppress TRPM4 under normal cellular conditions[39].

To identify inhibitors that remain effective within the physiological conformational

landscape, we turned to NBA and its analog CBA[37]. In contrast to ATP and other temperature-

sensitive ligands, NBA and CBA inhibit TRPM4 in a temperature-independent manner,

maintaining high potency at 37°C (Extended Data Fig. 2a). This makes them ideal probes for

uncovering how TRPM4 inhibition can be achieved under physiological conditions through a

temperature-insensitive mechanism.

At 37 °C in the presence of Ca^{2+}, cryo-EM analysis of TRPM4 in complex with NBA or CBA

uncovered a previously unknown ligand binding site within the S1–S4 domain in both datasets,

present in both cold and warm conformations (Extended Data Figs. 6,9). This site lies below the

Ca_{TMD} site and adjacent to the S1–S4$_{Upper}$ pocket that accommodates TPPO and NC1, and is

therefore termed S1–S4$_{Lower}$ site (Fig. 5a,b). The ligand densities were of excellent quality (local

resolution ~2.6 Å) and matched the chemical structures of NBA and CBA, respectively, allowing

unambiguous ligand placement (Extended Data Figs. 6; 10a,b; Extended Data Tables 2, 3). In

both inhibitors, the chlorophenyl-carboxyl group points upward within the S1–S4 domain, with

its carbonyl stabilized by two positively charged residues, R905 on S4 and R1072 on the TRP

helix (Fig. 5a,b).

Surprisingly, this S1–S4$_{Lower}$ site differs from the NBA binding site previously reported based

on lower-resolution cryo-EM maps[40], which placed the inhibitor ~10 Å away on the opposite

face of the S4 helix (Extended Data Fig. 10c)—within the cleft between the S1–S4 domain and

the pore domain (Extended Data Fig. 10d), analogous to the vanilloid pocket in TRPV channels

and the NDNA site in TRPM5[41,42]. Careful inspection of both regions in our high-resolution

maps revealed no evidence supporting that earlier assignment. Instead, in our NBA- and CBA-

bound structures, the vanilloid pocket consistently displayed the same Y-shaped density seen in

apo, TPPO-, and NC1-bound structures, indicating a tightly bound endogenous lipid rather than

inhibitor occupancy (Extended Data Fig. 10e).

To validate the S1–S4$_{Lower}$ pocket functionally, we introduced mutations at key interacting

residues and examined their effects using patch-clamp recordings. Substitution of either R905 or

R1072 by alanine completely abolished inhibition (Fig. 5c,d). Moreover, the second aromatic

group, a chlorophenyl in CBA or the bulkier naphthyl in NBA, extends downward into a cavity

formed between the S4–S5 linker and the TRP helix (Fig. 5a,b). Mutation of Q1061, positioned

near this distal aromatic moiety, also markedly reduced inhibition (Fig. 5c,d).

Together, these results establish the S1–S4$_{Lower}$ pocket as the functional binding site for NBA

and CBA, defining a second pharmacologically active locus within the S1–S4 domain. This

domain thus emerges as a bifunctional pharmacological hub of TRPM4, accommodating

both activators (TPPO, NC1, Ca^{2+}) and inhibitors (NBA, CBA) through spatially segregated

upper and lower sites that converge on shared gating residues such as R1072 on the TRP helix

(Fig. 5e).

NBA and CBA inhibit TRPM4 by locking a pre-open conformation

To elucidate how NBA and CBA inhibit Ca^{2+}-dependent activation, we first examined the

conformational transitions that underlie canonical TRPM4 gating. In the Ca^{2+}-free cold state,

W864 on S3 and H908 on S4 form a stabilizing π–π stacking interaction[43-45]. Upon Ca^{2+} binding,

coordinated engagement of the Ca_{TMD} and Ca_{ICD} triggers rearrangements in the S1–S4 bundle

and TRP helix, repositioning W864 and H908 and thereby disrupting the contact between the

intracellular tips of the S3 and S4 helices (Fig. 5f, the gray structure)—a hallmark of Ca^{2+}-gated

TRPM channel activation[42,46]. This frees the S4–S5 linker, allowing the S5–S6 pore domain to

open[10].

In the Ca^{2+}/NBA- and Ca^{2+}/CBA-bound structures, this transition is blocked. The distal

aromatic moiety of NBA or CBA occupies the position normally adopted by H908 in the

activated state, thereby forcing H908 to remain in its apo-like conformation. As a result, H908,

W864, and the ligand form a π–π–π triple stacking interaction that restrains the S4–S5 linker and

prevents pore opening (Fig. 5f, the purple structure). The overall architecture of the inhibited

state closely matches the Ca^{2+}-bound warm closed conformation (backbone RMSD 0.7 Å),

indicating that NBA and CBA trap TRPM4 in a Ca^{2+}-primed but non-conductive pre-open state.

If this model were correct, NBA and CBA should prevent the transition to the fully open state

normally induced by positive modulators such as DVT[10,47]. To test this, we determined the

structure of TRPM4 in the presence of Ca^{2+}, DVT, and CBA at 37 °C (Extended Data Figs.

6,11a; Extended Data Table 3). The resulting map contained densities only for Ca^{2+} and CBA,

with no evidence of DVT binding, and the protein conformation was identical to the Ca^{2+}/CBA-

bound closed state (Extended Data Fig. 11b). These data provide direct structural evidence that

DVT cannot override NBA/CBA inhibition, therefore supporting our hypothesis.

Interestingly, this mechanism converges with that of ATP, which also prevents opening by

stabilizing the pre-open state, albeit through a distinct intracellular binding site[10]. Unlike ATP,

however, NBA and CBA act from the extracellular side and maintain high potency at

physiological temperature, offering both accessibility and stability within the native

conformational landscape. Thus, NBA and CBA represent promising scaffolds for the design of

TRPM4 inhibitors with therapeutic potential.

Discussion

This study reveals that protein pharmacology is not static but dynamically shaped by the

physiological environment, with temperature and intracellular Ca^{2+} acting as active determinants

of ligand binding, efficacy and mechanism. By integrating single-particle cryo-EM and

electrophysiology, we reveal that these intrinsic factors cooperatively remodel the

conformational landscape of TRPM4, exposing latent binding sites and revealing context-

dependent pharmacological behaviors that are obscured under non-physiological conditions.

Physiological context unmasks the true pharmacological profiles of ligands. The small

molecule TPPO, long regarded as a TRPM5-specific inhibitor, emerges here as a potent activator

of TRPM4, but only when both temperature and intracellular Ca^{2+} reach physiological levels

(Fig. 6a). Conversely, NC1, previously characterized as a Ca^{2+}-independent agonist, undergoes a

functional reversal: at basal Ca^{2+} it activates TRPM4, whereas at elevated Ca^{2+} it becomes

antagonized, revealing an unexpected inhibitory facet of Ca^{2+} regulation (Fig. 6b).

Mechanistically, these contrasting behaviors arise because temperature and Ca^{2+} cooperatively

determine the equilibrium between cold and warm conformations of TRPM4, thereby governing

access to key residues within the S1–S4 regulatory domain. Thus, ligand efficacy in TRPM4 is

not an intrinsic property of the compound alone, but an emergent consequence of its interplay

with the conformational and thermodynamic landscape of the channel.

Moreover, our structural analyses establish the S1–S4 domain as a versatile pharmacological

hub that integrates both environmental and chemical inputs. Within this domain, around the

Ca_{TMD} site, two spatially distinct but functionally coupled sites control opposite gating outcomes:

the S1–S4$_{Upper}$ pocket, occupied by activators TPPO, NC1, and deacetyl bisacodyl[25]; and the S1–

S4$_{Lower}$ pocket, targeted by inhibitors NBA and CBA (Fig. 5e). Both pockets converge on the

TRP helix through the conserved residue like R1072 on the TRP helix, which serves as a central

allosteric node linking environmental cues to gating transitions. Ligand engagement at the upper

pocket promotes activation by coupling between the S1–S4 domain and the pore domain,

whereas ligand binding to the lower pocket disrupts this coupling, stabilizing a non-conductive state. This dual-site organization provides a unified explanation for how temperature, Ca^{2+} and chemically diverse ligands—activators and inhibitors alike—modulate TRPM4 through a shared structural framework.

This regulatory logic uncovered here may represent a general feature of TPRM channels. In TRPM3, both agonists and antagonists occupy the upper pocket[48,49]; and in TRPM8, ligands engage both[50-52]. The S1–S4 domain thus exemplifies an evolutionary design in which temperature and chemical cues dynamically reconfigure a common structural scaffold to yield distinct, context-dependent functional outcomes.

Notably, we found that temperature-dependent ligand recognition in TRPM4 is not limited to regions that undergo large-scale rearrangements. Even sites within structurally stable domains—such as the S1–S4 transmembrane bundle, which exhibits only subtle temperature-driven movements—can undergo marked shifts in ligand accessibility and efficacy. This principle likely extends to many drug targets, highlighting that seemingly rigid structural regions may harbor cryptic, environment-sensitive pharmacology.

More broadly, these findings advocate for an environment-aware framework of pharmacology. Temperature and intracellular Ca^{2+} are not passive background factors but active regulators that reshape ligand binding, potency, and efficacy. The discrepancy between in vitro screening and in in vivo performance may arise from ignoring such physiological modulators, which can redefine both the apparent selectivity and the mechanism of drug action. By revealing how intrinsic cofactors govern the accessibility and stability of ligand-binding pockets, TRPM4 highlights a general principle: pharmacological specificity emerges from the interplay between ligand chemistry and the dynamic structural state of the target protein.

For ubiquitously expressed proteins such as TRPM4, these principles carry important therapeutic implications. By exploiting local physiological changes—such as elevated intracellular Ca^{2+} during stress or altered thermal states in disease—future drugs could be tailored to act selectively in pathological contexts while sparing normal function. Such environment-sensitive design strategies may reduce side effects and offer a new route toward precision pharmacology.

Figure 1. Three-way synergy between TPPO, temperature and Ca²⁺.

a–f, Whole-cell currents measured in tsA cells overexpressing wild-type TRPM4 at 22 °C (upper panels) and 37 °C (lower panels), with free intracellular Ca²⁺ concentrations of 0 µM (**a, d**), 0.1 µM (**b, e**) and 1 µM (**c, f**), in the presence of 50 µM extracellular TPPO. A voltage protocol stepping from −100 mV to +140 mV in 20-mV increments was applied, with each voltage step lasting 100 ms. The currents from all measured cells were then averaged and plotted versus voltage to generate the mean I–V curve, with the line representing the mean and the shaded indicating the s.e.m. The numbers of independently measured cells are as follows: (**a**) n=9 (−TPPO) / 9 (+TPPO), (**b**) n=7 (−TPPO) / 8 (+TPPO), (**c**) n=9 (−TPPO) / 9 (+TPPO), (**d**) n=7 (−TPPO) / 10 (+TPPO), (**e**) n=8 (−TPPO) / 8 (+TPPO), and (**f**) n=6 (−TPPO) / 6 (+TPPO). **g–i**,

TPPO dose-response measurements for wild-type TRPM4 with 0.1 µM free intracellular Ca^{2+} at

37 °C (**g**), 0.1 µM Ca^{2+} at 23 °C (**h**), and 0 µM Ca^{2+} at 37 °C (**i**). A voltage protocol was applied

every 10 s to monitor current changes: –100 mV for 200 ms, followed by +100 mV for 200 ms.

Once the current stabilized at each TPPO concentration, the current amplitudes at +100 mV from

all measured cells were averaged, and the mean amplitudes were plotted and fitted to obtain the

EC_{50}. Current amplitudes were normalized either to the mean at 1000 µM TPPO (**g, i**) or to the

fitted I_max from the Hill equation (**h**). Data are presented as mean ± s.e.m. The numbers of

independently measured cell are as follows: (**g**) n=7 for all concentrations, (**h**) n=5 at 1 µM and

n=7 for all other concentrations, and (**i**) n=7 for all concentrations.

Figure 2. Ca²⁺-dependent gating reversal of NC1.

a–f, Whole-cell currents measured in tsA cells overexpressing wild-type TRPM4 at 22 °C (upper panels) and 37 °C (lower panels), with free intracellular Ca^{2+} concentrations of 0 µM (**a, d**), 0.1 µM (**b, e**) and 1 µM (**c, f**). A voltage protocol was applied every 5 s to monitor current changes until reaching steady state: −100 mV for 50 ms, ramped to +100 mV over 200 ms and held at +100 mV for 50 ms. The currents without and with 1 µM extracellular NC1 in each condition were measured in a single cell. The currents from all measured cells were then averaged and plotted versus voltage to generate the mean I–V curve, with the line representing the mean and the shaded indicating the s.e.m. For clarity, the s.e.m. envelope was plotted in one direction. The numbers of independently measured cells are as follows: n=5 (**a**), n=7 (**b**), n=5 (**c**), n=5 (**d**), n=4

(**e**), and n=4 (**f**). **g–i**, NC1 dose-response measurements for wild-type TRPM4 at

22 °C with 0 μM free intracellular Ca^{2+} (**g**), 0.1 μM Ca^{2+} (**h**), and 1 μM Ca^{2+} (**i**). A voltage

protocol was applied every 5 s to monitor current changes: 0 mV for 50 ms, switch to +80 mV

for 50 ms, followed by −80 mV for 50 ms, and return to 0 mV for 50 ms. Once the current

stabilized at each NC1 concentration, the current amplitudes at +80 mV and −80 mV were

normalized to those at 10 μM NC1. Normalized currents from all measured cells were averaged

and plotted as mean ± s.e.m. EC_{50} values obtained from individual cells were averaged to yield

the final EC_{50} reported as mean ± s.e.m. (**g,h**). Data are presented as mean ± s.e.m. The numbers

of independently measured cells are as follows: n=5 (**g**), n=5 (**h**), and n=4 (**i**).

Figure 3. Structural basis of the three-way synergy between TPPO, temperature and Ca^{2+}.

a–c, Structures of TRPM4 in complex with Ca^{2+}/TPPO at 37 °C (**a**, PDB-ID 9Z1W), Ca^{2+}/TPPO at 18 °C (**b**, PDB-ID 9Z1Z), and EGTA/TPPO at 37 °C (**c**, PDB-ID 9Z2O). One subunit is shown in cartoon representation, while the remaining three subunits are displayed as the cryo-EM map, viewed parallel to the membrane. The transparent envelope outlines the detergent micelle and the disordered region of the C-terminal coiled-coil. TPPO is highlighted in yellow and Ca^{2+} in green. The resolution of the consensus map is indicated below. Bottom panels show the details of the

S1–S4 binding site in the consensus maps. The protein backbone is shown in cartoon

representation, with key interacting residues in stick representation. Densities at the TPPO

binding site (yellow) and at the Ca_{TMD} site (transparent in **a,b**) are contoured at 4.5 σ. **d**, Details

of the S1–S4 binding site in the warm (left, PDB-ID 9Z1X) and cold (right, PDB-ID 9Z1Y)

states, obtained from classification of the Ca^{2+}/TPPO dataset collected at 37 °C. The resolution

of each cryo-EM map is indicated. Densities at the TPPO binding site and the Ca_{TMD} site are

shown as transparent surface representations contoured at 9 σ. The fitted TPPO molecule is

shown in stick representation (only in the warm state), and Ca_{TMD} as a green sphere. **e**, Structural

comparison of TRPM4 with Ca^{2+}/TPPO at 37 °C (salmon, PDB-ID 9Z1X) and the Ca^{2+}-free apo

state (blue, PDB-ID 9B93), superimposed using the S1–S4 (residues 760–910). TPPO is shown

in stick representation with a transparent envelope, and Ca_{TMD} as a green sphere. Black arrows

indicate the approximate direction of movement of key residues and the TRP helix. The dashed

lines mark key interactions.

Figure 4. Mechanistic basis for Ca^{2+}-dependent gating reversal of NC1.

a,b The left panels show the structures of TRPM4 in complex with EGTA/NC1 at 37 °C (**a**, PDB-ID 9Z22) and Ca^{2+}/NC1 at 37 °C (**b**, PDB-ID 9Z21). One subunit is shown in cartoon representation, while the remaining three subunits are displayed as the cryo-EM map, viewed parallel to the membrane. The transparent envelope outlines the detergent micelle and the disordered region of the C-terminal coiled-coil. The middle panels show the details of the S1–S4 binding site, with the protein backbone in cartoon representation and key interacting residues in stick representation. Densities at the NC1 binding site, and at the Ca$_{TMD}$ site (only in **b**) are shown as transparent surface representations contoured at 10 σ. The fitted NC1 molecule in (only in **a**) is shown in stick representation, and Ca$_{TMD}$ as a green sphere (only in **b**). The right panels present the same regions as the middle panels but additionally display densities for well-defined

side chains within the binding site, contoured at 15 σ. **c**, NC1-induced whole-cell current

amplitudes at +100 mV in cells overexpressing wild-type TRPM4 and binding-site mutants,

calculated by subtracting the steady-state current recorded before NC1 application from that

recorded after application. Each point represents an independently measured cell (n = 5 (WT), 5

(N786A), 4 (S789A), 5 (Y790A), 5 (R905A), 4 (R1072A)), with bars indicating the

mean. Statistical analysis was performed using one-way ANOVA with Bonferroni's post hoc test,

comparing each mutant with wild type (*$p<0.05$, ***$p<0.001$). Averaged raw data are shown in

Extended Data Fig. 3b. **d**, Structural comparison of TRPM4 NC1-bound state (cyan, PDB-ID

9Z22) and the Ca^{2+}-bound warm state (blue, PDB-ID 9B8W) of TRPM4, superimposed using the

S1–S4 (residues 760–910). NC1 is shown in stick representation with a transparent envelope,

and Ca_{TMD} as a green sphere. Black arrows indicate the approximate direction of movement of

key residues. The dashed line marks a key interaction. **e**, Structural comparison of a single

subunit of the Ca^{2+}-free cold state (blue, PDB-ID 9B93) and the Ca^{2+}-bound warm state (grey,

PDB-ID 9B8W) of TRPM4, superimposed using the S1–S4 (residues 760–910). MHR3/4 is

outlined with a pink envelope and MHR1/2 with an orange envelope. The right panel shows

details of the S1–S4 binding site and its surroundings. Ca_{TMD} is shown in a green sphere, and the

black arrows indicate the movement of structural elements forming the binding site upon Ca^{2+}

binding. **f**, Structural comparison of the NC1-bound state (cyan, PDB-ID 9Z22) and the Ca^{2+}-

bound warm state (grey, PDB-ID 9B8W) of TRPM4. The structures are superimposed and

displayed similarly to panel (**e**). NC1 is shown in stick representation, and Ca_{TMD} as a green

sphere.

Figure 5. NBA and CBA inhibit TRPM4 through the S1–S4$_{Lower}$ site.

a,b Structure of TRPM4 in complex with Ca^{2+}/CBA at 37 °C (**a**, PDB-ID 9Z23) and Ca^{2+}/NBA at 37 °C (**b**, PDB-ID 9Z25). One subunit is shown in cartoon representation, while the remaining

three subunits are displayed as the cryo-EM map, viewed parallel to the membrane. The transparent envelope outlines the detergent micelle and the disordered region of the C-terminal coiled-coil. CBA and NBA are highlighted in yellow and Ca^{2+} in green. Bottom panels show the details of the S1–S4 binding site, with the protein backbone in cartoon representation and key interacting residues in stick representation. Densities for CBA and Ca_{TMD} (**a**) and for NBA and Ca_{TMD} (**b**) are shown as transparent surface representations contoured at 17 σ and 20 σ, respectively. The fitted CBA and NBA molecules are shown in stick representation, and Ca_{TMD} as a green sphere. **c,d**, CBA- and NBA-dependent inhibition at +100 mV in cells overexpressing wild-type TRPM4 and binding-site mutants. The percentage of remaining currents after the application of NBA or CBA at 22 °C was calculated by dividing the steady-state current recorded after NBA or CBA application by the steady-state current recorded before application. Each point represents an independently measured cell: NBA, n = 6 (WT), 5 (W864A), 5 (R905A), 5 (H908A), 5 (Q1061A), and 6 (R1072A); CBA, n = 7 (WT), 6 (W864A), 5 (R905A), 5 (H908A), 5 (Q1061A), and 4 (R1072A). Bars indicate the mean. Statistical analysis was performed using one-way ANOVA with Bonferroni's post hoc test, comparing each mutant with wild type. (*p<0.05, ***p<0.001). Averaged raw data are shown in Extended Data Fig. 2b. **e**, Structural comparison of the Ca^{2+}/CBA-bound state (blue, 9Z23) and the NC1-bound state (orange, PDB-ID 9Z21) of TRPM4, superimposed using S1–S4 (residues 760–910). CBA and NC1 are shown in stick representation with a transparent envelope, and Ca_{TMD} as a green sphere. The filled orange and blue ellipses mark the upper and lower pockets within the S1–S4 domain, respectively. Arg1072 interacts with both ligands, as indicated by the dashed lines. **f**, Structural comparison of the Ca^{2+}/CBA-bound state (blue, PDB-ID 9Z23) and the Ca^{2+}-bound warm state (grey, PDB-ID 9B8W) of TRPM4, superimposed using S1–S4 (residues 760–910). CBA and the side chains of H908 and W864 are shown as stick representation, forming a triple stacking interaction indicated by the transparent discs and the red dashed line.

Figure 6. Distinct temperature- and Ca²⁺-dependent activation mechanisms of TPPO and NC1 on TRPM4.

a, Schematic representation of the polymodal activation of TRPM4 by TPPO, Ca²⁺, and temperature. Neither resting Ca²⁺ levels nor physiological temperature alone is sufficient to enable TPPO-mediated activation. Only when both resting Ca²⁺ and physiological temperature are present can TPPO exert its activating effect. **b**, Schematic representation of NC1 activity, which functions independently of Ca²⁺ at resting levels but loses efficacy once intracellular Ca²⁺ rises to stimulatory concentrations.

Extended Data Figure 1. Electrophysiological characterization of TPPO-dependent activation of TRPM4.

a–f, Representative whole-cell current traces recorded at room temperature (23 °C) and 37 °C with free intracellular Ca^{2+} concentrations of 0 µM (**a,d**), 0.1 µM (**b,e**), and 1 µM (**c,f**), evoked by a voltage-step protocol from –100 mV to +140 mV in 20-mV increments (100 ms per step). The currents without and with 50 µM extracellular TPPO at each condition were measured from a single cell. **g–i**, Representative whole-cell current traces from the TPPO dose response recordings acquired using a two-step protocol (–100 mV followed by +100 mV for 200 ms each) under 0.1 µM Ca^{2+} at 37 °C (**g**), 0.1 µM Ca^{2+} at 23 °C (**h**), and 0 µM Ca^{2+} at 37 °C (**i**). **j**, Whole-cell currents of non-transfected tsA cells and tsA cells overexpressing wild-type TRPM4 and TPPO binding-site mutants evoked by 1 µM free intracellular Ca^{2+} and measured at +100 and −100 mV at room temperature. Each point represents an independently measured cell: non-transfected tsA201 (n=8), WT (n=10), R1072A (n=10), Y790A (n=7), W820A (n=10), W828A (n=9), S789A (n=12), R905A (n=7), F793A (n=12), F902A (n=6), D868A (n=5), E898A (n=3), F1069A (n=4), and L824A (n=5). Bars indicate the mean. **k**, Mean I–V relationships from tsA cells overexpressing wild-type TRPM4 or the R1072A mutant under the condition of 0.1 µM free intracellular Ca^{2+} and 50 µM extracellular TPPO, measured at 37 °C. Currents were elicited using a voltage-step protocol from –100 mV to +140 mV. Steady-state currents at each voltage were averaged and plotted (WT, n=15; R1072A, n=10). Data are presented as mean ± s.e.m.

Extended Data Figure 2. Electrophysiological characterization of TRPM4 inhibition by NBA and CBA.

a, Left and middle panels show whole-cell currents measured in tsA cells overexpressing wild-type TRPM4 at 22 °C and 37 °C, with 1 μM free intracellular Ca^{2+}. A voltage protocol was applied every 5 s to monitor current changes until reaching steady state: −100 mV for 50 ms, ramped to +100 mV over 200 ms, and held at +100 mV for 50 ms. The currents without and with 10 μM extracellular NBA or CBA at each condition were measured from a single cell. The currents from all measured cells were then averaged and plotted versus voltage to generate the mean I–V curve, with the line representing the mean and the shaded indicating the s.e.m. For clarity, the s.e.m. envelope was plotted in one direction. The numbers of independently measured

cells are as follows: n=6 (NBA, 22 °C), n=5 (NBA, 37 °C), n=7 (CBA, 22 °C), n=5 (CBA,

37 °C). Right panels show quantification of CBA- and NBA-dependent inhibition at +100 mV

for the recordings shown in the left and middle panels. The percentage of remaining currents

after the application of NBA or CBA was calculated by dividing the steady-state current recorded

after NBA or CBA application by the steady-state current recorded before application. Each

point represents an independently measured cell, and bars indicate the mean. A t-test was used to

compare inhibition at 22 °C versus 37 °C. **b**, Whole-cell currents measured in tsA cells

overexpressing wild-type TRPM4 or NBA/CBA binding-site mutants at 22 °C, with 1 µM free

intracellular Ca^{2+}, without and with 10 µM extracellular NBA or CBA. Measurements and

analysis were performed as in panel (**a**). The numbers of independently measured cells are as

follows. NBA: n=6 (WT), n=6 (W864A), n=5 (R905A), n=5 (H908A), n=6 (Q1061A), n=6

(R1072A); CBA: n=7 (WT), n=6 (W864A), n=5 (R905A), n=5 (H908A), n=5 (Q1061A), n=4

(R1072A).

Extended Data Figure 3. Electrophysiological characterization of NC1-dependent activation of TRPM4.

a, The representative current traces for the NC1 dose-response measurements for wild-type TRPM4 at 22 °C with free intracellular Ca^{2+} of 0 µM, 0.1 µM, and 1 µM. A voltage protocol was applied every 5 s to monitor current changes: 0 mV for 50 ms, switch to +80 mV for 50 ms, followed by −80 mV for 50 ms, and return to 0 mV for 50 ms. **b**, Whole-cell currents measured in tsA cells overexpressing wild-type TRPM4 or NC1 binding-site mutants at 37 °C, with 0.1 µM

free intracellular Ca^{2+}, without and with 1 μM extracellular NC1. A voltage protocol was applied

every 5 s to monitor current changes until reaching steady state: −100 mV for 50 ms, ramped to

+100 mV over 200 ms, and held at +100 mV for 50 ms. The currents without and with NC1 at

each condition were measured from a single cell. The currents from all measured cells were then

averaged and plotted versus voltage to generate the mean I–V curve, with the line representing

the mean and the shaded indicating the s.e.m. For clarity, the s.e.m. envelope was plotted in one

direction. The numbers of independently measured cells are as follows: n=5 (WT), n=5 (N786A),

n=4 (S789A), n=5 (Y790A), n=5 (R905A), n=4 (R1072A). **c**, Whole-cell currents were

measured in tsA cells overexpressing the E396A mutant of TRPM4 at 22 °C (upper panels) and

37 °C (lower panels), with intracellular free Ca^{2+} concentrations of 0 μM, 0.1 μM, and 1 μM. A

voltage protocol was applied every 5 s to monitor current changes: −100 mV for 50 ms, ramped

to +100 mV over 200 ms and held at +100 mV for 50 ms. The current without or with 1 μM

extracellular NC1 in each condition was measured in a single cell. The currents from all

measured cells were then averaged and plotted versus voltage to generate the mean I–V curve,

with the line representing the mean and the shaded indicating the s.e.m. For clarity, the s.e.m.

envelope was plotted in one direction. The numbers of independently measured cells are as

follows: n=5 (0 μM Ca^{2+}, 22 °C), n=5 (0.1 μM Ca^{2+}, 22 °C), n=5 (1 μM Ca^{2+}, 22 °C), n=5 (0 μM

Ca^{2+}, 37 °C), n=5 (0.1 μM Ca^{2+}, 37 °C) and n=5 (1 μM Ca^{2+}, 37 °C).

Extended Data Figure 4. Cryo-EM data processing workflow for the Ca²⁺/TPPO–TRPM4–

37 °C dataset. A representative micrograph is shown. The final consensus map and the TMD-

focused map with the best-resolved ligand density are displayed with their respective resolutions.

Extended Data Figure 5. Cryo-EM data processing workflow for the Ca²⁺/TPPO–TRPM4–

18 °C (a) and EGTA/TPPO–TRPM4–37 °C (b) datasets. Representative micrographs are

shown. The final maps are displayed with their respective resolutions.

644 **Extended Data Figure 6. Assessment of cryo-EM data quality.** For each map, left: cryo-EM

645 maps colored by local resolution; right: Fourier shell correlation (FSC) plot.

Extended Data Figure 7. Densities of the TPPO binding site from various datasets or conformations contoured at different σ levels.

Densities at the TPPO binding site are shown as yellow surfaces. For comparison, densities for Ca_{TMD} and surrounding interacting residues are displayed as transparent surfaces. The TPPO molecule in the pose fitted to the Ca^{2+}/TPPO–TRPM4–37 °C warm map is shown on the left in stick representation.

Extended Data Figure 8. Cryo-EM data processing workflow for the Ca²⁺/NC1–TRPM4–37 °C (a) and EGTA/NC1–TRPM4–18 °C (b) datasets. Representative micrographs are shown. The final maps are displayed with their respective resolutions.

Extended Data Figure 9. Cryo-EM data processing workflow for the Ca²⁺/CBA–TRPM4–18 °C (a) and Ca²⁺/CBA–TRPM4–37 °C (b) datasets. Representative micrographs are shown. The final consensus map and the TMD-focused map with the best-resolved ligand density are displayed with their respective resolutions.

Extended Data Figure 10. Comparison of the NBA and CBA sites identified in this study with the previously reported NBA site, along with the densities observed within these sites.

a,b, NBA and CBA densities contoured at different σ levels for the Ca^{2+}/NBA–TRPM4–37 °C

TMD map and the Ca^{2+}/CBA–TRPM4–37 °C TMD map in this study. Densities of NBA and

CBA are shown as yellow surfaces. For comparison, densities for Ca_{TMD} and surrounding

interacting residues are displayed as transparent surfaces. The NBA and CBA molecules in the

poses fitted to the maps are shown on the left in stick representation. **c**, Location of the NBA

binding site identified in this work (yellow; PDB-ID 9Z24) compared with the previously

reported site (light blue; PDB-ID 8RD9). **d**, The overall cryo-EM map (top) and the densities of

the putative NBA binding site (bottom) from the previously published TRPM4 structure in the

presence of NBA (EMD-19069), contoured at different σ levels. The putative NBA densities are

highlighted in yellow, and the protein densities are shown in gray. The NBA molecule in the pose

fitted to the map is shown on the left in stick representation. **e**, Densities at the previously

reported NBA binding site in our high-resolution cryo-EM maps across different ligand

conditions consistently reveal a Y-shaped density characteristic of a lipid molecule. The Y-shaped

densities are highlighted in yellow, and the protein densities are shown in gray.

Extended Data Figure 11. Cryo-EM data processing workflow for the Ca²⁺/DVT/CBA–

TRPM4–18 °C dataset. a, A representative micrograph is shown. The final map is displayed

with the resolution. **b**, Left: superimposition of the Ca²⁺/DVT/CBA–TRPM4 map and the

Ca²⁺/DVT–TRPM4$_{Warm}$ model (red; open state), focusing on the transmembrane domain, viewed

from the intracellular side. Middle: Superimposition of the Ca²⁺/DVT/CBA–TRPM4 map and

the Ca²⁺/CBA–TRPM4 model (purple), focusing on the transmembrane domain, viewed from the

intracellular side. Right: superimposition of the cryo-EM maps of Ca²⁺/DVT/–TRPM4$_{Warm}$

(transparent pink) and Ca²⁺/DVT/CBA–TRPM4 (solid purple), focusing on the DVT$_{Warm}$ site and

its surroundings.

	Ca^{2+}/TPPO–TRPM4–37°C (EMDB-73754) (PDB 9Z1W)	Ca^{2+}/TPPO–TRPM4–37°C warm TMD (EMDB-73755) (PDB 9Z1X)	Ca^{2+}/TPPO–TRPM4–37°C cold TMD (EMDB-73756) (PDB 9Z1Y)	Ca^{2+}/TPPO–TRPM4–18°C (EMDB-73757) (PDB 9Z1Z)	EGTA/TPPO–TRPM4–37°C (EMDB-73758) (PDB 9Z20)
Data collection and processing					
Magnification	135000	135000	135000	135000	135000
Voltage (kV)	200	200	200	200	200
Electron exposure (e–/Å^2)	40	40	40	40	40
Defocus range (μm)	-0.5 – -1.4	-0.5 – -1.4	-0.5 – -1.4	-0.5 – -1.4	-0.5 – -1.4
Pixel size (Å)	0.87	0.87	0.87	0.87	0.874
Symmetry imposed	C4	C1	C1	C1	C4
Initial particle images (no.)	2.6M			2.7M	4.9M
Final particle images (no.)	1180K	882K	294K	760K	1929K
Map resolution (Å)	2.9	3.0	3.3	2.6	2.5
FSC threshold	0.143	0.143	0.143	0.143	0.143
Map resolution range (Å)	2.9 – 246.2	3.0 – 246.2	3.2 – 246.2	2.6 – 246.2	2.5 – 246.2
Refinement					
Initial model used (PDB code)	5WP6	5WP6	5WP6	5WP6	5WP6
Model resolution (Å)	3.3	3.3	3.5	3.1	2.7
FSC threshold	0.5	0.5	0.5	0.5	0.5
Model resolution range (Å)	3.3 – 246.2	3.3 – 246.2	3.5 – 246.2	3.1 – 246.2	2.7 – 246.2
Map sharpening B factor (Å^2)	-120	-30	-30	-100	-100
Model composition					
Non-hydrogen atoms	30220	2912	2912	29272	29312
Protein residues	3976	365	365	3892	3896
Ligands	12	2	2	12	0
B factors (Å^2)					
Protein	133.28	83.39	82.41	123.92	103.06
Ligand	144.31	98.78	95.88	78.23	
R.m.s. deviations					
Bond lengths (Å)	0.004	0.003	0.002	0.006	0.003
Bond angles (°)	0.469	0.448	0.419	0.493	0.452
Validation					
MolProbity score	1.21	1.13	1.10	1.14	0.92
Clashscore	1.89	2.41	2.24	1.69	0.97
Poor rotamers (%)	0.68	0	0	0.5	1.13
Ramachandran plot					
Favored (%)	96.15	97.49	97.49	96.63	97.58
Allowed (%)	3.85	2.51	2.51	3.37	2.42
Disallowed (%)	0	0	0	0	0

696 **Extended Data Table 1: Cryo-EM data collection, refinement and validation statistics.** Data

697 are shown for Ca^{2+}/TPPO–TRPM4–37°C, Ca^{2+}/TPPO–TRPM4–37°C warm TMD, Ca^{2+}/TPPO–

698 TRPM4–37°C cold TMD, Ca^{2+}/TPPO–TRPM4–18°C, and EGTA/TPPO–TRPM4–37°C.

	Ca²⁺/NC1–TRPM4–37°C (EMDB-73759) (PDB 9Z21)	EGTA/NC1–TRPM4–37°C (EMDB-73760) (PDB 9Z22)	Ca²⁺/CBA–TRPM4–37°C (EMDB-73761) (PDB 9Z23)	Ca²⁺/CBA–TRPM4–37°C TMD (EMDB-73762) (PDB 9Z24)	Ca²⁺/NBA–TRPM4–37°C (EMDB-73763) (PDB 9Z25)
Data collection and processing					
Magnification	135000	135000	135000	135000	135000
Voltage (kV)	200	200	200	200	200
Electron exposure (e–/Å²)	40	40	40	40	40
Defocus range (μm)	-0.5 – -1.4	-0.5 – -1.4	-0.5 – -1.4	-0.5 – -1.4	-0.5 – -1.4
Pixel size (Å)	0.874	0.87	0.87	0.87	0.874
Symmetry imposed	C4	C4	C4	C1	C4
Initial particle images (no.)	5.6M	4.2M	3.3M		6.2M
Final particle images (no.)	1636K	1475K	1290K	160K	2049K
Map resolution (Å)	2.8	2.6	2.6	3.1	2.6
FSC threshold	0.143	0.143	0.143	0.143	0.143
Map resolution range (Å)	2.8 – 246.2	2.6 – 246.2	2.6 – 246.2	3.1 – 246.2	2.6 – 246.2
Refinement					
Initial model used (PDB code)	5WP6	5WP6	5WP6	5WP6	5WP6
Model resolution (Å)	3.1	2.9	3.2	3.3	2.9
FSC threshold	0.5	0.5	0.5	0.5	0.5
Model resolution range (Å)	3.1 – 246.2	2.9 – 246.2	3.2 – 246.2	3.3 – 246.2	2.9 – 246.2
Map sharpening B factor (Å²)	-117	-105	-102	-50	-106
Model composition					
Non-hydrogen atoms	30300	29396	30304	2740	30300
Protein residues	3976	3900	3972	338	3972
Ligands	12	4	12	2	12
B factors (Å²)					
Protein	107.36	100.17	108.2	101.63	114.79
Ligand	117.04	83.25	66.02	116.92	82.15
R.m.s. deviations					
Bond lengths (Å)	0.004	0.004	0.004	0.004	0.003
Bond angles (°)	0.488	0.439	0.460	0.468	0.440
Validation					
MolProbity score	1.36	1.44	1.22	1.2	1.34
Clashscore	2.14	4.57	1.6	2.19	2.35
Poor rotamers (%)	0.27	0.14	1.08	0.35	0.27
Ramachandran plot					
Favored (%)	94.62	96.74	95.79	96.69	95.28
Allowed (%)	5.38	3.26	4.21	3.31	4.72
Disallowed (%)	0	0	0	0	0

700 **Extended Data Table 1: Cryo-EM data collection, refinement and validation statistics.** Data

701 are shown for Ca²⁺/NC1–TRPM4–37°C, EGTA/NC1–TRPM4–37°C, Ca²⁺/CBA–TRPM4–37°C,

702 and Ca²⁺/NBA–TRPM4–37°C.

	Ca²⁺/NBA–TRPM4–37°C TMD (EMDB-73764) (PDB 9Z26)	Ca²⁺/CBA/DVT–TRPM4–37°C (EMDB-73765) (PDB 9Z27)
Data collection and processing		
Magnification	135000	135000
Voltage (kV)	200	200
Electron exposure (e–/Å²)	40	40
Defocus range (μm)	-0.5 – -1.4	-0.5 – -1.4
Pixel size (Å)	0.874	0.87
Symmetry imposed	C1	C4
Initial particle images (no.)		2M
Final particle images (no.)	239K	278K
Map resolution (Å)	2.9	2.8
FSC threshold	0.143	0.143
Map resolution range (Å)	2.9 – 246.2	3.8 – 246.2
Refinement		
Initial model used (PDB code)	5WP6	5WP6
Model resolution (Å)	3.2	3.2
FSC threshold	0.5	0.5
Model resolution range (Å)	3.2 – 246.2	3.2 – 246.2
Map sharpening B factor (Å²)	-50	-95
Model composition		
Non-hydrogen atoms	2945	30292
Protein residues	364	3972
Ligands	2	12
B factors (Å²)		
Protein	110.32	191.28
Ligand	125.37	118.86
R.m.s. deviations		
Bond lengths (Å)	0.004	0.002
Bond angles (°)	0.499	0.466
Validation		
MolProbity score	1.11	1.34
Clashscore	2.72	3.37
Poor rotamers (%)	0.97	0.54
Ramachandran plot		
Favored (%)	97.77	96.64
Allowed (%)	2.23	3.36
Disallowed (%)	0	0

704 **Extended Data Table 1: Cryo-EM data collection, refinement and validation statistics.** Data

705 are shown for Ca²⁺/NBA–TRPM4–37°C TMD, and Ca²⁺/DVT/CBA–TRPM4–37°C.

Data Availability

Cryo-EM density maps have been deposited at the EMDB (Electron Microscopy Data Bank) and the Research Collaboratory for Structural Bioinformatics Protein Data Bank (RCS-PDB), respectively.

Acknowledgements

We thank for the support with data collection at the Structural Biology Facility at Northwestern University. We appreciate the Structural Biology Facility and Northwestern IT Research Computing and Data Services for computational support. W.L. is supported by National Institutes of Health (NIH) grants (R01HL153219 and R01NS112363, and R35GM138321). J.D. is supported by a McKnight Scholar Award, a Klingenstein-Simon Scholar Award, a Sloan Research Fellowship in neuroscience, a Pew Scholar in the Biomedical Sciences award, and NIH grants (R01NS111031 and R01NS129804). J.H. is supported by American Heart Association postdoctoral fellowship (24POST1196982).

Author Contributions

J.D. and W.L. supervised the project. J.H. carried out mutagenesis, protein purification, cryo-EM data collection and processing, and structural analysis for all TRPM4 structures included in the study. S.I. and J.C. contributed to the initial cryo-EM studies of TRPM4 in complex with NBA, CBA, and NC1. S.P. and J.L. performed functional experiments. G.O. and J.S. carried out mutagenesis and expression test. J.H., J.D. and W.L. wrote the manuscript, with contributions from S.P. and J.L. All authors participated in manuscript preparation and approved the final version. The authors declare no conflicts of interest.

Methods

Human TRPM4 protein expression and purification

The gene encoding human full-length TRPM4 (UniProtKB: Q8TD43) was subcloned into pEG BacMam vector with a 2×Strep tag, GFP and a thrombin-cleavage site at the N terminus[53]. Bacmid and baculovirus of TRPM4 in a BacMam vector were generated, and P2 viruses were used to infect a suspension of tsA cells. Cells were incubated at 37 °C for 8 h and10 mM sodium butyrate was added to the culture and the temperature was changed to 30 °C. The cells were collected 72 h after infection and resuspended in a buffer containing 100 mM Tris pH 8.0 and 150 mM NaCl (TBS buffer) in the presence of 1 mM phenylmethylsulphonyl fluoride, 0.8 μM aprotinin, 2 μg ml^{-1} leupeptin and 2 mM pepstatin A. The cells were lysed by sonication and the membrane fraction was collected by centrifugation at 186,000g using a 45 Ti rotor (Beckman Coulter) for 1 h at 4 °C. The membrane then homogenized with a Dounce homogenizer in TBS buffer supplemented with protease inhibitors. The protein was extracted from the membrane with TBS buffer supplemented with 1% GDN and protease inhibitors for 3 h at 4 °C. The solubilized proteins were loaded to Strep-Tactin resin. After washing with TBS buffer supplemented with 0.02% GDN, TRPM4 was eluted with the same buffer, supplemented with 10 mM desthiobiotin. The GFP tag were cleaved, and proteins were concentrated and further purified by size-exclusion chromatography (Superose 6). The peak fractions containing the TRPM4 were pooled and concentrated to 8 mg ml^{-1}.

EM sample preparation and data acquisition

Purified TRPM4 was incubated with 5 mM calcium chloride or 5 mM EGTA, as required for each experiment, for 30 s at 18 or 37 °C. Then 0.1mM TPPO, 0.25mM CBA/NBA or 0.8mM NC1 was added and further incubation for 2 min at 18 or 37 °C respectively. 2.5 μl sample was applied to a glow-discharged Quantifoil holey carbon grid (gold, 2/1 μm size/hole space, 300 mesh). The grids were blotted for 1.5 s in the Vitrobot Mark III set to 100% humidity and 37 °C with a 15 s wait time before being plunge-frozen into liquid ethane cooled by liquid nitrogen.

For the all the sample, images were recorded using the FEI Glacios electron microscope at 200 kV and a nominal magnification of 135,000×. A Falcon 4i direct electron detector with

756   Selectris energy filter was used resulting a pixel size of 0.87 Å or 0.874 Å. EPU was used for

757   automated acquisition. Nominal defocus ranged from −0.5 to −1.4 μm.

Cryo-electron microscopy data analysis procedure

The detailed workflow for the data-processing procedure is summarized in Extended Data

Figs. 4–5, 8, 9, 11. In general, the raw movies for each dataset were motion-corrected using

MotionCor2 (v.1.1.0)[54]. The per-micrograph defocus values were estimated using ctffind

(v.4.1.10)[55]. Particle picking was performed using relion template picker[56] and topaz (v.0.2.4)[57].

Junk particles were removed by rounds of 3D heterogeneous refinement using CryoSPARC[58].

Good particles were selected for non-uniformed refinement with C_4 symmetry in CryoSPARC to

generate a 3D map. Multiple rounds of CTF refinement and Bayesian polishing were performed

in RELION and CryoSPARC to further improve the map resolution.

For 37 °C with Ca^{2+} and TPPO dataset, symmetry expansion at the single subunit of ICD level

was done from the map refined with C_4 symmetry, followed by monomer ICD subtraction. The

subtracted images of the monomer ICD underwent 3D classification without image alignment in

RELION to identify the warm and cold conformation. The correspond warm and cold monomer

were subsequently undergone similar TMD monomer 3D classification to identify the best the

local resolution map with the bound ligand.

For the rest of 37 °C with Ca^{2+} dataset, symmetry expansion at the single subunit of TMD level

was done from the map refined with C_4 symmetry, followed by monomer TMD subtraction. The

subtracted images of the monomer TMD underwent 3D classification without image alignment

in RELION to identify the class with the best local resolution map with the bound ligand.

Map resolution estimates were based on the gold standard Fourier shell correlation 0.143

criterion for all datasets.

Model building

Atomic models were generated by rigid-body fitting of the TMD, MHR1/2 and MHR3/4 and C-

terminal domains from a published human TRPM4 model (PDB: 5WP6) into the final cryo-EM

maps. Ligands were fitted into the density through real-space refinement using COOT[59]. The CIF

file of ligands was generated using grade web server (Global Phasing Ltd.). The models were

then manually adjusted in COOT and subjected to phenix.real_space_refine[60] to improve the

model metrics. The final models were validated using phenix.molprobity[61]. Figures were

generated using UCSF ChimeraX[62].

Electrophysiology

TsA201 cells expressing plasmids encoding N-terminal GFP-tagged human TRPM4 WT and

mutants were used. One day after transfection with plasmid DNA (100 ng ml^{-1}) and

Lipofectamine 2000 (Invitrogen, 11668019), the cells were trypsinized and replated onto poly-L-

lysine-coated (Sigma, P4707) glass coverslips. After cell attachment, the coverslip was

transferred to a recording chamber. Whole-cell patch-clamp recordings were performed at room

temperature (21–23 °C) or body temperature (36–38 °C). The temperature of perfusion solutions

was controlled by thermal control devices (SC-20/CL-100, Warner Instruments).

For whole-cell recordings, glass pipettes were pulled to 3–5 MΩ and filled with an internal

solution. Signals were amplified with a MultiClamp 700B amplifier (controlled by MultiClamp

Commander v2.2.2.2) and digitized using a Digidata 1550B A/D converter controlled by

Clampex v11.3 (Molecular Devices). Whole-cell currents were measured in cells with an access

resistance of < 10 MΩ after achieving whole-cell configuration. Whole-cell capacitance

was compensated by the amplifier circuitry. Whole-cell currents were recorded using a two-step

voltage protocol in which cells were held at –100 mV for 200 ms and then stepped to +100 mV

for another 200 ms, repeated every 10 s. An alternative voltage-step protocol was applied, in

which cells were stepped from –100 mV to +140 mV in 20-mV increments, with each step

lasting 100 ms. A voltage ramp from –100 to +100 mV (200 ms duration) was applied every 5 s

from a holding potential of 0 mV. Data were sampled at 10 kHz and low-pass filtered at 2 kHz.

Recordings were analyzed using Clampfit v11.3 (Axon Instruments), GraphPad Prism

v10.6.1 (GraphPad Software, Inc., La Jolla, CA), and OriginPro 2024 (OriginLab, Northampton,

MA). Data were also analyzed using custom Python scripts, including pyABF for loading and

processing ABF electrophysiology files. Electrophysiology files were parsed and analyzed in

Python v3.11.8 executed in Spyder v6.0.7 (Anaconda environment) on macOS v26.0.1 (ARM

architecture). Data processing used pandas v2.2.2 and numerical operations used NumPy

v1.26.4. All custom analysis scripts used in this study are available

at https://github.com/junuk861/ephys-analysis-scripts. The bath solution (pH 7.4) contained 150

mM NaCl, 3 mM KCl, 10 mM HEPES, 2 mM $CaCl_2$, 1 mM $MgCl_2$, and 12 mM mannitol. Patch

pipettes were filled with internal solutions containing 150 mM NaCl, 1 mM $MgCl_2$, 10 mM

HEPES, 5 mM EGTA, and $CaCl_2$ adjusted (0 or 1.69 mM) to yield 0 or 0.1 μM free Ca^{2+},

respectively (pH 7.3,

22 °C) (source: https://somapp.ucdmc.ucdavis.edu/pharmacology/bers/maxchelator/CaEGTA-

TS.htm). For whole-cell recordings of NC1-induced and NBA/CBA-inhibited currents, the

extracellular solution contained the following (in mM): 150 NaCl, 10 HEPES and 1 $MgCl_2$. The

intracellular solution contained (in mM): 150 NaCl, 10 HEPES and 5 EGTA. The intracellular

concentration of $CaCl_2$ was set to either 0, 1.69 or 4.45 mM (corresponding to 0, 0.1 or 1 μM of

free Ca^{2+}). The pH was adjusted to 7.4 using NaOH in both the extracellular and intracellular

solutions.

TPPO (Sigma-Aldrich, T84603) was prepared daily as a 70 mM stock solution in ethanol and

diluted to the desired final concentration in the bath solution immediately before use. NC1

(Necrocide 1, MCE, HY-14307), NBA (4-chloro-2-(2-(naphthalen-1-yloxy)acetamido)benzoic

acid, MCE, HY-128172), and CBA (4-chloro-2-[2-(2-chloro-phenoxy)-acetylamino]-benzoic

acid, Tocris, 6724) were dissolved in DMSO to a stock concentration of 10, 100, and 50 mM,

respectively.

References

1   Gao, Y., Cao, E., Julius, D. & Cheng, Y. TRPV1 structures in nanodiscs reveal
mechanisms of ligand and lipid action. *Nature* **534**, 347-351 (2016).
https://doi.org/10.1038/nature17964
2   Yang, X. *et al.* Structure deformation and curvature sensing of PIEZO1 in lipid
membranes. *Nature* **604**, 377-383 (2022). https://doi.org/10.1038/s41586-022-
04574-8
3   Yao, X., Fan, X. & Yan, N. Cryo-EM analysis of a membrane protein embedded in the
liposome. *Proc Natl Acad Sci U S A* **117**, 18497-18503 (2020).
https://doi.org/10.1073/pnas.2009385117
4   Tao, X., Zhao, C. & MacKinnon, R. Membrane protein isolation and structure
determination in cell-derived membrane vesicles. *Proc Natl Acad Sci U S A* **120**,
e2302325120 (2023). https://doi.org/10.1073/pnas.2302325120

5   Zhou, J. *et al*. Resolving native GABA(A) receptor structures from the human brain.
Nature **638**, 562-568 (2025). https://doi.org/10.1038/s41586-024-08454-1

6   Lin, S., Ke, M., Zhang, Y., Yan, Z. & Wu, J. Structure of a mammalian sperm cation
channel complex. *Nature* **595**, 746-750 (2021). https://doi.org/10.1038/s41586-021-
03742-6

7   Zhang, M. *et al*. Assembly and architecture of endogenous NMDA receptors in adult
cerebral cortex and hippocampus. *Cell* **188**, 1198-1207 e1113 (2025).
https://doi.org/10.1016/j.cell.2025.01.004

8   Jeong, H. *et al*. Structures of the TMC-1 complex illuminate mechanosensory
transduction. *Nature* **610**, 796-803 (2022). https://doi.org/10.1038/s41586-022-
05314-8

9   Kumar Mondal, A., Carrillo, E., Jayaraman, V. & Twomey, E. C. Glutamate gating of
AMPA-subtype iGluRs at physiological temperatures. *Nature* (2025).
https://doi.org/10.1038/s41586-025-08770-0

10   Hu, J. *et al*. Physiological temperature drives TRPM4 ligand recognition and gating.
Nature **630**, 509-515 (2024). https://doi.org/10.1038/s41586-024-07436-7

11   Bootman, M. D. & Bultynck, G. Fundamentals of Cellular Calcium Signaling: A
Primer. *Cold Spring Harb Perspect Biol* **12** (2020).
https://doi.org/10.1101/cshperspect.a038802

12   Zhang, W. H. *et al*. Association of concomitant H1 antihistamine and immune
checkpoint inhibitor therapy on survival outcome and safety in patients with
advanced primary lung cancer: a cohort study. *Transl Lung Cancer Res* **13**, 2787-
2801 (2024). https://doi.org/10.21037/tlcr-24-795

13   Launay, P. *et al*. TRPM4 is a Ca2+-activated nonselective cation channel mediating
cell membrane depolarization. *Cell* **109**, 397-407 (2002).
https://doi.org/10.1016/s0092-8674(02)00719-5

14   Nilius, B. *et al*. Voltage dependence of the Ca2+-activated cation channel TRPM4. *J
Biol Chem* **278**, 30813-30820 (2003). https://doi.org/10.1074/jbc.M305127200

15   Talavera, K. *et al*. Heat activation of TRPM5 underlies thermal sensitivity of sweet
taste. *Nature* **438**, 1022-1025 (2005). https://doi.org/10.1038/nature04248

16   Cheng, H. *et al*. TRPM4 controls insulin secretion in pancreatic beta-cells. *Cell
Calcium* **41**, 51-61 (2007). https://doi.org/10.1016/j.ceca.2006.04.032

17   Pironet, A., Vandewiele, F. & Vennekens, R. Exploring the role of TRPM4 in calcium-
dependent triggered activity and cardiac arrhythmias. *J Physiol* **602**, 1605-1621
(2024). https://doi.org/10.1113/JP283831

18   Vandewiele, F. *et al*. TRPM4 inhibition by meclofenamate suppresses Ca2+-
dependent triggered arrhythmias. *Eur Heart J* **43**, 4195-4207 (2022).
https://doi.org/10.1093/eurheartj/ehac354

19   Yan, J., Bengtson, C. P., Buchthal, B., Hagenston, A. M. & Bading, H. Coupling of
NMDA receptors and TRPM4 guides discovery of unconventional neuroprotectants.
Science **370** (2020). https://doi.org/10.1126/science.aay3302

20   Menigoz, A. *et al*. TRPM4-dependent post-synaptic depolarization is essential for
the induction of NMDA receptor-dependent LTP in CA1 hippocampal neurons.
Pflugers Arch **468**, 593-607 (2016). https://doi.org/10.1007/s00424-015-1764-7

Schattling, B. *et al*. TRPM4 cation channel mediates axonal and neuronal degeneration in experimental autoimmune encephalomyelitis and multiple sclerosis. *Nat Med* **18**, 1805-1811 (2012). https://doi.org/10.1038/nm.3015

Shimizu, T. *et al*. TRPM4 regulates migration of mast cells in mice. *Cell Calcium* **45**, 226-232 (2009). https://doi.org/10.1016/j.ceca.2008.10.005

Barbet, G. *et al*. The calcium-activated nonselective cation channel TRPM4 is essential for the migration but not the maturation of dendritic cells. *Nat Immunol* **9**, 1148-1156 (2008). https://doi.org/10.1038/ni.1648

Guinamard, R., Demion, M., Chatelier, A. & Bois, P. Calcium-activated nonselective cation channels in mammalian cardiomyocytes. *Trends Cardiovasc Med* **16**, 245-250 (2006). https://doi.org/10.1016/j.tcm.2006.04.007

Liu, Y. *et al*. Noncanonical calcium-independent TRPM4 activation governs intestinal fluid homeostasis. *Nat Commun* (2026). https://doi.org/10.1038/s41467-025-68014-7

Dai, Y., Wang, H., Wang, W. & Zhuang, X. Multidimensional pan cancer analysis of the sodium induced cell death gene TRPM4. *Sci Rep* **15**, 29768 (2025). https://doi.org/10.1038/s41598-025-15082-w

Guo, J. *et al*. TRPM4 channels contribute to Adriamycin chemoresistance in breast cancer cells. *Eur J Pharmacol* **998**, 177637 (2025). https://doi.org/10.1016/j.ejphar.2025.177637

Kappel, S. *et al*. TRPM4 is highly expressed in human colorectal tumor buds and contributes to proliferation, cell cycle, and invasion of colorectal cancer cells. *Mol Oncol* **13**, 2393-2405 (2019). https://doi.org/10.1002/1878-0261.12566

Sagredo, A. I. *et al*. TRPM4 regulates Akt/GSK3-beta activity and enhances beta-catenin signaling and cell proliferation in prostate cancer cells. *Mol Oncol* **12**, 151-165 (2018). https://doi.org/10.1002/1878-0261.12100

Gualandi, F. *et al*. Mutation Load of Multiple Ion Channel Gene Mutations in Brugada Syndrome. *Cardiology* **137**, 256-260 (2017). https://doi.org/10.1159/000471792

Saito, Y. *et al*. TRPM4 Mutation in Patients With Ventricular Noncompaction and Cardiac Conduction Disease. *Circ Genom Precis Med* **11**, e002103 (2018). https://doi.org/10.1161/CIRCGEN.118.002103

Hof, T. *et al*. TRPM4 non-selective cation channel variants in long QT syndrome. *BMC Med Genet* **18**, 31 (2017). https://doi.org/10.1186/s12881-017-0397-4

Liu, H. *et al*. Molecular genetics and functional anomalies in a series of 248 Brugada cases with 11 mutations in the TRPM4 channel. *PLoS One* **8**, e54131 (2013). https://doi.org/10.1371/journal.pone.0054131

Liu, H. *et al*. Gain-of-function mutations in TRPM4 cause autosomal dominant isolated cardiac conduction disease. *Circ Cardiovasc Genet* **3**, 374-385 (2010). https://doi.org/10.1161/CIRCGENETICS.109.930867

Palmer, R. K. *et al*. Triphenylphosphine oxide is a potent and selective inhibitor of the transient receptor potential melastatin-5 ion channel. *Assay Drug Dev Technol* **8**, 703-713 (2010). https://doi.org/10.1089/adt.2010.0334

36   Fu, W. *et al*. Persistent activation of TRPM4 triggers necrotic cell death characterized
by sodium overload. *Nat Chem Biol* **21**, 1238-1249 (2025).
https://doi.org/10.1038/s41589-025-01841-3

37   Ozhathil, L. C. *et al*. Identification of potent and selective small molecule inhibitors
of the cation channel TRPM4. *Br J Pharmacol* **175**, 2504-2519 (2018).
https://doi.org/10.1111/bph.14220

38   Bochen, F. *et al*. TRPM4 contributes to cell death in prostate cancer tumor
spheroids, and to extravasation and metastasis in a zebrafish xenograft model
system. *Mol Oncol* **19**, 1299-1309 (2025). https://doi.org/10.1002/1878-0261.13795

39   Nilius, B., Prenen, J., Voets, T. & Droogmans, G. Intracellular nucleotides and
polyamines inhibit the Ca2+-activated cation channel TRPM4b. *Pflugers Arch* **448**,
70-75 (2004). https://doi.org/10.1007/s00424-003-1221-x

40   Ekundayo, B. *et al*. Identification of a binding site for small molecule inhibitors
targeting human TRPM4. *Nat Commun* **16**, 833 (2025).
https://doi.org/10.1038/s41467-025-56131-2

41   Cao, E., Liao, M., Cheng, Y. & Julius, D. TRPV1 structures in distinct conformations
reveal activation mechanisms. *Nature* **504**, 113-118 (2013).
https://doi.org/10.1038/nature12823

42   Ruan, Z. *et al*. Structures of the TRPM5 channel elucidate mechanisms of activation
and inhibition. *Nat Struct Mol Biol* **28**, 604-613 (2021).
https://doi.org/10.1038/s41594-021-00607-4

43   Guo, J. *et al*. Structures of the calcium-activated, non-selective cation channel
TRPM4. *Nature* **552**, 205-209 (2017). https://doi.org/10.1038/nature24997

44   Autzen, H. E. *et al*. Structure of the human TRPM4 ion channel in a lipid nanodisc.
Science **359**, 228-232 (2018). https://doi.org/10.1126/science.aar4510

45   Winkler, P. A., Huang, Y., Sun, W., Du, J. & Lu, W. Electron cryo-microscopy structure
of a human TRPM4 channel. *Nature* **552**, 200-204 (2017).
https://doi.org/10.1038/nature24674

46   Huang, Y., Kumar, S., Lee, J., Lu, W. & Du, J. Coupling enzymatic activity and gating in
an ancient TRPM chanzyme and its molecular evolution. *Nat Struct Mol Biol* **31**,
1509-1521 (2024). https://doi.org/10.1038/s41594-024-01316-4

47   Nilius, B., Prenen, J., Janssens, A., Voets, T. & Droogmans, G. Decavanadate
modulates gating of TRPM4 cation channels. *J Physiol* **560**, 753-765 (2004).
https://doi.org/10.1113/jphysiol.2004.070839

48   Yin, Y. *et al*. Molecular basis of neurosteroid and anticonvulsant regulation of
TRPM3. *Nat Struct Mol Biol* **32**, 828-840 (2025). https://doi.org/10.1038/s41594-024-
01463-8

49   Kumar, S. *et al*. Structural basis for agonist and heat activation of nociceptor TRPM3.
Nat Struct Mol Biol (2025). https://doi.org/10.1038/s41594-025-01692-5

50   Diver, M. M., Cheng, Y. & Julius, D. Structural insights into TRPM8 inhibition and
desensitization. *Science* **365**, 1434-1440 (2019).
https://doi.org/10.1126/science.aax6672

51   Yin, Y. *et al*. Structural basis of cooling agent and lipid sensing by the cold-activated
TRPM8 channel. *Science* **363** (2019). https://doi.org/10.1126/science.aav9334

52   Zhao, C. *et al*. Structures of a mammalian TRPM8 in closed state. *Nat Commun* **13**,
3113 (2022). https://doi.org/10.1038/s41467-022-30919-y
53   Goehring, A. *et al*. Screening and large-scale expression of membrane proteins in
mammalian cells for structural studies. *Nat Protoc* **9**, 2574-2585 (2014).
https://doi.org/10.1038/nprot.2014.173
54   Zheng, S. Q. *et al*. MotionCor2: anisotropic correction of beam-induced motion for
improved cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017).
https://doi.org/10.1038/nmeth.4193
55   Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from
electron micrographs. *J Struct Biol* **192**, 216-221 (2015).
https://doi.org/10.1016/j.jsb.2015.08.008
56   Scheres, S. H. RELION: implementation of a Bayesian approach to cryo-EM
structure determination. *J Struct Biol* **180**, 519-530 (2012).
https://doi.org/10.1016/j.jsb.2012.09.006
57   Bepler, T., Kelley, K., Noble, A. J. & Berger, B. Topaz-Denoise: general deep denoising
models for cryoEM and cryoET. *Nat Commun* **11**, 5208 (2020).
https://doi.org/10.1038/s41467-020-18952-1
58   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for
rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296
(2017). https://doi.org/10.1038/nmeth.4169
59   Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of
Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501 (2010).
https://doi.org/10.1107/S0907444910007493
60   Afonine, P. V. *et al*. Real-space refinement in PHENIX for cryo-EM and
crystallography. *Acta Crystallogr D Struct Biol* **74**, 531-544 (2018).
https://doi.org/10.1107/S2059798318006551
61   Chen, V. B. *et al*. MolProbity: all-atom structure validation for macromolecular
crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12-21 (2010).
https://doi.org/10.1107/S0907444909042073
62   Pettersen, E. F. *et al*. UCSF Chimera--a visualization system for exploratory research
and analysis. *J Comput Chem* **25**, 1605-1612 (2004).
https://doi.org/10.1002/jcc.20084

1008