Leaving X: How scale insects evolved alternatives to chromosomal sex determination

Andrew J. Mongue[1], Erin C. Powell[2], Tracy Liesenfelt[1], Ethan Grebler[1], Amanda Markee[1,3]

[1]Department of Entomology and Nematology, University of Florida, Gainesville, FL 32611, USA
[2]Division of Plant Industry, Florida Department of Agriculture and Consumer Services, Florida State Collection of Arthropods, Gainesville, FL 32608, USA
[3]Richard Gilder Graduate School, Division of Invertebrate Zoology, American Museum of Natural History, New York, NY 10024, USA

Abstract

Reproduction is a core feature of life, which makes understanding the reproductive diversity of organisms a fundamental goal of evolutionary biology. It is still unclear why sex determination systems are faithfully conserved for hundreds of millions of years in some taxa but repeatedly turn over in others. A prime example is the true bugs, Hemiptera. The vast majority of families employ a conserved X sex chromosomal system, but in scale insects (Coccomorpha), sex determination systems are far more varied. Different families may retain X chromosomes, employ a rare form of haplodiploidy known as paternal genome elimination (PGE), or forgo sexual reproduction in favor of simultaneous hermaphroditism. While these systems have been known for decades, only recently has non-model genomics enabled their study with modern bioinformatic methods. We collected and sequenced representatives of rare, early-diverging scale insects to study genomic changes that coincide with transitions between sex determination systems. Using ancillary expression data, we also explored sex-biased gene expression in a subset of scale insects on both sides of one transition. In a newly updated molecular phylogeny, we recovered two independent losses of X chromosomal sex determination, once in the family Monophlebidae and a second time at the base of the families that employ PGE. In the former, genomic rearrangements begin before the X is lost and any previously X-linked orthologs have been absorbed into one of two massive, shuffled autosomes in the hermaphrodites. In the latter, the X chromosome itself is faithfully preserved in early diverging PGE taxa; it appears to have merely lost its sex determination function. We analyze ortholog conservation across these transitions and find few genes lost, with only one orthogroup lost in both transitions, and a number of orthogroups gained, suggesting these large transitions did not require large changes to the gene content of the X chromosome. Expression data suggests that extreme sexual dimorphism of gene expression predates sex determination turnover and may have helped enable it. Together these new observations illustrate the multiple routes to sex determination turnover, help refine hypotheses for its causes, and provide a wealth of new resources for the modern genomic study of scale insects.

Introduction

The core goal of evolutionary biology is to explain the patterns of diversity of life and the processes that generate these patterns. Across more than 150 years of research, traits involved in reproduction have been of particular interest (starting with Darwin 1871), but the true extent of their diversity was not fully apparent until scientists began to study microscopic traits, including the chromosomes themselves. The discovery of chromosomal sex determination grew out of

observations of what were initially known as "accessory chromosomes," so termed because
they varied between males and females of the same species of insects and were thus initially
inferred to not be strictly necessary for the organism (McClung 1901, 1902). In subsequent
research, their true importance in sex determination has been recognized, as have the
numerous variations on these systems (Bachtrog et al. 2014), including some that defy
Mendel's familiar laws of inheritance (Ross et al. 2022). This staggering array of reproductive
diversity belies something of a paradox, however. Despite undeniable overall diversity, many
ancient, speciose clades have sex determination systems that are faithfully conserved across
hundreds of thousands of species and hundreds of millions of years of evolution (Rodríguez
Delgado et al. 2009; Fraïsse et al. 2017; Vicoso 2019). Thus, a more specific goal of
evolutionary reproductive biology is to understand how and under what conditions sex
determination systems turn over.
58
To reach this goal requires systematic knowledge of sex determination systems across species,
a way to track the genomic and genetic changes that enable it, and a suitable system in which
to study the variation. Tools to assess the former have been available for decades thanks to
microscopy and cytogenetic analyses. Note that the observations of sex chromosomes
referenced above are more than a century old (McClung 1901) and subsequent studies have
surveyed such traits broadly in the intervening decades (Hughes-Schrader 1925; Núñez 1962;
White 1973). It is the second piece, a deeper understanding of the genomic and genetic
changes that underlie differences in karyotype, that has lagged behind. The field of genome
sequencing is barely more than 30 years old (Olson 1993) and extension of these approaches
beyond model species is roughly half that age (Ekblom and Galindo 2011; Unamba et al. 2015),
so it is only recently that it has become tractable to ask broadly how turnover of sex
determination occurs at the genomic and genetic levels.
71
Finally, with both types of evidence cataloged or easily attainable, it is worth carefully
considering the choice of study taxa. As already alluded to, sex determination is variable across
the tree of life but tends to be conserved across large taxonomic groups. For instance,
mammals generally share the same orthologous X sex chromosome across over 100 million
years of evolution (Rodríguez Delgado et al. 2009) and, thus, offer limited insight into sex
determination turnover. And even where sex chromosome systems have turned over, like-for-
like transitions are the most commonly studied (Vicoso and Bachtrog 2015), for example the X
chromosome being replaced by a new X chromosome that was formerly an autosome (Vicoso
and Bachtrog 2013). While there is still more to learn in these systems, researchers have
already done much to characterize these turnovers. For instance, theory predicts that sexual
conflict can favor the evolution of new sex determination genes on autosomes, which can
supplant the existing sex determination mechanism (Van Doorn and Kirkpatrick 2007). Likewise
genomic analysis of like-for-like transitions in *Drosophila* flies has shown that such turnovers are
associated with the movement of genes expressed in males off of new X sex chromosomes
(Layana et al. 2025), as predicted based on the fact that X chromosomes spend more time in
females than males (Klein et al. 2021). Results from these systems do little to explain other,
non-like-for-like turnovers of sex determination, however.
89

Scale insects and the diversity of reproductive strategies

Scale insects (Hemiptera: Coccomorpha) are a uniquely well-suited choice for the study of
drastic sex determination turnover (Gavrilov 2007). This diverse infraorder, consisting of
approximately 32 extant and 19 extinct families (Vea and Grimaldi 2016; Powell et al. 2024) of
plant-feeding insects, belongs to the order Hemiptera (true bugs), which generally have well-
conserved XX/X0 or XY sex determination (Pal and Vicoso 2015); in other words, most
Hemiptera share X chromosomal sex determination and the same orthologous chromosome
serves as the X across species. However, within scale insects, different families may retain
chromosomal XX/X0 sex determination (Nur 1980), employ a rare form of haplodiploidy known
as paternal genome elimination (Nur 1966; de la Filia et al. 2021), or even largely forgo sexual
reproduction, as in the simultaneous hermaphrodite giant scale insects (Hughes-Schrader
1925), at least some of which still produce rare haploid males (Royer 1975; Hodgson and Foldi
2006; Mongue et al. 2021). In both cases, the X chromosome system has been replaced with a
fully autosomal genome in which the dynamics of the female/hermaphrodite genome (*i.e.*, full
diploidy) did not change across these transitions. Males, in contrast, have transitioned from
mostly diploid with a single haploid (X) chromosome, to genome-wide haploidy, albeit in two
very different contexts. Under paternal genome elimination, offspring sex ratio may fluctuate
over a mother's lifespan but overall remains at the classical 50-50 male-to-female ratio (Ross et
al. 2012a), but in the hermaphrodites, male production is rare, with the vast majority of
individuals being self-fertile hermaphrodites that produce more self-fertile offspring (Mongue et
al. 2021). Uncovering when and how scale insects have transitioned between these systems
can shed crucial light on the causes and consequences of sex determination turnover occurs
more generally.

To do so, we have (1) sequenced 17 previously unsampled species and assembled genomes
for each, including chromosome-level genomes for five species of scale insects from different
families and sex determination systems, (2) confirmed their places in the evolution of this group
via phylogenomics, (3) performed macrosynteny analyses to understand patterns of
conservation or change of chromosome organization, examined (4) orthology across scales to
track conservation of X-linked genes, and (5) expression differences between the sexes of
some species for insight into the state of the X chromosome before and after transitions.
Together these results show that the otherwise conserved hemipteran X has been lost twice, in
one case absorbed into a much larger autosome in the hermaphroditic scales as part of a
massive condensing and shuffling of chromosomes and, in the other case of paternal genome
elimination, the X chromosome was retained essentially intact at first, but apparently without its
sex determining function. Thus, major turnovers can occur in distinct ways, even from the same
starting genomic architecture.

Methods
Sourcing representatives of different sex determination systems

We downloaded existing datasets for scale insects representing different sex determination and
reproductive systems. These data ranged from assembled chromosome-level genomes to

unassembled transcriptomic reads. Details can be found in Table S1. To complement these
data, we added new sequence data for another 17 species listed in Table S2. Most of these
taxa have well characterized reproductive biology (also listed in Tables S1&2), with some
exceptions. For the species we sequenced, we confirmed putative sex chromosomes via
coverage difference between male and female sequenced reads where possible. For existing
datasets, the least well characterized was the recently described *Coronaproctus castanopsis* Li,
Xu & Wu (Monophlebidae). The species description and genome assembly do not report the
sex determination system (Zheng et al. 2023; Huang et al. 2024); however, evidence strongly
suggests that it is XX/X0. First, a recent molecular phylogeny based on genomic and
transcriptomic data confirmed its status as a monophlebid and suggests that *C. castanopsis*
shares a more recent common ancestor *Drosicha* spp. (Kuwana) than it does with the
hermaphroditic, haplodiploid *Icerya* spp. (Song et al. 2025). *Drosicha* Walker species have been
previously reported as XX/X0 (Yokogawa and Yahara 2009) and in synteny analyses we found
orthology between the smallest chromosome of *C. castanopsis* and the smallest chromosome of
D. contrahens, which itself is synthetic to the newly identified X of *Neosteingelia texana*.
Additionally, in the haplodiploid hermaphrodites like *Icerya purchasi*, production of pure males is
known to be exceedingly rare (e.g., a recent sample of over 300 adults, collected from 27
populations across the world recovered no males: Mongue et al. 2021). The species
descriptions for *C. castanopsis* include both male and female morphology with no indication that
males were rare or difficult to obtain (Li et al. 2023; Zheng et al. 2023), suggesting roughly equal
sex ratios expected of chromosomal sex determination.
155
156 *Collecting samples from across sex determination systems in scale insects*
157
For newly generated data, we sampled species across sex determination systems of scale
insects, with special interests in targeting representatives of families that had no existing
sequence data and/or with documented X chromosomes (Gavrilov 2007), but previously
karyotyped species were not always readily available for collection. Many of these scale insects
have cryptic life cycles and spend much of the year under tree bark or underground (Kosztarab
and Watson 1994; Thomson et al. 2021). In several cases, we were able to obtain only a few
individuals from a target species and sometimes only females, not enough starting tissue to
generate chromosome-level genomes, perform the sex chromosome analyses, or differential
expression analyses described below. Still, these samples represent new sequencing resources
for an understudied group of insects and help corroborate existing hypotheses of phylogenetic
relationships.
169
We aimed to sequence DNA from a single individual where possible, but for some smaller taxa,
we resorted to pooling multiple individuals. We sequenced these samples for inclusion in the
larger phylogeny, but given the fragmented nature of their draft genomes, we did not perform
the synteny and orthology analyses described below. Species sequenced under this strategy
include: *Orthezia annae* Cockerell (Ortheziidae), *Insignorthezia insignis* (Browne) (Ortheziidae),
Neogreenia osmanthus (Yang & Hu) (Qinococcidae), *Eumargarodes laingi* Jakubski
(Margarodidae), *Stomacoccus platani* Ferris (Steingeliidae), *Puto decorosus* McKenzie
(Putoidae), *Puto cupressi* (Coleman) (Putoidae), *Nipaecoccus nipae* (Maskell)

(Pseudococcidae), *Nipaecoccus floridensis* Beardsley (Pseudococcidae), *Pseudophilippia*
quaintancii Cockerell (Coccidae), and *Pityococcus rugulosus* McKenzie (Pityococcidae). Of
these, *P. decorosus* was the only species for which we obtained a male. For more information
on collected species, please see Table S2.
182
We were able to find an abundant ensign scale (Ortheziidae): the Spanish moss ensign scale,
Graminorthezia tillandsiae (Morrison) and a member of Kuwaniidae: the giant pecan scale,
Neosteingelia texana Morrison. Because much is still unknown about the ecology and life
history of these rarely-studied scale insects, we provide brief observational notes on each in the
supplement. We also sequenced two hermaphroditic monophlebids, *Crypticerya genistae*
(Hempel) and *Crypticerya rileyi* (Cockerell). We followed the same general sequencing strategy
for these species, with modifications noted below. Because scale insects are highly sexually
dimorphic, including in body size (with females being the larger sex, Vea et al. 2016) and the
expected karyotypes are either fully haplodiploid or XX/X0, we based our primary assemblies on
females. To identify sex chromosomes, we used a typical coverage-based approach (Mongue et
al. 2017, 2022; Mongue and Baird 2024; Baird et al. 2025), and sequenced DNA from pooled
males. Slide-mounted voucher specimens associated with each of the aforementioned
collections are deposited in the Florida State Collection of Arthropods (FSCA), Gainesville,
Florida, USA. We additionally obtained a limited sample of an unidentified *Puto* species: one
adult male and five adult females. To make the most of this valuable XX/X0 sample, we chose a
distinct sequencing approach, as described below.
199
DNA extraction and sequencing approach
For all samples, we used a modified Omniprep (G-Biosciences, St. Louis, MO, USA) chloroform
extraction and ethanol precipitation protocol, following previously successful application in scale
insects (Mongue et al. 2024; Liesenfelt et al. 2025a). Specifically, we used a dounce
homogenizer to grind whole individuals, extended the lysis step overnight, and used 2 µL of
20mg/mL glycogen to facilitate DNA precipitation during a 1hr rest at -20C. In the case of *N.*
texana we used a single adult female, but because *G. tillandsiae* is much smaller (length
<1mm), we pooled seven females into one extraction. We sent DNA extractions from *N. texana,*
C. genistae, and *C. rileyi* to Arizona Genomics Institute (AGI) (Tucson, AZ, USA) for standard
PacBio HiFi library prep and sequencing on a Revio (Pacific Biosciences, Menlo Park, CA,
USA). We sequenced *G. tillandsiae* with the University of Florida's Interdisciplinary Center for
Biotechnology Research (Gainesville, FL, USA) where we used a low-input library approach and
sequenced on a Sequel IIe machine.
213
214 To identify the X sex chromosomes in these species, we separately sequenced DNA from
215 males of both species to assess differences between autosomal and sex chromosome
216 coverage. For male extractions, we used the same extraction protocol, but in both species we
217 pooled five individuals per extraction. Once extracted, we sent samples away for sequencing on
218 an Illumina Novaseq at Novogene (Sacramento, CA, USA).
219
220 Separately, we snap froze (from live to -80°C) 50mg of female tissue from each species for Hi-C
221 protocols. We sent frozen tissue to AGI for crosslinking and sequencing. For both this and the

222 male sequencing above, we waited until we received initial PacBio data and calculated genome
223 size to ensure we requested adequate coverage (~50x).
224
225 For the *Puto* sp. sample, we extracted DNA from the single male using the protocol above but
226 falling short of the input thresholds for standard HiFi library preparation, we opted to use the
227 PacBio AmpliFi whole-genome amplification kit. We then submitted the female tissue, preserved
228 in ethanol, for Hi-C library preparation with AGI. Thus, although we used a different strategy,
229 ultimately, we still had sequencing from a male to identify the X chromosome by coverage.
230
Genome assembly and validation
For all species, we received raw PacBio HiFi reads. We started by passing the reads to jellyfish
v2.3.0 (Marcais and Kingsford 2012) to count 21-mers and build a distribution of the frequency
of these sequences. Simple calculations on these frequencies gave us both an expectation for
genome size and depth of sequencing coverage. Next, we assembled raw reads with hifiasm
v0.8 (Cheng et al. 2021). For this step, we included the -l 3 parameter for minimizing duplicate
haplotig retention and set the --hom-cov parameter to the peak observed in our k-mer counting
step. For these initial assemblies, we calculated basic summary statistics (N50, sequence
count, etc.) and ran BUSCO v5.8.3 using the hemiptera_odb10 lineage to assess completeness
and duplication (Manni et al. 2021).
241
For each species, we took the primary assembly through the purge_dups pipeline published
only as a git repo (https://github.com/dfguan/purge_dups) to identify cases in which both
haplotypes were kept as duplicate sequences in the assembly via a mix of coverage and
sequence identity during self-alignment. Note that purging duplicates is an iterative process and
can be run in successive rounds. Indeed, because we pooled multiple *G. tillandsiae* as a single
sample, we expected retained haplotigs to be a larger issue in its assembly. We assessed
BUSCO scores and assembly stats after the first round and ran a second round of duplicate
purging on the *G. tillandsiae* assembly based on the high rate of duplication.
250
After purging, we aligned Hi-C sequencing reads following Arima's recommended pipeline
(detailed on their git repo: https://github.com/ArimaGenomics/mapping_pipeline). In brief, we
quality trimmed reads using the repo's custom scripts, separately mapped each of the two mate
pairs to the draft genome using BWA v7.9a (Vasimuddin et al. 2019), and combined alignments
with Arima's scripts. We used the merged alignments as inputs for YaHS v1.1 (Zhou et al.
2023). After scaffolding, we again assessed assembly statistics before manual curation. For this
step, we used the YaHS command "juicer pre" to generate inputs for manual review in Juicebox
v2.17.00 (Durand et al. 2016). We made minor adjustments based on contacts as needed,
output the updated scaffold relationships, and updated the assembly with YaHs "juicer post"
command (Zhou et al. 2023). For the final time, we calculated assembly statistics and BUSCO
completeness (Manni et al. 2021).
262
To identify the sex chromosomes, we aligned male short-read sequencing to the respective
finalized reference using Bowtie2 v2.5.4 (Langmead and Salzberg 2012) using the "--very-
sensitive-local" alignment parameters. We then coordinate-sorted the alignments with Picard

tools v3.2.0 (Wysoker et al. 2013) and used SAMtools v1.9's coverage function (Danecek et al.
2021) to calculate coverage across scaffolds. For the *Puto* sp., we mapped long reads to the
reference with minimap2 VERSION (Li 2018) using the "-x map-hifi" parameter to optimize for
that data type. Based on the expectation of X0 karyotype for males, the X chromosomal scaffold
should have 50% the coverage depth of the diploid autosomes.
271
Phylogenomic analysis
We followed the BUSCO_phylogenomics pipeline
(https://github.com/jamiemcg/BUSCO_phylogenomics). For each genome or transcriptome
included in the analysis, we assessed the BUSCO completeness as above (again using the
hemiptera_odb10 dataset). Then, using the scripts provided by the git repo, we parsed single-
copy BUSCO amino acid sequences, with a threshold of presence in 75% of samples for
inclusion. In the analysis presented here, that totaled to 1,477 single-copy protein sequences.
The pipeline produced trimmed protein alignments and individual gene trees, which we
analyzed in two ways. First, we concatenated the individual gene alignments with custom bash
scripts and used IQTree (Minh et al. 2020) v2.2.2.7 to compute the maximum likelihood tree
using the "Q.insect+FO+G8" model of substitution, with confidence assessed with 1000 ultrafast
bootstraps and Shimodaira-Hasegawa-like approximate likelihood ratio test (SH-aLRT)
(Guindon et al. 2010).
285
Second, to offer an additional phylogenetic hypothesis that accounts for incomplete lineage
sorting, we used ASTRAL (Zhang et al. 2018) to generate a consensus species tree from the
individual gene trees with confidence assessed as posterior quartet probabilities (Sayyari and
Mirarab 2016). As both trees showed agreement, but the species tree approach offers less
informative branch lengths, we present only the maximum likelihood approach in the main text
and the ASTRAL approach in the supplement. In both cases, we manually rooted trees with
Acyrthosiphon pisum (Harris) and *Adelges tsugae* (Annand) as the non-scale outgroups,
visualized the output with FigTree v1.4.4 (Rambaut 2010), and manually colored branches by
sex determination type.
295
Annotation and macrosynteny analysis
With finalized assemblies, we next generated gene annotations for downstream analyses. First,
we used RepeatModeler v2.0 (Flynn et al. 2020) to *de novo* identify repetitive sequences,
including the --LTR parameter to search for long terminal repeats. We passed the identified
repeats to RepeatMasker v4.0.9 (Smit et al. 2019) to generate a softmasked assembly ready for
annotation. We used a standard approach across species: BRAKER v3.0.8 (Hoff et al. 2019)
with protein evidence from collected BUSCO protein sequences across scale insects. To
minimize biases in downstream analyses arising from different annotation pipelines, we re-
annotated published assemblies with the BRAKER protein pipeline described above Then, we
used MCScanX v1.0.0 to infer synteny (Wang et al. 2024). This method uses BLAST similarity
to identify putative 1-to-1 orthologs between pairs of species and then takes advantage of the
scaffold coordinate information in gene annotations to identify regions of the genome in one
species that correspond to the orthologous region of the other. We based pairwise comparisons
on the phylogenetic relationships inferred above.

310
Conservation of X-linked genes
Once we established the identity of the X chromosomes, we examined in more detail the fate of
X-linked genes across the losses of the X in hermaphrodite and the paternal genome
elimination clades. In particular, we examined two patterns. First we subsetted our gene
annotations for *G. tillandsia*, *N. texana*, *C. castanopsis*, and *Puto* sp. to only compare genes on
the X chromosomal scaffolds of these species to each other and to the full genomes of either (1)
I. purchasi, *C. genistae,* and *C. rileyi* or (2) the chromosome level assemblies of the PGE
species to look for conserved, X-linked genes lost in the transition to other sex determination
systems. Second, we compared the full genomes of the X sex determination species to the full
genomes of groups (1) or (2) above to look for any genes conserved in the hermaphrodites or
paternal genome elimination species but missing from the X species. We used OrthoFinder
v2.5.2 (Emms and Kelly 2019) to identify orthogroups. We then searched NBCI's RefSeq
protein database to infer functional information via sequence homology.

Expression data generation and analysis
We sought to further characterize the state of the X chromosome both before and after its loss
by examining patterns of gene expression between males and females. In practice, we were
only able to obtain sufficient samples of both sexes for two species: *N. texana* and *T. liriodendri*.
For each of these species, we generated Illumina paired end RNAseq from four biological
replicates of individual adult females and four replicates of pooled males. For *N. texana*, we
used pooled replicates of 2–4 males per extraction and for *T. liriodendri* we used replicates of 5
males per extraction.

For RNA extractions, we utilized a RiboPure™ RNA Purification Kit (Invitrogen™) with slight
modification to the protocol to account for low tissue mass of our samples. We thawed
previously frozen tissue directly into 500 μL of the kit provided TRI reagent to account for tissue
mass < 50 mg. With a lower starting volume, the aqueous phase measured ~ 250 μL after
phase separation. To keep the 1:2 volume ratios consistent to those in the protocol, we added
125 μL of 100% ethanol to this aqueous phase for RNA purification. Extractions were then
eluted into 50 μL of buffer instead of the recommended 100 μL. Our extractions were finally sent
to Genewiz (South Plainfield, NJ, USA) to generate 30 million paired end Illumina reads per
sample.

Added to these newly sequenced samples, we downloaded expression data for one adult male
and two early adult female *Ericerus pela* (Chavannes) from the NCBI Sequence Read Archive
(SRA) (accessions:SRR1027692 and SRR9617905 + SRR9617914, respectively). For each
SRA accession, we quantified per-sample expression against the BRAKER annotation
generated above using Kallisto v0.51.1 (Bray et al. 2016). We analyzed expression as TPM
(transcripts per million) using R scripts to calculate SPM, the specificity metric, (Kryuchkova-
Mostacci and Robinson-Rechavi 2017), as the proportion of each gene's expression in one sex
compared to the other. We chose this approach as a calculated tradeoff to make the most of
available data, as the single male sample from *E. pela* would not allow for more traditional
differential expression analyses, at the cost of losing statistical evaluation of expression

difference. We plotted sex specificity as the amount of expression in females, a value ranging from zero (male-limited expression) to one (female-limited expression). Additionally, following previous evolutionary genetic analyses (Mongue et al. 2022), we binned genes into categories of male-biased (SPM < 0.3), unbiased (0.3 < SPM < 0.7), or female-biased (SPM > 0.7). With these categories, we used X^2 tests of independence to assess whether the sex-biased composition of genes different between X or former-X and autosomes in *N. texana*, *E. pela*, and *T. liriodendri*.

Finally, we explored dosage compensation in the one X chromosome species for which we have RNA data, *N. texana*. In particular, we took the subset of genes expressed only in males or only in females and calculated the mean of each gene's TPM expression across four biological replicates for that sex and compared it across autosomal and X-linked genes using a Wilcoxon test. Because we found a majority of genes with sex-limited expression in our SPM analyses, we did not directly compare expression between the sexes.

Results

Assemblies, X assignments, and annotations

We assembled 17 new scale insect genomes to various levels from basic Illumina-only unordered contigs to Hi-C scaffolded chromosome level assemblies. For brevity and to keep the focus on the evolutionary genomic findings, we describe the full assembly statistics and report accessions in the supplement. We used assemblies of all levels for the phylogeny to infer turnover of sex determination, but proceeded to further analysis of synteny, orthology, and gene expression for only the subset of chromosome-level assemblies.

An updated phylogeny of scale insects

We reconstructed the phylogeny of scale insects using a mix of publicly available and newly generated datasets. We used two different methodologies, a maximum likelihood inference of sequence evolution using concatenated protein sequences and a consensus species tree inferred from individual gene trees. We present the maximum likelihood approach in the main text (**Figure 1**) and the consensus species tree in the supplement (**Figure S1**). Both approaches yielded the same topology and had high support metrics for relationships;therefore, the following results summary applies to both analyses.

387
Figure 1. Maximum likelihood phylogeny of scale insects annotated with sex determination system. Relationships inferred from concatenated BUSCO amino acid sequences from genomes and transcriptomes. Support values follow the format Shimodaira-Hasegawa-like approximate likelihood ratio test (SH-aLRT) / Ultrafast Bootstrap values; all nodes have 100/100 support unless otherwise noted. Sex determination systems are color-coded as follows: red - X chromosomal sex determination, purple - haplodiploid hermaphroditism, black - paternal genome elimination. Species with an * next to their label have newly sequenced draft genomes, those with illustrations have newly sequenced chromosome-level assemblies for this study. Branch lengths are proportional to the expected number of amino acid changes per site as indicated by the scale bar.

We rooted the tree with known outgroups, the pea aphid, *Acyrthosiphon pisum* and hemlock
woolly adelgid, *Adelges tsugae* (Johnson et al. 2018). We recovered most relationships in
agreement with other published phylogenies (Gullan and Cook 2007a; Cook et al. 2002;
Hodgson and Hardy 2013; Vea and Grimaldi 2016), so we focus on the main differences and
additional insight gained from newly sequenced taxa here. First, we recovered Ortheziidae as a
monophyletic group and the earliest diverging family of scale insects. In fact, based on current
sampling, all other family level groupings are valid with the exception of Coccidae. Our sole
representative, a non-specified member of the genus *Aclerda*, is placed confidently in the
middle of Coccidae. Aclerdids could represent a lineage of grass-specialist soft scales (though
not all aclerdids are found on grasses, see García Morales et al. 2016 for a review) and its
validity as a family may need to be reevaluated. Putoidae is recovered as a valid family falling
out between what has historically been considered 'Archaeococcoidea' and 'Neococcoidea'.
However, we do not find evidence that these two superfamily groupings are monophyletic. We
lack resolution to validate most genus-level relationships, but members of *Ceroplastes*, *Coccus*,
Aspidiotus, *Pseudococcus*, *Icerya*, *Crypticerya*, and *Puto* all share a more recent common
ancestor with a congener than any other taxon. The one exception is the genus *Nipaecoccus*;
Nipaecoccus floridensis and *N. nipae* group together as expected, but *N. viridis* is confidently
placed closer to *Balanococcus diminutus* than to other *Nipaecoccus*, suggesting the generic
placement of *N. viridis* may need to be revisited.
418
Finally, most importantly for the purposes of this study, the relationships described above,
combined with existing knowledge of family level reproductive systems (reviewed in Gavrilov
2007), suggests a straightforward scenario for sex determination turnover. Parsimoniously, X
chromosomal sex determination was lost twice, once in the lineage of simultaneous
hermaphrodites within Monophlebidae (here represented by *Icerya* and *Crypticerya*), and a
second time independently at the base of the group of paternal genome elimination families
including Pseudococcidae, Coccidae, Diaspididae, and others. We also explored the evolution
of different types of paternal genome elimination, but lacking information for most sequenced
Coccidae species, we cannot infer the timing or number of transitions within paternal genome
elimination. The only straightforward inference, based on the state of early-diverging taxa, is
that lecanoid paternal genome elimination appears to have evolved first, with Comstockiella and
diaspidid types arising secondarily (Figure S2).
431
432 *Macrosynteny: two divergent paths to losing X sex determination*
433
We mapped 1-to-1 orthologs between species across the subset for which we have
chromosome-level assemblies. First, we consider the Monophlebidae (**Figure 2**). Starting with
the outgroups, despite the X having split into two chromosomes, an X_1X_2 system, in *Adelges*
tsugae, both of these sex chromosomes are syntenic to the single X chromosome of
Acyrthosiphon pisum, thus the X is conserved in the outgroups. However, we found overall very
little synteny between our earliest diverging scale insect, *G. tillandsiae*, and other
Sternorrhyncha. Interestingly, what little detectable orthology exists between the *G. tillandsiae*
and *Acyrthosiphon pisum* is autosome-to-autosome, suggesting that the X chromosome has

 highly diverged between the scale insects and other Sternorrhyncha, or potentially even turned
443 over.
444

445
Figure 2. Macrosynteny across the transition from X to simultaneous hermaphroditism.
Individual colored horizontal bars represent chromosomal scaffolds. Lines connecting them
represent conserved orthologs between a pair of species. For each species, where applicable,
the X chromosome(s) are sorted to the left-most side of the plot and labeled. The dashed box
highlights the fate of X-linked orthologs across the turnover in sex determination systems.
Phylogenetic relationships follow those recovered in Figure 1 with the same coloring for

branches. Species depicted: 1. *Adelges tsugae* 2. *Acyrthosiphon pisum* 3. *Graminorthezia*
tillandsiae 4. *Neosteingelia texana* 5. *Drosicha contrahens* 6. *Coronaproctus castanopsis* 7.
Icerya purchasi 8. *Crypticerya genistae* 9. *Crypticerya rileyi*. Illustrations courtesy TL.
455
Next, comparing *G. tillandsiae* to *N. texana*, we found clearer evidence for a conservation of the
X chromosome. Both species' X chromosomes were confirmed by coverage difference between
males and females within species (see Supplement), and the two chromosomes only share
orthologous hits with each other. There are some rearrangements to the karyotype, as expected
when comparing n = 9 vs. n = 7 chromosomes (based on Hi-C contacts) but overall much more
conservation of orthologs.
462
Regarding, *N. texana* and *D. contrahens*, we again found good evidence for conservation of the
X chromosomes, but note that the assembly of *D. contrahens* contains no sex linkage
information. This genus is known to be XX/X0 based on karyotype, thus we inferred the identity
of the X via synteny to the *N. texana* X. In contrast to the conserved X, we found substantial
reorganization of the autosomes. *Drosicha contrahens* has a mere n = 2 autosomes, which
appear to have resulted from simple fusions given that these two autosomes map cleanly to 3
and 4 of the *N. texana* autosomes, with only 1 inferred to have orthology to multiple *D.*
contrahens autosomes. Intriguingly, this suggests that the extreme karyotype reduction found in
the largely asexual *Icerya* and *Crypticerya* (n = 2) began *before* the transition to asexuality, not
as a consequence of it.
473
Considering *D. contrahens* and *C. castanopsis*, we recovered additional evidence that the latter
is an XX/X0 sex determination system. Namely, the smallest of the three chromosomes
(CM065054.1) shows fewer orthologs than the other chromosomes and only has hits to the X of
D. contrahens. There appears to have been autosomal rearrangement between lineages here,
as one end of each autosome maps to the opposite autosome in the other species. Given that
we recover the same pattern between *C. castanopsis* and *I. purchasi* below though, it is more
parsimonious to conclude that this rearrangement happened once in the lineage leading to
Coronaproctus rather than shuffling and and reverting between multiple lineages.
482
Finally, we recovered much more shuffling in the comparison that spans the loss of X sex
determination between *C. castanopsis* and *I. purchasi*. The little remaining orthology to the X
maps to the larger of the two autosomes, which seems indicative of an X-autosome fusion, but
this only captures a tiny fraction of the rearrangements. The two autosomes of *C. castanopsis*
do not cleanly correspond to the two autosomes of *I. purchasi*. The two large inter-chromosomal
rearrangements may reflect a derived karyotype in *Coronaproctus*, as mentioned above, but we
also recovered myriad intra-chromosomal rearrangements within the autosomes between these
species. Continuing further into the *Crypticerya* species, we found additional rearrangements
both within and between chromosomes with plenty of conserved genes but very short blocks of
synteny.
493

Figure 3. Macrosynteny across the transition from X to paternal genome elimination.
Individual horizontal bars represent chromosomal scaffolds. Lines connecting them represent
conserved orthologs between a pair of species. For each species, where applicable, the X
chromosome is sorted to the left-most side of the plot and labeled. The dashed box highlights
the fate of X-linked orthologs across the turnover in sex determination systems. Phylogenetic
relationships follow those recovered in Figure 1 with the same coloring of branches. Species
depicted: 1. *Neosteingelia texana* 2. *Puto* sp. 3. *Acanthococcus lagerstroemiae* 4. *Kerria lacca*
5. *Ericerus pela* 6. *Toumeyella liriodendri* 7. *Ceroplastes pseudoceriferus* 8. *Coccus hesperidum*
9. *Phenacoccus solenopsis* 10. *Planococcus citri* 11. *Pseudococcus jackbeardsleyi* 12.
Paracoccus marginatus 13. *Balanococcus diminutus* 14. *Nipaecoccus viridis*.

For the second loss of X sex determination, we explored the transition to paternal genome
elimination (**Figure 3**). First, we confirmed that the X of the closest sequenced outgroup to
paternal genome elimination, *Puto* sp., is syntenic to the scale insect X via comparison to *N.
texana.* Next, we traced the orthologous chromosome into the paternal genome elimination
clade and found, surprisingly, that it is essentially intact, albeit no longer sex-determining, in the
crapemyrtle bark scale, *Acanthococcus lagerstroemiae* (Kuwana), where it is orthologous to
chromosome 7 in the most complete assembly (scaffold CM063112.1). Indeed, this
chromosome is well conserved in *Kerria lacca* (Kerr) and *Ericerus pela* and only begins to break
apart as part of a broad, substantial increase in chromosome number in *Toumeyella liriodendri*
(Gmelin) and *Ceroplastes pseudoceriferus* Green (n = 17 and 18, respectively). These
karyotype changes appear to be simple fissions as most *E. pela* chromosomes correspond to
two *T. liriodendri* chromosomes with an additional translocation from an autosome. In *Ce.
pseudoceriferus*, the former X has split into three smaller chromosomes. Oddly, the closest
sequenced relative of *Ce. pseudoceriferus*, *Coccus hesperidum* (Linnaeus), has a reduced
karyotype (n = 7) that appears to have resulted from secondary fusion of the fragmented
chromosomes as the fewer chromosomes here incorporate similar, but sometimes distinct,
syntenic blocks compared to the *E. pela* genome. In this species the remnant X has fused with a
syntenic block that was previously autosomal in all other sequenced species. *Coccus
hesperidum* also marks the last point at which the former X is easily distinguishable in the
paternal genome elimination species.

There have been major rearrangements across all chromosomes leading to the true mealybugs
(Pseudococcidae) and formerly X-linked orthologs are scattered across all five chromosomes of
Phenacoccus solenopsis Tinsley. After this rearrangement, however, genome structure is
remarkably well conserved. Five of the six chromosome-level assemblies show the n = 5
karyotype, with only individual-level gene-trafficking events rearranging synteny for the most
part, barring two exceptions. First, *Paracoccus marginatus* has only n = 4 chromosomes, though
this reduction is the result of a single, simple fusion of the second and fourth linkage groups in
other mealybugs. Second, even microsynteny (*i.e.*, the individual ordering of orthologous genes
within chromosomes) is well preserved across most species, but in comparisons between *Ph.
solenopsis* and *Planococcus citri* (Risso), as well as *Pl. citri* and *Pseudococcus jackbeardsleyi*
Gimpel & Miller, we recovered large inversions on four of the five chromosomes.

539   *Conservation of genes across sex determination turnover*
540
541   We searched for orthologs with two distinct conservation patterns: first, those conserved on the
542   X chromosomes but absent from the non X-chromosomal scales and, second, those absent
543   from anywhere in the genome of X sex determination species but conserved across all
544   hermaphrodites or paternal genome elimination species.
545
Starting with the former, we searched across the X chromosomes of *G. tillandsiae* (n = 3,534
genes), *N. texana* (2,497 genes), *C. castanopsis* (1,967 genes), and *Puto* sp. (1,633) compared
to the entire genomes of *I. purchasi* (18,371 genes), *C. genistae* (20,330), and *C. rileyi* (19,871
genes). We recovered 410 orthogroups conserved across all the X chromosomes. Of these
conserved X-linked orthologs, only 8 were absent from all hermaphroditic species (**Table S5**).
Similarly, but even more extreme for the paternal genome elimination clade, we identified only
two conserved X-linked orthologs that were missing from all sampled paternal genome
elimination species (**Table S6**).
554
We also searched for the inverse pattern, novel gene gain in the hermaphrodites or paternal
genome elimination species. We found 664 orthogroups conserved in all three hermaphrodite
species that were missing from all of our annotated X sex determination species. A majority of
these (449, 68%) had no significant BLAST hits and a further 54 matched to "hypothetical" or
"uncharacterized" proteins, leaving 161 with functional hits. There are too many to parse here,
and none appear as immediately crucial to sex determination, but all of these orthogroups, as
well as their top BLAST hits are listed in **Tables S5–6**.
562
Turning to paternal genome elimination, we found 49 orthogroups that were gained and
conserved across all sampled paternal genome elimination species. Of these, 18 (37%)
returned "hypothetical protein" hits or no significant matches. Given the unique nature of
paternal genome elimination, its molecular functions may lie in these poorly characterized
proteins, but we can do no more with this information at present. We identified one better
characterized orthogroup as a DDB1- and CUL4-associated factor 8, which has been implicated
in cancer in humans via changes in methylation regulation (Yang et al. 2015). Less directly
related to reproduction, but interesting nonetheless, we also identified an orthogroup related to
fatty acyl-CoA reductase, which is conserved as a single copy ortholog in *A. lagerstroemiae* and
K. lacca, but in multiple copies, from 2 to 16, in other paternal genome elimination species. This
apparent gene family expansion may be related to the diversity of wax production across the
group (Foldi 1991; Ding et al. 2022). These genes and BLAST hits are cataloged in **Table S7.**
575
576   *Patterns of gene expression and sex-biased genes before and after a loss of X*
577
Finally, we generated RNA sequencing from whole male and female samples of *N. texana* and
T. liriodendri, downloaded public data for *E. pela*, and analyzed patterns of sex-specific gene
expression. First, we explored gene expression levels, as measured by TPM (transcripts per
million), between the X and autosomes of *N. texana* with males and females separately (**Figure**
4, left). As expected, expression levels did not differ between the autosomes and diploid X of

females (W = 61230, p-value = 0.880). The haploid X in males also expressed genes at a
comparable level to the autosomes; in fact, expression trended higher for X-linked genes here
(W = 29963, p-value = 0.060). As such, it appears dosage of the haploid X is upregulated in
females. Owing to extremely dimorphic gene expression and few genes expressed roughly
equally between the sexes, we did not directly test expression levels between sexes.
588
589

590
Figure 4. Gene expression patterns within and between sexes. Left: Dosage
compensation in *Neosteingelia texana*. Genes on the X and autosomes show similar
expression levels (**top**) in females as well as (**bottom**) in males, as expected if the haploid X is
compensated to express genes at a higher level in males.The dot represents a trend for higher
expression on the male X compared to autosomes. **Right: Patterns of sex-biased gene**
expression across species. In each row, individual plots show the SPM (specificity metric) of
expression in male vs. female tissue. In each case, lower values represent male-biased

expression, with SPM = 0 being male-limited expression; SPM = 1 female-limited
expression.The **left** histogram shows the distribution of expression patterns for the X or its
orthologous chromosome(s) after the transition to paternal genome elimination. The **right**
histogram shows the same for (all other) autosomes.
602
Second, we explored the distribution of genes with sex-biased expression. Starting with *N.*
texana, surprisingly, we found that while the X still has a plurality of female-biased genes, it
holds proportionally fewer than the autosomes, as well as proportionally more male-biased
genes (X^2_2 = 93.15, p < 0.0001, **Figure 4, right, top; Table 1**). Even after the loss of the sex
determining function in *E. pela*, the former-X chromosome still has a different composition than
the autosomes (X^2_2 = 100.13, p < 0.0001, **Figure 4, right, middle; Table 1**), with fewer female-
biased and more male-biased and unexpressed genes. In *T. liriodendri*, which has had both
fissions and fusions of the former-X, there is no difference in the proportions of sex-biased
genes between the former-Xs and autosomes (X^2_2 = 1.91, p = 0.591, **Figure 4, right, bottom;**
Table 1).
613

Species	Linkage	Male-biased	Unbiased	Female-biased	No expression
Neosteingelia texana	**X**	**864 (37%)**	**301 (13%)**	**971 (42%)**	**194 (8%)**
	Autosomes	4,651 (29%)	1,849 (11%)	8,544 (52%)	1,323 (8%)
Ericerus pela	**Former-X**	**513 (24%)**	**226 (10%)**	**808 (38%)**	**3401 (28%)**
	Autosomes	3,637 (21%)	2,151 (13%)	7,847 (46%)	599 (20%)
Toumeyella liriodendri	Former-Xs	772 (34%)	449 (20%)	691 (30%)	365 (16%)
	Autosomes	6,282 (33%)	3,878 (20%)	5,882 (31%)	2,900 (16%)

614
615   **Table 1. Distribution of sex-biased genes across the genomes of X and paternal genome**
616   **elimination species.** Numbers represent raw counts of genes by category, parentheses give
617   the percentage of the total genes in that group. Bolding highlights significant differences
618   between linkage categories for *N. texana* and *E. pela*.
619
Discussion
We studied the turnover of reproductive systems across scale insects. To do so, we generated
new chromosome-level references for the Spanish moss ensign scale (*Graminorthezia*
tillandsiae), the giant pecan scale (*Neosteingelia texana*), a giant mealybug (*Puto* sp.), as well
as two giant scales (*Crypticerya genistae* and *C. rileyi*) in service of understanding the changes
to genome architecture associated with change in sex determination. However, before
considering sex determination further, these new genomic resources offer a chance to update

and confirm phylogenetic relationships across a taxonomic group that had, until recently, been
represented by a dearth of genomic data.
629
Insights from a new phylogeny of scale insects
Overall, our results agree with previous findings, with a few notable exceptions and additions.
First, the largest mainly morphological phylogeny includes an early split in scale insects
between Neococcoidea (containing true mealybugs and allies) and a clade including
Monophlebidae, Kuwaniiade, and Ortheziidae (Vea and Grimaldi 2016). A more recent genomic
and transcriptomic study included only monophlebid members of the latter group, and thus
could not evaluate this relationship (Song et al. 2025). With our sampling, we confidently place
Ortheziidae as the earliest diverging family of sequenced scale insects and place Kuwaniidae
sister to Monophlebidae among represented families. Our tree topology recovers the group of
Ortheziidae, Monophlebidae, and Kuwaniidae (sometimes grouped as part of Archaeococcoidea
(e.g., Foldi 2009; Gavrilov-Zimin 2018) as a paraphyletic group.
641
Putoidae is recovered as monophyletic falling out between what has been considered the
'Archaeococcoidea' and 'Neococcoidea' groups. Moreover, Putoidae is confirmed to have an X
chromosomal sex determination, further supporting morphological and molecular evidence that
they should not be placed within PGE Pseudococcide (Downie and Gullan 2004; Hardy et al.
2008; Vea and Grimaldi 2016; Williams et al. 2011; Choi and Lee 2022; Powell and Miller 2025)
despite contested historical placement (McKenzie 1967; Gavrilov-Zimin and Danzig 2012;
Danzig and Gavrilov-Zimin 2014). We also place the recently described *Neogreenia osmanthus*
(Qinococcidae, Zheng and Wu 2023) as the sister taxon to Kuwaniidae, Margarodidae, and
Monophlebidae. Despite superficial similarities to and previous placement within Kuwaniidae
(Wu and Nan 2012), we recover the ground pearls, Margarodidae, as more closely related to
Kuwaniidae than Qinococcidae is. We also sequenced and placed *Pityococcus rugulosus*, the
only sequenced member of Pityococcidae, which has been the subject of some disagreement
previously. One study placed them as sister to Margarodidae (Gullan and Cook 2007), another
as sister to Steingeliidae (in our tree, *Stomacoccus*, Hodgson and Hardy 2013), and yet another
placed them close to Diaspididae (Vea and Grimaldi 2016), the armored scales. Our analyses
agree with the latter placement. While there is no documentation of Pityococcidae's genetic
system, this placement suggests it employs paternal genome elimination. Cytogenetics could
easily confirm or refute this prediction, but our samples were not suitable for these analyses.
Clearly, there is more work to be done on scale insects, especially the early-diverging and
species-poor families; within the scope of this study, however, our phylogenetic tree resolves
the history of sex determination turnover. In our new phylogeny, the most parsimonious
scenario for sex determination turnover is quite straightforward.
664
Studies have shown that the X sex chromosome has been well-conserved across the order
Hemiptera (Pal and Vicoso 2015). Even in the suborder Sternorrhyncha, the group containing
scale insects, the X chromosome is conserved across Aphididae, Adelgidae, and Phylloxeridae
(Li et al. 2023). Particularly in Aphididae, the X chromosome has been preserved despite
extensive autosomal rearrangements (Huang et al. 2025). In contrast, we found very little
orthology between the scale insect X and that of non-scale insect Sternorrhyncha. In fact, we

failed to recover any blocks of synteny long enough to be visible between *G. tillandsiae* and
Acyrthosiphon pisum; the only detectable synteny from the X of *G. tillandsiae* mapped to
autosomes of the aphid. This lack of conservation raises the possibility that the X chromosome
has turned over between lineages. Certainly, there has been enough time, as Vea and
Grimaldi's (2016) phylogeny used fossil calibration to date nodes and placed the split between
scale insects and other Sternorrhyncha in the Permian.
677
Either way, within the scale insects, the X chromosome has been largely conserved between X
sex determination species, with the exception of *G. tillandsiae*. Our data suggest there has been
an X-autosome fusion in the lineage leading to this species, as it is syntenic to both the X of *N.*
texana and one of the autosomes. Consistent with this scenario, we note that Brown et al.
observed another ortheziid, *Praelongorthezia praelonga*, had the same diploid chromosome
count between males and females (1958), which could occur in an X system with a neo-Y
remnant from the X-autosome fusion.
685
More broadly, we recover two losses of X chromosomal sex determination as the simplest
scenario: once in the family Monophlebidae and again in the group leading to the paternal
genome elimination clade. In fact, this scenario also holds as most parsimonious in the other
recent phylogenies (Vea and Grimaldi 2016; Song et al. 2025), even with the slight differences
in tree topology and taxon sampling, so this finding is quite robust. What is novel is the ability to
track genetic and genomic changes across these transitions, which stand in contrast to better
studied turnover events, in which one sex chromosome is supplanted for another (Vicoso and
Bachtrog 2015; Jeffries et al. 2018). In both scale insect turnovers, the X sex chromosome has
been replaced by genome-wide haplodiploidy, but in very different forms, as we explore below.
695
696   *Leaving X for hermaphroditism*
697
Starting with the Monophlebidae, some members of this group still employ X chromosomes,
such as *Drosicha* spp. (Yokogawa and Yahara 2009) and, most likely, *C. castanopsis*. The
others in the tribe Iceryini, represented by the genera *Icerya* and *Crypticerya* in our analyses,
are fully diploid simultaneous hermaphrodites (Hughes-Schrader 1925; Hughes□Schrader
1963; Kokilamani et al. 2014). These individuals are phenotypically female in all ways except for
the presence of ancillary sperm-producing tissue in their gonads (Johnston 1912). It has been
proposed that the sperm producing cells are actually a distinct, self-perpetuating cell lineage
(Royer 1975); if true, it is perhaps more correct to think of hermaphrodites as individuals with a
diploid somatic and ovarian female genotype with a distinct, selfish sperm cell line. Adding to
the complexity, in the best studied species, *Icerya purchasi*, rare pure males have also been
observed (Royer and Delavaul 1974; Royer 1975; Kim et al. 2011). There is evidence that these
males, when present, still mate with hermaphrodites and genetically contribute to offspring
(Mongue et al. 2021), making the species an androdioecious reproductive system (Pannell
2002). At present it is unclear how common androdioecy is among the hermaphroditic
Monophlebidae (Mongue et al. 2021), but, uniqueness of this system notwithstanding, the most
directly pertinent feature is that males are fully haploid (Hughes-Schrader 1925; Kokilamani et

al. 2014). So, at the genomic level, sex determination (hermaphrodite or male) presents as
genome-wide haplodiploidy.
716
This transition to this lifestyle is associated with a reduction in karyotype, as all hermaphroditic
species studied so far have a mere n = 2 autosomes (Hughes-Schrader 1925;
Hughes☐Schrader 1963; Hughes-Schrader and Monahan 1966). Based on our analyses, this
reduction karyotype actually precedes the change in sex determination, as the more distantly
related *N. texana* has n = X + 8 autosomes and the closer relatives to hermaphrodites, *C.*
castanopsis and *D. contrahens*, have n = X + 2 autosomes. Despite the large difference in
chromosome count, the pattern of rearrangement appears to be relatively straightforward. The X
is conserved between the *N. texana* and *D. contrahens*, while seven of the eight autosomes of
N. texana essentially map entirely to one of the two autosomes of *D. contrahens*, suggesting
whole-chromosome fusion events explain most of the difference. Between *D. contrahens* and *C.*
castanopsis the X is still conserved but there has been some additional autosomal
rearrangement, namely a swap between the ends of the two autosomes. The rearrangements
that span the loss of X sex determination between *C. castanopsis* and *I. purchasi* are not nearly
as clean, however.
731
The X chromosome shows orthologous hits only to a portion of the larger of the two *I. purchasi*
autosomes but these appear scattered between autosomal orthologs on that chromosome.
Genes from two autosomes of *C. castanopsis* have been rearranged both within and between
the autosomes of *I. purchasi*. Other asexual lineages are known to exhibit elevated rates of
genome rearrangements, often enabled by mobilization of transposable elements (Ferreira de
Carvalho et al. 2016), and such a scenario may have occurred here as well. Indeed between the
I. purchasi and the two *Crypticerya* species, rearrangements both within and between
chromosomes have continued at an elevated rate and these genomes are large (>1Gb) and
have the high proportions of repetitive sequence masked in our annotation analyses. That said,
the sexually-producing *N. texana* had the highest proportion of repetitive sequence in our
analysis and a ~950Mb genome, so large, repeat-rich genomes also predate the transition to
hermaphrodites.
744
To get a more granular sense of changes associated with sex determination turnover, we also
performed orthology analyses, searching for conserved X-linked orthologs that were lost from
the genomes of the hermaphrodites, as well as genes conserved in the hermaphrodites but
missing from the genomes of the X sex determination species. Starting with the former, we
found only eight X-linked orthologs missing from all three hermaphrodite species, including: a
serine protease stubble-like, a digestive cysteine proteinase-like, a tumor protein p63 regulated
gene-like ortholog, and an ortholog of CCN family member 1. Stubble proteins are hypothesized
to be secreted signal proteins in another true bug (Bao et al. 2014); cysteine proteinases
canonically play roles in metabolism (Goulet et al. 2008), and p63 related genes have a suite of
functions from histone interaction to senescence (Sadu Murari et al. 2025). Proteins in the CCN
family have functions in neuronal development and gastrulation in vertebrate models (Katsube
et al. 2009), but are less well characterized in invertebrates (Hu et al. 2019). In summary, none
of these is an obvious candidate for sex determination. It could be that they perform non-

canonical functions in scale insects, but it is also possible that sex determination turnover has
occurred in a similar way to that proposed for like-for-like turnovers, through addition rather than
subtraction of genetic elements (van Doorn and Kirkpatrick 2007). Indeed, we found over 600
orthogroups conserved across the hermaphrodites but missing from the X sex determination
species. A majority of these had no database hits, as expected of true novelty. This number
could be inflated relative to the paternal genome comparison below for the simple reason that
we searched for conservation across fewer hermaphrodite species than paternal genome
elimination species, but it is also consistent with the overall more dramatic change during sex
determination turnover here. Further sequencing of Monophlebidae can hopefully narrow down
a more tractable set of candidate genes for more extensive study.
768
Leaving X for paternal genome elimination
The second loss of X chromosomal sex determination gave rise to the paternal genome
elimination clade of scale insects. Based on our recovered phylogenetic relationships, the
different forms of paternal genome elimination found across scale insects (reviewed in Gavrilov
2007) represent elaborations from a single origin rather than independent evolutions. Based on
data from early-diverging paternal genome elimination species, we infer that the ancestral type
of paternal genome elimination is lecanoid, in which males keep but do not express paternal
chromosomes in the soma (de la Filia et al. 2021) and only truly eliminate them during
spermatogenesis (Hughes-Schrader 1948; Nur 1980). This sequence of paternal genome
elimination evolution makes intuitive sense. All types of scale insect paternal genome
elimination are united by haploid male expression and elimination of paternal chromosomes
from sperm, but in the Comstockiella type some paternal chromosomes are lost from the soma
and under the diaspidid type, the full paternal haplotype is discarded early in embryonic
development (Brown 1965). In other words, in the earliest form of paternal genome elimination,
all paternal chromosomes are maintained in the soma but in later evolving types, some or all
paternal chromosomes are eliminated early on in development. Early molecular work suggested
that there have been multiple independent transitions in type of paternal genome elimination
within armored scales (Morse and Normark 2006), but we lack the taxon sampling of this group
to confirm or refute this specific result. We are, however, able to track the fate of the X
chromosome across this transition.
789
In sharp contrast to the loss of the X in Monophlebidae, the X chromosome is almost entirely
intact in the early diverging paternal genome elimination species, starting with the crapemyrtle
bark scale, *Acanthococcus lagerstroemiae*. Indeed, while there have been some autosome-
autosome rearrangements between these two species, the overall karyotype has not changed
drastically (n = 10 vs. n = 9) and at least four chromosomes, including the former-X, are largely
intact across the transition. Within the paternal genome elimination clade, complex
rearrangements appear even less common. The former-X in particular does not undergo any
major changes until the soft scale *Toumeyella liriodendri*, in which it splits into two
chromosomes along with most other chromosomes as part of a large increase in chromosome
number. Only in the family Pseudococcidae, the true mealybugs, has there been substantial
rearrangement to the chromosomes, including former-X-linked orthologs, which are scattered
across all five autosomes.

802
In keeping with the modest chromosomal changes, we recovered far fewer differences at the
level of individual gene conservation or novelty. Indeed, there were only two reliably X-linked
orthologs lost in the transition to paternal genome elimination, and fewer than 100 orthogroups
gained. The two lost orthologs were a PITH domain-containing protein and CCN family member
1. Intriguingly, the latter was also identified as lost from the hermaphrodite group. This pattern
could hint at a shared molecular mechanism across the two losses or be mere coincidence. As
discussed above,there is currently no evidence for orthologs of this protein playing a sex
determining role in other insects, though there is little work on invertebrates. The ortholog ID
came from a BLAST hit to the whitefly *Bemesia tabaci*, a haplodiploid hemipteran pest
(Blackman and Cahill 1998). A targeted BLAST of the *A. pisum* annotation recovered two
similarly annotated orthologs, but both are located on autosomes. In other words, an ortholog of
this protein was conserved across another loss of X sex determination and is not X-linked in
other Sternorrhyncha, so its status as a sex determination gene in scale insects is dubious. Still,
as a singular target for further study, it is quite tractable.
817
Turning to orthogroups gained in the transition, many of these have no currently defined
function, so their significance to sex determination and reproduction remains to be seen. In
contrast to the orthogroups gained in the hermaphrodite lineage, far fewer had no BLAST hits at
all. Part of this can be attributed to the number of species compared, but it is also worth noting
that mealybugs are by far the best-studied scale insects, so there is more genetic data for them
(de la Filia et al. 2021; Liesenfelt et al. 2025b; Mongue et al. 2025; Vea et al. 2025). While most
genes have yet to be characterized, these putatively unique genes make an intriguing set of
targets for future study with the rise of functional genomics techniques like RNAi in mealybugs
(Khan et al. 2018; Bansal and Fernandez 2025). For now, we can conclude that like in
Monophlebidae, sex determination turnover is not associated with any appreciable gene loss.
This pattern is also visible in the distribution of sex-biased genes across the genome, as
discussed below.
830
831 *State of the X before turnover and the drivers of its change*
832
A study on other independently evolved paternal genome elimination systems (Herbette and
Ross 2023) has suggested that transitions from X sex determination may be easier if a larger
portion of the genome is X-linked (Haig 1993). Previous research has shown a lower
chromosome count is associated with the evolution of paternal genome elimination in mites
(Blackmon et al. 2015) and more recent research has shown an association between a larger X
and paternal genome elimination in fungus gnats and springtails, in which males are somatically
diploid with the exception of the X and paternal chromosomes are only eliminated in the
germline (Anderson et al. 2022). At first glance, this scenario seems plausible with the
hermaphroditic scales; the number of chromosomes decreases in closer relatives to the
hermaphrodites. However, the reduction in karyotype is not associated with an increase in X-
linked genes, as *C. castanopsis* has a roughly comparable, but slightly lower proportion of X-
linked genes than *N. texana*. The pattern is even less convincing for the paternal genome
elimination scale insects, which did not see a substantial chromosome number change across

the transition, nor is the former-X a particularly large or gene-rich chromosome. So what
distinguishes turnover in scale insects from other systems?
848
To better understand the dynamics of sex determination turnover, we further characterized the
X chromosome of *N. texana.* First, we looked for evidence of how the X chromosome is
regulated in scale insects. Briefly, because the autosomal genome is diploid in both sexes but
the X is diploid in females and haploid in males, gene expression levels may need to be
compensated in males (Disteche 2012; Gu and Walters 2017). We found that the haploid X in
males expresses genes at the same level as the diploid autosomes; in other words, the dosage
is compensated, as seen in other X sex chromosome Hemiptera (Jaquiéry et al. 2013; Pal and
Vicoso 2015). This simple observation has two implications. To start, under haplodiploidy, the
autosomes and former-X have the same ploidy, so the gene dosage is the same across the
genome. Second, thanks to this new parity, the former-X may have evolved to be down-
regulated to match the expression levels of the now single-copy autosomes in males. If so
though, it would require changes in regulation only on the former-X.
861
Next, we explored the sex-biased expression profiles of genes on the X and autosomes. Sex
chromosome theory offers two distinct predictions for such genes. In general, because of the
ploidy difference between the sexes, the X chromosome spends more time over the generations
in females than in males, and is predicted to accumulate female-biased genes (Klein et al.
2021). However, because genes on the X expressed in males are expressed in a haploid state,
they are exposed to more efficient selection (Charlesworth et al. 1987), so male-biased genes
may accumulate on the X, especially if their fitness effects are recessive (Rice 1984). In
practice, the former outcome seems more common, as X chromosomes are often feminized
(Meisel et al. 2012; Allen et al. 2013; Albritton et al. 2014), and only rarely observed to be
masculinized (Jaquiéry et al. 2013; Robert B. Baird et al. 2025). Our observations from *N.*
texana fall into the latter set: a masculinized X chromosome relative to autosomes. It is worth
noting that both of the other examples occur in systems with unusual transmission of the
chromosomes (Jaquiéry et al. 2013; Baird et al. 2023), which differ from the presumed
Mendelian inheritance patterns of *N. texana.*
876
It may simply be that male-benefitting alleles are, on average, more recessive than female-
benefitting alleles in this species, as Rice (1984) would predict, but there may be other
contributing factors as well. Chiefly, sex-biased gene expression is extreme across the entire
genome, with even the autosomes of *N. texana* having a majority of female-biased genes based
on our analyses. This pattern itself is not unprecedented, as other insects with extreme
morphological sexual dimorphism also have highly dimorphic patterns of gene expression
(Mongue et al. 2025; Mongue et al. 2025), but it may complicate selective dynamics. If sexual
dimorphism is high and more genes are female-biased in expression (a proxy for fitness
benefit), then many male-biased alleles will be in close proximity to sexually-antagonistic loci by
chance. This could hinder the fixation of genes with sex-limited or sexually-antagonistic fitness
effects if they end up locked in inversions together (McAllester and Pool 2025) or even merely in
close proximity and in linkage disequilibrium (Patten et al. 2010). This potential problem should
be present across all chromosomes, but the X sex chromosome could offer some reprieve.

Because of its haploid expression in males, the X should offer more efficient selection on male-
biased alleles than any other part of the genome (Charlesworth et al. 1987). Under these
conditions, the X may be the best place in the genome for selection on male-biased traits.
Additional gene expression analyses will be required to test this prediction across scale insects
and other taxa, as this logic should hold for any highly sexually dimorphic system with sex
chromosomes.
896
Regardless of the cause of the gene expression patterns above, they help explain the loss of X
sex determination. If sex determination system turnover is driven by sexual conflict between
chromosomes (van Doorn and Kirkpatrick 2007) and haplodiploidy represents a state in which
the whole genome is biased towards females (Blackmon et al. 2015; Klein et al. 2021; Hitchcock
et al. 2022), then it stands to reason that its evolution would be more favored in systems with a
preponderance of female-biased genes distributed throughout the genome. In other words, it
may be that sex determination turnover in scale insects was driven, or at least enabled, by
extreme sexual dimorphism, which is a common feature across this superfamily (Vea et al.
2016). Once this transition occurred, however, there would be no further pressure for a
preferential accumulation of sex-biased alleles in one part of the genome in particular, as every
chromosome has the same transmission dynamics. Fitting with this prediction, the biased
distribution of genes does appear to persist on the former-X of *E. pela*, with a continued excess
of male-biased genes and relative dearth of female-biased genes. Only later, in *T. liriodendri*,
does this pattern break down as the former-Xs undergo more rearrangement and the
composition of sex-biased genes evens out. Future expression analyses across other X
chromosome scale insects and paternal genome elimination species will be necessary to
confirm the robustness of this pattern.
914
915 *Other potential drivers of change*
916
One final consideration is that scale insects, like other phloem-feeding Hemiptera, employ
endosymbiotic organisms, often bacteria, to provision essential nutrients (Baumann 2005).
These symbionts are transmitted from mothers to offspring but are purged from males before
adulthood (Kono et al. 2008). In other words, males are an evolutionary dead end for their
symbionts. Because of this transmission dynamic, symbionts could increase their evolutionary
fitness by skewing sex ratios towards females (Ross et al. 2012b). Indeed, consider that
hermaphroditic monophlebids are phenotypically female (with the addition of testis tissue,
Johnston 1912) and pass on endosymbiotic bacteria to their hermaphroditic offspring
(Szklarzewicz et al. 2020). Though the exact developmental sequence remains unclear, the
sperm-producing portion of the ovotestis gonad of hermaphrodites appears to arise from
supernumerary sperm nuclei that persist in the embryonic cytoplasm (Royer 1975). Royer
(1975) also observed that the endosymbiotic bacteria physically associate with these sperm
nuclei, as if supporting them. Thus, the endosymbiotic bacteria appear to have both plausible
selective pressure and opportunity to capture sex determination in Monophlebidae. In fact, this
hypothesis was first suggested some 20 years ago (Normark 2004). Similar arguments apply to
paternal genome elimination species, in which sex determination of embryos appears to be
labile, even across an individual's lifespan (Ross et al. 2011). We leave deeper exploration of

symbionts for a future study, but our new sequencing data offers new chances to characterize
symbionts in previously unstudied scale insects.
936

Conclusions

We generated 17 assemblies for previously unsequenced scale insects to explore the change in sex determination across this group. We found two transitions away from X chromosome sex determination, once for simultaneous hermaphroditism and once for paternal genome elimination haplodiploidy. These two transitions took very different forms at the genomic level, with the former associated with a reduction in karyotype and extensive rearrangements and the latter transition involving minimal change to the X chromosome at first. We found few to no individual X-linked genes lost in these transitions, suggesting that the genetic machinery for sex determination was supplanted rather than lost outright. The one exception, an ortholog of CCN family 1 proteins, does not appear crucial to sex determination in other Hemiptera but makes a suitable target for future study. Finally, in characterizing the X chromosome before the transition, we found a surprisingly masculinized chromosome compared to an overall feminized genome. We suggest that extreme sexual dimorphism and widespread sexual conflict across the genome may be a driving force of sex determination turnover in scale insects. We hope these new results and data are useful for the study of scale insects in both basic and applied settings, and more generally, that they encourage other studies of sex determination turnovers across non-model systems.

Acknowledgements

We are grateful to Douglass Miller and Zachary Lahey for their constructive comments on earlier drafts of the manuscript. We thank the following contributors for supplying invaluable material: Douglass Miller and Barbara Denno for *Crypticerya rileyi, Orthezia annae*, *Puto cupressi*, *Puto decorosus*, and *Stomacoccus platani*, Mark Zenoble for *Crypticerya genistae*, Elena Oey for the *Puto* sp., Edward Ruden for *Pityococcus rugulosus*, Xinyi Zheng for *Neogreenia osmanthus*, and Lyle Buss for *Eumargarodes laingi*. The Florida Department of Agriculture and Consumer Services, Division of Plant Industry (FDACS-DPI) supported this work. These analyses were enabled by the infrastructure of the University of Florida Information Technology Research Computing (UFIT-RC)'s HiPerGator computing cluster.

References

Albritton, S. E., A. L. Kranz, P. Rao, M. Kramer, C. Dieterich, and S. Ercan. 2014. Sex-biased

gene expression and evolution of the X chromosome in nematodes. Genetics 197:865–

883.

Allen, S. L., R. Bonduriansky, and S. F. Chenoweth. 2013. The genomic distribution of sex-

biased genes in *Drosophila serrata*: X chromosome demasculinization, feminization, and

hyperexpression in both sexes. Genome Biol. Evol. 5:1986–1994.

Anderson, N., K. S. Jaron, C. N. Hodson, M. B. Couger, J. Ševčík, B. Weinstein, S. Pirro, L.

Ross, and S. W. Roy. 2022. Gene-rich X chromosomes implicate intragenomic conflict in

the evolution of bizarre genetic systems. Proc. Natl. Acad. Sci. 119:e2122580119.

National Academy of Sciences.

Bachtrog, D., J. E. Mank, C. L. Peichel, M. Kirkpatrick, S. P. Otto, T.-L. Ashman, M. W. Hahn, J.

Kitano, I. Mayrose, and R. Ming. 2014. Sex determination: why so many ways of doing

it? PLoS Biol 12:e1001899. Public Library of Science.

Baird, R. B., J. M. Urban, A. J. Mongue, K. S. Jaron, C. N. Hodson, M. Grewoldt, S. H. Martin,

and L. Ross. 2023. Recent evolution of a maternally acting sex-determining supergene

in a fly with single-sex broods. Mol. Biol. Evol. 40:msad148.

Bansal, R., and E. Fernandez. 2025. Injection, soaking, and oral delivery systems induce RNA

interference☐mediated silencing in the mealybug, Ferrisia gilli Gullan (Hemiptera:

Pseudococcidae). Pest Manag. Sci. Wiley Online Library.

Bao, Y.-Y., X. Qin, B. Yu, L.-B. Chen, Z.-C. Wang, and C.-X. Zhang. 2014. Genomic insights

into the serine protease gene family and expression profile analysis in the planthopper,

Nilaparvata lugens. BMC Genomics 15:507. Springer.

Baumann, P. 2005. Biology of bacteriocyte-associated endosymbionts of plant sap-sucking

insects. Annu. Rev. Microbiol. 59:155–189.

Blackman, R., and M. Cahill. 1998. The karyotype of Bemisia tabaci (Hemiptera: Aleyrodidae).

Bull. Entomol. Res. 88:213–215. Cambridge University Press.

Blackmon, H., N. B. Hardy, and L. Ross. 2015. The evolutionary dynamics of haplodiploidy:

genome architecture and haploid viability. Evolution 69:2971–2978. Blackwell Publishing

Inc Malden, USA.

Bray, N. L., H. Pimentel, P. Melsted, and L. Pachter. 2016. Near-optimal probabilistic RNA-seq

quantification. Nat. Biotechnol. 34:525–527. Nature Publishing Group.

Brown, S. 1965. Chromosomal survey of the armored and palm scale insects (Coccoidea:

Diaspididae and Phoenicococcidae). Hilgardia 36:189–294. University of California,

Agriculture and Natural Resources.

Charlesworth, B., J. A. Coyne, and N. H. Barton. 1987. The relative rates of evolution of sex

chromosomes and autosomes. Am. Nat. 130:113–146.

Cheng, H., G. T. Concepcion, X. Feng, H. Zhang, and H. Li. 2021. Haplotype-resolved de novo

assembly using phased assembly graphs with hifiasm. Nat. Methods 18:170–175.

Nature Publishing Group US New York.

Danecek, P., J. K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M. O. Pollard, A. Whitwham, T.

Keane, S. A. McCarthy, and R. M. Davies. 2021. Twelve years of SAMtools and

BCFtools. Gigascience 10:giab008. Oxford University Press.

Darwin, C. 1871. The descent of man and selection in relation to sex.

de la Filia, A. G., A. J. Mongue, J. Dorrens, H. Lemon, D. R. Laetsch, and L. Ross. 2021. Males

That Silence Their Father's Genes: Genomic Imprinting of a Complete Haploid Genome.

Mol. Biol. Evol. 38:2566–2581.

Ding, W.-F., X.-F. Ling, Q. Lu, W.-W. Wang, X. Zhang, Y. Feng, X.-M. Chen, and H. Chen.

2022. Identification of the key pathways and genes involved in the wax biosynthesis of

the Chinese white wax scale insect (Ericerus pela Chavannes) by integrated weighted

gene coexpression network analysis. Genes 13:1364. MDPI.

Disteche, C. M. 2012. Dosage Compensation of the Sex Chromosomes. Annu. Rev. Genet.

46:537–560.

Durand, N. C., J. T. Robinson, M. S. Shamim, I. Machol, J. P. Mesirov, E. S. Lander, and E. L.

Aiden. 2016. Juicebox provides a visualization system for Hi-C contact maps with

unlimited zoom. Cell Syst. 3:99–101. Elsevier.

Ekblom, R., and J. Galindo. 2011. Applications of next generation sequencing in molecular

ecology of non-model organisms. Heredity 107:1–15. Nature Publishing Group.

Emms, D. M., and S. Kelly. 2019. OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome Biol. 20:238. Springer.

Ferreira de Carvalho, J., V. de Jager, T. P. van Gurp, N. C. Wagemaker, and K. J. Verhoeven.

2016. Recent and dynamic transposable elements contribute to genomic divergence

under asexuality. BMC Genomics 17:884. Springer.

Flynn, J. M., R. Hubley, C. Goubert, J. Rosen, A. G. Clark, C. Feschotte, and A. F. Smit. 2020.

RepeatModeler2 for automated genomic discovery of transposable element families.

Proc. Natl. Acad. Sci. 117:9451–9457. National Acad Sciences.

Foldi, I. 2009. Archaeococcoid scale insects (Hemiptera: Coccoidea) from the tropical high

mountains of the Andean Cordillera, South America. Zootaxa 2300:1–38.

Foldi, I. 1991. The wax glands in scale insects: comparative ultrastructure, secretion, function

and evolution (Homoptera: Coccoidea). Pp. 163–188 *in*. Taylor & Francis.

Fraïsse, C., M. A. Picard, and B. Vicoso. 2017. The deep conservation of the Lepidoptera Z

chromosome suggests a non-canonical origin of the W. Nat. Commun. 8:1486.

García Morales, M., B. D. Denno, D. R. Miller, G. L. Miller, Y. Ben-Dov, and N. B. Hardy. 2016.

ScaleNet: a literature-based model of scale insect biology and systematics. Database

2016:bav118. Oxford University Press.

Gavrilov, I. A. 2007. A catalog of chromosome numbers and genetic systems of scale insects

(Homoptera: Coccinea) of the world. Isr. J. Entomol. 37:1–45.

Gavrilov-Zimin, I. A. 2018. Ontogenesis, morphology and higher classification of archaeococcids

(Homoptera: Coccinea: Orthezioidea). Zoosystematica Ross. 1–264. Федеральное

государственное бюджетное учреждение науки Зоологический ….

Goulet, M.-C., C. Dallaire, L.-P. Vaillancourt, M. Khalf, A. M. Badri, A. Preradov, M.-O. Duceppe,

C. Goulet, C. Cloutier, and D. Michaud. 2008. Tailoring the specificity of a plant cystatin

toward herbivorous insect digestive cysteine proteases by single mutations at positively

selected amino acid sites. Plant Physiol. 146:1010–1019. American Society of Plant

Biologists.

Gu, L., and J. R. Walters. 2017. Evolution of Sex Chromosome Dosage Compensation in

Animals: A Beautiful Theory, Undermined by Facts and Bedeviled by Details. Genome

Biol. Evol. 9:2461–2476.

Guindon, S., J.-F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk, and O. Gascuel. 2010. New

algorithms and methods to estimate maximum-likelihood phylogenies: assessing the

performance of PhyML 3.0. Syst. Biol. 59:307–321. Oxford University Press.

Gullan, P., and L. Cook. 2007. Phylogeny and higher classification of the scale insects

(Hemiptera: Sternorrhyncha: Coccoidea). Zootaxa 1668:413–425.

Haig, D. 1993. The evolution of unusual chromosomal systems in sciarid flies: intragenomic

conflict and the sex ratio. J. Evol. Biol. 6:249–261. Blackwell Publishing Ltd Oxford, UK.

Herbette, M., and L. Ross. 2023. Paternal genome elimination: patterns and mechanisms of

drive and silencing. Curr. Opin. Genet. Dev. 81:102065.

Hitchcock, T. J., A. Gardner, and L. Ross. 2022. Sexual antagonism in haplodiploids. Evolution

76:292–309. John Wiley & Sons, Ltd.

Hodgson, C., and I. Foldi. 2006. A review of the Margarodidae sensu Morrison (Hemiptera:

Coccoidea) and some related taxa based on the morphology of adult males. Zootaxa

1263:1–250.

Hodgson, C. J., and N. B. Hardy. 2013. The phylogeny of the superfamily C occoidea (H

emiptera: S ternorrhyncha) based on the morphology of extant and extinct macropterous

males. Syst. Entomol. 38:794–804. Wiley Online Library.

Hoff, K. J., A. Lomsadze, M. Borodovsky, and M. Stanke. 2019. Whole-genome annotation with

BRAKER. Pp. 65–95 *in* Gene prediction. Springer.

Hu, K., Y. Tao, J. Li, Z. Liu, X. Zhu, and Z. Wang. 2019. A comparative genomic and

phylogenetic analysis of the origin and evolution of the CCN gene family. BioMed Res.

Int. 2019:8620878. Wiley Online Library.

Huang, Y.-X., X.-S. Zhu, X.-N. Chen, X.-Y. Zheng, B.-S. Su, X.-Y. Shi, X. Wang, S.-A. Wu, H.-Y.

Hu, and J.-P. Yu. 2024. A chromosome-level genome assembly of the forestry pest

Coronaproctus castanopsis. Sci. Data 11:218. Nature Publishing Group UK London.

Hughes-Schrader, S. 1948. Cytology of coccids (Coccoidea-Homoptera). Adv. Genet. 35:127–

203.

Hughes-Schrader, S. 1925. Cytology of hermaphroditism in Icerya purchasi (Coccidae). Z. Für

Zellforsch. Mikrosk. Anat. 2:264–290. Springer.

Hughes☐Schrader, S. 1963. Hermaphroditism in an African coccid, with notes on other

margarodids (Coccoidea—Homoptera). J. Morphol. 113:173–184. Wiley Online Library.

Hughes-Schrader, S., and D. F. Monahan. 1966. Hermaphroditism in Icerya zeteki Cockerell,

and the mechanism of gonial reduction in iceryine coccids (Coccoidea: Margarodidae

Morrison). Chromosoma 20:15–31. Springer.

Jaquiéry, J., C. Rispe, D. Roze, F. Legeai, G. Le Trionnaire, S. Stoeckel, L. Mieuzet, C. Da

Silva, J. Poulain, and N. Prunier-Leterme. 2013. Masculinization of the X chromosome in

the pea aphid. PLoS Genet. 9:e1003690. Public Library of Science San Francisco, USA.

Jeffries, D. L., G. Lavanchy, R. Sermier, M. J. Sredl, I. Miura, A. Borzée, L. N. Barrow, D.

Canestrelli, P.-A. Crochet, and C. Dufresnes. 2018. A rapid rate of sex-chromosome

turnover and non-random transitions in true frogs. Nat. Commun. 9:4088. Nature

Publishing Group UK London.

Johnson, K. P., C. H. Dietrich, F. Friedrich, R. G. Beutel, B. Wipfler, R. S. Peters, J. M. Allen, M.

Petersen, A. Donath, and K. K. Walden. 2018. Phylogenomics and the evolution of

hemipteroid insects. Proc. Natl. Acad. Sci. 115:12775–12780. National Academy of

Sciences.

Johnston, C. E. 1912. The Internal Anatomy of Icerya Purchasi. Ann. Entomol. Soc. Am. 5:383–

388.

Katsube, K., K. Sakamoto, Y. Tamamura, and A. Yamaguchi. 2009. Role of CCN, a vertebrate

specific gene family, in development. Dev. Growth Differ. 51:55–67. Wiley Online

Library.

Khan, A. M., M. Ashfaq, A. A. Khan, M. T. Naseem, and S. Mansoor. 2018. Evaluation of

potential RNA☐interference☐target genes to control cotton mealybug, Phenacoccus

solenopsis (Hemiptera: Pseudococcuidae). Insect Sci. 25:778–786. Wiley Online Library.

Kim, D. H., J. Y. Yang, Y. S. Jang, K. San Choi, H. N. Hyun, and D.-S. Kim. 2011. Stage-

specific population dynamics of cottony cushion scale, Icerya purchasi (Hemiptera:

Monophlebidae), in citrus orchards in Jeju, Korea. J. Asia-Pac. Entomol. 14:305–309.

Elsevier.

Klein, K., H. Kokko, and H. Ten Brink. 2021. Disentangling verbal arguments: intralocus sexual

conflict in haplodiploids. Am. Nat. 198:678–693. The University of Chicago Press

Chicago, IL.

Kokilamani, A., S. Ramakrishna, G. Venu, and G. Venkatachalaiah. 2014. Karyotypic studies in

Margarodid Coccoid, *Icerya aegyptiaca* (Hempitera: Coccoidea). J. Cytol. Genet. 15:15–

21.

Kono, M., R. Koga, M. Shimada, and T. Fukatsu. 2008. Infection dynamics of coexisting beta-

and gammaproteobacteria in the nested endosymbiotic system of mealybugs. Appl.

Environ. Microbiol. 74:4175–4184. Am Soc Microbiol.

Kosztarab, M., and E. Watson. 1994. Phenological, Behavioral, and Biological Observations on

a Giant Scale, Neosteingelia texana Morrison, in Blacksburg, Virginia (Homoptera:

Coccinea: Margarodidae).

Kryuchkova-Mostacci, N., and M. Robinson-Rechavi. 2017. A benchmark of gene expression

tissue-specificity metrics. Brief. Bioinform. 18:205–214.

Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nat

Methods 9:357–359.

Layana, L., M. A. Toups, and B. Vicoso. 2025. Causes and consequences of sex-chromosome

turnovers in Diptera. bioRxiv 2025–10. Cold Spring Harbor Laboratory.

Li, H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34:3094–

3100. Oxford University Press.

Li, J., H. Xu, and S.-A. Wu. 2023. A new genus and species of giant mealybugs (Hemiptera:

Coccomorpha: Monophlebidae) from eastern China. Zootaxa 5254:434–442.

Liesenfelt, T., A. Markee, E. P. Demard, L. M. Diepenbrock, and A. J. Mongue. 2025a. Genome

report: Genome sequence of the hibiscus mealybug, Nipaecoccus viridis (Newstead), an

invasive pest of citrus. bioRxiv 2025–02. Cold Spring Harbor Laboratory.

Liesenfelt, T., A. Markee, E. P. Demard, L. M. Diepenbrock, and A. J. Mongue. 2025b. Genome

report: Genome sequence of the hibiscus mealybug, Nipaecoccus viridis (Newstead), an

invasive pest of citrus. G3 Genes Genomes Genet. jkaf154. Oxford University Press.

Manni, M., M. R. Berkeley, M. Seppey, F. A. Simão, and E. M. Zdobnov. 2021. BUSCO update:

novel and streamlined workflows along with broader and deeper phylogenetic coverage

for scoring of eukaryotic, prokaryotic, and viral genomes. Mol. Biol. Evol. 38:4647–4654.

Oxford University Press.

Marcais, G., and C. Kingsford. 2012. Jellyfish: A fast k-mer counter. Tutorialis E Manuais 1:1–8.

McAllester, C. S., and J. E. Pool. 2025. The potential of inversions to accumulate balanced

sexual antagonism is supported by simulations and Drosophila experiments. eLife

12:RP93338. eLife Sciences Publications Limited.

McClung, C. E. 1901. Notes on the accessory chromosome. Anat. Anz. 20:220–226.

McClung, C. E. 1902. The accessory chromosome—sex determinant? Biol. Bull. 3:43–84.

Marine Biological Laboratory.

Meisel, R. P., J. H. Malone, and A. G. Clark. 2012. Disentangling the relationship between sex-

biased gene expression and X-linkage. Genome Res. 22:1255–1265.

Minh, B. Q., H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. Von Haeseler,

and R. Lanfear. 2020. IQ-TREE 2: new models and efficient methods for phylogenetic

inference in the genomic era. Mol. Biol. Evol. 37:1530–1534. Oxford University Press.

Mongue, A. J., and R. B. Baird. 2024. Genetic drift drives faster-Z evolution in the salmon louse
Lepeophtheirus salmonis. Evolution qpae090.

Mongue, A. J., M. E. Hansen, and J. R. Walters. 2022. Support for faster and more adaptive Z
chromosome evolution in two divergent lepidopteran lineages*. Evolution 76:332–345.
John Wiley & Sons, Ltd.

Mongue, A. J., A. Markee, E. Grebler, T. Liesenfelt, and E. C. Powell. 2024. Genome report:
Genome sequence of tuliptree scale, Toumeyella liriodendri (Gmelin), an ornamental
pest insect. G3 Genes Genomes Genet. jkae231. Oxford University Press US.

Mongue, A. J., S. Michaelides, O. Coombe, A. Tena, D.-S. Kim, B. B. Normark, A. Gardner, M.
S. Hoddle, and L. Ross. 2021. Sex, males, and hermaphrodites in the scale insect Icerya
purchasi*. Evolution 75:2972–2983. John Wiley & Sons, Ltd.

Mongue, A. J., P. Nguyen, A. Volenikova, and J. R. Walters. 2017. Neo-sex chromosomes in
the Monarch butterfly, Danaus plexippus. G3 GenesGenomesGenetics 7:3281–3294.

Mongue, A. J., T. E. Woodman, H. Marshall, A. Garber, J. C. Franco, J. P. McCutcheon, and L.
Ross. 2025. Contrasting Evolutionary Trajectories Under Paternal Genome Elimination
in Male and Female Citrus Mealybugs. Mol. Ecol. e17826. Wiley Online Library.

Mongue, Andrew J., Johnson, Thomas W., Liesenfelt, Tracy, Amanda Markee, Nguyen, Petr,
and Hospodářská, Monika. 2025. Insights from the lack of an enigmatic trait:
monomorphic sperm in evergreen bagworm moths and the evolution of sperm
dimorphism. BioRxiv, doi: https://doi.org/10.64898/2025.12.05.692592.

Morse, G. E., and B. B. Normark. 2006. A molecular phylogenetic study of armoured scale
insects (Hemiptera: Diaspididae). Syst. Entomol. 31:338–349. Wiley Online Library.

Normark, B. B. 2004. Haplodiploidy as an outcome of coevolution between male□killing
cytoplasmic elements and their hosts. Evolution 58:790–798. Blackwell Publishing Ltd
Oxford, UK.

Núñez, O. 1962. Cytology of Collembola. Nature 194:946–947. Nature Publishing Group.

Nur, U. 1980. Evolution of unusual chromosome systems in scale insects (Coccoidea:

Homoptera). Insect Cytogenet. 97–117. Blackwell Scientific Publications.

Nur, U. 1966. Nonreplication of heterochromatic chromosomes in a mealy bug Planococcus citri

(Coccoidea - Homoptera). Chromosoma 19:439–448.

Olson, M. V. 1993. The human genome project. Proc. Natl. Acad. Sci. 90:4338–4344.

Pal, A., and B. Vicoso. 2015. The X Chromosome of Hemipteran Insects: Conservation, Dosage

Compensation and Sex-Biased Expression. Genome Biol. Evol. 7:3259–3268.

Pannell, J. R. 2002. The evolution and maintenance of androdioecy. Annu. Rev. Ecol. Syst.

33:397–425. Annual Reviews 4139 El Camino Way, PO Box 10139, Palo Alto, CA

94303-0139, USA.

Patten, M. M., D. Haig, and F. Ubeda. 2010. Fitness variation due to sexual antagonism and

linkage disequilibrium. Evolution 64:3638–3642. Blackwell Publishing Inc Malden, USA.

Powell, E. C., D. R. Miller, and O. Keller. 2024. Barbenigma Powell & Miller, a bizarre new

genus and two new species of scale insects (Hemiptera: Coccomorpha: Ortheziidae)

from the United States. Zootaxa 5529:293–317.

Rambaut, A. 2010. FigTree v1.3.1. Institute of Evolutionary Biology, University of Edinburgh,

Edinburgh.

Rice, W. R. 1984. Sex chromosomes and the evolution of sexual dimorphism. Evolution

38:735–742.

Robert B. Baird, Thomas J. Hitchcock, Jan Ševčík, Katy M. Monteith, Andy Gardner, Laura

Ross, and Andrew J. Mongue. 2025. Faster adaptation but slower divergence of X

chromosomes under paternal genome elimination. Nat. Commun.

Rodríguez Delgado, C. L., P. D. Waters, C. Gilbert, T. J. Robinson, and J. A. M. Graves. 2009.

Physical mapping of the elephant X chromosome: conservation of gene order over 105

million years. Chromosome Res. 17:917–926. Springer.

Ross, L., E. J. Dealey, L. W. Beukeboom, and D. M. Shuker. 2011. Temperature, age of mating

and starvation determine the role of maternal effects on sex allocation in the mealybug

Planococcus citri. Behav. Ecol. Sociobiol. 65:909–919. Springer.

Ross, L., M. B. Langenhof, I. Pen, and D. M. Shuker. 2012a. Temporal variation in sex

allocation in the mealybug Planococcus citri: adaptation, constraint, or both? Evol. Ecol.

26:1481–1496. Springer.

Ross, L., A. J. Mongue, C. N. Hodson, and T. Schwander. 2022. Asymmetric Inheritance: The

Diversity and Evolution of Non-Mendelian Reproductive Strategies. Annu. Rev. Ecol.

Evol. Syst. 53:1–23. Annual Reviews.

Ross, L., D. M. Shuker, B. B. Normark, and I. Pen. 2012b. The role of endosymbionts in the

evolution of haploid☐male genetic systems in scale insects (Coccoidea). Ecol. Evol.

2:1071–1081. Wiley Online Library.

Royer, M. 1975. Hermaphroditism in insects. Studies on Icerya purchasi. Pp. 135–145 *in*

Intersexuality in the animal kingdom. Springer.

Royer, M. and DELAVAUL. R. 1974. DEVELOPMENT OF MALES IN ICERYA-PURCHASI,

HERMAPHRODITIC INSECT (HOMOPTERA, COCCIDAE). COMPTES RENDUS Hebd.

SEANCES Acad. Sci. Ser. D 278:2205–2208. GAUTHIER-VILLARS 120 BLVD SAINT-

GERMAIN, 75280 PARIS, FRANCE.

Sadu Murari, L. S., S. Kunkel, A. Shetty, A. Bents, A. Bhandary, and J. C. Rivera-Mulia. 2025.

p63: A Master Regulator at the Crossroads Between Development, Senescence, Aging,

and Cancer. Cells 14:43. MDPI.

Sayyari, E., and S. Mirarab. 2016. Fast Coalescent-Based Computation of Local Branch

Support from Quartet Frequencies. Mol. Biol. Evol. 33:1654–1668.

Smit, A., R. Hubley, and P. Green. 2019. 2013–2015. RepeatMasker Open-4.0.

Song, N., M. Wang, H. Tang, and Z. Dang. 2025. A phylogenetic analysis of scale insects

(Hemiptera, Coccoidea) based on genomic and transcriptomic data. J. Syst. Evol.

63:693–707. Wiley Online Library.

Szklarzewicz, T., A. Michalik, and K. Michalik. 2020. The diversity of symbiotic systems in scale

insects. Pp. 469–495 *in* Symbiosis: cellular, molecular, medical and evolutionary

aspects. Springer.

Thomson, M. B., S. D. Campbell, and A. J. Young. 2021. Ground pearls (Hemiptera:

Margarodidae) in crops and pastures: biology and options for management. Crop

Pasture Sci. CSIRO Publishing.

Unamba, C. I., A. Nag, and R. K. Sharma. 2015. Next generation sequencing technologies: the

doorway to the unexplored genomics of non-model plants. Front. Plant Sci. 6:1074.

Frontiers Media SA.

Van Doorn, G., and M. Kirkpatrick. 2007. Turnover of sex chromosomes induced by sexual

conflict. Nature 449:909–912. Nature Publishing Group UK London.

Vasimuddin, M., S. Misra, H. Li, and S. Aluru. 2019. Efficient architecture-aware acceleration of

BWA-MEM for multicore systems. Pp. 314–324 *in*. IEEE.

Vea, I. M., A. G. de la Filia, K. S. Jaron, S. E. Barlow, M. Herbette, A. J. Mongue, R. Nelson, F.

J. Ruiz-Ruano, and L. Ross. 2025. The B chromosome of Pseudococcus viburni: a

selfish chromosome that exploits whole-genome meiotic drive. Genome Biol. Evol.

17:evae257. Oxford University Press UK.

Vea, I. M., and D. A. Grimaldi. 2016. Putting scales into evolutionary time: the divergence of

major scale insect lineages (Hemiptera) predates the radiation of modern angiosperm

hosts. Sci. Rep. 6:23487. Nature Publishing Group UK London.

Vea, I. M., S. Tanaka, T. Shiotsuki, A. Jouraku, T. Tanaka, and C. Minakuchi. 2016. Differential

juvenile hormone variations in scale insect extreme sexual dimorphism. PLoS One

11:e0149459. Public Library of Science San Francisco, CA USA.

Vicoso, B. 2019. Molecular and evolutionary dynamics of animal sex-chromosome turnover.

Nat. Ecol. Evol. 3:1632–1641. Nature Publishing Group UK London.

Vicoso, B., and D. Bachtrog. 2015. Numerous Transitions of Sex Chromosomes in Diptera.

PLoS Biol. 13:e1002078.

Vicoso, B., and D. Bachtrog. 2013. Reversal of an ancient sex chromosome to an autosome in

Drosophila. Nature 499:332–335. Nature Publishing Group UK London.

Wang, Y., H. Tang, X. Wang, Y. Sun, P. V. Joseph, and A. H. Paterson. 2024. Detection of

colinear blocks and synteny and evolutionary analyses based on utilization of MCScanX.

Nat. Protoc. 19:2206–2229. Nature Publishing Group UK London.

White, M. J. D. 1973. Animal Cytology and Evolution (Cam. bridge Univ. Press, Cambridge, UK.

WU, S., and N. Nan. 2012. Neogreenia lonicera sp. nov., a new species of Margarodidae sensu

lato (Hemiptera: Coccoidea) from China, with a key to species of Neogreenia

MacGillivray and placement of the genus in the family Kuwaniidae. Zootaxa 3274:43–54.

Wysoker, A., K. Tibbetts, and T. Fennell. 2013. Picard tools version 1.90. Httppicard

Sourceforge Net Accessed 14 Dec. 2016 107:308.

Yang, Y., R. Liu, R. Qiu, Y. Zheng, W. Huang, H. Hu, Q. Ji, H. He, Y. Shang, and Y. Gong.

2015. CRL4B promotes tumorigenesis by coordinating with SUV39H1/HP1/DNMT3A in

DNA methylation-based epigenetic silencing. Oncogene 34:104–118. Nature Publishing

Group.

Yokogawa, T., and T. Yahara. 2009. Mitochondrial phylogeny certified PGL (Paternal Genome

Loss) is of single origin and haplodiploidy sensu stricto (arrhenotoky) did not evolve from

PGL in the scale insects (Hemiptera: Coccoidea). Genes Genet. Syst. 84:57–66. The

Genetics Society of Japan.

Zhang, C., M. Rabiee, E. Sayyari, and S. Mirarab. 2018. ASTRAL-III: polynomial time species

tree reconstruction from partially resolved gene trees. BMC Bioinformatics 19:15–30.

BioMed Central.

Zheng, X., J. Li, and S. Wu. 2023. Morphology of the preadult stages of Coronaproctus

castanopsis Li, Xu & Wu (Hemiptera: Coccomorpha: Monophlebidae). Zootaxa

5352:537–552.

1286 Zheng, X., and S. Wu. 2023. Morphology of all the developmental stages of Neogreenia

1287    osmanthus (Yang & Hu)(Hemiptera: Coccomorpha), and transfer of the genera

1288    Neogreenia MacGillivray and Jansenus Foldi to the family Qinococcidae. Zootaxa

1289    5244:101–122.

1290 Zhou, C., S. A. McCarthy, and R. Durbin. 2023. YaHS: yet another Hi-C scaffolding tool.

1291    Bioinformatics 39:btac808. Oxford University Press.

1297