Genomic analysis of *Klebsiella pneumoniae* causing community-acquired respiratory deaths among Zambian infants and children using targeted RNA-probe hybridization-capture metagenomics

(short title) *Klebsiella pneumoniae* fatal community-acquired pneumonia in Zambian children

Kenneth Lindstedt[1,2,3,*], Alyse Wheelock[4,*], Mulemba Samutela[5], Wasifa Kabir[6], Mwelwa Chasaya[7], Natasha Namuziya[8], Emilia Jumbe Marsden[9], Monica Kapasa[8], Chibamba Mumba[10], Bwalya Mulenga[10], Lisa Nkole[8], Rachel Pieciak[11], Victor Mudenda[10], Chilufya Chikoti[7], Benard Ngoma[7], Charles Chimoga[7], Sarah Chirwa[7], Lilian Pemba[7], Diana Nzara[7], James Lungu[7], Leah Forman[12], Edgar Simulundu[13, 14], William MacLeod[11], Crispin Moyo[7], Somwe Wa Somwe[5,8], Kathryn E. Holt[15,16], Arnfinn Sundsfjord[1,2,3], Christopher J. Gill[11]

Norwegian Centre for Detection of Antimicrobial Resistance, Department of Microbiology and Infection Control, University Hospital of North Norway, Norway

Department of Medical Biology, Faculty of Health Sciences, UiT the Arctic University of Norway, Norway

Centre for New Antibacterial Strategies, UiT the Arctic University of Norway, Norway

Section of Preventive Medicine and Epidemiology, Chobanian & Avedisian School of Medicine, Boston University, Boston, MA, USA

School of Health Sciences, University of Zambia, Lusaka, Zambia

Aquaculture and Environment Group, UiT the Arctic University of Norway, Norway

Avencion Limited, Lusaka, Zambia

University Teaching Hospital – Children's Hospital, Lusaka, Zambia

Pendleton Children's Clinic, Lusaka, Zambia

Department of Pathology, University Teaching Hospital, Lusaka, Zambia

11 Department of Global Health, Boston University School of Public Health, Boston, MA, USA

12 Center for Health Data Science (CHDS), Boston University School of Public Health, Boston,
MA, USA

13 Macha Research Trust, Choma, Zambia

14 Zambia National Public Health Institute, Lusaka, Zambia

15 Department of Infection Biology, London School of Hygiene and Tropical Medicine, London,
UK

16 Department of Infectious Diseases, School of Translational Medicine, Monash University,
Melbourne, Victoria 3004, Australia

*Co-first authors, contributed equally

Corresponding author (technical questions): Kenneth Lindstedt, MBBS, PhD, Norwegian Centre

for Detection of Antimicrobial Resistance, Department of Microbiology and Infection Control,

University Hospital of North Norway, Norway. E-mail: kenneth.w.lindstedt@uit.no

Corresponding author (study information): Alyse Wheelock, MD, Section of Preventive Medicine

and Epidemiology, Chobanian & Avedisian School of Medicine, Boston University, Boston, MA,

USA. E-mail: alyse.wheelock@gmail.com

Abstract

Klebsiella pneumoniae (Kp) is a leading cause of neonatal and infant deaths in sub-Saharan

Africa and frequently associated with antimicrobial resistance. Previously, we identified Kp as a

major cause of fatal community-associated lower respiratory infections among infants and

children under five years in Lusaka, Zambia, using postmortem tissue sampling and pathogen

specific multiplex qPCR. In this follow-up study, we employed a novel culture-independent RNA-

probe hybridization-capture metagenomic sequencing approach, targeting Kp pan-genome core

and accessory genes, to perform in-depth genomic analysis of Kp from eleven post-mortem lung

biopsy samples from seven of these children. Analysis detected Kp in all cases except one, which

identified *Klebsiella quasipneumoniae* subspecies *similipneumoniae*. Core-genome multi-locus

sequence typing (cgMLST) revealed six clonal groups (CG607, CG1123, CG10072, CG280,

CG3648, and CG10344) belonging to five sublineages (SL607, SL17, SL280, SL37, and

SL10072), with perfect concordance between paired samples from the same case. Two infants

sampled the same month harbored SL607 lineages sharing 621 out of 629 cgMLST alleles,

suggesting clonal spread. Kp capsule (K) loci were detected in all but one case and included

potential vaccine targets KL25, KL23, and KL122. Antimicrobial resistance genes were

widespread among samples, particularly encoding resistance toward aminoglycosides, β-

lactams, sulphonamides, tetracyclines, and trimethoprim. Extended spectrum β-lactamases were

identified in four cases, three of which were $bla_{\text{CTX-M-15}}$. The acquired Kp sideophore yersiniabactin

(lineage *ybt14*) was identified in both cases associated with SL607, and the acquired siderophore

aerobactin (lineage *iuc5*) was identified in one of these, suggesting possible convergence of

antimicrobial resistance and hypervirulence. The detection of Kp with extensive antimicrobial

resistance causing fatal community acquired pneumonia signals a deeply concerning

epidemiologic shift from a largely nosocomial pathogen. This calls for urgent epidemiological

investigations to better understand the burden, transmission dynamics, antimicrobial resistances,

and potential vaccine targets for Kp in other community settings across sub-Saharan Africa.

Author Summary

Klebsiella pneumoniae is a major cause of infections and death among newborns and young

children, particularly in low-income countries, where it is frequently resistant to antibiotics. While

well-known as a hospital-associated pathogen, we previously showed *K. pneumoniae* is also a

leading cause of fatal community lung infections among infants and children in Lusaka, Zambia.

In this follow-on analysis, we performed deeper genetic analysis of *K. pneumoniae* detected from

the cluster of community pneumonia deaths using lung tissue samples from seven of these

children. Since traditional bacterial cultures were unavailable, we instead used a novel approach

that enriched and sequenced specific regions of the *K. pneumoniae* genome directly from the

biopsy samples without culturing bacterial isolates. We identified five different *K. pneumoniae*

genetic subtypes, known as sublineages. Two sublineages, which came from children sampled

the same month, were highly similar, suggesting clonal spread. Multiple acquired antimicrobial

resistance genes were detected across all sublineages. Acquired virulence factors, which may

cause more aggressive infections, were also detected in two cases. We also identified capsule

types previously suggested as potential vaccine targets. This study underscores the urgent need

to better understand and address the emerging burden of antibiotic-resistant *K. pneumoniae*

pneumonia and other invasive infections among infants and children in community settings in sub-

Saharan Africa.

Introduction

Historically, *Klebsiella pneumoniae* (Kp) has not been recognized as a significant cause of

community-acquired acute lower respiratory infection (ALRI) among children, as informed by

landmark epidemiologic research conducted in high-resource settings (1, 2). Previous large, multi-

center studies of ALRI etiologies in low- to- middle-income countries (LMICs) have systematically

excluded Kp from their analyses (3, 4). However, prior research suggests that nasal carriage and

subsequent ALRI from enteric Gram-negative bacilli such as Kp vary geographically, with these

pathogens featuring more prominently as causes of ALRI in warmer and more humid climates (5-

9). More recent data from the Drakenstein birth cohort in South Africa strongly implicate Kp as an

important cause of childhood community-acquired ALRI in certain settings (10). Moreover, Kp has

emerged as a frequent pathogenic cause of childhood mortality in the Child Health and Mortality

Prevention Surveillance (CHAMPS) network across sub-Saharan Africa and south Asia (11-13),

including in deaths due to pneumonia (14). While the majority (86%) of deaths investigated

through the CHAMPS network occurred in-hospital, of 329 deaths occurring in the community, 47

(14%) featured Kp in the causal chain, raising concern for possible community-acquired infection.

We previously reported on a post-mortem study investigating 106 community deaths among

children under age five in Lusaka, Zambia. Using multiplex PCR testing of lung samples collected

via minimally invasive tissue sampling (MITS) (15-17), we were concerned to identify a cluster of

community-acquired ALRI caused by Kp in 13 of 49 (27%) respiratory-related deaths.

Kp as a potential cause of childhood ALRI in community settings is alarming because 1) this

pathogen is not well-covered by current treatment guidelines (18) and 2) global molecular

epidemiology of Kp has been characterized by rising prevalence of multidrug resistant (MDR) and

hypervirulent sublineages, as well as convergence of MDR and hypervirulent phenotypes (19).

Molecular epidemiology data are sparser in sub-Saharan Africa than other regions of the world,

but evidence indicates widespread MDR among clinical Kp isolates (13, 20-22). Genotypic and

phenotypic studies have shown common to near-universal levels of extended-spectrum β-

lactamase (ESBL) carriage among Kp in African settings, and molecular studies have revealed

frequent detections of virulence factors, particularly the acquired siderophore yersiniabactin (23-

27).

Previous in-depth genomic analysis has revealed Kp phylogeny is composed of a highly diverse

population structure, consisting of hundreds of deep branching lineages (28). While acquisition of

resistance and virulence determinants is widespread throughout the phylogeny, the majority of

AMR and hypervirulence within Kp has been disproportionately driven by the expansion of a

distinct subset of globally disseminated lineages, termed 'global problem clones' (28). Monitoring

Kp genomic epidemiology is therefore of high importance to understand regional and international

patterns of clonal spread, identify high-risk lineages, and inform infection control and intervention

strategies, including vaccine development.

The finding of community deaths due to Kp ALRI in our original analysis was surprising and carried

important clinical and public health implications. We therefore sought to better characterize the

causative Kp in terms of antimicrobial resistance genes, virulence factors, lineage and capsule

serotype by sequencing Kp DNA detected from the preserved lung tissue biopsies. Due to the

unavailability of bacterial culture from these samples, we utilized culture-independent targeted

hybridization-capture metagenomic sequencing to achieve these aims.

Results

Description of cases

Our previous post-mortem surveillance study enrolled 121 deceased infants and children between

January 2020 to December 2022, with pauses in enrollment due to the COVID-19 pandemic. As

previously reported, a total of 106 cases had complete molecular and histopathologic data and

were adjudicated by the DeCoDe panel, of which 49 (46%) deaths were determined to be due to

respiratory causes (15). Among these, Kp pneumonia was the most common respiratory cause

of death, accounting for 13 (27%) of the respiratory deaths. For Kp targeted hybridization-capture

analysis, DNA was re-extracted from 16 paired samples of 8 selected cases with the strongest

molecular signal supporting Kp pneumonia. Of these, 5 samples proved insufficient for target

hybridization-capture due to low DNA concentration (<10 ng/μL) and poor-quality metrics

(260/280 ratio <1.5 or >2.2; 260/230 ratio <0.5 or >2.3). The remaining 11 samples from 7

deceased infants and children (4 paired samples and 3 individual samples) had sufficient quality

and proceeded to target metagenomic hybridization-capture sequencing (**Supp. Table 1**).

The 7 deceased infants and children whose samples underwent Kp probe hybridization-capture

analysis ranged in age from 7 days to 20 months and were enrolled in the study between February

2020 and December 2022. Based on enrollment criteria, none were hospitalized at the time of

death; the deaths all occurred at home, en route to care, or while awaiting medical evaluation at

a healthcare facility. All the deceased had a cough preceding their death, some had notable

shortness of breath, and other symptoms reported included fever, vomiting, and diarrhea (**Table**

1). From the narratives provided by family members describing the events leading up to the death,

some of the deceased had a noted prior interaction with a health facility, including prior

hospitalizations (**Supp. Table 2**). In all but one case (where the information was insufficient), the

DeCoDe panel determined that the death had been preventable, such as through earlier access

to care, earlier escalation of care, avoiding inappropriate discharge, or via infection prevention

measures during a prior hospitalization.

Table 1: Demographics of the deceased, case characteristics, *Klebsiella pneumoniae*

sublineages and clonal groups detected among 7 infants and children who died of suspected

Klebsiella pneumoniae acute lower respiratory tract infection in the community in Lusaka, Zambia.

Kp, *Klebsiella pneumoniae*; Kqs, *Klebsiella quasipneumoniae* subsp. *similipneumoniae*.

Case	Age	Sex	Month and year	Location of death	Cough/ severe cough	Non-respiratory	Species	Sub lineage	Clonal Group

			of death		(duration in days)	symptoms			
2337	2 weeks	Female	Dec 2022	En route to a facility	Yes (1)	Fever	Kp	SL607	CG607
4803	7 months	Male	Dec 2022	At home	Yes (2)	Fever, vomiting, diarrhea	Kp	SL607	CG607
4771	8 months	Male	Oct 2022	Waiting to be seen by a doctor	Yes (2)	Vomiting, diarrhea	Kqs	SL10072	CG10344
4796	17 months	Male	Nov 2022	At home	Yes (1)	Fever	Kp	SL17	CG1123
4201	7 days	Female	Feb 2020	En route to hospital or other health facility	Yes (2)	Fever, vomiting, diarrhea	Kp	SL17	CG10072
4220	18 weeks	Male	Jun 2020	En route to hospital or other health facility	Yes (7)	Fever	Kp	SL280	CG280
4774	20 months	Male	Oct 2022	Waiting to be seen by a doctor	Yes (1)	Fever	Kp	SL37	CG3648

Histopathology findings varied across participants as well as across the 6 biopsies obtained from

each of the deceased, from interstitial expansion to histologic pneumonia. Most of the 7

individuals included in this study had Kp detected by FTD Respiratory Pathogens 33 multiplex

PCR in 5 or 6 out of all 6 biopsies sent for molecular testing (**Supp. Table 3**). Most cases had

other pathogens detected by the multiplex PCR on one or more biopsies; based on the totality of

evidence (including number of detecting biopsies, qPCR cycle thresholds, host and pathogen

characteristics), these were considered not to be the primary driver of the infection. Reanalysis of

the 11 biopsy samples using the highly Kp-specific ZKIR-qPCR (29, 30) confirmed the presence

of Kp in all samples, ranging from 45.5 to 6586.4 genome copies/ng DNA (median 147.6) (**Supp.**

Table 1). A strong correlation was also observed between the ZKIR-qPCR quantified abundances

and the FTD Respiratory Pathogens 33 multiplex PCR Kp Cq values (Pearson's r = -0.94, *p* <

0.001, **Supp. Fig 1**), demonstrating high concordance between these two qPCR assays.

Hybridization-capture of *K. pneumoniae* by targeted RNA-probe metagenomics

We performed in-depth genomic analysis of Kp lineages from lung biopsy samples by employing

RNA-probe hybridization-capture metagenomics targeting Kp core and accessory genes (see

Methods). This targeted Kp hybridization-capture was able to overcome considerable residual

human DNA contamination.

Short-read sequencing of the 11 target-enriched samples produced an average of 30.3×10^6 75

bp paired-end reads per sample (range: $5.7 \times 10^6 – 51.2 \times 10^6$). Following removal of human DNA,

an average of 2.4×10^6 bacterial reads per sample remained (range: $6.4 \times 10^4 - 13.5 \times 10^6$),

indicating considerable residual human DNA contamination in samples despite targeted Kp

hybridization-capture. Of these bacterial reads, an average of 57.8% were assigned to Kp (range:

30.1% - 89.6%), with the number of on-target Kp reads correlating strongly with the abundance

of Kp in samples pre-hybridization-capture (Pearson's r = 0.91, p < 0.001, **Supp. Fig 2**) (**Supp.**

Table 4). With these limitations noted, our analysis was able to detect key genomic features of

the Kp lineages, including sublineage (SL), clonal group (CG), capsule serotype, as well as

acquired AMR genes and virulence determinants (**Table 2**).

Table 2. Key genomic features of *Klebsiella pneumoniae* sequenced from post-mortem lung

biopsies from 7 infants and children who died of suspected *Klebsiella pneumoniae* acute lower

respiratory infection in Lusaka, Zambia

	Phylogenetic Classification			Capsule	Acquired Antimicrobial Resistance										Acquired Virulence	
Sample	Species	SL	CG	K-locus	AGly	Flq	Phe	Rif	Sul	Tet	Tmt	Bla	Bla_ESBL	Bla_chr	Ybt	Iuc
2337-V2A	Kp	SL607	CG607	KL25	AAC(3)-IIe, AAC(6')-Ib-cr4*, AAC(6')-Iz , aadA16, aadA5, APH(6)-Id	AAC(6')-Ib-cr4*, QnrB17	-	arr-3	sul2	tetD	dfrA27	TEM-239	CTX-M-15	SHV-1	ybt 14; ICEKp12	
4803-V1A	Kp	SL607	CG607	KL25	AAC(3)-IIe, AAC(6')-Ib-cr4*, aadA16, aadA5, APH(6)-Id	AAC(6')-Ib-cr4*, QnrB17	-	arr-3	sul2	tetA, tetD	dfrA14, dfrA27	TEM-239	CTX-M-15	SHV-1	ybt 14; ICEKp12	iuc 5
4803-V2B	Kp	SL607	CG607	KL55	AAC(3)-IIe, AAC(3)-IId, AAC(6')-Ib-cr4*, aadA16, aadA5, APH(6)-Id	AAC(6')-Ib-cr4*	-	-	sul2	tetA, tetB	dfrA17	TEM-239, OXA-1	CTX-M-55	-	-	iuc 5
4201-V2B	Kp	SL17	CG10098	KL128	APH(6)-Id		-	-	sul2	tetB	dfrA8, dfrA14	TEM-239	-	SHV-155	-	-
4201-V2C	Kp	SL17	CG10098	KL128	aadA, aadA5, APH(6)-Id		-	-	sul2	tetB	dfrA8, dfrA14	TEM-239	-	SHV-97	-	-
4796-V2C	Kp	SL17	CG1123	KL122	AAC(6')-Ib-cr4*, aadA16, APH(6)-Id	AAC(6')-Ib-cr4*, QnrB6	-	arr-3	sul2	tetD	dfrA27	-	TEM-116	SHV-155	-	-
4771-V2A	Kqs	10072	CG10344	KL46	aadA, APH(6)-Id, AAC(6')-Ib7		-	-	sul2	tetB, tetD	dfrA14	SHV-217, TEM-239	-	OKP-B-22	-	-
4771-V2B	Kqs	10072	CG10344	KL46	APH(6)-Id		-	-	sul2	-	dfrA14	-	-	-	-	-
4774-V1A	Kp	SL37	CG3648	KL23	AAC(6')-Ib-cr4*, AAC(3)-IId	AAC(6')-Ib-cr4*	-	-	-	-	-	-	-	SHV-85	-	-
4774-V1C	Kp	SL37	CG3648	-	AAC(6')-Ib-cr4*	AAC(6')-Ib-cr4*	-	-	-	-	-	-	-	-	-	-

| 4220-V1A | | Kp | SL280 | CG280 | KL23 | AAC(3)-IIe, AAC(6')-Ib-cr4*, APH(3'')-Ib, APH(6)-Id | AAC(6')-Ib-cr4*, QnrB1 | catA1, catB3 | - | - | tetA, tetD | dfrA14 | OXA-1, TEM-239 | CTX-M-15 | SHV-27 | | - | - |

199 Kp = *Klebsiella pneumoniae*, Kqs = *Klebsiella quasipneumoniae* subspecies *similipneumoniae*, SL = sublineage, CG =

200 clonal group, AGly = aminoglycoside, Flq = fluoroquinolone, Phe = phenicols, Rif = rifampicin, Sul = sulfomamides, Tet

201 = tetracycline, Tmt = trimethoprim, Bla = β-lactamase, Bla_ESBL = extrended spectrum β-lactamase, Bla_chr =

202 chromosomal β-lactamase, Ybt = yersiniabactin, Iuc = aerobactin. *bifunctional ARG: catalyzes acetylation of both

203 fluoroquinolones and aminoglycosides

205 Determination of *K. pneumoniae* cgMLST and MLST from enriched samples.

Targeted RNA-probe metagenomics allowed for sequence typing according to current

conventions. SL and CG analysis of Kp from the target enriched samples was performed using

the previously described 629-gene core genome multilocus sequence typing (cgMLST) scheme

and dual bar-coding system (31, 32). A median of 627 out of 629 cgMLST genes were detected

per sample (IQR 625.5 - 628.5) (**Supp. Table 5**). When profiled against the Kp cgMLST profile

database available through BIGSdb (https://bigsdb.pasteur.fr/klebsiella/), five SLs and six CGs

were identified from the 11 enriched metagenomic samples: SL607 (CG607), SL17 (CG10098),

SL17 (CG1123), SL37 (CG3648), SL10072 (CG10344), and SL280 (CG280) (**Table 1**). cgMLST

profiles from enriched samples matched to cgMLST profiles within the database between 159 to

615 out of the 629 allelic positions (median 564) (**Supp. Table 5**). A strong correlation was

observed between the number of matching alleles and on-target Kp reads in samples post-

hybridization-capture (Spearman's rho = 0.95, $p < 0.001$) (**Supp. Fig 3**), suggesting this was a

major factor determining correct allele detection. All paired samples from the same case had

identical SLs and CGs identified, supporting accurate cgMLST reconstruction. Additionally, all

identified SLs corresponded to *K. pneumoniae sensu stricto*, except SL10072 (case 4771), which

was identified as *Klebsiella quasipneumoniae* subspecies *similipneumoniae* (Kqs). Notably, the

SL607 cgMLST allele profiles detected in samples 4803-V1A and 2237-V2A, which were taken

during the same month from different individuals, were perfectly concordant at 621 out of 629

detected cgMLST allelic positions, indicating a very high degree of phylogenetic relatedness

between the Kp strains present in these two samples (**Supp. Table 6**).

No cgMLST genes were detected in the negative control sample above our cut-off limit (≥80%

gene coverage with ≥90% sequence identity). This confirmed no significant Kp DNA

contamination of samples occurred during the hybridization-capture procedure, either through

background laboratory contamination or through cross-contamination between samples, giving

confidence in our cgMLST gene detection.

To complement our cgMLST analysis, reconstruction of sequence types (STs) using the

previously described seven-gene Kp MLST scheme was also performed (33). Although this

approach produced less consistent ST matches between sample pairs, the closest matching

MLST allele profile was still concordant with our cgMLST analysis in 7 out of 11 samples (**Supp.**

Table 7). Notably, 6 out of the 11 samples detected either 6 or 7 out of 7 MLST genes with 100%

gene coverage with 100% sequence identity. All these alleles matched their expected MLST

profiles positions based on our cgMLST analysis, except samples 4796-V2C and 4771-V1A which

had single allele mismatches. Like our cgMLST analysis, the number of detected MLST alleles

that matched with the expected MLST allele profile correlated strongly with the number of on-

target Kp reads in samples post-hybridization-capture (Spearman's rho = 0.86, $p < 0.001$, **Supp.**

Fig 4). Moreover, all samples that produced over 800,000 on-target Kp reads following

hybridization-capture produced an MLST profile matching that predicted by cgMLST analysis.

Thus, although less precise than cgMLST reconstruction, MLST analysis was able to produce

accurate ST profiles in the majority of samples, supporting our cgMLST analysis. The strong

dependence on the number of on-target Kp reads indicated MLST analysis is best suited to

samples that produce a higher yield of reads from the targeted hybridization-capture method.

Capsule type (K-locus) prediction

Prediction of the polysaccharide capsule synthesis locus (K-locus), a key potential target for future

development of Kp vaccines, was successfully performed using hybridization-capture. K-locus

prediction was performed by aligning reads to alleles corresponding to the K locus marker genes

wzi, *wzy*, and *wzx*, using allelic sequences from the Kaptive database (34). Using this approach,

K-locus marker genes were detected in all samples except 4774-V1C (**Table 2**). The K-loci

predicted from these marker genes were concordant between both paired samples from cases

4771 (SL10072, KL46) and 4201 (SL17, KL128), while unpaired samples 4220–V1A (SL280),

4796-V2C (SL17), and 2337-V2A (SL607) predicted K-loci KL23, KL122, and KL25, respectively.

A mismatch occurred between paired samples from case 4803 (SL607), with sample 4803-V1A

predicting KL25, in agreement with SL607 and KL25 detected in sample 2337-V2A, while 4803-

V2B predicted KL55. Notably, samples 4803-V2B and 4774-V1C had the two lowest number of

Kp-assigned reads of all samples, both less than 100,000 reads, which may suggest these were

below the detection threshold for accurate K-locus detection by this method.

Acquired antimicrobial resistance genes

We next analyzed samples for the presence of antimicrobial resistance genes (ARGs) known to

be acquired by Kp, which constrain effective options for clinical treatment. A high prevalence of

acquired ARGs was observed in all samples, with a median of 9 acquired ARGs from a median

of 5 resistance classes detected per sample (**Table 2**). A strong concordance was observed in

ARG profiles detected between paired samples from the same case. SL607 from samples 4803-

V1A, 4803-V2B, and 2337-V2A, as well as SL280 from sample 4220-V1A, were associated with

the highest number of acquired ARGs (n = 13, 13, 12, and 12, respectively). ARGs encoding

acquired resistance to aminoglycosides, trimethoprim, β-lactams, tetracyclines, and

sulphonamides were highly prevalent, detected in n = 11, 9, 8, 8, and 8 samples, respectively.

Transferable fluoroquinolone resistance determinants were also detected in seven samples;

AAC(6')-Ib-cr4 plus *Qnr* in four samples and *AAC(6')-Ib-cr4* only in three samples.

Within the β-lactamase class, the globally disseminated extended-spectrum β-lactamase (ESBL),

bla$_{CTX-M-15}$, was detected in three samples; 2337-V2A and 4803-V1A associated with SL607, and

4220-V1A associated with SL280. Notably, the other sample from case 4803 associated with

SL607 (4803-V2B) had a different CTX-M allele detected, *bla*$_{CTX-M-55}$, possibly a misidentification

due to low recovered reads within this sample. Another ESBL, *bla*$_{TEM-116}$, was detected in sample

4796-V2C associated with SL17. The additional acquired broad-spectrum β-lactamases, *bla*$_{TEM-}$

$_{239}$ and *bla*$_{OXA-1}$, were detected in 7 and 2 samples, respectively.

The vast majority of Kp and Kqs lineages carry the intrinsic broad-spectrum β-lactamases *bla*$_{SHV}$

and *bla*$_{OKP}$, respectively, providing resistance to amoxicillin/ampicillin (35). This is of high clinical

importance, as amoxicillin is the empiric therapy for outpatient community acquired pneumonia in

Lusaka, Zambia (18). These β-lactamases were detected in all samples containing Kp (*bla*$_{SHV}$) or

Kqs (*bla*$_{OKP}$) except 4774-V1C, 4803-V2B, and 4771-V2B. These samples had the lowest, second

lowest, and fifth lowest number of on-target Kp reads recovered (median 62,020 reads), providing

further support for a minimum threshold of approximately 100,000 reads for reliable gene

detection. Thus, the Kp lineages associated with these cases of fatal community-acquired LRTIs

were highly resistant, with most harboring multiple ARGs affecting clinically important drug

classes.

Acquired virulence determinants

Analysis of known acquired virulence loci amongst the identified Kp lineages revealed the presence of virulence factors in two cases. This analysis showed that Kp sublineage SL607 from samples 4803-V1A and 2337-V2A were associated with the acquired siderophore yersiniabactin (*ybt*), specifically lineage *ybt* 14 which is associated with the ICEKp12 mobile element (**Table 2**). Additionally, nine out of eleven alleles matching this *ybt* lineage were also detected in the other

sample from case 4803 (4803-V2B) at ≥ 80% coverage and ≥ 99% sequence identity, highly

suggestive of presence within this sample. None of these alleles, however, reached our minimum

threshold of 50x coverage depth required for reliable detection. Similarly, samples 4774-V1A and

4774-V1C associated with SL37 had 8 and 6 of the 11 alleles matching the *ybt 9* lineage

associated with ICEKp3, detected at ≥ 80% coverage and ≥ 95% sequence identity, suggestive

of the presence of this locus within these samples, however these also did not meet our minimum

threshold of coverage depth.

Both samples from case 4803 associated with SL607 also had the acquired siderophore aerobactin (*iuc*), specifically lineage *iuc5,* detected above the minimum identity and coverage thresholds. The other major known virulence loci which can be acquired by Kp (siderophore salmochelin, hypermucoidy locus *rmp*, and genotoxin colibactin) were not detected in any samples.

Detection of possible convergence of MDR and hypervirulence in sublineage SL607 in case 4803

by the acquisition of *iuc5* is a potentially important finding with significant potential public health

implications. It was therefore important to investigate whether *iuc5* detected in this case may have

been due to contamination with *Escherichia coli*, which is also a known carrier of this *iuc* lineage.

Unique reads were assigned to *E. coli* in all 11 samples, ranging from 161 to 30896 (median:

13637, IQR: 3791 – 23003), suggesting either spurious hits during taxonomy profiling or low-level

background contamination (**Supp. Table 8**). Notably, the two samples with *iuc5* detected, 4803-

V1A and 4803-V2B, contained among the highest levels of *E. coli* reads detected, with 25012 and

22469 reads, respectively, ranking second and fourth highest overall. Thus, although detection of

iuc5 in case 4803 may suggest acquisition by Kp, resulting in a higher propensity to cause

invasive infection, spurious detection from contaminating *E. coli* could not be ruled out.

Plasmid content

Identification of multiple MDR-associated plasmid replicon markers by hybridization-capture

provided insights into potential mechanisms of ARG acquisition and dissemination among Kp

lineages. Analysis of plasmid content associated with Kp lineages identified seven different

plasmid replicon types, with a median of four detected per sample (range 0 – 8) (**Supp. Table 9**).

Detected plasmid replicon families included IncFIA, IncFIB, IncFII, IncI, IncN, IncR, and Col, with

similar replicons detected between paired samples. Of these, the commonly Kp- and MDR-

associated replicons Col(pHAD28), IncFIB(K), IncFIA(HI1), and IncR were most prevalent,

detected in 9, 5, 5, and 5 samples, respectively. Other detected plasmid replicons associated with

high-risk Kp clones and MDR carriage included IncFII(K), IncFII(pKP91), IncFII(pHN7A8),

IncColRNAI, and IncN (28, 36, 37).

Analysis of *K. pneumoniae* lineages from assembled contigs

To complement our read-based analysis, reads were assembled into longer contiguous

sequences (contigs) and Kp lineages were analyzed for MLST, serotyping, acquired ARGs, and

virulence determinants. Although contig-based analysis can have reduced detection sensitivity,

reconstructing entire gene sequences can offer improved precision and confidence of results

compared to read-based analysis alone. This analysis resulted in complete MLST profiles,

containing perfect allele matches for all seven genes, in four samples (2337-V2A, 4803-V2B,

4796-V2C, and 4220-V1A) (**Supp. Table 10**). All these samples identified the same ST lineage

as our read-based cgMLST and MLST analysis (ST607, ST607, ST17, and ST280, respectively).

Of the remaining samples, all alleles with a 100% identical match to the MLST database produced

the expected allele for that position based on our previous read-based analysis, except for one

(*pgi* in sample 4771-V2B).

K-locus prediction from contigs was also highly concordant with our read-based analysis, with the

best match K-locus agreeing with the K-locus detected from reads in almost all samples. Notably,

the K-locus predicted in SL607 from case 4803 that had a mismatch between sample pairs from

read-based analysis was concordant when performed on contigs. Both samples from this case

predicted KL25, making it also concordant with SL607 detected in sample 2337-V2A. Additionally,

paired samples 4774-V1A that detected KL23 and 4774-V1C which did not detect a K-locus from

read-based analysis had a mismatch from contig-based analysis, predicting KL15 and KL51,

respectively.

Analysis of acquired ARGs also produced highly concordant ARG profiles to our read-based

analysis, with strong agreement between sample pairs. Similar to our read-based analysis, this

demonstrated the presence of genes associated with resistance to β-lactams, aminoglycosides,

trimethoprim, sulphonamides, and tetracycline was widespread among samples, as well as

resistance to fluoroquinolones, MLS, and chloramphenicol in the majority of samples. Within the

β-lactamase class, all ESBLs detected from contig analysis were concordant with our previous

read-based analysis, with an additional detection of $bla_{CTX-M-14}$ and $bla_{CTX-M-15}$ in sample 4774-V1A

which were undetected from reads.

Finally, analysis of acquired virulence from contigs also showed good agreement with read-based

detection, with samples 2337-V2A and 4803-V1A associated with SL607 matching *ybt14* at 11

out of 11 allelic positions, while sample 4803-V1A also matched *iuc5* at four out of five allelic

positions. Additionally, the acquired siderophore salmochelin (*iro*), lineage *iro5*, was also detected

in sample 4803-V1A with two out of four allelic matches.

Discussion

This analysis of a series of Kp pneumonia cases among children who died in the community in

Lusaka, Zambia, revealed several previously identified Kp genetic lineages that had been

observed in NICU settings and other environments. Notably, the sequencing revealed high rates

of AMR, global problem clones SL17 and SL37 (36, 38), multiple virulence factors, and multiple

capsular types. Two of the cases from the same month were both found to have SL607

sublineages that were highly phylogenetically similar, suggestive of clonal spread. Given that the

majority of the deceased had no apparent contact with the hospital setting, the most parsimonious

explanation for this cluster is that they represent spillover of hospital isolates that are now

circulating outside the facility and causing community acquired infections. This has significant

implications as it reinforces observations that the epidemiology of Kp has evolved and is no longer

exclusively a hospital acquired pathogen. Given the lack of AMR testing, and the ubiquitous use

of amoxicillin as first line therapy for pneumonia in community settings such as Lusaka, these

findings should be viewed with significant alarm.

RNA-probe hybridization-capture metagenomic sequencing adds to evidence generated by our

prior MITS study implicating Kp ALRI as the cause of deaths in these infants and children. While

detection of Kp does not equate to invasive infection or cause of death, the histopathologic

evidence and PCR data across multiple (six) samples adjudicated by an expert panel strongly

supported Kp as the causative pathogen in these deaths. These findings, though limited by a

small sample size, add to mounting evidence of a community burden of severe Kp, including as

a cause of ALRI, in parts of sub-Saharan Africa and other LMIC settings. Of note, this study did

not collect sufficient information to prove that the site of acquisition occurred in the community;

narrative verbal autopsies did not mention any such hospital contact in the majority of cases.

If reproduced within larger studies, the implications of a shift in the epidemiology of Kp towards

causing severe and fatal pediatric ALRI in the community should be viewed with extreme concern;

this is especially true in resource-limited settings, where ALRI treatment is algorithmic and

empiric, and almost never guided by pathogen diagnostics, let alone antimicrobial resistance

testing. Current WHO guidelines for childhood pneumonia treatment at health facilities

recommend amoxicillin for pneumonia without warning signs and ampicillin with gentamicin in

severe pneumonia (18). Due to the chromosomally encoded broad-spectrum β-lactamase bla_{SHV},

Kp is intrinsically resistant to amoxicillin/ampicillin (35). Among the 7 Kp ALRI deaths we

investigated, aminoglycoside resistance genes were universal. Extended-spectrum β-lactamase,

fluoroquinolone, and trimethoprim-sulphamethoxazole resistance determinants were also

prevalent in our study. This leaves the carbapenem class, which is rarely available even in tertiary

care settings in Lusaka, although notably, acquired resistance to this class has been detected

recently within Kp isolated from Zambian hospitals (39, 40).

Several genomic findings bear highlighting. While most of the sublineages identified between

children were distinct, two of the cases from the same month contained highly phylogenetically

similar SL607 sublineages which shared 621 out of 629 cgMLST alleles, likely representing clonal

spread. The biopsies associated with these related sublineages came from a neonate of 2 weeks

and a 7-month-old infant, whose verbal autopsy described a hospital admission the previous

week. Further information that could point to a common transmission source is not available. In

contrast, the SL17 sublineages detected from two separate cases were less closely

phylogenetically related, belonging to two different CGs, and thus were unlikely to represent a

direct epidemiologic link.

Regarding the specific SLs identified in this study, SL607, SL17, SL10072, SL37, and SL280,

each has been previously identified as known MDR-associated sublineages, and most have been

reported to be important causes of hospital-acquired infections and outbreaks among neonates

(27, 38, 41, 42). SL17 in particular is recognized as a MDR global problem clone that is prevalent

throughout sub-Saharan Africa and has been reported as a cause of neonatal infections both

within hospitals and the community (38, 43). SL37 is also a global problem clone that was

identified among the most prominent Kp lineages responsible for neonatal sepsis in low- and

middle-income countries within the BARNARDS study (23, 36). One detected sublineage,

SL10072, was additionally found to belong to the species *K. quasipneumoniae subsp.*

similipneumoniae, and encoded MDR. *K. quasipneumoniae* is increasingly recognized as a cause

of invasive infection capable of acquiring MDR, including in sub-Saharan Africa (44, 45),

supporting a potential need for increased surveillance of this species.

All SLs identified in this study, with the exception of SL10072, were also identified in a recent

hospital-based study of Kp isolates from Malawi (27). Among these overlapping SLs, all except

SL17 from case 4201 had matching K-loci with those predicted in our study and exhibited highly

similar profiles of acquired ARGs. These findings suggest that the lineages identified in our cohort

may form part of a broader circulating Kp population spanning both community and hospital

settings across sub-Saharan Africa. Notably however, only a single sublineage (SL280)

overlapped with a recent hospital-based study of Kp isolates from Zambia, which also had a

matching K-locus (KL23) (46). While this limited overlap could indicate differences between

community- and hospital-associated Kp populations within Zambia, it may also reflect temporal

differences in sampling between studies, with several years separating sample collection resulting

in shifts in the circulating lineage composition.

Another key finding was the detection of the acquired siderophores *ybt14* associated with SL607

in cases 2337 and 4803, as well as *iuc5* in case 4803. Analysis of all ST607 isolates available

through PathogenWatch via Kleborate (n = 39, downloaded July 2025) (47) demonstrated that

66.7% of these were associated with carriage of *ybt14*, consistent with our findings. Notably,

however, none of these isolates were associated with *iuc5* carriage. Acquisition of these

siderophores may suggest convergence of MDR genes and hypervirulence factors within this

lineage. Converged MDR-hypervirulent lineages may cause highly aggressive infections with

limited treatment options that many primary and secondary health settings in limited-resource

settings are unequipped to address. This finding therefore places the community death associated

with this case within a well-documented and concerning trend of high mortality related to

converged MDR-hypervirulent lineages of Kp, which is being observed globally, including other

countries across sub-Saharan Africa (27, 48).

Serotyping of lineages causing community-acquired pneumonia may be an important contribution

to choosing capsule targets for vaccine development, after further study in larger populations (49).

Five different K-loci were identified within this study, three of which (KL25, KL23, and KL122)

were present among the 30 most prevalent identified globally as potential vaccine targets in a

recent metanalysis of 1930 Kp neonatal blood isolates from 13 countries, with KL25 ranking as

the third most prevalent (49). Although our study consisted of only a small sample size, this

nevertheless suggests these K loci may be important targets for vaccine development within the

community as well as the hospital setting. There is interest in development of Kp vaccines for the

prevention of neonatal sepsis through maternal vaccination as well as for other high-risk groups

(50). Importantly, however, assuming a duration of effect of a maternal vaccination candidate of

three months, only 2 of the 7 infants and children would have been protected by a hypothetical

maternal vaccine in this small study of ALRI deaths in the community. We believe our findings

call for larger and more systematic studies of Kp causing severe community ALRI in infants and

children to inform vaccine target profiles and policy.

Plasmids are major vehicles through which Kp acquires and disseminates ARGs and virulence

determinants via horizontal gene transfer. MDR-carrying plasmids can circulate among Kp

lineages and even transfer to other species, facilitating ARG dissemination and contributing to

AMR spread and difficult-to-control outbreaks (51, 52). In this study, several clinically important

plasmid replicons were detected, including FIB(K), IncFII(K), Inc(R), ColRNAI, IncFII(pKP91),

IncFIAHI1, IncN, IncFII(pHN7A8), and Col(pHAD28). These replicons are frequently associated

with MDR plasmids and high-risk global Kp clones (28, 36, 37). Although detection of the same

plasmid replicon across multiple Kp sublineages suggests possible horizontal gene transfer

between these lineages, further investigation using long-read sequencing would be needed to

confirm this.

Several data points support the validity of the genomic analysis generated through this novel

application of RNA-probe hybridization-capture of Kp from lung tissue samples. Kp C_q values from

the FTD-32 multiplex PCR correlated highly to a singleplex Kp qPCR and to the on-target Kp

reads from RNA-probe hybridization-capture sequencing. We observed high concordance in

detected SLs, CGs, K-loci, ARGs, and acquired virulence determinants between paired samples

obtained from the same deceased individual. Previously identified SL and K-locus associations

were detected, including the association between SL280 and KL23, SL607 and KL25, SL17 and

KL122, as well as SL37 and KL23 (27, 41, 42, 46). Additionally, SLs from our study that

overlapped with strains reported in other recent sub-Saharan African studies in Zambia and

Malawi showed highly similar ARG profiles, including acquired aminoglycoside, β-lactamase,

fluoroquinolone, sulphonamide, tetracycline, and trimethoprim resistance genes (27, 46). Our

results thus demonstrate this method allows detailed analysis of genetic epidemiology including

STs (MLST and cgMLST), serotyping, acquired AMR and virulence of Kp lineages from tissue

samples. RNA-probe hybridization-capture may be useful for other settings, including post-

mortem surveillance studies, where genetic epidemiology investigation is warranted but cultures

are not available. The costs, probe-target development needs, and inability to perform phenotypic

testing, however, may limit its broader applicability.

This study was not without limitations. First, the small sample size limits the generalizability of the

sequencing findings. Larger, representative studies incorporating genetic analyses of Kp causing

severe pediatric ALRI in the community are urgently needed. Second, determining the pathogenic

cause in ALRI is complex and the sequencing findings in our study do not remove uncertainty

from the true cause of death for the infants and children in this study. To incorporate multiple

modalities of evidence into this determination, we relied on the collective clinical reasoning of a

DeCoDe panel of experts. The biopsies that underwent Kp sequencing in this study had multiple

pieces of evidence in support, first, of a respiratory cause of death and, second, of Kp as the

causative pathogen. However, some cases contained multiple conflicting pieces of evidence, for

example one case (**Table 1**), where the same three virulent pathogens were detected on multiple

biopsies, leading the DeCoDe panel to give a designation of moderate certainty for the diagnosis

of Kp ALRI.

The RNA-probe hybridization-capture technique also had limitations. High human DNA

contamination despite hybridization-capture resulted in low Kp reads in several samples, which

limited the resolution of lineage typing and detection of ARGs. While sufficient for analysis, higher

coverage would have given confidence to some of the lower coverage hits. In addition, the method

may have benefited from a second round of hybridization-capture as with other high sensitivity

RNA-probe hybridization-capture approaches. Moreover, while the acquired ARGs and virulence

determinants we detected frequently associate with Kp, especially within sub-Saharan Africa, we

could not definitively attribute these to Kp over other species potentially present at low

abundances within this study. For example, *iuc5* which was detected in case 4803, is also

commonly associated with *E. coli*, of which reads were detected in all samples. Although it is

possible these *E. coli* reads may have arisen from misclassification during taxonomy profiling,

due to shared homology between Kp and *E. coli*, they may also suggest background

contamination, possibly due to increased gut barrier permeability post-mortem. Given these

uncertainties, future culture-based isolation and whole genome sequencing studies would be

required to confirm the association of these genes with the detected Kp lineages.

Conclusions

This small study characterizing Kp genomes from Zambian children who died of suspected Kp

ALRI in the community warrants further study into the prevalence of severe and fatal Kp ALRI and

the underlying genomic epidemiology. This study investigated Kp sequenced from preserved lung

biopsies in 7 suspected Kp ALRI deaths identified through a post-mortem MITS study of under-5

community respiratory deaths in Lusaka, Zambia. ALRI remains the second most common cause

of under-5 mortality globally and growing evidence suggesting Kp as a contributing pathogen

raises concerns around current clinical management approaches. Across the samples, we

detected several distinct sublineages of global importance, genes encoding acquired resistance

to important antimicrobial classes in all cases and acquired virulence factors in two cases. Given

the critical implications for prevention and treatment of severe under-5 ALRI in the community,

additional, larger scale genomic epidemiology studies should be carried out in similar populations.

Methods

This study used a target hybridization-capture metagenomic approach to sequence Kp from metagenomic DNA extracted from post-mortem lung biopsies of infants and children who died outside of hospital settings in Lusaka, Zambia. The methods of the original post-mortem study have been previously detailed (15). Briefly, enrollments occurred at the University Teaching Hospital morgue, which serves the city of Lusaka. All next-of-kin of the deceased approached for the study were offered grief counseling, and, if interested in study participation, provided informed consent followed by a verbal autopsy. The body of the deceased was then brought to the autopsy theater, the thorax was cleaned with ethanol and iodine, and six minimally invasive tissue samples (MITS) were obtained from upper, middle, and lower lung zones from the left and right side along the axillary line. Molecular testing and histopathology was performed on specimens from each of the six lung zones sampled. The molecular testing performed was a 32-respiratory pathogen PCR panel (FTD-33 kit, Fast-Track Diagnostics). For histopathology, lung samples were fixed in neutral buffered formalin, followed by paraffin embedding, microtomy, hematoxylin and eosin staining, and finally, reviewed by two experienced pathologists.

A panel adjudication approach modeled after the Child Health and Mortality Prevention Surveillance Determination of Cause of Death (DeCoDe) Process was used to determine the causes of death in this post-mortem study, based on the totality of subjective and objective data. The panel convened for this study included 7 clinicians with infectious diseases and pediatrics experience practicing in Lusaka. The panel met in person and reviewed case dossiers, including representative photographs of the histopathology for each of the deceased, to achieve a consensus determination of whether the death was due to respiratory or non-respiratory causes, the cause of death, and whether the death was preventable or not. For the present sequencing study, we selected 8 cases with the strongest molecular data (at least two biopsies with Kp detected at qPCR cycle thresholds of less than 30) from among the 13 deceased infants and

children adjudicated to have died from Kp pneumonia. We attempted to perform target

hybridization-capture sequencing on two separate lung biopsy samples from each of these select

cases.

DNA extraction, quality control, and *K. pneumoniae*-specific ZKIR-qPCR analysis

DNA was extracted using the InviSorb® Spin Universal Kit (Invitek Diagnostics, Germany)

following the manufacturer's protocol for tissue biopsy samples in final elution volume of 60 μL

DNA concentration. The concentration and quality of the extracted DNA from each of the lung

biopsy samples was measured by Quibit 3.0 fluorometer using dsDNA Narrow Range Assay

(Thermo Fisher Scientific, Waltham, USA) and Nanodrop 1000 Spectrophotometer (Thermo

Fisher Scientific), respectively. All DNA samples which passed quality control assessment

underwent analysis using the highly Kp sensitive and specific ZKIR-qPCR assay as previously

described (29, 30). Briefly, 2.5 ng of 10 ng/μL DNA (25 ng total) was used as input for each 25 μL

qPCR reaction, with reaction mixture and cycling conditions as described (30). All samples were

assayed in triplicate using an Applied Biosystems 7500 Real-Time PCR System (Thermo Fisher

Scientific), and criteria used for positive Kp detection were as described previously. Kp genome

copy number was calculated from the standard curve using 7500 Real-Time PCR Analysis

Software v2.3 (Applied Biosystems, Life Technologies, Waltham, USA) as described (30).

Due to the very high detection sensitivity of the RNA-probe target hybridization-capture method

employed in this study, a blank negative control was used to account for background/cross-

contamination during the hybridization-capture process which consisted of DNA elution buffer

only. This blank sample had no detectable DNA by qubit analysis and was negative for Kp by the

ZKIR-qPCR.

Library construction, targeted RNA probe hybridization-capture, and sequencing

Hybridization-capture of Kp from lung biopsy metagenomic samples was performed by SureSelect XT HS2 targeted RNA probe hybridisation-capture (Agilent, Santa Clara, USA). A custom panel of 145,787 120bp RNA probes (total size: 2.635 Mbp) was designed targeting core and acquired genes from the *Klebsiella pneumoniae* pan-genome, including cgMLST, MLST, K- and O- loci, ARGs, virulence loci, and plasmid replicons (**Supp. Table 11**). Probes were designed with at least 90% sequence homology to the reference sequences with 1–2x tiling density, by an in-house developed algorithm by Agilent Technologies. For regions with ambiguous bases, among all possible sequences, representative probes with a 90% base homology threshold were selected. The algorithm reduces the number of relevant baits to ensure cost efficiency. The final bait sets comprised baits (Design ID S3491653 and 61512) of 120bp RNA probes (total size: 2.635 Mbp).

Sequencing library construction and targeted RNA probe hybridization-capture of samples was performed as per the SureSelect XT HS2 DNA Kit Library Preparation (+/- MBC)/Fast-Hyb Target Hybridization-capture/Post-hybridization-capture Pooling Workflow protocol (Agilent; catalog nr G9982A) with modifications based on the Utilization of Agilent SureSelect Target Hybridization-capture for Whole Genome Sequencing of Viruses and Bacteria protocol (Agilent; catalog nr PR7000-2035). Briefly, 150 - 200 ng DNA was used as input for library construction. Samples with <150 ng DNA were bulked with human genomic DNA (Promega, Madison, USA). Following enzymatic fragmentation, end repair and A-tailing, libraries underwent PCR amplification and indexing, followed by purification on AMPure Beads. 1000 ng of DNA library were hybridized with 1:10 diluted probes for 90 mins according to the protocol. After hybridization, a hybridization-capture step using Streptavidin beads, followed by several steps of washing were done according

to the protocol. A final post-hybridization-capture PCR was performed with 19 cycles for samples

with > 200 Kp genome copies/ng DNA and 21 cycles for samples with < 200 Kp genome copies/ng

DNA.

All target enriched samples were sequenced on the Illumina NextSeq 500 platform at 75 bp paired

end reads. Processing of FASTQ files and removal of residual human DNA was performed as

previously described (30).

Analysis of Target-enriched samples

Taxonomy profiling

Taxonomic profiling, including total bacterial read count as well as Kp read count and relative

abundance estimates were performed using Kraken2 v2.1.2 and Bracken v2.6.1 with

MiniKrakenDB_8GB v202003 (53, 54). Quantification of unique Kp and *E. coli* reads in samples

was performed using Centrifuge v1.0.4 using the standard p_compressed+h+v database (55).

cgMLST and MLST analysis

Kp lineage analysis was performed using both the previously described 7-gene multi-locus

sequence typing (MLST) scheme and the core genome multi-locus sequence typing (cgMLST)

scheme consisting of 629 core genome genes (32, 33). Both MLST and cgMLST alleles were

identified from sequenced reads using the aligner KMA v1.4.18 (56) against either the MLST or

cgMLST allelic nucleotide sequence databases downloaded from BIGsdb

(https://bigsdb.pasteur.fr/klebsiella/). Detected MLST/cgMLST alleles were filtered to include only

those with ≥80% gene coverage and ≥90% sequence identity. The allele corresponding to each

MLST/cgMLST gene detected above this threshold that had the highest ConClave score

(accumulated alignment score) was then used to construct the final Kp MLST/cgMLST profile for

each sample. The resulting detected MLST/cgMLST profiles were compared to the Kp

MLST/cgMLST allelic profile database available at BIGSdb to identify the closest matching

ST/cgST lineage.

K-locus prediction

Within our custom RNA-probe panel were probes targeting the K-locus genes *wzi*, *wzx*, and *wzy*.

To predict the Kp K-locus from our samples we used KMA v1.4.18 (56) to align sequence reads

to the K-locus gene nucleotide sequence database downloaded from Kaptive v3.0.0 (34) using

the command 'kaptive extract kpsc_k --ffn'. Detected alleles were filtered to include only those

with ≥80% gene coverage and ≥90% sequence identity. The allele for each K-locus gene with the

highest ConClave score was then retained. K-locus was called if at least two out of the three

detected K-locus genes (*wzi*, *wzx*, and *wzy*) corresponded to the same K-locus.

Detection of acquired ARGs, virulence loci, and plasmid replicons

Detection of ARGs, virulence loci, and plasmid replicons was performed using the read alignment

tool KMA v1.4.18 (56). For ARGs, reads were aligned to the CARD v3.0.8 database. For virulence

loci, allelic sequences were downloaded from BIGSdb (https://bigsdb.pasteur.fr/klebsiella/) and

ST profiles for yersinabactin (ybt), colibactin (clb), aerobactin (iuc), salmochelin (iro) and rmpADC

were analyzed using schemes described previously (57-59). For detection of plasmid replicon

sequences, the PlasmidFinder database was used. Presence of a gene was called if it was

detected with ≥ 90% sequence identity, ≥ 80% gene coverage, and ≥ 50x coverage depth.

Detection of an acquired virulence loci was called if ≥ 50% of the expected alleles for that loci

were detected above the minimum thresholds. Lineage was reported if ≥ 50% of the detected

alleles matched with an allelic profile when queried against the BIGSdb database.

Contig-based analysis

Paired-end reads were assembled into contigs using MegaHit v1.2.9 and --presets meta-sensitive
parameter. Assembled contigs were analyzed for ST, acquired virulence loci and ARGs via
Kleborate v3.1.2 (19) and capsule and O-antigen loci via Kaptive v3.0.0 (34) using default
parameters.

Statistical analysis and data visualization

Statistical analysis and data visualization was performed in R v4.5.0 using RStudio
2025.05.01+513.

Computational resources

Bioinformatic analysis of target enriched metagenomic samples was performed on the Norwegian
academic high-performance computing and storage services maintained by Sigma2 Norwegian
Research Infrastructure (NRIS) (http://www.sigma2.no/) (Sigma2 project number: nn9794k).

Data availability

Raw Illumina sequence reads from target enriched samples (human DNA removed) are available

on ENA, Bioproject PRJEB105599.

Disclosure Statement

All authors report no conflict of interest.

Ethics statement

Ethical oversight for the ZPRIME and LCIS studies were provided by institutional review boards

at Boston University and the University of Zambia. Written informed consent was obtained from

the deceased's family members or representatives.

Acknowledgements

The authors thank Sylvain Brisse (Institut Pasteur, France), Kelly Wyres (Monash University,

Australia), Erik Hjerde (UiT The Arctic University of Norway), and Ryan Wick (The University of

Melbourne, Australia) for assistance in RNA probe target selection and design. We thank Even

Østli and his team Agilent for their assistance in RNA-probe production and technical support with

the hybridization-capture. We thank Ruth Paulsson and Hagar Taman (the Genome Sequencing

Support Centre, Norway) for their assistance with library construction and sequencing. We also

thank Tom Stanton (Monash University, Australia) for his advice when analysing Kp K-loci.

Funding sources

The ZPRIME and LCIS studies, DeCoDe process, and RNA probe design and hybridization-

capture reagent kit costs were funded by the Bill and Melinda Gates Foundation, grant number

OPP1163027. The salary of KL, target enrichment library construction, and sequencing costs

were funded by the Trond Mohn Foundation, grant number TMF2019TMT03 and the Northern

Norway Regional Health Authority, grant numbers HNF1756-25 and HNF1589-21. AW received

salary support through National Institutes of Health 2T32HL125232-07.

Author contributions

The study was conceptualized by CG, AS, KH, AW, KL, and MS. Funding acquisition was performed by CG, AS, and KL. Participant recruitment, grief counseling, and sample acquisition were performed by BN, CC, SC, LP, DN, and JL. Adjudications were performed by NN, EJM, MK, CM, BM, LN, VM, SWS, and coordinated by MC. Lung biopsy DNA sample preparation and shipping was organized by AW and MS. RNA-probe synthesis was coordinated by KL and AW. RNA-probe hybridization capture of samples was performed by WK. DNA sequencing and library preparation was arranged by KL and WK. Data analysis was performed by KL with clinical summaries compiled by AW. KH and AS provided expertise on *Klebsiella pneumoniae* genomics and clinical aspects. AW and KL prepared the first manuscript draft. All authorships contributed with reviewing and editing the manuscript and approved the final version.

Abbreviations

ALRI = acute lower respiratory infection

AMR = antimicrobial resistance

ARG = antimicrobial resistance gene

CHAMPS = Child Health and Mortality Prevention Surveillance

CG = clonal group

cgMLST = core genome multilocus sequence typing

cgST = core genome sequence type

ESBL = Extended spectrum beta-lactamase

Kp = *Klebsiella pneumoniae*

LMIC = low- to- middle-income countries

MDR = multidrug resistant

MITS = minimally invasive tissue sampling

MLST = multilocus sequence typing

SL = sublineage

ST = sequence type

References

1.   Jain S, Williams DJ, Arnold SR, Ampofo K, Bramley AM, Reed C, et al. Community-
acquired pneumonia requiring hospitalization among U.S. children. N Engl J Med.
2015;372(9):835–45.

2.   Madhi SA, De Wals P, Grijalva CG, Grimwood K, Grossman R, Ishiwada N, et al. The
burden of childhood pneumonia in the developed world: a review of the literature. Pediatr Infect
Dis J. 2013;32(3):e119–27.

3.   Benet T, Sanchez Picot V, Messaoudi M, Chou M, Eap T, Wang J, et al. Microorganisms
Associated With Pneumonia in Children <5 Years of Age in Developing and Emerging
Countries: The GABRIEL Pneumonia Multicenter, Prospective, Case-Control Study. Clin Infect
Dis. 2017;65(4):604–12.

4.   Pneumonia Etiology Research for Child Health Study G. Causes of severe pneumonia
requiring hospital admission in children without HIV infection from Africa and Asia: the PERCH
multi-country case-control study. Lancet. 2019;394(10200):757–79.

5.   Wolf B, Gama A, Rey L, Fonseca W, Roord J, Fleer A, et al. Striking differences in the
nasopharyngeal flora of healthy Angolan, Brazilian and Dutch children less than 5 years old.
Ann Trop Paediatr. 1999;19(3):287–92.

6.   Wolf B, Rey LC, Moreira LB, Milatovic D, Fleer A, Verhoef J, et al. Carriage of gram-
negative bacilli in young Brazilian children with community-acquired pneumonia. Int J Infect Dis.
2001;5(3):155–9.

7.   Ghafoor A, Nomani NK, Ishaq Z, Zaidi SZ, Anwar F, Burney MI, et al. Diagnoses of
acute lower respiratory tract infections in children in Rawalpindi and Islamabad, Pakistan. Rev
Infect Dis. 1990;12 Suppl 8:S907–14.

8.   Ikeogu MO, Wolf B, Mathe S. Pulmonary manifestations in HIV seropositivity and
malnutrition in Zimbabwe. Arch Dis Child. 1997;76(2):124–8.

9.   El Seify MY, Fouda EM, Ibrahim HM, Fathy MM, Husseiny Ahmed AA, Khater WS, et al.
Microbial Etiology of Community-Acquired Pneumonia Among Infants and Children Admitted to
the Pediatric Hospital, Ain Shams University. Eur J Microbiol Immunol (Bp). 2016;6(3):206–14.

10.   Zar HJ, MacGinty R, Workman L, Burd T, Smith G, Myer L, et al. Klebsiella pneumoniae
Lower Respiratory Tract Infection in a South African Birth Cohort: a Longitudinal Study. Int J
Infect Dis. 2022;121:31–8.

11.   Taylor AW, Blau DM, Bassat Q, Onyango D, Kotloff KL, Arifeen SE, et al. Initial findings
from a novel population-based child mortality surveillance approach: a descriptive study. Lancet
Glob Health. 2020;8(7):e909–e19.

12.   Bassat Q, Blau DM, Ogbuanu IU, Samura S, Kaluma E, Bassey IA, et al. Causes of

Death Among Infants and Children in the Child Health and Mortality Prevention Surveillance

(CHAMPS) Network. JAMA Netw Open. 2023;6(7):e2322494.

13.   Verani JR, Blau DM, Gurley ES, Akelo V, Assefa N, Baillie V, et al. Child deaths caused

by Klebsiella pneumoniae in sub-Saharan Africa and south Asia: a secondary analysis of Child

Health and Mortality Prevention Surveillance (CHAMPS) data. Lancet Microbe. 2024;5(2):e131–

e41.

14.   Mahtab S, Blau DM, Madewell ZJ, Ogbuanu I, Ojulong J, Lako S, et al. Post-mortem

investigation of deaths due to pneumonia in children aged 1-59 months in sub-Saharan Africa

and South Asia from 2016 to 2022: an observational study. Lancet Child Adolesc Health.

2024;8(3):201–13.

15.   Wheelock A, Chasaya M, Namuziya N, Jumbe Marsden E, Kapasa M, Mumba C, et al.

Using Minimally Invasive Tissue Sampling and Determination of Cause of Death to Establish

Etiologies of Community Respiratory Deaths Among Zambian Infants and Children. J Pediatric

Infect Dis Soc. 2025;14(1).

16.   Gill CJ, Mwananyanda L, MacLeod WB, Kwenda G, Pieciak R, Mupila Z, et al. Infant

deaths from respiratory syncytial virus in Lusaka, Zambia from the ZPRIME study: a 3-year,

systematic, post-mortem surveillance project. Lancet Glob Health. 2022;10(2):e269–e77.

17.   Mwananyanda L, Gill CJ, MacLeod W, Kwenda G, Pieciak R, Mupila Z, et al. Covid-19

deaths in Africa: prospective systematic postmortem surveillance study. BMJ. 2021;372:n334.

18.   Revised WHO Classification and Treatment of Pneumonia in Children at Health

Facilities: Evidence Summaries. Geneva: The World Health Organization, ISBN: 978-92-4-

150781-3; 2014 Available from https://www.ncbi.nlm.nih.gov/books/NBK264162/.

19.   Lam MMC, Wick RR, Watts SC, Cerdeira LT, Wyres KL, Holt KE. A genomic
surveillance framework and genotyping tool for Klebsiella pneumoniae and its related species
complex. Nat Commun. 2021;12(1):4188.

20.   Thomson KM, Dyer C, Liu F, Sands K, Portal E, Carvalho MJ, et al. Effects of antibiotic
resistance, drug target attainment, bacterial pathogenicity and virulence, and antibiotic access
and affordability on outcomes in neonatal sepsis: an international microbiology and drug
evaluation prospective substudy (BARNARDS). Lancet Infect Dis. 2021;21(12):1677–88.

21.   Collaborators GAR. Global burden of bacterial antimicrobial resistance 1990-2021: a
systematic analysis with forecasts to 2050. Lancet. 2024;404(10459):1199–226.

22.   Global antibiotic resistance surveillance report 2025: WHO Global Antimicrobial
Resistance and Use Surveillance System (GLASS). Geneva: World Health Organization; 2025.
Licence: CC BY-NC-SA 3.0 IGO. ISBN 978-92-4-011633-7.

23.   Sands K, Carvalho MJ, Portal E, Thomson K, Dyer C, Akpulu C, et al. Characterization
of antimicrobial-resistant Gram-negative bacteria that cause neonatal sepsis in seven low- and
middle-income countries. Nat Microbiol. 2021;6(4):512–23.

24.   Mills RO, Dadzie I, Le-Viet T, Baker DJ, Addy HPK, Akwetey SA, et al. Genomic
diversity and antimicrobial resistance in clinical Klebsiella pneumoniae isolates from tertiary
hospitals in Southern Ghana. J Antimicrob Chemother. 2024;79(7):1529–39.

25.   Muraya A, Kyany'a C, Kiyaga S, Smith HJ, Kibet C, Martin MJ, et al. Antimicrobial
Resistance and Virulence Characteristics of Klebsiella pneumoniae Isolates in Kenya by Whole-
Genome Sequencing. Pathogens. 2022;11(5).

26.   Amulele AV, Orindi B, Ndumba M, Tigoi C, Omuoyo D, Kahindi E, et al. Genomic
epidemiology of Klebsiella pneumoniae neonatal and infant sepsis in Kenyan hospitals.
MedRxiv. 2025.

27.   Heinz E, Pearse O, Zuza A, Bilima S, Msefula C, Musicha P, et al. Longitudinal analysis
within one hospital in sub-Saharan Africa over 20 years reveals repeated replacements of
dominant clones of Klebsiella pneumoniae and stresses the importance to include temporal
patterns for vaccine design considerations. Genome Med. 2024;16(1):67.

28.   Wyres KL, Lam MMC, Holt KE. Population genomics of Klebsiella pneumoniae. Nat Rev
Microbiol. 2020;18(6):344–59.

29.   Barbier E, Rodrigues C, Depret G, Passet V, Gal L, Piveteau P, et al. The ZKIR Assay, a
Real-Time PCR Method for the Detection of Klebsiella pneumoniae and Closely Related
Species in Environmental Samples. Appl Environ Microbiol. 2020;86(7).

30.   Lindstedt K, Buczek D, Pedersen T, Hjerde E, Raffelsberger N, Suzuki Y, et al.
Detection of Klebsiella pneumoniae human gut carriage: a comparison of culture, qPCR, and
whole metagenomic sequencing methods. Gut Microbes. 2022;14(1):2118500.

31.   Hennart M, Guglielmini J, Bridel S, Maiden MCJ, Jolley KA, Criscuolo A, et al. A Dual
Barcoding Approach to Bacterial Strain Nomenclature: Genomic Taxonomy of Klebsiella
pneumoniae Strains. Mol Biol Evol. 2022;39(7).

32.   Bialek-Davenet S, Criscuolo A, Ailloud F, Passet V, Jones L, Delannoy-Vieillard AS, et
al. Genomic definition of hypervirulent and multidrug-resistant Klebsiella pneumoniae clonal
groups. Emerg Infect Dis. 2014;20(11):1812–20.

33.   Diancourt L, Passet V, Verhoef J, Grimont PA, Brisse S. Multilocus sequence typing of
Klebsiella pneumoniae nosocomial isolates. J Clin Microbiol. 2005;43(8):4178–82.

34.   Stanton TD, Hetland MAK, Lohr IH, Holt KE, Wyres KL. Fast and accurate in silico
antigen typing with Kaptive 3. Microb Genom. 2025;11(6).

35.   Holt KE, Wertheim H, Zadoks RN, Baker S, Whitehouse CA, Dance D, et al. Genomic
analysis of diversity, population structure, virulence, and antimicrobial resistance in Klebsiella

pneumoniae, an urgent threat to public health. Proc Natl Acad Sci U S A. 2015;112(27):E3574–
81.

36.   Navon-Venezia S, Kondratyeva K, Carattoli A. Klebsiella pneumoniae: a major
worldwide source and shuttle for antibiotic resistance. Fems Microbiol Rev. 2017;41(3):252–75.

37.   Spadar A, Perdigao J, Campino S, Clark TG. Large-scale genomic analysis of global
Klebsiella pneumoniae plasmids reveals multiple simultaneous clusters of carbapenem-resistant
hypervirulent strains. Genome Med. 2023;15(1):3.

38.   Hetland MAK, Hawkey J, Bernhoff E, Bakksjo RJ, Kaspersen H, Rettedal SI, et al.
Within-patient and global evolutionary dynamics of Klebsiella pneumoniae ST17. Microb
Genom. 2023;9(5).

39.   Chizimu JY, Kalungia AC, Mudenda S, Muhimba Z, Lukwesa C, Mufwambi W, et al.
Diagnostic Stewardship Trends and Antimicrobial Resistance Profiles of Bacteria Isolated in
Zambia: A Five-Year Retrospective Study (2020-2024). Antibiotics (Basel). 2025;14(11).

40.   Namukonda S, Shawa M, Siame A, Mwansa J, Mulundu G. Prevalence and antibiotic
resistance profiles of ESKAPE pathogens in the neonatal intensive care unit of the women and
newborn hospital in Lusaka, Zambia. Antimicrob Resist Infect Control. 2025;14(1):96.

41.   Peltier F, Choquet M, Decroix V, Adjide CC, Castelain S, Guiheneuf R, et al.
Characterization of a multidrug-resistant Klebsiella pneumoniae ST607-K25 clone responsible
for a nosocomial outbreak in a neonatal intensive care unit. J Med Microbiol. 2019;68(1):67–76.

42.   Castro J, Araujo D, Oliveira H, Fernandes L, Oliveira R, Brinks E, et al. Multidrug-
resistant Klebsiella pneumoniae and Klebsiella variicola isolated from patients in Portuguese
hospitals: Genomic and phenotypic characterization. The Microbe. 2024;5:100172.

43.   Henson SP, Boinett CJ, Ellington MJ, Kagia N, Mwarumba S, Nyongesa S, et al.
Molecular epidemiology of Klebsiella pneumoniae invasive infections over a decade at Kilifi
County Hospital in Kenya. Int J Med Microbiol. 2017;307(7):422–9.

44.   Brinkac LM, White R, D'Souza R, Nguyen K, Obaro SK, Fouts DE. Emergence of New
Delhi Metallo-beta-Lactamase (NDM-5) in Klebsiella quasipneumoniae from Neonates in a
Nigerian Hospital. mSphere. 2019;4(2).

45.   Mishra K, Banerjee T, Yadav G, Kumar A, Pratap A, Chaurasiya S, et al. Emergence of
drug-resistant Klebsiella pneumoniae phylogroups (K. quasipneumoniae and K. variicola)
causing human infections. Microbiol Spectr. 2025;13(9):e0019825.

46.   Phillips LT, Bates M, Coffin SE, Foster-Nyarko E, Kapasa M, Machona S, et al.
Transmission dynamics of Klebsiella pneumoniae in a neonatal intensive care unit in Zambia
before and after an infection control bundle. PLOS Glob Public Health. 2026;6(2):e0005965.

47.   Argimon S, David S, Underwood A, Abrudan M, Wheeler NE, Kekre M, et al. Rapid
Genomic Characterization and Global Surveillance of Klebsiella Using Pathogenwatch. Clin
Infect Dis. 2021;73(Suppl_4):S325–S35.

48.   Lan P, Jiang Y, Zhou J, Yu Y. A global perspective on the convergence of
hypervirulence and carbapenem resistance in Klebsiella pneumoniae. J Glob Antimicrob Resist.
2021;25:26–34.

49.   Stanton TD, Keegan SP, Abdulahi JA, Amulele AV, Bates M, Heinz E, et al. Distribution
of capsule and O types in Klebsiella pneumoniae causing neonatal sepsis in Africa and South
Asia: meta-analysis of genome-predicted serotype prevalence and potential vaccine coverage.
MedRxiv. 2025. https://doi.org/10.1101/2025.06.28.25330253.

50.   Dangor Z, Benson N, Berkley JA, Bielicki J, Bijsma MW, Broad J, et al. Vaccine value
profile for Klebsiella pneumoniae. Vaccine. 2024;42(19 Suppl 1):S125–S41.

51.   David S, Cohen V, Reuter S, Sheppard AE, Giani T, Parkhill J, et al. Integrated
chromosomal and plasmid sequence analyses reveal diverse modes of carbapenemase gene
spread among Klebsiella pneumoniae. Proc Natl Acad Sci U S A. 2020;117(40):25043–54.

52.   Wyres KL, Holt KE. Klebsiella pneumoniae as a key trafficker of drug resistance genes
from environmental to clinically important bacteria. Curr Opin Microbiol. 2018;45:131–9.

53.   Lu J, Breitwieser FP, Thielen P, Salzberg SL. Bracken: estimating species abundance in
metagenomics data. Peerj Comput Sci. 2017.

54.   Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome
Biol. 2019;20(1):257.

55.   Kim D, Song L, Breitwieser FP, Salzberg SL. Centrifuge: rapid and sensitive
classification of metagenomic sequences. Genome Res. 2016;26(12):1721–9.

56.   Clausen P, Aarestrup FM, Lund O. Rapid and precise alignment of raw reads against
redundant databases with KMA. BMC Bioinformatics. 2018;19(1):307.

57.   Lam MMC, Wyres KL, Judd LM, Wick RR, Jenney A, Brisse S, et al. Tracking key
virulence loci encoding aerobactin and salmochelin siderophore synthesis in Klebsiella
pneumoniae. Genome Med. 2018;10(1):77.

58.   Lam MMC, Salisbury SM, Treat LP, Wick RR, Judd LM, Wyres KL, et al. Genomic and
functional analysis of rmp locus variants in Klebsiella pneumoniae. Genome Med.
2025;17(1):36.

59.   Lam MMC, Wick RR, Wyres KL, Gorrie CL, Judd LM, Jenney AWJ, et al. Genetic
diversity, mobilisation and spread of the yersiniabactin-encoding mobile element ICEKp in
Klebsiella pneumoniae populations. Microb Genom. 2018;4(9).
886