1

ACE-2-like Enzymatic Activity in Anti-SARS-CoV-2 Spike Protein Monoclonal Antibodies

Yufeng Song[1, #], Frances Mehl[1, #], Tom No[1,¶], Lauren Livingston[1], Juan Sebastian

Quintero Barbosa[1], Jun Hayashi[2], Ginette Serrero[2], Pamela Schoppee Bortz[3], Jeffrey M

Wilson[3], James E Crowe Jr[4], David D Ho[5], Michael T Yin[6], Joshua Tan[7], Steven L

Zeichner[1, 8*]

[1] Department of Pediatrics, University of Virginia, Charlottesville, Virginia, USA.

[2] Precision Antibody Service, 9130 Red Branch Rd., Suite X Columbia, Maryland

21045, United States.

[3] Division of Allergy and Clinical Immunology, Department of Medicine, University of

Virginia, Charlottesville, Virginia, USA.

[4] Howard Hughes Medical Institute, Seattle, WA 98109, USA; Department of Pathology,

Microbiology, and Immunology, Vanderbilt University Medical Center, Nashville, TN

37232, USA; Department of Pediatrics, Vanderbilt University Medical Center, Nashville,

TN 37232, USA.

[5] Aaron Diamond AIDS Research Center, Columbia University Vagelos College of

Physicians and Surgeons, New York, NY 10032, USA; Division of Infectious Diseases,

Department of Medicine, Columbia University Vagelos College of Physicians and

Surgeons, New York, NY 10032, USA; Department of Microbiology and Immunology,

Columbia University Vagelos College of Physicians and Surgeons, New York, NY

10032, USA.

[6] Columbia University Vagelos College of Physicians and Surgeons, New York, New

York, USA.

[7] Antibody Biology Unit, Laboratory of Immunogenetics, National Institute of Allergy and

[8] Vaccine Research Center, National Institute of Allergy and Infectious Diseases,

National Institutes of Health, Bethesda, MD 20892, USA; Aaron Diamond AIDS

Research Center, Columbia University Vagelos College of Physicians and Surgeons,

and Department of Biochemistry and Molecular Biophysics, Columbia University, New

York, NY 10032, USA.

[9] Department of Microbiology, Immunology, and Cancer Biology, University of Virginia,

Charlottesville, Virginia, USA.

38   [#]Contributed equally to this work.

40   [¶]Present address: Department of Urology, University of California, Irvine, CA, USA.

42   *Correspondence: zeichner@virginia.edu

Abstract

Many people are affected by post-acute sequelae of COVID-19 (PASC/long COVID, LC). LC has severely affected public health. Features of LC including blood pressure dysregulation, coagulopathies, hyperinflammation, and neuropsychiatric complaints. Mechanisms responsible for LC pathogenesis are not clear. The receptor for SARS-CoV-2 is human angiotensin converting enzyme 2 (ACE2), which binds SARS-CoV-2 spike protein receptor-binding domain (RBD) to initiate infection. We hypothesized that some people produce anti-RBD antibodies that sufficiently resemble ACE2 structure to have ACE2-like catalytic activity. Those antibodies, ACE2-like abzymes, may contribute to LC pathogenesis. We previously showed that ACE2-like activity was associated with immunoglobulin in some people with acute and convalescent COVID-19. ACE2-like catalytic activity correlated with blood pressure changes following moderate exercise challenge in convalescents. We screened human monoclonal antibodies (mAbs) against SARS-CoV-2 spike protein from 4 sources. We identified 4 human mAbs with ACE2-like catalytic activity. The activity was not inhibited by MLN-4760, a compound that inhibits native human ACE2, nor by EDTA, unlike native ACE2, a Zinc metalloprotease, but was inhibited by an overlapping pool of Spike peptides. Enzyme kinetic studies showed that the mAbs had lower Vmax and *Km* values than ACE2. The data suggested that the antibodies cleave angiotensin II via a different mechanism than ACE2. Identification of mAbs with ACE2-like catalytic activity supports the hypothesis that antibodies induced by SARS-CoV-2 infection could help mediate the pathogenesis of COVID-19 and LC, and more generally, the hypothesis that catalytic antibodies induced by infectious agents can contribute to disease pathogenesis.

Introduction

COVID-19 and Post-Acute Sequelae of COVID-19 (PASC, or Long COVID, LC), as reviewed in [1-3], can affect up to 7% of COVID-19 patients[4]. LC patients can have many unusual symptoms such as dizziness, fatigue, malaise, "brain fog", alterations in smell or taste, abnormal movements, libido changes, and other neuropsychiatric symptoms, GI symptoms, cardiovascular complaints such as palpitations, blood pressure dysregulation, inflammation, coagulopathies, cough, and chest pain[5 6, 7]. While the clinical features, laboratory findings, and epidemiology of LC have been extensively described, clear primary causal etiologies have yet to be determined [8].

Antibodies with catalytic activity, also called "abzymes" have been known for some time [9, 10]. Investigators initially produced abzymes by generating anti-idiotypic antibodies against antibodies targeting enzymes' active sites[11, 12]. In some instances, the anti-idiotypic antibodies had sufficient resemblance to the original enzyme's active site, retaining catalytic activity similar to that of the initially targeted enzyme[13, 14]. However, further research showed that the abzymes had decreased substrate specificity and less favorable enzyme constants compared to the targeted enzymes (*e.g.,* $k$cat/Km values ~10^2 - 10^4 s^{-1}•M s^{-1} [15] vs. $k$cat/Km values of ~10^5 s^{-1}•M s^{-1})[16 12, 13, 17]. These limitations led to a decline in enthusiasm for the use of abzymes in biotechnological applications.

In an attempt to identify potential mechanisms responsible for the unusual features of COVID-19 and LC, we considered that many of these aberrant features could be viewed as consequences of pathologic derangements of proteolytic regulatory cascades, for example, blood pressure regulation (the renin-angiotensin system), inflammation (C', kallikrein-kinin)[18-20], and coagulation. For LC, the kinetics of symptom development would be consistent with antibody production, rather than the pathology directly produced by viral replication[21, 22]. While the benefits of COVID-19 vaccination vastly outweigh the risks, the fact that a very small, but still non-zero, fraction of patients who received SARS-CoV-2 vaccines had symptoms including hyperinflammation, blood pressure dysregulation, or coagulopathies [23]further suggests that, in

some cases, an antibody response could play a role in pathogenesis.

Angiotensin Converting Enzyme 2 (ACE2) on the host cell surface serves as the principal

receptor for the SARS-CoV-2 Receptor Binding Domain (RBD). Angiotensin Converting Enzyme

(ACE) catalyzes the cleavage of angiotensin I to angiotensin II, which acts as vasoconstrictor to

increase blood pressure [24]. ACE2 cleaves angiotensin II to produce angiotensin 1-7, which has

vasodilatory activity that lowers blood pressure, an activity that is counterregulatory to AC[25]. We

hypothesized that the pathogenesis of some of the unusual features of LC might be caused by

the induction of antibodies against the SARS-CoV-2 Spike protein (S) that resembled the

structure of ACE2 closely enough to have ACE2-like catalytic activity.

In our previous studies[26, 27], we found that some patients' plasma with ACE2-like activity did

not require divalent cations, unlike native ACE2, a Zinc metalloprotease. This activity co-

depleted with immunoglobulin and was inhibited by a tiled overlapping RBD peptide pool. These

findings suggested that the ACE2-like activity observed in patients' plasma differed from native

ACE2 and was the result of anti-RBD immunoglobulins exhibiting ACE2-like activity.

Additionally, ACE-2 like catalytic activity found in plasma collected from convalescent

patients correlated with changes in blood pressure following a moderate exercise challenge[26],

suggesting that the catalytic antibodies could have physiologic consequences.

However, to definitively establish that COVID-19 patients produce antibodies with catalytic

activity, it is essential to identify a pure monoclonal antibody derived from COVID-19 patients

that exhibits the hypothesized catalytic activity. Here, we survey several sets of monoclonal

antibodies (mAbs) from 4 different sources.

While we found that most mAbs lacked catalytic activity, two mAbs from one set showed

detectable catalytic activity, and two others showed borderline detectable activity. These

observations show that some patients develop antibodies with catalytic activity following SARS-

CoV-2 infection, with potential clinical consequences.

Materials and Methods

Monoclonal Antibody Collections

We received 129 purified human anti-SARS-CoV-2 RBD mAbs, 1 purified anti-SARS-CoV-2 fusion peptide (FP) mAb, 7 human anti-SARS-CoV-2 RBD mAb hybridoma supernatants, 3 irrelevant mAb hybridoma supernatants, and 1 empty hybridoma culture media as a negative control from 4 different collaborator groups (Table 1). Some of the anti-RBD antibodies were further characterized for their ability to block or enhance ACE2 binding to SARS-CoV-2 spike protein. The hybridoma culture medium is 1/3 Medium E (Stem Cell Technologies) and 2/3 DMEM + 10 % FBS.

Monoclonal Antibody for Screening

All antibodies were assessed for binding to SARS-CoV-2 RBD (with the exceptions of the negative control mAbs) as described[28-34]. The mAbs purchased from Precision Antibodies targeting the RBD (but not those from other sources) were further screened for interactions with ACE2. This screening used (spike) S and (receptor binding domain) RBD recombinant proteins with HIS tag and an ACE2 human Fc fusion protein. Ni-NTA plates were used to capture both S and RBD proteins. ACE2 binding to both S and RBD proteins was confirmed to ensure that both S and RBD retained the ability to bind ACE2. The S protein captured on Ni-NTA plates was used for the initial screening. All the clones that showed binding to S-protein were subsequently screened for specific RBD binding using RBD-captured Ni-NTA plates. ACE2 was added to the plate to see if

any of the anti-RBD antibodies blocked ACE2 binding to the S or RBD proteins captured

on the plate. The binding of ACE2 was detected using HRP-anti-human Fc.

Angiotensin II Converting Enzyme (ACE2)-like Activity Assays

For the ACE2 activity assay, 25 µg purified antibody or 50 µL hybridoma supernatant

was combined with 50 µM MCA-APK(Dnp) fluorogenic ACE2 substrate (AAT Bioquest,

Pleasanton CA. Cat. #1355) in assay buffer, 50 mM MES assay buffer (Thermo Fisher,

Waltham MA. Cat. # J62231.AK) supplemented with 300 mM NaCl and 10 µM $ZnCl_2$, in

a non-binding black bottom plate (Corning, Glendale AZ. Cat. #3991) [35, 36] with total

100µL reaction volume[37]. Empty hybridoma culture media was used as a negative

control. Fluorescence levels were measured with excitation/emission wavelengths

320/405nm and a 325 nm emission filter on the SpectraMax® M5 multi-mode microplate

reader (Molecular Devices, San Jose CA.)[36]. Measurements were taken every five

minutes for 16 hours, while the plate was kept at a constant temperature of 37°C. 5 mM

EDTA was added to the hybridoma culture supernatants to inhibit the ACE2 present in

10% FBS medium[38]. Specificity of ACE2 substrate cleavage was determined by parallel

comparisons of samples with 25 mM OMNIMMP fluorogenic control substrate (Enzo

Life Sciences, Farmingdale, NY. Cat. #MML-P127), MCA-Pro-Leu-OH, according to

previous studies [26, 27, 39].

ACE2 Substrate Peptide Cleavage Competition Inhibition by SARS-CoV-2 Spike

RBD Peptide Pools

Twelve peptides (SB-Peptide, SmartBioscience SAS, France) covering the SARS-CoV-

2 RBD and RBM were reconstituted in ddH2O or DMSO to a concentration of 10 µg/µL

(Table 2) [26, 27]. The peptides were combined in equal amounts. The ACE2 activity assay

was set up as described above, and 2.4 µL of the peptide pool was added so that there

was 2 µg of each peptide per well. 2.4 µl of peptide diluent (50% ddH$_2$O and 50%

DMSO) was added to other wells as a control.

ACE2 Substrate Peptide Cleavage Competition Inhibition by ACE2 Inhibitor

The ACE2 activity assay was run as described above, with the addition of 100 µM MLN-

4760 (Bio-Techne, Minneapolis, MN, Cat. #3345), a small-molecule ACE2 active-site

inhibitor[37, 40]. This concentration is recommended in commercial ACE2 activity kits

(Abcam, Waltham, MA. Cat. #ab273297).

Enzyme Kinetics Assay

The ACE2 activity assay was set up as described, but with substrate dilutions raging

from 12.5 µM to 200 µM. An MCA standard curve (AAT Bioquest, Pleasanton CA. Cat.

#557) was set up with 500, 250, 200, 150, 100, 50, 25, and 0 pmol MCA/well in 100 µL

assay buffer. This curve enabled the calculation of pmol of MCA-APK(Dnp) substrate

cleaved per minute.

RBD-specific Binding Assays by ImmunoCAP

Following our previous studies[26, 27], S-RBD binding was measured by using 50 µL of

hybridoma supernatant with a quantitative ImmunoCAP system applying Phadia 250

(Thermo Fisher Scientific, Waltham MA. Thermo-Fisher/Phadia). Briefly, biotinylated S-

RBD (RayBiotech, Peachtree Corners, GA) was conjugated to streptavidin-coated

ImmunoCAPs (Thermo Fisher Scientific, Waltham MA. Thermo-Fisher/Phadia).

Unconjugated streptavidin ImmunoCAP was run in parallel with its signal as a

background to be subtracted from the singles of other samples.

Human IgG ELISA

The total amount of human IgG in hybridoma supernatant was measured by quantitative

human IgG ELISA following the kit manufacturer's instructions (Abcam, Waltham, MA.

Cat. #ab100547) as described[26, 27].

Data Analysis

ACE2 substrate cleavage was monitored by measuring RFU for 16 hours with 5-minute

intervals. The RFU values were corrected for background fluorescence by subtracting

the minimum RFU value from the remaining RFU values for each well in the assay

plate. Linear regression analysis was done on the first 5-120 minutes to determine the

increase in RFU over time. This slope (RFU/min) was used as an indicator of ACE2-like

activity. A slope of 0.5 RFU/min (the slope of 0.08 ng rhACE2) was determined to be the

baseline detection level of ACE2 activity, which is two-fold greater than the lower limit of

detection, the activity of 0.04 ng rhACE2[26, 27]. Monoclonal antibodies with a slope

greater than 0.5 RFU/min were identified as positive for ACE2-like abzyme activity. For

the MCA standard curve, the average RFU over 2 hours was plotted against pmol

MCA/well. Linear regression was done to determine the number of RFU per pmol MCA.

This value allowed for the calculation of pmols substrate cleaved per minute from the

slope data. The supernatant IgG concentration was used to determine the *Kcat* of mAb

catalytic activity. All data analysis and visualization were performed with R (version

2022-06-23) using R Studio and packages including drc, ggh4x, ggplot2, patchwork,

tidyverse, readr, GGally, corrplot and stringr.

Results

We screened 129 purified human anti-RBD mAbs (Table 1) [28, 29, 31, 32, 34, 41] with

one purified broadly neutralizing mAb against SARS-CoV-2 fusion peptide[33] as an

irrelevant control (Table 1). We also screened 10 human mAb hybridoma culture

supernatants purchased from Precision Antibody, including 3 anti-RBD mAbs that

blocked ACE2 binding to RBD, 4 anti-RBD mAbs that enhanced ACE2 binding to RBD,

and 3 negative control mAbs (Table 1). 5mM EDTA was added to the hybridoma

supernatants to inhibit any native ACE2 protein in the media, following previous studies

that inhibited ACE2 in 10% FBS media[38]. A concentration of 5 mM EDTA was confirmed

to give complete inhibition of ACE2 in the hybridoma culture media serving as our

negative control (FIG 1).

We found 4 anti-RBD mAbs (AGRD013-4G9, AGRD016-2B3, AGRD016-4H2,

AGRD021-9A10), that had ACE2-like abzyme activity (FIG 2. A). Two of these

(AGRD013-4G9 and AGRD016-4H2) had previously been determined to block RBD

binding to ACE2, while the other two (AGRD016-2B3 and AGRD021-9A10) had been

determined to enhance binding of RBD to ACE2. An additional ~~997~~ mAbs had catalytic

activity higher than the lower limit of detection, but lower than our predetermined

threshold for significant ACE2 activity and thus were classified as negative for ACE2-

like activity (FIG 2. B). All negative control mAbs had slopes below the lower limit (FIG

2. B).

Antibody concentrations in hybridoma culture supernatants were measured by a

total human IgG ELISA[26, 27]. The antibody levels ranged from 16 to 28 μg/ml. Anti-RBD

binding activity in those hybridoma culture supernatants was assessed using an

ImmunoCAP assay[26, 27], with RBD antibody concentrations ranging from 8 to 13 μg/mL.

Significant correlation (R = 0.92, p = 0.0035, Pearson correlation coefficient) was

observed between total human IgG concentration and anti-RBD IgG (FIG 3), indicating

that ACE2-like activity correlates with anti-RBD antibody, consistent with our previous

findings[26, 27].

Analysis of the enzymatic activities of those mAbs, ACE2-like activity positive

samples, as well as the rhACE2 positive control, was conducted using a Michaelis–

Menten analysis[42]. IgG concentrations were determined by human IgG ELISA (FIG 3,

Table 3). The ACE2-like abzymes have lower K_ms, and ~4 $\log_{10}$ lower catalytic

efficiency (Table 3, FIG 4). The reduced catalytic efficiency (K_{cat}/K_m) is consistent with

the data obtained for other abzymes[15, 16]. Our calculated K_{cat}/K_m for rhACE2 at 37°C is

0.8 $M^{-1}s^{-1}$ higher than the reported value for rhACE2 at room temperature[43].

Substrate specificity of abzyme catalytic activity was determined by comparing

the cleavage of ACE2 substrate (MCA-APK(Dnp)) and control substrate (MCA-Pro-Leu-

OH), as in prior studies[26, 27, 44]. Abzyme catalytic activities with control substrate had

negative slopes (FIG 5), demonstrating that the abzyme activity is ACE2 substrate

specific.

To further confirm the specificity of the catalytic activity, we conducted ACE2

substrate cleavage peptide inhibition assays using 12 tiled overlapping peptides that

cover the SARS-CoV-2 RBD and RBM amino acid sequences (Table 2). The RBD

peptide pools significantly inhibited the ACE2-like activity of two mAbs, AGRD013-4G9

and AGRD016-4H2, which block RBD binding to ACE2, but had overall limited inhibition

effect on two other mAbs, AGRD016-2B3 and AGRD021-9A10, which enhance RBD

binding to ACE2 (FIG 6). The RBD peptide inhibition results show that the mAb's

catalytic activity could be inhibited by peptide competition, and that there is a difference

in the ability of the peptides to inhibit activity between mAbs which block or enhance

receptor binding[26, 27].

We also tested the effect of a small molecule ACE2 inhibitor, MLN-4760, a well-

documented inhibitor of ACE2 catalytic activity[37, 40]. MLN-4760 completely inhibits

rhACE2, consistent with previous findings[37, 40], but does not inhibit ACE2-like abzyme

activity (FIG 7). As we previously showed, the divalent cation chelator, EDTA does not

affect the mAbs' substrate cleavage activity, but inhibits native ACE2 catalytic activity,

since native ACE2 is a Zn^{2+} metalloprotease[26, 27]. These findings suggest that the mAbs

with ACE2-like activity employ different catalytic mechanisms than native ACE2.

Discussion

The COVID-19 pandemic has claimed around 7 million lives, according to the World
Health Organization as of January 2026[45], with about 65 million patients experiencing
lingering symptoms long after their initial infection[3]. Long COVID, characterized by
persistent symptomatology following an initial COVID-19 infection, includes symptoms
consistent with dysregulated proteolytic regulatory cascades[18-20] such as the
angiotensin-renin system [24].

Our previous research showed that some patients suffering from acute SARS-CoV-2
infection or convalescing from COVID-19 produced antibodies with ACE2-like catalytic
activity [26 , 27]. In this study, to further explore our hypothesis that some anti-spike
antibodies exhibit ACE2-like catalytic activity, we screened 129 anti-SARS-CoV-2
mAbs, 7 anti-spike mAb hybridoma supernatants; and 3 irrelevant mAb hybridoma
supernatants and an anti-SARS-CoV-2 FP mAb as controls (Table 1). Although most
anti-RBD mAbs exhibited no ACE2-like activity, we identified 4 anti-RBD mAb
hybridoma supernatants that had ACE2-like activity. This finding is consistent with our
hypothesis that only a small fraction of elicited RBD antibodies exhibit ACE2 substrate-
cleavage activity after infection [24, 25].

ACE2-like catalytic activity, whether native ACE2 or other metalloproteases, is inhibited
by 5 mM EDTA in the assay buffer. The ACE2-like catalytic activity of the mAbs was not
inhibited by 5 mM EDTA (FIG 1), just as we observed directly in samples from research
volunteers with both acute COVID-19 and convalescing from COVID-19[26, 27]. ACE2
substrate cleavage specificity was further confirmed by conducting assays using either
a native substrate or a negative control ACE2 substate widely used to establish

specificity in ACE2 assays (FIG 5)[26, 27, 44], and by using a SARS-CoV-2 spike peptide

pool in a competition/inhibition assay (FIG 6), which has been used to demonstrate that

the specificity of immunoglobulin-associated ACE2-like activity in research subject

samples[26, 27]. The peptide pool inhibited the substrate cleavage of two mAbs

(AGRD013-4G9 and AGRD016-4H2), which blocked ACE2 binding to the virus spike

protein, suggesting that the catalytic sites of the mAbs have sufficient resemblance to

the ACE2 active site to be inhibited by the overlapping spike-derived peptide pool.

However, two mAbs (AGRD016-2B3 and AGRD021-9A10) enhance spike protein

binding to ACE2, indicating the mAbs have distinct structural and functional features.

The four mAbs with ACE2-like activity exhibit low catalytic efficiency (Table 3 and FIG

4), a known characteristic of other abzymes[16], so the characteristics of the isolated

mAbs with ACE2-like catalytic activity fit well within those described for other abzymes.

The inability of the native ACE2-specific small-molecule inhibitor, MLN-4760, to inhibit

the ACE2-like activity of abzymes (FIG. 7) suggests that the active site structures of the

mAbs, and perhaps their catalytic mechanisms, differ from those of native ACE2[12, 13, 17],

further defining the unique features of the anti-spike mAbs.

Our study had several strengths. We screened many mAbs from four laboratories for

ACE2-like catalytic activity. Although a large fraction of the tested mAbs did not have

ACE2-like catalytic activity, we found that four isolated anti-SARS-CoV-2 spike mAbs

have ACE2-like catalytic activity. Our previous studies describing immunoglobulin-

associated ACE2-like activity in clinical samples could have had other, unappreciated

substances other than Abs responsible for the ACE2-like catalytic activity. The

identification of ACE2-like catalytic activity in mAbs uncontaminated by the many

substances present in clinical plasma samples strongly supports the existence of Abs

with ACE2-like catalytic activity.

However, our study had certain limitations. By screening pre-established mAb

collections, the mAbs were produced without clear linkage to patients with symptoms

associated with COVID-19. In particular, the mAbs were not produced from LC patients

with some of the most problematic symptoms. This prevents correlation of abzymes

characteristics with clinical symptoms. In addition, the motivation behind the production

of most of the mAbs was to obtain mAbs capable of inhibiting SARS-CoV-2 infection of

cells. Selecting for a mAb that is highly effective at blocking the virus from infecting a

cell may bias the mAb production process toward selecting mAbs with catalytic activity,

since those mAbs may not be particularly effective at blocking cell infection. In the

future, we hope to acquire and screen mAbs isolated from research subjects with

defined clinical symptoms and amplify and characterize clones that are not specifically

biased toward highly neutralizing mAbs. In patients, highly neutralizing Abs may not

dominate the total pool of Abs made in response to SARS-CoV-2 infection[46, 47].

Identified mAbs with ACE2-like catalytic activity are from hybridoma culture

supernatants, of which the mAbs are in a condition close to natural physiological

environment, while other mAbs screened in this study are purified IgGs. Nevertheless,

the ACE2-like catalytic activity of the hybridoma supernatants did not require divalent

cations (was not inhibited by EDTA), so the activity is distinct from native ACE2. The

catalytic activities of the identified abzymes are much lower than native ACE2. In future

studies, to detect these low catalytic activities it may be important to ensure that the

antibodies are maintained in an physiological intact state during isolation, since it is

conceivable that binding or infection inhibition may be less fragile properties that

catalytic activity[48-51], which could be a conceivable reason why catalytic activity was not

detected in the purified mAbs. Nevertheless, the identification of any mAbs with ACE2-

like catalytic activity supports the hypothesis underlying this study.

An essential next step involves elucidating the mechanistic properties of the abzymes'

catalytic activity. Structural analyses aimed at delineating these activities could provide

critical insight into the mechanisms of ACE2-like catalytic activity. A clear understanding

of the catalytic mechanisms of the mAbs with ACE2-like catalytic activity could enable

the development of inhibitors against that activity. Alternatively, the cells producing the

pathologic abzymes could be targeted.

Overall, our study further confirms the existence of anti-spike antibodies with ACE2-like

activity that can plausibly contribute to the pathogenesis of COVID-19 and LC. While

much work remains to be done, including the production and study of new anti-spike

mAbs not selected specifically for SARS-CoV-2 neutralizing activity, study of catalytic

activities other than ACE2 in Abs made after exposure to SARS-CoV-2 that could

potentially mediate the pathogenesis of COVID-19 and LC, studies associating different

clinical features with abzymes with particular characteristics, structural and mechanistic

studies of the abzymes themselves, and larger epidemiologic survey studies to better

determine the rates of abzyme production in persons exposed to SARS-CoV-2. There

are also longer-term translational implications. Other viruses, in addition to SARS-CoV-

2, use cell surface enzymes as their receptors [52-57]. If pathologic abzymes induced by

exposure to SARS-CoV-2 are responsible for some of the pathologic features of

COVID-19 and LC, then abzymes induced by other viruses, including viruses causing

subclinical infections, could be responsible for other disorders.

Institutional Review Board Statement: The UVA IRB approved enrollment of all

subjects and collection of specimens and related metadata (HSR #HSR13166).

Informed Consent Statement: Informed consent was obtained from the subjects in the

study.

Data Availability Statement: Data is presented in the manuscript. Raw data is

available upon request.

Acknowledgments and Funding: The study was supported by University of Virginia

internal funding, including the Pendleton Laboratory Fund for Pediatric Infectious

Disease Research, the Manning Fund for COVID-19 Research at UVA, the Ivy

Foundation, University of Virginia Summer Research Internship Program and grants

from NIAID, NIH (R01 AI176515).

Conflicts of Interest: The authors declare no conflict of interest.

Tables
414
415

416   **Table 1:** Summary of Monoclonal Antibody Characteristics Used in the Study

Source	Total Number	Form	Characteristics	Number with Abzyme Activity	Reference
Dr. James Crowe, Vanderbilt University Medical Center	2	Purified	Human Anti SARS-CoV-2 RBM	0	28, 29
Dr. Jun Hayashi, Precision Antibody	3	Supernatant	Human Anti SARS-CoV-2 Spike and RBD, block RBD binding to ACE2	2	
	4	Supernatant	Human Anti SARS-CoV-2 Spike and RBD, enhance RBD binding to ACE2	2	
	3	Supernatant	Negative Controls	0	
Dr. David Ho, Columbia University	39	Purified	Human Anti SARS-CoV-2 RBD	0	31, 41, 58

| Dr. Joshua Tan at NIH | 88 | Purified | Human Anti SARS-CoV-2 RBD | 0 | 34 |
| | 1 | Purified | Human Anti SARS-CoV-2 FP (Negative Control) | 0 | 33 |

419
420

421 **Table 2:** Synthesized SARS-CoV-2 Spike RBD Peptides for ACE2 Substrate
422 Competition Assay

Cat #	Domain	Sequence (Amino Acid Position)	Concentration (mg/mL)	Reconstitution
SB086-1MG	**RBD**	319-RVQPTESIVRFPNITNL-335	10	**DMSO**
SB087-1MG	**RBD**	336-CPFGEVFNATRF-347	10	**DMSO**
SB088-1MG	**RBD**	348-ASVYAWNRKR-357	10	**ddH$_2$O**
SB089-1MG	**RBD**	352-AWNRKRISNCVADY-365	10	**ddH$_2$O**
SB090-1MG	**RBD**	371-SASFSTFKCYGVSPTKLNDLCFTN-394	10	**DMSO**
SB091-1MG	**RBD**	395-VYADSFVIRGDEVRQIAPGQTGKIADYNYKLPDDFT-430	10	**ddH$_2$O**
SB092-1MG	**RBM**	438-SNNLDSKVGGNYNYLYRLFRK-458	10	**ddH$_2$O**
SB093-1MG	**RBM**	450-NYLYRLFRKSNLKPFERDISTEIY-473	10	**ddH$_2$O**
SB094-1MG	**RBM**	480-CNGVEGFNCYFPLQSYG-496	10	**DMSO**
SB095-1MG	**RBM**	500-TNGVGYQPYR-509	10	**DMSO**
SB096-1MG	**RBD**	513-LSFELLHA-520	10	**DMSO**
SB097-1MG	**RBD**	523-TVCGPKKSTNLVKNKCVNF-541	10	**ddH$_2$O**

423
424

425
426
427
428
429 **Table 3:** Enzyme Kinetic Studies of ACE2-like Abzymes and Purified Recombinant
430 Human ACE2 Molecule (From FIG 4)
431

Sample	Assay Conc	K_m (µM)	V_{max} (pmol/min)	V_{max} (mol/s)	K_{cat} (s^{-1})	K_{cat}/K_m (M^{-1}s^{-1})
rhACE2[a]	25 ng/mL	43.7	66.9	1.12E-12	3.79E+01	8.68E+05
AGRD013-4G9	5.8 µg/mL	26.9	0.192	3.20E-15	8.27E-04	3.07E+01
AGRD016-2B3	5.6 µg/mL	34.3	0.147	2.45E-15	6.57E-04	1.91E+01
AGRD016-4H2	7.9 µg/mL	38.9	0.295	4.92E-15	9.33E-04	2.40E+01
AGRD021-9A10	5.3 µg/mL	33.9	0.253	4.22E-15	1.19E-03	3.52E+01

432
433 a: Purified recombinant human ACE2 molecule as positive control.
434
435
436

437   **FIGURES**
438

439
FIG 1. ACE2 Background Activity Inhibition by EDTA in Supernatant Control
Media. ACE2 background activity in hybridoma supernatant control media (empty
culture media without hybridoma) was detected by the ACE2 assay, indicated by the
slope calculated from the kinetic curve. ACE2 background activity was effectively
inhibited by adding EDTA to the media. The dashed line shows the cutoff, which was
the limitation of the ACE2 assay and was determined using a purified recombinant
human ACE2 positive control to call a sample positive for ACE2 activity. 5mM EDTA
(**Triangle**) were used for following hybridoma culture supernatants screening as
reported[38].
449

FIG 2. Initial Screening of mAbs for ACE2-like Activity. ACE2 substrate cleavage
451

activity of mAbs. Catalytic activity was determined by calculating the slope of RFU from
452

the initial 2 hours of the kinetics curve. 0.08 ng of purified recombinant human ACE2
453

served as a positive control and standard to define ACE2-positive activity. mAbs with
454

ACE2-like activity greater than or equal to 0.08ng rhACE2 were identified as having
455

ACE2-like catalytic activity. (A) Representative ACE2 activity screening kinetic curves
456

from the Precision Antibody mAbs and ACE2 activity determination by slope. 0.08 ng
457

rhACE2 served as a positive and standard control, and medium (empty hybridoma
458

culture supernatant) served as a negative control. (B) Summary of average ACE2-like
459

activity of all mAbs screened in this study. Results were from at least 2 rounds of
460

screening. ACE2 substrate cleavage activity was determined by the slope of the RFU
461

from the initial 2 hours of the kinetic curve. The dashed line shows the slope calculated
462

from 0.08 ng of purified recombinant human ACE2, the initial 2-hour kinetic curve.
463

467 **FIG 3. Correlation of total IgG and anti-RBD antibody.** Total IgG concentrations in

468 mAb hybridoma culture supernatants were measured by human IgG ELISA. Anti-RBD

469 IgG concentrations of mAb hybridoma culture supernatants were determined by

470 ImmunoCAP antibody-binding assay. (Pearson correlation test).

FIG 4. Michaelis-Menten Analysis of mAb Catalytic Activity. Michaelis-Menten plots

for (A) Positive control (2.5ng rhACE2) and (B) mAbs that had ACE2 substrate cleavage

activities. Abzyme IgG amount for Michaelis-Menten kinetics was from human IgG

ELISA.

481 **FIG 5. Enzyme Substrate Catalytic Specificity.** The substrate cleavage specificity of

482 the mAbs was characterized by comparing the activity of the mAbs using an ACE2

483 substrate and a control non-cleavable substrate. The slope of the RFU from the initial 2

484 hours of the kinetic curve was calculated to determine ACE2-like activity. Activity

485 observed with the control substrate was substantially diminished.

FIG 6. SARS-CoV-2 RBD Peptide Competitive Inhibition of mAb ACE2 Substrate

Cleavage Activity. mAbs ACE2-like activity inhibition by RBD peptide competition was

performed by adding RBD peptide pools to ACE2 assay reactions containing mAbs

defined as ACE2-like activity-positive. Peptide pool diluent, with an equal volume of

peptide pool added to the assay reactions, served as a comparison. 2.5 ng purified

recombinant human ACE2 was used as a positive control. ACE2-like activity was

determined by calculation of the slope of the RFU from the initial 2 hours of the kinetic

curve.

FIG 7. Effect of ACE2 Inhibitor, MLN-4760, on ACE2 Substrate Cleavage Activity of

mAbs and rhACE2. 100 μM ACE2 inhibitor, MLN-4760, was added to ACE2 assay

reactions for comparison of ACE2 activity with ACE2 assay reactions that had no

inhibitor. 2.5 ng purified recombinant human ACE2 was the positive control, and control

media (empty hybridoma culture supernatant) was the negative control. ACE2-like

activity was determined by calculation of the slope of the RFU from the initial 2 hours of

the kinetic curve.

References

1.   Altmann DM, Whettlock EM, Liu S, Arachchillage DJ, Boyton RJ. The immunology of long COVID. Nature Reviews Immunology. 2023;23(10):618-34.

2.   Perumal R, Shunmugam L, Naidoo K, Abdool Karim SS, Wilkins D, Garzino-Demo A, Brechot C, Parthasarathy S, Vahlne A, Nikolich JŽ. Long COVID: a review and proposed visualization of the complexity of long COVID. Frontiers in immunology. 2023;14:1117464.

3.   Davis HE, McCorkell L, Vogel JM, Topol EJ. Long COVID: major findings, mechanisms and recommendations. Nature Reviews Microbiology. 2023;21(3):133-46.

4.   Adjaye-Gbewonyo D, Vahratian A, Perrine CG, Bertolli J. Long COVID in Adults: United States, 2022. 26/2023.

5.   Marchi M, Grenzi P, Serafini V, Capoccia F, Rossi F, Marrino P, Pingani L, Galeazzi GM, Ferrari S. Psychiatric symptoms in Long-COVID patients: a systematic review. Frontiers in Psychiatry. 2023;14:1138389.

6.   Thaweethai T, Jolley SE, Karlson EW, Levitan EB, Levy B, McComsey GA, McCorkell L, Nadkarni GN, Parthasarathy S, Singh U, Walker TA, Selvaggi CA, Shinnick DJ, Schulte CCM, Atchley-Challenner R, Horwitz LI, Foulkes AS, Authors RC, Consortium R. Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection. JAMA. 2023;329(22):1934-46.

7.   Soriano JB, Murthy S, Marshall JC, Relan P, Diaz JV. A clinical case definition of post-COVID-19 condition by a Delphi consensus. Lancet Infect Dis. 2022 Apr;22(4):e102-e7. PMC8691845

8.   Castanares-Zapatero D, Chalon P, Kohn L, Dauvrin M, Detollenaere J, Maertens de Noordhout C, Primus-de Jong C, Cleemput I, Van den Heede K. Pathophysiology and mechanism of long COVID: a comprehensive review. Ann Med. 2022 Dec;54(1):1473-87. PMC9132392

9.   Tramontano A, Janda KD, Lerner RA. Catalytic antibodies. Science. 1986 Dec 19;234(4783):1566-70.

10.   Pollack SJ, Jacobs JW, Schultz PG. Selective chemical catalysis by an antibody. Science. 1986 Dec 19;234(4783):1570-3.

11.   Avalle B, Mistro D, Thomas D, Friboulet A. Polyclonal catalytic anti-idiotypic antibodies with a beta-lactamase activity. Ann N Y Acad Sci. 1996 Oct 12;799:172-5.

12.   Friboulet A, Izadyar L, Avalle B, Roseto A, Thomas D. Abzyme generation using an anti-idiotypic antibody as the "internal image" of an enzyme active site. Appl Biochem Biotechnol. 1994 May-Jun;47(2-3):229-37; discussion 37-9.

13.   Izadyar L, Friboulet A, Remy MH, Roseto A, Thomas D. Monoclonal anti-idiotypic
antibodies as functional internal images of enzyme active sites: production of a catalytic
antibody with a cholinesterase activity. Proc Natl Acad Sci U S A. 1993 Oct
1;90(19):8876-80. PMC47463

14.   Friboulet A, Izadyar L, Avalle B, Roseto A, Thomas D. Antiidiotypic antibodies as
functional internal images of enzyme-active sites. Ann N Y Acad Sci. 1995 Mar
31;750:265-70.

15.   Hilvert D. Critical analysis of antibody catalysis. Annu Rev Biochem.
2000;69:751-93.

16.   Bar-Even A, Noor E, Savir Y, Liebermeister W, Davidi D, Tawfik DS, Milo R. The
moderately efficient enzyme: evolutionary and physicochemical trends shaping enzyme
parameters. Biochemistry. 2011 May 31;50(21):4402-10.

17.   Ponomarenko NA, Pillet D, Paon M, Vorobiev, II, Smirnov IV, Adenier H, Avalle
B, Kolesnikov AV, Kozyr AV, Thomas D, Gabibov AG, Friboulet A. Anti-idiotypic
antibody mimics proteolytic function of parent antigen. Biochemistry. 2007 Dec
18;46(50):14598-609.

18.   Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X,
Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L,
Xie J, Wang G, Jiang R, Gao Z, Jin Q, Wang J, Cao B. Clinical features of patients
infected with 2019 novel coronavirus in Wuhan, China. Lancet. 2020 Feb
15;395(10223):497-506. PMC7159299

19.   Liao D, Zhou F, Luo L, Xu M, Wang H, Xia J, Gao Y, Cai L, Wang Z, Yin P, Wang
Y, Tang L, Deng J, Mei H, Hu Y. Haematological characteristics and risk factors in the
classification and prognosis evaluation of COVID-19: a retrospective cohort study.
Lancet Haematol. 2020 Sep;7(9):e671-e8. PMC7351397

20.   Del Valle DM, Kim-Schulze S, Huang HH, Beckmann ND, Nirenberg S, Wang B,
Lavin Y, Swartz TH, Madduri D, Stock A, Marron TU, Xie H, Patel M, Tuballes K, Van
Oekelen O, Rahman A, Kovatch P, Aberg JA, Schadt E, Jagannath S, Mazumdar M,
Charney AW, Firpo-Betancourt A, Mendu DR, Jhang J, Reich D, Sigel K, Cordon-Cardo
C, Feldmann M, Parekh S, Merad M, Gnjatic S. An inflammatory cytokine signature
predicts COVID-19 severity and survival. Nat Med. 2020 Oct;26(10):1636-43.
PMC7869028

21.   Sun DS, Lien TS, Chang HH. Virus-Induced Pathogenic Antibodies: Lessons
from Long COVID and Dengue Hemorrhage Fever. Int J Mol Sci. 2025 Feb 22;26(5).
PMC11899886

22.   Santos Guedes de Sa K, Silva J, Bayarri-Olmos R, Brinda R, Alec Rath
Constable R, Colom Diaz PA, Kwon DI, Rodrigues G, Wenxue L, Baker C,
Bhattacharjee B, Wood J, Tabacof L, Liu Y, Putrino D, Horvath TL, Iwasaki A. A causal
link between autoantibodies and neurological symptoms in long COVID. medRxiv. 2024
Jun 19. PMC11213106

23.   Yousaf AR, Cortese MM, Taylor AW, Broder KR, Oster ME, Wong JM, Guh AY,
McCormick DW, Kamidani S, Schlaudecker EP, Edwards KM, Creech CB, Staat MA,
Belay ED, Marquez P, Su JR, Salzman MB, Thompson D, Campbell AP. Reported
cases of multisystem inflammatory syndrome in children aged 12-20 years in the USA
who received a COVID-19 vaccine, December, 2020, through August, 2021: a
surveillance investigation. Lancet Child Adolesc Health. 2022 Feb 22. PMC8864018

24.   Khurana V, Goswami B. Angiotensin converting enzyme (ACE). Clin Chim Acta.
2022 Jan 1;524:113-22.

25.   Turner AJ, Nalivaeva NN. Angiotensin-converting enzyme 2 (ACE2): Two
decades of revelations and re-evaluation. Peptides. 2022 May;151:170766.
PMC8830188

26.   Song Y, Mehl F, Muehling LM, Canderan G, Enfield K, Sun J, Yin MT, Ratcliffe
SJ, Wilson JM, Kadl A, Woodfolk JA, Zeichner SL. ACE-2-like enzymatic activity in
COVID-19 convalescents with persistent pulmonary symptoms associated with
immunoglobulin. mBio. 2025 Jul 22:e0173525.

27.   Song Y, Myers R, Mehl F, Murphy L, Brooks B, Wilson JM, Kadl A, Woodfolk J,
Zeichner SL. ACE-2-like enzymatic activity is associated with immunoglobulin in
COVID-19 patients. mBio. 2024 Apr 10;15(4):e0054124. PMC11005375

28.   Zost SJ, Gilchuk P, Case JB, Binshtein E, Chen RE, Nkolola JP, Schäfer A,
Reidy JX, Trivette A, Nargi RS, Sutton RE, Suryadevara N, Martinez DR, Williamson
LE, Chen EC, Jones T, Day S, Myers L, Hassan AO, Kafai NM, Winkler ES, Fox JM,
Shrihari S, Mueller BK, Meiler J, Chandrashekar A, Mercado NB, Steinhardt JJ, Ren K,
Loo YM, Kallewaard NL, McCune BT, Keeler SP, Holtzman MJ, Barouch DH, Gralinski
LE, Baric RS, Thackray LB, Diamond MS, Carnahan RH, Crowe JE, Jr. Potently
neutralizing and protective human antibodies against SARS-CoV-2. Nature. 2020
Aug;584(7821):443-9. PMC7584396

29.   Greaney AJ, Starr TN, Gilchuk P, Zost SJ, Binshtein E, Loes AN, Hilton SK,
Huddleston J, Eguia R, Crawford KHD, Dingens AS, Nargi RS, Sutton RE, Suryadevara
N, Rothlauf PW, Liu Z, Whelan SPJ, Carnahan RH, Crowe JE, Jr., Bloom JD. Complete
Mapping of Mutations to the SARS-CoV-2 Spike Receptor-Binding Domain that Escape
Antibody Recognition. Cell Host Microbe. 2021 Jan 13;29(1):44-57.e9. PMC7676316

30.   Cerutti G, Guo Y, Liu L, Liu L, Zhang Z, Luo Y, Huang Y, Wang HH, Ho DD,
Sheng Z, Shapiro L. Cryo-EM structure of the SARS-CoV-2 Omicron spike. Cell Rep.
2022 Mar 1;38(9):110428. PMC8818377

31.   Liu L, Iketani S, Guo Y, Reddem ER, Casner RG, Nair MS, Yu J, Chan JF, Wang
M, Cerutti G, Li Z, Morano NC, Castagna CD, Corredor L, Chu H, Yuan S, Poon VK,
Chan CC, Chen Z, Luo Y, Cunningham M, Chavez A, Yin MT, Perlin DS, Tsuji M, Yuen
KY, Kwong PD, Sheng Z, Huang Y, Shapiro L, Ho DD. An antibody class with a
common CDRH3 motif broadly neutralizes sarbecoviruses. Sci Transl Med. 2022 May
25;14(646):eabn6859. PMC9017343

32.   Liu L, Casner RG, Guo Y, Wang Q, Iketani S, Chan JF, Yu J, Dadonaite B, Nair
MS, Mohri H, Reddem ER, Yuan S, Poon VK, Chan CC, Yuen KY, Sheng Z, Huang Y,
Bloom JD, Shapiro L, Ho DD. Antibodies targeting a quaternary site on SARS-CoV-2
spike glycoprotein prevent viral receptor engagement by conformational locking.
Immunity. 2023 Oct 10;56(10):2442-55.e8. PMC10588992

33.   Dacon C, Tucker C, Peng L, Lee CD, Lin TH, Yuan M, Cong Y, Wang L, Purser
L, Williams JK, Pyo CW, Kosik I, Hu Z, Zhao M, Mohan D, Cooper A, Peterson M,
Skinner J, Dixit S, Kollins E, Huzella L, Perry D, Byrum R, Lembirik S, Zhang Y, Yang
ES, Chen M, Leung K, Weinberg RS, Pegu A, Geraghty DE, Davidson E, Douagi I, Moir
S, Yewdell JW, Schmaljohn C, Crompton PD, Holbrook MR, Nemazee D, Mascola JR,
Wilson IA, Tan J. Broadly neutralizing antibodies target the coronavirus fusion peptide.
bioRxiv. 2022 Apr 12:2022.04.11.487879. PMC9016638

34.   Cho H, Gonzales-Wartz KK, Huang D, Yuan M, Peterson M, Liang J, Beutler N,
Torres JL, Cong Y, Postnikova E, Bangaru S, Talana CA, Shi W, Yang ES, Zhang Y,
Leung K, Wang L, Peng L, Skinner J, Li S, Wu NC, Liu H, Dacon C, Moyer T, Cohen M,
Zhao M, Lee FE, Weinberg RS, Douagi I, Gross R, Schmaljohn C, Pegu A, Mascola JR,
Holbrook M, Nemazee D, Rogers TF, Ward AB, Wilson IA, Crompton PD, Tan J.
Bispecific antibodies targeting distinct regions of the spike protein potently neutralize
SARS-CoV-2 variants of concern. Sci Transl Med. 2021 Oct 20;13(616):eabj5413.
PMC8651051

35.   Hifumi E, Ito Y, Tsujita M, Taguchi H, Uda T. Enzymatization of mouse
monoclonal antibodies to the corresponding catalytic antibodies. Sci Rep. 2024 May
28;14(1):12184. PMC11133420

36.   McConnell SA, Sachithanandham J, Mudrak NJ, Zhu X, Farhang PA, Cordero
RJB, Wear MP, Shapiro JR, Park HS, Klein SL, Tobian AAR, Bloch EM, Sullivan DJ,
Pekosz A, Casadevall A. Spike-protein proteolytic antibodies in COVID-19 convalescent
plasma contribute to SARS-CoV-2 neutralization. Cell Chem Biol. 2023 Jul
20;30(7):726-38.e4. PMC10288624

37.   Xiao F, Burns KD. Measurement of Angiotensin Converting Enzyme 2 Activity in
Biological Fluid (ACE2). Methods Mol Biol. 2017;1527:101-15. PMC7121061

38.   Lubel JS, Herath CB, Velkoska E, Casley DJ, Burrell LM, Angus PW. Angiotensin
converting enzyme 2 (ACE2) activity in fetal calf serum: implications for cell culture
research. Cytotechnology. 2008 Nov;58(3):119-26. PMC2652558

39.   Ferrari M, Mekkaoui L, Ilca FT, Akbar Z, Bughda R, Lamb K, Ward K, Parekh F,
Karattil R, Allen C, Wu P, Baldan V, Mattiuzzo G, Bentley EM, Takeuchi Y, Sillibourne J,
Datta P, Kinna A, Pule M, Onuoha SC. Characterization of a Novel ACE2-Based
Therapeutic with Enhanced Rather than Reduced Activity against SARS-CoV-2
Variants. Journal of Virology. 2021;95(19):10.1128/jvi.00685-21.

40.   Svilenov HL, Delhommel F, Siebenmorgen T, Rührnößl F, Popowicz GM, Reiter
A, Sattler M, Brockmeyer C, Buchner J. Extrinsic stabilization of antiviral ACE2-Fc

fusion proteins targeting SARS-CoV-2. Commun Biol. 2023 Apr 8;6(1):386.
PMC10082628
706
707   41.   Liu L, Wang P, Nair MS, Yu J, Rapp M, Wang Q, Luo Y, Chan JF, Sahi V,
708   Figueroa A, Guo XV, Cerutti G, Bimela J, Gorman J, Zhou T, Chen Z, Yuen KY, Kwong
709   PD, Sodroski JG, Yin MT, Sheng Z, Huang Y, Shapiro L, Ho DD. Potent neutralizing
710   antibodies against multiple epitopes on SARS-CoV-2 spike. Nature. 2020
711   Aug;584(7821):450-6.
712
713   42.   Michaelis L, Menten ML, Johnson KA, Goody RS. The original Michaelis
714   constant: translation of the 1913 Michaelis-Menten paper. Biochemistry. 2011 Oct
715   4;50(39):8264-9. PMC3381512
716
717   43.   Rice GI, Thomas DA, Grant PJ, Turner AJ, Hooper NM. Evaluation of
718   angiotensin-converting enzyme (ACE), its homologue ACE2 and neprilysin in
719   angiotensin peptide metabolism. Biochem J. 2004 Oct 1;383(Pt 1):45-51. PMC1134042
720
721   44.   Ferrari M, Mekkaoui L, Ilca FT, Akbar Z, Bughda R, Lamb K, Ward K, Parekh F,
722   Karattil R, Allen C, Wu P, Baldan V, Mattiuzzo G, Bentley EM, Takeuchi Y, Sillibourne J,
723   Datta P, Kinna A, Pule M, Onuoha SC. Characterization of a Novel ACE2-Based
724   Therapeutic with Enhanced Rather than Reduced Activity against SARS-CoV-2
725   Variants. J Virol. 2021 Sep 9;95(19):e0068521. PMC8432736
726
45.   Organization WH. WHO COVID-19 Dashboard. World Health Organization; 2026
[cited 2026 14 January 2026]; https://data.who.int/dashboards/covid19/deaths.
46.   de Souza MO, Madan B, Teng IT, Huang A, Liu L, Fahad AS, Lopez Acevedo
SN, Pan X, Sastry M, Gutierrez-Gonzalez M, Yin MT, Zhou T, Ho DD, Kwong PD,
DeKosky BJ. Mapping monoclonal anti-SARS-CoV-2 antibody repertoires against
diverse coronavirus antigens. Front Immunol. 2022;13:977064. PMC9478573
733
47.   Amanat F, Thapa M, Lei T, Sayed Ahmed SM, Adelsberg DC, Carreno JM,
Strohmeier S, Schmitz AJ, Zafar S, Zhou JQ, Rijnink W, Alshammary H, Borcherding N,
Reiche AG, Srivastava K, Sordillo EM, van Bakel H, Turner JS, Bajic G, Simon V,
Ellebedy AH, Krammer F. The plasmablast response to SARS-CoV-2 mRNA
vaccination is dominated by non-neutralizing antibodies and targets both the NTD and
the RBD. medRxiv. 2021 May 1. PMC7987037
740
741   48.   Cesaro-Tadic S, Lagos D, Honegger A, Rickard JH, Partridge LJ, Blackburn GM,
742   Plückthun A. Turnover-based in vitro selection and evolution of biocatalysts from a fully
743   synthetic antibody library. Nat Biotechnol. 2003 Jun;21(6):679-85.
744
745   49.   Ermakov EA, Kabirova EM, Buneva VN, Nevinsky GA. IgGs-Abzymes from the
746   Sera of Patients with Multiple Sclerosis Recognize and Hydrolyze miRNAs. Int J Mol
747   Sci. 2021 Mar 10;22(6). PMC8000798
748
749   50.   Li JW, Xia L, Su Y, Liu H, Xia X, Lu Q, Yang C, Reheman K. Molecular imprint of
750   enzyme active site by camel nanobodies: rapid and efficient approach to produce
751   abzymes with alliinase activity. J Biol Chem. 2012 Apr 20;287(17):13713-21.
752   PMC3340132

753
754 51.   Richard JP. Protein Flexibility and Stiffness Enable Efficient Enzymatic Catalysis.
755 J Am Chem Soc. 2019 Feb 27;141(8):3320-31. PMC6396832
756
757 52.   Jin M, Jefferson VA, Zhao Z, Catanzaro NJ, De Sabato L, Keller EL, Menasche
758 BL, Hoffman C, Castelli A, Di Bartolo I, Vaccari G, Starr TN, Seifert SN, Castellanos AA,
759 Han BA, Serebryannyy L, Spaulding AB, Douek DC, Moreno A, Baric RS, Wilen CB,
760 Rini JM, Letko M. Aminopeptidase N is a receptor for hedgehog merbecoviruses.
761 bioRxiv. 2025 Sep 3. PMC12425009
762
763 53.   Dufloo J, Fernández I, Arbabian A, Haouz A, Temperton N, Gimenez-Lirola LG,
764 Rey FA, Sanjuán R. Dipeptidase 1 is a functional receptor for a porcine coronavirus. Nat
765 Microbiol. 2025 Nov;10(11):2981-96. PMC12578638
766
767 54.   Dufloo J, Andreu-Moreno I, Moreno-García J, Valero-Rello A, Sanjuán R.
768 Receptor-binding proteins from animal viruses are broadly compatible with human cell
769 entry factors. Nat Microbiol. 2025 Feb;10(2):405-19. PMC11790484
770
771 55.   Simmons G, Zmora P, Gierer S, Heurich A, Pöhlmann S. Proteolytic activation of
772 the SARS-coronavirus spike protein: cutting enzymes at the cutting edge of antiviral
773 research. Antiviral Res. 2013 Dec;100(3):605-14. PMC3889862
774
775 56.   López-Cortés GI, Palacios-Pérez M, Hernández-Aguilar MM, Veledíaz HF, José
776 MV. Human Coronavirus Cell Receptors Provide Challenging Therapeutic Targets.
777 Vaccines (Basel). 2023 Jan 13;11(1). PMC9862439
778
779 57.   Li F. Receptor recognition mechanisms of coronaviruses: a decade of structural
780 studies. J Virol. 2015 Feb;89(4):1954-64. PMC4338876
781
782 58.   Liu L, Casner RG, Guo Y, Wang Q, Iketani S, Chan JF, Yu J, Dadonaite B, Nair
783 MS, Mohri H, Reddem ER, Yuan S, Poon VK, Chan CC, Yuen KY, Sheng Z, Huang Y,
784 Bloom JD, Shapiro L, Ho DD. Antibodies that neutralize all current SARS-CoV-2
785 variants of concern by conformational locking. bioRxiv. 2023 Apr 10. PMC10120718
786
787