1

ACE-2-like Enzymatic Activity in Anti-SARS-CoV-2 Spike Protein Monoclonal Antibodies

Yufeng Song[1, #], Frances Mehl[1, #], Tom No[1,¶], Lauren Livingston[1], Juan Sebastian

Quintero Barbosa[1], Jun Hayashi[2], Ginette Serrero[2], Pamela Schoppee Bortz[3], Jeffrey M

Wilson[3], James E Crowe Jr[4], David D Ho[5], Michael T Yin[6], Joshua Tan[7], Peter Kwong[8],

Steven L Zeichner[1,9,*]

[1] Department of Pediatrics, University of Virginia, Charlottesville, Virginia, USA.

[2] Precision Antibody, 9130 Red Branch Rd., Suite X Columbia, Maryland 21045, United

States.

[3] Division of Allergy and Clinical Immunology, Department of Medicine, University of

Virginia, Charlottesville, Virginia, USA.

[4] Howard Hughes Medical Institute, Seattle, WA 98109, USA; Department of Pathology,

Microbiology, and Immunology, Vanderbilt University Medical Center, Nashville, TN

37232, USA; Department of Pediatrics, Vanderbilt University Medical Center, Nashville,

TN 37232, USA.

[5] Aaron Diamond AIDS Research Center, Columbia University Vagelos College of

Physicians and Surgeons, New York, NY 10032, USA; Division of Infectious Diseases,

Department of Medicine, Columbia University Vagelos College of Physicians and

Surgeons, New York, NY 10032, USA; Department of Microbiology and Immunology,

Columbia University Vagelos College of Physicians and Surgeons, New York, NY

10032, USA.

[6] Columbia University Vagelos College of Physicians and Surgeons, New York, New

York, USA.

[7] Antibody Biology Unit, Laboratory of Immunogenetics, National Institute of Allergy and

[8] Vaccine Research Center, National Institute of Allergy and Infectious Diseases,

National Institutes of Health, Bethesda, MD 20892, USA; Aaron Diamond AIDS

Research Center, Columbia University Vagelos College of Physicians and Surgeons,

and Department of Biochemistry and Molecular Biophysics, Columbia University, New

York, NY 10032, USA.

[9] Department of Microbiology, Immunology, and Cancer Biology, University of Virginia,

Charlottesville, Virginia, USA.

38   [#]Contributed equally to this work.

40   [¶]Present address: Department of Urology, University of California, Irvine, CA, USA.

42   *Correspondence: zeichner@virginia.edu

Abstract

Many people with acute COVID-19 have clinical disease not clearly attributable to viral replication and many COVID-19 convalescents are affected by post-acute sequelae of COVID-19 (PASC, or long COVID, LC). LC has severely affected public health and economies worldwide. Features of LC including blood pressure dysregulation, coagulopathies, high levels of inflammation, and neuropsychiatric complaints. The mechanisms responsible for the pathogenesis of some COVID-19 clinical features and LC have not been well established. The host cell receptor for SARS-CoV-2 is human angiotensin converting enzyme 2 (ACE2), which binds the SARS-CoV-2 spike protein receptor-binding domain (RBD) to initiate infection. We hypothesized that some people may produce anti-RBD antibodies that sufficiently resemble ACE2 structure to have ACE2-like catalytic activity after infection. Those antibodies, ACE2-like abzymes, may contribute to the pathogenesis of LC. Our previous studies showed that ACE2-like activity was associated with immunoglobulin in some acute and convalescent COVID-19 patients. ACE2-like catalytic activity correlated with blood pressure changes following a moderate exercise challenge in people convalescing from COVID-19. To further establish that ACE2-like activity could be attributed to antibodies, we screened human monoclonal antibodies (mAbs) against SARS-CoV-2 spike protein from 3 different research centers and others purchased from a commercial source for ACE2-like catalytic activity. We identified 4 human monoclonal antibodies with ACE2-like catalytic activity. The ACE2-like catalytic activity of these mAbs was not inhibited by MLN-4760, a compound that inhibits native human ACE2 catalytic activity, nor by EDTA, unlike native ACE2, a Zinc metalloprotease, but was inhibited by an overlapping pool of spike peptides. Enzyme kinetic studies showed that the mAbs had substantially lower Vmax and *Km* values than native ACE2. The data therefore suggested that the antibodies cleave ACE2 substrate via a different mechanism than native ACE2. The identification of specific mAbs with ACE2-like catalytic activity supports the hypothesis that antibodies induced by SARS-CoV-2 infection could help mediate the pathogenesis of COVID-19 and LC, and more generally, the hypothesis that catalytic antibodies induced by infectious agents can contribute to disease pathogenesis.

Introduction

COVID-19 and Post-Acute Sequelae of COVID-19 (PASC, or Long COVID, LC), as reviewed in [1-3], can affect up to 7% of COVID-19 patients[4]. LC patients can have many unusual symptoms such as dizziness, fatigue, malaise, "brain fog", alterations in smell or taste, abnormal movements, libido changes, and other neuropsychiatric symptoms, GI symptoms, cardiovascular complaints such as palpitations, blood pressure dysregulation, inflammation, coagulopathies, cough, and chest pain[5 6, 7]. While the clinical features, laboratory findings, and epidemiology of LC have been extensively described, clear primary causal etiologies have yet to be determined [8].

Antibodies with catalytic activity, also called "abzymes" have been known for some time [9, 10]. Investigators initially produced abzymes by generating anti-idiotypic antibodies against antibodies targeting enzymes' active sites[11, 12]. In some instances, the anti-idiotypic antibodies had sufficient resemblance to the original enzyme's active site so as to exhibit catalytic activity similar to that of the initially targeted enzyme[13, 14]. However, further research showed that the abzymes had decreased substrate specificity and less favorable enzyme constants compared to the targeted enzymes (*e.g.,* $k$cat/Km values $\sim 10^2$ - 10^4 s^{-1}•M s^{-1} [15] vs. $k$cat/Km values of $\sim 10^5$ s^{-1}•M s^{-1})[16 12, 13, 17]. These limitations led to a decline in enthusiasm for the use of abzymes in biotechnological applications.

Antibodies are traditionally understood as high affinity binding molecules. However, a subset of antibodies with intrinsic catalytic activity, commonly referred to as natural abzymes, has been described in a variety of immune contexts, including autoimmune disease, chronic viral infection, and exposure to persistent antigens[18-20]. Such antibodies typically exhibit low catalytic turnover relative to canonical enzymes, have heterogeneous substrate preferences, and occur at low frequencies within polyclonal repertoires[18, 20-22]. While the catalytic efficiency of such abzymes is lower than typical enzymes, their biological significance lies in their persistence, the potentially high concentration of immunoglobulin molecules in the circulation, and systemic distribution[23]. With respect to SARS-CoV-2, anti-spike protein antibodies have been observed in

the nanomolar range [24].

Abzymes have been shown to exert measurable biochemical activity and, in certain

contexts, have been implicated in immune-mediated modulation of host physiology[18, 20, 25, 26].

Given that homeostatic regulatory mechanisms are finely balanced, low-level hydrolysis of

regulatory peptide, like angiotensin II, could shift the homeostatic "proteolytic tone"[27], although

the short half-life of regulatory peptides, like angiotensin II (less than 1 minute) [28] make

meaningful *in vivo* measurements difficult .

In an attempt to identify potential mechanisms responsible for the unusual features of

COVID-19 and LC, we considered that many of these aberrant clinical features could be viewed

as consequences of pathologic derangements of proteolytic regulatory cascades, for example,

blood pressure regulation (the renin-angiotensin system), inflammation (C', kallikrein-kinin)[29-31],

and coagulation. For LC, some most recent studies found the kinetics of symptom development

would be consistent with antibody production, rather than the pathology directly produced by

viral replication[32, 33] . In addition, a recent report showed that immunoglobulins transferred from

some LC patients to mice can induce symptoms, further suggesting that immunoglobulin may

help mediate some LC symptoms[34].

While the benefits of COVID-19 vaccination vastly outweigh the risks, the fact that a very

small, but still non-zero, fraction of patients who received SARS-CoV-2 vaccines had symptoms

including hyperinflammation, blood pressure dysregulation, or coagulopathies [35-37] further

suggests that, in some cases, an antibody response could play a role in pathogenesis of LC, as

opposed to lasting sequelae of active viral replication.

Angiotensin Converting Enzyme 2 (ACE2) on the host cell surface serves as the principal

receptor for the SARS-CoV-2 Receptor Binding Domain (RBD). Angiotensin Converting Enzyme

(ACE) catalyzes the cleavage of angiotensin I to angiotensin II, which acts as vasoconstrictor to

increase blood pressure [38]. ACE2 cleaves angiotensin II to produce angiotensin 1-7, which has

vasodilatory activity that lowers blood pressure, an activity that ACE2 counteracts the activity of

ACE in the renin–angiotensin system by converting angiotensin II to angiotensin (1–7),

opposing ACE-driven signaling[39]. While interactions between the SARS-CoV-2 Spike protein

and the human ACE2 receptor are well-documented at the binding interface, the functional

consequences of this interaction remain under-explored[40]. We hypothesized that the

pathogenesis of some of the unusual features of LC might be caused by the induction of

antibodies against the SARS-CoV-2 Spike protein (S) that resembled the structure of ACE2

closely enough to have ACE2-like catalytic activity.

In our previous studies[41, 42], we found that some patients' plasma with ACE2-like activity did

not require divalent cations, unlike native ACE2, a Zinc metalloprotease. This activity co-

depleted with immunoglobulin and was inhibited by a tiled overlapping RBD peptide pool. These

findings suggested that the ACE2-like activity observed in patients' plasma differed from native

ACE2 and was the result of anti-RBD immunoglobulins exhibiting ACE2-like activity.

Additionally, ACE-2 like catalytic activity found in plasma collected from convalescent

patients correlated with changes in blood pressure following a moderate exercise challenge[41],

suggesting that the catalytic antibodies could have physiologic consequences.

However, to definitively establish that COVID-19 patients produce antibodies with catalytic

activity, it is important to identify a pure monoclonal antibody derived from COVID-19 patients

that exhibits the hypothesized catalytic activity. Here, we survey several sets of monoclonal

antibodies (mAbs) from 4 different research centers.

We utilized a widely used surrogate substrate which enables sensitive, real-time detection

of catalytic activity and is commonly employed in ACE2 activity and inhibitor screening assays

[43-45] to monitor ACE2-like activity. While we found that most mAbs lacked catalytic activity, two

mAbs from one set showed catalytic activity higher than the level from the standard positive

control, and two others showed borderline detectable activity. These observations show that

some patients develop antibodies with catalytic activity following SARS-CoV-2 infection, with

potential clinical consequences, providing a biochemical basis for how a virus-induced immune

response might transition into a chronic, autoimmune catalytic process.

Materials and Methods

Monoclonal Antibody Collections

We received 129 purified human anti-SARS-CoV-2 RBD mAbs, 1 purified anti-SARS-CoV-2 fusion peptide (FP) mAb, 7 human anti-SARS-CoV-2 RBD mAb hybridoma supernatants, 3 irrelevant mAb hybridoma supernatants, and 1 empty hybridoma culture media as a negative control from 4 different collaborator groups (Table 1). Some of the anti-RBD antibodies were further characterized for their ability to block or enhance ACE2 binding to SARS-CoV-2 spike protein. The hybridoma culture medium is 1/3 Medium E (Stem Cell Technologies) and defined 2/3 Dulbecco's Modified Eagle Medium (DMEM) + 10 % fetal bovine serum (FBS).

Monoclonal Antibodies for Screening

All antibodies were assessed for binding to SARS-CoV-2 RBD (with the exceptions of the negative control mAbs) as described[46-52]. The mAbs purchased from Precision Antibodies targeting the RBD (but not those from other sources) were further screened for interactions with ACE2. This screening used (spike) S and (receptor binding domain) RBD recombinant proteins with HIS tag and an ACE2 human Fc fusion protein. Ni-NTA plates were used to capture both S and RBD proteins. ACE2 binding to both S and RBD proteins was confirmed to ensure that both S and RBD retained the ability to bind ACE2. The S protein captured on Ni-NTA plates was used for the initial screening. All the clones that showed binding to S-protein were subsequently screened for specific RBD binding using RBD-captured Ni-NTA plates. ACE2 was added to the plate to see if

any of the anti-RBD antibodies blocked ACE2 binding to the S or RBD proteins captured

on the plate. The binding of ACE2 was detected using HRP-anti-human Fc.

Angiotensin II Converting Enzyme (ACE2)-like Activity Assays

For the ACE2 activity assay, 25 µg purified antibody or 50 µL hybridoma supernatant

before and after immunoglobulin depletion was combined with 50 µM MCA-APK(Dnp)

fluorogenic ACE2 substrate (AAT Bioquest, Pleasanton CA. Cat. #1355) in assay buffer,

50 mM MES assay buffer (Thermo Fisher, Waltham MA. Cat. # J62231.AK)

supplemented with 300 mM NaCl and 10 µM $ZnCl_2$, in a non-binding black bottom plate

(Corning, Glendale AZ. Cat. #3991) [53, 54] with total 100µL reaction volume[55]. Empty

hybridoma culture media was used as a negative control. Antibody concentration in

hybridoma supernatant was determined by IgG ELISA (Table 4) and antibody

concentration in substrate catalytic reaction is shown in Table 3. Fluorescence levels

were measured with excitation/emission wavelengths 320/405nm and a 325 nm

emission filter on the SpectraMax® M5 multi-mode microplate reader (Molecular

Devices, San Jose CA.)[54]. Measurements were taken every five minutes for 16 hours,

while the plate was kept at a constant temperature of 37°C. 5 mM EDTA was added to

the hybridoma culture supernatants to inhibit the ACE2 present in 10% FBS medium[56].

Specificity of ACE2 substrate cleavage was determined by parallel comparisons of

samples with 50 µM OMNIMMP fluorogenic control substrate (Enzo Life Sciences,

Farmingdale, NY. Cat. #MML-P127), MCA-Pro-Leu-OH, according to previous studies

[41, 42, 57].

ACE2 Substrate Peptide Cleavage Competition Inhibition by SARS-CoV-2 Spike

RBD Peptide Pools

Twelve peptides (SB-Peptide, SmartBioscience SAS, France) covering the SARS-CoV-2 RBD (including RBM) were reconstituted in ddH2O or DMSO to a concentration of 10 µg/µL (Table 2) [41, 42]. The peptides were combined in equal amounts. The ACE2 activity assay was set up as described above, and 2.4 µL of the peptide pool was added so that there was 2 µg of each peptide per well. 2.4 µL of peptide diluent (50% ddH$_2$O and 50% DMSO) was added to other wells as a control.

ACE2 Substrate Peptide Cleavage Competition Inhibition by ACE2 Inhibitor

The ACE2 activity assay was run as described above, with the addition of 100 µM MLN-4760 (Bio-Techne, Minneapolis, MN, Cat. #3345), a small-molecule ACE2 active-site inhibitor[55, 58]. This concentration is recommended in commercial ACE2 activity kits (Abcam, Waltham, MA. Cat. #ab273297).

Enzyme Kinetics Assay

The ACE2 activity assay was set up as described, but with substrate dilutions ranging from 12.5 µM to 200 µM. An MCA standard curve (AAT Bioquest, Pleasanton CA. Cat. #557) was set up with 500, 250, 200, 150, 100, 50, 25, and 0 pmol MCA/well in 100 µL assay buffer. This curve enabled the calculation of pmol of MCA-APK(Dnp) substrate cleaved per minute.

RBD-specific Binding Assays by ImmunoCAP

Following our previous studies[41, 42], S-RBD binding was measured by using 50 µL of hybridoma supernatant with a quantitative ImmunoCAP system applying Phadia 250

(Thermo Fisher Scientific, Waltham MA. Thermo-Fisher/Phadia). Briefly, biotinylated S-

RBD (RayBiotech, Peachtree Corners, GA) was conjugated to streptavidin-coated

ImmunoCAPs (Thermo Fisher Scientific, Waltham MA. Thermo-Fisher/Phadia).

Unconjugated streptavidin ImmunoCAP was run in parallel with its signal as a

background to be subtracted from the singles of other samples.

Immunoglobulin Depletion in Hybridoma Supernatants

Human immunoglobulin depletion was performed following pre-established methods [41,

42]. Hybridoma culture supernatants were centrifuged at 12,000 × g at 4°C for 15 min,

and supernatants were passed through a 0.45 µm syringe filter (FisherScientific, Cat.

#97204). 500 µL of the filtered samples were added to ultrafiltration columns with a 100

kDa cutoff ultrafiltration membrane (Pierce Cat. # 88503) and centrifuged at 12,000 × g

at 4°C until >400 µL of filtrate was produced. Then filtrated samples were incubated with

30 µL pre-washed Protein A/G Magnetic Beads (Life Technologies, Cat. # 88803, 10

mg/mL) at room temperature for 1 hour, followed by magnetic removal of the beads,

followed by a second round of Protein A/G Magnetic Bead depletion with another freshly

prepared aliquot of 30 µL magnetic beads. ACE2 substrate cleavage activity before and

after immunoglobulin depletion was measured as described above. Immunoglobulin

depletion efficiency was determined by human IgG ELISA, as described[41, 42].

Human IgG ELISA

The total amount of human IgG in hybridoma supernatant was measured by quantitative

human IgG ELISA following the kit manufacturer's instructions (Abcam, Waltham, MA.

Cat. #ab100547) as described[41, 42].

Data Analysis

ACE2 substrate cleavage was monitored by measuring relative fluorescence units (RFU) for 16 hours with 5-minute intervals. The RFU values were corrected for background fluorescence by subtracting the minimum RFU value from the remaining RFU values for each well in the assay plate. Linear regression analysis was done on the first 5-120 minutes to determine the increase in RFU over time. This slope (RFU/min) was used as an indicator of ACE2-like activity. A slope of 0.5 RFU/min (the slope of 0.08 ng rhACE2) was determined to be the baseline detection level of ACE2 activity, which is two-fold greater than the lower limit of detection, the activity of 0.04 ng rhACE2[41, 42]. Monoclonal antibodies with a slope greater than 0.5 RFU/min were identified as positive for ACE2-like abzyme activity. For the MCA standard curve, the average RFU over 2 hours was plotted against pmol MCA/well. Linear regression was done to determine the number of RFU per pmol MCA. This value allowed for the calculation of pmols substrate cleaved per minute from the slope data. The supernatant IgG concentration was used to determine the *Kcat* of mAb catalytic activity. All data analysis and visualization were performed with R (version 2022-06-23) using R Studio user interface and packages including drc, ggh4x, ggplot2, patchwork, tidyverse, readr, GGally, corrplot and stringr.

Results

We screened 129 purified human anti-RBD mAbs (Table 1) [46, 47, 49, 50, 52, 59] with

one purified broadly neutralizing mAb against SARS-CoV-2 fusion peptide[51] as an

irrelevant control (Table 1). We also screened 10 human mAb hybridoma culture

supernatants purchased from Precision Antibody, including 3 anti-RBD mAbs that

blocked ACE2 binding to RBD, 4 anti-RBD mAbs that enhanced ACE2 binding to RBD,

and 3 negative control irrelevant mAbs (Table 1). 5mM EDTA was added to the

hybridoma supernatants to inhibit any native ACE2 protein in the media, following

previous studies that inhibited ACE2 in 10% FBS media[56]. A concentration of 5 mM

EDTA was confirmed to give complete inhibition of ACE2 in the hybridoma culture

media serving as our negative control (FIG 1).

We found 4 anti-RBD mAbs (AGRD013-4G9, AGRD016-2B3, AGRD016-4H2,

AGRD021-9A10), that had ACE2-like abzyme activity (FIG 2. A). Two of these

(AGRD013-4G9 and AGRD016-4H2) had previously been determined to block RBD

binding to ACE2, while the other two (AGRD016-2B3 and AGRD021-9A10) had been

determined to enhance binding of RBD to ACE2. An additional 9 mAbs had catalytic

activity higher than the lower limit of detection, but lower than our predetermined

threshold for significant ACE2 activity and thus were classified as negative for ACE2-

like activity (FIG 2. B). All negative controls had slopes below the lower limit (FIG 2. B).

Antibody concentrations in hybridoma culture supernatants were measured by a

total human IgG ELISA[41, 42]. The antibody levels ranged from 16 to 28 μg/ml. Anti-RBD

binding activity in those hybridoma culture supernatants was assessed using an

ImmunoCAP assay[41, 42], with RBD antibody concentrations ranging from 8 to 13 μg/mL.

Significant correlation (R = 0.92, p = 0.0035, Pearson correlation coefficient) was

observed between total human IgG concentration and anti-RBD IgG (FIG 3), indicating

that ACE2-like activity correlates with anti-RBD antibody, consistent with our previous

findings[41, 42].

Studies of the enzymatic activities of those mAbs, ACE2-like activity positive

samples, and the rhACE2 positive control, was conducted using Michaelis–Menten and

Lineweaver-Burk analyses[60]. IgG concentrations were determined by human IgG ELISA

(FIG 3, Table 3). The ACE2-like abzymes have lower K_ms, and ~4 $\log_{10}$ lower catalytic

efficiency (Table 3, FIG 4). The low catalytic efficiency (K_{cat}/K_m) is consistent with the

data obtained for other abzymes[15, 16]. Our calculated K_{cat}/K_m for rhACE2 at 37^{o}C is 0.8

$M^{-1}s^{-1}$ higher than the reported value for rhACE2 at room temperature[61].

Substrate preference of abzyme catalytic activity was determined by comparing

the cleavage of ACE2 substrate (MCA-APK(Dnp)) and control substrate (MCA-Pro-Leu-

OH), as in prior studies[41, 42, 62]. In contrast to robust signal generation with the ACE2

substrate, reactions containing the control substrate produced negative slopes (FIG 5),

consistent with the absence of measurable cleavage. These findings indicate that the

antibodies with identified catalytic activity preferentially cleaved the ACE2 substrate

rather than nonspecifically cleaving the control substrate.

To further confirm the catalytic activity of the monoclonal antibodies, we

conducted ACE2 substrate cleavage peptide inhibition assays using 12 tiled overlapping

peptides that cover the SARS-CoV-2 RBD and RBM amino acid sequences (Table 2).

The RBD peptide pools significantly inhibited the ACE2-like activity of two mAbs,

AGRD013-4G9 and AGRD016-4H2, which block RBD binding to ACE2, but had overall

limited inhibition effect on two other mAbs, AGRD016-2B3 and AGRD021-9A10, which

enhance RBD binding to ACE2 (FIG 6). The RBD peptide inhibition results show that

the mAb catalytic activity could be inhibited by peptide competition, and that there is a

difference in the ability of the peptides to inhibit activity between mAbs which block or

enhance receptor binding[41, 42].

To confirm the ACE2 substrate cleavage activity was immunoglobulin related, we

completely depleted the immunoglobulin from mAb hybridoma culture supernatants

(depletion efficiency >99.99%, Table 4) by previously established IgG depletion assay

[41, 42]. Elimination of ACE2 substrate cleavage activity was observed with

immunoglobulin depletion (FIG 7) indicating the ACE2 substrate cleavage activity is

associated with immunoglobulin.

We also tested the effect of a small molecule ACE2 inhibitor, MLN-4760, a well-

documented inhibitor of ACE2 catalytic activity[55, 58]. MLN-4760 completely inhibits

rhACE2, consistent with previous findings[55, 58], but did not inhibit ACE2-like abzyme

activity (FIG 8). As we previously showed, the divalent cation chelator, EDTA did not

affect the mAbs' substrate cleavage activity, but inhibited native ACE2 catalytic activity,

since native ACE2 is a Zn^{2+} metalloprotease[41, 42]. These findings suggest that the mAbs

with ACE2-like activity employ different catalytic mechanisms than native ACE2.

Discussion

The COVID-19 pandemic has claimed around 7 million lives, according to the World Health Organization, as of January 2026[63], with about 65 million patients experiencing lingering symptoms long after their initial infection[3]. Long COVID, characterized by persistent symptomatology following an initial COVID-19 infection, includes symptoms consistent with dysregulated proteolytic regulatory cascades[29-31] such as the angiotensin-renin system [38].

Our previous research showed that some patients suffering from acute SARS-CoV-2 infection or convalescing from COVID-19 produced antibodies with ACE2-like catalytic activity [41, 42]. In this study, to further explore our hypothesis that some anti-spike antibodies exhibit ACE2-like catalytic activity, we screened 129 anti-SARS-CoV-2 mAbs, 7 anti-spike mAb hybridoma supernatants; and 3 irrelevant mAb hybridoma supernatants and an anti-SARS-CoV-2 FP mAb as controls (Table 1). Although most anti-RBD mAbs exhibited no ACE2-like activity, we identified 4 anti-RBD mAb hybridoma supernatants that exhibited measurable catalytic activity against a synthetic fluorogenic peptide substrate commonly used to assess metalloprotease function. This finding is consistent with our hypothesis that only a small fraction of elicited RBD antibodies exhibit a widely used surrogate ACE2 substrate cleavage activity after infection underscoring the rarity of this phenomenon [24, 25].

The catalytic activity of metalloprotease ACE2 is inhibited by 5 mM EDTA in the assay buffer. The ACE2-like catalytic activity of the mAbs was not inhibited by 5 mM EDTA (FIG 1), just as we observed directly in samples from research volunteers with both acute COVID-19 and convalescing from COVID-19[41, 42]. ACE2 substrate cleavage was

further confirmed by conducting assays using either a native substrate or a negative

control ACE2 substrate widely used in established ACE2 activity detection assays (FIG

5)[41, 42, 62], and a SARS-CoV-2 spike peptide pool in a competition/inhibition assay (FIG

6), which has been used to demonstrate immunoglobulin-associated ACE2-like activity

in clinical samples[41, 42.]. The peptide pool inhibited the substrate cleavage of two mAbs

(AGRD013-4G9 and AGRD016-4H2), which blocked ACE2 binding to the virus spike

protein, suggesting that the catalytic sites of the mAbs have sufficient resemblance to

the ACE2 to be inhibited by the overlapping spike-derived peptide pool. However, two

mAbs (AGRD016-2B3 and AGRD021-9A10) enhance spike protein binding to ACE2,

indicating the mAbs have distinct structural and functional features. ACE2 substate

cleavage activity was removed after IgG complete depletion (Table 4 and FIG 7) further

confirmed the association of catalytic activity with immunoglobulin.

The four mAbs with ACE2-like activity exhibit catalytic efficiency (Table 3 and FIG 4) 4

$\log_{10}$ lower than human ACE2 molecule, a known characteristic of other abzymes[16], so

the characteristics of the isolated mAbs with ACE2-like catalytic activity fit well within

those described for other abzymes. The inability of the native ACE2-specific small-

molecule inhibitor, MLN-4760, to inhibit the ACE2-like activity of abzymes (FIG. 8)

suggests that the active site structures of the mAbs differ from those of native ACE2[12,

13, 17]. These experiments contribute to further characterizing the magnitude and

biochemical properties of antibody associated catalytic activity.

Our study had several strengths. We screened many mAbs from four sources for ACE2-

like catalytic activity. Although a large fraction of the tested mAbs did not have ACE2-

like catalytic activity, we found that four isolated anti-SARS-CoV-2 spike mAbs have

ACE2-like catalytic activity. Our previous studies describing immunoglobulin-associated

ACE2-like activity in clinical samples could have had other, unappreciated substances,

other than Abs, responsible for the ACE2-like catalytic activity. The identification of

ACE2-like catalytic activity in mAbs uncontaminated by the many substances present in

clinical plasma samples strongly supports the existence of Abs with ACE2-like catalytic

activity.

It is important to note that the Spike RBD binds to a region on ACE2 that does not

physically fully overlap with its catalytic cleft[64]. This suggests that the ACE2-like activity

we observed is likely not a result of a complete mimicry of the ACE2 active site. It is

more likely that some Spike components may structurally resemble the transition state

of ACE2-mediated peptide hydrolysis. Consequently, antibodies templated by these

Spike motifs, along with the regions that bind ACE2, may possess adventitious catalytic

pockets—a phenomenon previously observed in other autoimmune pathologies such as

systemic lupus erythematosus (SLE) and Hashimoto's thyroiditis[18, 65].

However, our study had certain limitations. By screening pre-established mAb

collections from our collaborators[46, 49-52, 59, 66], the mAbs were produced without clear

linkage to patients with symptoms associated with COVID-19. In particular, the mAbs

were not produced from LC patients with some of the most problematic symptoms. This

prevents correlation of abzymes characteristics with clinical symptoms. In addition, the

motivation behind the production of most of the mAbs was to obtain mAbs capable of

inhibiting SARS-CoV-2 infection of cells. Selecting for a mAb that is highly effective at

blocking the virus from infecting a cell may bias the mAb production process away from

selecting mAbs with catalytic activity, since those mAbs may not be particularly effective

at blocking cell infection. In patients, highly neutralizing Abs may not dominate the total

pool of Abs made in response to SARS-CoV-2 infection[67, 68].

All four mAbs with ACE2-like activity were identified in hybridoma culture supernatants

from a single source; the 129 purified mAbs from the other three sources were showed

no ACE2-like catalytic activity. Several hypotheses, which are not mutually exclusive,

could account for this pattern. The most straightforward is that differences in screening

and selection methodology enriched the Precision Antibody panel for clones

predisposed to abzyme activity. Those mAbs were specifically selected for their ability

to modulate ACE2 binding to the spike RBD — a criterion that favors isolation of

antibodies whose combining sites are structurally complementary to the ACE2-RBD

interface, which is precisely the structural prerequisite for ACE2-like catalytic activity.

The mAbs from the other three sources were selected under different criteria, primarily

neutralization potency or epitope coverage, which optimize for different features, which

may be less likely to include catalytic sites. A second, independent hypothesis is that

mAb purification procedures disrupted catalytic activity in the other panels. Low-pH

Protein A/G elution steps can alter antibody conformation, and a catalytic pocket, which

depends on precise active-site geometry, may be more conformationally labile than an

antigen-binding site, which is governed primarily by CDR contacts that are generally

robust in purification. Finally, the differences in panel size and format across sources

mean that statistical sampling cannot be excluded as a partial contributor — the

probability of detecting rare abzyme-positive clones is lower in small panels, and is

further reduced if purification suppresses activity. Importantly, none of these

considerations undermine the validity of the positive supernatant results, which satisfy

the same criteria used in our prior clinical studies: the activity co-depletes with

immunoglobulin at greater than 99.99% efficiency and is inhibited by competition with a

tiled overlapping RBD peptide pool, establishing that it is antibody-associated and

antigen-specific. The ACE2-like catalytic activity of the hybridoma supernatants did not

require divalent cations (was not inhibited by EDTA), so the activity is distinct from

native ACE2. The low catalytic efficiency observed here is consistent with the broader

abzyme literature, in which antibody associated catalysis typically displays slow

turnover, modest substrate affinity, and considerable heterogeneity[22, 69-71]. Previous

well-documented abzymes from autoimmune diseases like Systemic Lupus

Erythematosus (SLE) and Multiple Sclerosis exhibited significantly lower catalytic

efficiency than that of dedicated nucleases or proteases, yet the abzymes remain

fundamentally pathogenic and leading to chronic tissue damage due to their ability to

work under physiological conditions[65, 72]. Future studies should prioritize production and

testing of mAbs from LC patients not selected for neutralization activity, evaluated in

both supernatant and purified formats under varying purification conditions, to

systematically disentangle these hypotheses.

Native enzymes are strictly regulated by localization and inhibitors, in contrast, IgG-

based abzymes circulate with a long biological half-life and lack the features that help

control the physiological regulatory cascade pathways like the renin-angiotensin

system[20, 22, 23]. The cumulative effect over months of disease progression as occurs in

LC or autoimmune diseases could be substantial If these abzymes similarly cleave

native substrate under physiological condition, even at a fraction of the rate of native

enzymes[73-76]. In a patient with high titers of these catalytic IgGs, the constant,

unregulated processing of signaling peptides leading to a significant integrated deficit in

homeostatic peptide levels over months could contribute to the vascular and

inflammatory dysregulation frequently observed in Long COVID[77] [34].

The data presented here established that catalytic activity can be detected in rare anti-

Spike antibodies under defined *in vitro* conditions using a highly sensitive model

substrate. In future studies, to detect these low catalytic activities it may be important to

ensure that the antibodies are maintained in an physiological intact state during isolation,

since it is conceivable that binding or infection inhibition may be less fragile properties

that catalytic activity[78-81], which could be a conceivable reason why catalytic activity was

not detected in the purified mAbs. Nevertheless, the identification of any mAbs with

ACE2-like catalytic activity supports the hypothesis underlying this study.

Another limitation of this work is that catalytic activity was assessed using a short

synthetic fluorogenic peptide, Mca-APK(Dnp), rather than a native biological substrate.

While this synthetic substrate is cleaved by ACE2 and is useful for detecting low level

metalloprotease like activity, it differs substantially from endogenous ACE2 substrates

such as angiotensin II in size, sequence context, and structural constraints [82]. This

substrate was specifically engineered to mimic the C-terminal sequence of natural

ACE2 substrates and has been validated as a highly sensitive and selective surrogate

for ACE2 carboxypeptidase activity[44, 45]. Mca-APK(Dnp) hydrolysis has been shown to

follow the same zinc- and chloride-dependent mechanisms as the natural substrate,

Angiotensin II[83, 84]. Its use in this study reflects its established utility as a sensitive

surrogate for real-time and continuous kinetic monitoring ACE2 class metalloprotease

activity, which is essential for the precise determination of *Vmax* and *Kcat* in low-activity

antibody preparations where endpoint-based HPLC methods may lack sufficient

sensitivity[84-90]. We noted that these analyses were designed to assess substrate

preference rather than absolute specificity and did not exclude the possibility that

additional substrates may also be cleaved at lower efficiency. A comprehensive

definition of substrate specificity would require systematic testing across diverse peptide

libraries and protein substrates in future studies, which was beyond the scope of this

study.

An essential next step involves elucidating the mechanistic properties of the abzymes'

catalytic activity. Structural analyses aimed at delineating these activities could provide

critical insight into the mechanisms of ACE2-like catalytic activity. A clear understanding

of the catalytic mechanisms of the mAbs with ACE2-like catalytic activity could enable

the development of inhibitors against that activity, which would represent a potential

novel avenue for LC therapeutics development. Alternatively, the source of pathologic

antibodies, the cells producing the pathologic abzymes, could be targeted.

Overall, our study further confirms the existence of anti-spike antibodies with ACE2-like

activity that can plausibly contribute to the pathogenesis of COVID-19 and LC. While

much work remains to be done, including the production and study of new anti-spike

mAbs not selected specifically for SARS-CoV-2 neutralizing activity, structural studies of

mAbs with catalytic activity, and the investigation of catalytic activities other than ACE2

in Abs made after exposure to SARS-CoV-2 that could potentially mediate the

pathogenesis of COVID-19 and LC. Additional studies in understand COVD-19 and LC

pathogenesis would include work aimed at associating different clinical features of

disease with abzymes that may have particular characteristics, structural and

mechanistic studies of the abzymes themselves, and larger epidemiologic survey

studies to better determine the rates of abzyme production in persons exposed to

SARS-CoV-2.

There are also longer-term translational implications. Other viruses, in addition to

SARS-CoV-2, use cell surface enzymes as their receptors [91-96]. If pathologic abzymes

induced by exposure to SARS-CoV-2 are responsible for some of the pathologic

features of COVID-19 and LC, then abzymes induced by other viruses, including viruses

causing subclinical infections, could be responsible for other disorders.

Institutional Review Board Statement: The UVA IRB approved enrollment of all

subjects and collection of specimens and related metadata (HSR #HSR13166).

Informed Consent Statement: Informed consent was obtained from the subjects in the

study.

Data Availability Statement: Data is presented in the manuscript. Raw data is

available upon request.

Acknowledgments and Funding: The study was supported by University of Virginia

internal funding, including the Pendleton Laboratory Fund for Pediatric Infectious

Disease Research, the Manning Fund for COVID-19 Research at UVA, the Ivy

Foundation, University of Virginia Summer Research Internship Program and grants

from NIAID, NIH (R01 AI176515). Some data for this study were generated using

instrumentation supported by the UVA Comprehensive Cancer Center (P30-

CA044579). The authors acknowledge Research Computing at The University of

Virginia for providing computational resources and technical support that have

contributed to the results reported within this publication. URL: https://rc.virginia.edu.

This research was supported in part by the Intramural Research Program of the

National Institutes of Health (NIH). The contributions of the NIH author(s) were made as

part of their official duties as NIH federal employees, are in compliance with agency

policy requirements, and are considered Works of the United States Government.

However, the findings and conclusions presented in this paper are those of the

author(s) and do not necessarily reflect the views of the NIH or the U.S. Department of

Health and Human Services. AI agents (Claude, Copilot, Edison Scientific, Gemini)

were use code-writing, debugging, and optimization; literature searches and finding

source identification, manuscript editing, cross-checking, and proof-reading.

Conflicts of Interest: J.T. is a coinventor on a patent filed on the mAb COV44-79 used

in this study.

Tables

Table 1: Summary of Monoclonal Antibody Characteristics Used in the Study

Source	Total Number	Form	Characteristics	Number with Abzyme Activity	Reference
Dr. James Crowe, Vanderbilt University Medical Center	2	Purified	Human Anti SARS-CoV-2 RBM	0	46, 47
Dr. Jun Hayashi, Precision Antibody	3	Supernatant	Human Anti SARS-CoV-2 Spike and RBD, block RBD binding to ACE2	2	
	4	Supernatant	Human Anti SARS-CoV-2 Spike and RBD, enhance RBD binding to ACE2	2	
	3	Supernatant	Negative Controls	0	
Dr. David Ho, Columbia University	39	Purified	Human Anti SARS-CoV-2 RBD	0	49, 59, 97

Dr. Joshua Tan at NIH	88	Purified	Human Anti SARS-CoV-2 RBD	0	52
	1	Purified	Human Anti SARS-CoV-2 FP (Negative Control)	0	51

577
578

579 **Table 2:** Synthesized SARS-CoV-2 Spike RBD Peptides for ACE2 Substrate
580 Competition Assay

Cat #	Domain	Sequence (Amino Acid Position)	Concentration (mg/mL)	Reconstitution
SB086-1MG	RBD	319-RVQPTESIVRFPNITNL-335	10	DMSO
SB087-1MG	RBD	336-CPFGEVFNATRF-347	10	DMSO
SB088-1MG	RBD	348-ASVYAWNRKR-357	10	ddH$_2$O
SB089-1MG	RBD	352-AWNRKRISNCVADY-365	10	ddH$_2$O
SB090-1MG	RBD	371-SASFSTFKCYGVSPTKLNDLCFTN-394	10	DMSO
SB091-1MG	RBD	395-VYADSFVIRGDEVRQIAPGQTGKIAD YNYKLPDDFT-430	10	ddH$_2$O
SB092-1MG	RBM	438-SNNLDSKVGGNYNYLYRLFRK-458	10	ddH$_2$O
SB093-1MG	RBM	450-NYLYRLFRKSNLKPFERDISTEIY-473	10	ddH$_2$O
SB094-1MG	RBM	480-CNGVEGFNCYFPLQSYG-496	10	DMSO
SB095-1MG	RBM	500-TNGVGYQPYR-509	10	DMSO
SB096-1MG	RBD	513-LSFELLHA-520	10	DMSO
SB097-1MG	RBD	523-TVCGPKKSTNLVKNKCVNF-541	10	ddH$_2$O

581
582

583
584
585
586
587 **Table 3:** Enzyme Kinetic Studies of ACE2-like Abzymes and Purified Recombinant
588 Human ACE2 Molecule (From FIG 4)
589

Sample	Assay Conc	K_m (μM)	V_{max} (pmol/min)	V_{max} (mol/s)	K_{cat} (s^{-1})	K_{cat}/K_m (M^{-1}s^{-1})
rhACE2[a]	25 ng/mL	43.7	66.9	1.12E-12	3.79E+01	8.68E+05
AGRD013-4G9	5.8 µg/mL	26.9	0.192	3.20E-15	8.27E-04	3.07E+01
AGRD016-2B3	5.6 µg/mL	34.3	0.147	2.45E-15	6.57E-04	1.91E+01
AGRD016-4H2	7.9 µg/mL	38.9	0.295	4.92E-15	9.33E-04	2.40E+01
AGRD021-9A10	5.3 µg/mL	33.9	0.253	4.22E-15	1.19E-03	3.52E+01

590
591 a: Purified recombinant human ACE2 molecule as positive control.
592
593
594

597 **Table 4:** Human IgG Depletion Efficiency by IgG ELISA

| Antibody | Hybridoma culture supernatant IgG concentration (mean ± SD) | | IgG depletion efficiency |
	Pre-IgG depletion (μg/mL)	Post-IgG depletion (ng/mL)	
AGRD013-4A4	32.06 ± 1.17	0.22 ± 0.01	>99.99%
AGRD013-4G9	11.89 ± 0.92	0.20 ± 0.002	>99.99%
AGRD016-2B3	9.20 ± 0.34	0.20 ± 0.019	>99.99%
AGRD016-3H2	12.78 ± 0.93	0.19 ± 0.002	>99.99%
AGRD016-4H2	11.66 ± 0.92	0.25 ± 0.018	>99.99%
AGRD021-9A10	8.33 ± 0.27	0.24 ± 0.033	>99.99%
AGRD021-19C11	9.11 ± 0.56	0.22 ± 0.002	>99.99%

 FIGURES

601
FIG 1. ACE2 Background Activity Inhibition by EDTA in Supernatant Control
602
Media. ACE2 background activity in hybridoma supernatant control media (empty
603
culture media without hybridoma) was detected by the ACE2 assay, indicated by the
604
slope calculated from the kinetic curve. ACE2 background activity was effectively
605
inhibited by adding EDTA to the media. The dashed line shows the cutoff, which was
606
the limitation of the ACE2 assay and was determined using a purified recombinant
607
human ACE2 positive control to call a sample positive for ACE2 activity. 5mM EDTA
608
(**Triangle**) were used for following hybridoma culture supernatants screening as
609
reported[56].
610

FIG 2. Initial Screening of mAbs for ACE2-like Activity. ACE2 substrate cleavage

activity of mAbs. Catalytic activity was determined by calculating the slope of RFU from

the initial 2 hours of the kinetics curve. 0.08 ng of purified recombinant human ACE2

served as a positive control and standard to define ACE2-positive activity. mAbs with

ACE2-like activity greater than or equal to 0.08ng rhACE2 were identified as having

ACE2-like catalytic activity. (A) Representative ACE2 activity screening kinetic curves

from the Precision Antibody mAbs and ACE2 activity determination by slope. 0.08 ng

rhACE2 served as a positive and standard control, and medium (empty hybridoma

culture supernatant) served as a negative control. (B) Summary of average ACE2-like

activity of all mAbs screened in this study. Results were from at least 2 rounds of

screening. ACE2 substrate cleavage activity was determined by the slope of the RFU

from the initial 2 hours of the kinetic curve. The dashed line shows the slope calculated

from 0.08 ng of purified recombinant human ACE2, the initial 2-hour kinetic curve.

FIG 3. Correlation of total IgG and anti-RBD antibody. Total IgG concentrations in

mAb hybridoma culture supernatants were measured by human IgG ELISA. Anti-RBD

IgG concentrations of mAb hybridoma culture supernatants were determined by

ImmunoCAP antibody-binding assay. (Pearson correlation test).

FIG 4. Michaelis-Menten and Lineweaver-Burk Analysis of mAb Catalytic Activity.

Michaelis-Menten plots for (A) Positive control (2.5ng rhACE2) and (B) mAbs that had

fluorogenic ACE2 surrogate peptide substrate cleavage activities. Lineweaver-Burk plot

for (C) Positive control (2.5ng rhACE2) and (D) mAbs that had ACE2 substrate

cleavage activities. Abzyme IgG amount for catalytic kinetics was from human IgG

ELISA.

FIG 5. Comparative Substrate Preference of Abzymes. The substrate cleavage

activity of the mAbs was characterized by comparing using a fluorogenic ACE2 peptide

substrate, MCA⬚APK(Dnp), and a control non-cleavable substrate. The slope of the

RFU from the initial 2 hours of the kinetic curve was calculated to determine ACE2-like

activity. No catalytic activity was observed with the control substrate indicated by

negative slope.

FIG 6. SARS-CoV-2 RBD Peptide Competitive Inhibition of mAb ACE2 Substrate

Cleavage Activity. mAbs ACE2-like activity inhibition by RBD peptide competition was

performed by adding RBD peptide pools to ACE2 assay reactions containing mAbs

defined as ACE2-like activity-positive. Peptide pool diluent, with an equal volume of

peptide pool added to the assay reactions, served as a comparison. 2.5 ng purified

recombinant human ACE2 was used as a positive control. ACE2-like activity was

determined by calculation of the slope of the RFU from the initial 2 hours of the kinetic

curve.

664
**FIG 7. ACE2 Substrate Cleavage Activity of mAb Hybridoma Culture Supernatants
before and after Immunoglobulin Depletion.** ACE2 substrate cleavage activities of
mAb hybridoma culture supernatants before (Pre) and after (Post) immunoglobulin
depletion were parallel compared. The slope of the RFU from the initial 2 hours of the
kinetic curve was calculated to determine ACE2-like activity. The dashed line shows the
cutoff, which was the limitation of the ACE2 assay and was determined using a purified
recombinant human ACE2 positive control to call a sample positive for ACE2 activity.
IgG depletion efficiency was determined by human IgG ELISA (Table 4). No catalytic
activity was observed from IgG complete depleted samples indicated by negative slope
of kinetic curve.

FIG 8. Effect of ACE2 Inhibitor, MLN-4760, on ACE2 Substrate Cleavage Activity of mAbs and rhACE2. 100 M ACE2 inhibitor, MLN-4760, was added to ACE2 assay reactions for comparison of ACE2 activity with ACE2 assay reactions that had no inhibitor. 2.5 ng purified recombinant human ACE2 was the positive control, and control media (empty hybridoma culture supernatant) was the negative control. ACE2-like activity was determined by calculation of the slope of the RFU from the initial 2 hours of the kinetic curve.

References

1. Altmann DM, Whettlock EM, Liu S, Arachchillage DJ, Boyton RJ. The immunology of long COVID. Nature Reviews Immunology. 2023;23(10):618-34.

2. Perumal R, Shunmugam L, Naidoo K, Abdool Karim SS, Wilkins D, Garzino-Demo A, Brechot C, Parthasarathy S, Vahlne A, Nikolich JŽ. Long COVID: a review and proposed visualization of the complexity of long COVID. Frontiers in immunology. 2023;14:1117464.

3. Davis HE, McCorkell L, Vogel JM, Topol EJ. Long COVID: major findings, mechanisms and recommendations. Nature Reviews Microbiology. 2023;21(3):133-46.

4. Adjaye-Gbewonyo D, Vahratian A, Perrine CG, Bertolli J. Long COVID in Adults: United States, 2022. 26/2023.

5. Marchi M, Grenzi P, Serafini V, Capoccia F, Rossi F, Marrino P, Pingani L, Galeazzi GM, Ferrari S. Psychiatric symptoms in Long-COVID patients: a systematic review. Frontiers in Psychiatry. 2023;14:1138389.

6. Thaweethai T, Jolley SE, Karlson EW, Levitan EB, Levy B, McComsey GA, McCorkell L, Nadkarni GN, Parthasarathy S, Singh U, Walker TA, Selvaggi CA, Shinnick DJ, Schulte CCM, Atchley-Challenner R, Horwitz LI, Foulkes AS, Authors RC, Consortium R. Development of a Definition of Postacute Sequelae of SARS-CoV-2 Infection. JAMA. 2023;329(22):1934-46.

7. Soriano JB, Murthy S, Marshall JC, Relan P, Diaz JV. A clinical case definition of post-COVID-19 condition by a Delphi consensus. Lancet Infect Dis. 2022 Apr;22(4):e102-e7. PMC8691845

8. Castanares-Zapatero D, Chalon P, Kohn L, Dauvrin M, Detollenaere J, Maertens de Noordhout C, Primus-de Jong C, Cleemput I, Van den Heede K. Pathophysiology and mechanism of long COVID: a comprehensive review. Ann Med. 2022 Dec;54(1):1473-87. PMC9132392

9. Tramontano A, Janda KD, Lerner RA. Catalytic antibodies. Science. 1986 Dec 19;234(4783):1566-70.

10. Pollack SJ, Jacobs JW, Schultz PG. Selective chemical catalysis by an antibody. Science. 1986 Dec 19;234(4783):1570-3.

11. Avalle B, Mistro D, Thomas D, Friboulet A. Polyclonal catalytic anti-idiotypic antibodies with a beta-lactamase activity. Ann N Y Acad Sci. 1996 Oct 12;799:172-5.

12. Friboulet A, Izadyar L, Avalle B, Roseto A, Thomas D. Abzyme generation using an anti-idiotypic antibody as the "internal image" of an enzyme active site. Appl Biochem Biotechnol. 1994 May-Jun;47(2-3):229-37; discussion 37-9.

734 13.   Izadyar L, Friboulet A, Remy MH, Roseto A, Thomas D. Monoclonal anti-idiotypic
735 antibodies as functional internal images of enzyme active sites: production of a catalytic
736 antibody with a cholinesterase activity. Proc Natl Acad Sci U S A. 1993 Oct
737 1;90(19):8876-80. PMC47463
738
739 14.   Friboulet A, Izadyar L, Avalle B, Roseto A, Thomas D. Antiidiotypic antibodies as
740 functional internal images of enzyme-active sites. Ann N Y Acad Sci. 1995 Mar
741 31;750:265-70.
742
743 15.   Hilvert D. Critical analysis of antibody catalysis. Annu Rev Biochem.
744 2000;69:751-93.
745
746 16.   Bar-Even A, Noor E, Savir Y, Liebermeister W, Davidi D, Tawfik DS, Milo R. The
747 moderately efficient enzyme: evolutionary and physicochemical trends shaping enzyme
748 parameters. Biochemistry. 2011 May 31;50(21):4402-10.
749
750 17.   Ponomarenko NA, Pillet D, Paon M, Vorobiev, II, Smirnov IV, Adenier H, Avalle
751 B, Kolesnikov AV, Kozyr AV, Thomas D, Gabibov AG, Friboulet A. Anti-idiotypic
752 antibody mimics proteolytic function of parent antigen. Biochemistry. 2007 Dec
753 18;46(50):14598-609.
754
755 18.   Nevinsky GA, Buneva VN. Catalytic antibodies in healthy humans and patients
756 with autoimmune and viral diseases. J Cell Mol Med. 2003 Jul-Sep;7(3):265-76.
757 PMC6741318
758
759 19.   Paul S, Planque SA, Nishiyama Y, Hanson CV, Massey RJ. Nature and nurture
760 of catalytic antibodies. Adv Exp Med Biol. 2012;750:56-75.
761
762 20.   Nevinsky GA. Immune Response and Production of Abzymes in Patients with
763 Autoimmune and Neurodegenerative Diseases. Biochemistry (Mosc). 2025
764 Jan;90(Suppl 1):S373-s400.
765
766 21.   Shokat KM, Schultz PG. Catalytic antibodies. Annu Rev Immunol. 1990;8:335-
767 63.
768
769 22.   Nevinsky GA, Kanyshkova TG, Buneva VN. Natural catalytic antibodies
770 (abzymes) in normalcy and pathology. Biochemistry (Mosc). 2000 Nov;65(11):1245-55.
771
772 23.   Zhao D, Chen J, Hu X, Zhang S. Catalytic Antibodies: Design, Expression, and
773 Their Applications in Medicine. Appl Biochem Biotechnol. 2023 Feb;195(2):1514-40.
774 PMC9554387
775
776 24.   Klegerman ME, Peng T, Seferovich I, Rahbar MH, Hessabi M, Tahanan A,
777 Wanger A, Grimes CZ, Ostrosky-Zeichner LZ, Koster K, Cirillo JD, Abeydeera D, De
778 Lira S, McPherson DD. Absolute concentration estimation of COVID-19 convalescent
779 and post-vaccination IgG antibodies. PLoS One. 2024;19(11):e0311777. PMC11530011
780
25.   Kuah E, Toh S, Yee J, Ma Q, Gao Z. Enzyme Mimics: Advances and
Applications. Chemistry. 2016 Jun 13;22(25):8404-30.

26.   Urusov AE, Tolmacheva AS, Aulova KS, Nevinsky GA. Autoantibody-Abzymes
with Catalase Activity in Experimental Autoimmune Encephalomyelitis Mice. Molecules.
2023 Jan 30;28(3). PMC9921166

27.   South AM, Tomlinson L, Edmonston D, Hiremath S, Sparks MA. Controversies of
renin-angiotensin system inhibition during the COVID-19 pandemic. Nat Rev Nephrol.
2020 Jun;16(6):305-7. PMC7118703

28.   Al-Merani SA, Brooks DP, Chapman BJ, Munday KA. The half-lives of
angiotensin II, angiotensin II-amide, angiotensin III, Sar1-Ala8-angiotensin II and renin
in the circulatory system of the rat. J Physiol. 1978 May;278:471-90. PMC1282363

29.   Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X,
Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L,
Xie J, Wang G, Jiang R, Gao Z, Jin Q, Wang J, Cao B. Clinical features of patients
infected with 2019 novel coronavirus in Wuhan, China. Lancet. 2020 Feb
15;395(10223):497-506. PMC7159299

30.   Liao D, Zhou F, Luo L, Xu M, Wang H, Xia J, Gao Y, Cai L, Wang Z, Yin P, Wang
Y, Tang L, Deng J, Mei H, Hu Y. Haematological characteristics and risk factors in the
classification and prognosis evaluation of COVID-19: a retrospective cohort study.
Lancet Haematol. 2020 Sep;7(9):e671-e8. PMC7351397

31.   Del Valle DM, Kim-Schulze S, Huang HH, Beckmann ND, Nirenberg S, Wang B,
Lavin Y, Swartz TH, Madduri D, Stock A, Marron TU, Xie H, Patel M, Tuballes K, Van
Oekelen O, Rahman A, Kovatch P, Aberg JA, Schadt E, Jagannath S, Mazumdar M,
Charney AW, Firpo-Betancourt A, Mendu DR, Jhang J, Reich D, Sigel K, Cordon-Cardo
C, Feldmann M, Parekh S, Merad M, Gnjatic S. An inflammatory cytokine signature
predicts COVID-19 severity and survival. Nat Med. 2020 Oct;26(10):1636-43.
PMC7869028

32.   Sun DS, Lien TS, Chang HH. Virus-Induced Pathogenic Antibodies: Lessons
from Long COVID and Dengue Hemorrhage Fever. Int J Mol Sci. 2025 Feb 22;26(5).
PMC11899886

33.   Santos Guedes de Sa K, Silva J, Bayarri-Olmos R, Brinda R, Alec Rath
Constable R, Colom Diaz PA, Kwon DI, Rodrigues G, Wenxue L, Baker C,
Bhattacharjee B, Wood J, Tabacof L, Liu Y, Putrino D, Horvath TL, Iwasaki A. A causal
link between autoantibodies and neurological symptoms in long COVID. medRxiv. 2024
Jun 19. PMC11213106

34.   Chen H-J, Appelman B, Willemen HLDM, Bos A, Prado J, Mak WA, Keijzer N,
Santos Ribeiro PS, Goncalves SV, Versteeg S, Geyer CE, Larsen M, Schüchner E,
Bomers MK, Lavell AHA, Charlton B, Wüst R, Wiersinga WJ, van Vugt M, Vidarsson G,
Eijkelkamp N, den Dunnen J. Transfer of IgG from long COVID patients induces
symptomology in mice. Cell Reports Medicine. 2026 2026/03/24/:102693.

35.   Yousaf AR, Cortese MM, Taylor AW, Broder KR, Oster ME, Wong JM, Guh AY,
McCormick DW, Kamidani S, Schlaudecker EP, Edwards KM, Creech CB, Staat MA,
Belay ED, Marquez P, Su JR, Salzman MB, Thompson D, Campbell AP. Reported
cases of multisystem inflammatory syndrome in children aged 12-20 years in the USA
who received a COVID-19 vaccine, December, 2020, through August, 2021: a
surveillance investigation. Lancet Child Adolesc Health. 2022 Feb 22. PMC8864018
837
838   36.   Wilhelm F, Cadamuro J, Mink S. Autoantibodies in long COVID: a systematic
839   review. Lancet Infect Dis. 2026 Apr;26(4):e220-e30.
840
841   37.   Talwar S, Harker JA, Openshaw PJM, Thwaites RS. Autoimmunity in long
842   COVID. J Allergy Clin Immunol. 2025 Apr;155(4):1082-94.
843
844   38.   Khurana V, Goswami B. Angiotensin converting enzyme (ACE). Clin Chim Acta.
845   2022 Jan 1;524:113-22.
846
847   39.   Turner AJ, Nalivaeva NN. Angiotensin-converting enzyme 2 (ACE2): Two
848   decades of revelations and re-evaluation. Peptides. 2022 May;151:170766.
849   PMC8830188
850
851   40.   Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition
852   of SARS-CoV-2 by full-length human ACE2. Science. 2020 Mar 27;367(6485):1444-8.
853   PMC7164635
854
855   41.   Song Y, Mehl F, Muehling LM, Canderan G, Enfield K, Sun J, Yin MT, Ratcliffe
856   SJ, Wilson JM, Kadl A, Woodfolk JA, Zeichner SL. ACE-2-like enzymatic activity in
857   COVID-19 convalescents with persistent pulmonary symptoms associated with
858   immunoglobulin. mBio. 2025 Jul 22:e0173525.
859
860   42.   Song Y, Myers R, Mehl F, Murphy L, Brooks B, Wilson JM, Kadl A, Woodfolk J,
861   Zeichner SL. ACE-2-like enzymatic activity is associated with immunoglobulin in
862   COVID-19 patients. mBio. 2024 Apr 10;15(4):e0054124. PMC11005375
863
864   43.   Mendiola-Salazar XA, Munguía-Laguna MA, Franco M, Cano-Martínez A,
865   Santamaría Sosa J, Bautista-Pérez R. SARS-CoV-2 Spike Protein Enhances
866   Carboxypeptidase Activity of Angiotensin-Converting Enzyme 2. Int J Mol Sci. 2024 Jun
867   6;25(11). PMC11172802
868
869   44.   Vickers C, Hales P, Kaushik V, Dick L, Gavin J, Tang J, Godbout K, Parsons T,
870   Baronas E, Hsieh F, Acton S, Patane M, Nichols A, Tummino P. Hydrolysis of biological
871   peptides by human angiotensin-converting enzyme-related carboxypeptidase. J Biol
872   Chem. 2002 Apr 26;277(17):14838-43.
873
874   45.   Dales NA, Gould AE, Brown JA, Calderwood EF, Guan B, Minor CA, Gavin JM,
875   Hales P, Kaushik VK, Stewart M, Tummino PJ, Vickers CS, Ocain TD, Patane MA.
876   Substrate-based design of the first class of angiotensin-converting enzyme-related
877   carboxypeptidase (ACE2) inhibitors. J Am Chem Soc. 2002 Oct 9;124(40):11852-3.
878

46.   Zost SJ, Gilchuk P, Case JB, Binshtein E, Chen RE, Nkolola JP, Schäfer A,
Reidy JX, Trivette A, Nargi RS, Sutton RE, Suryadevara N, Martinez DR, Williamson
LE, Chen EC, Jones T, Day S, Myers L, Hassan AO, Kafai NM, Winkler ES, Fox JM,
Shrihari S, Mueller BK, Meiler J, Chandrashekar A, Mercado NB, Steinhardt JJ, Ren K,
Loo YM, Kallewaard NL, McCune BT, Keeler SP, Holtzman MJ, Barouch DH, Gralinski
LE, Baric RS, Thackray LB, Diamond MS, Carnahan RH, Crowe JE, Jr. Potently
neutralizing and protective human antibodies against SARS-CoV-2. Nature. 2020
Aug;584(7821):443-9. PMC7584396
887
888   47.   Greaney AJ, Starr TN, Gilchuk P, Zost SJ, Binshtein E, Loes AN, Hilton SK,
889   Huddleston J, Eguia R, Crawford KHD, Dingens AS, Nargi RS, Sutton RE, Suryadevara
890   N, Rothlauf PW, Liu Z, Whelan SPJ, Carnahan RH, Crowe JE, Jr., Bloom JD. Complete
891   Mapping of Mutations to the SARS-CoV-2 Spike Receptor-Binding Domain that Escape
892   Antibody Recognition. Cell Host Microbe. 2021 Jan 13;29(1):44-57.e9. PMC7676316
893
894   48.   Cerutti G, Guo Y, Liu L, Liu L, Zhang Z, Luo Y, Huang Y, Wang HH, Ho DD,
895   Sheng Z, Shapiro L. Cryo-EM structure of the SARS-CoV-2 Omicron spike. Cell Rep.
896   2022 Mar 1;38(9):110428. PMC8818377
897
49.   Liu L, Iketani S, Guo Y, Reddem ER, Casner RG, Nair MS, Yu J, Chan JF, Wang
M, Cerutti G, Li Z, Morano NC, Castagna CD, Corredor L, Chu H, Yuan S, Poon VK,
Chan CC, Chen Z, Luo Y, Cunningham M, Chavez A, Yin MT, Perlin DS, Tsuji M, Yuen
KY, Kwong PD, Sheng Z, Huang Y, Shapiro L, Ho DD. An antibody class with a
common CDRH3 motif broadly neutralizes sarbecoviruses. Sci Transl Med. 2022 May
25;14(646):eabn6859. PMC9017343
904
905   50.   Liu L, Casner RG, Guo Y, Wang Q, Iketani S, Chan JF, Yu J, Dadonaite B, Nair
906   MS, Mohri H, Reddem ER, Yuan S, Poon VK, Chan CC, Yuen KY, Sheng Z, Huang Y,
907   Bloom JD, Shapiro L, Ho DD. Antibodies targeting a quaternary site on SARS-CoV-2
908   spike glycoprotein prevent viral receptor engagement by conformational locking.
909   Immunity. 2023 Oct 10;56(10):2442-55.e8. PMC10588992
910
51.   Dacon C, Tucker C, Peng L, Lee CD, Lin TH, Yuan M, Cong Y, Wang L, Purser
L, Williams JK, Pyo CW, Kosik I, Hu Z, Zhao M, Mohan D, Cooper A, Peterson M,
Skinner J, Dixit S, Kollins E, Huzella L, Perry D, Byrum R, Lembirik S, Zhang Y, Yang
ES, Chen M, Leung K, Weinberg RS, Pegu A, Geraghty DE, Davidson E, Douagi I, Moir
S, Yewdell JW, Schmaljohn C, Crompton PD, Holbrook MR, Nemazee D, Mascola JR,
Wilson IA, Tan J. Broadly neutralizing antibodies target the coronavirus fusion peptide.
bioRxiv. 2022 Apr 12:2022.04.11.487879. PMC9016638
918
52.   Cho H, Gonzales-Wartz KK, Huang D, Yuan M, Peterson M, Liang J, Beutler N,
Torres JL, Cong Y, Postnikova E, Bangaru S, Talana CA, Shi W, Yang ES, Zhang Y,
Leung K, Wang L, Peng L, Skinner J, Li S, Wu NC, Liu H, Dacon C, Moyer T, Cohen M,
Zhao M, Lee FE, Weinberg RS, Douagi I, Gross R, Schmaljohn C, Pegu A, Mascola JR,
Holbrook M, Nemazee D, Rogers TF, Ward AB, Wilson IA, Crompton PD, Tan J.
Bispecific antibodies targeting distinct regions of the spike protein potently neutralize
SARS-CoV-2 variants of concern. Sci Transl Med. 2021 Oct 20;13(616):eabj5413.
PMC8651051
927

53.   Hifumi E, Ito Y, Tsujita M, Taguchi H, Uda T. Enzymatization of mouse
monoclonal antibodies to the corresponding catalytic antibodies. Sci Rep. 2024 May
28;14(1):12184. PMC11133420

54.   McConnell SA, Sachithanandham J, Mudrak NJ, Zhu X, Farhang PA, Cordero
RJB, Wear MP, Shapiro JR, Park HS, Klein SL, Tobian AAR, Bloch EM, Sullivan DJ,
Pekosz A, Casadevall A. Spike-protein proteolytic antibodies in COVID-19 convalescent
plasma contribute to SARS-CoV-2 neutralization. Cell Chem Biol. 2023 Jul
20;30(7):726-38.e4. PMC10288624

55.   Xiao F, Burns KD. Measurement of Angiotensin Converting Enzyme 2 Activity in
Biological Fluid (ACE2). Methods Mol Biol. 2017;1527:101-15. PMC7121061

56.   Lubel JS, Herath CB, Velkoska E, Casley DJ, Burrell LM, Angus PW. Angiotensin
converting enzyme 2 (ACE2) activity in fetal calf serum: implications for cell culture
research. Cytotechnology. 2008 Nov;58(3):119-26. PMC2652558

57.   Ferrari M, Mekkaoui L, Ilca FT, Akbar Z, Bughda R, Lamb K, Ward K, Parekh F,
Karattil R, Allen C, Wu P, Baldan V, Mattiuzzo G, Bentley EM, Takeuchi Y, Sillibourne J,
Datta P, Kinna A, Pule M, Onuoha SC. Characterization of a Novel ACE2-Based
Therapeutic with Enhanced Rather than Reduced Activity against SARS-CoV-2
Variants. Journal of Virology. 2021;95(19):10.1128/jvi.00685-21.

58.   Svilenov HL, Delhommel F, Siebenmorgen T, Rührnößl F, Popowicz GM, Reiter
A, Sattler M, Brockmeyer C, Buchner J. Extrinsic stabilization of antiviral ACE2-Fc
fusion proteins targeting SARS-CoV-2. Commun Biol. 2023 Apr 8;6(1):386.
PMC10082628

59.   Liu L, Wang P, Nair MS, Yu J, Rapp M, Wang Q, Luo Y, Chan JF, Sahi V,
Figueroa A, Guo XV, Cerutti G, Bimela J, Gorman J, Zhou T, Chen Z, Yuen KY, Kwong
PD, Sodroski JG, Yin MT, Sheng Z, Huang Y, Shapiro L, Ho DD. Potent neutralizing
antibodies against multiple epitopes on SARS-CoV-2 spike. Nature. 2020
Aug;584(7821):450-6.

60.   Michaelis L, Menten ML, Johnson KA, Goody RS. The original Michaelis
constant: translation of the 1913 Michaelis-Menten paper. Biochemistry. 2011 Oct
4;50(39):8264-9. PMC3381512

61.   Rice GI, Thomas DA, Grant PJ, Turner AJ, Hooper NM. Evaluation of
angiotensin-converting enzyme (ACE), its homologue ACE2 and neprilysin in
angiotensin peptide metabolism. Biochem J. 2004 Oct 1;383(Pt 1):45-51. PMC1134042

62.   Ferrari M, Mekkaoui L, Ilca FT, Akbar Z, Bughda R, Lamb K, Ward K, Parekh F,
Karattil R, Allen C, Wu P, Baldan V, Mattiuzzo G, Bentley EM, Takeuchi Y, Sillibourne J,
Datta P, Kinna A, Pule M, Onuoha SC. Characterization of a Novel ACE2-Based
Therapeutic with Enhanced Rather than Reduced Activity against SARS-CoV-2
Variants. J Virol. 2021 Sep 9;95(19):e0068521. PMC8432736
975

976 63.   Organization WH. WHO COVID-19 Dashboard. World Health Organization; 2026
977 [cited 2026 14 January 2026]; https://data.who.int/dashboards/covid19/deaths.
978 64.   Lan J, Ge J, Yu J, Shan S, Zhou H, Fan S, Zhang Q, Shi X, Wang Q, Zhang L,
979 Wang X. Structure of the SARS-CoV-2 spike receptor-binding domain bound to the
980 ACE2 receptor. Nature. 2020 May;581(7807):215-20.
981
65.   Ponomarenko NA, Durova OM, Vorobiev, II, Belogurov AA, Jr., Kurkova IN,
Petrenko AG, Telegin GB, Suchkov SV, Kiselev SL, Lagarkova MA, Govorun VM,
Serebryakova MV, Avalle B, Tornatore P, Karavanov A, Morse HC, 3rd, Thomas D,
Friboulet A, Gabibov AG. Autoantibodies to myelin basic protein catalyze site-specific
degradation of their antigen. Proc Natl Acad Sci U S A. 2006 Jan 10;103(2):281-6.
PMC1324791
988
989 66.   Greaney AJ, Starr TN, Gilchuk P, Zost SJ, Binshtein E, Loes AN, Hilton SK,
990 Huddleston J, Eguia R, Crawford KHD, Dingens AS, Nargi RS, Sutton RE, Suryadevara
991 N, Rothlauf PW, Liu Z, Whelan SPJ, Carnahan RH, Crowe JE, Jr., Bloom JD. Complete
992 Mapping of Mutations to the SARS-CoV-2 Spike Receptor-Binding Domain that Escape
993 Antibody Recognition. Cell Host Microbe. 2020 Nov 19. PMC7676316
994
995 67.   de Souza MO, Madan B, Teng IT, Huang A, Liu L, Fahad AS, Lopez Acevedo
996 SN, Pan X, Sastry M, Gutierrez-Gonzalez M, Yin MT, Zhou T, Ho DD, Kwong PD,
997 DeKosky BJ. Mapping monoclonal anti-SARS-CoV-2 antibody repertoires against
998 diverse coronavirus antigens. Front Immunol. 2022;13:977064. PMC9478573
999
68.   Amanat F, Thapa M, Lei T, Sayed Ahmed SM, Adelsberg DC, Carreno JM,
Strohmeier S, Schmitz AJ, Zafar S, Zhou JQ, Rijnink W, Alshammary H, Borcherding N,
Reiche AG, Srivastava K, Sordillo EM, van Bakel H, Turner JS, Bajic G, Simon V,
Ellebedy AH, Krammer F. The plasmablast response to SARS-CoV-2 mRNA
vaccination is dominated by non-neutralizing antibodies and targets both the NTD and
the RBD. medRxiv. 2021 May 1. PMC7987037
1006
1007 69.   Tawfik DS, Eshhar Z, Green BS. Catalytic antibodies: a critical assessment. Mol
1008 Biotechnol. 1994 Feb;1(1):87-103.
1009
1010 70.   Lacroix-Desmazes S, Wootla B, Delignat S, Dasgupta S, Nagaraja V,
1011 Kazatchkine MD, Kaveri SV. Pathophysiology of catalytic antibodies. Immunol Lett.
1012 2006 Feb 28;103(1):3-7.
1013
1014 71.   Paul S. Natural catalytic antibodies. Mol Biotechnol. 1996 Jun;5(3):197-207.
1015
1016 72.   Baranovskii AG, Odintsova ES, Buneva VN, Doronin BM, Nevinsky GA.
1017 Comparison of enzymatic properties of DNA-abzymes and human DNAses.
1018 Nucleosides Nucleotides Nucleic Acids. 2004 Oct;23(6-7):1053-6.
1019
1020 73.   Arthur JM, Forrest JC, Boehme KW, Kennedy JL, Owens S, Herzog C, Liu J,
1021 Harville TO. Development of ACE2 autoantibodies after SARS-CoV-2 infection. PLoS
1022 One. 2021;16(9):e0257016. PMC8415618
1023

74.   Sodhi PV, Sidime F, Tarazona DD, Valdivia F, Levano KS. A Closer Look at ACE2 Signaling Pathway and Processing during COVID-19 Infection: Identifying Possible Targets. Vaccines (Basel). 2022 Dec 21;11(1). PMC9867515

75.   Nevinsky GA, Buneva VN. Natural catalytic antibodies in norm, autoimmune, viral, and bacterial diseases. ScientificWorldJournal. 2010 Jun 29;10:1203-33. PMC5763945

76.   Wootla B, Lacroix-Desmazes S, Warrington AE, Bieber AJ, Kaveri SV, Rodriguez M. Autoantibodies with enzymatic properties in human autoimmune diseases. J Autoimmun. 2011 Sep;37(2):144-50. PMC3173522

77.   Santos Guedes de Sa K, Silva J, Bayarri-Olmos R, Brinda R, Alec Rath Constable R, Colom Diaz PA, Kwon D-i, Rodrigues G, Wenxue L, Baker C, Bhattacharjee B, Wood J, Tabacof L, Liu Y, Putrino D, Horvath TL, Iwasaki A. A causal link between autoantibodies and neurological symptoms in long COVID. medRxiv. 2024:2024.06.18.24309100.

78.   Cesaro-Tadic S, Lagos D, Honegger A, Rickard JH, Partridge LJ, Blackburn GM, Plückthun A. Turnover-based in vitro selection and evolution of biocatalysts from a fully synthetic antibody library. Nat Biotechnol. 2003 Jun;21(6):679-85.

79.   Ermakov EA, Kabirova EM, Buneva VN, Nevinsky GA. IgGs-Abzymes from the Sera of Patients with Multiple Sclerosis Recognize and Hydrolyze miRNAs. Int J Mol Sci. 2021 Mar 10;22(6). PMC8000798

80.   Li JW, Xia L, Su Y, Liu H, Xia X, Lu Q, Yang C, Reheman K. Molecular imprint of enzyme active site by camel nanobodies: rapid and efficient approach to produce abzymes with alliinase activity. J Biol Chem. 2012 Apr 20;287(17):13713-21. PMC3340132

81.   Richard JP. Protein Flexibility and Stiffness Enable Efficient Enzymatic Catalysis. J Am Chem Soc. 2019 Feb 27;141(8):3320-31. PMC6396832

82.   Carrera E, Linares A, Joachim A, Jean-Baptiste M, Speth R. MCA-APK(Dnp) is not a selective substrate of angiotensin-converting enzyme-2 (1067.6). The FASEB Journal. 2014 2014/04/01;28(S1):1067.6.

83.   Guy JL, Jackson RM, Acharya KR, Sturrock ED, Hooper NM, Turner AJ. Angiotensin-converting enzyme-2 (ACE2): comparative modeling of the active site, specificity requirements, and chloride dependence. Biochemistry. 2003 Nov 18;42(45):13185-92.

84.   Pedersen KB, Sriramula S, Chhabra KH, Xia H, Lazartigues E. Species-specific inhibitor sensitivity of angiotensin-converting enzyme 2 (ACE2) and its implication for ACE2 activity assays. Am J Physiol Regul Integr Comp Physiol. 2011 Nov;301(5):R1293-9. PMC3213941

85.   Sentandreu MA, Toldrá F. A fluorescence-based protocol for quantifying angiotensin-converting enzyme activity. Nat Protoc. 2006;1(5):2423-7.

86.   Elased KM, Cunha TS, Gurley SB, Coffman TM, Morris M. New mass spectrometric assay for angiotensin-converting enzyme 2 activity. Hypertension. 2006 May;47(5):1010-7.

87.   Meng QC, Berecek KH. Quantification of Angiotensin-Converting Enzyme (ACE) Activity. Methods Mol Med. 2001;51:257-66.

88.   Cruz-Diaz N, Wilson BA, Pirro NT, Brosnihan KB, Marshall AC, Chappell MC. Identification of dipeptidyl peptidase 3 as the Angiotensin-(1-7) degrading peptidase in human HK-2 renal epithelial cells. Peptides. 2016 Sep;83:29-37. PMC5500233

89.   Cheng XC, Fang H, Xu WF. Advances in assays of matrix metalloproteinases (MMPs) and their inhibitors. J Enzyme Inhib Med Chem. 2008 Apr;23(2):154-67.

90.   Haber PK, Ye M, Wysocki J, Maier C, Haque SK, Batlle D. Angiotensin-converting enzyme 2-independent action of presumed angiotensin-converting enzyme 2 activators: studies in vivo, ex vivo, and in vitro. Hypertension. 2014 Apr;63(4):774-82. PMC4004079

91.   Jin M, Jefferson VA, Zhao Z, Catanzaro NJ, De Sabato L, Keller EL, Menasche BL, Hoffman C, Castelli A, Di Bartolo I, Vaccari G, Starr TN, Seifert SN, Castellanos AA, Han BA, Serebryannyy L, Spaulding AB, Douek DC, Moreno A, Baric RS, Wilen CB, Rini JM, Letko M. Aminopeptidase N is a receptor for hedgehog merbecoviruses. bioRxiv. 2025 Sep 3. PMC12425009

92.   Dufloo J, Fernández I, Arbabian A, Haouz A, Temperton N, Gimenez-Lirola LG, Rey FA, Sanjuán R. Dipeptidase 1 is a functional receptor for a porcine coronavirus. Nat Microbiol. 2025 Nov;10(11):2981-96. PMC12578638

93.   Dufloo J, Andreu-Moreno I, Moreno-García J, Valero-Rello A, Sanjuán R. Receptor-binding proteins from animal viruses are broadly compatible with human cell entry factors. Nat Microbiol. 2025 Feb;10(2):405-19. PMC11790484

94.   Simmons G, Zmora P, Gierer S, Heurich A, Pöhlmann S. Proteolytic activation of the SARS-coronavirus spike protein: cutting enzymes at the cutting edge of antiviral research. Antiviral Res. 2013 Dec;100(3):605-14. PMC3889862

95.   López-Cortés GI, Palacios-Pérez M, Hernández-Aguilar MM, Veledíaz HF, José MV. Human Coronavirus Cell Receptors Provide Challenging Therapeutic Targets. Vaccines (Basel). 2023 Jan 13;11(1). PMC9862439

96.   Li F. Receptor recognition mechanisms of coronaviruses: a decade of structural studies. J Virol. 2015 Feb;89(4):1954-64. PMC4338876

97.   Liu L, Casner RG, Guo Y, Wang Q, Iketani S, Chan JF, Yu J, Dadonaite B, Nair MS, Mohri H, Reddem ER, Yuan S, Poon VK, Chan CC, Yuen KY, Sheng Z, Huang Y,

1121   Bloom JD, Shapiro L, Ho DD. Antibodies that neutralize all current SARS-CoV-2
1122   variants of concern by conformational locking. bioRxiv. 2023 Apr 10. PMC10120718
1123
1124