Revegetation following grazing cessation drives sequential shifts in soil

microbial functions and life-history strategies over decadal time scales

Timothy M. Ghaly[1,2]*, Vanessa J. McPherson[1], Vaheesan Rajabal[1,2], Mary E. Ghaly[1], Nicki

Taws[3], Rachael V. Gallagher[4], Johannes J. Le Roux[1], Sasha G. Tetu[1,2]*

[1]School of Natural Sciences, Macquarie University, New South Wales, Australia

[2]ARC Centre of Excellence in Synthetic Biology, New South Wales, Australia

[3]Greening Australia, Australian Capital Territory, Australia

[4]Hawkesbury Institute for the Environment, Western Sydney University, New South Wales,

Australia

*Corresponding Authors:

Timothy M. Ghaly: timothy.ghaly@mq.edu.au

Sasha G. Tetu: sasha.tetu@mq.edu.au

Keywords: biogeochemical cycling; carbon fixation; ecological restoration; land use change;

microbial life history strategies; plant-microbe interactions; soil carbon storage

Abstract

Revegetation is a key strategy for restoring degraded lands globally. While this process can

reshape belowground microbial communities, the extent to which such changes restore soil

ecosystem functions remains less clear. To address this, we applied deep metagenomic

sequencing to characterise soil microbial responses following revegetation (spanning 1-31

years prior to sampling) on grazing agricultural lands. We find that revegetation following the

cessation of livestock grazing was associated with significant shifts in soil microbiome

structure, functional potential, and dominant life history strategies, which were clearly

detectable as early as three years, and persisted >30 years after revegetation. Revegetated

soils were enriched in traits involved in nutrient retention, carbon sequestration, and plant

growth promotion, albeit with possible livestock-specific legacy effects. While shifts in key

biogeochemical cycling functions occurred rapidly (~3 years) and then stabilised, the

enrichment of plant growth-promoting traits was more gradual, suggesting that soil

microbiome responses involve an early restructuring of soil health processes, including

nutrient retention and carbon fixation, followed by a more gradual progression of plant-

microbe interaction traits as the plant community matures. Further, our analyses of over 450

high-purity metagenome-assembled genomes revealed a shift in the prevailing microbial life

history strategies: revegetated soils favoured more diverse biosynthesis, growth, and carbon

fixation strategies, whereas grazing soils selected resource-scavenging and stress tolerance

strategies. Together, our findings suggest that soil microbial communities not only respond to,

but may facilitate, ecological restoration following revegetation.

Introduction

Human land use is a major driver of environmental degradation and ecosystem service loss. Agricultural grazing land, which covers more than 25% of Earth's land surface, is subject to significant degradation through soil compaction and altered nutrient fluxes, both of which reduce plant diversity, and collectively alter ecosystem processes and functioning [1, 2]. Revegetation, the process of restoring plant species in disturbed areas, is a widely adopted strategy for the restoration of degraded lands, with agricultural land representing a key global target for this style of restoration [3]. Although the aboveground outcomes of revegetation are often visible, the impacts on belowground ecosystems, and the bi-directional feedbacks between them, are less clear.

Soil microbiomes are fundamental mediators of ecosystem processes from local to global scales, governing biogeochemical cycles, soil structure and moisture storage, and climate regulation [4-6]. Soil microbiomes drive the cycling of nutrients that underpin ecosystem health, with microbial metabolism tightly linked to aboveground biodiversity [7]. Soil microbial communities also exert strong control over carbon storage and soil-atmosphere gas fluxes through metabolic processes such as carbon and nitrogen fixation, decomposition, and the production or consumption of greenhouse gases [8, 9]. Microbial life history strategies, defined by trade-offs between growth yield, resource acquisition, and stress tolerance, also impact soil carbon dynamics [10]. Specifically, a high *growth yield* strategy promotes soil carbon storage by increasing biomass and subsequently trapping much of this carbon as stable 'necromass'. In contrast, a focus on *resource acquisition* can drive decomposition and carbon loss by breaking down complex organic matter; while greater investment in *stress response* strategies can increase CO_2 respiration, as energy is redirected from biomass production to maintenance and repair. In addition to impacts on

biogeochemical cycling, soil microbiomes can also have significant influence on plant health

and productivity, with some beneficial members having the capacity to facilitate plant

nutrient uptake, abiotic stress tolerance, and disease suppression [6].

Agricultural management practices induce substantial physical, biological and

chemical changes in soil that alter microbial communities [11-13]. Understanding how

microbes respond to subsequent revegetation practices is needed to evaluate ecosystem-level

outcomes of such efforts. Previous studies have linked revegetation of agricultural land to

clear shifts in microbial community composition [14-16]. However, whether these

compositional changes translate to changes in soil microbiome functional and ecological

processes remain less clear. In particular, understanding the impacts of revegetation on

microbial metabolism and life history strategies represent a critical knowledge gap. Without

this understanding, we cannot meaningfully predict or verify the long-term success of

revegetation, particularly its capacity to sequester carbon and build resilient ecosystems.

To address this, we used deep shotgun metagenomics and genome-resolved analyses

to examine how revegetation, over decadal time scales, influences the structure, functional

potential, and life history strategies of soil microbial communities in former grazing

landscapes. We find that revegetation reshapes both the composition and metabolic potential

of soil microbiomes in ways likely to support improved soil, plant, and ecosystem health.

Methods

Sample collection and processing

Sampling was conducted on six livestock grazing farms (five sheep pastures and one cattle

pasture) within the Southern Tablelands and South West Slopes of New South Wales,

Australia. All farms are working farms that have had plots restricted from grazing and

revegetated at time scales ranging from 1 to 31 years prior to our field collections. Thus, each

farm contained a paired site: an active livestock pasture and an adjacent revegetated plot.

Revegetation was carried out by the not-for-profit organisation, Greening Australia. Sites

were revegetated either via seeding or direct plantings, covering 8 to 20 species per site

(Supplementary Table S1). Specific coordinates and establishments rates for each farm can be

found in Supplementary Table S1.

Soil sampling was conducted within a 24-hour period (29-30th May 2024). Within

each farm, we collected triplicate soil cores from both revegetated and grazing areas. Cores

were taken 10 m apart along 20 m transects. Within each farm, the paired (grazing versus

revegetated) transects were located within 50 m of one another. For each sample, the top ~1

cm of material was removed, and a soil core (5 cm wide × 10 cm deep) was collected. Cores

were passed through a 2 mm sieve and homogenised in the field using ethanol-sterilised

equipment. From each homogenised core, we retained four subsamples in 5 mL sterile tubes

for independent DNA extractions, plus additional soil filled in a sterile 50 mL tube for

physicochemical analyses. All samples were immediately placed on ice in a cooler box.

Material for DNA extraction was stored at −20°C until processing, while samples for soil

physicochemical measurements were kept at 4°C and processed within one week at the

Environmental Analysis Laboratory (Southern Cross University, Lismore, Australia).

Measured physicochemical properties included pH, electrical conductivity, available

phosphorus, calcium, magnesium, potassium, effective cation exchange capacity, total carbon

and total nitrogen.

Total metagenomic DNA was extracted from 0.25 g of each quadruplicate soil

subsample using a previously established bead-beating protocol [17], and subsequently

pooled by sample at equimolar concentrations. DNA libraries were prepared for shotgun

metagenomic sequencing and run on an Illumina NovaSeq X Plus 10B (300-cycle) platform

at the Ramaciotti Centre for Genomics (UNSW, Sydney, Australia), producing, on average,

137.6 million × 150-bp paired-end reads (equivalent to 20.6 gigabases) per sample. All

sequence data have been made available via the European Nucleotide Archive (ENA) under

project accession PRJEB97193 (ENA sample accessions: ERS26780466 – ERS26780501).

Sequence read processing and metagenomic assembly

Paired-end metagenomic reads underwent adapter-clipping and quality-trimming using fastp

v0.24.1 [18, 19], employing a 4bp sliding window, from 5'- to 3'-end, with a mean quality

threshold of 20 [parameters: --cut_right --cut_right_window_size 4 --cut_right_mean_quality

20 --thread 16]. Reads were then cleaned to remove Illumina PhiX control spike-in and

contaminating human sequences. Contaminant reads were identified by mapping the trimmed

reads to the human telomere-to-telomere (T2T-CHM13+Y v2.0) [20, 21] and *Escherichia*

phage phiX174 (NC_001422.1) genomes, using Strobealign v0.16.0 [22]. Paired-end reads

where both read pairs mapped to a reference genome were identified with SAMtools [23],

and removed using the *filterbyname* program from the BBTools software suite

(https://sourceforge.net/projects/bbmap/). Contaminant reads accounted for 0.03% – 0.1%

(median=0.06%) of reads. The final set of filtered reads were then assembled for each sample

using MetaHipMer2 [24], with default settings. Each single-sample assembly was run across

40 HPC nodes on the Gadi supercomputer (National Computational Infrastructure Australia),

with 1,920 CPU cores and 7.6 Tb of available memory.

Taxonomic profiling of the metagenomes

Taxonomic profiling of the soil metagenomes was performed with SingleM v0.19.0 [25], run

using the *pipe* subcommand. The SingleM *summarise* subcommand was used to obtain

relative abundances for each taxonomic rank. To calculate beta-diversity between samples,

we used read counts of the universal marker, S3.5 ribosomal protein S2, clustered at 96.67%

using the SingleM *summarise* subcommand. The OTU counts were normalised by total-sum

scaling (TSS) using the *transform_sample_counts* function from the phyloseq v1.3.2 R

package [26]. Non-metric multidimensional scaling (NMDS) was used to visualise sample-

level differences in community composition. First, all pairwise Bray-Curtis dissimilarities

[27] were calculated from the normalised OTU table using the *distance* function from

phyloseq, and ordinated with the *metaMDS* function from the vegan v2.5.7 R package [28] to

generate a two-dimensional NMDS plot [parameters: distance = "bray", k = 2, maxit = 999,

trymax = 1000, autotransform = FALSE, wascores = FALSE]. Taxonomic alpha-diversity

was measured using the Shannon diversity index [29], calculated from the TSS-normalised

OTU table with the *diversity* function from vegan.

We used an eDNA approach to profile the plant community in each site via targeted

assembly of full-length ITS sequences from the metagenomes. First, we merged paired-end

reads with FLASH [30] [parameters: --max-overlap 150], which, together with the unmerged

forward and reverse reads, were profiled against the EUKARYOME v1.9.4 ITS database [31]

using Kraken 2 [32]. All reads classified by Kraken 2 as putative eukaryote ITS sequences

were extracted using the *extract_kraken_reads.py* utility script, and co-assembled with the

overlap-based assembler PenguiN Release 5-cf8933 [33] [parameters: penguin nuclassemble

--min-contig-len 350 --num-iterations 50]. Full-length ITS sequences (ITS1-5.8S-ITS2) were

identified and extracted using ITSx v1.1.3 [34] [parameters: -t all --only_full T]. ITS

sequences were clustered into OTUs at 97% identity and 80% coverage using Vclust v1.2.3

[35], applying the Leiden clustering algorithm [36]. Plant OTUs were identified using the

SINTAX function [37] of VSEARCH v2.29.1 [38] against the EUKARYOME ITS database.

The presence or absence of plant ITS sequences was determined in each sample using

CoverM, with a minimum read alignment identity of 97% and ITS sequence coverage of

50%. The ITS presence/absence table was collapsed at the transect-level, and putative plant

community compositional differences were visualised using NMDS ordination as described

above. We acknowledge limitations in using eDNA to infer plant community composition,

including potential contamination from wind-blown pollen or leaves, and an inability to

quantify relative abundances due to biased DNA representation from sources like fine roots.

Future work will include vegetation surveys at soil sampling locations.

Functional profiling of the metagenomes

Functional profiling of the soil metagenomes was performed using EcoFoldDB v2.0 [39], a

database and pipeline for protein structure-guided annotations of ecologically relevant

microbial functions. First, proteins were predicted using Pyrodigal v3.6.3 [40], a Python

library binding to Prodigal [41] [parameters: -p meta -j 96]. This resulted in a total of

56,154,166 predicted proteins. We removed protein redundancy from this set, including

fragments, by clustering the 56 million proteins with MMSeqs2 Linclust [42] at >95%

identity and >99% coverage of the smaller sequence [parameters: mmseqs linclust --min-seq-

id 0.95 --cov-mode 1 --cluster-mode 2 -c 0.99]. The resulting dataset was further reduced by

clustering with MMSeqs2 cluster [43] with 90% identity and 80% bidirectional coverage

thresholds [parameters: mmseqs cluster --min-seq-id 0.9 --cov-mode 0 -c 0.8], generating a

non-redundant set of 29,725,552 proteins. These protein sequences were functionally

annotated using the *EcoFoldDB-annotate* v2.0 pipeline with default settings, which leverages

the ProstT5 protein language model [44] and Foldseek [45] to allow rapid protein structure-

based functional annotations against EcoFoldDB directly from sequence data.

Genome binning and analysis of metagenome-assembled genomes

Assembled contigs were binned into clusters of metagenome-assembled genomes (MAGs)

using GenomeFace [46]. Prior to binning, contig coverage calculation was performed using

CoverM v0.7.0 [47], which mapped the filtered reads from each sample to the complete set of

pooled contigs using Strobealign [parameters: contig -p strobealign --strobealign-use-index --

min-read-percent-identity 97 -m metabat]. The contig coverage information was provided to

GenomeFace for binning in 'single sample assembly' mode, and with a minimum contig

length of 1,000 bp [parameters: -s -m 1000]. Genome quality of the resulting MAGs was

assessed using CheckM2 [48]. Only high-purity MAGs, with CheckM2-estimated

contamination <5% and completeness > 50%, were retained for downstream analysis.

Filtered MAGs were then dereplicated at 99% average nucleotide identity (ANI) using Galah

[49], as part of the CoverM package. Galah was run using skani [50] for ANI calculations,

and the Checkm2 output for genome quality information [parameters: coverm cluster --ani 99

--checkm2-quality-report --precluster-method skani --cluster-method skani].

214   MAGs were taxonomically classified using GTDB-Tk v2.4.1 [51, 52] against GTDB

release 226 [53-56] [parameters: classify_wf --skip_ani_screen]. Domain-specific

phylogenies were inferred from the concatenated protein alignments produced by GTDB-Tk,

based on the *bac120* [57] and *ar53* [58] marker sets. Approximate maximum-likelihood

phylogenetic trees were constructed from the alignments using FastTree2 v2.1.11 [59],

applying the WAG amino acid substitution model [60], and with branch lengths rescaled to

optimise the Gamma20 likelihood [parameters: -gamma -wag]. The trees were rooted with

minimal ancestor deviation (MAD) [61] using MADroot [62]

(https://github.com/davidjamesbryant/MADroot), and visualised using the *ggtree* [63] and

ggtreeExtra [64] R packages.

Metabolic pathways encoded by the MAGs were identified using MetaPathPredict

[65], which takes KofamScan annotations [66], and applies machine learning models to

predict the presence or absence of KEGG modules [67] within incomplete genomes. For

trophic-lifestyle inference, each MAG was evaluated for carbon-fixation pathways (based on

MetaPathPredict and EcoFoldDB annotations), and for substrate profiles of carbohydrate-

active enzyme (CAZyme) gene clusters (CGCs) using the run_dbcan v5.1.2 [68]

easy_substrate pipeline. MAGs encoding carbon-fixation pathways but lacking detectable

CGCs were classified as autotrophs; MAGs with CGCs but no carbon-fixation pathways were

classified as heterotrophs; and MAGs with both features were classified as mixotrophs.

Genome-based maximum growth rates were predicted for each MAG using Phydon

[69], which incorporates codon usage biases and phylogenetic signal for robust predictions of

minimum doubling times. Genome annotation data used by Phydon was generated using

Prokka v1.14.6 [70], while the evolutionary relationship data were based on the phylogenetic

placements of each MAG in the GTDB bacterial and archaeal trees, produced by GTDB-Tk.

Statistical analyses

All statistical analyses were conducted in R [71]. To control the false discovery rate (FDR)

arising from multiple hypothesis testing, all *p*-values were adjusted using the Benjamini-

Hochberg method [72] where applicable.

We used the MaAsLin 2 R package [73] to identify OTUs differentially abundant

between grazing and revegetated soils. Linear models were fit with land-use (grazing vs

revegetated) as a fixed effect, and farm as a random effect. OTU counts were normalised by

total-sum scaling (TSS) and log-transformed. OTUs present in fewer than 10% of samples

were excluded.

MaAslin 2 was also used to identify MAGs enriched in either grazing or revegetated

soils. MAG per-base abundances were estimated with CoverM in genome mode (parameters:

genome -p strobealign -m mean), and then scaled for sequencing depth by the average

number of read pairs across all samples. These normalised MAG abundances were analysed

with MaAsLin2 using the same model structure as the OTU analysis, but with MaAsLin2

count normalisation disabled.

We used linear modelling to identify microbial functions enriched in revegetated

versus grazing soils. Gene relative abundances were taken from the CoverM-calculated

coverage of their contigs (i.e., per-base contig abundances scaled for sequencing depth by the

average read-pair count), and summed for each EcoFoldDB functional category. We fit the

model: *Abundance ~ Farm + Land-use + Land-use × Time*, for each function where,

Abundance = summed normalised abundance of that function; *Farm* = blocking covariate

accounting for paired sampling design (i.e., grazing and revegetated transects from the same

farm); *Land-use* = grazing versus revegetated; and *Land-use × Time* = interaction term of

years since revegetation (mean-centred).

Alpha-diversity (Shannon index) differences were tested with Wilcoxon rank-sum

tests [74]. Differences in beta-diversity (Bray-Curtis dissimilarities) were tested with

PERMANOVAs using the *adonis2* function from the vegan R package. We applied

environmental vector fitting to examine any associations between measured soil abiotic

variables and microbial community composition using the *envfit* function from vegan.

Fisher's exact tests [75] were used to identify KEGG modules and trophic-lifestyle categories

over-represented in revegetated- versus grazing-enriched MAGs. Linear models were used to

test for correlations between MaAsLin2 MAG enrichment coefficients and Phydon-predicted

maximum growth rates.

Results and Discussion

Shifts in the compositional structure of soil microbiomes

We used deep shotgun metagenomic sequencing to assess soil microbiome responses to

revegetation following the cessation of livestock grazing. Average soil microbial biodiversity,

assessed using the Shannon diversity index (alpha-diversity), did not differ significantly

between grazed and revegetated sites, although, the variation in diversity was greater in

revegetated soils (Fig. 1b; Supplementary Fig. S1). In contrast, beta-diversity was found to be

significantly different (Fig. 1b, c), indicating that revegetation following the cessation of

grazing significantly altered microbial community composition relative to grazed soils.

Notably, this restructuring occurred independently of major abiotic soil properties, as no

measured physicochemical parameter was significantly correlated with community

composition via environmental fitting (FDR-adjusted p-values > 0.05), although a trend

towards lower pH was observed in revegetated soils (Supplementary Fig. S2). We found that

soils from sheep-grazed pastures shifted as early as 3 years post-revegetation, forming a

distinct cluster across the 3-to-31-year range (PERMANOVA: sheep-grazed versus

revegetated samples, $p = 0.001$). In contrast, the only cattle-grazed pasture (F3) exhibited no

compositional shift, even after 11 years since revegetation (Fig. 1c; PERMANOVA: cattle-

grazed versus revegetated samples, $p = 0.1$). This differential response might indicate that the

restoration of grazing soil microbiomes is livestock-specific; an effect possibly stemming

from cattle causing a higher degree of initial soil degradation than sheep, though further

investigation is required to confirm this. In contrast to these observed soil microbiome shifts,

we found a consistent convergence in putative plant community composition (inferred from

eDNA) regardless of previous land use, with all revegetated sites >3 years old clustering

together and differing significantly from grazed pastures (PERMANOVA, $p = 0.008$;

Supplementary Fig. S3).

301
Fig. 1. The impacts of revegetation following grazing cessation on soil microbial
community diversity. a, Soil was sampled from six active livestock farms in NSW,
Australia. Revegetated areas had been restored 1-31 years prior. At each farm, triplicate soil
cores were sampled from both grazing and adjacent revegetated areas. **b**, Soil microbial
alpha-diversity (Shannon index) comparisons between grazing (blue) and revegetated soils
(orange). See Supplementary Fig. S1 for farm-wise comparisons. **c**, Non-metric
multidimensional scaling (NMDS) visualising sample-level differences in soil community
composition (i.e., beta-diversity). Community dissimilarities are derived from pairwise Bray-
Curtis distances. Point shapes indicate land-use type: grazing (triangles) versus revegetated
(stars). Points are coloured by years since revegetation. Text labels indicate the farm from
which the samples derived, as shown in panel **a**. Note, F3 represents the cattle-grazed farm,
while all others are sheep-grazed.
314

Revegetated soil microbiomes were characterised by a consistent increase in the relative

abundance of Actinomycetota (Fig. 2a), the dominant phylum across all samples. This shift

was driven by the enrichment of specific Actinomycetota taxa at the OTU level (Fig. 2b),

particularly from the family Streptosporangiaceae, as well as members of the

Solirubrobacteraceae, Pseudonocardiaceae and Nocardioidaceae families. Other significantly

enriched OTUs belonged to the genus *Bradyrhizobium* (class Alphaproteobacteria) (Fig. 2b),

the predominant rhizobial genus that forms symbioses with many Australian native legumes

[76].

323
Fig. 2. Taxonomic shifts in soil microbiomes following grazing cessation and
revegetation. a, Stacked bar chart shows phylum-level composition of soil microbial
communities from each sample. The corresponding farm and years since revegetation are
shown above. **b**, Microbial OTUs significantly enriched in grazing versus revegetated soils.
The most significant OTUs, with an enrichment coefficient > 0.5 and *q*-value (FDR-adjusted
p-value) < 0.01, are labelled with their lowest taxonomic classification, and coloured by
phylum.

Shifts in the functional potential of soil microbiomes

Cessation of grazing and revegetation drove significant shifts in the functional potential of

the soil microbial communities (Fig. 3). Using protein structure-based functional profiling

with EcoFoldDB [39], we identified genes linked to key ecological traits. Soils revegetated

after cessation of grazing were enriched relative to grazed soils in functions associated with a

suite of diverse biogeochemical processes, including carbon fixation, nutrient mineralisation

and retention (N, P and S) and plant-microbe interactions (Fig. 3).

Fig. 3. Functional shifts in soil microbiomes following revegetation. Community functions
that are significantly enriched (FDR-adjusted *p*-values < 0.05) in grazing versus revegetated
soils. Bars are coloured according to their broader functional category.

All of the core carbon fixation pathways present in prokaryotes [77] were

significantly enriched in the soil microbial communities of revegetated sites, indicating a

heightened potential for soil carbon sequestration. As revegetation is strongly associated with

increased soil organic carbon storage [78, 79], our findings suggest that the enrichment of

diverse carbon-fixing taxa may play a key role in this re-carbonisation of soils. In addition,

inorganic phosphorus solubilisation and organic phosphorus mineralisation were enriched in

revegetated soils, suggesting a community with a greater potential for increasing plant-

available phosphorus. We also observed nitrogen cycling pathways enriched in the

revegetated communities, involving nitrogen acquisition and recycling. These included

assimilatory nitrate reduction (incorporating nitrogen into biomass), nitrogen fixation

(introducing new nitrogen), and the recycling of organic nitrogen compounds (e.g., nucleic

acids, amides, urea) into plant-available ammonia. Revegetated soils were also enriched in

pathways representing complete sulphur cycling, including the mineralisation of organic

sulphur, reduction of sulphate, and the re-oxidation back to sulphate. This sulphur cycling is

likely to sustain plant-available sulphur and prevent the accumulation of toxic hydrogen

sulphide.

In contrast, soils which continue to be grazed were significantly enriched in nitric

oxide reduction, relative to those revegetated, a central step in denitrification that promotes

the direct loss of nitrogen from soil systems through the emission of nitrous oxide (N_2O).

This also highlights a potential climate change related consequence associated with grazed

soils since N_2O is a potent greenhouse gas with a warming potential approximately 300

times that of CO_2 [80]. Additionally, grazing soils were enriched in sulphur reduction to

hydrogen sulphide (without the corresponding enrichment of the complementary oxidative

pathways required for a complete sulphur cycle). This has the potential to increase hydrogen

sulphide accumulation, which can inhibit other microbial processes as well as plant growth.

Soil microbiomes within revegetated areas were also significantly enriched in genes

involved in beneficial plant-microbe interactions (Fig. 3). Key enriched functions included

ACC deaminase activity, which modulates plant ethylene levels to alleviate abiotic stress

[81]; the synthesis of phytohormones such as indole-3-acetic acid (auxin), jasmonic acid, and

gibberellins to directly stimulate plant growth and defence [82-84]; and the production of

polyamines (e.g., putrescine and spermidine) and gamma-aminobutyric acid (GABA), which

contribute to stress resilience and root development [85, 86]. Furthermore, the enrichment of

root nodulation genes highlights the shift in microbial community to support mutualistic

symbioses between rhizobia and native legumes. Collectively, these findings indicate that the

cessation of grazing together with revegetation drive functional shifts in soil microbiomes

towards processes important for soil, plant, and ecosystem health.

Temporal dynamics of shifts in microbial function

A key finding of our study is the distinct chronology of microbial functional shifts following

the cessation of grazing and revegetation. The time-series analysis revealed that soil

microbiome responses to grazing cessation and revegetation occurs in two broad temporal

phases. We found that many of the functions observed to be enriched in the revegetated soils

had no correlation with time, indicating that fundamental shifts in biogeochemical cycling

occur rapidly (within ~3 years) and then stabilise over longer time scales. This suggests a

rapid initial restructuring of core ecosystem processes, such as carbon fixation and nutrient

mineralisation, establishing a new, stable functional baseline soon after land use changes.

However, a second set of functional shifts occurred more gradually following grazing

cessation and revegetation (Fig. 4). Among the functions that were significantly correlated

with age in these sites, those facilitating plant-microbe interactions represented the most

abundant category. This pattern suggests that as the plant community grows and matures over

time, selection for more specialised plant-microbe interactions increases along with it.

Consequently, the development of a native plant-adapted microbiome appears to be an

ongoing process that continues to develop over longer (decadal) periods following

revegetation. This decadal-scale maturation may reflect a gradual fine-tuning of the soil

microbiome, aligning with the establishing plant community, however, specific tests of this

are needed. Beyond plant-associated functions, several other metabolic processes showed

progressive enrichment with time since grazing cessation and revegetation, including urease

activity, phospholipid degradation, thiosulphate oxidation, and methane and carbon monoxide

oxidation, suggesting community adaptation to more diverse metabolic niches as revegetated

sites develop.

408
409 **Fig. 4. Functional shifts correlated with time since revegetation.** Individual panels show
410 community functions (summed normalised gene abundances) that are significantly correlated
411 (FDR-adjusted p-values < 0.05) with years since revegetation and cessation of grazing.
412 Panels are coloured according to their broader functional category.
413

Together, these findings suggest that the restoration of soil microbiome function after grazing

cessation and revegetation follows a clear temporal sequence: an initial, rapid transformation

of core biogeochemical cycles, followed by a prolonged, gradual specialisation towards more

complex plant-microbe interactions and niche adaptation.

Shifts in microbial life history strategies

We next performed genome-resolved analyses to assess the impact of grazing cessation along

with revegetation on genomic traits linked to microbial growth and life history strategies.

From the assemblies, we reconstructed 442 bacterial and 12 archaeal MAGs of high purity

(>50% completeness and >95% purity), primarily representing the phyla Actinomycetota,

Acidobacteriota, and Pseudomonadota. We observed a phylogenetic signal in the patterns of

genome enrichment, indicating that responses to grazing cessation and revegetation is often

lineage-specific (Fig. 5). This suggests that evolutionary history influences microbial

enrichment, likely stemming from certain clades sharing conserved functional or

physiological traits that drive their success in either grazing or revegetated soils.

Fig. 5. Enrichment of metagenome-assembled genomes (MAGs) in grazing and revegetated soils. Phylogeny of **a**, 442 bacterial and **b**, 12 archaeal high-purity (>95% purity, >50% completeness) MAGs. Tree tip points are coloured by GTDB phylum. Scale bars indicate amino acid substitutions per site. Colour scale bars indicate enrichment in grazing (blue) versus revegetated (red) soils, and asterisks indicate statistical significance (FDR-adjusted p-values < 0.05).

Grazing cessation coupled with revegetation significantly altered patterns of microbial life history traits. Growth rate predictions, based on codon usage bias and phylogenetic signal (which serve as robust predictors for microbial growth potential [69, 87]), revealed a significant enrichment of faster-growing taxa in revegetated soils (Fig. 6a). This aligns with previous observations of copiotrophic (faster-growing) taxa enriched in soils following revegetation [88, 89], potentially driven by increased inputs of plant-derived organic matter. Our results, consistent with these studies, suggest that the cessation of grazing and revegetation select for microbial communities with greater investment in growth capacity. We also found that revegetated soils favoured a higher abundance of carbon-fixing taxa (Fig. 6b), key drivers of soil carbon sequestration, as well as genomes enriched in diverse biosynthetic pathways (Fig. 6c). These included the *de novo* synthesis of amino acids (e.g., tryptophan, lysine, valine, leucine), nucleotides (deoxyribonucleotides, pyrimidines), and essential cofactors (riboflavin, NAD, Coenzyme A, betaine). Collectively, this profile indicates communities geared toward enhanced synthesis of complex macromolecules, growth, and high biomass yield. This is supported by previous studies involving empirical measurements of soils, which show strong associations between revegetation and increased soil microbial biomass and soil organic carbon [78, 79, 88-92]. Our findings suggest that these increases could be driven by shifts in life history strategies that prioritise carbon fixation, biosynthesis and growth yields. This could have significant implications for enhancing the carbon sink potential of soils, warranting more targeted investigations into the effects of land use and revegetation on microbiome-mediated carbon dynamics.

460
Fig. 6. Genome-resolved enrichment of microbial life history traits. a, Relationship
between enrichment coefficients for the metagenome-assembled genomes (MAGs) (positive
= enriched in revegetated soil; negative = enriched in grazing soils) and their predicted
maximum growth rates. The black line shows the linear regression with 95% confidence
interval (shaded). The *p*-value for the association is displayed on the panel. **b**, Proportion of
MAGs enriched in revegetated (red) or grazing (blue) samples classified into trophic
categories. Bars show the percentage of MAGs in each category. The *p*-value for the
difference between groups is indicated on the panel. **c**, Enrichment of metabolic pathways
(KEGG modules) significantly enriched (FDR-adjusted *p*-value < 0.05) in revegetated-
associated (red) versus grazing-associated (blue) MAGs. KEGG module annotations are
displayed on the plot.
472

Conversely, grazing soils were enriched in slower-growing taxa (Fig. 6a) with a

greater investment in catabolic and stress resistance mechanisms (Fig. 6b,c), suggesting a

trade-off in life history strategies, favouring resource scavenging and stress tolerance, over

biomass production. Grazing-enriched metabolic pathways included energy-harvesting (e.g.,

pyruvate oxidation, histidine degradation) and processing one-carbon compounds (C1-unit

interconversion). Metabolic pathways often associated with stress response strategies were

also enriched. These included beta-lactam resistance (a maker for antimicrobial and general

cell envelope stress response [93]), fatty acid biosynthesis (important for maintaining cellular

integrity and cell membrane repair during stress), and cysteine biosynthesis (a key pathway

for oxidative stress resistance through its role in antioxidant production [94, 95]). Such

investments into cellular maintenance and stress tolerance mechanisms, generally reduce

microbial carbon use efficiency, diverting carbon away from growth and limiting the

formation of microbial biomass and soil organic carbon [10].

Conclusion

Here, we demonstrate that the cessation of grazing, together with revegetation, drives

significant shifts in the structure, function and prevailing life history strategies of soil

microbial communities. Microbiome responses were consistent across all sites where

revegetation and the cessation of grazing occurred at least three years prior, with one notable

exception. We note that this outlier site, which showed no significant response after 11 years,

was the only previously cattle-grazed pasture, with all others being previously sheep-grazed.

We hypothesise that degradation associated with cattle grazing may be more impactful on soil

microbiomes, imposing a longer-lasting negative legacy that impedes microbial recovery,

though further research is needed to confirm this. Despite this exception, soil microbiomes

from revegetation sites were consistently associated with clear shifts relative to the sheep-

grazed soil microbiomes.

This grazing-to-revegetation transition was also marked by fundamental functional

shifts in the soil microbiome. In grazed soils, the microbial community was enriched in

pathways involved in nitrogen loss, emissions of the potent greenhouse gas, nitrous oxide,

and sulphur reduction, which can lead to the accumulation of toxic hydrogen sulphide. These

enriched functions were accompanied in grazed soils by prevailing life history strategies

focused on catabolic resource scavenging and stress-response. In contrast, revegetated soils

were enriched in communities geared toward higher growth yield strategies and metabolic

processes that underpin biosynthesis, carbon sequestration, nutrient retention, and plant

mutualisms.

Crucially, while shifts in key biogeochemical functions linked to soil health occurred

rapidly and stabilised following the cessation of grazing and revegetation, plant growth-

promoting traits increased more gradually, likely in response to the progressive maturity of

the plant community over time. This indicates a multi-stage microbial successional process:

an early restructuring of core soil health processes, followed by a progressive development of

plant-microbe interactions as the plant community matures. Together, our findings suggest

that soil microbial communities not only respond to but may also facilitate long-term

ecosystem recovery following revegetation.

Data availability

Raw shotgun metagenomic sequence data generated in this study have been made available

via the European Nucleotide Archive (ENA), under project accession PRJEB97193 (ENA

sample accessions ERS26780466 – ERS26780501).

Competing interests

The authors declare that they have no competing interests.

Funding

527 This work was supported by the Macquarie University Research Fellowship (TMG) and the

528 ARC Centre of Excellence in Synthetic Biology (SGT). JLR acknowledges funding from the

529 Australian Research Council (grant no. DP2401012143).

Acknowledgements

532 The authors would like to thank all landholders for permitting access to collection sites.

References

1.   Asner GP, Elmore AJ, Olander LP, Martin RE, Harris AT: **Grazing systems,**
ecosystem responses, and global change. *Annual Review of Environment*
and Resources 2004, **29:**261-299.
2.   Eldridge DJ, Poore AGB, Ruiz-Colmenero M, Letnic M, Soliveres S:
Ecosystem structure, function, and composition in rangelands are
negatively affected by livestock grazing. *Ecological Applications* 2016,
26:1273-1283.
3.   Wolff S, Schrammeijer EA, Schulp CJE, Verburg PH: **Meeting global land**
restoration and protection targets: What would the world look like in
2050? *Global Environmental Change* 2018, **52:**259-272.
4.   Coban O, De Deyn GB, van der Ploeg M: **Soil microbiota as game-**
changers in restoration of degraded lands. *Science* 2022, **375:**abe0725.
5.   Sokol NW, Slessarev E, Marschmann GL, Nicolas A, Blazewicz SJ, Brodie
EL, Firestone MK, Foley MM, Hestrin R, Hungate BA, et al: **Life and death in**
the soil microbiome: how ecological processes influence
biogeochemistry. *Nature Reviews Microbiology* 2022, **20:**415-430.
6.   Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK: **Plant–microbiome**
interactions: from community assembly to plant health. *Nature reviews*
microbiology 2020, **18:**607-621.
7.   Wardle DA, Bardgett RD, Klironomos JN, Setala H, Van Der Putten WH, Wall
DH: **Ecological linkages between aboveground and belowground biota.**
Science 2004, **304:**1629-1633.
8.   Jansson JK, Hofmockel KS: **Soil microbiomes and climate change.** *Nature*
Reviews Microbiology 2020, **18:**35-46.
9.   Cavicchioli R, Ripple WJ, Timmis KN, Azam F, Bakken LR, Baylis M,
Behrenfeld MJ, Boetius A, Boyd PW, Classen AT, et al: **Scientists' warning**
to humanity: microorganisms and climate change. *Nature Reviews*
Microbiology 2019, **17:**569-586.

10.   Malik AA, Martiny JBH, Brodie EL, Martiny AC, Treseder KK, Allison SD:
564   **Defining trait-based microbial strategies with consequences for soil**
565   **carbon cycling under climate change.** *The ISME Journal* 2019, **14:**1-9.
566   11.   Ghaly TM, Shah BS, Coleman NV, Elbourne LD, Le Roux JJ, Gillings MR,
567   Paulsen IT, Tetu SG: **Agriculture alters protein evolution of nitrogen**
568   **cycling genes in soil bacteria at a global scale.** *bioRxiv* 2025**:**2025.2005.
569   2007.652758.
570   12.   Hartmann M, Six J: **Soil structure and microbiome functions in**
571   **agroecosystems.** *Nature Reviews Earth & Environment* 2023, **4:**4-18.
572   13.   Peng Z, Qian X, Liu Y, Li X, Gao H, An Y, Qi J, Jiang L, Zhang Y, Chen S:
573   **Land conversion to agriculture induces taxonomic homogenization of**
574   **soil microbial communities globally.** *Nature Communications* 2024,
575   **15:**3624.
576   14.   Xiao H, Li Z, Dong Y, Chang X, Deng L, Huang J, Nie X, Liu C, Liu L, Wang D,
577   et al: **Changes in microbial communities and respiration following the**
578   **revegetation of eroded soil.** *Agriculture, Ecosystems & Environment* 2017,
579   **246:**30-37.
580   15.   Guo Y, Chen X, Wu Y, Zhang L, Cheng J, Wei G, Lin Y: **Natural revegetation**
581   **of a semiarid habitat alters taxonomic and functional diversity of soil**
582   **microbial communities.** *Science of The Total Environment* 2018, **635:**598-
583   606.
584   16.   Gellie NJC, Mills JG, Breed MF, Lowe AJ: **Revegetation rewilds the soil**
585   **bacterial microbiome of an old field.** *Molecular Ecology* 2017, **26:**2895-
586   2904.
587   17.   Yeates C, Gillings M, Davison A, Altavilla N, Veal D: **Methods for microbial**
588   **DNA extraction from soil for PCR amplification.** *Biological procedures*
589   *online* 1998, **1:**40-47.
590   18.   Chen S: **Ultrafast one-pass FASTQ data preprocessing, quality control,**
591   **and deduplication using fastp.** *iMeta* 2023, **2:**e107.
592   19.   Chen S, Zhou Y, Chen Y, Gu J: **fastp: an ultra-fast all-in-one FASTQ**
593   **preprocessor.** *Bioinformatics* 2018, **34:**i884-i890.
594   20.   Nurk S, Koren S, Rhie A, Rautiainen M, Bzikadze AV, Mikheenko A, Vollger
595   MR, Altemose N, Uralsky L, Gershman A, et al: **The complete sequence of a**
596   **human genome.** *Science* 2022, **376:**44-53.
597   21.   Rhie A, Nurk S, Cechova M, Hoyt SJ, Taylor DJ, Altemose N, Hook PW, Koren
598   S, Rautiainen M, Alexandrov IA, et al: **The complete sequence of a human**
599   **Y chromosome.** *Nature* 2023, **621:**344-354.
600   22.   Sahlin K: **Strobealign: flexible seed size enables ultra-fast and accurate**
601   **read alignment.** *Genome Biology* 2022, **23:**260.
602   23.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G,
603   Abecasis G, Durbin R, Subgroup GPDP: **The Sequence Alignment/Map**
604   **format and SAMtools.** *Bioinformatics* 2009, **25:**2078-2079.
605   24.   Hofmeyr S, Egan R, Georganas E, Copeland AC, Riley R, Clum A, Eloe-
606   Fadrosh E, Roux S, Goltsman E, Buluç A, et al: **Terabase-scale**
607   **metagenome coassembly with MetaHipMer.** *Scientific Reports* 2020,
608   **10:**10689.
609   25.   Woodcroft BJ, Aroney STN, Zhao R, Cunningham M, Mitchell JAM,
610   Nurdiansyah R, Blackall L, Tyson GW: **Comprehensive taxonomic**
611   **identification of microbial species in metagenomic data using SingleM**

and Sandpiper. *Nature Biotechnology* 2025:https://doi.org/10.1038/s41587-
41025-02738-41581.
26.   McMurdie PJ, Holmes S: **phyloseq: An R package for reproducible**
interactive analysis and graphics of microbiome census data. *PLoS One*
2013, **8:**e61217.
27.   Bray JR, Curtis JT: **An ordination of the upland forest communities of**
southern Wisconsin. *Ecological Monographs* 1957, **27:**326-349.
28.   Jari Oksanen, F. Guillaume Blanchet, Michael Friendly, Roeland Kindt, Pierre
Legendre, Dan McGlinn, Peter R. Minchin, R. B. O'Hara, Gavin L. Simpson,
Peter Solymos, et al: **vegan: Community Ecology Package.** *R package*
version 25-7 https://CRANR-projectorg/package=vegan 2020.
29.   Shannon CE: **A mathematical theory of communication.** *The Bell System*
Technical Journal 1948, **27:**379-423.
30.   Magoč T, Salzberg SL: **FLASH: fast length adjustment of short reads to**
improve genome assemblies. *Bioinformatics* 2011, **27:**2957-2963.
31.   Tedersoo L, Hosseyni Moghaddam MS, Mikryukov V, Hakimzadeh A, Bahram
M, Nilsson RH, Yatsiuk I, Geisen S, Schwelm A, Piwosz K, et al:
EUKARYOME: the rRNA gene reference database for identification of all
eukaryotes. *Database (Oxford)* 2024, **2024:**baae043.
32.   Wood DE, Lu J, Langmead B: **Improved metagenomic analysis with**
Kraken 2. *Genome Biology* 2019, **20:**257.
33.   Jochheim A, Jochheim FA, Kolodyazhnaya A, Morice É, Steinegger M, Söding
J: **Strain-resolved de-novo metagenomic assembly of viral genomes and**
microbial 16S rRNAs. *Microbiome* 2024, **12:**187.
34.   Bengtsson-Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, De
Wit P, Sánchez-García M, Ebersberger I, de Sousa F: **Improved software**
detection and extraction of ITS1 and ITS 2 from ribosomal ITS
sequences of fungi and other eukaryotes for analysis of environmental
sequencing data. *Methods in Ecology and Evolution* 2013, **4:**914-919.
35.   Zielezinski A, Gudyś A, Barylski J, Siminski K, Rozwalak P, Dutilh BE,
Deorowicz S: **Ultrafast and accurate sequence alignment and clustering**
of viral genomes. *Nature Methods* 2025, **22:**1191-1194.
36.   Traag VA, Waltman L, Van Eck NJ: **From Louvain to Leiden: guaranteeing**
well-connected communities. *Scientific reports* 2019, **9:**1-12.
37.   Edgar RC: **SINTAX: a simple non-Bayesian taxonomy classifier for 16S**
and ITS sequences. *bioRxiv* 2016:074161.
38.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F: **VSEARCH: a versatile**
open source tool for metagenomics. *PeerJ* 2016, **4:**e2584.
39.   Ghaly TM, Rajabal V, Russell D, Colombi E, Tetu SG: **EcoFoldDB: Protein**
structure-guided functional profiling of ecologically relevant microbial
traits at the metagenome scale. *Environmental Microbiology* 2025,
27:e70178.
40.   Larralde M: **Pyrodigal: Python bindings and interface to Prodigal, an**
efficient method for gene prediction in prokaryotes. *Journal of Open*
Source Software 2022, **7:**4296.
41.   Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ: **Prodigal:**
prokaryotic gene recognition and translation initiation site identification.
BMC Bioinformatics 2010, **11:**119.
42.   Steinegger M, Söding J: **Clustering huge protein sequence sets in linear**
time. *Nature Communications* 2018, **9:**2542.

43. Steinegger M, Söding J: **MMseqs2 enables sensitive protein sequence
searching for the analysis of massive data sets.** *Nature Biotechnology*
2017, **35:**1026-1028.
44. Heinzinger M, Weissenow K, Sanchez Joaquin G, Henkel A, Mirdita M,
Steinegger M, Rost B: **Bilingual language model for protein sequence and
structure.** *NAR Genomics and Bioinformatics* 2024, **6:**lqae150.
45. van Kempen M, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchrist CLM,
Söding J, Steinegger M: **Fast and accurate protein structure search with
Foldseek.** *Nature Biotechnology* 2024, **42:**243-246.
46. Lettich R, Egan R, Riley R, Wang Z, Tritt A, Oliker L, Yelick K, Buluç A:
**GenomeFace: a deep learning-based metagenome binner trained on
43,000 microbial genomes.** *bioRxiv* 2024**:**2024.2002.2007.579326.
47. Aroney STN, Newell RJP, Nissen JN, Camargo AP, Tyson GW, Woodcroft BJ:
CoverM: read alignment statistics for metagenomics. *Bioinformatics* 2025,
41:btaf147.
48. Chklovski A, Parks DH, Woodcroft BJ, Tyson GW: **CheckM2: a rapid,
scalable and accurate tool for assessing microbial genome quality using
machine learning.** *Nature Methods* 2023, **20:**1203-1212.
49. Aroney STN, Camargo AP, Tyson GW, Woodcroft B: **Galah: More scalable
dereplication for metagenome assembled genomes. Zenodo
10.5281/zenodo.10526086.** 2024.
50. Shaw J, Yu YW: **Fast and robust metagenomic sequence comparison
through sparse chaining with skani.** *Nature Methods* 2023, **20:**1661-1665.
51. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH: **GTDB-Tk: a toolkit to
classify genomes with the Genome Taxonomy Database.** Oxford
University Press; 2020.
52. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH: **GTDB-Tk v2: memory
friendly classification with the genome taxonomy database.**
Bioinformatics 2022, **38:**5315-5316.
53. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A,
Hugenholtz P: **A standardized bacterial taxonomy based on genome
phylogeny substantially revises the tree of life.** *Nature biotechnology*
2018, **36:**996-1004.
54. Parks DH, Chuvochina M, Chaumeil P-A, Rinke C, Mussig AJ, Hugenholtz P:
A complete domain-to-species taxonomy for Bacteria and Archaea.
Nature biotechnology 2020, **38:**1079-1086.
55. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P:
**GTDB: an ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-
based taxonomy.** *Nucleic acids research* 2022, **50:**D785-D794.
56. Rinke C, Chuvochina M, Mussig AJ, Chaumeil P-A, Davín AA, Waite DW,
Whitman WB, Parks DH, Hugenholtz P: **A standardized archaeal taxonomy
for the Genome Taxonomy Database.** *Nature microbiology* 2021, **6:**946-
959.
57. Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Woodcroft BJ, Evans PN,
Hugenholtz P, Tyson GW: **Recovery of nearly 8,000 metagenome-
assembled genomes substantially expands the tree of life.** *Nature
microbiology* 2017, **2:**1533-1542.
58. Dombrowski N, Williams TA, Sun J, Woodcroft BJ, Lee J-H, Minh BQ, Rinke
C, Spang A: **Undinarchaeota illuminate DPANN phylogeny and the impact**

of gene transfer on archaeal evolution. *Nature Communications* 2020,
11:3939.

59.   Price MN, Dehal PS, Arkin AP: **FastTree 2–approximately maximum-**
likelihood trees for large alignments. *PLoS One* 2010, **5:**e9490.

60.   Whelan S, Goldman N: **A general empirical model of protein evolution**
derived from multiple protein families using a maximum-likelihood
approach. *Molecular Biology and Evolution* 2001, **18:**691-699.

61.   Tria FDK, Landan G, Dagan T: **Phylogenetic rooting using minimal**
ancestor deviation. *Nature Ecology & Evolution* 2017, **1:**0193.

62.   Bryant D, Charleston M: **MAD roots for large trees.** *arXiv* 2018:1811.03174.

63.   Yu G, Smith DK, Zhu H, Guan Y, Lam TTY: **ggtree: an R package for**
visualization and annotation of phylogenetic trees with their covariates
and other associated data. *Methods in Ecology and Evolution* 2017, **8:**28-
36.

64.   Xu S, Dai Z, Guo P, Fu X, Liu S, Zhou L, Tang W, Feng T, Chen M, Zhan L:
ggtreeExtra: compact visualization of richly annotated phylogenetic
data. *Molecular Biology and Evolution* 2021, **38:**4039-4042.

65.   Geller-McGrath D, Konwar KM, Edgcomb VP, Pachiadaki M, Roddy JW,
Wheeler TJ, McDermott JE: **Predicting metabolic modules in incomplete**
bacterial genomes with MetaPathPredict. *eLife* 2024, **13:**e85749.

66.   Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata
H: **KofamKOALA: KEGG Ortholog assignment based on profile HMM and**
adaptive score threshold. *Bioinformatics* 2020, **36:**2251-2252.

67.   Kanehisa M, Goto S: **KEGG: kyoto encyclopedia of genes and genomes.**
Nucleic acids research 2000, **28:**27-30.

68.   Zheng J, Ge Q, Yan Y, Zhang X, Huang L, Yin Y: **dbCAN3: automated**
carbohydrate-active enzyme and substrate annotation. *Nucleic Acids*
Research 2023, **51:**W115-W121.

69.   Xu L, Zakem E, Weissman JL: **Improved maximum growth rate prediction**
from microbial genomes by integrating phylogenetic information. *Nature*
Communications 2025, **16:**4256.

70.   Seemann T: **Prokka: rapid prokaryotic genome annotation.** *Bioinformatics*
2014, **30:**2068-2069.

71.   **R Core Team (2020). R: A language and environment for statistical**
computing. R Foundation for Statistical Computing, Vienna, Austria.
URL https://www.R-project.org/.

72.   Benjamini Y, Hochberg Y: **Controlling the false discovery rate: A practical**
and powerful approach to multiple testing. *Journal of the Royal Statistical*
Society: Series B (Methodological) 1995, **57:**289-300.

73.   Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, Tickle TL,
Weingart G, Ren B, Schwager EH, et al: **Multivariable association**
discovery in population-scale meta-omics studies. *PLoS Computational*
Biology 2021, **17:**e1009442.

74.   Wilcoxon F: **Individual comparisons by ranking methods.** *Biometrics*
Bulletin 1945, **1:**80-83.

75.   Fisher RA: **Statistical methods for research workers.** In *Breakthroughs in*
statistics: Methodology and distribution. Springer; 1970: 66-70

76.   Lafay B, Burdon JJ: **Molecular diversity of rhizobia occurring on native**
shrubby legumes in southeastern Australia. *Applied and Environmental*
Microbiology 1998, **64:**3989-3997.

77.   Garritano AN, Song W, Thomas T: **Carbon fixation pathways across the**
763   **bacterial and archaeal tree of life.** *PNAS Nexus* 2022, **1:**pgac226.
764   78.   Bell SM, Barriocanal C, Terrer C, Rosell-Melé A: **Management opportunities**
765   **for soil carbon sequestration following agricultural land abandonment.**
766   *Environmental Science & Policy* 2020, **108:**104-111.
767   79.   Zhang Q, Hu Y, Shao M, Jia X, Wei X: **Revegetation re-carbonizes soil:**
768   **Patterns, mechanisms, and challenges.** *Fundamental Research* 2024, **In**
769   **Press, Corrected Proof**.
770   80.   Arias PA, Bellouin N, Coppola E, Jones RG, Krinner G, Marotzke J, Naik V,
771   Palmer MD, Plattner GK, Rogelj J, et al: **Technical Summary.** In *Climate*
772   *Change 2021: The Physical Science Basis Contribution of Working Group I to*
773   *the Sixth Assessment Report of the Intergovernmental Panel on Climate*
774   *Change.* Edited by Masson-Delmotte V, Zhai P, Pirani A, Connors SL, Péan C,
775   Berger S, Caud N, Chen Y, Goldfarb L, Gomis MI*, et al*. Cambridge, United
776   Kingdom and New York, NY, USA: Cambridge University Press; 2021: 33–144
777   81.   Glick BR, Todorovic B, Czarny J, Cheng Z, Duan J, McConkey B: **Promotion**
778   **of plant growth by bacterial ACC deaminase.** *Critical Reviews in Plant*
779   *Sciences* 2007, **26:**227-242.
780   82.   Tang J, Li Y, Zhang L, Mu J, Jiang Y, Fu H, Zhang Y, Cui H, Yu X, Ye Z:
781   **Biosynthetic pathways and functions of indole-3-acetic acid in**
782   **microorganisms.** *Microorganisms* 2023, **11:**2077.
783   83.   Li C, Schilmiller AL, Liu G, Lee GI, Jayanty S, Sageman C, Vrebalov J,
784   Giovannoni JJ, Yagi K, Kobayashi Y: **Role of β-oxidation in jasmonate**
785   **biosynthesis and systemic wound signaling in tomato.** *The Plant Cell*
786   2005, **17:**971-986.
787   84.   Bottini R, Cassán F, Piccoli P: **Gibberellin production by bacteria and its**
788   **involvement in plant growth promotion and yield increase.** *Applied*
789   *Microbiology and Biotechnology* 2004, **65:**497-503.
790   85.   Dunn MF, Becerra-Rivera VA: **The biosynthesis and functions of**
791   **polyamines in the interaction of plant growth-promoting rhizobacteria**
792   **with plants.** *Plants* 2023, **12:**2671.
793   86.   Kinnersley AM, Turano FJ: **Gamma aminobutyric acid (GABA) and plant**
794   **responses to stress.** *Critical Reviews in Plant Sciences* 2000, **19:**479-509.
795   87.   Weissman JL, Hou S, Fuhrman JA: **Estimating maximal microbial growth**
796   **rates from cultures, metagenomes, and single cells via codon usage**
797   **patterns.** *Proceedings of the National Academy of Sciences* 2021,
798   **118:**e2016810118.
799   88.   Gao X, Dong S, Xu Y, Li Y, Li S, Wu S, Shen H, Liu S, Fry EL: **Revegetation**
800   **significantly increased the bacterial-fungal interactions in different**
801   **successional stages of alpine grasslands on the Qinghai-Tibetan**
802   **Plateau.** *CATENA* 2021, **205:**105385.
803   89.   Yang Y, Dou Y, Wang B, Xue Z, Wang Y, An S, Chang SX: **Deciphering**
804   **factors driving soil microbial life-history strategies in restored**
805   **grasslands.** *iMeta* 2023, **2:**e66.
806   90.   Xiao L, Liu G-b, Zhang J-y, Xue S: **Long-term effects of vegetational**
807   **restoration on soil microbial communities on the Loess Plateau of**
808   **China.** *Restoration Ecology* 2016, **24:**794-804.
809   91.   Hu L, Li Q, Yan J, Liu C, Zhong J: **Vegetation restoration facilitates**
810   **belowground microbial network complexity and recalcitrant soil organic**

carbon storage in southwest China karst region. *Science of The Total*
Environment 2022, **820:**153137.
92. Sha G, Chen Y, Wei T, Guo X, Yu H, Jiang S, Xin P, Ren K: **Responses of**
soil microbial communities to vegetation restoration on the Loess
Plateau of China: A meta-analysis. *Applied Soil Ecology* 2023, **189:**104910.
93. Bucher T, Keren-Paz A, Hausser J, Olender T, Cytryn E, Kolodkin-Gal I: **An**
active β-lactamase is a part of an orchestrated cell wall stress resistance
network of Bacillus subtilis and related rhizosphere species.
Environmental Microbiology 2019, **21:**1068-1085.
94. Turnbull AL, Surette MG: **Cysteine biosynthesis, oxidative stress and**
antibiotic resistance in *Salmonella typhimurium***.** *Research in Microbiology*
2010, **161:**643-650.
95. Tikhomirova A, Rahman MM, Kidd SP, Ferrero RL, Roujeinikova A: **Cysteine**
and resistance to oxidative stress: implications for virulence and
antibiotic resistance. *Trends in Microbiology* 2024, **32:**93-104.
826