Soil microbial traits shift on contrasting timescales following revegetation

of former grazing lands

Timothy M. Ghaly[1,2]*, Vanessa J. McPherson[1], Vaheesan Rajabal[1,2], Mary E. Ghaly[1], Nicki

Taws[3], Rachael V. Gallagher[4], Johannes J. Le Roux[1], Sasha G. Tetu[1,2]*

[1]School of Natural Sciences, Macquarie University, New South Wales, Australia

[2]ARC Centre of Excellence in Synthetic Biology, New South Wales, Australia

[3]Greening Australia, Australian Capital Territory, Australia

[4]Hawkesbury Institute for the Environment, Western Sydney University, New South Wales,

Australia

*Corresponding Authors:

Timothy M. Ghaly: timothy.ghaly@mq.edu.au

Sasha G. Tetu: sasha.tetu@mq.edu.au

Keywords: biogeochemical cycling; carbon fixation; ecological restoration; land use change;

microbial life history strategies; plant-microbe interactions; soil carbon storage

Abstract

Revegetation is a key strategy for restoring degraded lands globally. While this process can reshape belowground microbial communities, the extent to which such changes restore soil ecosystem functions, and whether different microbial traits recover synchronously or on distinct timescales remains less clear. Understanding this is essential for evaluating restoration success, as different microbial traits underpin distinct ecosystem services, from carbon storage to plant growth promotion, and the sequence in which these are restored can inform restoration targets and provide meaningful indicators of long-term ecological recovery. To address this, we applied deep metagenomic sequencing to characterise soil microbial responses following revegetation (spanning 1-31 years prior to sampling) on grazing agricultural lands. We find that revegetation following grazing cessation drove significant and sequential shifts in dominant microbial functions and life-history strategies. Functional changes occurred in distinct phases: an early, rapid restructuring of core soil health processes, detectable as early as three years, including enrichment of nutrient retention and carbon fixation pathways, followed by a more gradual development of plant growth-promoting traits as the plant community matures. Genome-resolved analyses of nearly 500 metagenome-assembled genomes revealed a fundamental shift in dominant microbial life history strategies: from a resource-scavenging and stress tolerance profile in grazing soils to strategies that prioritise biosynthesis and growth yields in revegetated soils. These lifestyle shifts have important implications for enhancing the microbial biomass and carbon sink potential of soils. Together, these findings show that microbial functional shifts following revegetation are temporally structured – informing expectations for when key restoration targets may be achieved, and providing a practical framework for monitoring ecosystem recovery.

Introduction

Human land use is a major driver of environmental degradation and ecosystem service loss.

Agricultural grazing land, which covers more than 25% of Earth's land surface, is subject to

significant degradation through soil compaction and altered nutrient fluxes, both of which

reduce plant diversity, and collectively alter ecosystem processes and functioning [1, 2].

Revegetation, the process of restoring plant species in disturbed areas, is a widely adopted

strategy for the restoration of degraded lands, with agricultural land representing a key global

target for this style of restoration [3]. Although the aboveground outcomes of revegetation are

often visible, the impacts on belowground ecosystems, and the bi-directional feedbacks

between them, are less clear.

Soil microbiomes are fundamental mediators of ecosystem processes from local to

global scales, governing biogeochemical cycles, soil structure and moisture storage, and

climate regulation [4-6]. Soil microbiomes drive the cycling of nutrients that underpin

ecosystem health, with microbial metabolism tightly linked to aboveground biodiversity [7].

Soil microbial communities also exert strong control over carbon storage and soil-atmosphere

gas fluxes through metabolic processes such as carbon and nitrogen fixation, decomposition,

and the production or consumption of greenhouse gases [8, 9]. Microbial life history

strategies, defined by trade-offs between growth yield, resource acquisition, and stress

tolerance, also impact soil carbon dynamics [10]. Specifically, a high *growth yield* strategy

promotes soil carbon storage by increasing biomass and subsequently trapping much of this

carbon as stable 'necromass'. In contrast, a focus on *resource acquisition* can drive

decomposition and carbon loss by breaking down complex organic matter; while greater

investment in *stress response* strategies can increase CO_2 respiration, as energy is redirected

from biomass production to maintenance and repair. In addition to impacts on

biogeochemical cycling, soil microbiomes can also have significant influence on plant health

and productivity, with some beneficial members having the capacity to facilitate plant

nutrient uptake, abiotic stress tolerance, and disease suppression [6].

Agricultural management practices induce substantial physical, biological and

chemical changes in soil that alter microbial communities [11-13]. Understanding how

microbes respond to subsequent revegetation practices is needed to evaluate ecosystem-level

outcomes of such efforts. Prior work has firmly established that the composition of soil

microbial communities shift in response to revegetation [14-16]. Studies using 16S rRNA

amplicon profiling have shown that revegetation alters the relative abundances of dominant

bacterial and fungal phyla, and that these shifts broadly parallel aboveground plant

community recovery [14-16]. What remains poorly resolved, however, is whether

compositional change translates into functional recovery, and whether functional shifts occur

synchronously across different microbial traits or on distinct timescales. This distinction

matters practically, as it provides critical information regarding when soil functional capacity

is restored and whether different management targets (e.g., carbon sequestration, nutrient

cycling, plant growth promotion) are achieved simultaneously or sequentially.

A further unresolved question concerns the ecological strategies of microorganisms in

revegetated soils. Revegetation, through the progressive restoration of plant-derived carbon

inputs, root exudate diversity, and reduced disturbance, is predicted to shift the selective

landscape toward yield-oriented life history strategies over time. However, whether these

predicted shifts in strategies do indeed occur post-revegetation, and at what time scales

remains unclear. Understanding the dominant life history strategies of revegetated soil

microbiomes has direct consequences for carbon storage (e.g., via necromass production by

high-yield strategists) and nutrient cycling rates, and therefore provides an ecologically

meaningful indicator of restoration trajectory. Addressing these questions requires moving

beyond profiling compositional change to testing mechanistic predictions about how

microbial life-history strategies and ecosystem functions respond to land-use change over

decadal timescales.

Here, we used deep shotgun metagenomics and genome-resolved analyses to examine

how revegetation across decadal timescales reshapes the functional potential and life history

strategies of soil microbial communities on former livestock grazing lands. We tested the

central hypothesis that revegetation following grazing cessation alters soil microbiome

functional capacity relative to actively grazed soils. We further predicted that distinct classes

of functional traits and life history strategies would shift on different timescales, reflecting

the sequential nature of ecosystem recovery. Our findings reveal a temporally structured

microbiome response to revegetation, with implications for how belowground recovery can

be monitored and interpreted in the context of broader ecosystem restoration.

Methods

Sample collection and processing

Sampling was conducted on six livestock grazing farms (five sheep pastures and one cattle

pasture) within the Southern Tablelands and South West Slopes of New South Wales,

Australia. All farms are working farms that have had plots restricted from grazing and

revegetated at time scales ranging from 1 to 31 years prior to our field collections. Thus, each

farm contained a paired site: an active livestock pasture and an adjacent revegetated plot.

Revegetation was carried out by the not-for-profit organisation, Greening Australia. Sites

were revegetated either via seeding or direct plantings, covering 8 to 20 species per site

(Supplementary Table S1). Specific coordinates and establishments rates for each farm can be

found in Supplementary Table S1.

Soil sampling was conducted within a 24-hour period (29-30[th] May 2024). Within

each farm, we collected triplicate soil cores from both revegetated and grazing areas. Cores

were taken 10 m apart along 20 m transects. Within each farm, the paired (grazing versus

revegetated) transects were located within 50 m of one another. For each sample, the top ~1

cm of material was removed, and a soil core (5 cm wide $\times$ 10 cm deep) was collected. Cores

were passed through a 2 mm sieve and homogenised in the field using ethanol-sterilised

equipment. From each homogenised core, we retained four subsamples in 5 mL sterile tubes

for independent DNA extractions, plus additional soil filled in a sterile 50 mL tube for

physicochemical analyses. All samples were immediately placed on ice in a cooler box.

Material for DNA extraction was stored at $-20°C$ until processing, while samples for soil

physicochemical measurements were kept at 4°C and processed within one week at the

Environmental Analysis Laboratory (Southern Cross University, Lismore, Australia).

Measured physicochemical properties included pH, electrical conductivity, available

phosphorus, calcium, magnesium, potassium, effective cation exchange capacity, total carbon

and total nitrogen.

Total metagenomic DNA was extracted from 0.25 g of each quadruplicate soil

subsample using a previously established bead-beating protocol [17], and subsequently

pooled by sample at equimolar concentrations. DNA libraries were prepared for shotgun

metagenomic sequencing and run on an Illumina NovaSeq X Plus 10B (300-cycle) platform

at the Ramaciotti Centre for Genomics (UNSW, Sydney, Australia), producing, on average,

137.6 million $\times$ 150-bp paired-end reads (equivalent to 20.6 gigabases) per sample. All

sequence data have been made available via the European Nucleotide Archive (ENA) under

project accession PRJEB97193 (ENA sample accessions: ERS26780466 – ERS26780501).

Sequence read processing and metagenomic assembly

Paired-end metagenomic reads underwent adapter-clipping and quality-trimming using fastp

v0.24.1 [18, 19], employing a 4bp sliding window, from 5'- to 3'-end, with a mean quality

threshold of 20 [parameters: --cut_right --cut_right_window_size 4 --cut_right_mean_quality

20 --thread 16]. Reads were then cleaned to remove Illumina PhiX control spike-in and

contaminating human sequences. Contaminant reads were identified by mapping the trimmed

reads to the human telomere-to-telomere (T2T-CHM13+Y v2.0) [20, 21] and *Escherichia*

phage phiX174 (NC_001422.1) genomes, using Strobealign v0.16.0 [22]. Paired-end reads

where both read pairs mapped to a reference genome were identified with SAMtools [23],

and removed using the *filterbyname* program from the BBTools software suite

(https://sourceforge.net/projects/bbmap/). Contaminant reads accounted for 0.03% – 0.1%

(median=0.06%) of reads. The final set of filtered reads were then assembled for each sample

using MetaHipMer2 [24], with default settings. Each single-sample assembly was run across

40 HPC nodes on the Gadi supercomputer (National Computational Infrastructure Australia),

with 1,920 CPU cores and 7.6 Tb of available memory.

Taxonomic profiling of the metagenomes

Taxonomic profiling of the soil metagenomes was performed with SingleM v0.19.0 [25], run

using the *pipe* subcommand. The SingleM *summarise* subcommand was used to obtain

relative abundances for each taxonomic rank. To calculate beta-diversity between samples,

we used read counts of the universal marker, S3.5 ribosomal protein S2, clustered at 96.67%

using the SingleM *summarise* subcommand. The OTU counts were normalised by total-sum

scaling (TSS) using the *transform_sample_counts* function from the phyloseq v1.3.2 R

package [26]. Non-metric multidimensional scaling (NMDS) was used to visualise sample-

level differences in community composition. First, all pairwise Bray-Curtis dissimilarities

[27] were calculated from the normalised OTU table using the *distance* function from

phyloseq, and ordinated with the *metaMDS* function from the vegan v2.5.7 R package [28] to

generate a two-dimensional NMDS plot [parameters: distance = "bray", k = 2, maxit = 999,

trymax = 1000, autotransform = FALSE, wascores = FALSE]. Taxonomic alpha-diversity

was measured using the Shannon diversity index [29], calculated from the TSS-normalised

OTU table with the *diversity* function from vegan.

182   We used an eDNA approach to profile the plant community in each site via targeted

assembly of full-length ITS sequences from the metagenomes. First, we merged paired-end

reads with FLASH [30] [parameters: --max-overlap 150], which, together with the unmerged

forward and reverse reads, were profiled against the EUKARYOME v1.9.4 ITS database [31]

using Kraken 2 [32]. All reads classified by Kraken 2 as putative eukaryote ITS sequences

were extracted using the *extract_kraken_reads.py* utility script, and co-assembled with the

overlap-based assembler PenguiN Release 5-cf8933 [33] [parameters: penguin nuclassemble

--min-contig-len 350 --num-iterations 50]. Full-length ITS sequences (ITS1-5.8S-ITS2) were

identified and extracted using ITSx v1.1.3 [34] [parameters: -t all --only_full T]. ITS

sequences were clustered into OTUs at 97% identity and 80% coverage using Vclust v1.2.3

[35], applying the Leiden clustering algorithm [36]. Plant OTUs were identified using the

SINTAX function [37] of VSEARCH v2.29.1 [38] against the EUKARYOME ITS database.

The presence or absence of plant ITS sequences was determined in each sample using

CoverM, with a minimum read alignment identity of 97% and ITS sequence coverage of

50%. The ITS presence/absence table was collapsed at the transect-level, and putative plant

community compositional differences were visualised using NMDS ordination as described

above. We acknowledge limitations in using eDNA to infer plant community composition,

including potential contamination from wind-blown pollen or leaves, and an inability to

quantify relative abundances due to biased DNA representation from sources like fine roots.

Future work will include vegetation surveys at soil sampling locations.

Functional profiling of the metagenomes

Functional profiling of the soil metagenomes was performed using EcoFoldDB v2.0 [39], a

database and pipeline for protein structure-guided annotations of ecologically relevant

microbial functions. First, proteins were predicted using Pyrodigal v3.6.3 [40], a Python

library binding to Prodigal [41] [parameters: -p meta -j 96]. This resulted in a total of

56,154,166 predicted proteins. We removed protein redundancy from this set, including

fragments, by clustering the 56 million proteins with MMSeqs2 *Linclust* [42] at >95%

identity and >99% coverage of the smaller sequence [parameters: mmseqs linclust --min-seq-

id 0.95 --cov-mode 1 --cluster-mode 2 -c 0.99]. The resulting dataset was further reduced by

clustering with MMSeqs2 *cluster* [43] with 90% identity and 80% bidirectional coverage

thresholds [parameters: mmseqs cluster --min-seq-id 0.9 --cov-mode 0 -c 0.8], generating a

non-redundant set of 29,725,552 proteins. These protein sequences were functionally

annotated using the *EcoFoldDB-annotate* v2.0 pipeline with default settings, which leverages

the ProstT5 protein language model [44] and Foldseek [45] to allow rapid protein structure-

based functional annotations against EcoFoldDB directly from sequence data.

Genome binning and analysis of metagenome-assembled genomes

Assembled contigs were binned into clusters of metagenome-assembled genomes (MAGs)

using GenomeFace [46]. Prior to binning, contig coverage calculation was performed using

CoverM v0.7.0 [47], which mapped the filtered reads from each sample to the complete set of

pooled contigs using Strobealign [parameters: contig -p strobealign --strobealign-use-index --

min-read-percent-identity 97 -m metabat]. The contig coverage information was provided to

GenomeFace for binning in 'single sample assembly' mode, and with a minimum contig

length of 1,000 bp [parameters: -s -m 1000]. Genome quality of the resulting MAGs was

assessed using CheckM2 [48]. Only high-purity MAGs, with CheckM2-estimated

contamination <5% and completeness > 50%, were retained for downstream analysis.

Filtered MAGs were then dereplicated at 99% average nucleotide identity (ANI) using Galah

[49], as part of the CoverM package. Galah was run using skani [50] for ANI calculations,

and the Checkm2 output for genome quality information [parameters: coverm cluster --ani 99

--checkm2-quality-report --precluster-method skani --cluster-method skani].

MAGs were taxonomically classified using GTDB-Tk v2.4.1 [51, 52] against GTDB

release 226 [53-56] [parameters: classify_wf --skip_ani_screen]. Domain-specific

phylogenies were inferred from the concatenated protein alignments produced by GTDB-Tk,

based on the *bac120* [57] and *ar53* [58] marker sets. Approximate maximum-likelihood

phylogenetic trees were constructed from the alignments using FastTree2 v2.1.11 [59],

applying the WAG amino acid substitution model [60], and with branch lengths rescaled to

optimise the Gamma20 likelihood [parameters: -gamma -wag]. The trees were rooted with

minimal ancestor deviation (MAD) [61] using MADroot [62]

(https://github.com/davidjamesbryant/MADroot), and visualised using the *ggtree* [63] and

ggtreeExtra [64] R packages.

Metabolic pathways encoded by the MAGs were identified using MetaPathPredict

[65], which takes KofamScan annotations [66], and applies machine learning models to

predict the presence or absence of KEGG modules [67] within incomplete genomes. For

trophic-lifestyle inference, each MAG was evaluated for carbon-fixation pathways (based on

MetaPathPredict and EcoFoldDB annotations), and for substrate profiles of carbohydrate-

active enzyme (CAZyme) gene clusters (CGCs) using the run_dbcan v5.1.2 [68]

easy_substrate pipeline. MAGs encoding carbon-fixation pathways but lacking detectable

CGCs were classified as autotrophs; MAGs with CGCs but no carbon-fixation pathways were

classified as heterotrophs; and MAGs with both features were classified as mixotrophs.

254   Genome-based maximum growth rates were predicted for each MAG using Phydon

[69], which incorporates codon usage biases and phylogenetic signal for robust predictions of

minimum doubling times. Genome annotation data used by Phydon was generated using

Prokka v1.14.6 [70], while the evolutionary relationship data were based on the phylogenetic

placements of each MAG in the GTDB bacterial and archaeal trees, produced by GTDB-Tk.

Statistical analyses

All statistical analyses were conducted in R [71]. To control the false discovery rate (FDR)

arising from multiple hypothesis testing, all *p*-values were adjusted using the Benjamini-

Hochberg method [72] where applicable.

265   We used the MaAsLin 2 R package [73] to identify OTUs differentially abundant

266   between grazing and revegetated soils. Linear models were fit with land-use (grazing vs

267   revegetated) as a fixed effect, and farm as a random effect. OTU counts were normalised by

268   total-sum scaling (TSS) and log-transformed. OTUs present in fewer than 10% of samples

269   were excluded.

270   MaAslin 2 was also used to identify MAGs enriched in either grazing or revegetated

soils. MAG per-base abundances were estimated with CoverM in genome mode (parameters:

genome -p strobealign -m mean), and then scaled for sequencing depth by the average

number of read pairs across all samples. These normalised MAG abundances were analysed

with MaAsLin2 using the same model structure as the OTU analysis, but with MaAsLin2

count normalisation disabled.

We used linear modelling to identify microbial functions enriched in revegetated

versus grazing soils. Gene relative abundances were taken from the CoverM-calculated

coverage of their contigs (i.e., per-base contig abundances scaled for sequencing depth by the

average read-pair count), and summed for each EcoFoldDB functional category. We fit the

model: *Abundance ~ Farm + Land-use + Land-use × Time*, for each function category

where, *Abundance* = summed normalised abundance of that function; *Farm* = blocking

covariate accounting for paired sampling design (i.e., grazing and revegetated transects from

the same farm); *Land-use* = grazing versus revegetated; and *Land-use × Time* = interaction

term of years since revegetation (mean-centred).

Alpha-diversity (Shannon index) differences were tested with Wilcoxon rank-sum

tests [74]. Differences in beta-diversity (Bray-Curtis dissimilarities) were tested with

PERMANOVAs using the *adonis2* function from the vegan R package. We applied

environmental vector fitting to examine any associations between measured soil abiotic

variables and microbial community composition using the *envfit* function from vegan.

Fisher's exact tests [75] were used to identify KEGG modules and trophic-lifestyle categories

over-represented in revegetated- versus grazing-enriched MAGs. Linear models were used to

test for correlations between MaAsLin2 MAG enrichment coefficients and Phydon-predicted

maximum growth rates.

Results and Discussion

Shifts in the compositional structure of soil microbiomes

We used deep shotgun metagenomic sequencing to assess soil microbiome responses to

revegetation following the cessation of livestock grazing. Average soil microbial biodiversity,

assessed using the Shannon diversity index (alpha-diversity), did not differ significantly

between grazed and revegetated sites, although, the variation in diversity was greater in

revegetated soils (Fig. 1b; Supplementary Fig. S1). In contrast, beta-diversity was found to be

significantly different (Fig. 1b, c), indicating that revegetation following the cessation of

grazing significantly altered microbial community composition relative to grazed soils.

Notably, this restructuring occurred independently of major abiotic soil properties, as no

measured physicochemical parameter was significantly correlated with community

composition via environmental fitting (FDR-adjusted p-values > 0.05), although a trend

towards lower pH was observed in revegetated soils (Supplementary Fig. S2). We found that

soils from sheep-grazed pastures shifted as early as 3 years post-revegetation, forming a

distinct cluster across the 3-to-31-year range (PERMANOVA: sheep-grazed versus

revegetated samples, $p = 0.001$). In contrast, the only cattle-grazed pasture (F3) exhibited no

compositional shift, even after 11 years since revegetation (Fig. 1c; PERMANOVA: cattle-

grazed versus revegetated samples, $p = 0.1$). This differential response might indicate that the

restoration of grazing soil microbiomes is livestock-specific; an effect possibly stemming

from cattle causing a higher degree of initial soil degradation than sheep, though further

investigation is required to confirm this. In contrast to these observed soil microbiome shifts,

we found a consistent convergence in putative plant community composition (inferred from

eDNA) regardless of previous land use, with all revegetated sites >3 years old clustering

together and differing significantly from grazed pastures (PERMANOVA, $p = 0.008$;

Supplementary Fig. S3).

322
Fig. 1. The impacts of revegetation following grazing cessation on soil microbial
community diversity. a, Soil was sampled from six active livestock farms in NSW,
Australia. Revegetated areas had been restored 1-31 years prior. At each farm, triplicate soil
cores were sampled from both grazing and adjacent revegetated areas. **b**, Soil microbial
alpha-diversity (Shannon index) comparisons between grazing (blue) and revegetated soils
(orange). See Supplementary Fig. S1 for farm-wise comparisons. **c**, Non-metric
multidimensional scaling (NMDS) visualising sample-level differences in soil community
composition (i.e., beta-diversity). Community dissimilarities are derived from pairwise Bray-
Curtis distances. Point shapes indicate land-use type: grazing (triangles) versus revegetated
(stars). Points are coloured by years since revegetation. Text labels indicate the farm from
which the samples derived, as shown in panel **a**. Note, F3 represents the cattle-grazed farm,
while all others are sheep-grazed.
335

Revegetated soil microbiomes were characterised by a consistent increase in the relative

abundance of Actinomycetota (Fig. 2a), the dominant phylum across all samples. This shift

was driven by the enrichment of specific Actinomycetota taxa at the OTU level (Fig. 2b),

particularly from the family Streptosporangiaceae, as well as members of the

Solirubrobacteraceae, Pseudonocardiaceae and Nocardioidaceae families. Other significantly

enriched OTUs belonged to the genus *Bradyrhizobium* (class Alphaproteobacteria) (Fig. 2b),

the predominant rhizobial genus that forms symbioses with many Australian native legumes

[76].

344
Fig. 2. Taxonomic shifts in soil microbiomes following grazing cessation and
revegetation. a, Stacked bar chart shows phylum-level composition of soil microbial
communities from each sample. The corresponding farm and years since revegetation are
shown above. **b**, Microbial OTUs significantly enriched in grazing versus revegetated soils.
The most significant OTUs, with an enrichment coefficient > 0.5 and *q*-value (FDR-adjusted
p-value) < 0.01, are labelled with their lowest taxonomic classification, and coloured by
phylum.
352

Shifts in the functional potential of soil microbiomes

Cessation of grazing and revegetation drove significant shifts in the functional potential of

the soil microbial communities (Fig. 3). Using protein structure-based functional profiling

with EcoFoldDB [39], we identified genes linked to key ecological traits. Soils revegetated

after cessation of grazing were enriched relative to grazed soils in functions associated with a

suite of diverse biogeochemical processes, including carbon fixation, nutrient mineralisation

and retention (N, P and S) and plant-microbe interactions (Fig. 3).

362
363 **Fig. 3. Functional shifts in soil microbiomes following revegetation.** Community functions
364 that are significantly enriched (FDR-adjusted p-values < 0.05) in grazing versus revegetated
365 soils. Bars are coloured according to their broader functional category.
366

All of the core carbon fixation pathways present in prokaryotes [77] were

significantly enriched in the soil microbial communities of revegetated sites, indicating a

heightened potential for soil carbon sequestration. As revegetation is strongly associated with

increased soil organic carbon storage [78, 79], our findings suggest that the enrichment of

diverse carbon-fixing taxa may play a key role in this re-carbonisation of soils. In addition,

inorganic phosphorus solubilisation and organic phosphorus mineralisation were enriched in

revegetated soils, suggesting a community with a greater potential for increasing plant-

available phosphorus. We also observed nitrogen cycling pathways enriched in the

revegetated communities, involving nitrogen acquisition and recycling. These included

assimilatory nitrate reduction (incorporating nitrogen into biomass), nitrogen fixation

(introducing new nitrogen), and the recycling of organic nitrogen compounds (e.g., nucleic

acids, amides, urea) into plant-available ammonia. Revegetated soils were also enriched in

pathways representing complete sulphur cycling, including the mineralisation of organic

sulphur, reduction of sulphate, and the re-oxidation back to sulphate. This sulphur cycling is

likely to sustain plant-available sulphur and prevent the accumulation of toxic hydrogen

sulphide.

In contrast, soils which continue to be grazed were significantly enriched in nitric

oxide reduction, relative to those revegetated, a central step in denitrification that promotes

the direct loss of nitrogen from soil systems through the emission of nitrous oxide (N_2O).

This also highlights a potential climate change related consequence associated with grazed

soils since N_2O is a potent greenhouse gas with a warming potential approximately 300

times that of CO_2 [80]. Additionally, grazing soils were enriched in sulphur reduction to

hydrogen sulphide (without the corresponding enrichment of the complementary oxidative

pathways required for a complete sulphur cycle). This has the potential to increase hydrogen

sulphide accumulation, which can inhibit other microbial processes as well as plant growth.

Soil microbiomes within revegetated areas were also significantly enriched in genes

involved in beneficial plant-microbe interactions (Fig. 3). Key enriched functions included

ACC deaminase activity, which modulates plant ethylene levels to alleviate abiotic stress

[81]; the synthesis of phytohormones such as indole-3-acetic acid (auxin), jasmonic acid, and

gibberellins to directly stimulate plant growth and defence [82-84]; and the production of

polyamines (e.g., putrescine and spermidine) and gamma-aminobutyric acid (GABA), which

contribute to stress resilience and root development [85, 86]. Furthermore, the enrichment of

root nodulation genes highlights the shift in microbial community to support mutualistic

symbioses between rhizobia and native legumes. Collectively, these findings indicate that the

cessation of grazing together with revegetation drive functional shifts in soil microbiomes

towards processes important for soil, plant, and ecosystem health.

Temporal dynamics of shifts in microbial function

A key finding of our study is the distinct chronology of microbial functional shifts following

the cessation of grazing and revegetation. The time-series analysis revealed that soil

microbiome responses to grazing cessation and revegetation occurs in two broad temporal

phases. We found that many of the functions observed to be enriched in the revegetated soils

had no correlation with time, indicating that fundamental shifts in biogeochemical cycling

occur rapidly (within ~3 years) and then stabilise over longer time scales. This suggests a

rapid initial restructuring of core ecosystem processes, such as carbon fixation and nutrient

mineralisation, establishing a new, stable functional baseline soon after land use changes.

However, a second set of functional shifts occurred more gradually following grazing

cessation and revegetation (Fig. 4). Among the functions that were significantly correlated

with age in these sites, those facilitating plant-microbe interactions represented the most

abundant category. This pattern suggests that as the plant community grows and matures over

time, selection for more specialised plant-microbe interactions increases along with it.

Consequently, the development of a native plant-adapted microbiome appears to be an

ongoing process that continues to develop over longer (decadal) periods following

revegetation. This decadal-scale maturation may reflect a gradual fine-tuning of the soil

microbiome, aligning with the establishing plant community, however, specific tests of this

are needed. Beyond plant-associated functions, several other metabolic processes showed

progressive enrichment with time since grazing cessation and revegetation, including urease

activity, phospholipid degradation, thiosulphate oxidation, and methane and carbon monoxide

oxidation, suggesting community adaptation to more diverse metabolic niches as revegetated

sites develop.

429
430   **Fig. 4. Functional shifts correlated with time since revegetation.** Individual panels show
431   community functions (summed normalised gene abundances) that are significantly correlated
432   (FDR-adjusted p-values < 0.05) with years since revegetation and cessation of grazing.
433   Panels are coloured according to their broader functional category.
434

Together, these findings suggest that the restoration of soil microbiome function after grazing

cessation and revegetation follows a clear temporal sequence: an initial, rapid transformation

of core biogeochemical cycles, followed by a prolonged, gradual specialisation towards more

complex plant-microbe interactions and niche adaptation.

Shifts in microbial life history strategies

We next performed genome-resolved analyses to assess the impact of grazing cessation along

with revegetation on genomic traits linked to microbial growth and life history strategies.

From the assemblies, we reconstructed 442 bacterial and 12 archaeal MAGs of high purity

(>50% completeness and >95% purity), primarily representing the phyla Actinomycetota,

Acidobacteriota, and Pseudomonadota. We observed a phylogenetic signal in the patterns of

genome enrichment, indicating that responses to grazing cessation and revegetation is often

lineage-specific (Fig. 5). This suggests that evolutionary history influences microbial

enrichment, likely stemming from certain clades sharing conserved functional or

physiological traits that drive their success in either grazing or revegetated soils.

452
Fig. 5. Enrichment of metagenome-assembled genomes (MAGs) in grazing and
revegetated soils. Phylogeny of **a**, 442 bacterial and **b**, 12 archaeal high-purity (>95% purity,
>50% completeness) MAGs. Tree tip points are coloured by GTDB phylum. Scale bars
indicate amino acid substitutions per site. Colour scale bars indicate enrichment in grazing
(blue) versus revegetated (red) soils, and asterisks indicate statistical significance (FDR-
adjusted *p*-values < 0.05).

Grazing cessation coupled with revegetation significantly altered patterns of microbial

life history traits. Growth rate predictions, based on codon usage bias and phylogenetic signal

(which serve as robust predictors for microbial growth potential [69, 87]), revealed a

significant enrichment of faster-growing taxa in revegetated soils (Fig. 6a). This aligns with

previous observations of copiotrophic (faster-growing) taxa enriched in soils following

revegetation [88, 89], potentially driven by increased inputs of plant-derived organic matter.

Our results, consistent with these studies, suggest that the cessation of grazing and

revegetation select for microbial communities with greater investment in growth capacity. We

also found that revegetated soils favoured a higher abundance of carbon-fixing taxa (Fig. 6b),

key drivers of soil carbon sequestration, as well as genomes enriched in diverse biosynthetic

pathways (Fig. 6c). These included the *de novo* synthesis of amino acids (e.g., tryptophan,

lysine, valine, leucine), nucleotides (deoxyribonucleotides, pyrimidines), and essential

cofactors (riboflavin, NAD, Coenzyme A, betaine). Collectively, this profile indicates

communities geared toward enhanced synthesis of complex macromolecules, growth, and

high biomass yield. This is supported by previous studies involving empirical measurements

of soils, which show strong associations between revegetation and increased soil microbial

biomass and soil organic carbon [78, 79, 88-92]. Our findings suggest that these increases

could be driven by shifts in life history strategies that prioritise carbon fixation, biosynthesis

and growth yields. This could have significant implications for enhancing the carbon sink

potential of soils, warranting more targeted investigations into the effects of land use and

revegetation on microbiome-mediated carbon dynamics.

Fig. 6. Genome-resolved enrichment of microbial life history traits. a, Relationship between enrichment coefficients for the metagenome-assembled genomes (MAGs) (positive = enriched in revegetated soil; negative = enriched in grazing soils) and their predicted maximum growth rates. The black line shows the linear regression with 95% confidence interval (shaded). The p-value for the association is displayed on the panel. **b**, Proportion of MAGs enriched in revegetated (red) or grazing (blue) samples classified into trophic categories. Bars show the percentage of MAGs in each category. The p-value for the difference between groups is indicated on the panel. **c**, Enrichment of metabolic pathways (KEGG modules) significantly enriched (FDR-adjusted p-value < 0.05) in revegetated-associated (red) versus grazing-associated (blue) MAGs. KEGG module annotations are displayed on the plot.

Conversely, grazing soils were enriched in slower-growing taxa (Fig. 6a) with a greater investment in catabolic and stress resistance mechanisms (Fig. 6b,c), suggesting a

trade-off in life history strategies, favouring resource scavenging and stress tolerance, over

biomass production. Grazing-enriched metabolic pathways included energy-harvesting (e.g.,

pyruvate oxidation, histidine degradation) and processing one-carbon compounds (C1-unit

interconversion). Metabolic pathways often associated with stress response strategies were

also enriched. These included beta-lactam resistance (a maker for antimicrobial and general

cell envelope stress response [93]), fatty acid biosynthesis (important for maintaining cellular

integrity and cell membrane repair during stress), and cysteine biosynthesis (a key pathway

for oxidative stress resistance through its role in antioxidant production [94, 95]). Such

investments into cellular maintenance and stress tolerance mechanisms, generally reduce

microbial carbon use efficiency, diverting carbon away from growth and limiting the

formation of microbial biomass and soil organic carbon [10].

Conclusion

Here, we demonstrate that the cessation of grazing, together with revegetation, drives

significant shifts in the structure, function and prevailing life history strategies of soil

microbial communities. Microbiome responses were consistent across all sites where

revegetation and the cessation of grazing occurred at least three years prior, with one notable

exception. We note that this outlier site, which showed no significant response after 11 years,

was the only previously cattle-grazed pasture, with all others being previously sheep-grazed.

We hypothesise that degradation associated with cattle grazing may be more impactful on soil

microbiomes, imposing a longer-lasting negative legacy that impedes microbial recovery,

though further research is needed to confirm this. Despite this exception, soil microbiomes

from revegetation sites were consistently associated with clear shifts relative to the sheep-

grazed soil microbiomes.

This grazing-to-revegetation transition was also marked by fundamental functional

shifts in the soil microbiome. In grazed soils, the microbial community was enriched in

pathways involved in nitrogen loss, production of the potent greenhouse gas, nitrous oxide,

and sulphur reduction, which can lead to the accumulation of toxic hydrogen sulphide. These

enriched functions were accompanied in grazed soils by prevailing life history strategies

focused on catabolic resource scavenging and stress-response. In contrast, revegetated soils

were enriched in communities geared toward higher growth yield strategies and metabolic

processes that underpin biosynthesis, carbon sequestration, nutrient retention, and plant

mutualisms.

Crucially, while shifts in key biogeochemical functions linked to soil health occurred

rapidly and stabilised following the cessation of grazing and revegetation, plant growth-

promoting traits increased more gradually, likely in response to the progressive maturity of

the plant community over time. This indicates a multi-stage microbial successional process:

an early restructuring of core soil health processes, followed by a progressive development of

plant-microbe interactions as the plant community matures. This multi-stage trajectory not

only shows that different microbial functions shift on distinct timescales, but also carries

practical implications: it informs expectations for when different management targets (such as

carbon sequestration, nutrient cycling, and plant growth promotion) may be achieved

following restoration efforts. This can provide meaningful indicators of restoration trajectory,

informing how ecosystem restoration outcomes are monitored and interpreted.

Availability of data and material

Raw shotgun metagenomic sequence data generated in this study have been made available

via the European Nucleotide Archive (ENA), under project accession PRJEB97193 (ENA

sample accessions ERS26780466 – ERS26780501).

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Macquarie University Research Fellowship (TMG) and the

ARC Centre of Excellence in Synthetic Biology (SGT). JLR acknowledges funding from the

Australian Research Council (grant no. DP2401012143).

Authors' contributions

T.M.G. conceived and designed the study, secured funding, contributed to fieldwork,

conducted data analysis, and wrote the original manuscript draft. V.M.J. contributed to

fieldwork and performed laboratory analyses. V.R. conducted laboratory work. N.T. identified

sampling sites. R.V.G., J.L.R., M.E.G., and S.G.T. contributed to project design and

management. All authors reviewed and edited the final manuscript.

Acknowledgements

TMG and MEG would like to thank Saoirse and Maebh Ghaly for their loving support. The

authors would also like to thank all landholders for permitting access to collection sites. We

acknowledge the Traditional Custodians of the lands on which soils were sampled, the

Ngunnawal and Gundungurra peoples, and pay our respects to their Elders past, present and

emerging. We recognise their continuing connection to Country and the role of Indigenous

land stewardship in maintaining healthy soils and ecosystems for millennia.

References

571   1.   Asner GP, Elmore AJ, Olander LP, Martin RE, Harris AT: **Grazing systems,**
572   **ecosystem responses, and global change.** *Annual Review of Environment*
573   *and Resources* 2004, **29:**261-299.
574   2.   Eldridge DJ, Poore AGB, Ruiz-Colmenero M, Letnic M, Soliveres S:
575   **Ecosystem structure, function, and composition in rangelands are**
576   **negatively affected by livestock grazing.** *Ecological Applications* 2016,
577   **26:**1273-1283.
578   3.   Wolff S, Schrammeijer EA, Schulp CJE, Verburg PH: **Meeting global land**
579   **restoration and protection targets: What would the world look like in**
580   **2050?** *Global Environmental Change* 2018, **52:**259-272.
581   4.   Coban O, De Deyn GB, van der Ploeg M: **Soil microbiota as game-**
582   **changers in restoration of degraded lands.** *Science* 2022, **375:**abe0725.
583   5.   Sokol NW, Slessarev E, Marschmann GL, Nicolas A, Blazewicz SJ, Brodie
584   EL, Firestone MK, Foley MM, Hestrin R, Hungate BA, et al: **Life and death in**
585   **the soil microbiome: how ecological processes influence**
586   **biogeochemistry.** *Nature Reviews Microbiology* 2022, **20:**415-430.
587   6.   Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK: **Plant–microbiome**
588   **interactions: from community assembly to plant health.** *Nature reviews*
589   *microbiology* 2020, **18:**607-621.
590   7.   Wardle DA, Bardgett RD, Klironomos JN, Setala H, Van Der Putten WH, Wall
591   DH: **Ecological linkages between aboveground and belowground biota.**
592   *Science* 2004, **304:**1629-1633.
593   8.   Jansson JK, Hofmockel KS: **Soil microbiomes and climate change.** *Nature*
594   *Reviews Microbiology* 2020, **18:**35-46.
595   9.   Cavicchioli R, Ripple WJ, Timmis KN, Azam F, Bakken LR, Baylis M,
596   Behrenfeld MJ, Boetius A, Boyd PW, Classen AT, et al: **Scientists' warning**
597   **to humanity: microorganisms and climate change.** *Nature Reviews*
598   *Microbiology* 2019, **17:**569-586.
599   10.   Malik AA, Martiny JBH, Brodie EL, Martiny AC, Treseder KK, Allison SD:
600   **Defining trait-based microbial strategies with consequences for soil**
601   **carbon cycling under climate change.** *The ISME Journal* 2019, **14:**1-9.
602   11.   Ghaly TM, Shah BS, Coleman NV, Elbourne LD, Le Roux JJ, Gillings MR,
603   Paulsen IT, Tetu SG: **Agriculture alters protein evolution of nitrogen**
604   **cycling genes in soil bacteria at a global scale.** *bioRxiv* 2025**:**2025.2005.
605   2007.652758.
606   12.   Hartmann M, Six J: **Soil structure and microbiome functions in**
607   **agroecosystems.** *Nature Reviews Earth & Environment* 2023, **4:**4-18.
608   13.   Peng Z, Qian X, Liu Y, Li X, Gao H, An Y, Qi J, Jiang L, Zhang Y, Chen S:
609   **Land conversion to agriculture induces taxonomic homogenization of**
610   **soil microbial communities globally.** *Nature Communications* 2024,
611   **15:**3624.
612   14.   Xiao H, Li Z, Dong Y, Chang X, Deng L, Huang J, Nie X, Liu C, Liu L, Wang D,
613   et al: **Changes in microbial communities and respiration following the**
614   **revegetation of eroded soil.** *Agriculture, Ecosystems & Environment* 2017,
615   **246:**30-37.
616   15.   Guo Y, Chen X, Wu Y, Zhang L, Cheng J, Wei G, Lin Y: **Natural revegetation**
617   **of a semiarid habitat alters taxonomic and functional diversity of soil**
618   **microbial communities.** *Science of The Total Environment* 2018, **635:**598-
619   606.

16. Gellie NJC, Mills JG, Breed MF, Lowe AJ: **Revegetation rewilds the soil**
bacterial microbiome of an old field. *Molecular Ecology* 2017, **26:**2895-
2904.
17. Yeates C, Gillings M, Davison A, Altavilla N, Veal D: **Methods for microbial**
DNA extraction from soil for PCR amplification. *Biological procedures*
online 1998, **1:**40-47.
18. Chen S: **Ultrafast one-pass FASTQ data preprocessing, quality control,**
and deduplication using fastp. *iMeta* 2023, **2:**e107.
19. Chen S, Zhou Y, Chen Y, Gu J: **fastp: an ultra-fast all-in-one FASTQ**
preprocessor. *Bioinformatics* 2018, **34:**i884-i890.
20. Nurk S, Koren S, Rhie A, Rautiainen M, Bzikadze AV, Mikheenko A, Vollger
MR, Altemose N, Uralsky L, Gershman A, et al: **The complete sequence of a**
human genome. *Science* 2022, **376:**44-53.
21. Rhie A, Nurk S, Cechova M, Hoyt SJ, Taylor DJ, Altemose N, Hook PW, Koren
S, Rautiainen M, Alexandrov IA, et al: **The complete sequence of a human**
Y chromosome. *Nature* 2023, **621:**344-354.
22. Sahlin K: **Strobealign: flexible seed size enables ultra-fast and accurate**
read alignment. *Genome Biology* 2022, **23:**260.
23. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G,
Abecasis G, Durbin R, Subgroup GPDP: **The Sequence Alignment/Map**
format and SAMtools. *Bioinformatics* 2009, **25:**2078-2079.
24. Hofmeyr S, Egan R, Georganas E, Copeland AC, Riley R, Clum A, Eloe-
Fadrosh E, Roux S, Goltsman E, Buluç A, et al: **Terabase-scale**
metagenome coassembly with MetaHipMer. *Scientific Reports* 2020,
10:10689.
25. Woodcroft BJ, Aroney STN, Zhao R, Cunningham M, Mitchell JAM,
Nurdiansyah R, Blackall L, Tyson GW: **Comprehensive taxonomic**
identification of microbial species in metagenomic data using SingleM
and Sandpiper. *Nature Biotechnology* 2025**:**https://doi.org/10.1038/s41587-
41025-02738-41581.
26. McMurdie PJ, Holmes S: **phyloseq: An R package for reproducible**
interactive analysis and graphics of microbiome census data. *PLoS One*
2013, **8:**e61217.
27. Bray JR, Curtis JT: **An ordination of the upland forest communities of**
southern Wisconsin. *Ecological Monographs* 1957, **27:**326-349.
28. Jari Oksanen, F. Guillaume Blanchet, Michael Friendly, Roeland Kindt, Pierre
Legendre, Dan McGlinn, Peter R. Minchin, R. B. O'Hara, Gavin L. Simpson,
Peter Solymos, et al: **vegan: Community Ecology Package.** *R package*
version 25-7 https://CRANR-projectorg/package=vegan 2020.
29. Shannon CE: **A mathematical theory of communication.** *The Bell System*
Technical Journal 1948, **27:**379-423.
30. Magoč T, Salzberg SL: **FLASH: fast length adjustment of short reads to**
improve genome assemblies. *Bioinformatics* 2011, **27:**2957-2963.
31. Tedersoo L, Hosseyni Moghaddam MS, Mikryukov V, Hakimzadeh A, Bahram
M, Nilsson RH, Yatsiuk I, Geisen S, Schwelm A, Piwosz K, et al:
EUKARYOME: the rRNA gene reference database for identification of all
eukaryotes. *Database (Oxford)* 2024, **2024:**baae043.
32. Wood DE, Lu J, Langmead B: **Improved metagenomic analysis with**
Kraken 2. *Genome Biology* 2019, **20:**257.

33.   Jochheim A, Jochheim FA, Kolodyazhnaya A, Morice É, Steinegger M, Söding
670   J: **Strain-resolved de-novo metagenomic assembly of viral genomes and**
671   **microbial 16S rRNAs.** *Microbiome* 2024, **12:**187.
672   34.   Bengtsson-Palme J, Ryberg M, Hartmann M, Branco S, Wang Z, Godhe A, De
673   Wit P, Sánchez-García M, Ebersberger I, de Sousa F: **Improved software**
674   **detection and extraction of ITS1 and ITS 2 from ribosomal ITS**
675   **sequences of fungi and other eukaryotes for analysis of environmental**
676   **sequencing data.** *Methods in Ecology and Evolution* 2013, **4:**914-919.
677   35.   Zielezinski A, Gudyś A, Barylski J, Siminski K, Rozwalak P, Dutilh BE,
678   Deorowicz S: **Ultrafast and accurate sequence alignment and clustering**
679   **of viral genomes.** *Nature Methods* 2025, **22:**1191-1194.
680   36.   Traag VA, Waltman L, Van Eck NJ: **From Louvain to Leiden: guaranteeing**
681   **well-connected communities.** *Scientific reports* 2019, **9:**1-12.
682   37.   Edgar RC: **SINTAX: a simple non-Bayesian taxonomy classifier for 16S**
683   **and ITS sequences.** *bioRxiv* 2016**:**074161.
684   38.   Rognes T, Flouri T, Nichols B, Quince C, Mahé F: **VSEARCH: a versatile**
685   **open source tool for metagenomics.** *PeerJ* 2016, **4:**e2584.
686   39.   Ghaly TM, Rajabal V, Russell D, Colombi E, Tetu SG: **EcoFoldDB: Protein**
687   **structure-guided functional profiling of ecologically relevant microbial**
688   **traits at the metagenome scale.** *Environmental Microbiology* 2025,
689   **27:**e70178.
690   40.   Larralde M: **Pyrodigal: Python bindings and interface to Prodigal, an**
691   **efficient method for gene prediction in prokaryotes.** *Journal of Open*
692   *Source Software* 2022, **7:**4296.
693   41.   Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ: **Prodigal:**
694   **prokaryotic gene recognition and translation initiation site identification.**
695   *BMC Bioinformatics* 2010, **11:**119.
696   42.   Steinegger M, Söding J: **Clustering huge protein sequence sets in linear**
697   **time.** *Nature Communications* 2018, **9:**2542.
698   43.   Steinegger M, Söding J: **MMseqs2 enables sensitive protein sequence**
699   **searching for the analysis of massive data sets.** *Nature Biotechnology*
700   2017, **35:**1026-1028.
701   44.   Heinzinger M, Weissenow K, Sanchez Joaquin G, Henkel A, Mirdita M,
702   Steinegger M, Rost B: **Bilingual language model for protein sequence and**
703   **structure.** *NAR Genomics and Bioinformatics* 2024, **6:**lqae150.
704   45.   van Kempen M, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchrist CLM,
705   Söding J, Steinegger M: **Fast and accurate protein structure search with**
706   **Foldseek.** *Nature Biotechnology* 2024, **42:**243-246.
707   46.   Lettich R, Egan R, Riley R, Wang Z, Tritt A, Oliker L, Yelick K, Buluç A:
708   **GenomeFace: a deep learning-based metagenome binner trained on**
709   **43,000 microbial genomes.** *bioRxiv* 2024**:**2024.2002.2007.579326.
710   47.   Aroney STN, Newell RJP, Nissen JN, Camargo AP, Tyson GW, Woodcroft BJ:
711   **CoverM: read alignment statistics for metagenomics.** *Bioinformatics* 2025,
712   **41:**btaf147.
713   48.   Chklovski A, Parks DH, Woodcroft BJ, Tyson GW: **CheckM2: a rapid,**
714   **scalable and accurate tool for assessing microbial genome quality using**
715   **machine learning.** *Nature Methods* 2023, **20:**1203-1212.
716   49.   Aroney STN, Camargo AP, Tyson GW, Woodcroft B: **Galah: More scalable**
717   **dereplication for metagenome assembled genomes. Zenodo**
718   **10.5281/zenodo.10526086.** 2024.

50. Shaw J, Yu YW: **Fast and robust metagenomic sequence comparison through sparse chaining with skani.** *Nature Methods* 2023, **20:**1661-1665.

51. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH: **GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database.** Oxford University Press; 2020.

52. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH: **GTDB-Tk v2: memory friendly classification with the genome taxonomy database.** *Bioinformatics* 2022, **38:**5315-5316.

53. Parks DH, Chuvochina M, Waite DW, Rinke C, Skarshewski A, Chaumeil P-A, Hugenholtz P: **A standardized bacterial taxonomy based on genome phylogeny substantially revises the tree of life.** *Nature biotechnology* 2018, **36:**996-1004.

54. Parks DH, Chuvochina M, Chaumeil P-A, Rinke C, Mussig AJ, Hugenholtz P: **A complete domain-to-species taxonomy for Bacteria and Archaea.** *Nature biotechnology* 2020, **38:**1079-1086.

55. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P: **GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy.** *Nucleic acids research* 2022, **50:**D785-D794.

56. Rinke C, Chuvochina M, Mussig AJ, Chaumeil P-A, Davín AA, Waite DW, Whitman WB, Parks DH, Hugenholtz P: **A standardized archaeal taxonomy for the Genome Taxonomy Database.** *Nature microbiology* 2021, **6:**946-959.

57. Parks DH, Rinke C, Chuvochina M, Chaumeil P-A, Woodcroft BJ, Evans PN, Hugenholtz P, Tyson GW: **Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the tree of life.** *Nature microbiology* 2017, **2:**1533-1542.

58. Dombrowski N, Williams TA, Sun J, Woodcroft BJ, Lee J-H, Minh BQ, Rinke C, Spang A: **Undinarchaeota illuminate DPANN phylogeny and the impact of gene transfer on archaeal evolution.** *Nature Communications* 2020, **11:**3939.

59. Price MN, Dehal PS, Arkin AP: **FastTree 2–approximately maximum-likelihood trees for large alignments.** *PLoS One* 2010, **5:**e9490.

60. Whelan S, Goldman N: **A general empirical model of protein evolution derived from multiple protein families using a maximum-likelihood approach.** *Molecular Biology and Evolution* 2001, **18:**691-699.

61. Tria FDK, Landan G, Dagan T: **Phylogenetic rooting using minimal ancestor deviation.** *Nature Ecology & Evolution* 2017, **1:**0193.

62. Bryant D, Charleston M: **MAD roots for large trees.** *arXiv* 2018**:**1811.03174.

63. Yu G, Smith DK, Zhu H, Guan Y, Lam TTY: **ggtree: an R package for visualization and annotation of phylogenetic trees with their covariates and other associated data.** *Methods in Ecology and Evolution* 2017, **8:**28-36.

64. Xu S, Dai Z, Guo P, Fu X, Liu S, Zhou L, Tang W, Feng T, Chen M, Zhan L: **ggtreeExtra: compact visualization of richly annotated phylogenetic data.** *Molecular Biology and Evolution* 2021, **38:**4039-4042.

65. Geller-McGrath D, Konwar KM, Edgcomb VP, Pachiadaki M, Roddy JW, Wheeler TJ, McDermott JE: **Predicting metabolic modules in incomplete bacterial genomes with MetaPathPredict.** *eLife* 2024, **13:**e85749.

66.   Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata
769   H: **KofamKOALA: KEGG Ortholog assignment based on profile HMM and**
770   **adaptive score threshold.** *Bioinformatics* 2020, **36:**2251-2252.
771   67.   Kanehisa M, Goto S: **KEGG: kyoto encyclopedia of genes and genomes.**
772   *Nucleic acids research* 2000, **28:**27-30.
773   68.   Zheng J, Ge Q, Yan Y, Zhang X, Huang L, Yin Y: **dbCAN3: automated**
774   **carbohydrate-active enzyme and substrate annotation.** *Nucleic Acids*
775   *Research* 2023, **51:**W115-W121.
776   69.   Xu L, Zakem E, Weissman JL: **Improved maximum growth rate prediction**
777   **from microbial genomes by integrating phylogenetic information.** *Nature*
778   *Communications* 2025, **16:**4256.
779   70.   Seemann T: **Prokka: rapid prokaryotic genome annotation.** *Bioinformatics*
780   2014, **30:**2068-2069.
781   71.   **R Core Team (2020). R: A language and environment for statistical**
782   **computing. R Foundation for Statistical Computing, Vienna, Austria.**
783   **URL https://www.R-project.org/.**
784   72.   Benjamini Y, Hochberg Y: **Controlling the false discovery rate: A practical**
785   **and powerful approach to multiple testing.** *Journal of the Royal Statistical*
786   *Society: Series B (Methodological)* 1995, **57:**289-300.
787   73.   Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, Tickle TL,
788   Weingart G, Ren B, Schwager EH, et al: **Multivariable association**
789   **discovery in population-scale meta-omics studies.** *PLoS Computational*
790   *Biology* 2021, **17:**e1009442.
791   74.   Wilcoxon F: **Individual comparisons by ranking methods.** *Biometrics*
792   *Bulletin* 1945, **1:**80-83.
793   75.   Fisher RA: **Statistical methods for research workers.** In *Breakthroughs in*
794   *statistics: Methodology and distribution.* Springer; 1970: 66-70
795   76.   Lafay B, Burdon JJ: **Molecular diversity of rhizobia occurring on native**
796   **shrubby legumes in southeastern Australia.** *Applied and Environmental*
797   *Microbiology* 1998, **64:**3989-3997.
798   77.   Garritano AN, Song W, Thomas T: **Carbon fixation pathways across the**
799   **bacterial and archaeal tree of life.** *PNAS Nexus* 2022, **1:**pgac226.
800   78.   Bell SM, Barriocanal C, Terrer C, Rosell-Melé A: **Management opportunities**
801   **for soil carbon sequestration following agricultural land abandonment.**
802   *Environmental Science & Policy* 2020, **108:**104-111.
803   79.   Zhang Q, Hu Y, Shao M, Jia X, Wei X: **Revegetation re-carbonizes soil:**
804   **Patterns, mechanisms, and challenges.** *Fundamental Research* 2024, **In**
805   **Press, Corrected Proof**.
806   80.   Arias PA, Bellouin N, Coppola E, Jones RG, Krinner G, Marotzke J, Naik V,
807   Palmer MD, Plattner GK, Rogelj J, et al: **Technical Summary.** In *Climate*
808   *Change 2021: The Physical Science Basis Contribution of Working Group I to*
809   *the Sixth Assessment Report of the Intergovernmental Panel on Climate*
810   *Change.* Edited by Masson-Delmotte V, Zhai P, Pirani A, Connors SL, Péan C,
811   Berger S, Caud N, Chen Y, Goldfarb L, Gomis MI*, et al*. Cambridge, United
812   Kingdom and New York, NY, USA: Cambridge University Press; 2021: 33–144
813   81.   Glick BR, Todorovic B, Czarny J, Cheng Z, Duan J, McConkey B: **Promotion**
814   **of plant growth by bacterial ACC deaminase.** *Critical Reviews in Plant*
815   *Sciences* 2007, **26:**227-242.

82.   Tang J, Li Y, Zhang L, Mu J, Jiang Y, Fu H, Zhang Y, Cui H, Yu X, Ye Z:
Biosynthetic pathways and functions of indole-3-acetic acid in
microorganisms. *Microorganisms* 2023, **11:**2077.
83.   Li C, Schilmiller AL, Liu G, Lee GI, Jayanty S, Sageman C, Vrebalov J,
Giovannoni JJ, Yagi K, Kobayashi Y: **Role of β-oxidation in jasmonate**
biosynthesis and systemic wound signaling in tomato. *The Plant Cell*
2005, **17:**971-986.
84.   Bottini R, Cassán F, Piccoli P: **Gibberellin production by bacteria and its**
involvement in plant growth promotion and yield increase. *Applied*
Microbiology and Biotechnology 2004, **65:**497-503.
85.   Dunn MF, Becerra-Rivera VA: **The biosynthesis and functions of**
polyamines in the interaction of plant growth-promoting rhizobacteria
with plants. *Plants* 2023, **12:**2671.
86.   Kinnersley AM, Turano FJ: **Gamma aminobutyric acid (GABA) and plant**
responses to stress. *Critical Reviews in Plant Sciences* 2000, **19:**479-509.
87.   Weissman JL, Hou S, Fuhrman JA: **Estimating maximal microbial growth**
rates from cultures, metagenomes, and single cells via codon usage
patterns. *Proceedings of the National Academy of Sciences* 2021,
118:e2016810118.
88.   Gao X, Dong S, Xu Y, Li Y, Li S, Wu S, Shen H, Liu S, Fry EL: **Revegetation**
significantly increased the bacterial-fungal interactions in different
successional stages of alpine grasslands on the Qinghai-Tibetan
Plateau. *CATENA* 2021, **205:**105385.
89.   Yang Y, Dou Y, Wang B, Xue Z, Wang Y, An S, Chang SX: **Deciphering**
factors driving soil microbial life-history strategies in restored
grasslands. *iMeta* 2023, **2:**e66.
90.   Xiao L, Liu G-b, Zhang J-y, Xue S: **Long-term effects of vegetational**
restoration on soil microbial communities on the Loess Plateau of
China. *Restoration Ecology* 2016, **24:**794-804.
91.   Hu L, Li Q, Yan J, Liu C, Zhong J: **Vegetation restoration facilitates**
belowground microbial network complexity and recalcitrant soil organic
carbon storage in southwest China karst region. *Science of The Total*
Environment 2022, **820:**153137.
92.   Sha G, Chen Y, Wei T, Guo X, Yu H, Jiang S, Xin P, Ren K: **Responses of**
soil microbial communities to vegetation restoration on the Loess
Plateau of China: A meta-analysis. *Applied Soil Ecology* 2023, **189:**104910.
93.   Bucher T, Keren-Paz A, Hausser J, Olender T, Cytryn E, Kolodkin-Gal I: **An**
active β-lactamase is a part of an orchestrated cell wall stress resistance
network of Bacillus subtilis and related rhizosphere species.
Environmental Microbiology 2019, **21:**1068-1085.
94.   Turnbull AL, Surette MG: **Cysteine biosynthesis, oxidative stress and**
antibiotic resistance in *Salmonella typhimurium*. *Research in Microbiology*
2010, **161:**643-650.
95.   Tikhomirova A, Rahman MM, Kidd SP, Ferrero RL, Roujeinikova A: **Cysteine**
and resistance to oxidative stress: implications for virulence and
antibiotic resistance. *Trends in Microbiology* 2024, **32:**93-104.
862