Title: Frogs uncouple neural activity from oxygen consumption after hibernation

Authors: Hafsa Yaseen[1], Joseph M. Santin[1*]

Affiliations: [1]University of Missouri-Columbia, Division of Biological Sciences, Missouri, United States of America.

***Email for correspondence**: santinj@missouri.edu

Funding statement: This work was supported by the following grants: National Science Foundation-Award # 2515635 (to JMS) and National Institutes of Health- R01NS114514 (to JMS).

Ethics approval statement: Approval for the use of animals was approved by the Animal Care and Use Committee at the University of Missouri (protocol #39264).

Data availability statement: The data that support the findings of this study are available from the corresponding author upon reasonable request.

Abstract

Aim: Aerobic metabolism supplies ~90% of the ATP for neural activity. In frogs, activity has large aerobic needs typical of an average vertebrate, but surprisingly, can shift to using only glycolysis upon emergence from hibernation. We hypothesized that hibernation triggers a global reduction in the aerobic cost of neural function.

Methods: We simultaneously measured activity of the brainstem respiratory network via motor nerves and tissue oxygen partial pressure (pO_2) *in vitro* from control and hibernated bullfrogs (4 weeks cold submergence; 4°C). To identify which functions differentially consume O_2, we sequentially blocked activity and various cellular processes requiring activity-independent ion regulation and used the resulting tissue pO_2 change (ΔpO_2) as an index of O_2 consumed. We further assessed how activity varies as a function of tissue pO_2 and how O_2 consumption varies across network activity levels.

Results: Despite similar network activity levels, we provide three lines of evidence that hibernation reduces its aerobic requirement. First, hibernators consume less oxygen for baseline activity. Second, network output remains stable from baseline to anoxia, while moderate hypoxia disrupts controls. Finally, accelerating activity does not enhance oxygen consumption as in controls, but aerobic metabolism ultimately increases during seizure-like activity.

Conclusion: Hibernating frogs reduce aerobic needs for sustaining physiological levels of neural activity, revealing how they overcome the challenge of restarting motor circuits on the background of hypoxia during emergence from hibernation. More broadly,

vertebrate neural circuits seemingly constrained by aerobic metabolism can exhibit substantial plasticity in the aerobic requirements for function.

Keywords: Oxygen consumption, neural activity, hibernation, hypoxia, metabolic plasticity, breathing network, frog

Practitioner points:

1. Hibernation in bullfrogs reduces the oxygen consumed by neural activity while maintaining normal network output, demonstrating that the aerobic requirements of brain function are not fixed in the vertebrate brain.

2. After hibernation, brainstem motor circuits maintain stable function from high levels of O_2 to anoxia and do not increase O_2 consumption when activity is elevated within the physiological range.

3. These findings reveal that vertebrate neural circuits can enter metabolic states requiring far less aerobic respiration, which may inform strategies for improving metabolic resilience in the brain during conditions of impaired oxygen delivery and other metabolic dysfunction.

1   **INTRODUCTION**

2   Brain function is energetically expensive[1,2]. At the circuit level, synaptic

transmission and the restoration of ion gradients following action potentials have high

ATP demands, the majority of which is supplied by oxidative phosphorylation[3,4]. Given

the ATP yield of aerobic respiration (30-32 moles ATP/1 mole glucose) compared to

glycolysis (2 moles of ATP/1 mole of glucose), neural activity and oxygen consumption

are tightly coupled, whereby increases in activity drive aerobic metabolism to sustain

neuronal performance[5–7]. This relationship is especially critical, as disruptions to aerobic

metabolism impair neural output and lead to neurological disease[8]. The requirement for

aerobic respiration is conserved across most vertebrate nervous systems and has

shaped prevailing views that sustained neural computation requires oxygen

consumption[9].

Despite the view that neural activity's large energy demands remain fixed, the

American bullfrog (*Aquarana catesbeiana*) provides a unique opportunity to address

how neural circuits can enter states that operate with far less energy. Under normal

circumstances frog circuits exhibit typical metabolic demands and rely heavily on

oxidative metabolism[10,11]. However, activity of brainstem motor circuits that generate

breathing improve function during anoxia from ~10 minutes to upwards of 4 hours under

otherwise identical experimental conditions after frogs emerge from hibernation[12,13].

This represents a striking display of metabolic plasticity, as this network has high

metabolic demands, with rhythm generating circuits that use glutamatergic synaptic

transmission, which is thought to incur large energy costs, and motor pools with

neurons that fire action potentials at rates >100 Hz during rhythmic bursting[14–16]. This

form of metabolic plasticity is physiologically significant because arterial pO_2 can fall as

low as 1–3 mmHg during submergence[17]. While low pO_2 does not present a problem

due to low metabolic rates and decreased need for many behaviors in the cold, the

problem is much more apparent upon emergence. Indeed, since neural activity typically

has oxygen demands, frogs emerging from hibernation must resume neural activity

under conditions when pO_2 is expected to be similarly low[18], which would be difficult if

they did not become able to function with a lower requirement for aerobic ATP

production.

Mechanistically, there are two non-mutually exclusive possibilities that could

explain how hibernation alters the brainstem to function with less O_2: increasing ATP

supply during hypoxia and/or reducing the aerobic metabolism required to sustain

activity. First, while glycolysis produces ~15-fold less ATP than oxidative

phosphorylation, hypoxia can increase glycolysis through the Pasteur effect. Thus,

hibernation may lead to cellular changes that bolster flux through glycolysis[19] to sustain

neural activity without oxygen. On the other hand, hibernation may reduce the

dependence of neural activity on aerobic metabolism making while maintaining similar

network output. While there are several ways neural circuits could in principle enhance

their energy efficiency[20–22], hibernation reduces the Ca^{2+} permeability of NMDA-

glutamate receptors and enhances receptor desensitization[23], which is likely to reduce

the energetic burden of ion regulation at synapses. Resolving these possibilities may

provide new insight into how neural circuits can shift between states with vastly different

aerobic requirements while maintaining similar function, as well as leading to new ideas

for improving metabolic resilience in the brain.

In this study, we assessed the extent to which hibernation alters activity's aerobic

requirements in the brainstem respiratory network. To estimate the aerobic metabolic

cost of activity under normal (control) and hypoxia-tolerant (after hibernation) network

states, we simultaneously measured network activity through motor nerve rootlets and

tissue oxygen partial pressure (pO_2) within the rhythm-generating circuit *in vitro*. This

approach allows us to estimate aerobic metabolism because tissue pO_2 reflects

consumption by mitochondria without the confound of cerebral blood flow, where pO_2

decreases relate to greater rates of aerobic metabolism and increases reflect less

oxygen consumption[5,24]. We then assessed which cellular processes alter their energy

requirements, the relationship between tissue pO_2 and activity, and how oxygen

consumption varies over a range of network intensities. Collectively, results here

indicate that hibernation uncouples physiological activity from its typical aerobic

requirement, demonstrating that the same neural circuit within a single species can

enter states that require less aerobic respiration while upholding physiological function.

RESULTS

The amphibian respiratory network *in vitro* generates output that occurs as

rhythmic bursts, which are periodically interrupted by high-drive breathing events

termed episodes[25]. While the electrochemical sensor used in this study could not

resolve O_2 changes associated with single bursts which last ~1s, when breathing

episodes spanned several seconds, pO_2 transients of a few mmHg occurred in phase

(Figure 1A). This supports the detection of oxygen consumption within or around the

active respiratory network. Despite having similar baseline network outputs (Figure 1B),

hibernators had higher average tissue pO_2 across the range of depths from the surface

of the brainstem to 350 µm at the baseline external O_2 levels (Figure 1C), and also in 6

additional preparations that were measured down to 650 µm near the center of the

tissue (Supplemental Figure 1). Moreover, lowering the oxygen in the bathing solution

decreased tissue pO_2 as expected, but values were higher in hibernators at most fixed

external pO_2 (Figure 1D). These observations indicate a reduction in brainstem O_2

consumption in hibernators when oxygen is present.

We then used a pharmacological approach to identify which of the energy-

consuming processes use less oxygen. Our rationale was to first block network activity

and then apply a variety of antagonists that inhibit physiological processes involved in

ion regulation that occur in the absence of ongoing network output. For this, we

measured changes in averaged tissue pO_2 during the sequential application of inhibitors

(Supplemental Figure 2) that block network activity (tetrodotoxin; TTX), action potential-

independent spontaneous excitatory postsynaptic potentials (i.e., "minis"; DNQX and

APV), the vacuolar ATPase that pump H^+ into vesicles (V-ATPase; bafilomycin), and the

Na^+-K^+ ATPase that maintains resting ion gradients (ouabain), each of which has the

potential to represent energy sinks in the brain as a result of activity or activity-

independent housekeeping[22,26] (Figure 1E). Finally, to account for remaining O_2

consumption that would arise due to various other activity-independent physiological

processes, we blocked complex IV of the mitochondrial respiratory chain (sodium

cyanide).

In controls, TTX increased tissue pO_2, demonstrating the consumption of O_2 to

power baseline activity. Blocking spontaneous excitatory transmission and then the V-

ATPase had no further effects on tissue pO_2. Inhibition of the Na^+-K^+ ATPase then

raised tissue pO_2, demonstrating a metabolic cost of regulating resting Na^+ and K^+

gradients. Poisoning mitochondria with cyanide led to a large increase in tissue pO_2.

When performing the same protocol on hibernators, less O_2 was consumed by activity

than controls, as TTX led to a significantly smaller rise in tissue pO_2 (p = 0.0198). The

other activity-independent processes consumed similar amounts of O_2 as controls

(DNQX+APV, p = 0.8966; bafilomycin, p = 0.5048; ouabain, p = 0.7841). However,

hibernators showed a significantly smaller pO_2 increase upon blocking mitochondrial

Complex IV with sodium cyanide (p = 0.0003; Figure 1F). While these responses are

presented as ΔpO_2 before and after the sequential application of each drug, they are

corroborated when analyzed absolutely (Supplemental Figure 2). Indeed, hibernators

have a higher starting pO_2 than controls, but no difference in TTX, indicating that TTX

had a smaller effect in hibernators. After this, hibernators and controls had similar tissue

pO_2, indicating no differences across groups with respect to the O_2 consumed by

spontaneous synaptic postsynaptic events, the V-ATPase, and the Na^+-K^+ ATPase.

Cyanide application had a smaller effect in hibernators, reaching lower absolute pO_2

values. Finally bath O_2 did not differ across groups. These results indicate that reduced

oxygen consumption in hibernators is driven primarily by decreasing the cost of network

activity and additional activity-independent mitochondrial O_2 consumption that does not

arise from the Na^+-K^+ ATPase, the V-ATPase, or spontaneous synaptic postsynaptic

events.

Since hibernators had a reduced reliance on aerobic metabolism, we predicted

that activity would be less sensitive to oxygen reductions and the associated decrease

in aerobic metabolism. In control brainstems (n = 6), network activity decreased with

tissue pO_2 during graded reduction in bath O_2. Each control preparation reached a

critical tissue pO_2 (p_{crit}), beyond which the output either lost its rhythmicity or assumed a

slow disrupted pattern of output while decreasing in average frequency (Figure 2A). In

contrast, hibernators maintained normal activity throughout the entire range of graded

hypoxia even at 0 mmHg (Figure 2B & 2C) with no apparent p_{crit} (Figure 2D). These

results indicate that decreasing the reliance on aerobic metabolism allows hibernators

to generate normal network activity across a wide range of tissue oxygen levels, from

supraphysiological to anoxia.

Beyond baseline activity, increased network activity drives oxygen consumption

through synaptic vesicle cycling and restoration of ion gradients for action potentials and

synaptic transmission, which is needed to sustain activity[3,7]. To test whether increased

network activity would accelerate aerobic metabolism, we bath-applied the β-adrenergic

receptor agonist, isoproterenol[27]. Example traces show that increases in activity were

associated with a significant decline in average tissue pO_2 in controls over the 30-

minute application, indicating increased oxygen consumption to meet the demands of

elevated activity (Figure 3A). In contrast, hibernators maintained a stable tissue pO_2

despite achieving comparable increases in burst frequency (Figures 3B-3E). These

results further support that metabolic adaptation in hibernators decouples neural activity

from its typical aerobic support, allowing them to increase motor output without

accelerating oxygen consumption.

Finally, we asked if this metabolic decoupling extends to extreme

hyperexcitability. We induced pathological network activity by blocking fast GABAergic

inhibition with bicuculline, previously shown to generate high-amplitude bursts that

resemble seizure-like activity in controls and hibernators[28]. In controls, bicuculline

triggered abnormal bursts, which were accompanied by a large drop in tissue pO_2,

indicating substantial increases in aerobic respiration (Figures 4A, 4C-4D). Hibernators

also exhibited an initial, large drop in pO_2 following the first abnormal burst,

demonstrating the ability to increase aerobic metabolism when activity goes beyond the

physiological range (Figures 4B-4D). While the absolute decrease in pO_2 from baseline

was the same in controls ($\Delta92 \pm 36.4$ mmHg) and hibernators ($\Delta120 \pm 22.4$ mmHg;

unpaired t-test, $p = 0.31$), after this initial consumption event, hibernators stabilized at a

tissue pO_2 approximately double that of controls (Figures 4B and 4C), indicating that

aerobic metabolism was less under the same hyperexcitable conditions. Despite less

oxygen consumption in the presence of bicuculine, hibernators maintained normal

respiratory network output, while controls produced mainly seizure-like bursts (Figure

4A inset iii, 4D). Therefore, when pushed to the extreme, hibernators meet these

demands by consuming O_2 but ultimately maintain activity homeostasis during

conditions of severe hyperexcitability with less aerobic metabolism.

DISCUSSION

Owing to the high ATP yield of aerobic respiration, increases in oxidative

metabolism are thought to be essential for meeting the energy costs incurred by neural

activity, from flies to mammals[6,29]. Energy limitations in neurological disorders have

prompted interest in animal adaptations as sources of novel mechanisms for improving

metabolic robustness in neural systems[18,30–33]. In frogs, hibernation transforms

brainstem circuits to function during lack of oxygen by altering synapses such that

glycolysis supports activity as a lone ATP source[12,13]. While impressive, we now show

that neural activity can be uncoupled from its support by aerobic metabolism, even

when oxygen is readily available. These results introduce that circuits with a seemingly

obligate requirement for aerobic metabolism can enter a low-cost state that achieves

the same functional output over a wide range of energetic challenges, including hypoxia,

elevated activity, and pathological hyperexcitability, all while consuming less O_2.

Based on the cellular processes addressed, neural activity, activity-independent

Na^+/K^+ ATPase turnover, and additional housekeeping processes that are activity-

independent are the main contributors of O_2 consumption in the brainstem, with minimal

input from spontaneous excitatory postsynaptic currents and the V-ATPase involved in

filling vesicles with neurotransmitter (Figure 1F, Supplemental Figure 2). Despite similar

levels of neural activity, hibernator brainstems exhibited a reduction in O_2 consumption,

driven by less O_2 consumed by neural activity and less spent on other activity-

independent processes. While our results point to a lower cost of activity, they do not

allow us to address which aspects of network activity decrease energetic needs.

Synaptic transmission incurs a greater metabolic cost than action potentials in active

networks[34,35], positioning them as a critical site for reducing energy consumption.

Indeed, NMDA-glutamate receptors, which are involved in generating respiratory motor

bursts, maintain normal postsynaptic current amplitude after hibernation but adopt a

receptor profile that is less permeable to calcium and strongly desensitize, suggesting a

lower cost of ion regulation[23].

We also observed an apparent reduction in the O_2 consumed by activity-

independent processes beyond those contributing to spontaneous synaptic

transmission, the Na^+-K^+ ATPase, and the V-ATPase. As we could not comprehensively

block all additional activity-independent housekeeping process within the brain, applying

cyanide to stop remaining respiration allowed us to assess potential reduction in energy

use that was not captured with inhibitors of specific physiological processes. Therefore,

a smaller change in tissue O_2 upon the application of cyanide suggests less O_2

consumption for activity-independent processes in hibernators. There are two non-

mutually exclusive possibilities that may explain this finding. First, hibernation may

reduce the energy demands of housekeeping processes such as lipid turnover, cellular

cargo transport, and other maintenance functions, each of which can incur substantial

energy costs[24,36]. Second, these results may reflect alterations in mitochondrial function

per se, such as tighter coupling between electron transport and ATP synthesis due to

reduced proton leak across the inner membrane, potentially driven by altered

uncoupling protein function, membrane lipid composition, or cristae structure[37–41].

Therefore, future work directly assessing mitochondrial function and energy spent on

housekeeping physiology will help to discern between these possibilities. Overall,

hibernation leads to a reduction in O_2 consumed in the brainstem through both activity-

dependent and activity-independent physiological processes.

We present three additional lines of evidence that neuronal activity incurs a lower

aerobic cost while maintaining normal function. In controls, activity declined with tissue

pO_2 and had a critical threshold where rhythmic output became disrupted. In contrast,

hibernators were stable from baseline to anoxia and lacked a measurable p_{crit}. This

shows that network function persists as aerobic metabolism decreases, and even

ceases, corroborating our previous reports[12,13]. Further supporting the reduced reliance

on aerobic respiration, increasing activity within the physiological range normally

elevates O_2 consumption, but after hibernation, comparable increases in activity are met

without consuming further O_2. While the specific mechanisms for this are not yet known,

we hypothesize that glycolysis may be upregulated to provide ATP that supports the

increased demands of activity without supplying end-products to the mitochondria

and/or network efficiency increases to such a degree where enhanced activity requires

minimal additional ATP input. Finally, we demonstrate that disinhibition leads to

substantial oxygen consumption in both controls and hibernators, indicating that

although the normal dynamic range of activity is supported with less O_2 consumption,

they retain a capacity to increase aerobic metabolism during pathological

hyperexcitability. Yet, following this initial consumption event, hibernators stabilized at

higher pO_2 values than controls while maintaining normal activity. While reductions in

GABA's contribution to rhythm generation likely play a role[14], these results suggest that

activity may be preserved because of its reduced energetic requirements, improving the

capacity to "share" energetic resources with the elevated metabolic demands of seizure-

like bursts. We present multiple lines of evidence that these circuits shift to consume

less O_2, and we suggest that similar adaptations may extend to other species. For

example, turtles and diving mammals and birds exhibit neural networks capable of

maintaining function under low-oxygen conditions[42,43], suggesting that energy-efficient

network function, as described here, may also apply to species traditionally viewed as

hypoxia-tolerant. Overall, these results challenge the assumption that neuronal output is

constrained by aerobic respiration, highlighting high-efficiency modes of neural

communication as a plastic trait.

One additional intriguing feature of these results is that while activity consumed

less O_2 across a range of intensities (Figure 1-3), hibernator mitochondria do not appear

to be dysfunctional since they retain a capacity to aggressively consume O_2 during

pathological hyperexcitability (Figure 4). This indicates the relationship between activity

and O_2 operates with a reduced sensitivity, rather than being eliminated. The sensitivity

of mitochondrial respiration to neural activity occurs through calcium-dependent

mechanisms[44]. During presynaptic firing, calcium influx from extracellular sources enter

mitochondria through mitochondrial calcium uniporter (MCU) to stimulate oxidative

phosphorylation. MCU's high sensitivity for calcium uptake in neurons is controlled by a

regulatory protein, mitochondrial calcium uptake 3 (MICU3), where its deletion reduces

mitochondrial Ca^{2+} uptake and aerobic metabolism[6,45]. In contrast, MICU1 keeps MCU

in a closed state, while MICU2 permits less calcium entry than MICU3, leading to less

activity-dependent aerobic respiration[6,46,47]. While factors that control the calcium

permeability of the MCU in frogs are not known, it presents a biologically plausible

explanation for alterations in the relationship between activity and O_2 consumption as

we show: Hibernation may reduce mitochondrial calcium permeability, which prevents

neural activity from increasing aerobic metabolism within the physiological range.

However, after crossing a calcium threshold during pathological hyperexcitability, O_2

consumption accelerates. Future work should address the possibility of plasticity in

coupling between activity and calcium-dependent activation of mitochondrial respiration.

MATERIALS AND METHODS

Experimental model

All procedures were approved by the Animal Care and Use Committee at the University

of Missouri (protocol #39264). Adult female American bullfrogs (*Aquarana catesbeiana*,

~100 g; n = 51) were obtained from Rana Ranch (Twin Falls, ID, USA) and housed in

20-gallon tanks containing dechlorinated, aerated water at room temperature on a 12-h

light/dark cycle. Frogs had access to wet and dry platforms, were fed weekly, along with

daily water checks and weekly water changes. Control animals were acclimated for ≥1

week before experiments. A subset of frogs was subjected to a simulated hibernation

protocol. Food was withheld and water temperature was decreased from room

temperature to 4°C over 10 days in a temperature-controlled chamber. Once at 4°C, a

perforated screen was placed below the water surface to enforce continuous

submergence while allowing gas exchange at the air-water interface, and frogs were

maintained for 4 weeks before experimentation. Water pO_2 was approximately 155

mmHg in both control (~22°C) and hibernation (4°C) tanks, consistent with air-

equilibrated water. Frogs subjected to the cold-submergence treatment are referred to

here as "hibernators," following prior work showing that similar protocols induces

metabolic suppression, consistent with the definitions of hibernation that involve

reduced whole-animal metabolism beyond that which is driven by reduced temperature

alone[48].

Brainstem–Spinal Cord Preparation

Brainstem–spinal cord preparations were obtained as described previously[49]. Frogs

were anesthetized with isoflurane (~1 ml/L; v/v) until the toe-pinch reflex was no longer

present, then rapidly decapitated with a guillotine. The head was submerged in ice-cold

bullfrog artificial cerebrospinal fluid (aCSF; in mM: 104 NaCl, 4 KCl, 1.4 $MgCl_2$, 7.5

glucose, 40 $NaHCO_3$, 2.5 $CaCl_2$, and 1 NaH_2PO_4; bubbled with 98.5% O_2/ 1.5% CO_2;

pH 7.85). The forebrain was removed, and the brainstem–spinal cord preparation

(including the midbrain, brainstem, cut caudal to the hypoglossal nerve roots) was

isolated with nerve roots intact and transferred to a recording chamber continuously

superfused with oxygenated aCSF. Chamber temperature was maintained at 22°C

using a Warner Instruments CL-100 bipolar temperature controller with the thermistor

placed adjacent to the tissue. Motor output was recorded from the vagus nerve (cranial

nerve X) using glass suction electrodes. Vagus nerve recordings provide a reliable

index of the overall motor pattern generated by the respiratory network and has been

widely used to assess circuit function in this and similar preparations[15,50,51]. Signals

were AC-amplified (1000×, Model 1700; A-M Systems, Carlsborg, WA), filtered (10Hz-

5kHz), digitized (PowerLab 8/35, ADInstruments, Colorado Springs, CO), rectified, and

integrated online (time constant = 100 ms) to visualize rhythmic motor bursts. Vagus

nerve output was allowed to stabilize for ≥1h in aCSF bubbled with 98% O_2/1.5% CO_2

prior to any experimental manipulation.

Oxygen Profiling

Tissue oxygen partial pressure (pO_2) was measured using a Clark-type microsensor

(OX-50; Unisense A/S, Denmark) with a 50 µm tip diameter, defining a local detection

volume on the order of tens of microns. The sensor has a response time of ~5 s,

resulting in a ~5s temporal averaging of the pO_2 signal with a negligible oxygen

consumption relative to tissue, drawing $4x10^{-4}$-$5×10^{-3}$ nmol O_2/hr during operation. The

sensor was calibrated at 22 °C in air-saturated water and in an anoxic solution (0.1 M

sodium ascorbate in 0.1 M NaOH). The detection limit of the OX-50 sensor is ~0.3

µmol/L (~0.165 mmHg under our experimental conditions; 22°C, ~755 mmHg

atmospheric pressure), with a noise floor of approximately 0.2 mmHg. The analog signal

from a PA2000 amplifier (Unisense A/S) was digitized through an ADC-216USB

converter. For all experiments, the sensor was inserted slightly lateral to the midline of

the brainstem in between the vagus and trigeminal nerves. The tissue pO_2 was

recorded in 50 µm vertical steps from the surface to a depth of 350 µm, an area that

encompasses part of the respiratory network[52]. At each 50 µm vertical step, the sensor

was allowed to stabilize for 20 seconds before recording the pO_2 value, which was

taken as the tissue pO_2 at that depth. In 6 experiments for each group, the sensor was

advanced beyond 350 µM to up to 650 µm, near the center of the tissue.

The amphibian respiratory network tends to produce rhythmic output with bursts often

clustered into stronger multi-breath episodes but can also assume a variety of output

patterns[51]. While the respiratory network is distributed throughout the brainstem[52],

tissue pO_2 fluctuations detected by the microsensor typically occurred in phase with

"crescendo-like" breathing episodes when these occurred (Figure 1A), confirming that

the sensor reported oxygen dynamics within or around the region of the active

respiratory network. The sensor was only able to detect oxygen fluctuations associated

with these high-drive episodes that spanned the 5 s detection duration of the sensor, as

individual, single bursts that are most dominant in this preparation are 1s in duration,

and therefore, likely produce oxygen transients that are faster than the response time of

the sensor. Because the oxygen microsensor reports pO_2 from spatially limited region

around the tip, we also cannot assign the precise anatomical compartment neurons

contributing to each measurement (*e.g.*, in regions densely packed with cell bodies,

synapses, or a combination of the two). The electrode tip may lie closer to somatic

regions, axon fiber tracts, or synaptically dense neuropil, all of which have the potential

for different degrees of oxygen consumption[53]. Thus, the recorded pO_2 reflects the

integrated activity of multiple cellular processes within that microenvironment, which

was recorded as consistently as possible in the same location across experiments.

Targeted assessments of oxygen consumption in sub-cellular compartments require

further investigation.

Pharmacological Experiments

To estimate the contribution of aerobic metabolism to different ATP-consuming

processes, inhibitors specific to each process were sequentially blocked while

monitoring tissue pO_2 at a fixed electrode depth (350 µm). Drugs were applied

cumulatively, i.e., each subsequent inhibitor was added to the superfusate without

washing out the previous drug(s). The following final bath concentrations were applied

in sequence: Tetrodotoxin (TTX, voltage-gated Na^+ channel blocker, 500nM);

DNQX+APV (AMPA receptor antagonist, 10µM; APV (NMDA receptor antagonist,

50µM); Bafilomycin A1 (V-ATPase inhibitor, 1µM); Ouabain (Na^+/K^+-ATPase pump

inhibitor, 500μM); Sodium cyanide (mitochondrial Complex IV inhibitor, 5mM). Each

drug was superfused until a stable plateau in pO_2 was achieved (~15–20 min), and then

the next was applied. Because all previously applied drugs remained in the bath, the

ΔpO_2 following each new inhibitor (the steady-state pO_2 reading before the drug

subtracted from the steady-state reading after) reflects the loss of the O_2 that was being

consumed specifically by the newly blocked process, serving as a functional readout of

the relative metabolic contribution of the targeted pathway. Seven control and seven

hibernated brainstems were used. Although our pharmacological approach isolates

specific pathways powered by aerobic metabolism, the relative magnitude of their pO_2

changes depends on their local abundance within the sampled volume. As a result,

oxygen consumption associated with spontaneous synaptic transmission or vesicle

recycling may be underestimated if the sensor was positioned nearer to neuronal cell

bodies than to synapse-rich zones[53]. These factors are inherent to single-point oxygen

microsensors and reflect the spatial averaging of pO_2 over the sensor's diffusion field

and require further assessments targeting specific sub-cellular recordings.

Graded Hypoxia and Increased Activity

Graded hypoxia was induced by progressively replacing O_2 with N_2 in the aCSF using a

mass-flow controller (MFC-4, Sable Systems International). The N_2 fraction was

increased by 10% every 10 min until anoxia (0% O_2 in the reservoir) was reached.

When bubbled with 98.5% N_2/ 1.5% CO_2, bath pO_2 surrounding the tissue was ~15

mmHg due to mixing at the liquid-air interface. In all controls and hibernators, tissue pO_2

at the sensor reported 0 mmHg; thus, we refer this outcome as "anoxia." Therefore,

while the sensor reported 0 mmHg inside the tissue when bathed with solution gased

with 98.5% N_2/ 1.5% CO_2 (and in some cases at even higher fractions of O_2), it is not

possible to resolve differences in tissue pO_2 below the sensor's detection limit (see

Oxygen Profiling). Regardless, a zero reading under these conditions reflects an

infinitesimal amount of O_2 should any exist in the tissue. Six control and six hibernated

brainstems were tested in this series. Tissue pO_2 and respiratory burst frequency were

recorded continuously (n = 6 controls; n = 6 hibernators). We interpret p_{crit} as the O_2

tension where oxidative phosphorylation decreases to a point that can no longer

preserve normal neural activity, beyond which activity was either chaotic or irregular.

To assess the metabolic cost of elevated physiological activity, isoproterenol (10 µM)

was bath-applied after ≥1 h of stable baseline activity and superfused for 30min (n=7

controls; n=6 hibernators). While other stimuli such as hypercapnia could be used to

stimulate respiratory frequency, in our hands, the responses can be inconsistent from

preparation to preparation, while isoproterenol produces consistent increases in

frequency. To evaluate responses to pathological hyperexcitability, bicuculline

methiodide (10 µM) was superfused for 30 min to induce non-respiratory, seizure-like

bursts (n=6 per group). Non-respiratory bursts were manually counted based on their

prolonged duration and distinct waveform morphology, as previously described[28]. The

oxygen microsensor integrates over ~5s which does not resolve the rapid pO_2 transients

associated with individual respiratory bursts (~1s duration). Therefore, pO_2 data for

these experiments reflect longer-timescale changes in tissue oxygen dynamics, rather

than instantaneous fluctuations tied to single bursts. This averaging is appropriate for

quantifying sustained differences in oxygen consumption over minutes, which were

apparent in these experiments.

Drugs

Tetrodotoxin (TTX), DNQX, APV, and bicuculline were obtained from Hello Bio.

Bafilomycin A1 was purchased from Cayman Chemical. Ouabain was obtained from

Thermo Scientific, sodium cyanide from MP Biomedicals, and isoproterenol from Tokyo

Chemical Industry. All drugs were prepared as stock solutions according to

manufacturer recommendations and diluted to final concentrations in artificial

cerebrospinal fluid immediately prior to use.

Data Analysis

Respiratory burst frequency was determined from rectified and integrated vagus

recordings using the Peak Analysis function in LabChart 8 (ADInstruments). Bursts

were identified by standard criteria (~1 s duration) with start/stop thresholds at 5% of

peak height[50]. For graded hypoxia experiments, mean burst frequency and tissue pO_2

were calculated for every 10 mmHg drop in tissue pO_2 relative to baseline at 98% O_2.

For experiments involving sequential drug exposures, mean pO_2 values were obtained

over the final 10 min of each treatment at steady-state. During the application of

isoproterenol, pO_2 values were taken at baseline and then at 10 minute intervals

sampling for 2 minutes during the 30 minute exposure. For bicuculine, pO_2 values were

taken at baseline, at the trough of the large dip caused by the large non-respiratory

burst following bicuculine, and then the last 5 minutes of the 30-minute exposure

Statistical analysis

Data are presented as mean ± SD or box and whisker plots unless otherwise stated.

Statistical comparisons were performed in GraphPad Prism v10.6.1 (San Diego, CA,

USA). Two-way repeated measures ANOVA was applied with two independent

variables and one dependent variable (e.g., the effect of depth and group on tissue pO_2).

In these cases, Holm–Šidák multiple comparisons test was applied after if there were

any interaction or group effects. To assess the effects of tissue pO_2 and burst frequency,

we used a mixed linear model with restricted maximum likelihood, as this dataset had

many missing values due to the differing starting points of pO_2 from preparation to

preparation. Unpaired t tests were used to compare two independent groups. Individual

data points are shown where appropriate to show variability among preparations.

CONCLUSION

Our findings reveal that neural circuits in the vertebrate brain can undergo a

fundamental reorganization of the relationship between activity and O_2 consumption. By

reducing the aerobic requirement for neural activity, the frog brainstem maintains stable

network output under metabolic conditions that normally induce severe perturbations.

The relevance of this reduced aerobic cost is likely for emergence from hibernation,

when frogs must restart the neural circuits that drive breathing before the first breath is

taken, at which point brain oxygenation has not yet been restored by lung ventilation. A

central element of this shift appears to be the enhanced efficiency of network function,

which allows hibernators to generate normal output with less ongoing aerobic

metabolism. These results raise the question of how neural circuits can transition

between vastly different aerobic requirements while maintaining similar functional output.

Understanding these mechanisms may be particularly relevant for neurological

disorders such as Alzheimer's and Parkinson's disease, where impaired aerobic

metabolism compromises circuit function[8]. By identifying mechanisms that allow circuits

transition between states with vastly different aerobic demands needed to produce

similar function, our work highlights how natural adaptation in diverse animal models

can serve as valuable frameworks for exploring strategies to improve metabolic

resilience in the mammalian brain.

Acknowledgements: The authors would like to thank Nik Bueschke and Jose Viteri for

comments on the manuscript.

Conflict of interest disclosure: The authors do have conflicts of interest to declare.

REFERENCES

1.   Attwell, D., and Laughlin, S.B. (2001). An Energy Budget for Signaling in the Grey

Matter of the Brain. Journal of Cerebral Blood Flow & Metabolism *21*, 1133–1145.

https://doi.org/10.1097/00004647-200110000-00001.

2.   Niven, J.E., and Laughlin, S.B. (2008). Energy limitation as a selective pressure

on the evolution of sensory systems. Journal of Experimental Biology *211*, 1792–

1804. https://doi.org/10.1242/jeb.017574.

3.   Trevisiol, A., Saab, A.S., Winkler, U., Marx, G., Imamura, H., Möbius, W., Kusch,

K., Nave, K.-A., and Hirrlinger, J. (2017). Monitoring ATP dynamics in electrically

active white matter tracts. Elife *6*. https://doi.org/10.7554/eLife.24241.

4.   Hall, C.N., Klein-Flugge, M.C., Howarth, C., and Attwell, D. (2012). Oxidative

Phosphorylation, Not Glycolysis, Powers Presynaptic and Postsynaptic

Mechanisms Underlying Brain Information Processing. Journal of Neuroscience

32, 8940–8951. https://doi.org/10.1523/JNEUROSCI.0026-12.2012.

5.   Özugur, S., Kunz, L., and Straka, H. (2020). Relationship between oxygen
consumption and neuronal activity in a defined neural circuit. BMC Biol. *18*, 76.
https://doi.org/10.1186/s12915-020-00811-6.

6.   Ashrafi, G., de Juan-Sanz, J., Farrell, R.J., and Ryan, T.A. (2020). Molecular
Tuning of the Axonal Mitochondrial Ca2+ Uniporter Ensures Metabolic Flexibility
of Neurotransmission. Neuron *105,* 678-687.e5.
https://doi.org/10.1016/j.neuron.2019.11.020.

7.   Rangaraju, V., Calloway, N., and Ryan, T.A. (2014). Activity-Driven Local ATP
Synthesis Is Required for Synaptic Function. Cell *156*, 825–835.
https://doi.org/10.1016/j.cell.2013.12.042.

8.   Li, S., and Sheng, Z.-H. (2022). Energy matters: presynaptic metabolism and the
maintenance of synaptic transmission. Nat. Rev. Neurosci. *23*, 4–22.
https://doi.org/10.1038/s41583-021-00535-8.

9.   Chklovskii, D.B., Schikorski, T., and Stevens, C.F. (2002). Wiring Optimization in
Cortical Circuits. Neuron *34,* 341–347. https://doi.org/10.1016/S0896-
6273(02)00679-7.

10. Adams, S., Zubov, T., Bueschke, N., and Santin, J.M. (2021). Neuromodulation or
energy failure? Metabolic limitations silence network output in the hypoxic
amphibian brainstem. Am. J. Physiol. Regul. Integr. Comp. Physiol. *320*, R105–
R115. https://doi.org/10.1152/AJPREGU.00209.2020.

11. Winmill, R.E., Chen, A.K., and Hedrick, M.S. (2005). Development of the
respiratory response to hypoxia in the isolated brainstem of the bullfrog Rana

catesbeiana. Journal of experimental biology *208*, 213–222.

https://doi.org/10.1242/jeb.01399.

12. Bueschke, N., Do Amaral-Silva, L., Adams, S., and Santin, J.M. (2021).

Transforming a neural circuit to function without oxygen and glucose delivery.

Current Biology *31*, R1564–R1565. https://doi.org/10.1016/j.

13. Amaral-Silva, L., and Santin, J.M. (2023). Synaptic modifications transform neural

networks to function without oxygen. BMC Biol. *21*, 54.

https://doi.org/10.1186/s12915-023-01518-0.

14. Saunders, S.E., and Santin, J.M. (2024). Hibernation reduces GABA signaling in

the brainstem to enhance motor activity of breathing at cool temperatures. BMC

Biol. *22*, 251. https://doi.org/10.1186/s12915-024-02050-5.

15. Kottick, A., Baghdadwala, M.I., Ferguson, E. V., and Wilson, R.J.A. (2013).

Transmission of the respiratory rhythm to trigeminal and hypoglossal motor

neurons in the American Bullfrog (Lithobates catesbeiana). Respir. Physiol.

Neurobiol. *188*, 180–191. https://doi.org/10.1016/j.resp.2013.06.008.

16. Chen, A.K., and Hedrick, M.S. (2008). Role of glutamate and substance P in the

amphibian respiratory network during development. Respir. Physiol. Neurobiol.

162, 24–31. https://doi.org/10.1016/j.resp.2008.03.010.

17. Ultsch, G.R., Reese, S.A., and Stewart, E.R. (2004). Physiology of hibernation in

Rana pipiens: metabolic rate, critical oxygen tension, and the effects of hypoxia

on several plasma variables. J. Exp. Zool. A Comp. Exp. Biol. *301A*, 169–176.

https://doi.org/10.1002/jez.a.20014.

18.   Amaral-Silva, L., and Santin, J. (2024). Neural Processing without O2 and

Glucose Delivery: Lessons from the Pond to the Clinic. Physiology *39*, 360–372.

https://doi.org/10.1152/physiol.00030.2023.

19.   Hardie, D.G. (2000). Metabolic control: A new solution to an old problem. Current

Biology *10*, R757–R759. https://doi.org/10.1016/S0960-9822(00)00744-2.

20.   Sengupta, B., Stemmler, M., Laughlin, S.B., and Niven, J.E. (2010). Action

Potential Energy Efficiency Varies Among Neuron Types in Vertebrates and

Invertebrates.   PLoS   Comput.   Biol.   *6*,   e1000840.

https://doi.org/10.1371/journal.pcbi.1000840.

21.   Harris, J.J., Jolivet, R., Engl, E., and Attwell, D. (2015). Energy-Efficient

Information Transfer by Visual Pathway Synapses. Current Biology *25*, 3151–

3160. https://doi.org/10.1016/j.cub.2015.10.063.

22.   Alle, H., Roth, A., and Geiger, J.R.P. (2009). Energy-Efficient Action Potentials in

Hippocampal   Mossy   Fibers.   Science   (1979).   *325*,   1405–1408.

https://doi.org/10.1126/science.1174331.

23.   Bueschke, N., Amaral-Silva, L., Hu, M., Alvarez, A., and Santin, J.M. (2024).

Plasticity in the Functional Properties of NMDA Receptors Improves Network

Stability during Severe Energy Stress. Journal of Endocrinology *44*, 1–12.

https://doi.org/10.1523/JNEUROSCI.0502-23.2024.

24.   Harris, J.J., and Attwell, D. (2012). The Energetics of CNS White Matter. The

Journal of Neuroscience *32*, 356–371. https://doi.org/10.1523/JNEUROSCI.3430-

11.2012.

25.   Milsom, W.K., Reid, S.G., Meier, J.T., and Kinkead, R. (1999). Central respiratory
pattern generation in the bullfrog, Rana catesbeiana. Comparative Biochemistry
and   Physiology   Part   A   *124*,   253–264.   https://doi.org/10.1016/S1095-
6433(99)00113-0.

26.   Pulido, C., and Ryan, T.A. (2021). Synaptic vesicle pools are a major hidden
resting   metabolic   burden   of   nerve   terminals.   Sci.   Adv.   *7*.
https://doi.org/10.1126/sciadv.abi9027.

27.   Cannon, D.J., and Santin, J.M. (2025). Homeostatic regulation of a motor circuit
through temperature sensing rather than activity sensing. Current Biology *35*,
2256-2265.e3. https://doi.org/10.1016/j.cub.2025.03.054.

28.   Broch, L., Morales, R., Sandoval, A., and Hedrick, M. (2002). Regulation of the
respiratory central pattern generator by chloride-dependent inhibition during
development in the bullfrog (Rana catesbeiana). Journal of Experimental Biology
205, 1161–1169. https://doi.org/10.1242/jeb.205.8.1161.

29.   Justs, K.A., Lu, Z., Chouhan, A.K., Borycz, J.A., Lu, Z., Meinertzhagen, I.A., and
Macleod, G.T. (2022). Presynaptic Mitochondrial Volume and Packing Density
Scale with Presynaptic Power Demand. The Journal of Neuroscience *42*, 954–
967. https://doi.org/10.1523/JNEUROSCI.1236-21.2021.

30.   Meir, J.U., Champagne, C.D., Costa, D.P., Williams, C.L., and Ponganis, P.J.
(2009). Extreme hypoxemic tolerance and blood oxygen depletion in diving
elephant seals. Am J Physiol Regul Integr Comp Physiol *297*, 927–939.
https://doi.org/10.1152/ajpregu.00247.2009.-Species.

31.   Buck, L.T., and Pamenter, M.E. (2018). The hypoxia-tolerant vertebrate brain:

538   Arresting synaptic activity. Comp. Biochem. Physiol. B Biochem. Mol. Biol. *224*,

539   61–70. https://doi.org/10.1016/j.cbpb.2017.11.015.

540   32.   Cunnane, S.C., Trushina, E., Morland, C., Prigione, A., Casadesus, G., Andrews,

541   Z.B., Beal, M.F., Bergersen, L.H., Brinton, R.D., de la Monte, S., et al. (2020).

542   Brain energy rescue: an emerging therapeutic concept for neurodegenerative

543   disorders of ageing. Nat. Rev. Drug Discov. *19,* 609–633.

544   https://doi.org/10.1038/s41573-020-0072-x.

545   33.   Larson, J., Drew, K.L., Folkow, L.P., Milton, S.L., and Park, T.J. (2014). No

546   oxygen? No problem! intrinsic brain tolerance to hypoxia in vertebrates. Journal of

547   Experimental Biology *217*, 1024–1039. https://doi.org/10.1242/jeb.085381.

548   34.   Harris, J.J., Jolivet, R., and Attwell, D. (2012). Synaptic Energy Use and Supply.

549   Neuron *75*, 762–777. https://doi.org/10.1016/j.neuron.2012.08.019.

550   35.   Howarth, C., Gleeson, P., and Attwell, D. (2012). Updated Energy Budgets for

551   Neural Computation in the Neocortex and Cerebellum. Journal of Cerebral Blood

552   Flow & Metabolism *32*, 1222–1232. https://doi.org/10.1038/jcbfm.2012.35.

553   36.   Engl, E., and Attwell, D. (2015). Non-signalling energy use in the brain. J. Physiol.

554   *593*, 3417–3429. https://doi.org/10.1113/jphysiol.2014.282517.

555   37.   Ballinger, M.A., Schwartz, C., and Andrews, M.T. (2017). Enhanced oxidative

556   capacity of ground squirrel brain mitochondria during hibernation. J Physiol Regul

557   Integr Comp Physiol *312,* 301–310. https://doi.org/10.1152/ajpregu.00314.2016.-

558   Dur.

38.   Salin, K., Auer, S.K., Rey, B., Selman, C., and Metcalfe, N.B. (2015). Variation in
the link between oxygen consumption and ATP production, and its relevance for
animal performance. Proceedings of the Royal Society B: Biological Sciences *282*,
20151028. https://doi.org/10.1098/rspb.2015.1028.

39.   Divakaruni, A.S., and Brand, M.D. (2011). The Regulation and Physiology of
Mitochondrial Proton Leak. Physiology *26*, 192–205.
https://doi.org/10.1152/physiol.00046.2010.

40.   Hoang, T., Kuljanin, M., Smith, M.D., and Jelokhani-Niaraki, M. (2015). A
biophysical study on molecular physiology of the uncoupling proteins of the
central nervous system. Biosci. Rep. *35*, e00226.
https://doi.org/10.1042/BSR20150130.

41.   Decker, S.T., and Funai, K. (2024). Mitochondrial membrane lipids in the
regulation of bioenergetic flux. Cell Metab. *36*, 1963–1978.
https://doi.org/10.1016/j.cmet.2024.07.024.

42.   Ramirez, J.-M., Folkow, L.P., and Blix, A.S. (2007). Hypoxia Tolerance in
Mammals and Birds: From the Wilderness to the Clinic. Annu. Rev. Physiol. *69*,
113–143. https://doi.org/10.1146/annurev.physiol.69.031905.163111.

43.   Ariel, M., Ahuja, S., and Warren, D.E. (2023). Responses of the in vitro turtle brain
to visual and auditory stimuli during severe hypoxia. Journal of Experimental
Biology *226*, jeb244687. https://doi.org/10.1242/jeb.244687.

44.   Hayakawa, Y., Nemoto, T., Iino, M., and Kasai, H. (2005). Rapid Ca2+-dependent

580   increase in oxygen consumption by mitochondria in single mammalian central

581   neurons. Cell Calcium *37*, 359–370. https://doi.org/10.1016/j.ceca.2004.11.005.

582   45.   Patron, M., Granatiero, V., Espino, J., Rizzuto, R., and De Stefani, D. (2019).

583   MICU3 is a tissue-specific enhancer of mitochondrial calcium uptake. Cell Death

584   Differ. *26*, 179–195. https://doi.org/10.1038/s41418-018-0113-8.

585   46.   Mallilankaraman, K., Doonan, P., Cárdenas, C., Chandramoorthy, H.C., Müller, M.,

586   Miller, R., Hoffman, N.E., Gandhirajan, R.K., Molgó, J., Birnbaum, M.J., et al.

587   (2012). MICU1 Is an Essential Gatekeeper for MCU-Mediated Mitochondrial Ca2+

588   Uptake   that   Regulates   Cell   Survival.   Cell   *151*,   630–644.

589   https://doi.org/10.1016/j.cell.2012.10.011.

590   47.   Patron, M., Checchetto, V., Raffaello, A., Teardo, E., Vecellio Reane, D., Mantoan,

591   M., Granatiero, V., Szabò, I., De Stefani, D., and Rizzuto, R. (2014). MICU1 and

592   MICU2 Finely Tune the Mitochondrial Ca2+ Uniporter by Exerting Opposite

593   Effects   on   MCU   Activity.   Mol.   Cell   *53*,   726–737.

594   https://doi.org/10.1016/j.molcel.2014.01.013.

595   48.   Donohoe, P.H., West, T.G., and Boutilier, R.G. (1998). Respiratory, metabolic,

596   and acid-base correlates of aerobic metabolic rate reduction in overwintering

597   frogs. American Journal of Physiology-Regulatory, Integrative and Comparative

598   Physiology *274*, R704–R710. https://doi.org/10.1152/ajpregu.1998.274.3.R704.

49.   Burton, M.T., and Santin, J.M. (2020). A direct excitatory action of lactate ions in

the central respiratory network of bullfrogs, Lithobates catesbeianus. Journal of

Experimental Biology *223*, jeb235705. https://doi.org/10.1242/jeb.235705.

50.   Kogo, N., Perry, S.F., and Remmers, J.E. (1994). Neural organization of the

ventilatory activity in the frog, Rana catesbeiana. J. Neurobiol. *25*, 1067–1079.

https://doi.org/10.1002/neu.480250904.

51.   Reid, S.G., and Milsom, W.K. (1998). Respiratory pattern formation in the isolated

bullfrog (Rana catesbeiana) brainstem-spinal cord. Respir. Physiol. *114*, 239–255.

https://doi.org/10.1016/S0034-5687(98)00091-7.

52.   Oku, Y., Kimura, N., Masumiya, H., and Okada, Y. (2008). Spatiotemporal

organization of frog respiratory neurons visualized on the ventral medullary

surface. Respir. Physiol. Neurobiol. *161*, 281–290.

https://doi.org/10.1016/j.resp.2008.03.002.

53.   Lecoq, J., Tiret, P., Najac, M., Shepherd, G.M., Greer, C.A., and Charpak, S.

(2009). Odor-Evoked Oxygen Consumption by Action Potential and Synaptic

Transmission in the Olfactory Bulb. The Journal of Neuroscience *29*, 1424–1433.

https://doi.org/10.1523/JNEUROSCI.4817-08.2009.

Figures and Legends

Figure 1. Hibernators have lower apparent oxygen consumption due to reduced

energy demands of network activity and basal mitochondrial respiration.

A. Schematic of the brainstem preparation with an oxygen sensor placed in the

proposed rhythm-generating region. Example recording shows synchronized

vagus nerve activity, network frequency, and corresponding tissue pO_2

fluctuations.

B. Control and hibernator brainstems showed similar baseline burst frequency

(Welch's unpaired t-test, p = 0.852; n = 19/group).

C. Tissue pO_2 depth profiles up to 350 μm showed significantly higher pO_2 in

hibernators (n=19) when compared to controls (n=18); two-way repeated

measures ANOVA; main effect of group, $F_{(1,35)}$ = 31.18, p< 0.0001; main effect

of depth, $F_{(1.964, 68.74)}$ = 590.9, p < 0.0001; depth x group interaction, $F_{(7,245)}$

= 0.4117, p = 0.8946; Holm–Šídák multiple comparisons test; ****p < 0.0001 at

all depths.

D. During gradual external pO_2 reduction, hibernator brainstems maintained higher

tissue pO_2 throughout hypoxia induction (n = 6/group; two-way repeated

measures ANOVA; main effect of group, $F_{(1, 10)}$ = 9.826, p = 0.0106 ; main

effect of pO_2, $F_{(1,10)}$=145.2, p<0.0001; pO_2 × group interaction, $F_{(1.70, 11.7)}$ =

4.929, p = 0.0426. Holm–Šídák multiple comparisons test, **p < 0.01, *p < 0.05).

E. An illustration of major energy consuming pathways at the synapse with the

blocked pathways numbered (made in Biorender).

F. Summary of ΔpO_2 after blocking voltage-gated Na^+ channels, AMPA and NMDA

receptors, vesicular ATPase, Na^+-K^+ ATPase and mitochondrial Complex IV (n =

7/group). Two-way repeated measures ANOVA revealed a significant main effect

of group, ($F_{(1, 12)}$ = 122.3, p < 0.0001); main effect of drug, ($F_{(4, 48)}$ = 139.9, p<

0.0001 and drug × group interaction ($F_{(4, 48)}$ = 6.213, p = 0.0004). Hibernators

showed significantly smaller ΔpO_2 with TTX (p = 0.01) and sodium cyanide (p =

0.0003; Holm–Šídák multiple comparisons test).

Figure 2. Brainstem motor activity no longer requires aerobic respiration after

hibernation

A. Example of graded hypoxia experiment in a control, where nerve burst frequency

declined strongly at a critical threshold (p_{crit}).

B. In hibernators (n = 6), neural activity remained stable even as tissue pO_2 at ~0

mmHg.

C. Mean ± SD of fold change in burst frequency are shown for control and

hibernator (n = 6 each) groups. Controls show a lower starting pO_2 with an

inflection point and subsequent decline in burst frequency as pO_2 decreases,

whereas hibernators have higher starting pO_2 and remain relatively stable across

the pO_2 range. P values of group and pO_2 from linear mixed model are shown in

the figure.

663   D. Critical pO_2 thresholds: controls lost rhythmic activity at significantly higher pO_2

664   than hibernators, which showed no apparent critical pO_2 threshold as all

665   preparations were active at 0 mmHg, and therefore assigned a p_{crit} value of 0

666   mmHg (Welch's unpaired t-test, $t(5) = 6.847$, $p = 0.001$; $n = 6$ per group).

Figure 3. Hibernators sustain elevated neural activity without consuming more

oxygen.

A. Representative traces showing changes in tissue pO_2 and vagus nerve activity

during isoproterenol exposure in a control brainstem. Increased burst frequency

caused a drop in tissue pO_2. The insets show nerve activity of control brainstem

at baseline and during isoproterenol exposure.

B. Tissue pO_2 in hibernators remained stable during elevated burst frequency. The

insets show nerve activity of hibernator brainstem at baseline and during

isoproterenol exposure.

C. Fold change in burst frequency during isoproterenol application for controls (n=7)

and hibernators (n=6; Welch's unpaired t-test, t(7.931) = 1.302, p = 0.229).

D. Absolute tissue pO_2 at different timepoints of isoproterenol application (n = 7

controls, n = 6 hibernators). Two-way repeated measures ANOVA; main effect of

group, F(1, 11) = 51.09, p < 0.001; main effect of time, F(2.038, 22.42) = 10.92, p

= 0.0005; time × group interaction, F(2.038, 22.42) = 7.1, p = 0.0037. ****

indicates p<0.0001 between controls and hibernators at all time points, and

■■p<0.01, and ■ p<0.05 indicate differences from baseline within the control

group (Holm–Šídák multiple comparisons test).

E. ΔpO_2 at 10, 20, and 30 minutes of isoproterenol application for controls and

hibernators n = 6). Two-way repeated measures ANOVA; main effect of group,

F(1, 11) = 16.2, p = 0.0020; main effect of time, F(1.338,14.71) = 4.9, p = 0.338;

time × group interaction, F(2, 22) = 1.6, p = 0.2173; Holm–Šídák multiple

comparisons test, **p < 0.01, *p < 0.05.

Figure 4. Hibernators rely on aerobic metabolism during pathological hyperexcitability but maintain elevated tissue oxygen.

A. Representative recordings of tissue pO$_2$ and vagus nerve activity in a control brainstem during application of the GABA$_A$ receptor antagonist, bicuculline. Expanded nerve traces show baseline respiratory activity (RESP), the first large chaotic burst during bicuculline application, and then at the end of the 30 minute bicuculline exposure.

B. Representative recordings of tissue pO_2 and nerve activity in a hibernator

brainstem during application of bicuculline. Expanded nerve traces show

baseline respiratory activity (RESP), the first large chaotic burst during

bicuculline application, and then at the end of the 30-minute bicuculline exposure.

C. Absolute tissue pO_2 at baseline, immediately after the first major non-respiratory

burst (post-NR), and at the end of bicuculline treatment in control and hibernator

brainstems (n = 6 per group). Both controls and hibernators face a significant

drop in tissue pO_2 post-NR, however tissue pO_2 settles at a higher level in

hibernators as compared to controls. Two-way repeated measures; main effect of

group, $F(1, 10) = 29$, $p = 0.0003$; main effect of time, $F(1.438, 14.38) = 39244$, p

< 0.0001; time × group interaction, $F(2, 20) = 1.6$, $p = 0.2224$; Holm–Šídák

multiple comparisons test, $***p < 0.001$, $**p < 0.01$.

D. Number of normal bursts per minute after 30 minutes of bicuculline application in

control and hibernator brainstem (n=6 per group; Welch's unpaired t-test, $t(5) =$

11.15, $p < 0.0001$).