Title: Mutations and predicted glycosylation patterns in respiratory syncytial virus isolates correlate with

disease severity.

Authors: Hunte Morgan L[1,2], Herbst Katherine W [3], Michelow Ian C[** 3,4,5], Szczepanek Steven M[* 1,2]

1. Department of Pathobiology, University of Connecticut, Storrs, CT, USA

2. Center of Excellence for Vaccine Research, University of Connecticut, Storrs, CT, USA

3. Connecticut Children's Research Institute, Connecticut Children's, Hartford, CT, USA

4. Division of Infectious Disease and Immunology, Connecticut Children's, Hartford, CT, USA

5. Department of Pediatrics, University of Connecticut School of Medicine, Farmington, CT, USA

*Corresponding author:

Steven M. Szczepanek, Ph.D.
University of Connecticut
61 North Eagleville Road (Atwater Laboratory)
Storrs, CT 06238
Ph: 860-486-8101
Steven.Szczepanek@uconn.edu
18

**Co-Corresponding author:

Ian C. Michelow, M.D
Connecticut Children's
85 Seymour Street, Suite 816
Hartford, CT 06106
Ph: 860-545-9490
Fax: 860-545-9371
IMichelow@connecticutchildrens.org

Abstract

Respiratory syncytial virus (RSV) remains an important cause of lower respiratory tract infections in

young children, producing mild to life-threatening disease. Although rapid viral evolution through genetic

drift is well established, the structural and functional impacts of specific pathoadaptive mutations linked

to enhanced virulence are poorly defined. We investigated these relationships by isolating RSV from

available nasal swabs of five hospitalized infants during the 2022–2023 winter season and conducting

comparative viral genomic analysis. Severity of disease was evaluated using a validated clinical scoring

system. Whole-genome sequencing followed by reference-guided assembly and structural modeling

revealed distinct amino acid polymorphisms correlating with disease severity. Phylogenetic analysis

placed all isolates within the RSV-A GA2.3.5 G clade. Isolates from mild moderate and severe cases

clustered in A.D.1.5 and A.D.1.8 subclades. Nineteen amino acid differences were associated with clinical

severity and isolates from moderate or severe cases replicated more rapidly in vitro than mild isolates.

Computational glycosylation predictions indicated an increasing number of glycosylation sites in the G

protein corresponding with greater disease severity. Together, these data suggest that specific

pathoadaptive mutations may contribute to enhanced viral replication and severity, and are relevant for

future surveillance efforts and the development of immune-based strategies targeting virulence-associated

residues.

Key Words: Respiratory syncytial virus, genetic mutations, clinical severity, attachment glycoprotein,

RNA polymerase

Introduction

Respiratory syncytial virus (RSV) remains a leading global cause of lower respiratory tract infections in infants and young children (1–3). Virtually all children acquire RSV infection at least once by 2 years of age (4). Although premature neonates, immunocompromised individuals and those with congenital heart, chronic lung or neuromuscular disorders are at great risk for severe disease, most severe cases occur in previously healthy, full-term infants. Factors contributing to severe disease in healthy children remain unknown (5,6).

The intrinsic mutability of RSV, owing to its RNA-dependent RNA polymerase, allows it to explore a vast sequence space, generating variants with potential fitness advantages (7,8). While the fundamental biology of RSV remains conserved, multiple studies have reported that clinical outcomes may be influenced by specific, clade-defining mutations (9–16). However, this association is subject to debate owing to the influence of confounding factors such as patient age, prematurity, comorbidities, prior immunity, environmental factors, demographic and geographic variability, as well as heterogeneity in severity metrics and study methodologies (17–20).

Upon entering the human respiratory tract, RSV initiates a highly coordinated infection process driven by its eleven proteins encoded in a ~15,000 base pair negative-sense RNA genome. Collectively, the concerted actions of these eleven proteins – NS1, NS2, N, P, M, SH, G, F, M2-1, M2-2, and L – enable RSV to induce the profound airway inflammation and epithelial injury frequently associated with RSV disease. There is a critical knowledge gap in linking specific RSV genotypes to clinical severity through integrated genomic, structural, and functional analyses (9,19). A barrier to further reducing the burden of RSV disease is the incomplete understanding of the viral and host determinants that drive the heterogeneity of clinical outcomes. This proof-of-concept study investigated the hypothesis that specific amino acid polymorphisms in RSV are associated with clinical disease severity by means of comparative genomics. We sequenced RSV isolates from children enrolled at Connecticut Children's during a single

season and correlated differences in the viral protein sequences with validated clinical disease severity

scores.

Methods

Subjects and biospecimens

We selected eligible subjects with RSV who had been enrolled as controls in a single-center COVID-19

biorepository from November 2022 to March 2023 (IRB approval #20-055). Inclusion criteria for the

current study were defined as subjects under 24 months of age who were hospitalized for treatment of

RSV mono-infection confirmed by a commercial RT-PCR assay. Demographic data were collected via in-

person surveys and clinical data were abstracted from inpatient records. The primary outcome was

defined by the previously validated Clinical Disease Severity Score (CDSS) based on the subject's most

severe condition during their hospitalization (**S1 Table**) (22). The CDSS is an objective measure

comprised of five parameters: 1) respiratory rate, 2) auscultation, 3) transcutaneous O_2 saturation, 4)

retraction, and 5) activity level (mainly assessed by feeding difficulty). Each parameter ranks from 0

(normal) to 3 (most abnormal), and subjects were classified as having mild (0-5), moderate (6-10), or

severe (11-15) RSV (22). Mid-turbinate nasal fluid was collected at time of enrollment using BD

Universal Viral Transport Collection Kits (ThermoFisher Scientific, Cat #B220529). Vials were vortexed

for 20 seconds, media was divided into 500 µL aliquots, and samples were stored at -80 °C until used.

Isolation of RNA from Mid-turbinate nasal swabs and total RNA QC

RNA was isolated from mid-turbinate nasal swab specimens using the Direct-zol™ RNA Miniprep Kit

(Zymo Research) according to the manufacturer's instructions. Total RNA was quantified and purity

ratios determined for each sample using the NanoDrop 2000 spectrophotometer (ThermoFisher Scientific,

Waltham, MA, USA). To further assess RNA quality, total RNA was analyzed on the Agilent TapeStation

4200 (Agilent Technologies, Santa Clara, CA, USA) using the RNA High Sensitivity assay.

Illumina RNA Enrichment Library preparation and Sequencing

Total RNA samples were prepared for RNA enrichment using the Illumina RNA for Enrichment Sample

Preparation with the Respiratory Virus Oligo Panel (RVOP) following the manufacturer's protocol

(Illumina, San Diego, CA, USA). Libraries were validated for length and adapter dimer removal using

the Agilent TapeStation 4200 D1000 High Sensitivity assay (Agilent Technologies, Santa Clara, CA,

USA), then quantified and normalized using the dsDNA High Sensitivity Assay for Qubit 3.0 (Life

Technologies, Carlsbad, CA, USA). Sample libraries were prepared for Illumina sequencing by

denaturing and diluting the libraries per manufacturer's protocol (Illumina, San Diego, CA, USA). All

samples were pooled into one sequencing pool, equally normalized, and run as one sample pool across the

Illumina NovaSeq 6000 using version 1.5 chemistry. Target read depth of 2M reads was achieved per

sample with paired end 100bp reads.

Sequence analysis pipeline

Viral genomic analysis was performed using Illumina paired-end sequencing data processed through a

reference-guided assembly pipeline. Raw reads underwent quality control with FastP and FastQC

software, which trimmed low-quality bases (Q-score <20) and adapter sequences. To exclude non-RSV

genetic material, unassembled but trimmed reads were taxonomically classified with Kraken2 software

(23) against the standard Kraken Database. Only sequences assigned to Human orthopneumovirus (RSV

taxon ID: 208895) were filtered and retained using SeqTK software. Filtered RSV-specific reads were

aligned to the RSV-A reference genome (NCBI accession: KT992094.1) using BWA-MEM (v0.7.17) [28]

with default settings. Consensus RSV genomes generated through reference-guided assembly were

annotated using VIGOR4 (24).

Sequence comparisons and analysis of point mutations

Genomes of the RSV clinical isolates were analyzed using Nextstrain's web-based platform, Nextclade

(v3.13.2) (25) to assign clades, identify mutations, and perform additional quality checks. Sequences were

aligned with global RSV-A genomes from GenBank via Nextstrain's automated pipeline. For mutation

profiling, Nextstrain's variant-calling module identified amino acid polymorphisms relative to the RSV A

reference (EPI_ISL_412866). Clade classification followed Nextstrain's nomenclature, supported by

bootstrap values >70%.

Culture of RSV from clinical isolates

A549 cells (ATCC CC-185) were seeded in DMEM (Gibco, Ref # 31966-021) + 1% GlutaMAX (Gibco,

35050-061) supplemented with 10% fetal bovine serum (Gibco, Ref# A56707-01) and 1% Anti-Anti

(Gibco, Ref # 15240-062). Cells were allowed to adhere overnight. After 12 hours post-seed

(approximately 95% confluent cells) were infected with each clinical isolate at an MOI of 0.01 for 2

hours. Inoculum was removed, cells were washed with PBS (Gibco, Ref# 2085290), and cell culture

media was added. Plates were incubated in 5% CO_2 at 37 °C for 24, 48, 72, or 96 hours post-infection

(HPI).

Determination of viral load

Extracellular viral load was determined using the 50% tissue culture infectious dose ($TCID_{50}$) assay. The

$TCID_{50}$ assay was performed by infecting approximately 95% confluent A549 cell cultures cells in 96-

well plates. Serial tenfold dilutions of each sample were prepared in eight replicates in infection medium.

After washing the cell monolayers, 100 µL of each dilution was added per well and incubated at 37°C for

2 hours. An additional 75 µL of infection medium was then added, and plates were incubated for 4–5

days. After incubation, infection medium was aspirated then cells were fixed and stained with crystal

violet. $TCID_{50}$ was calculated using the Reed-Muench method.

Structural Analyses

L polymerase structure predictions were obtained using homology-based modelling based on RSV RNA

polymerase (PDB: 6UEN). N-linked and O-linked glycosylation predictions were performed using

NetNGlyc 1.0 (30) and NetOGlyc 4.0 (31), respectively.

Results/Discussion

This proof-of-concept study provides insights into molecular determinants of RSV virulence by

integrating genomic and phenotypic analyses with measures of clinical disease severity. After excluding

three subjects due to co-detection of SARS-CoV-2 (n=1), adenovirus (n=1), rhinovirus/enterovirus (n=1),

we analyzed RSV isolates from five infants who had no comorbidities (**Table 1**). Nasal samples from the

five subjects were collected a median of five days after start of symptoms (IQR, 4 - 6 days). Two subjects

were scored as having severe RSV disease (Isolate 090, Isolate 104), one scored as moderate RSV disease

(Is. 101), and two scored as mild RSV disease (Is. 081, Is. 084). One subject with severe disease was

admitted to the intensive care unit. All subjects received oxygen therapy via nasal cannula.

Genome sequencing and comparative analysis of the five RSV isolates revealed that all belonged to RSV-

A and the GA2.3.5 G clade. Isolates from mild cases clustered in the A.D clade with moderate and severe

cases clustering in the A.D.1.5 and A.D.1.8 subclades (**Figure 1**). Comparative analysis of amino acid

polymorphisms in RSV between mild and moderate/severe disease isolates revealed multiple non-

synonymous substitutions. Amino acid differences at 19 positions were identified that distinguished all

mild isolates from all moderate/severe isolates (**Table 2**). Of the 19 identified, the majority were found to

be in the G protein (n=8) and the L polymerase (n=5). Others were found in the F protein (n=1), M

protein (n=1), P protein (n=1), M2-1 protein (n=1), and M2-2 (n=2). Many are associated with altered

physicochemical properties as shown in **Table 2**, which may contribute to differences in RSV disease

severity observed in hospitalized children.

In vitro growth kinetics of all RSV isolates in A549 cells demonstrated that isolates from moderate/severe

cases replicated more rapidly than those from mild cases; isolates from moderate/severe cases showed

significantly greater mean levels of extracellular viral titers than those from mild cases at 72 and 96-hours

post-infection (two-way ANOVA with Tukey's multiple-comparison test, $p<0.05$) (**Figure 2**).

Importantly, the phenotypic differences in viral replication in the present study were observed in a

controlled cell culture system, which reflects intrinsic viral properties because the influence of host-

related confounding variables was removed.

Given the evidence that isolates from moderate/severe RSV disease replicate to higher titers than those

from mild cases (**Figure 2)**, we employed predictive protein modeling to elucidate the structural

consequences of associated mutations. Structural modeling suggested that these mutations, particularly in

the L and G proteins, may enhance viral fitness by altering protein stability, dimerization, and host

interaction. All isolates encoded a 321-aa G protein. N-glycosylation prediction showed a stepwise

increase with severity: mild isolates had two sites, moderate/severe had three to four (**S1 Figure**). A

previous report demonstrated that glycosylation of the C-terminal domain of the G protein reduced

reactivity with specific antibodies, suggesting that various glycosylation patterns of RSV may contribute

to evasion of the host's humoral immune response (26).

L polymerase sequences revealed five amino acid differences between mild and moderate/severe isolates:

two in the RdRp domain (143, 179) and three within the connector region (1653, 1661, 1715) (**Table 2**).

Structural modeling based on Protein Data Bank 6UEN (27) highlighted these substitutions but limited

interpretation to the experimentally resolved RdRp and capping regions (**S2 Figure**). An additional

aspartate residue in the finger domain of the RSV L protein, specifically at position AA143, may have

significant implications for polymerase function. Introduction of an additional aspartate in this region

could alter the electrostatics of the template entry pore or influence RNA binding affinity (28,29). Such

changes may impact processivity or regulation of RNA synthesis, thereby affecting viral replication

efficiency and, consequently, clinical outcomes.

Reported associations between RSV viral load and disease severity in young children have been

inconsistent, likely due to variations in study design, analytical methods, host immune responses, and

intrinsic viral factors. Although our sample size is small, the compelling molecular findings presented

here highlight the potential for specific pathoadaptive mutations to influence evolving RSV virulence and

clinical outcomes. These results may help to explain previously reported inconsistencies and highlight the

need for larger studies, reverse genetics, and in vivo models to elucidate the specific mechanisms by

which RSV phenotype shapes disease severity.

Acknowledgments: We would like to express our immense gratitude to our colleagues Carlie DeFelice,

Dr. Kelly Hawley, Judy Kalinowski, Carson Karanian, Stephanie Lesmes, Dr. Bo Reese, and Jordan Wolf

for their vital contribution to the project, including participant recruitment, biospecimen collection,

processing and testing, and data capture.

References

1. Mazur NI, Caballero MT, Nunes MC. Severe respiratory syncytial virus infection in children: burden,
management, and emerging therapies. The Lancet. 2024 Sept 21;404(10458):1143–56.

2. Jain S, Williams DJ, Arnold SR, Ampofo K, Bramley AM, Reed C, et al. Community-acquired
pneumonia requiring hospitalization among U.S. children. N Engl J Med. 2015 Feb 26;372(9):835–
45.

3. Li Y, Wang X, Blau DM, Caballero MT, Feikin DR, Gill CJ, et al. Global, regional, and national
disease burden estimates of acute lower respiratory infections due to respiratory syncytial virus in
children younger than 5 years in 2019: a systematic analysis. The Lancet. 2022 May
28;399(10340):2047–64.

4. Andeweg S, Schepp R, van de Kassteele J, Mollema L, Berbers G, van Boven M. Population-based
serology reveals risk factors for RSV infection in children younger than 5 years. Sci Rep.
2021;11(1):8953.

5. Abrams EM, Doyon-Plourde P, Davis P, Brousseau N, Irwin A, Siu W, et al. Burden of disease of
respiratory syncytial virus in infants, young children and pregnant women and people. Can Commun
Dis Rep. 50(1–2):1–15.

6. Halasa N, Zambrano L, Amarin J, Stewart L, Newhams M, Levy E, et al. Infants admitted to US
intensive care units for RSV infection during the 2022 seasonal peak. JAMA Netw Open.
2023;6(8):e2328950.

7. Shishir TA, Saha O, Rajia S, Mondol SM, Masum MHU, Rahaman MM, et al. Genome-wide study of
globally distributed respiratory syncytial virus (RSV) strains implicates diversification utilizing
phylodynamics and mutational analysis. Sci Rep. 2023 Aug 19;13(1):13531.

8. Yu JM, Fu YH, Peng XL, Zheng YP, He JS. Genetic diversity and molecular evolution of human
respiratory syncytial virus A and B. Sci Rep. 2021 June 21;11(1):12941.

9. Bender W, Zhang Y, Corbett A, Chu C, Grier A, Wang L, et al. Association of disease severity and
genetic variation during primary Respiratory Syncytial Virus infections. BMC Medical Genomics.
2024 June 19;17(1):165.

10. Fletcher J, Smyth R, Thomas H, Ashby D, Hart C. Respiratory syncytial virus genotypes and disease
severity among children in hospital. Arch Dis Child. 1997 Dec;77(6):508–11.

11. Human S, Hotard A, Rostad C, Lee S, McCormick L, Larkin E, et al. A respiratory syncytial virus
attachment gene variant associated with more severe disease in infants decreases fusion protein
expression, which may facilitate immune evasion. J Virol. 2020;95(2).

12. Li W, Wang Y, Yu B, Tan Q, Zhou J, Hu J, et al. Disease severity of respiratory syncytial virus (RSV)
infection correlate to a novel set of five amino acid substitutions in the RSV attachment glycoprotein
(G) in China. Virus Research. 2020 May 1;281:197937.

13. Midulla F, Nenna R, Scagnolari C, Petrarca L, Frassanito A, Viscido A, et al. How Respiratory
Syncytial Virus Genotypes Influence the Clinical Course in Infants Hospitalized for Bronchiolitis.
The Journal of Infectious Diseases. 2019 Jan 29;219(4):526–34.

14. Comas-García A, Noyola DE, Cadena-Mota S, Rico-Hernández M, Bernal-Silva S. Respiratory
Syncytial Virus-A ON1 Genotype Emergence in Central Mexico in 2009 and Evidence of Multiple
Duplication Events. J Infect Dis. 2018 Mar 13;217(7):1089–98.

15. Esposito S, Piralla A, Zampiero A, Bianchini S, Di Pietro G, Scala A, et al. Characteristics and Their
Clinical Relevance of Respiratory Syncytial Virus Types and Genotypes Circulating in Northern Italy
in Five Consecutive Winter Seasons. PLoS One. 2015;10(6):e0129369.

16. Oraby AK, Stojic A, Elawar F, Bilawchuk LM, McClelland RD, Erwin K, et al. A single amino acid
mutation alters multiple neutralization epitopes in the respiratory syncytial virus fusion glycoprotein.
npj Viruses. 2025 Apr 22;3(1):1–14.

17. Bøås H, Havdal LB, Størdal K, Døllner H, Leegaard TM, Bekkevold T, et al. No association between
disease severity and respiratory syncytial virus subtypes RSV-A and RSV-B in hospitalized young
children in Norway. PLoS One. 2024;19(3):e0298104.

18. Lin GL, Drysdale SB, Snape MD, O'Connor D, Brown A, MacIntyre-Cockett G, et al. Targeted
metagenomics reveals association between severity and pathogen co-detection in infants with
respiratory syncytial virus. Nat Commun. 2024 Mar 16;15(1):2379.

19. Anderson LJ, Peret TC, Piedra PA. RSV Strains and Disease Severity. The Journal of Infectious
Diseases. 2019 Jan 29;219(4):514–6.

20. Vandini S, Biagi C, Lanari M. Respiratory syncytial virus: The influence of serotype and genotype
variability on clinical course of infection. Int J Mol Sci. 2017;18(8).

21. CDC. RESP-NET. 2025 [cited 2025 Nov 18]. Respiratory Virus Hospitalization Surveillance
Network (RESP-NET). Available from: https://www.cdc.gov/resp-net/dashboard/index.html

22. Mejias A, Dimo B, Suarez NM, Garcia C, Suarez-Arrabal MC, Jartti T, et al. Whole blood gene
expression profiles to assess pathogenesis and disease severity in infants with respiratory syncytial
virus infection. PLoS Med. 2013 Nov;10(11):e1001549.

23. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome Biology.
2019 Nov 28;20(1):257.

24. Wang S, Sundaram JP, Stockwell TB. VIGOR extended to annotate genomes for additional 12
different viruses. Nucleic Acids Res. 2012 July;40(Web Server issue):W186–92.

25. Aksamentov I, Roemer C, Hodcroft EB, Neher RA. Nextclade: clade assignment, mutation calling
and quality control for viral genomes. Journal of Open Source Software. 2021 Nov 30;6(67):3773.

26. Rawling J, Melero JA. The Use of Monoclonal Antibodies and Lectins to Identify Changes in Viral
Glycoproteins That are Influenced by Glycosylation. In: Sugrue RJ, editor. Glycovirology Protocols
[Internet]. Totowa, NJ: Humana Press; 2007 [cited 2025 Nov 18]. p. 109–25. Available from:
https://doi.org/10.1007/978-1-59745-393-6_8

27. Bank RPD. RCSB PDB - 6UEN: Cryo-EM structure of the respiratory syncytial virus RNA
polymerase [Internet]. [cited 2025 Nov 18]. Available from: https://www.rcsb.org/structure/6UEN

28. Sutto-Ortiz P, Eléouët JF, Ferron F, Decroly E. Biochemistry of the Respiratory Syncytial Virus L
Protein Embedding RNA Polymerase and Capping Activities. Viruses. 2023 Jan 25;15(2):341.

29. Liang B. Structures of the Mononegavirales Polymerases. J Virol. 2020 Oct 27;94(22):e00175-20.

30. Gupta R, Brunak S. Prediction of glycosylation across the human proteome and the correlation to
protein function. Pac Symp Biocomput. 2002;310–22.

31. Steentoft C, Vakhrushev SY, Joshi HJ, Kong Y, Vester-Christensen MB, Schjoldager KTBG, et al.
Precision mapping of the human O-GalNAc glycoproteome through SimpleCell technology. EMBO
J. 2013 May 15;32(10):1478–88.

Table 1. Cohort Characteristics

Characteristic	RSV Cohort (n=5)
Enrollment Age (mos)	9 (3-9)
Male	4 (80%)
Race	
White	2 (40%)
Black	1 (20%)
Other	2 (40%)
Hispanic	3 (60%)
Length of Stay (days)	2 (1-3)
CDSS Category	
Mild	2 (40%)
Moderate	1 (20%)
Severe	2 (40%)

Data given as median (IQR) or n (%)

Table 2. Amino Acid Differences and side chain characteristics distinguishing Mild from Moderate/Severe

Protein	AA #		Mild (Is. 081, Is. 084)		Moderate/severe (Is. 090, Is. 101, Is. 104)
L	143	N	HYDROPHILIC	D	NEGATIVE
L	179	T	HYDROPHILIC	S	HYDROPHILIC
L	1653	I	HYDROPHILIC	V	HYDROPHILIC
L	1661	K	POSITIVE	N	HYDROPHILIC
L	1715	S	HYDROPHILIC	N	HYDROPHILIC
G	105	T	HYDROPHILIC	S	HYDROPHILIC
G	123	K	POSITIVE	E	NEGATIVE
G	142	L	HYDROPHOBIC	S	HYDROPHILIC
G	281	K	POSITIVE	T	HYDROPHILIC
G	283	F	AROMATIC	S	HYDROPHILIC
G	294	S	HYDROPHILIC	P	PROLINE
G	309	H	POSITIVE	Y	HYDROPHOBIC
G	310	P	PROLINE	L	HYDROPHOBIC
M2-1	172	A	HYDROPHOBIC	T	HYDROPHILIC
M2-2	9	E	NEGATIVE	D	NEGATIVE
M2-2	49	T	HYDROPHILIC	P	PROLINE
F	555	A	HYDROPHOBIC	T	HYDROPHILIC
M	43	M	HYDROPHOBIC	I	HYDROPHOBIC
P	62	T	HYDROPHILIC	I	HYDROPHOBIC

Figure 1. Phylogenetic analysis of RSV clinical isolates. CDSS, Clinical Disease Severity Score

A

B

24 HPI

24 HPI		Is. 090 (Severe)	Is. 101 (Moderate)	Is. 104 (Severe)
	Is. 081 (Mild)	ns	ns	ns
	Is. 084 (Mild)	ns	ns	ns

48 HPI

48 HPI		Is. 090 (Severe)	Is. 101 (Moderate)	Is. 104 (Severe)
	Is. 081 (Mild)	ns	ns	ns
	Is. 084 (Mild)	ns	ns	ns

72 HPI

72 HPI		Is. 090 (Severe)	Is. 101 (Moderate)	Is. 104 (Severe)
	Is. 081 (Mild)	*	**	**
	Is. 084 (Mild)	*	*	*

96 HPI

96 HPI		Is. 090 (Severe)	Is. 101 (Moderate)	Is. 104 (Severe)
	Is. 081 (Mild)	*	ns	ns
	Is. 084 (Mild)	****	ns	ns

Figure 2. Temporal analysis of RSV isolate replication **A)** A549 cells were infected with RSV isolates at an MOI of 0.01. Extracellular RSV was isolated, and viral titer was determined by $TCID_{50}$. Samples were collected at 24H intervals over 96 hours. Data shown represent the mean of duplicate measurements +/- SD **B)** Table summarizing the statistical significance of differences in viral titer between mild, moderate, and severe strains at each timepoint. ns (not significant), * ($p < 0.05$), ** ($p < 0.01$), *** ($p < 0.001$), **** ($p < 0.0001$). Abbreviations: $TCID_{50}$, 50% tissue culture infectious dose; MOI, multiplicity of infection; Is, Isolate; HPI, Hours post-infection.

Supplemental Information

S1 Table. Clinical Disease Severity Score (CDSS)

	0	1	2	3
O₂ Saturation	>95% in RA	92%-95% in RA	<92% in RA or suppl O₂ ≤35%	Intubated/CPAP or suppl O2>35%
Respiratory Rate	≤6 m <40/min >6 m <30/min	≤6 m 41-55/min >6 m 31-45/min	≤6 m 56-70 >6 m 46-60	≤6 m >70 >6 m >60
Retractions	No	Subcostal +	Subcostal ++ or intercostal	Subcostal +++ or intercostal and suprasternal
Auscultation	Normal	Crackles/ ronchi	Any wheezing	Exp/insp wheezing without stethoscope or no breath sounds heard
Activity	Normal, alert	Abnormal, fussy, or decreased oral intake	Unable to maintain hydration (IV fluids or NG feeds)	Intubated

[a]Score from 0-15: Mild (0-5), Moderate (6-9), Severe (≥10).
[b]CPAP, continuous positive airway pressure; exp, expiratory; insp, inspiratory; IV, intravenous; m, months; min, minute; NG, nasogastric tube; O₂, oxygen; RA, room air; suppl, supplemental; +, mild; ++, moderate; +++, severe.

S1 Figure. Predicted N-glycosylation sites in RSV Attachment G protein A) in isolates from cases of mild disease (Is. 081, Is. 084), B) in isolates from moderate cases of disease (Is. 101) and C) in isolates from severe cases of disease (Is. 090, Is. 104). N-glycosylation potential threshold = 0.5 (purple line). Potential is the predicted likelihood of glycosylation (green). Jury is the number of prediction tools in agreement out of 9.

S2 Figure. Structural model of RSV L proteins of representative isolates from mild and moderate/severe cases using homology modelling based on the RSV polymerase complex structure (PDB: 6UEN). Ribbon representation of L proteins are shown on left (yellow, pathoadaptive mutations; blue, RNA-dependent RNA polymerase domain; grey, PRNTase capping domain). Ramachandran plots are shown for each model (center) followed by quality parameters of each model. T, Threonine; N, Asparagine; S, Serine; D, Aspartate: