Slow-growing human cell lines are a suitable alternative to rabbit Sf1Ep cells for *in vitro*
cultivation of *Treponema pallidum*

Katia Capuccini[1], Divenita Govender[1], David Goulding[1], Cecilia Kyanya[1], Shivani Pasricha[2], Lorenzo
Giacani[3, 4], Nichola R Thomson*[1, 5], Linda Grillová*[1#]

[1]Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, UK.

[2] Department of Infectious Diseases, The University of Melbourne at the Peter Doherty Institute for Infection
and Immunity, Melbourne, Australia

[3]Department of Medicine, Division of Allergy and Infectious Diseases, University of Washington, Seattle (WA),
USA

[4]Department of Global Health, University of Washington, Seattle, WA, 98104, USA

[5]London School of Hygiene & Tropical Medicine, London, UK

*These authors contributed equally and share senior authorship

Corresponding author, correspondence to lg22@sanger. ac.uk; lindagrillova@gmail.com

Abstract

Treponema pallidum subsp. *pallidum*, the causative agent of syphilis, remains difficult to study due to long-standing limitations in *in vitro* cultivation. Although a rabbit epithelial cell co-culture system (Sf1Ep) enabled major advances in recent years, the lack of human cell-based models restricts clinical relevance and mechanistic insight into host–pathogen interactions. Here, we sought to establish human epithelial cell lines capable of supporting *T. pallidum* growth *in vitro*. Six human epithelial or epithelial-like cell lines from diverse tissue origins were evaluated under microaerophilic conditions using the standard *T. pallidum* cultivation medium. Among these, CAL-39 (vulva) and HepG2 (liver) supported *T. pallidum* survival, replication, characteristic growth behaviours, and long-term passage at levels comparable to Sf1Ep cells. Growth kinetics, attachment dynamics, and motility of *T. pallidum* were quantified over extended culture periods. Using live-cell imaging, we define for the first time two distinct *T. pallidum* host-cell interaction behaviors—surface-associated crawling and stable single-polar attachment. Both behaviours were observed across all tested host cell lines and persisted over time only in cell lines permissive for sustained growth. Together, our findings establish clinically relevant human epithelial co-culture models for *T. pallidum*, provide new insights into host-cell-dependent growth and motility, and create a platform for future mechanistic studies of syphilis pathogenesis and vaccine target discovery.

Introduction

Syphilis, caused by *Treponema pallidum* subsp. *pallidum* (*T. pallidum*), typically begins with a localized chancre at the site of infection, and rapidly disseminates in multiple tissues and organs, leading to a wide spectrum of clinical manifestations across the primary, secondary, and tertiary stages. If left untreated, syphilis can result in severe and irreversible neurological and cardiovascular complications and remains a leading cause of adverse pregnancy outcomes worldwide, including stillbirths and neonatal deaths when congenital infection occurs (1). Although syphilis is readily curable with appropriate antibiotic therapy, its incidence has increased substantially over the past two decades, particularly among specific communities, including men who have sex with men. The WHO estimates approximately 8 million new syphilis infections worldwide in 2022, compared to 7.1 million cases in 2020 (2). Together, the rising incidence and burden of preventable morbidity and mortality highlight the urgent need to strengthen syphilis prevention, screening, and control strategies through a deeper understanding of base biology as well as testing.

While the development of an effective vaccine would significantly help control the spread of syphilis, there is currently little consensus on target selection. Importantly, while there is a strong focus on syphilis vaccine development, progress is hindered by a lack of understanding of the pathogenesis and basic biology of *T. pallidum*. For decades investigators have been unable to steadily propagate this

bacterium *in vitro,* needing to rely on *in vivo* propagation in the rabbit infection model. However, this
changed in 2018, when a system for successful long-term *in vitro* co-cultivation of *T. pallidum* with
rabbit skin epithelial cells (Sf1Ep) was established (3). This breakthrough enabled accelerated progress
in the field, with significant advancements represented by methods for genetic modification of *T.*
pallidum (4), high-throughput antibiotic sensitivity/resistance testing (5), and proteomics and
transcriptomic studies (6, 7), among other tools and procedures that are standard in traditional
microbiology for different bacteria.

The existing Sf1Ep co-culture system is valuable for propagating *T. pallidum* and has provided
an accessible model of early host-pathogen interactions at epithelial surfaces, where the bacterium first
encounters the human host. However, because *T. pallidum* disseminates widely in the host and
interactions with the pathogen are likely to vary between cells from different anatomical sites, there is
a need for infection models that use human rather than rabbit cells. Accordingly, the goal of this study
was to establish more clinically relevant *in vitro* infection models by identifying suitable human cell
lines to replace rabbit epithelial Sf1Ep cells as foundation for future studies and mechanistic work.

Here, we identified two human epithelial cell lines from distinct body sites—vulva and liver—
that support sustained growth and the complete growth cycle of *T. pallidum* in a manner comparable to
Sf1Ep cells. For the first time, using these more clinically relevant novel co-culture models in
combination advance microscopy platforms, we describe a duality in *T. pallidum* behaviour when
interacting with human and rabbit epithelia, observed using both live imaging of replicating cultures,
as well as in in fixed samples. It is anticipated that identifying human cell models that support *T.*
pallidum growth will enable a new generation of clinically grounded, mechanistic discoveries,
accelerating vaccine target discovery and closing critical knowledge gaps in *T. pallidum*-host
interactions, ultimately transforming control of syphilis—a disease that remains poorly understood
despite its long history.

Results

Selection of human cell lines

Currently, *T. pallidum* is cultured using Sf1Ep rabbit epithelial cells under low-oxygen
conditions (1.5% O_2) in *T. pallidum* cultivation medium (TPCM-2) for 7 days, reflecting its long
generation time (>35 h)(3). To identify human alternatives to rabbit cells, we selected six well-
characterized human epithelial or epithelial-like cell lines prioritizing hypoxia-tolerant cell lines with
longer doubling times (>30 h), replicating key characteristics of Sf1Ep cells, while including a faster-
growing cell line (HEK-293, <20 h) as a comparator (Table 1). The selected cell lines span diverse
tissue origins: vulva (CAL-39, SW954), placenta (JAR, JEG-3), liver (HepG2), and kidney (HEK-293).
All cell lines were first assessed for growth in 1.5% O_2 and in TPCM-2 over a 7-day period, replicating

the established Sf1Ep co-culture workflow (3). While some cell lines exhibited longer doubling times

under these conditions compared to optimal normoxic conditions typical of human cell culture (Table

1), none showed morphological abnormalities or signs of significant cell damage when cultured under

low oxygen (see Method section for details).

Testing human cell lines for T. pallidum in vitro growth support

We first assessed *T. pallidum* growth using SS14 strain across the six selected human cell lines

and compared these to *T. pallidum* cultured without mammalian cells (negative control, NC) or

cocultured with the reference supporting cell line Sf1Ep (positive control, PC), over a 4-week period

(Fig. 1A). At each weekly passage, cultures were reseeded with a fixed inoculum of 2.5×10^6 *T.*

pallidum cells that were attached on to the host cell monolayer. We have demonstrated that both *T.*

pallidum cells attached to the surface of mammalian cells and non-attached cells freely floating in the

medium are viable under standard culture conditions. This was shown by re-seeding experiments in

which *T. pallidum* cells collected exclusively from the culture medium re-attached to fresh monolayers

and completed a full growth cycle (Supplementary material). Therefore, both attached and non-attached

T. pallidum populations were considered throughout the characterization of growth dynamics in this

study. The only exception was this specific experiment, in which—apart from the negative control—

only monolayer-attached *T. pallidum* cells were used to estimate growth kinetics, in accordance with

standard *T. pallidum* passage recommendations. (8). For the HEK-293 and JAR cell lines, *T. pallidum*

cell numbers fell below this threshold during the first and third passages, respectively, and the cultures

were terminated after weeks 1 and 3, respectively. As expected, *T. pallidum* negative control cultures

showed markedly reduced total *T. pallidum* yields over time. Among the six candidate cell lines tested,

CAL-39 and HEPG2 displayed *T. pallidum* growth profiles that were comparable to the Sf1Ep standard

(Fig.1A). Comparison of total *T. pallidum* counts showed no significant difference in total *T. pallidum*

counts between *T. pallidum* co-cultured with CAL-39 or HepG2 and *T. pallidum* co-cultured with Sf1Ep

($p > 0.05$). In contrast, *T. pallidum* co-cultured with either CAL-39 or HepG2 cells showed significantly

higher total *T. pallidum* counts compared to *T. pallidum* NC ($p < 0.05$; see Fig. 1B for p-values).

Across all time points, *T. pallidum* motility remained high (~90%) in CAL-39, HEPG2, and

Sf1Ep, but was significantly reduced in the other tested cell lines (Fig. 1C; $p = 0.0055$). Motility did

not differ significantly among CAL-39, HEPG2, and Sf1Ep. All cell lines increased motility relative to

the no-cell control, reaching statistical significance for HEPG2 and Sf1Ep, while CAL-39 showed a

consistent but non-significant increase, supporting its suitability as a co-culture model.

Relationship between mammalian cells division time and T. pallidum yield

To explore the relationship between human cell division time and *T. pallidum* yield, *T. pallidum*

fold change was plotted against the division time of each human cell line, with Sf1Ep cells included as

a reference (Fig. 1D). *T. pallidum* yield, expressed as $\log_{10}$ fold change, increased with longer host cell

division times in both human and rabbit cells (slope = 0.034 $\log_{10}$ units per hour). This relationship

explained 37% of the variance but did not reach statistical significance (R^2 = 0.37, p = 0.147).

Longitudinal T. pallidum growth dynamics in *in vitro* co-culture

To characterize *T. pallidum* growth across the different host cell lines and to determine the

timing of exponential growth of *T. pallidum*, cultures were monitored daily for 10-days. Each 24-hour

time point was analysed using a dedicated culture plate, with all plates inoculated simultaneously on

day 0 with the same starting inoculum; at each time point a single plate was terminated and processed

for *T. pallidum* quantification. In this experiment, growth was estimated from the host cell attached as

well as the non-attached *T. pallidum* cells floating in the culture medium to provide a more detailed

view of *T. pallidum* growth dynamics.

Cell lines Sf1Ep, CAL-39, and HepG2 all supported similar *T. pallidum* growth trajectories,

with comparable increases in total *T. pallidum* cell count over the 10-day period (Fig. 2A). The relative

proportions of attached and floating *T. pallidum* populations in these cultures are shown in (Fig. 2B).

After day 9 of culture, the medium in CAL-39 and HepG2 co-cultures became acidified, as indicated

by a colour change to yellow, and *T. pallidum* cell numbers in both attached and floating fractions began

to decline. Intact and motile treponemes were still detectable by microscopy, but at lower abundance,

suggesting that cell lysis may contribute to the reduced counts under these conditions. In contrast, no

comparable decline was observed in Sf1Ep co-cultures. Therefore, currently culturing *T. pallidum* with

these human cell lines would not be recommended beyond 9 days, while the standard 7-day culture

period is recommended for consistency and convenience.

In parallel, *T. pallidum* growth was monitored in the non-supporting cell lines HEK-293,

SW954, JAR, and JEG-3 at days 1, 3, 5, and 7 (Fig. 2C). Total *T. pallidum* counts in these non-

supporting cell lines were substantially lower than those observed in the supporting cell lines, with a

maximum of 3.5×10^6 cells. Moreover, a higher proportion of *T. pallidum* remained in the floating

fraction when co-culturing with non-supporting cell lines, consistent with reduced attachment to the

host-cell monolayer (Fig. 2D). Wilcoxon rank-sum tests conducted independently for each day yielded

p-values of 0.057 (day 1), 0.057 (day 2), 0.114 (day 3), 0.229 (day 5), and 0.229 (day 7). Corresponding

boxplots and statistical comparisons are presented in (Supplementary Fig. 1).

Real time interaction of T. pallidum with host cell lines

To visualize *T. pallidum* interactions with host cells in real time, we established the first live-cell imaging platform for *T. pallidum*, enabling continuous monitoring of bacterial behaviour during co-culture. This approach was first used to characterise host–pathogen interactions in the standard Sf1Ep co-culture system. Two distinct populations of *T. pallidum* interacting differently with host cells were identified. First, *T. pallidum* exhibited highly dynamic, surface-associated motility characterized by close contact with host cell membranes and movement along the cellular surface. This behaviour resembles the "crawling" movement, or "translocation via an undulating waveform," previously described for other human pathogenic spirochetes such as *Leptospira* and *Borrelia* (Video file, Fig. 3A). *T. pallidum* cells displayed back-and-forth (reversal) movement using alternating poles and, in some cases, turned in the opposite direction using the same pole, likely due to transient self-entanglement (Video file, Fig. 3B). Second, a distinct population of *T. pallidum* cells exhibited stable single-polar attachment to the host cell surface, with the opposite pole remaining free (Video file, Fig. 3C).

Next, real-time imaging of *T. pallidum* interacting with all tested host cell lines was performed, and these interactions were monitored over a 10-day period to complement the full picture of *T. pallidum* growth dynamics. Interestingly, the characteristic behaviours described in this study, represented by two motility types—crawling-like and single–pole–attached—, were observed in all host cell lines tested during the first 3 days (including rabbit Sf1Ep cells and human Cal-39, HepG2, JEG-3, JAR, HEK-293, and SW954 cells). In Sf1Ep, Cal-39, and HepG2 cells, both motility types persisted throughout the full 10 days of observation. In contrast, in human cell lines described as non-supportive for *T. pallidum* growth (JEG-3, JAR, HEK-293, and SW954), *T. pallidum* cells exhibiting crawling-like behaviour gradually disappeared; overall, cells became increasingly non-motile, with the only remaining motile cells showing single-polar attachment.

Discussion

Since 2018, *T. pallidum* is cultured *in vitro* using Sf1Ep rabbit epithelial cells in *T. pallidum* cultivation medium (TPCM-2; media (3)) and under low-oxygen conditions (1,5% O_2) for 7 days, a duration required to allow *T. pallidum* multiplication due to its long generation time (more than 35 hours). Over the past several decades, numerous culture media formulations and multiple cell lines were tested, but only the combination of TPCM-2 and Sf1Ep cells proved effective (3). Historically tested human, rabbit, and rat cell types included those from testis, kidney, spleen, lung, epidermis, cervix, urethra, and nerve tissues. Although *T. pallidum* showed affinity for these cells, it did not survive long term, persisting for no more than a few days. (9). Importantly, the human cell lines tested were predominantly fast-growing (with doubling time less than 20 hours in optimal conditions), in contrast to the much slower-growing Sf1Ep cells (with doubling time more than 30 hours in optimal conditions). Moreover, using higher passage (>60) Sf1Ep cells in the co-culture system results in a decreased treponemal yield and motility (8). This is likely because high-passage Sf1Ep cells tend to divide more rapidly. Similarly, starting the culture with more than 10^5 Sf1Ep cells per well of a 6-well plate have

been found to result in lower culture yield and decreased treponemal motility (8). Therefore, a high density of Sf1Ep cells appears to negatively affect treponemal growth, likely due to competition between Sf1Ep and treponemes for shared nutrients, accumulation of secondary metabolites, and consequent acidification of the culture medium.

Given the above, we hypothesized that slow-growing human cell lines with longer generation times may have greater potential to support a slow-growing bacterium such as *T. pallidum*. Hence, our human cell line selection was guided by the premise that reduced host-cell proliferation may limit nutrient competition and media acidification, and that adaptation to low-oxygen environments may make certain human cell lines more permissive under the hypoxic conditions (1.5% O_2) required for *T. pallidum* cultivation. Among the six host cell lines tested, CAL-39 and HepG2 supported *T. pallidum* growth at levels comparable to the reference Sf1Ep cell line. In contrast, but consistent with previous studies, *T. pallidum* cultured without any eukaryotic cell showed markedly reduced total cell counts over time, consistent with a stress condition leading to progressive cell loss (10). A similar outcome was observed with non-supporting cell lines, in which *T. pallidum* cell numbers frequently declined below the threshold required for continued passaging, indicating that a specific host–cell interaction is necessary to sustain *T. pallidum* growth *in vitro*.

The establishment of a live-cell imaging platform enabled, for the first time, direct visualization of *T. pallidum* interactions with host cells under physiologically relevant, microaerophilic conditions. It showed that T. pallidum can initially colonize all tested cell lines, regardless of their ability to support long-term growth, at least during the first 3 days of co-culture. This observation is consistent with extensive clinical and historical evidence demonstrating dissemination of *T. pallidum* to a wide range of human tissues in atypical and systemic syphilis, including bone (11), gastric mucosa (12), liver (13), extragenital derma (14), lungs (15), thyroid gland (16), eyes (17) and even multiple sites simultaneously (18). However, *in vitro, T. pallidum* is likely to engage in distinct molecular interactions with host cells that support sustained bacterial growth compared with cells that do not permit long-term culture or completion of the growth cycle. This is reflected in the distinct growth dynamics observed between supporting and non-supporting cell lines, particularly in the differing proportions of bacteria interacting directly with the monolayer versus those remaining unattached in the culture medium (Supplementary Figure 1). Consequently, human monolayers that enable long-term maintenance of *T. pallidum* may represent a more appropriate model for studying *T. pallidum*–host interactions than cell lines used only for short-term exposure. Our findings also give light to the fact that successful long-term *in vitro* cultivation appears to be constrained by the interplay between host cell growth and that of the infecting bacterium with a clear positive trend toward increased *T. pallidum* yield with longer host cell division times, albeit in a limited number of cell lines. In addition, live-cell imaging also demonstrated that *T. pallidum* is capable not only of polar attachment but also of active surface crawling, offering a new

perspective on mechanisms underlying tissue invasion and expanding our understanding of *T. pallidum*

motility and early host interactions.

Despite our co-culture experiments being limited to four weeks in duration, at the time of

writing, we have maintained *T. pallidum* in CAL-39 and HepG2 for 11 and 6 weeks, respectively,

suggesting longer-term co-cultivation with human cell lines is feasible and will be important for

dissecting additional aspects of *T. pallidum* stability and host-cell adaptation. While cell systems cannot

fully recapitulate the complexity of infection in the human body, this work represents an important step

toward replacing rabbit Sf1Ep cells in the existing cultivation platform for *T. pallidum*, with the

potential to increase clinical relevance and enable mechanistic studies of human–pathogen interactions

central to syphilis pathogenesis, particularly through long-term culture approaches that may ultimately

capture stages of the bacterial growth cycle that remain unexplored. Our discovery will likely enable

and guide additional research laboratories that have these and other similar human cell lines already

available to adopt the *in vitro* cultivation system for *T. pallidum*.

Methods

Human cell lines

Human cell lines to be tested were selected from the Cell Model Passports database (https://cellmodelpassports.sanger.ac.uk/), which contains more

than 2000 well-characterized cell lines from diverse tissues. Lines with known tolerance for hypoxia tolerance and slow division rate were prioritized, while

the faster-growing HEK-293 cell line was included as a reference (Table 1). Cell lines were also selected to represent anatomically relevant tissue sites

associated with *T. pallidum* infection and systemic dissemination.

Table 1: Human cell lines tested for *T. pallidum* growth support.

Cell line	Repository ID	Tissue of origin	Cancer / origin	Doubling time (optimal conditions.)	Hypoxia tolerance	Doubling time (low O_2 and TPCM-2)
CAL-39	DSMZ ACC 187	Vulva	Squamous cell carcinoma	40–60 h	Moderate	50 h
SW954	ATCC HTB-117™	Vulva	Squamous cell carcinoma	~50 h	Moderate	52 h
JAR	ATCC HTB-144™	Placenta	Choriocarcinoma	36–48 h	High	39 h
JEG-3	ATCC HTB-36™	Placenta	Choriocarcinoma	48–72 h	High	37 h
HepG2	ATCC HB-8065™	Liver	Hepatocellular carcinoma	~48 h	Moderate	46 h
HEK-293	ATCC CRL-1573™	Embryonic kidney	Adenovirus-transformed	24–30 h	Moderate	40 h

 Culture conditions and T. pallidum strain

CAL-39, SW954, JEG-3, HepG2, and HEK-293 cells were cultured in filtered, sterile
DMEM/F-12 (Gibco 31330-038, ThermoFisher Scientific, UK) supplemented with 10% FBS. JAR cells
were cultured in RPMI (Gibco 52400-025, ThermoFisher Scientific, UK) supplemented with 10% FBS
and 4.5 mg/mL glucose. Sf1Ep cells were maintained in medium as described by Edmondson et al.,
2018 (3). All cell lines were passaged at confluence (every 3–7 days), and media were changed every
3–5 days depending on cell line and degree of medium acidification. For experimental wells used for *T.*
pallidum culture, 50,000 cells were seeded per well in 6-well plates one day prior to inoculation. The
T. pallidum SS14 strain engineered to express GFP (19) was used for all experiments and was grown *in*
vitro as previously described (3) either with standard Sf1Ep cells or with the indicated human cell lines
(Table 1). The growth of *T. pallidum* was monitored by dark-field microscopy by assessing both cell
counts and motility. *T. pallidum* cell counts were quantified at multiple time points, and mean values
with associated variability were plotted over time. Monitoring was performed either at 7-day intervals,
as recommended, or daily to generate *T. pallidum* growth curves, with each 24-hour time point analysed
from a separate plate that was inoculated simultaneously with an identical inoculum. Growth was
assessed in both *T. pallidum* cells attached to the monolayer, following standard culture evaluation
procedures (3), and in non-attached, floating cells present in the culture medium. For analysis of floating
cells, 4 mL of culture medium were centrifuged at $10,000 \times g$ for 20 min to pellet *T. pallidum* cells, and
the pellet was resuspended in 1 mL of TPCM-2 prior to microscopy. While high-speed centrifugation
is likely to disrupt the fragile membrane of *T. pallidum* and reduce cell viability, this was not critical in
this context, as the goal was solely to count cells and they were not intended for further sub-passaging.

Statistical analysis

All statistical analyses and figure generation were performed in R (version 4.5.2). Total *T.*
pallidum counts in different cell lines were compared using one-way ANOVA with Tukey's HSD post
hoc test. Temporal *T. pallidum* growth across all cell lines was analysed with two-way ANOVA for the
main effects of cell line, day, and their interaction. When significant differences were found, a linear
model was used to determine pairwise comparison for each day p-values. *T. pallidum* motility was
compared using the Kruskal–Wallis's test, with Dunn's post hoc test and Benjamini–Hochberg
correction for pairwise comparisons. The relationship between mammalian cell division time and *T.*
pallidum yield ($\log_{10}$ fold change) was assessed by linear regression. *T. pallidum* floating in the media
were compared between supporting and non-supporting cell lines using Wilcoxon rank-sum tests
performed separately for each day.

Each measurement was performed using three biological replicates. Although this limited
statistical power for some comparisons, the number of replicates was dictated by experimental
complexity and the need to complete treponemal counting and motility assessments within a narrow

time window. Increasing the number of replicates could have been detrimental to data reliability, as

treponemes do not maintain viability indefinitely after harvest.

Live cell imaging

For live-cell imaging, a Leica THUNDER Imager (Leica Microsystems, Milton Keynes, UK)

was used, selected for its computational background removal and low phototoxicity. To maintain

appropriate culture conditions during imaging, a cage incubator, an on-stage gas-tight chamber, and gas

and temperature controllers (Okolab, Rovereto, Italy) were employed. These were connected to a

premixed gas cylinder (1.5% O_2, 5% CO_2, balance N_2; BOC, Cambridge, UK) to provide a

microaerophilic atmosphere suitable for *T. pallidum* growth and to maintain the temperature at 34 °C.

T. pallidum was co-cultured with different cell lines in microscopy dishes compatible with the gas-tight

chamber (Nunc™ Glass Bottom Dishes, cat. no. 150680, Thermo Fisher Scientific, UK). A 63× oil-

immersion objective was used to monitor *T. pallidum* motility and interactions.

Scanning and transmission electron microscopy (SEM and TEM)

T. pallidum was cultured under standard conditions for 7 days in the presence of monolayers of

various mammalian cells grown on sterile circular coverslips (12 mm; Paul Marienfeld GmbH & Co.

KG, Lauda-Königshofen, Germany). Coverslips were placed one per well in 24-well plates and used

for SEM or TEM sample preparation. Immediately after removal of the plates from the incubator, cells

were inactivated and fixed by exchanging the culture medium for 2% glutaraldehyde and 4%

paraformaldehyde (PFA) in 0.1M sodium cacodylate (pH 7.42) for 2 hours (all steps at room

temperature). They were then rinsed in cacodylate buffer (3 x 10 minutes), post-fixed in cacodylate

buffered 1% osmium tetroxide for a further 2 hours, rinsed in buffer again and mordanted in either 1%

buffered tannic acid (TEM) or 1% thiocarbohydrazide (SEM) for 1 hour. Then, for TEM, cells were

washed in sodium sulfate for 10 minutes, rinsed 6 x 2 minutes in distilled water, dehydrated in an ethanol

series, stained *en bloc* in 1% uranyl acetate at the 20% stage, embedded using an Epoxy resin embedding

kit (Epoxy embedding medium 45345, Sigma-Aldrich – Merck, Gilingham, UK), and cured in an oven

at 60°C for 48 hours. Areas containing cells were cut from the well and ultrathin-sectioned at 60nm on

a Leica UC6 ultramicrotome. Sections were collected onto EM grids, contrasted again with uranyl

acetate and lead citrate and analysed on a 120kV Tecnai Spirit Biotwin TEM with a Tietz F4.16 CCD.

For SEM, samples were further impregnated with osmium tetroxide for 1 hour, rinsed in buffer for 3 x

10 minutes and distilled water for 6 x 2 minutes, dehydrated in an ethanol series, the coverslips removed

and critical point dried in a Leica EM CPD300, mounted onto SEM stubs with conducting paste, sputter-

coated with 5nm of gold in a Leica ACE600 sputter-coater and analysed on a Hitachi SU8030 SEM.

Figure 1. Human epithelial cell line–dependent growth, motility, and yield of *T. pallidum* in long-term co-culture. (A) *T. pallidum* growth in the presence of different human cell lines over 28 days. Connected scatterplots depict mean *T. pallidum* cell counts from three biological replicates, measured at weekly harvests, with lines connecting mean yields across passages. For each weekly passage, a standardized treponemal inoculum (2.5×10^6 cells) was used to subculture into wells freshly seeded with human cells. For HEK-293 (green) and JAR (teal) cell lines, cultures were terminated early due to insufficient *T. pallidum* inoculum ($<2.5 \times 10^6$ cells) at passage 1 and passage 3, respectively. NC, negative control (*T. pallidum* cultured without host cells); PC, positive control (*T. pallidum* cultured with rabbit epithelial Sf1Ep cells). (B) Triangular heatmap of pairwise TukeyHSD post hoc comparisons of *T. pallidum* counts between cell lines. Each cell line shows the adjusted p-value for a given pairwise comparison; blue tiles indicate statistically significant differences ($p < 0.05$). (C) Percentage of motile *T. pallidum* cells across co-cultures over 4 weeks. Points represent mean values from biological replicates. (D) Relationship between mammalian cell division time and *T. pallidum* yield. Scatterplot shows $\log_{10}$ *T. pallidum* cell number fold change versus mean mammalian cell division time (hours), with the solid line representing linear regression and shaded 95% confidence interval. Regression analysis indicated a positive but non-significant association (slope = 0.034; R^2 = 0.37; p = 0.147).

Figure 2. Growth of *T. pallidum* in supporting and non-supporting mammalian cell lines. (A) *T. pallidum* counts from the *T. pallidum* adherent fraction in supporting cell lines (Sf1Ep, CAL-39, and HEPG2). (B) Total *T. pallidum* counts over time in supporting cell lines, overlaid with stacked barplots showing percentages of attached (light grey) and floating (dark grey) fractions. (C) Attached *T. pallidum* counts in non-supporting cell lines (HEK-293, SW954, JAR, JEG-3). (D) Total *T. pallidum* counts in non-supporting cell lines with stacked bars indicating attached and floating fractions. Data points represent mean values across biological replicates. For supporting cell lines, there is no significant difference between conditions between day 1 and 7 (p-values for all comparisons for each day reported in Supplementary Table 1). For non-supporting cell lines, differences between cell lines were also not significant across the whole experiment (ANOVA p = 0.142).

Figure 3. Interaction of *T. pallidum* with host cell lines. (A) Snapshot from live-cell imaging (Video file) showing two populations of *T. pallidum* interacting with the host cell surface. The population marked by the circle in the top right represents cells exhibiting single polar attachment, while the population marked by the circle in the bottom left represents *T. pallidum* cells displaying crawling-like motility. (B) SEM of crawling-like *T. pallidum* populations. The top panel shows a *T. pallidum* cell fully extended on the host cell surface, remaining attached along its length while moving across the cell. The bottom panel shows a *T. pallidum* cell captured at the moment of turning in the opposite direction, also observed in the Video file. (C) TEM of *T. pallidum* cells attached to the host cell surface by a single pole.

Video file link (real speed *T. pallidum* interaction with the host cell)

 https://drive.google.com/file/d/14sPVNNl79a9wy-LJo6AXBiy3GagYyBuB/view?usp=drive_link

Acknowledgments

This work was supported by the Biotechnology and Biological Sciences Research Council (BBSRC) (grant number BB/010589), the Bill & Melinda Gates Foundation (INV-051483), and the Wellcome funding (grant 220540/Z/20/A). We also thank Dr. Matthew Garnett (Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, UK) for providing access to cell lines in the Cell Model Passports dataset.

Supplementary material

Supplementary Figure 1. Floating fraction of *T. pallidum* in supporting (Sf1Ep, CAL-39, HEPG2) and
non-supporting (HEK-293, SW954, JAR, JEG-3) cell lines across days 1, 2, 3, 5, and 7. Boxplots show
the distribution of floating *T. pallidum* fractions for each group, with individual data points overlaid.
Wilcoxon rank-sum tests were performed separately for each day (day 1, p = 0.057; day 2, p = 0.057;
day 3, p = 0.114; day 5, p = 0.229; day 7, p = 0.229).

Supplementary table 1. Pairwise contrasts from a linear model comparing CAL-39, HepG2, and Sf1Ep across 10 days. Reported are estimated differences, standard errors, degrees of freedom, t-ratios, adjusted p-values. Significant differences after adjustment were observed primarily at later time points, including CAL-39 vs HepG2 on days 6 and 8, and contrasts involving Sf1EP on days 9 and 10, indicating time-dependent divergence between cell lines.

Contrast	Day	Difference	Se	Df	T_Ratio	P_Adjusted
CAL-39 - HEPG2	day_1	3746000	5254010	58	0.712979	0.756833
CAL-39 - SF1EP	day_1	2313333	5254010	58	0.440299	0.898848
HEPG2 - SF1EP	day_1	-1432667	5254010	58	-0.27268	0.959869
CAL-39 - HEPG2	day_2	9020667	5254010	58	1.716911	0.207578
CAL-39 - SF1EP	day_2	6734000	5254010	58	1.281688	0.411135
HEPG2 - SF1EP	day_2	-2286667	5254010	58	-0.43522	0.901043
CAL-39 - HEPG2	day_3	10901333	5254010	58	2.07486	0.104008
CAL-39 - SF1EP	day_3	7410000	5254010	58	1.410351	0.342288
HEPG2 - SF1EP	day_3	-3491333	5254010	58	-0.66451	0.784895
CAL-39 - HEPG2	day_4	8213333	5254010	58	1.563251	0.269663
CAL-39 - SF1EP	day_4	5490000	5254010	58	1.044916	0.551868
HEPG2 - SF1EP	day_4	-2723333	5254010	58	-0.51833	0.862722
CAL-39 - HEPG2	day_5	6898667	5254010	58	1.313029	0.393771
CAL-39 - SF1EP	day_5	3283333	5254010	58	0.62492	0.80708
HEPG2 - SF1EP	day_5	-3615333	5254010	58	-0.68811	0.771345
CAL-39 - HEPG2	day_6	13632667	5254010	58	2.594717	0.031641
CAL-39 - SF1EP	day_6	2283333	5254010	58	0.434589	0.901316
HEPG2 - SF1EP	day_6	-1.1E+07	5254010	58	-2.16013	0.086819
CAL-39 - HEPG2	day_7	11850000	5254010	58	2.25542	0.070455
CAL-39 - SF1EP	day_7	2673333	5254010	58	0.508818	0.867355
HEPG2 - SF1EP	day_7	-9176667	5254010	58	-1.7466	0.196854
CAL-39 - HEPG2	day_8	16310000	5254010	58	3.104296	0.00817
CAL-39 - SF1EP	day_8	4830000	5254010	58	0.919298	0.630397
HEPG2 - SF1EP	day_8	-1.1E+07	5254010	58	-2.185	0.082271
CAL-39 - HEPG2	day_9	5723333	5254010	58	1.089327	0.524476
CAL-39 - SF1EP	day_9	-1.7E+07	5254010	58	-3.32698	0.004297
HEPG2 - SF1EP	day_9	-2.3E+07	5254010	58	-4.41631	0.00013
CAL-39 - HEPG2	day_10	2703333	5254010	58	0.514528	0.864582
CAL-39 - SF1EP	day_10	-2.8E+07	5254010	58	-5.41935	3.58E-06
HEPG2 - SF1EP	day_10	-3.1E+07	5254010	58	-5.93388	5.22E-07

References

1. Gomez GB, Kamb ML, Newman LM, Mark J, Broutet N, Hawkes SJ. 2013. Untreated maternal

syphilis and adverse outcomes of pregnancy: a systematic review and meta-analysis. Bull World

Health Organ 91:217–226.

2. World Health Organization. Implementing the global health sector strategies on HIV, viral

hepatitis and sexually transmitted infections, 2022–2030 |report on progress and gaps 2024.

https://www.who.int/publications/i/item/9789240094925. Retrieved 22 December 2025.

3. Edmondson DG, Hu B, Norris SJ. 2018. Long-Term In Vitro Culture of the Syphilis Spirochete

Treponema pallidum subsp. pallidum. mBio 9:e01153-18.

4. Romeis E, Tantalo L, Lieberman N, Phung Q, Greninger A, Giacani L. 2021. Genetic

engineering of Treponema pallidum subsp. pallidum, the Syphilis Spirochete. PLoS Pathog

17:e1009612.

5. Tantalo LC, Lieberman NAP, Pérez-Mañá C, Suñer C, Mayans MV, Ubals M, González-Beiras

C, Rodríguez-Gascón A, Canut A, González-Candelas F, Mueller J, Tapia K, Greninger AL,

Giacani L, Mitjà O. 2023. Antimicrobial susceptibility of Treponema pallidum subspecies

pallidum: an in-vitro study. The Lancet Microbe 4:e994–e1004.

6. Lay BDD, Cameron TA, Lay NRD, Norris SJ, Edmondson DG. 2021. Comparison of

transcriptional profiles of Treponema pallidum during experimental infection of rabbits and in

vitro culture: Highly similar, yet different. PLOS Pathogens 17:e1009949.

7. 2024. In-Depth Proteome Coverage of In Vitro-Cultured Treponema pallidum and Quantitative

Comparison Analyses with In Vivo-Grown Treponemes. Journal of Proteome Research

23:1725–1743.

8. Edmondson DG, Norris SJ. 2021. In Vitro Cultivation of the Syphilis Spirochete Treponema

pallidum. Curr Protoc 1:e44.

9.   Fitzgerald T. 1981. In vitro cultivation of Treponema pallidum: a review. Bull World Health

Organ 59:787–812.

10.   Edmondson DG, DeLay BD, Kowis LE, Norris SJ. Parameters Affecting Continuous In Vitro

Culture of Treponema pallidum Strains. mBio 12:e03536-20.

11.   Trujillo D, Agudelo CA, Chavarriaga A, Villa P, Vélez A, Cardona R, Hidrón A. 2023. Bone

involvement in non-congenital syphilis. Biomedica 43:157–163.

12.   Okamoto K, Hatakeyama S, Umezawa M, Hayashi S. 2018. Gastric syphilis: The great imitator

in the stomach. IDCases 12:97–98.

13.   Al Dallal HA, Narayanan S, Alley HF, Eiswerth MJ, Arnold FW, Martin BA, Shandiz AE. 2021.

Case Report: Syphilitic Hepatitis–A Rare and Underrecognized Etiology of Liver Disease With

Potential for Misdiagnosis. Front Med 8.

14.   Santos-Coelho M, Alves Barbosa J, Valejo Coelho MM, João A. 2024. An unusual case of

extragenital primary syphilis. An Bras Dermatol 99:289–290.

15.   Arzoun H, Mohammed F, Saeed S, Khan TMA, Singh K. 2024. An Unusual Case of Syphilis

With Pulmonary Involvement. Cureus 16:e63758.

16.   Strypens T, Alliet G, Roef G, Winne L. 2022. Thyroid gland involvement in secondary syphilis:

a case report. Acta Clin Belg 77:137–141.

17.   Teixeira AM, Meireles E, Pereira Fontes C, Manuel M. Ocular Syphilis: A Case Report. Cureus

14:e23509.

18.   Adachi E, Koibuchi T, Okame M, Sato H, Imai K, Shimizu S, Tsurita G, Oyaizu N, Iwamoto A,

Fujii T. 2011. Case of Secondary Syphilis Presenting with Unusual Complications: Syphilitic

Proctitis, Gastritis, and Hepatitis. Journal of Clinical Microbiology 49:4394–4396.

19.   Grillová L, Romeis E, Lieberman NAP, Tantalo LC, Xu LH, Molini B, Trejos AT, Lacey G,

Goulding D, Thomson NR, Greninger AL, Giacani L. 2024. Bright New Resources for Syphilis

Research: Genetically Encoded Fluorescent Tags for Treponema pallidum and Sf1Ep Cells. Mol

Microbiol 122:455–464.