Chromatin architecture and physical constriction cooperate in phenotype switching and cancer cell dissemination

Pietro Berico[1,2,#], Cody Dunton[3], Luay Almassalha[4], Amanda Flores-Yanke[1,2], Karla I. Medina[3], Nicolas Acosta[3], Tara Muijlwijk[2,5], Catherine Do[1,6], Soobeom Lee[1,6], Sharon N Edmiston[7,8,9], David L. Corcoran[7,8,9], Allison Reiner[10], Caroline Kostrzewa[10], Kathy Dorsey[7,8,9], Milad Ibrahim[2,5], Shi Qui[2,5], Ronglai Shen[10], Nancy E. Thomas[7,8,9], Amanda W. Lund[2,5], Ata S. Moshiri[5], Iman Osman[2,5], Iannis Aifantis[1,6], Jane A. Skok[1,6], Vadim Backman[3,#], Eva Hernando[1,2,#]

[1] Department of Pathology, NYU Grossman School of Medicine, New York, NY, USA.

[2] Interdisciplinary Melanoma Cooperative Group, Perlmutter Cancer Center, NYU Langone Health, New York, NY, USA.

[3] Department of Biomedical Engineering, Northwestern University, Evanston, IL 60208, USA.

[4] Medical Scientist Training Program, Feinberg School of Medicine, Northwestern University, Evanston, IL 60611, USA.

[5] The Ronald O. Perelman Department of Dermatology, New York University School of Medicine, New York, NY, USA.

[6] Perlmutter Cancer Center, NYU Langone Health, New York, NY, USA.

[7] Department of Dermatology, School of Medicine, University of North Carolina, Chapel Hill, NC.

[8] Lineberger Comprehensive Cancer Center (LCCC), University of North Carolina at Chapel Hill, Chapel Hill, NC.

[9] Department of Genetics, School of Medicine, University of North Carolina, Chapel Hill, NC.

[10] Memorial Sloan Kettering Cancer Center, New York, NY, USA.

#Corresponding authors: Eva.Hernando-Monge@nyulangone.org, v-backman@northwestern.edu, Pietro.Berico@nyulangone.org

Abstract

Phenotypic plasticity is a prominent cancer feature that contributes to metastatic potential and resistance to therapy across multiple cancer types. Cancer cell state transitions have been attributed to transcriptional programs, such as the AP1/TEAD-regulated gene network driving the mesenchymal-like (MES) phenotype. In addition, during dissemination, tumor cells are subjected to variable loads of physical mechanical pressure and constriction across transited tissue, which are thought to impact nuclear molecular crowding. How the interplay between mechanical pressure, global 3D nuclear architecture and transcriptional programs contributes to MES identity and metastatic adaptation remains unclear. Using cutaneous melanoma as a model for early dissemination, we integrate in vitro and in vivo epigenomic profiling with nanoscale imaging of cell lines and patient samples to investigate chromatin organization features underlying the MES phenotype. We find that in MES cells, CTCF is relocated from domain boundaries to regulatory regions of EMT-like genes, leading to reduced insulation, extended topological associated domains (TADs) and increased inter-domain contacts, and de novo formation of chromatin hubs. This conformational rewiring, along with loss of heterochromatin, supports nuclear deformability during invasion and dissemination. Conversely, physical constriction of melanocytic cells induces MES-like chromatin features —including CTCF repositioning and heterochromatin loss— and promotes metastasis in vivo. Similarly, pharmacological inhibition of the heterochromatin mark H3K9me3 triggers MES characteristics and increases invasiveness. These results demonstrate that metastatic competency involves both epigenetic and structural nuclear reprogramming, enabling shifts in gene networks and physical adaptability. Our findings reveal mechanistic links between nuclear architecture and aggressive tumor behavior, identifying potential biomarkers and therapeutic targets to intercept metastatic progression.

Introduction

Metastasis is the leading cause of death among cancer patients[1-3]. Still, the mechanisms behind cancer dissemination remain largely unknown[4] and an unmet need to advance therapeutic options. Modern omics technologies have proven that (irreversible) genetic mutations do not fully explain cancer cell metastatic behavior nor patients' outcome, while 'reversible' epigenetic mechanisms can also confer aggressive potential[5, 6]. Cancer cells acquire transient phenotypic states through heritability and plasticity of gene regulatory networks (GRNs) led by different transcription factors (TFs)[7], as an adaptive response to the tumor microenvironment stimuli[8]. Among the cues, tissue cellular density and matrix composition exert biodynamic forces directly toward tumor cells which can increase molecular crowding within the nuclear compartment[9-11].

CTCF is the master regulator of nuclear architecture and, together with cohesin complex, nuclear lamina, non-coding RNAs, and other DNA and RNA-binding proteins pack the two meters long human DNA into chromatin, which expands and contracts to regulate gene expression, DNA replication and repair, RNA transcription and splicing[12]. Chromatin is organized into thousands of nanoscale chromatin packing domains (CPDs) that are only vizualied by super-resolution imaging. These are typically up to 200 nm in size and containing 10 kb to 1 Mbp of DNA. These domains represent a distinct organization layer in addition to compartments, TADs and loops that are defined by contacts through chomatin conformation capture or similar approaches, and may function as transcriptional memory elements[13-16]. Not surprisingly, alterations in nuclear architecture can lead to cancer[17]. For instance, genetic and epigenetic aberrations can disrupt compartments, domains and structural loops to rewire cis-regulatory elements (CRE)[18-22] and establish oncogenic chromatin hubs[23, 24], composed of CREs simultaneously contacting multiple genes involved in pro-tumorigenic biological processes.

Recently, it has also emerged that chromatin regulation provides nuclear physical structure,

mechanical rigidity[25, 26] and shape[27] which steer microenvironment adaptability[28, 29] and may ease

tumor cell compression during intra- and extravasation in the metastatic cascade[30].

Concomitantly, mechanical compression can directly rewire 3D chromatin, establishing an

intricated bi-directional regulatory feedback between the two processes[28, 31, 32,33] . The amount

and maturation of CPDs within the nucleus changes the mechanical stability of a cell; for instance,

stable mature domains have been shown to lower the probability of blebbing during cell migration,

with predominantly 'domain-less' chromatin forming the blebs.[34]   However, the impact of

constriction-induced changes on nuclear architecture and phenotype switch during metastasis

remains unclear.

A mesenchymal or dedifferentiated cancer cell state (MES), enriched on an epithelial-to-

mesenchymal-transition (EMT) program, is thought to be the founder of dissemination in multiple

cancer types[35-39]. Skin cutaneous melanoma (SKCM) represents an exquisite model of metastasis

because it can disseminate from thin primary lesions[40] —which already display intra-tumor

heterogeneity including MES cells[39, 41, 42]— irrespective of the genetic drivers. In response to

microenvironmental cues[43], melanoma cells 'switch' from a differentiated melanocytic-like state

(MEL) led by MITF/SOX10 GRNs, to an EMT-like, AP1/TEADs-dependent MES phenotype[44, 45].

However, MITF/SOX10 and AP1/TEADs silencing are insufficient to trigger a complete switch

toward a MES state[46, 47], suggesting other changes in addition to differential gene expression are

required [45, 47-53]. We propose that the bi-directional interplay between nuclear 3D chromatin

organization and physical compression critically shapes cancer cell plasticity by impacting both

gene expression and mechanical properties. In the context of melanoma, our understanding of

global nuclear architecture and its contribution to phenotypic plasticity and metastasis is largely

unknown.

Combining multiple omics techniques, we profiled the nuclear physical properties and global 3D

chromatin architecture of patient samples, and of a large panel of melanoma cell lines

representative of distinct phenotypic states (MEL, intermediate —INT— and MES) or MEL upon

silencing of MITF and SOX10 (siMS), the key GRNs of the MEL state. Compared to MEL, siMS

and INT cells, we observed a dramatic rewiring of 3D chromatin in MES cells and a preferential

binding of CTCF to CRE of EMT-related genes and chromatin hubs over TAD boundaries.

Accordingly, silencing of CTCF re-activates MEL-like GRNs and disrupts the activity of a subset

of chromatin hubs, some of which support MES cell migration. While melanoma states exhibit

extensive differences in nuclear topology, these only result in limited transcriptional changes,

suggesting that MES-like 3D chromatin features confer additional biological properties

contributing to metastasis beyond gene expression, such as nuclear deformability and flexibility.

Supporting this hypothesis, we find MES cells display higher nuclear flexibility and lower stiffness

by mathematical polymer simulations, Partial Wave Spectroscopy (PWS) reduced H3K9me3

packing scale domains by spectroscopic single-molecule localization microscopy (sSMLM) and

increased cis-long interactions by HiC. Strikingly, these molecular features are partially adopted

by MEL cells undergoing constriction during transwell migration or upon H3K9me3 inhibition,

which are both independently able to increase their metastatic potential. Altogether, our study

shows that nuclear architecture and cell compression co-regulate gene expression and

mechanical properties unleashing the MES identity and metastatic capacity.

Methods

Cell culture

All melanoma cell lines were grown at 37°C in 5% CO_2 and routinely tested for mycoplasma

contamination using MycoAlert® PLUS Mycoplasma Detection Kit (Lonza, #LT07-703). SKMEL-

5, SKMEL-28, SKMEL-239 and SKMEL-147 were kindly provided by Alan Houghton (Memorial

Sloan-Kettering Cancer Center, New York, NY, USA); 501mel were purchased from Yale

University, MM cell lines (MM057, MM074, MM087, MM029 and MM099) were obtained from Prof.

Ghanem Ghanem (Institute Jules Bordet, Bruxelles, BE). MM047 cells were obtained from

Frédéric Coin and Irwin Davidson (IGBMC, Illkirch, France). 501mel and SK-MEL cell lines were

grown in 10% Fetal Bovine Serum (FBS) in DMEM and penicillin-streptomycin (PS). MM cell lines

were grown in Ham-F10 supplemented with 10% FBS, PS and 2mM L-Glutamine.

Cells were transfected with 25nM of siNTC (Horizon Discovery, #D-001810-10), siMITF (Horizon

Discovery, #L-008674-00) and siCTCF (Horizon Discovery, #L-020165-00) using Lipofectamine

RNAiMAX (Thermo Scientific Scientific, 13778150) following manufacturer's instructions.

Transwell migration assay

For migration under constriction, $1x10^6$ 501mel and MM099 were seeded into cell culture inserts

with 3.0μm pore size (Corning, #353091) into a dedicated 6 well plate (Corning, #353502) and

grown in complete media. 72hrs later, cells were gently detached from both sides of the insert

using trypsin and processed for HiC, CUT&Tag (CTCF, H3K9me3 and H3K27ac), ATAC and

RNAseq.

To assess the effects of H3K9me3 perturbation, $1-2x10^5$ 501mel and SKMEL-28 were seeded

into cell culture inserts with 8.0μm pore size (Corning, #353097) placed into a dedicated 24 well

plate (Corning, #353504) and grown in complete media supplemented with DMSO or 5nM

Chaetocin (MedChem Express, #HY-N2019) for 12-24hrs. Next, inserts were rinsed twice in PBS

and fixed with 4% PFA-PBS solution (Electron Microscopy Sciences, #15733-10) for 10min at

room temperature. Next, inserts were stained in 0.2% crystal violet solution for 15min at room

temperature, rinsed in distilled water by simultaneously cleaning the inner side with a cotton swab.

After drying at room temperature, inserts were imaged using a digital brightfield microscope

(Invitrogen, EVOS core) and quantified with ImageJ.

ChIP and HiC

Melanoma cell lines were double crosslinked for 30min in 25nM EGS (Thermo Scientific Scientific, #21565) and for 10min in 1% FA (Sigma-Aldrich, #47608) at room temperature and then processed for ChIP and HiC.

For ChIP, 1×10^7 cells were lysed in ChIP lysis buffer (CLB = 10mM Tris HCl pH8, 100mM NaCl, 1mM EDTA pH8, 0.5mM EGTA pH8, 0.1% Sodium Deoxycholate, 0.5% N-lauroylsarcosine, 1% Triton X-100) and sonicated with Diagenode Bioruptor for 15-30 cycles (30" ON +30" OFF max intensity). 5ug/IP of H3K27ac antibody (Active Motif, #91193), H3K9me3 (Active Motif, #61013) or IgG control (Cell Signaling Technology, #3900) were prebound to 50ul of Protein A/G magnetic beads (Pierce Thermo Scientific Scientific, #88803) in PBS+BSA 0.5% for 1hr @RT. Chromatin was diluted to 1ml in CLB and magnetic beads-antibody complex was added to each IP ON at 4°C on revolver. IP were washed twice in Wash Buffer I (WBI, 20mM Tris HCl pH7.5, 150mM NaCl, 2mM EDTA pH8, 0.1% SDS, 1% Triton X100), Wash Buffer II (WBII, 10mM Tris HCl pH8, 250mM LiCl, 1mM EDTA, 1% Triton X100, 0.7% Sodium Deoxycholate) and TET Buffer (10mM Trist HCl pH8, 1mM EDTA, 0.2% Tween20) and eluted in Elution buffer (EB, 10mM Tris HCl pH8, 300mM NaCl, 5mM EDTA pH8, 0.5% SDS, 10ug RNAse A, 100ug proteinase K) 30min at 37°C + ON at 55°C. IP elutions were purified using MinElute Reaction Cleanup kit (Qiagen, #28204) and libraries were prepared using NEBNext Ultra II DNA Library Prep Kit for Illumina (NEB, #E7600S) following manufacturer's instructions.

For HiC, 2.5×10^6 cells were processed using Arima-HiC+ kit (Arima Genomics, #A101020) following the manufacturer instructions. HiC libraries were generated using Arima-HiC+ kit (Arima Genomics, #A510008, #A303011) following manufacturer's instructions.

ChIP and HiC libraries were pooled together at 2nM concentration and sequenced paired-end 2x150bp using Illumina NovaSeqX+.

HiC in DNA from FFPE tissues

Between 5-10 Formalin Fixed Paraffin Embedded (FFPE) patient tumor tissue sections of 10μm per sample were cut and pooled in a 1.5mL tube for de-waxing and re-hydration using xylene, ethanol and molecular grade water. Next, FFPE samples were processed with the Arima-HiC+ FFPE Kit (Arima Genomics, #A101060) following manufacturer's instructions. HiC FFPE libraries were generated using Arima-HiC+ kit (Arima Genomics, #A510008, #A303011) following manufacturer's instructions.

Omni-ATACseq

$5x10^5$ cells were harvested and assessed to not have more than 15% of cell death using trypan blue. Next, dead cells were additionally removed treating cells with 200U/ml of DNAse (Worthington, #LS002007). Cells were resuspended in 500ul of cold buffer 1 (10mM Tris-HCl pH 7.5, 10mM NaCl, 3mM $MgCl_2$, 0.1% NP-40, 0.1% Tween-20) and the equivalent volume for $5x10^4$ cells (50ul) was kept for the next steps. A final concentration of 0.1% of digitonin was added to the cells and incubated on ice for 3 minutes. Digitonin permeabilization was quenched adding 1ml cold buffer 2 (10mM Tris-HCl pH 7.5, 10mM NaCl, 3mM $MgCl_2$, 0.1% Tween-20). Afterward, cells were pelleted and resuspended in 50ul of transposition mixture (20mM Tris HCl pH 7.5, 10mM $MgCl_2$, 20% dimethyl formamide, PBS, 1% digitonin, 10% tween-20) supplemented with 2.5-5ul of Transposase (Illumina, #20034197) and incubated in thermomixer at 37°C for 30 minutes with 1000rpm. Transposition mixture was purified using DNA Clean and Concentrator-5 kit (Zymo, #D4014) for library preparation. 20ul of transposed DNA was processed for Nextera library using the same protocol described in Schmidl work[54]. OmniATAC libraries were pooled together at 2nM concentration and sequenced paired end 2x150 using Illumina NovaSeqX+.

CUT&Tag

For CTCF genomic occupancy was assessed using CUT&Tag-IT™ (Active Motif, #53160) following manufacture instructions. Briefly, 1.5-5x10[5] cells were captured with concanavalin A beads and 1ug of IgG control (Cell Signaling Technology, #3900) or CTCF (Active Motif, #61311) antibodies were incubated 2hrs at 4°C on revolver in cell lysis buffer. For H3K9me3 and H3K27ac Pre and Post migration, 1ug of antibodies H3K9me3 (Active Motif, #39062) and H3K27ac (Active Motif, #39133) were incubated overnight at 4°C on revolver in cell lysis buffer. Afterward, 1ul of guinea pig anti-rabbit antibody was added to the reaction followed by 1hr of tagmentation at room temperature with 1ul of Tn5 transposase enzyme. Tagmentation was stopped with EDTA, SDS and proteinase K incubation for 1hr at 55°C. Tagmented DNA was purified using DNA purification reagents within the kit. Finally, tagmented DNA was processed for Nextera library preparation using i7 and i5 indexed primers and Q5 TAQ polymerase. CUT&Tag libraries were pooled together at 2nM concentration and sequenced paired end 2x50 using Illumina NovaSeqX+.

RNAseq

Total RNA libraries were prepared using automated TruSeq stranded total RNA and RiboZero Gold kit (Illumina, #RS-122-2301) according to the manufacturer's instructions. RNA libraries were sequenced paired end 2x50 using Illumina NovaSeqX+.

Methylation array

DNA methylation profiling was performed using the genome-wide Infinium MethylationEPIC array (Illumina) as described previously[55]. Briefly, DNA (250ng/ sample) from melanoma cell lines was extracted with QIAamp DNA Micro Kit (Qiagen, #56304) and bisulfite-treated using EZ DNA Methylation GOLD kits (Zymo Research, #11-335) according to manufacturer's protocol. BeadArray processing and scanning was performed in the UNC-CH Mammalian Genotyping

Core. Methylated and non-methylated control samples (Zymo Research, #50-444-307) were

included in the experiment. BeadArray data was assembled using Illumina GenomeStudio

methylation module (v3.2). Background intensity from negative controls was subtracted from each

data point.

Tumor dissociation

Freshly dissected tumors from mice were dissociated at single cell by first chopping the tissue in

a petri dish with a scalpel. Then, whole tumors were resuspended in 4mL/gr of HBSS (Sigma-

Aldrich, #H6648) supplemented with 750U/ml of Collagenase I (Sigma-Aldrich, #C2674) and

625U/ml of DNAse I (Worthington, #LS002138) and incubated for 50min at 37°C upon strong

shaking. Cell suspension was sequentially filtered through a first 70μm cell strainer (Fisher

Scientific, #22363548) and then a 40μm cell strainer (Fisher Scientific, #22363547). Next, single

cell suspensions were depleted of immune and dead cells by flow cytometry sorting (Sony

Biotechnology, SY3200 HAPS) using PE-conjugated CD45 (BioLegend, mouse #111103) and

DAPI staining.

PIPseq

Before processing, cell number and viability were assessed using a TC20 Cell Automated Cell

Counter (BioRad). Briefly, cells were partitioned into PIPs (PIPseq T20 3' Single Cell RNA Kits,

V4.0PLUS, Fluent BioScience) with subsequent cDNA generation and final library preparation

following manufacturer's instructions. Sample input varied from ~80,000 to 200,000 per

reaction. cDNA quality and concentration were evaluated on a BioAnalyzer 2100 using a High

Sensitivity DNA Kit (Agilent Technologies). Final gene expression libraries were amplified between

6 and 15 cycles and visualized on an Agilent TapeStation 4200 using High Sensitivity D1000

ScreenTape (Agilent Technologies). Sequencing was performed using a NovaSeq X+ 10B 200

Cycle Flowcell (Illumina).

Low input HiC (LoHiC)

Between $0.5\text{-}1\times10^6$ tumor cells were double crosslinked and subjected to HiC as previously

described[56]. Then, single cells were processed with Chromium Next GEM single Cell ATAC

Reagent Kits v2 (10X Genomics) following manufacture instructions.

CRISPR/Cas9 screen

MM099 cells were transduced with lentivirus carrying a doxycycline inducible dCas9-KRAB-

MeCP2 (Addgene, #140690) and a custom CRISPR library pRSGScribe1 (Cellecta) targeting 43

MES hubs plus 2 positive (*ACTB*, *GAPDH*) and 5 negative controls[57] (5 sgRNAs/target) for a total

of 230 sgRNAs. Single guide RNAs (sgRNAs) were designed using CRISPick[58]. After viral library

titration, cells were infected with a 0.3 MOI to prevent multi-copy insertion and have a maximum

of one sgRNA/cell. Finally, cells were collected at day 0 and 21 upon doxycycline induction

(1μg/ml). Cell pellets were lysed to retrieve genomic DNA using QIAamp DNA Micro Kit (Qiagen,

#56304). sgRNA sequencing libraries for t0 and t21 were prepared using NGS prep kit for sgRNA

Libraries with supplementary primers (Cellecta, #LNGS-360, #LNGS-360-SP) following

manufacturer's instructions.

Spectral partial wave spectroscopy (PWS) Imaging

PWS imaging was performed under physiological conditions (37ºC, 5% CO_2) using a stage-top

incubation system (In Vivo Scientific). Cells were seeded on 8-chamber cover glass slides and

allowed to adhere for 24 hours prior to imaging. Measurements were acquired on an inverted

microscope (Leica DMIRB) equipped with a 63x oil-immersion objective (HCX PL APO, Leica), a

CCD camera (ImageEM C9100-13, Hamamatsu), and a liquid crystal tunable filter (LCTF, Cri

VariSpec) to enable spectrally resolved imaging. Broadband illumination was provided by a white-

light LED source (X-Cite 120 LED, Excelitas), and a long-pass filter (Semrock BLP01-405R-25)

was used to reject short-wavelength excitation light. Spectra image stacks were collected from

500 to 700 nm at 2-nm intervals, as previously described[59-61].

PWS imaging quantifies wavelength-dependent interference patterns arising from multiple light

scattering events within the sample, which are sensitive to nanoscale fluctuations in the local

refractive index. Because refractive index is proportional to macromolecular density, PWS

provides a label-free optical readout of chromatin nanoscale organization. For each pixel in the

lateral image plane (x,y), the standard deviation of the backscattered intensity across the spectral

dimension (λ) was computed, yielding a local chromatin packing scaling parameter, $D_a(x, y)$. The

average nuclear chromatin packing scaling, D_n, was obtained by averaging D_a across all pixels

within the nuclear region of interest. D_n reflects the organization of chromatin into spatially

segregated packing domains and is derived from optical modeling and chromatin density

autocorrelation functions established in prior chromatin tomography and PWS studies[61-63]. For

each experimental condition, D_n values were calculated for individual nuclei across biological

replicates, and group-level statistics were used to quantify treatment-associated changes in

chromatin packing.

In addition to live-cell measurements, PWS imaging was also performed on formalin-fixed

paraffin-embedded tumor sections. OPAL-stained tumor sections were imaged by PWS and multi-

channel fluorescence imaging. PWS and OPAL images were spatially registered, enabling

chromatin packing metrics to be associated with immunophenotypically defined cell populations

within intact tumor tissue.

Dynamic PWS Measurements

Dynamic PWS imaging was performed as previously described[61, 64] to quantify nanoscale

chromatin motion in live cells. In this mode, wide-field backscattered intensity was recorded at a

fixed wavelength (550 nm) over time, generating a three-dimensional data cube with dimensions

(x, y, t). Temporal intensity fluctuations at each pixel arise from the movement of intracellular

macromolecular structures, including chromatin. Fractional moving mass (FMM) was computed

from the temporal variance of the backscattered intensity, σ_t^2, and normalized using a biophysical

scattering model to convert raw fluctuations into a quantitative measure of nanoscale mass

motion. Following Gladstein _et. al._ (2019)[61], FMM is defined as:

309
$$\sigma_t^2 \cdot \frac{\pi \rho_0}{2\Gamma^2 k^3 n_i} \left(\frac{NA_i}{NA_c}\right)^2 \left(\frac{n_1}{n_m - n_1}\right)^2 = \rho_0 V_{cm} \phi = m_c \phi = m_f$$

Where m_f represents the mass undergoing motion within the sample and is given by the product

of the mass of a typical moving cluster (m_c) and the volume fraction of the mobile mass (ϕ). Here,

$m_c = \rho_0 V_{cm}$, with V_{cm} denoting the volume of a characteristic moving macromolecular cluster and

ρ_0 the dry mass density.

For normalization, the refractive index of a nucleosome was approximated as $n_m = 1.43$, the

nuclear refractive index as $n_1 = 1.37$, and the refractive index of the immersion medium as $n_i =$

1.518. The dry density of a nucleosome was taken as $\rho_0 = 0.55\ g\ cm^{-3}$. The scalar wavenumber

of the illumination light was $k = 1.57 \times 10^5\ cm^{-1}$, Γ denotes the Fresnel intensity coefficient for

normal incidence, and the numerical apertures of collection and illumination were $NA_c = 1.49$ and

$NA_i = 0.52$, respectively. As previously reported[61], σ_t^2 is sensitive to instrument-specific

parameters, including depth of field and substrate refractive index. These effects were corrected

using empirically determined calibration constants. Pixelwise FMM values were calculated for

each nucleus, and distributions were compared across experimental conditions to quantify

treatment-induced changes in chromatin mobility.

sSMLM

Cells were seeded on 8-chamber cover glass slides (Cellvis, #C8-1.5H-N) and cultured to

approximately 30% confluence. Samples were fixed in 4% paraformaldehyde prepared in

phosphate-buffered saline (PBS) for 10 minutes at room temperature, followed by two PBS

washes. After fixation, cells were quenched in 0.1% sodium borohydride in PBS for 7 minutes.

After three PBS washes, cells were permeabilized and blocked in PBS containing 0.2% Triton X-

100 and 3% bovine serum albumin (BSA) for 2hrs at room temperature. To label heterochromatic

regions of chromatin, cells were incubated overnight at 4ºC with anti-H3K9me3 antibody (Abcam,

ab176916) at 1.2 µg/mL in blocking buffer. After 3 washes in PBS with 0.1% Triton X-100 and

0.2% BSA, cells were incubated with Alexa Fluor 647-conjugated secondary antibody (Thermo

Scientific, #A-21245) at 2 µg/mL for 1 hour at room temperature. Cells were washed twice in PBS;

to stain euchromatic regions, the labeling steps aforementioned were repeated using anti-

H3K27Ac antibody (Thermo Scientific, #MA5-23516) and tagged with fluorophore Alexa Fluor

488-conjugated secondary antibody (Thermo Scientific, #A-11001) at 4 µg/mL in blocking buffer.

To conduct imaging in STORM buffer containing 0.5 mg/mL glucose oxidase (Sigma-Aldrich), 40

µg/mL catalase (Roche), and 100 mg/mL glucose in TN buffer (50 mM Tris, pH 8.0, and 10 mM

NaCl). Super-resolution imaging was performed using a Nikon Eclipse Ti-U inverted microscope

equipped with a 100×/1.49 NA oil immersion objective (SR APO TIRF, Nikon) and an iXon Ultra

888 electron-multiplying CCD camera (Andor). A 637-nm laser (Obis, Coherent) provided

excitation at the sample plane with an average power of 3–10 kW/cm² and using type F immersion

oil (Nikon, #MXA22168). At least 10,000 frames were acquired per field of view with a 30 ms

exposure time.

Western blotting

Cells were harvested and lysed in LSDB buffer (0.5M KCl, 50mM Tris HCl pH 8.0, 20% Glycerol,

1% NP40, 1mM DTT) through three cycle of snap freezing in liquid nitrogen and thermobath at

37°C. Cell pellet debris was removed with the help of a pipette tip following centrifugation at

20.000g for 15min at 4°C. Between 10-20µg of proteins were run into pre-casted gels Nupage™

4-12% Bis-Tris (Thermo Scientific) using the dedicated apparatus (Thermo Scientific, #EI0001).

Next, separated proteins were transfer into nitrocellulose membranes (Cytiva, #10600011) which

were subsequently blocked into 5% milk TBST solution. Primary antibody staining was conducted

by diluting antibodies in TBST (between 1:1000-5000 dilution, Supplemental reagents) and

incubated overnight at 4°C with the corresponding membranes upon gentle mix. Secondary

antibodies HRP-conjugated were diluted 1:10.000 in TBST and incubated 1hr at room

temperature with gentle shaking. Protein signal was revealed using luminol substrate mix

(Millipore, #WBLUR0500) and detected with an Odyssey XF Imaging System (LICORbio). Protein

quantification was conducted using ImageJ.

Immunoprecipitation (IP)

Protein immunoprecipitation was conducted as previously described[46]. Briefly, cells were

harvested, pellet and gently lysed into 2 pellet volume of hypotonic buffer (10mM Tris HCl pH 7.6,

1.5mM $MgCl_2$, 10mM KCl, 0.1% NP40) using mechanical shear with a loose Dounce to release

the cytoplasmic fraction. Next, nuclei were pellet and resuspended in 1 volume of sucrose buffer

(20mM Tris HCl PH 7.6, 15mM KCl, 60mM NaCl, 0.34M Sucrose) followed by a drop-by-drop

gradual integration of 0.3 volumes of high salt buffer (20mM Tris HCl pH 7.6, 25% glycerol, 1.5mM

$MgCl_2$, 0.2mM EDTA, 0.9M NaCl) upon gentle vortex. After 30 min incubation at 4°C on constant

slow spinning, ruptured nuclei were pellet, nuclear soluble fraction was collected and nuclei were

further resuspended in 1 volume of sucrose buffer supplied with 1mM $CaCl_2$ and 20U/μl of

Micrococcal Nuclease (NEB, #M0247S), and incubated in thermoblock for 10min at 37°C.

Micrococcal digestion was stopped by adding EDTA (final concentration 4mM) and lysate was

centrifuged 20.000g for 30min at 4°C to obtain the nuclear insoluble fraction.

For CTCF co-IP, 500μg of nuclear insoluble fraction was diluted in 1mL of TGEN buffer (10mM

Tris HCl pH 7.6, 150mM NaCl, 3mM $MgCl_2$, 0.1mM EDTA, 10% glycerol, 0.01% NP40) for each

IP. 1% of diluted extract was kept as western blot input. Next, between 5-10μg of primary

antibodies against CTCF (Cell Signaling Technology, #2899), FOSL1 (Cell Signaling Technology,

#5281), FOSL2 (Cell Signaling Technology, #19967), c-JUN (Cell Signaling Technology, #9165),

pan-TEAD (Cell Signaling Technology, #13295) and IgG control (Cell Signaling Technology,

#3900) were added in each IP and incubated overnight at 4°C on constant slow spinning. The

day after, 25μl of magnetic beads A/G (Thermo Scientific, #88803) previously rinsed in PBS +

0.5% BSA were added to each IP and incubated 4hrs at 4°C upon slow spinning. Next, beads-

antibody-proteins complexes were gently washed five times in cold TGEN buffer and eluted in

LSDB and 1X Laemmli buffer in a thermoblock for 10min at 98°C before proceeding with western

blot.

Immunofluorescence (IF)

Tissue microarrays (TMAs) were sectioned at 5 μm sections and immunostained on a Leica

BondRx auto-stainer according to the manufacturer's instructions. In brief, sections were

deparaffinized online and underwent antigen retrieval with either ER1 (Leica, AR9961; pH6) or

ER2 (Leica, AR9640; pH9) retrieval buffer at 100o for either 20 minutes. Sections were then

treated with 3% H2O2 to inhibit endogenous peroxidases. After blocking with Primary Antibody

Diluent (Leica, AR93520), slides were incubated with the first primary antibody and secondary

HRP polymer pair. Following secondary incubation, slides underwent HRP-mediated tyramide

signal amplification with a specific Opal® fluorophore. Once the Opal® fluorophore was covalently

linked to the antigen, primary and secondary antibodies were removed with a 95o retrieval step.

This sequence was repeated 5 more times with subsequent primary and secondary antibody

pairs, using a different Opal fluorophore for each primary antibody against human MITF (Sigma-

Aldrich, #HPA003259), TCF4 (Abcam, #ab223073), S100B (Abcam, #ab115803), p-EGFR (Cell

Signaling Technology, #4407), AXL (Cell Signaling Technology, #8661), and SOX10 (Sigma-

Aldrich, #383R-15). After antibody staining, sections were counterstained with spectral DAPI

(Akoya Biosciences, FP1490) and mounted with ProLong Gold Antifade (ThermoFisher Scientific,

P36935).

Semi-automated image acquisition was performed on an Akoya Vectra Polaris (PhenoImagerHT)

multispectral imaging system. Slides were scanned at 40X magnification using PhenoImagerHT

2.0 software in conjunction with Phenochart 2.0 and InForm 3.0 to generate unmixed whole slide

qptiff scans. Image files were uploaded to the NYUGSoM's OMERO Plus image data

management system (Glencoe Software).

Computational analysis

ChIPseq, CUT&Tag and ATACseq

Sequencing results were demultiplexed and converted in fastq using Illumina bcl2fastq software.

Next, fastq files were trimmed and filtered according to the qc report obtained with TrimGalore

(0.6.6,   https://github.com/FelixKrueger/TrimGalore)   and   fastqc   (0.11.7,

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/). For ChIP samples, fastq trimming

was set for Illumina adaptors; for ATACseq and CUT&Tag, it was set for Nextera adaptors.

Alignment to Hg38 human genome was performed using bowtie2 (2.4.1)[65] and sam files were

converted into bam using samtools (1.16)[66]. Bigwig files were generated from bam files using

bamcoverage (deeptools 3.5.6)[67] and filtered out for duplicate and bad genomic regions. Peak

calling was performed using MACS2 (2.2.9.1)[68] using paired-end mode, IgG (ChIP and CUT&Tag)

was used as control, a q-value 0.05 threshold was set and both, narrowPeak and summit peaks

were filtered for blacklisted regions using bedtools (2.31.0)[69]. An Hg38 CTCF whole genome track

generated by JASPAR[70] was used to filter CTCF peaks and maintain only the ones containing the

CTCF motif. Peak annotation was performed with homer (4.10)[71] using narrowPeak files and

Hg38 genome parameters. DNA motif analysis of CTCF was done by first expanding of 200bp

MACS2 summit peaks with bedtools slop, then fasta sequence was obtained using bedtools

getfasta which were processed using MEME-ChIP[72]. DiffBind[73] was used to identify state-specific

peaks for CTCF, H3K27ac, ATAC and H3K9me3.

HiC

For HiC samples, fastq trimming, qc, alignment, valid pairs identification and matrix generation

were performed using HiC-Pro workflow (3.1.0)[74] as previously described[75].

Briefly, valid pairs outputs were converted into juicer and cool formats using hicpro2juicebox and

hicpro2higlass respectively (HiC-Pro). Matrix normalization was performed using cooler balance

(cooltools)[76] and addNorm (juicer tools)[77]. Copy number variations (CNVs) and structural variants

(SVs) were detected using HiNT[78] cnv and tl functions at 50kb BIN resolution and chromosomes'

segmentation were pooled into one file and converted in Log2 using python. Segments with a

$Log_2 \geq 0.5$ and $Log_2 \leq -0.5$ were considered amplified and deleted respectively. HiC-Pro valid pairs

were filtered for CNVs and SVs using bedtools pairtobed. Compartment calling was conducted

using eigs-cis function (cooltools) and 100kb BIN resolution matrixes. For compartment switch

analysis, compartment score data from multiple cell lines were combined in a single text-

delimited-file and delta switch was calculated as described in Narang et al[19]: compartments

flipping sign with an absolute difference ≥ 0.5 and an absolute delta ≥ 1.2 were considered

switched. Compartments' saddle-plot was generated with cooltools. To measure H3K9me3,

H3K27ac, ATAC and RNA RPKM enrichment in the compartments, bigwig files from these

markers were segmented in 100KB BIN and averaged in each compartment using deeptools.

Topological Associated Domains (TADs) identification was conducted using cooltools insulation

providing an input matrix of 10kb BIN resolution, a 120kb window and Li threshold. From the tsv

output, i) TADs boundaries were identified if considered 'true' under the 'is_boundary' column and

ii) if the distance between two boundaries is less or equal to 3Mb was considered as a domain[79].

Loops were called using both 10kb and 5kb BIN matrixes with Peakachu[80] and 10kb and 5kb

were merged using hicMergeLoops (HiCexplorer). TADs boundaries and loops anchors co-

occupied by CTCF were retained for downstream analysis using bedtools intersect and

hicValidateLocation (HiCexplorer) respectively. HiC pile up maps, intra-TAD and inter-TAD

contacts data were generated using GENOVA r package[81]. CTCF occupancy at loops anchors

and TADs boundaries was calculated and visualized using deeptools. Loops and TADs were

annotated using bedtools intersect and pairtobed against GENCODEv44 Hg38 bed file

downloaded from UCSC Genome Browser. Graphs were generated combining Prism10, R and

phyton.

ABC-score model

Active-by-Contact (ABC) model[82] was generated following the authors guidelines

(https://github.com/broadinstitute/ABC-Enhancer-Gene-Prediction) as previously described[75].

Briefly, cCRE were called and measured in activity using makeCandidateRegions and

run.neighborhoods scripts in which ATACseq and H3K27ac bam and narrow.Peaks are

integrated. Next, HiC matrix were processed using juicebox_dump and

compute_powerlaw_fit_from_hic to extrapolate genome wide contacts. ABCscore for each active

cCRE was measured using predict function in which cCRE, HiC contacts and RNAseq are

integrated. Finally, cCRE were filtered using an ABCscore threshold of 0.02 and self-promoter

contacts were excluded for downstream analysis. DESeq2[83] was used to identify differentially

active cCRE across melanoma cell lines.

To identify chromatin hubs, sumABC (ΣABC) was calculated for each cCRE as the sum of all

contacted genes per cCRE and it was calculated with R (4.4.1) using the corresponding formulas:

$$\Sigma ABC = \sum_{j=1}^{g_i} ABC_{i,j}$$

ABC = score

g = gene

Next, each cCRE was ranked based on the number of contacting genes with an expression level

of Transcript Per Million (TPM) ≥1.

Polychrom

Chromatin 3D dynamics were simulated using a modified version of the Polychrom engine

(OpenMM-based molecular dynamics, https://github.com/open2c/polychrom), implemented in our

custom pipeline. Briefly, each chromosome is represented as a coarse-grained polymer, where

individual 10 kb bins from Hi-C data correspond to "beads" connected by harmonic bonds (rest

length = 1). Angular stiffness of each bead (k_θ) is computed from integrated epigenetic, Hi-C, and

TAD information:

• Epigenetic terms: local enrichment of H3K27ac and H3K9me3 (BigWig tracks)

• Compartment identity (A/B): sign(E1) from eigenvector decomposition

• TAD topology: TAD size-normalized scaling factor

• Hi-C contact environment: short-range, long-range, and inter-TAD interactions

Stiffness was modeled as:

$$k_i = \exp\left(-0.40 Z_{27} + 0.28 Z_9 - 0.25 sign(E1)\right) \times TAD_{factor} \times f(HiC)$$

followed by clipping to a biophysically stable range (0.3–4.0). Softness is defined as:

$$s_i = \frac{1}{k_i}$$

All forces used in the simulation include harmonic bonds, angular stiffness (per-bead k_θ), excluded

volume via polynomial repulsion and high-confidence Hi-C–derived long-range harmonic

restraints (above a quantile threshold). An initial random-walk polymer is energy minimized and

then propagated in Langevin dynamics for 150k integration steps (timestep 40 fs). Snapshots are

saved every 5,000 steps using Polychrom's HDF5Reporter, generating bead-track annotation, full

3D trajectory, contact map, per-frame radius of gyration (Rg) and curvature-based softness.

Outputs include genome-wide stiffness tracks, compartment-stratified distributions, TAD-level

summaries, and effective chromosome stiffness calculated from mean curvature across all

trajectory frames. To quantify chromatin deformability under nuclear constriction, we implemented

an additional energy-minimization pipeline. For each sample, the final equilibrium chromatin

configuration from the Polychrom simulation is subjected to increasing levels of axial compression

inside a narrowing slit. Given a slit half-height h, all beads satisfying $|z| > h$ are energetically

penalized by a soft-wall potential:

510
$$U_{slit} = \frac{1}{2} k_{wall}(|z| - h)^2$$

For each constriction level (typically h =1.0, 0.8, 0.6, 0.4, 0.3), the system undergoes NaN-safe

local energy minimization and extraction of total system energy and principal radii of gyration (Rg1

≤ Rg2 ≤ Rg3). This yields per-sample tables of Energy(h) and Rg(h). For each sample,

deformability metrics were computed from the relaxed (largest slit size) and tight (smallest slit

size) states:

•   Constriction energy cost:

517
$$\Delta E = E_{tight} - E_{relaxed}$$

518   •   Axial deformability (compression along Rg1):

519
$$frac\_Rg1 = \frac{Rg1_{tight}}{Rg1_{relaxed}}$$

520   •   Changes in chromatin anisotropy:

521
$$\Delta Flattening = (\frac{Rg3}{Rg1})_{tight} - (\frac{Rg3}{Rg1})_{relaxed}$$

These metrics quantify whether chromatin becomes shorter, flatter, or more anisotropic under

confinement. Statistical comparisons were performed in R using group-wise ANOVA and Kruskal–

Wallis tests.

scRNAseq

PIPseeker v1.0.0 (Fluent Biosciences) was used to generate single cell matrix, barcode and

features for each sample against Hg38 (human). Separate contig for GFP cDNAs was added to

the genomes to detect malignant cells expressing a reporter. Cell calling, qc (including doublets

removal) and clustering were done using Seurat5[84]. Melanoma cell states were assigned based

on the score of previously established signatures[39, 85] using AUCell[86].

scATAC

The complete processing of fastq from alignment to single cell ATAC peak calling was performed

using cellranger-atac-2.1.0 (https://github.com/10XGenomics/cellranger-atac). Next, scATACseq

cellranger matrix was processed using Signac[87] to analyze the overall ATAC distribution around

TSS, fragments length, number of peaks per cell. Given that, cells with at least 100-3000

fragments, a blacklist ratio < 0.4, a nucleosome signal < 2 and a TSS enrichment > 1 were retained

for downstream analysis. Malignant states were assigned to scATACseq object using Seurat5

FindTransferAnchors function and the scRNAseq matched object as a reference. Cell barcodes

passing the qc and belonging to the assigned states were used to filter the paired fastq files which

were then processed using the HiC analysis described above.

CRISPR/Cas9 screen

sgRNAs counts and dropout enrichment analysis comparing t0 with t21 were conducted using

MAGeCK (v0.5.9.5)[88].

RNAseq

Fastq were trimmed and evaluated for qc using TrimGalore (0.6.6) and fastqc (0.11.7). Then,

trimmed fastq were aligned to Hg38 human genome and annotated using STAR (2.7.7a)[89] using

–quantMode TranscriptomeSAM GeneCounts and –sjdbGTFfile GENCODEv44. Using R (4.4.3),

ReadsPerGene.out.tab were combined in a single table using data.table and processed for

normalization and differential expression analysis using respectively edgeR[90] and DESeq2[83].

Transcript per million (TPM) gene expression normalization was performed with rsem[91] using

Aligned.toTranscriptome.out.bam from STAR alignment as input.

Methylation

Raw IDAT files with 853,307 probes were imported and normalized in several consecutive steps

using *SeSAMe* (version 1.12.9)[92]. A signal intensity-associated P-value was calculated based on

out-of-band array hybridization (*pOOBAH*) and a cutoff of 0.05 was imposed. Background signal

and dye-bias were corrected with *noob* and *dyeBiasCorr,* respectively. The methylation level of

individual CpG sites was determined by calculating the beta (β) value, defined as the ratio of the

fluorescent signal from the methylated allele to the sum of the fluorescent signals of both the

methylated and unmethylated alleles. β values range from 0 (completely unmethylated) to 1.0

(fully methylated). β values were calculated using *getBetas.* Next, HM450 Hg38 probe manifest

was downloaded from TCGAbiolinks[93] and filtered for blacklisted regions. Genomic βvalues of

each cell line were then inferred at bed regions of interests using GenomicRanges[94]. Analysis

was conducted using R (4.4.1).

Mouse experiments

12-273BM GFPLuc cell derived xenografts (CDXs) intended for loHiC were generated by injecting

$1x10^4$ cells in 50% PBS + 50% Matrigel (Corning, #354234) intradermally into 8 weeks old

NOD/SCID/IL2yR- /- male mice (RRID:IMSR_JAX:005557). Tumors were harvested once a

~500mm^3 volume was reached.

For metastatic dissemination studies, $1x10^5$ 501mel GFPLuc were resuspended in 100μl PBS

and instilled by ultrasound-guided (Visualsonics Vevo 770 Ultrasound Imaging System)

intracardiac injection into NOD/SCID/IL2yR- /- male mice 8-12 weeks old. Mouse distress and

body weight were monitored over the course of the experiment until the established endpoint of

~20% weight loss. After euthanasia, mice organs were dissected and imaged for GFP

fluorescence on a Odissey M (LICORbio). GFP foci and signal intensity were quantified with

ImageJ.

All mice were purchased from Jackson Laboratory and maintained in the NYULH animal housing

according to the ethical compliance established by the NYU Institutional Animal Care and Use

Committee (IACUC). All the experiments were approved and conducted under the IACUC

protocol # IA16-00051.

Data mining public datasets

TCGA SKCM transcriptomic data were downloaded from cBioPortal[95], while SKCM

HumanMethylation450 array data were obtained through Bioconductor package TCGAbiolinks.

SKCM patients were classified as Melanocytic, Transitory, Neural-Crest and Undifferentiated

based on a previous study[51]. Analysis was conducted using R (4.4.1).

DepMap Public 25Q2[96] data was downloaded from the official website (https://depmap.org/portal).

scRNAseq expression values from melanoma states in human[85] were generated using a Seurat

object retrieved from a public drive of the Marine Lab

(https://rdr.kuleuven.be/dataset.xhtml?persistentId=doi%3A10.48804/GSAXBN).

ATACseq generated in TCGA tumors was retrieved from XenaBrowser[97].

H3K27ac Hi-ChIP ValidPairs derived from TCGA tumors[98] were downloaded from the NCI

repository (https://gdc.cancer.gov/about-data/publications/TCGA-HiChIP-2024).

Data and Code Availability

Supplementary data, raw data and the code to process the data generated in this work will be

available in Gene Expression Omnibus and in a dedicated Github link after peer reviewed

publication.

Results

Melanoma states exhibit distinct chromatin diffusion, density and domain maturation

At the nanoscale level (~10nm), super-resolution imaging has revealed that chromatin is

organized into CPDs, whose functions are determined by size (50-200nm) and packing scaling

(D_n), which we previously showed influence the probability of a cell to adopt a certain

transcriptional state and nuclear physical properties[16, 62,99]. To investigate potential differences in

CPDs across melanoma cell states as a measure of their nuclear physical properties, we

performed high-resolution imaging on a panel of melanoma cell lines representing three main

phenotypic states: melanocytic (MEL, n = 4), intermediate (INT, n = 3), mesenchymal (MES, n =

4), and of melanocytic cells upon MITF silencing (siMS, n = 2) vs. non-targeting control (siNTC)

(**Fig. 1a**). Given that MITF directly regulates SOX10, siMS simultaneously abrogates the

expression of both master regulators of melanocytic identity, and we tested whether their

concomitant silencing elicits nuclear 3D changes. Partial Wave Spectroscopy (PWS) showed that

MES cells exhibit lower CPD scaling (measured as D_n) and more chromatin diffusion (measured

as D_e) than MEL cells, changes that were only partially recapitulated by siMS (**Fig. 1b**). As

expected, INT cells displayed intermediate values between MEL and MES (**Fig. 1b**). To

investigate if these differences were also observed in patient tissues, we integrated PWS with

Opal multiplexed immunofluorescence (mIF) staining for melanoma cells (S100B+) in melanocytic

(MITF+SOX10+) and mesenchymal (AXL, p-EGFR, TCF4) states in a patient tissue microarray

(TMA) (**Fig. 1c**) containing cores representative of 20 unique primary melanoma cells

(**Supplemental Table 1**). Following MEL and MES cell identification within the TMA (**Fig. 1d**,

Supp Fig. 1a), melanoma cells classified as MES-like showed decreased CPD scaling compared

to MEL-like cells (**Fig. 1e**) suggesting that CPD properties within melanoma cell states are also

found in vivo.

Heterochromatin and euchromatin have been previously linked to higher and lower mechanical

stiffness, respectively[100]. CPD maturation relies on H3K9me3 heterochromatin accelerating

packing in the CPD core, while euchromatin, characterized by higher H3K27ac, stabilizes the

peripheral CPD zone[101]. To gain more insight into CPD maturation related to cell states, we

performed sSMLM for H3K9me3 and H3K27ac in our panel of cell lines. In line with our

hypothesis, CPD maturation is reduced in MES cells, as indicated by a significant decrease in

H3K9me3-positive CPD size and number relative to MEL cells (**Fig. 1f-g**, **Supp Fig. 1b**), findings

further supported by a global genome-wide loss of H3K9me3 by ChIP-seq (**Fig. 1h**). No significant

differences were observed in H3K27ac by sSMLM (**Fig. 1g**, **Supp Fig.1b**), ChIP-seq (**Supp Fig.**

1c) nor upon siMS (**Supp Fig. 1d-e**).

Altogether, we demonstrate that MES cells are characterized by increased chromatin diffusion

and reduced chromatin density, explained by a reduction in CPD number, size, and maturation,

as well as global loss of the H3K9me3 mark, suggesting decreased nuclear stiffness. These

differences are only partially mimicked by MITF and SOX10 silencing (siMS), suggesting that

additional mechanisms are required to fully recapitulate the nuclear topology and biophysical

properties of MES cells during phenotype switching.

Inactive compartments and heterochromatin are reshaped during melanoma phenotypic

switch.

The genome has two major, micron-sized spatial compartments, A and B, associated with active

and inactive chromatin respectively[102, 103] within which CPDs are embedded [16]. Using Cooltools[76],

we examined melanoma cell lines by sub-chromosomal compartments A and B, further validated

by integrating H3K27ac and H3K9me3 ChIPseq, ATACseq, and RNAseq in our panel of cell lines.

As expected, A-type compartments are enriched for H3K27ac, ATAC, and RNA signal compared

to B compartments, which instead are enriched in H3K9me3 (**Supp Fig. 2a**). To identify state-

specific changes in A and B compartments, we performed one minus cosine similarity Kmeans

clustering and found a subset of A compartments specific to each phenotype (MEL and MES; **Fig.**

2a). Although no change in the global proportion of A and B compartments was observed across

states or upon siMS (**Supp. Fig. 2b**), saddle *cis* compartment quantification shows an increase

in B-to-B interactions in INT, MES, and siMS cells relative to MEL and siNTC, respectively (**Fig.**

2b-c). No significant differences were found across *trans* compartments interactions (**Supp Fig.**

2c-d). We analyzed compartment switching from A-to-B and B-to-A between states using a

recently published model[19]. Consistent with a switch, epigenetic and transcriptomics marks were

found significantly changed from A-to-B and B-to-A comparing MEL, INT and MES, but not in siMS

(**Supp Fig. 2e**). We found < 400 compartment switches between MEL and INT, while almost 2000

compartments switched between MEL and MES, predominantly from B-to-A which is triggered as

well by siMS (**Fig. 2d**). This is in line with the observed decrease in H3K9me3 (**Fig. 1h-i**) and

CPDs maturation by sSMLM (**Fig. 1f-g**). In line with our previous observation of decreased

H3K9me3 from MEL to MES, our 3D chromatin data indicate that the melanoma phenotypic

transition from MEL to INT and MES is marked by a B-to-A compartment switch, mimicked by

siMS.

MES cells have a reduced number of topologically associated domains.

To further interrogate 3D chromatin changes associated to cell states, we examined the number,

size and distribution of TADs (contact hubs of genes, promoters, and enhancers) and quantified

insulation scores. We found MES cell lines have significantly fewer (**Fig. 3a***, top*), larger (**Fig. 3a***,*

bottom) TADs than MEL and INT cell lines. However, MITF and SOX silencing (siMS) did not

impact TAD size or number (**Fig. 3a**). Next, TAD activity was evaluated by first averaging the TAD

boundary strength, measured by insulation score. MES cells showed an overall decrease in

insulation and average CTCF occupancy at MEL and INT TADs boundaries (**Fig. 3b, Supp. Fig.**

3a-b), whereas siMS display only decreased CTCF binding at boundaries but no change in

insulation (**Supp. Fig. 3c**). TAD contact frequency was measured using Aggregate Topological

Analysis (ATA), which revealed a global increase of contacts in the inter-TAD surroundings in INT

and MES cells (**Fig. 3c**). In parallel, we observed significantly more inter-TAD contacts between

+1, +2 and +3 neighbor TADs in INT and MES cells than in MEL cells (**Fig. 3d**). siMS had no

effect on intra and inter-TADs contacts (**Supp. Fig. 3d-e**).

Next, we identified loops and filtered for structural loops occupied by CTCF at both anchors.

Notably, INT cells have the highest number of loops compared to other states, while siMS

significantly increased the number of loops relative to siNTC (**Fig. 3e**). Similarly to TADs'

boundaries, CTCF occupancy at loops' anchors is reduced in MES, INT and siMS cells (**Fig. 3f,**

Supp Fig. 3f). Aggregate Peak Analysis (APA) shows an overall enrichment of contacts at the

loops' foreground (#contacts foreground - #contacts background) of the corresponding state,

except for INT cells that have higher activity, mimicked by siMS cells compared to siNTC (**Fig. 3g,**

Supp. Fig. 3g).

Finally, we intersected boundaries and loop anchors to identify unique TADs and loops across

states and in siNTC vs siMS (**Supp. Fig. 3h**), and cross-referenced them with differentially

expressed genes (DEGs) identified by RNAseq (**Fig. 3h**, **Supp. Fig. 3i, Supplemental Table 2**).

Remarkably, only a modest fraction of state-specific TADs were found associated to DEGs for all

states and siMS; i.e., only 1-10% of state-specific TADs result in DEGs (**Fig. 3i**, *left*). Similarly,

structural loops were poorly associated to DEGs; except for siNTC, siMS and MEL, with ~20-30%

of specific loops resulting in DEGs (**Fig. 3i**, *right*). This indicates that differences in TAD and loop

activity between states are disproportional to the changes in gene expression—particularly in

MES cells, suggesting the possibility that changes in chromatin structure during the MES-to-MEL

phenotypic transition are not just responsible for gene expression changes but may confer some

other advantage to MEL cells.

In sum, melanoma dedifferentiation from MEL to MES is associated with enlarged domains and

increased inter-TAD communications, in line with potentially enhanced nuclear and chromatin

flexibility. Intermediate (INT) cells present multi-state loops and thus an overall higher number of

loops, suggesting a 'poised' 3D chromatin state, which is in part recapitulated by siMS.

State-specific chromatin architecture features are conserved *in vivo*.

We have shown that melanoma phenotypic states display specific 3D chromatin architectures that

correlate with nuclear physical properties, irrespective of mutant drivers (i.e., BRAF, NRAS). To

complement these *in vitro* findings, we asked if melanoma states adopt similar state-specific 3D

chromatin architecture *in vivo* using: i) scRNAseq combined with loHiC and ii) bulk HiC on

chromatin from FFPE specimens from melanoma patients. We performed loHiC on a cell-derived

xenograft (CDX) from a patient-derived melanoma brain metastasis short term culture (12-

273BM) [104, 105], which was able to recapitulate the intra-tumor heterogeneity observed in clinical

specimens. MEL and MES states were identified by scRNAseq through Seurat and integrated into

the loHiC Signac clusters using Seurat bridge integration[84] (**Fig. 4a, Supp. Fig. 4a**). Next, we

performed pseudo-bulk HiC analysis on the MEL and MES clusters for which we obtained a good,

genome-wide HiC contact coverage (**Supp. Fig. 4b**), and confirmed their malignant status using

copy number variation analysis (**Supp. Fig. 4c**). Remarkably, we validated most of our in vitro

observations, such as increase in BB compartment interactions (**Fig. 4b, Supp. Fig. 4d**), fewer

and larger TADs (**Fig. 4c**), weaker insulation (**Fig. 4d**) and decreased intra-TAD contacts in MES

vs MEL cell states (**Supp. Fig. 4e**). Similarly, bulk HiC in a small cohort of patients' FFPE tumor

samples broadly classified as MEL or MES based on histopathological characteristics

(**Supplemental Table 3**) showed a trend to a reduction in the number of TADs in MES tumors

(**Fig. 4e**). Overall, these data indicate that state-specific 3D chromatin features are not an artifact

of in vitro culture but an actual feature of melanoma tumors.

Compression during migration induces a MES-like rewiring of nuclear architecture in MEL

cells, suppresses heterochromatin levels, and enhances their metastatic potential.

While mechanical forces are known to modulate chromatin accessibility[31, 32,33] and viceversa[25, 100,]

[106-108], the reciprocal influence of 3D chromatin conformation on nuclear flexibility in the context of

phenotype switching is poorly understood. We have shown that MES cells, compared to MEL

cells, possess a more 'relaxed' nuclear architecture, characterized by a lower number and less

mature of CPDs **(Fig. 1f-g)**, B-to-A compartment switching (**Fig. 2d**), fewer number of TADs with

weaker insulation (**Fig. 3a-b**), and reduced heterochromatin (**Fig. 1g-i**). We hypothesized that this

'loose' nuclear conformation allows MES cells to better withstand the mechanical stress of

invasion or intra/extravasation, and that reduced physical constriction can in turn remodel the 3D

genome towards a pro-metastatic state.

To determine this, we integrated HiC, H3K27ac, and H3K9me3 profiles to model chromosome 1

chromatin as a Hamiltonian coarse-grained polymer bead chain using Polychrom (Open2C).

Here, we simulate angle bending upon artificial scaled compression forces to measure stiffness

and cost of motion of the polymer. The radius of gyration (Rg) measures the extension and

compaction capacity: stiffer regions will adopt extended and rigid conformations increasing Rg,

while softer regions can collapse reducing Rg and additionally facilitate *cis*-long interactions

(**Supp. Fig. 5a**). In line with our hypothesis, our simulation predicts that MES cells have lower

TAD and chromosome stiffness (**Fig. 5a),** as well as reduced energy cost upon simulated

constriction and a trend to increased deformability (**Fig. 5b**, **Supp. Fig. 5b**). To experimentally

prove this finding, MEL and MES cells were seeded in 3μm-pores transwells for 72hrs in complete

media to allow cells to migrate and constrict spontaneously. Next, both non-migrated (Pre = on

top side of the transwell) and migrated (Post = on lower side of the transwell) cells were processed

for HiC, RNAseq, CUT&Tag (H3K27ac, CTCF and H3K9me3), and ATACseq. Post-migration,

MEL cells were associated with gained MES-like features such as lower melanocytic gene

expression (i.e., MITF, SOX10; **Fig. 5c, Supp. Fig. 5c**) reduced H3K9me3 levels (**Fig. 5c-d**),

increased BB compartments interactions (**Fig. 5e, Supp. Fig. 5d-e**), fewer TADs (**Fig. 5f**), and

decreased insulation and CTCF binding at TAD boundaries (**Fig. 5g**In contrast, chromatin and

gene expression of MES cells were only modestly affected (i.e., EGFR, AXL; **Fig. 5c**), except for

an increased melanoma mesenchymal signature, as if migration 'exacerbates' MES cells identity

(**Supp. Fig. 5f**).

Polychrom simulation in Pre and Post cells predicted a significant decrease in stiffness in Post

MEL relative to Pre MEL, but not in Post MES cells relative to its Pre counterpart (**Fig. 5h**).

Consistent with this finding, CPD scaling (D_n) measured by PWS was significantly reduced in Post

cells, particularly in MEL cells (**Fig. 5i**, **Supp Fig. 5g**). In parallel, MEL Post cells displayed a

significant reduction in mature large H3K9me3 CPDs and a concomitant increase in small ones

(**Fig. 5j**) together with a global decrease in H3K9me3 by sSMLM (**Fig. 5k**) which were not

observed in MES Post cells (**Supp Fig. 5h**). While constriction did not change their proliferative

capacity (**Supp. Fig. 5i**), Post MEL cells exhibited increased metastatic potential to the liver upon

intracardiac injection relative to their Pre counterparts (**Fig. 5l, Supp. Fig. 5j**).

Next, we tested whether Post cells were a product of selection of a pre-existing subpopulation in

the initial culture, or whether physical constriction during migration actively results in cells with

Post features. To do that, we infected MEL cells with a doxycycline inducible CBX5-GFP construct

at 0.3 MOI (~ 0.3 plasmid per cell) as a reporter of H3K9me3 content within the cells. Confirming

the value of this construct as a marker of heterochromatin, we observed increased H3K9me3

levels in FACS sorted CBX5-GFP[High] cells compared to CBX5-GFP[Low], and a decrease of these

markers upon post migration (**Fig. 5m**). Flow cytometry on Pre and Post CBX5-GFP[High] cells

showed the emergence of a CBX5-GFP[Low] subpopulation Post migration (**Fig. 5n**). These data

suggest that physical confinement during migration does not passively select but *actively* triggers

a H3K9me3 low subpopulation.

We further tested whether reducing H3K9me3 global content in MEL cells is sufficient to increase

nuclear flexibility and cell migration observed in MES cells. Only 12-24hrs of treatment with the

histone methyltransferase (HMT) SUV39H1/2 inhibitor, Chaetocin increased cell migration (**Fig.**

5o, Supp Fig. 5k) and reduced H3K9me3 without affecting the levels of melanocytic markers

such as MITF and SOX10 (**Supp Fig. 5l**). In line with our hypothesis of heterochromatin being a

physical barrier to migratory constriction, a short pulse of Chaetocin (24hrs) was sufficient to

significantly decrease D_n as a proxy of nuclear stiffness (**Fig. 5p**), and sorted CBX5-GFP^{Low} (low

H3K9me3) cells displayed higher migratory capacity than CBX5-GFP^{High} (high H3K9me3) cells

relative to their non-migrated counterpart (PreM) (**Fig. 5q**).

These data indicate that MEL cells gain MES features as they intravasate and that treatments

that make MEL 3D chromatin ressemble the MES state are sufficient to confer increased

migratory potential before changes in gene expression are observed.

CTCF genomic occupancy reveals a transcriptional activator function specific to

mesenchymal cells.

We observed that MES cells have a global reduction of CTCF at TAD boundaries compared to

MEL cells (**Fig. 3b**), which is in line with an overall CTCF protein decrease in MES (**Supp. Fig.**

6a). However, mean CTCF occupancy at TSSs is higher in MES compared to MEL and INT cells

(**Fig. 6a**), suggesting a CTCF relocation from insulators to regulatory elements. To test this

hypothesis, we first identified all active candidate *cis* regulatory elements (cCREs) by integrating

ATACseq, H3K27ac ChIPseq, RNAseq and HiC using the Activity-by-Contact (ABC) model[82] in

all cell lines. Next, we intersected CTCF peaks with the ABCmodel identified cCREs[82] to define

the proportion of CTCF binding at structural-like (ABC-) vs. enhancer–promoter (ABC+) regions.

This analysis demonstrated a global increase of CTCF binding to cCRE in MES and siMS cells

compared to MEL cells (**Fig. 6b**). In line with the reduced insulator function in MES cells, HiC

pileup map analysis at CTCF peaks shows a strong increase in global contacts as well as between

adjacent 5' and 3' regions (span in graph) in MES cells, indicating a reduced level of conventional

insulation activity compared to MEL, INT and siMS cells (**Fig. 6c-d, Supp. Fig. 6b**). Conversely,

differential binding analysis of CTCF between MEL and MES cells (**Supp. Fig. 6c**) showed

increased occupancy of MES CTCF at cCREs relative to structural-like (CTCF-only) anchors (**Fig.**

6e). These MES-specific CTCF bound regions are co-occupied by MES TFs such as TEAD4 and

FOSL2 (**Fig. 6f**) and associated mostly with enhancers (n = 1,479 ABC cCREs) (**Supp. Fig. 6d**),

suggesting that CTCF may exert a direct gene regulatory function in concert with AP1/TEADs in

MES cells. Accordingly, CTCF and AP1/TEADs co-immunoprecipitated (**Fig. 6g**), further

supporting these TFs binding as part of CTCF complexes. Moreover, TF motif footprint analysis

demonstrates a significant increase in CTCF motif accessibility in MES and siMS compared to

MEL and siNTC (MEL vs MES, Lg_2FC = -0.35251; siNTC vs siMS, Lg_2FC = -0.40685) (**Fig. 6h**).

To identify CTCF regulated genes in MES cells, we looked at the activity and genes directly

contacted by the ABC cCREs overlapping with CTCF MES peaks (**Supp Fig. 6e**), finding 5,081

cCRE-associated genes (**Supplemental Table 4**). We confirmed that putative CTCF-regulated

MES genes are upregulated in MES cell lines[51] and patient samples[85] relative to MEL cells (**Supp.**

Fig. 6f-g). To determine if MES genes depend on CTCF, we performed a brief knockdown of

CTCF (24hrs) using siRNA followed by RNA-seq in MM099 (MES) (**Fig. 6i**, **Supp. Fig. 6h**).

Notably, we found a significant overlap (n = 852) between CTCF ABC-predicted contacted genes

(n= 5,081) and siRNA differentially expressed genes (n = 4,037) (**Supp. Fig. 6i**). Strikingly, CTCF

depletion leads to a significant increase in MEL and decrease in MES gene signatures[53] (**Fig. 6j**),

and siCTCF upregulated and downregulated genes are found significantly associated to MEL and

MES states respectively in patients (**Supp. Fig. 6j**), suggesting that CTCF is indeed required to

maintain MES identity.

Altogether, our data suggest that CTCF partially relocates from TAD boundaries to enhancer–

promoter regions in MES cells, where —together with MES TFs AP1/TEADs— it may directly

regulate a subset of MES relevant GRNs. Therefore, melanoma state transitions can involve

changes in CTCF deposition and activity.

State-specific chromatin hubs are highly active in mesenchymal cells, functionally

involved in invasion and associated with patients' survival.

We demonstrated that melanoma cells states have profound differences in nuclear architecture

irrespective of the mutational driver and burden (i.e., CNVs, SVs). These differences are

associated with changes in both the transcriptional output and physical properties (i.e., stiffness)

tuning metastatic potential. We hypothesized that MES cell increased chromatin flexibility and

inter-domain contacts may favor an increase in cis-CREs (cCREs) connectivity and the formation

of new chromatin hubs. To examine global connectivity, valid paired interactions were divided in

cis-short (distance < 20 Kb), cis-long (distance > 20 Kb), and trans (inter-chromosomal contacts).

To account for DNA mutations/arrangements, we ran EagleC[109] and HiNT[78] to detect structural

variants (SVs) and copy number variations (CNVs) respectively; valid DNA pairs spanning SVs

and CNVs breakpoints were excluded. No significant variations in the number of CNVs and SVs

were detected among states and upon siMS (**Supp Fig. 7a-b**).

In line with our hypothesis, MES cells have a significant increase in cis-long range interactions

compared to INT and MEL cells, and a reduction in *cis*-short range contacts (**Fig. 7a**). However,

siMS treatment does not result in significant variations in global contacts (**Supp. Fig. 7c**).

Moreover, HiC pileup map analysis at global TSS shows an augmented connectivity at promoters

in MES cells (**Fig. 7b**), an indication of enhanced regulatory activity, which is not mimicked by

siMS (**Supp. Fig. 7d**). Next, to map the global cCRE connectivity in melanoma states at the

granular level, we applied the previously described ABC-model[82] to our panel of cell lines, which

identified 215,858 cCREs after excluding self-promoter contacts and retaining cCREs with an

ABCscore $\geq$ 0.02 (**Supplemental Table 5**).

To identify state-specific cCREs, we performed DESeq2 analysis of cCREs based on the

ABCscore among the states to identify 11,581 MEL-specific cCRE, 3,987 INT-specific cCRE and

29,189 MES-specific cCRE (**Fig. 7c, Supp. Fig. 7e**). Within the state-specific cCRE, we identified

chromatin hubs, defined as cCREs able to connect multiple genes concurrently. We ranked

cCREs based on the number of genes contacted, the sum of ABCscore provided and the number

of contacted genes with a TPM expression $\geq$ of 1, identifying hubs specific of each state (**Fig. 7d,**

Supp. Fig. 7f, **Supplemental Table 6**). In line with our hypothesis, MES cells show the greatest

number of hubs compared to INT and MEL. State-specific hub anchors display significantly lower

DNA methylation in the corresponding state, suggesting they are epigenetically active (**Fig. 7e**).

Reduced DNA methylation was also observed at MES hubs in MES-like specimens of both

primary and metastatic melanoma (**Fig. 7f, Supp Fig. 7g**) and this reduction can be detected in

patients' samples[98] (**Fig. 7g, Supp. Fig. 7h-i**), suggesting that these hubs are active in patients'

tumors and from early tumor stages. To further support the in vivo conservation values of the

identified hubs, 12-273BM derived xenograft cells processed with loHiC (**Fig. 4a**) show ATAC

accessibility enrichment at hubs of the corresponding identified states (**Fig. 7h**). Next, DNA

footprinting of hub anchors revealed a significant enrichment in AP1/TEAD transcription factors

as well as CTCF (**Fig. 7i, Supp. Fig 7j, Supplemental Table 7**). Accordingly, MES hub

connectivity and activity are significantly affected by CTCF silencing (**Fig. 7j, Supp. Fig. 7k**),

further supporting CTCF-AP1/TEAD complexes as key regulators of MES specific GRNs.

Finally, we assessed the functional role of MES chromatin hubs by a drop-out screen using a

custom pooled CRISPR interference library against 43 top MES hubs. To test which chromatin

hubs are important for MES identity and migratory capacity, MES cells were transduced with a

doxycycline-inducible dCas9-KRAB-MeCP2 and the CRISPRi library and seeded for migration on

3.0μm pore transwell inserts. Post-migration cells were collected and re-seeded on a transwell

assay multiple times. Twenty-one days after consecutive migration cycles, DNA sequencing

revealed two hubs being significantly depleted by all 5 independent sgRNAs/hub (p-value<0.05)

compared to time zero (t0) (**Fig. 7k**). Strikingly, ATAC-seq accessibility of these two hubs is

significantly associated with shorter patient survival across various cancer types[110] (**Fig. 7l**).

Together, our results precisely map the most active cCREs in the main melanoma phenotypic

states. Particularly in MES cells, it revealed the bivalent function of chromatin architecture in which

CTCF gets recruited at hubs and cCREs of genes involved in MES identity at the expense of

domains' boundaries, simultaneously promoting a more flexible nuclear architecture which favors

cell migration and intravasation.

Discussion

State transitions and phenotypic plasticity have been linked to cancer metastasis and therapeutic

resistance. In particular, the acquisition of a de-differentiated, mesenchymal identity is a common

feature across tumors of various origins, such as carcinomas[111, 112], glioblastoma[113] or

melanoma[114]. To date, most studies have attributed associated transcriptional programs with

confer cancer cells with a selective advantage under multiple stressors as they invade, colonize

distal tissues, or survive treatment. While previous studies have been limited to a two-dimensional

analysis of state-specific epigenomes, we are the first to dissect the contribution of higher order

chromatin structures and nuclear architecture to mesenchymal (MES) identity, using melanoma

as a paradigmatic model of metastasis. Contrary to differentiated MEL state, we demonstrate that

the MES state encompasses a loose chromatin architecture and nuclear conformation which

supports not just a distinct transcriptional and epigenetic program but mechanical properties such

as nuclear deformability as an adaptive response to migratory constriction and metastasis. MES

nuclear topology results in part from a relocation of CTCF from TADs boundaries to MES-specific

chromatin hubs and cCREs, and enhanced cis-long interactions (larger TADs, more inter-TADs

contacts) and reduced H3K9me3 constitutive heterochromatin (small, immature CPDs) relative to

differentiated, MEL cells. MES cells represent an intriguing, de-differentiated cancer cell state

conserved across multiple cancer types[115] . Our data suggests that multiple tumor types from

various cellular origin may similarly converge from a differentiated state, with a 'rigid' chromatin

architecture locked by lineage-specific GRNs, to a pan-cancer undifferentiated (MEL-like) state,

broadly characterized by loose chromatin, nuclear elasticity and enhanced deformation upon

constriction, as we demonstrate in melanoma.

Supporting this model, silencing of the master melanocytic TFs MITF and SOX10 GRNs can

'push' melanoma cells toward an intermediate (INT-like) state as previously reported[46], but this is

insufficient in allowing cells to adopt MES nuclear architecture and mechanical features, which

suggests the requirement of additional stimuli. We showed that upon physical constriction such

as the one experienced during migration, melanocytic cells shift to a MES-like nuclear molecular

conformation, characterized by reduced CTCF occupancy and insulation at TADs' boundaries,

lower heterochromatin and stiffness, and decreased MITF and SOX10 levels, which confers them

with enhanced metastatic potential. This bidirectional crosstalk between 3D chromatin and

nuclear compression links for the first time epigenetic features, gene expression programs and

mechanical properties to cancer phenotypic plasticity and metastasis.

Traditionally, cancer cell phenotypes have been simply defined by differential transcriptional

states[116], where lack of genetic drivers indicates that, at least in part, they are epigenetically

driven[117, 118]. However, the role of epigenetic regulatory mechanisms in defining these

transcriptional states remains poorly understood, partly due to the limited application of in vivo

epigenetic technologies at single-cell resolution (mostly ATACseq)[119]. Although some state-

specific gene' functions clearly map to a corresponding state (e.g., ATF4 in stress response, E2Fs

in mitosis, MMPs in invasion, PD-L1 in immune evasion)[43], many others are harder to interpret

because their expression overlaps across states[116]. Instead, these overlapping genes may reflect

indirect downstream adaptations to upstream epigenetic changes, such as alterations in

chromatin architecture, rather than representing state-specific functional programs. In support of

this, we generated the first atlas of melanoma cell state regulatory element (n=215,858) activity

and their connections with other genes, which can inform future cancer epigenetic-related work.

Here, melanoma states exhibit profound, global differences in 3D chromatin organization that are

not explained solely by differential gene expression[120], but also by distinct nuclear organization

and physical properties *in vitro* and *in vivo*. These results contribute to the ongoing debate on the

role of chromatin architecture in global versus gene-specific transcriptional regulation[121]. Our

findings support a model of converging, transcription-dependent and independent roles of

chromatin architecture, exemplified by the repurposing of CTCF as both a structural and

transcription factor, and the emergence of chromatin hubs as regulatory network organizers.

Untangling CTCF gene regulatory function is demanding due to its polyhedric role as activator,

repressor and genome organization protein[122], as shown in mouse embryonic stem cells where it

acts as activator of a subset of genes together with its paralog CTCFL[123]. In cancer cells, CTCF

can shift from a structural to a gene-regulatory role, as seen in MES cells where it binds less to

insulators and more to CREs, promoting *cis*-long interactions, inter-TAD connectivity and

chromatin hubs formation, regulatory headquarters required for the expression of cancer cell

identity genes[23, 24, 57]. In agreement, a brief depletion of CTCF affects MES hub connectivity and

their target genes, and our CRISPRi screening revealed a previously unreported role of hubs in

controlling MES cells' biology. The two identified hubs control 72 genes involved in cell

proliferation (i.e., NCAPGP2, CTDP1, PARD6G)[124-126], migration (i.e., NFATC1, TJP1)[127, 128] and

survival (UBE3A)[129]. Yet, it is unclear if chromatin hubs solely control gene expression or also

nuclear flexibility. In fact, although physical constriction is sufficient to reduce CTCF at MEL

boundaries, it is insufficient to recruit it at MES CREs and hubs, which may require TF partners[120]

such as AP1/TEADs and CTCFL, all of which have been shown to promote a MES state in

melanoma[47, 130-132].

In addition, we demonstrate that MES cells display a reduced maturation and size of physical

CPDs, which directly correlates with mechanical stability. For example, stable mature domains

have been shown to lower the probability of blebbing during cell migration, with predominantly

'domain-less' chromatin forming the blebs.[34] In addition, mechanical stressors and forces acting

on cancer cells are expected to affect CPDs and their encoded transcriptional memory through a

selective process favoring immature and smaller CPDs (low H3K9me3) rather than mature (high

H3K9me3). Due to their greater steric hindrance, increased crowding upon constriction can

convert mature CPDs into immature ones, destabilizing H3K9me3, CBX5 (HP1a)[13, 14, 133] and

CTCF insulation which is required to contain heterochromatin nucleation[12]. Cell differentiation

requires H3K9me3 deposition to silence lineage unrelated genes expression[134] within mature

CPDs to allow transcriptional memory[16, 101]. Therefore, MES cells may represent a phenotypic

state in which mechanical confinement, coupled with the loss of lineage-determining GRNs (e.g.,

MITF and SOX10 in melanoma), leads to erosion of the transcriptional memory that encodes

tumor lineage identity (i.e., the melanocytic-like state), as reflected by CPD immaturity, increased

long-range cis contacts, TAD expansion, and reduced CTCF-mediated insulation. Based on this

model, cells in the MES state may lose plasticity by efficiently propagating a stabilized epigenetic

and 3D chromatin configuration through cell division, resulting in maintenance of MES identity

during tumor progression. This is consistent with observations in glioblastoma, where the MES

phenotype is highly heritable across cell generations[7]. Similarly, while MEL to MES

dedifferentiation has been extensively reported[51, 135], there is little to no evidence for the opposite

process (i.e., from a de-differentiated to a differentiated state).

Finally, our study provides the first evidence of a direct link between 3D chromatin and nuclear

biophysical properties and cancer cell states in actual tumor samples. By combining state-of-the-

art bulk and single cell high-resolution microscopy (PWS, sSMLM), epigenomics and molecular

and cell biology techniques, we demonstrate their potential for clinical translation. For instance,

we integrate PWS and multiplex IF (Opal) staining -which are relatively inexpensive and rapid

techniques- on a curated TMA containing. Our future work will integrate 3D genome and nuclear-property data across larger cohorts to train AI models on H&E images, enabling machine-learning–based identification of cancer cell states and linking intra-tumor heterogeneity to patient outcomes. Our findings identify nuclear chromatin organization as an essential driver of cancer cell plasticity, representing a novel intersection between mechanical properties, epigenetics and gene expression in tumor progression.

Acknowledgments

This work was supported by U54CA263001 (E.H., I.O and A.W.L.) and R01CA274100 (E.H.), from NCI/NIH. P.B. was awarded of a National Cancer Center (NCC), Melanoma Research Alliance (MRA) and Melanoma Research Foundation (MRF) fellowships. We thank the Genomics Technology Center (GTC), particularly Mr. Paul Zappile, for supervising all the sequencing; the Preclinical Imaging Core (Mr. Orlando Aristizabal), the Experimental Pathology Core (Dr. Cynthia Loomis and Mr. Gyles Ward for Opal panel optimization) and the Flow Cytometry core, all of which are supported by Cancer Center Support Grant (CCSG) from NCI/NIH to the NYULH Perlmutter Cancer Center (P30CA016087; PI: Alec Kimmelman). V.B. was supported by U54 (CA268084) and L.A. by a K23 (1K23DK144661) NIH/NCI awards. This research used resources of the Center for Physical Genomics and Engineering at Northwestern University, and philanthropic support from K. Hudson and R. Goldman, S. Brice and J. Esteve, M. E. Holliday and I. Schneider, the Christina Carinato Charitable Foundation, and D. Sachs. J.S. was supported by R35GM122525, and J.S. and I.A. by P01CA229086 NIH/NCI grants. T.M. was granted a Cancer Research Institute Dr. Keith Landesman Memorial Postdoctoral Fellowship (CRI14497).

Author contributions

1020

Figures

Figure 1: Melanoma cell states are characterized by distinguished chromatin nanoscale
structures.

a. Biorender illustration of the melanoma cell lines classified into states based on our and previous
studies[45].

b. PWS quantification of Chromatin Packing Domain scaling (D_n) and diffusion (D_e) in a large
panel of melanoma cell lines colored as melanocytic (MEL, red), intermediate (INT, grey) and
mesenchymal (MES, green). Each dot represents a single cell, in black is shown the median bar.
One-way ANOVA test p-values are shown and summarized as pairwise comparisons MEL vs
MES.

c. Whole slide snapshot of the Opal staining conducted in two TMAs (A-B) for S100B, SOX10,
MITF, AXL and phospho-EGFR (pEGFR). Scale bars are referred to 5mm size.

d. Representative MEL-like and MES-like cells identified in two melanoma TMAs by confocal
microscopy using a combination of markers to measure PWS in melanoma cell states.

e. Dot plot showing D_n measurements in MEL-like (n=109) and MES-like (n=67) cells in two
melanoma TMAs. Each dot represents a single cell. Student unpaired t-test p-value comparison
is shown.

f. Relative H3K9me3 cluster size in different melanoma cell lines representative of MEL, INT,
MES, siNTC and siMS groups. Each dot is the mean measurement in a single cell. One-way
ANOVA test was conducted. Significant p-value summarizing MEL vs MES is shown.

g. Representative sSMLM images for H3K9me3 and H3K27ac staining in three different
melanoma cell lines colored based on their state of belonging (MEL = red, INT = grey, MES =
green).

h. H3K9me3 ChIP-seq global metaprofile in MEL, INT and MES cells. Each profile is the average

of at least three biological replicates (different cell lines).

i. (Left) Immunoblot for different proteins in a large panel of melanoma cell lines marked in red

(MEL), grey (INT) and green (MES). MITF and EGFR are enriched in MEL/INT and MES cells

respectively. (Right) H3K9me3 relative quantification to total histone H3 based on the blot on the

left. Unpaired t-test was performed for pairwise comparisons. P-values are displayed.

1050

Figure 2: Nuclear compartments switch occurs during phenotype switching.

a. Whole genome *cis* compartments A and B clustering in melanoma cell lines. In red and green

are marked A compartments specific of melanocytic and mesenchymal states respectively. Color

scale represents compartment score z-score normalized by column. Ward hierarchical clustering

dendrogram is shown on the left.

b. States, siNTC and siMS *cis* compartments saddle plots. 20% quantile areas at the corners

were measured (as marked by dotted white squared) as representative of more radical

interactions across A and B.

c. Group histogram of *cis* AA, BB, and AB:BA interactions in states (left), siNTC and siMS (right).

Two-way ANOVA test was used, p-values are displayed.

d. Compartment switch analysis from BA, AB, AA and BB in melanoma states, siNTC and siMS.

Significant switches are shown in black, the number of switch events are displayed in the

corresponding areas.

1064

1065   **Figure 3: Structural domains and loops properties of melanoma cell states.**

a. Column dot plot of the number (top) and size (bottom) of TADs identified in melanoma states, siNTC and siMS. One-way ANOVA significant p-values are shown. Each dot represent a biological replicate.

b. Insulation score (top) and CTCF binding (bottom) profiles of melanoma states compared at melanocytic, intermediate and mesenchymal TAD boundaries. Summit values are quantified and statistically compared in the sided column dot plots. One-way ANOVA test p-values are displayed. Each dot is representative of a biological replicate.

c. Aggregate TAD analysis heatmap showing the relative difference in contacts of MES, INT and siMS cells against MEL/siNTC at melanocytic, intermediate and mesenchymal TADs.

d. Bar dot plot quantifying the *cis* interactions in INT and MES cells relative to MEL cells as intra-TADs (0) and inter-TADs between neighbor TADs (+1, +2, +3). One-way ANOVA test was conducted independently in each TAD group (0, +1, +2, +3). P-values are shown. Each dot represent an independent interaction.

e. Structural loop quantification in melanoma states (left), siNTC and siMS (right). One-way ANOVA and paired t-test were conducted for left and right plots respectively. Dots represent biological replicates.

f. CTCF mean genomic occupancy at melanocytic, intermediate and mesenchymal loops' anchors in melanoma states.

g. HiC contacts frequency at loops anchors (foreground) at melanocytic, intermediate and mesenchymal loops relative to the background (Foreground contacts) in melanoma states, siNTC and siMS. One-way and paired t-test analysis were performed for melanoma states, and siNTC/siMS plots respectively. P-values are shown.

h. Differentially expressed genes (DEGs) heatmap identified in melanoma states. Color bar represents z-score of TPM expression normalized by row. **i**. Bar plot quantification of the

1090 percentage of TADs (left) and loops (right) associated with DEGs that overlap with state-specific

1091 TADs and loops respectively.

1092

Figure 4: Melanoma states 3D chromatin differences are found *in vivo*

a. Seurat UMAP representing the bridge integration of melanoma cell states between scRNAseq

(left) and loHiC (right) obtained from a cell derived xenograft (CDX) of 12-273BM short-term

patients' culture.

b. cis compartment interactions quantified in MEL and MES states identified in CDX tumors.

c. Column dot plot showing the quantification of the number (left) and size (right) of TADs in MEL

and MES states by loHiC.

d. Genome-wide insulation score profile at TADs' boundaries in loHiC MEL and MES states.

e. TADs' number (left) and size (right) quantified in a small cohort of FFPE MEL (n=3) and MES

(n=4) tumors processed with HiC. Unpaired t-test was conducted and resulted p-values are

shown. Each dot represent a patient sample.

1104

Figure 5: Nuclear constriction actively rewire 3D chromatin in a state-specific manner.

a. Column dot plot quantification of chromatin stiffness predicted by polychorm in chromosome 1

(left) and the median stiffness per TAD of chromosome 1 (right) in melanoma states and siMS. P-

values obtained from one-way ANOVA study are shown. Each dot is a biological replicate.

b. Predicted energy cost of chromosome 1 subjected to a simulated constriction gradient. One-

way ANOVA test was conducted for statistical comparison. Significant p-values are shown.

c. Representative immunoblot for multiple targets detected in whole cell protein extracts from

501mel (MEL) and MM099 (MES) Pre and Post migration. ACTB and H3 serves as loading

controls. Relative molecular weights are shown on the right.

d. Bigwig comparison between Pre and Post genomic occupancy at chromosome 1 of H3K9me3

and H3K27ac in 501mel and MM099. Tracks are equally scaled spanning a range displayed on

top left.

e. Grouped histogram quantification of the 20% quantile *cis* compartments AA, BB, AB:BA

interactions in 501mel and MM099 Pre and Post migration. Brackets are shown to highlight BB

comparison between Pre and Post.

f. Quantification of the number (left) and size (right) of TADs in 501mel and MM099, Pre and Post

migration.

g. Insulation score (top) and CTCF occupancy (bottom) profiled in Pre and Post cells at TADs

boundaries of 501mel and MM099 cells.

h. Stiffness quantification of whole chromosome 1 (left) and median TAD stiffness of chromosome

1 (right) in Pre and Post 501mel and MM099 cells. One-way ANOVA test was conducted.

i. D_n measurement by PWS in 501mel and MM099 Pre and Post cells. Unpaired t-test

comparisons between Pre and Post were conducted and significant p-value is shown. Each dot

represents a single cell considered as biological replicate.

j. H3K9me3 domain proportion in 501mel Pre and Post cells. Based on their radii size, domains

were classified in small (≤0.55nm), medium (55-80nm) and large (>80nm). Each dot represents

the average of independent measurements in single cells. Chi-square test of independence was

conducted. Summary p-value is shown.

k. Representative images captured by sSMLM for H3K9me3 and H3K27ac in 501mel Pre and

Post migration.

l. Box dot plot quantification of liver metastasis area (%) derived from NSG animals injected systemically (intracardiac) with 501mel Pre and Post. Each dot represents a biological replicate. Unpaired t-test was performed and p-value is shown.

m. Immunoblot comparing Pre and Post 501mel$^{CBX5-GFP}$ whole cell protein extracts carrying low or high levels of CBX5-GFP. Protein targets and relative molecular weights are shown at left and right respectively. Vinculin and H3 serve as loading controls.

n. CBX5-GFP quantification by flow cytometry (FITC-A) in Pre and Post migration 501mel CBX5-GFP High. Dotted lines serve as visual reference to appreciate the appearance of low and decrease of high CBX5-GFP cells after migration. Three biological replicates are shown.

o. (Left) Representative image of post migration cells upon DMSO or Chaetocin. (Right) Bar dot plot quantification of 501mel migratory cells upon DMSO or Chaetocin (5nM) based on crystal violet staining of transwell. Paired t-test p-value is shown. Each dot represent a biological replicate, and each replicate is the average of three technical replicates.

p. Dot plot quantification of D_n measured by PWS in 501mel cells treated with DMSO (control) or Chaetocin. Each dot represent a single cell. Mean and standard deviations are shown in red. Unpaired t-test was performed, p-value is shown.

q. Relative cell number of post migratory cells to pre migration of 501mel$^{CBX5-GFP}$ High and Low. Each dot is a biological replicate. Paired t-test was conducted, p-value is displayed.

Figure 6: CTCF is a mesenchymal identity transcriptional regulator.

a. Metaprofile occupancy of CTCF at transcription start sites (TSS) in MEL, INT and MES cells.

b. Grouped bar plot representing CTCF occupancy at candidate *cis* regulatory elements (cCRE) or structural-like (CTCF-only) regions in melanoma states, siNTC and siMS. Each dot represents a biological replicate.

c. Aggregate region analysis at melanocytic, intermediate and mesenchymal CTCF peaks in INT and MES cells and their comparison against MEL, measured as Log_2 Fold Change of observed/expected HiC contacts. Span and center areas used for statistical comparisons are marked by dotted black squares.

d. Center and span areas quantification and statistical analysis by one-way ANOVA in MEL, siNTC, siMS, INT and MES cells at melanocytic, intermediate and mesenchymal CTCF peaks. Each dot represents a biological replicate.

e. Peak annotation of CTCF regions specific of MEL and MES states.

f. MITF/SOX10[46] and FOLS2/TEAD4[136] occupancy at CTCF MES-specific regions.

g. Immunoblot of CTCF, FOSL1, FOSL2, JUN and pan-TEAD co-immunoprecipitations using MM099 nuclear insoluble extracts. 1% of input and IgG were used as internal controls. Relative molecular weight are shown on the right.

h. Volcano plots representing the differential binding of transcription factors predicted by TOBIAS footprint analysis. Colored spots are the significantly enriched motifs in the colored matched conditions. Some of the motifs are shown.

i. siCTCF RNAseq volcano plot of DEGs found downregulated or upregulated in MM099. In red are the genes statistically significant (adjusted p-value <0.05).

j. Gene Set Enrichment Analysis plot of Mesenchymal and Melanocytic Widmer signatures in MM099 siCTCF. Adjusted p-values and normalized enrichment scores (NES) are shown.

Figure 7: Mesenchymal cells nuclear organization shape chromatin hubs important for migration and cell identity.

a. Quantification of cis-short (<20Kb), cis-long (>20Kb) and trans HiC contacts by column dot plot in MEL, INT and MES cell lines. One-way ANOVA test was performed, significant p-values are shown. Each dot represents a biological replicate.

b. (Left) HiC pileup region analysis of genome-wide TSS in INT and MES relative to MEL cells. (Right) Corresponding quantification at the center of the heatmap marked by the dotted black square area. Each dot is a biological replicate statistically evaluate by one-way ANOVA.

c. Heatmap of the cCRE differentially active in melanoma states. Color scale measure ABCscore z-score normalized per row.

d. Dot plot distribution of differentially active cCRE in MEL (n=11,581), INT (n=3,987) and MES (n=29,189) cells based on cCRE average sum of ABCscore relative to the number of contacted genes. Color bar measure the number of genes with a TPM expression ≥1 contacted by each cCRE.

e. Methylation quantification of 500 MES hubs in melanoma cell lines. Methylation levels were averaged from all 500 hubs and one-way ANOVA test was conducted.

f. Bar dot plot showing the average methylation of 500 MES hubs in primary melanoma samples (n=100) classified as MEL (melanocytic), mixed and NC/MES (neural-crest, mesenchymal). Each dot represent an independent biological sample, one-way ANOVA p-value comparison MEL vs NC/MES is shown.

g. H3K27ac Hi-ChIP contact aggregate analysis at MES hubs (n=500) in three melanoma patients from the TCGA consortium generated by Yost et al.[98] and classified in INT based on Tsoi et al.[51].

h. Pseudobulk ATACseq enrichment at MEL, INT and MES hubs regions in MEL and MES cells identified by loHiC in 12-273BM derived xenograft.

i. Volcano plot of the differential TF motifs activity in MES vs MEL/INT hubs according to TOBIAS. Some relevant motifs are shown, in red are the statistically significant motifs found.

j. HiC aggregate contact analysis at MES hubs (n=500) in MM099 siNTC and siCTCF.

k. Dot plot representing the CRISPR interfering MES hubs functional ranking in MM099 based on

their fold change enrichment from time zero (T0) to 21 days (T21) upon silencing. In red are

marked significantly dropped targets. GAPDH serves as internal positive control.

l. Kaplan-Meier showing the overall survival of TCGA patients (n=404) based on their levels of

ATACseq accessibility at two MES hubs found significantly depleted by a CRISPR screen. 50%

quartile was used to segregate high (n=202) vs low (n=202) accessibility. Mantel-Cox test p-

values are shown.

1213

Supplemental Figure 1:

a. Snapshot captured by confocal microscopy showing the staining of different markers using

Opal in cells classified as MEL-like and MES-like found in two melanoma TMAs.

b. sSMLM images of H3K9me3 and H3K27ac nuclear distribution in SKMEL-5 (MEL), MM057

(INT) and MM099 (MES).

c. Average H3K27ac distribution genome-wide in MEL, INT, MES. Each profile represents the

average of at least three independent cell lines.

d. H3K9me3 staining in 501mel siNTC and siMS captured by sSMLM.

e. H3K9me3 and H3K27ac ChIP-seq genome-wide distribution in 501mel siNTC and siMS.

1223

Supplemental Figure 2:

a. Normalized genomic occupancy (RPKM) of H3K9me3, H3K27ac, ATAC and RNA at active (A)

and inactive (B) compartments in MEL, INT, MES, siNTC and siMS cells. Standard deviations

bars are based on individual compartment BIN of 100kb spanning the whole genome. Unpaired t

test was performed comparing A vs B per each marker (H3K9me3, H3K27ac, ATAC and RNA).

b. Grouped column bar dot plot of the percentage of A and B compartments in *cis* and *trans*

identified in melanoma states (top) and siNTC vs siMS (bottom). Two-way ANOVA test was

performed to find significant differences. Each dot represents a biological replicate.

c. HiC *trans* compartment saddle plots of melanoma states (top) and siNTC vs siMS (bottom).

Values of 20% quantile areas are shown.

d. Quantification of trans compartment AA, AB:BA and BB interactions shown in panel c. Each

dot is a biological replicate. Unpaired t-test p-value of BB interactions in siNTC and siMS is shown.

e. Column dot plot showing the RPKM enrichment of H3K9me3, H3K27ac, ATAC (left) and RNA

(right) at compartments switching from A to B or B to A found in compartment switch analysis

between states, siNTC and siMS. (Left) Each dot represent a single compartment of 100kb BIN.

(Right) Each dot represent the average of normalized RNA reads, standard deviations bars are

generated by compartments replicates. Unpaired t-test comparison was used. Significant p-

values are displayed.

1242

Supplemental Figure 3:

a. Insulation score metaprofiles at MEL, INT and MES TADs boundaries across all melanoma cell

lines.

b. CTCF occupancy at MEL, INT and MES TADs boundaries in all cell lines. For a-b panels, red,

grey and green colors mark melanocytic, intermediate and mesenchymal cells respectively.

c. Insulation (top) and CTCF (bottom) profiles at siNTC, siMS, MEL, INT and MES TADs

boundaries.

d. HiC aggregate contact analysis of siMS cells at both siNTC and siMS TADs relative to siNTC

cells.

e. Relative intra-TAD (0) and inter-TAD (+1,+2,+3) HiC contacts in siMS relative to siNTC.

f. Mean CTCF distribution in siNTC and siMS cells at MEL, INT, MES, siNTC and siMS loops.

g. HiC aggregate peak analysis at MEL, INT and MES loops anchors in siMS, INT and MES cells

relative to MEL/siNTC cells.

h. Percentage of TADs (left) and loops (right) that are found common (shared) or specific of each

state, siNTC and siMS. **i**. Volcano plot showing the DEGs found upon siMS in 501mel. MITF and

SOX10 are shown. In red are DEGs with a significant adjusted p-value.

1259

Supplemental Figure 4:

a. Signac ATAC accessibility at *MITF* (top) and *EGFR* (bottom) loci in 12-273BM melanoma states

identified by loHiC.

b. HiC contact heatmap from chromosome 1 to X in MEL and MES cells profiled by loHiC. Color

scale represents the log1p of contacts.

c. Copy number variations (CNVs) analysis plot predicted by HiC in MEL and MES cells profiled

by loHiC. Pearson correlation coefficient and p-value comparing CNVs in MEL vs MES are shown.

d. Saddle plot of *cis* compartments in MEL and MES cells identified by loHiC.

e. Bar dot plot of the HiC contacts intra-TAD (o) and inter-TAD (+1,+2,+3) in MES cells relative to

MEL cells. Paired t-test was conducted comparing MEL vs MES at each TAD neighbor.

1270

Supplemental Figure 5:

a. 3D rendering of chromosome 1 modeled as a polymer chain using polychrom and based on

the three principal components. Each bead is a 10kb BIN color scaled based on the predicted

bead average softness. In grey are beads falling into blacklisted regions. The average

chromosome 1 radius of gyration (Rg) per cell line are shown.

b. Quantification of Rg changes upon a simulation of constriction of chr1 polymer chain. One-way

ANOVA test was performed. Each dot represent a biological replicate.

c. Column dot plot quantify the z-score enrichment of melanocytic and mesenchymal Widmer

signatures[53] in 501mel Pre and Post migration. Unpaired t-test comparing Pre and Post for each

signature was performed. Each dot is representative of an independent biological replicate.

d. *cis* compartments saddle plot of Pre and Post 501mel and MM099 cells. 20% quantiles are

shown.

e. Scatter plot of compartment switching between Pre and Post cells in 501mel (top) and MM099

(bottom). In black are highlighted significantly switching compartments.

f. Melanocytic and mesenchymal Widmer signatures enrichment in MM099 Pre and Post

quantified as averaged z-score. Three biological replicates were used. Unpaired statistical t-test

was conducted.

g. Chromatin diffusion (D_e) quantification in 501mel and MM099 Pre and Post migration. Unpaired

t-test was conducted in Pre vs Post and Pre vs Pre pairwise comparisons. Resulted p-values are

shown.

h. Representative images captured by sSMLM for H3K9me3 and H3K27ac staining in MM099

Pre and Post migration.

i. Proliferation curve of 501mel Pre and Post measured as percentage of well confluence from

time 0 till 96hrs. Each dot is the average of three biological replicates. Unpaired t-test was

conducted to compare the end point.

j. Snapshot of GFP fluorescence representing metastatic foci in livers of NSG injected intracardiac

with 501mel Pre (n=7) or Post (n=7).

k. (Left) Representative images of SKMEL-28 (MEL) post migratory cells upon 12hrs of DMSO

(control) or Chaetocin treatment. (Right) SKMEL-28 post migratory relative quantification to the

DMSO. Each dot represent an independent biological replicate. Paired t-test was performed. P-

values is shown.

l. Immunoblot showing H3K9me3, MITF and SOX10 variation upon 24hrs DMSO or Chaetocin

(CHAE) treatments in 501mel and SKMEL-28. H3 and VINCULIN serve as internal loading

controls. Relative molecular weights are shown on the right.

1305

Supplemental Figure 6:

a. (Left) Immunoblot for CTCF, MITF and EGFR across multiple melanoma cell lines used in this

study. Vinculin serve as loading controls. Relative molecular weights are shown on the left. MEL,

INT and MES cells are labeled in red, grey and green respectively. (Right) CTCF protein

quantification based on the immunoblot shown on the right. One-way ANOVA test was used to

assess statistical significance between MEL, INT and MES cells. Each dot represent a protein

band of a biological replicate.

b. Aggregate region analysis of siMS HiC contacts relative to siNTC at CTCF bound regions of

MEL, INT and MES cells. **c**. CTCF state-specific regions identified with Diffbind and profiled for

CTCF in all cell lines as averaged profile (top) and heatmap (bottom).

d. Pie chart representing the percentage of CTCF MES-specific regions (n=1,274) annotated as

CTCF-only (structural), intragenicEnh (intragenic enhancer), intergenicEnh (intergenic enhancer)

or promoter.

e. ABCscore quantified with a column dot plot. Each dot represent the average ABCscore value

of cCREs (n=1,479) identified with the ABC model found to overlap with CTCF MES-specific

regions. One-way ANOVA test was conducted, a summary p-value of the comparison MES vs all

is shown.

f. GSEA enrichment plot of the 5,081 genes found to be contacted by CTCF in MES cells and

scored in Tsoi et al.[51] short-term culture cells classified as MEL (n=17) and MES (n=10). NES and

adjusted p-values are shown.

g. 5,081 genes signature score (ABC_5k) in melanocytic and mesenchymal cells found in

melanoma patients by Pozniak et al.[85] Unpaired t-test p-value is displayed.

h. Immunoblot for CTCF and ACTB (loading control) in whole cell protein extracts of MM099

siNTC and siCTCF cells. Relative molecular weights are shown on the right.

i. Venn diagram of the overlap between significantly DEGs siCTCF and ABC-5k. Hypergeometric

test was conducted, representation factor and p-value are shown.

j. Violin plot of the downregulated (Down) and upregulated (Up) genes upon siCTCF in

melanocytic and mesenchymal melanoma cell states of patients' samples[85]. Unpaired t-test p-

values are shown.

1335

Supplemental Figure 7:

a. Quantification of the number of CNVs as a grouped column plot across MEL, siNTC, siMS, INT

and MES samples. Two-way ANOVA test was conducted.

b. Column dot plot of the number of structural variants (SVs) found in the melanoma cell lines

used in this study. One-way ANOVA test was performed.

c. Fraction of *cis*-short, *cis*-long and *trans* HiC contacts in siNTC and siMS cells. Paired t-test was

used.

d. HiC aggregate region analysis at all TSS in siMS relative to siNTC.

e. PCA plot of the most variable cCREs identified with DESeq2.

f. ABC quantification of the top 500 state-specific cCRE Hubs across all cell lines. Bar plots show

the average of all hubs. Standard error of the means bar are shown. Red, grey and green mark

melanocytic, intermediate and mesenchymal cell lines respectively.

g. Bar dot plot quantify the average methylation of 500 MES hubs in TCGA melanoma patients

(n=470) classified in MEL, INT, NC (neural crest) and MES according to Tsoi[51]. One-way ANOVA

p-value summarizing NC and MES vs INT and MEL comparisons is shown.

h. HiC aggregate pile up heatmap of contacts at MES hubs in FFPE patients' specimen classified

in MEL (n=3) and MES (n=4).

i. Quantification of HiC contacts at the foreground of MEL and MES hubs in MEL and MES

melanoma patients (n=7). Unpaired t-test p-values are shown.

j. CTCF average genomic occupancy at MES hubs (n=500) in individual mesenchymal cell lines.

k. Volcano plot of the differential MES cCREs activity in siCTCF MM099 cells predicted by

RNAseq. In blue are marked the significantly dysregulated cCREs, in red the hubs. Y-axis dotted

line show the significant p-value threshold. On top are marked the number of significantly active

MES hubs (42 down, 136 up). Multiple t-test with Benjamini/Hochberg correction was used.

For panels a,b,c,e,g and j, each dot is representative of a biological replicate.

1361

1362   **Supplementary information and material**

1363   **Supplemental Table 1:** Clinical and molecular information of the tumors' cores of the tissue

1364   microarray.

1365

1366   **Supplemental Table 2:** Differential expression analysis outputs in melanocytic, intermediate,

1367   mesenchymal and siMS. In yellow are marked the significantly positive associated gene to the

1368   corresponding condition.

1369

1370   **Supplemental Table 3:** Clinical and pathological information of the FFPE blocks processed for

1371   HiC.

1372

1373   **Supplemental Table 4:** List of genes contacted by CTCF in MES cells predicted by the ABC

1374   model.

1375

1376   **Supplemental Table 5:** List of all cCREs identified by the ABC model, their target genes and

1377   activity in melanoma cell states.

1378

1379   **Supplemental Table 6:** Ranked list of cCREs found significantly associated to MEL, INT and

1380   MES cells.

1381

1382   **Supplemental Table 7:** Transcription factor motif list analyzed for enrichment at hubs comparing

1383   MES vs MEL and INT cells.

1384

1385   **Supplemental reagents:** Complete list of reagents used in this work.

Bibliography

1. Hanahan D, Weinberg RA. The hallmarks of cancer. Cell. 2000;100(1):57-70. doi: 10.1016/s0092-8674(00)81683-9. PubMed PMID: 10647931.

2. Gupta GP, Massague J. Cancer metastasis: building a framework. Cell. 2006;127(4):679-95. doi: 10.1016/j.cell.2006.11.001. PubMed PMID: 17110329.

3. Lambert AW, Pattabiraman DR, Weinberg RA. Emerging Biological Principles of Metastasis. Cell. 2017;168(4):670-91. doi: 10.1016/j.cell.2016.11.037. PubMed PMID: 28187288; PMCID: PMC5308465.

4. Naxerova K. Evolutionary paths towards metastasis. Nat Rev Cancer. 2025;25(7):545-60. Epub 20250422. doi: 10.1038/s41568-025-00814-x. PubMed PMID: 40263543.

5. Hanahan D. Hallmarks of Cancer: New Dimensions. Cancer Discov. 2022;12(1):31-46. doi: 10.1158/2159-8290.CD-21-1059. PubMed PMID: 35022204.

6. Ciriello G, Magnani L, Aitken SJ, Akkari L, Behjati S, Hanahan D, Landau DA, Lopez-Bigas N, Lupianez DG, Marine JC, Martin-Villalba A, Natoli G, Obenauf AC, Oricchio E, Scaffidi P, Sottoriva A, Swarbrick A, Tonon G, Vanharanta S, Zuber J. Cancer Evolution: A Multifaceted Affair. Cancer Discov. 2024;14(1):36-48. doi: 10.1158/2159-8290.CD-23-0530. PubMed PMID: 38047596; PMCID: PMC10784746.

7. Schiffman JS, D'Avino AR, Prieto T, Pang Y, Fan Y, Rajagopalan S, Potenski C, Hara T, Suva ML, Gawad C, Landau DA. Defining heritability, plasticity, and transition dynamics of cellular phenotypes in somatic evolution. Nat Genet. 2024;56(10):2174-84. Epub 20240924. doi: 10.1038/s41588-024-01920-6. PubMed PMID: 39317739; PMCID: PMC11527590.

8. Esteller M, Dawson MA, Kadoch C, Rassool FV, Jones PA, Baylin SB. The Epigenetic Hallmarks of Cancer. Cancer Discov. 2024;14(10):1783-809. doi: 10.1158/2159-8290.CD-24-0296. PubMed PMID: 39363741.

9. Yan G, Monnier S, Mouelhi M, Dehoux T. Probing molecular crowding in compressed tissues with Brillouin light scattering. Proc Natl Acad Sci U S A. 2022;119(4). doi: 10.1073/pnas.2113614119. PubMed PMID: 35046032; PMCID: PMC8795543.

10. Li Y, Chen M, Hu J, Sheng R, Lin Q, He X, Guo M. Volumetric Compression Induces Intracellular Crowding to Control Intestinal Organoid Growth via Wnt/beta-Catenin Signaling. Cell Stem Cell. 2021;28(1):170-2. doi: 10.1016/j.stem.2020.12.003. PubMed PMID: 33417868.

11. Alric B, Formosa-Dague C, Dague E, Holt LJ, Delarue M. Macromolecular crowding limits growth under pressure. Nat Phys. 2022;18(4):411-6. Epub 20220224. doi: 10.1038/s41567-022-01506-1. PubMed PMID: 37152719; PMCID: PMC10162713.

12. Ong CT, Corces VG. CTCF: an architectural protein bridging genome topology and function. Nat Rev Genet. 2014;15(4):234-46. Epub 20140311. doi: 10.1038/nrg3663. PubMed PMID: 24614316; PMCID: PMC4610363.

13. Miron E, Oldenkamp R, Brown JM, Pinto DMS, Xu CS, Faria AR, Shaban HA, Rhodes JDP, Innocent C, de Ornellas S, Hess HF, Buckle V, Schermelleh L. Chromatin arranges in chains of mesoscale domains with nanoscale functional topography independent of cohesin. Science Advances. 2020;6(39). doi: 10.1126/sciadv.aba8811. PubMed PMID: WOS:000575531700008.

14. Li Y, Eshein A, Virk RKA, Eid A, Wu WL, Frederick J, VanDerway D, Gladstein S, Huang K, Shim AR, Anthony NM, Bauer GM, Zhou X, Agrawal V, Pujadas EM, Jain S, Esteve G, Chandler JE, Nguyen TQ, ..., Backman V. Nanoscale chromatin imaging and analysis platform bridges 4D chromatin organization with molecular function. Science Advances. 2021;7(1). doi: 10.1126/sciadv.abe4310. PubMed PMID: WOS:000605159200040.

15. Li WS, Carter LM, Almassalha LM, Gong R, Pujadas-Liwag LM, Kuo T, MacQuarrie KL, Carignano M, Dunton C, Dravid V, Kanemaki MT, Szleifer I, Backman V. Mature chromatin packing domains persist after RAD21 depletion in 3D. Science Advances. 2025;11:eadp0855.

16. Almassalha LM, Carignano M, Pujadas-Liwag EM, Li WS, Gong R, Acosta N, Dunton C, Gonzalez P, Carter L, Kakkaramadam R, Kröger M, Su P, Frederick J, Yu IC, Medina K, Pritchard

J, Skol A, Nap R, MacQuarrie KL, ..., Backman V. Chromatin conformation, gene transcription, and nucleosome remodeling as an emergent system. Science Advances. 2025;11:eadq6652.

17.	Deng S, Feng Y, Pauklin S. 3D chromatin architecture and transcription regulation in cancer. J Hematol Oncol. 2022;15(1):49. Epub 20220504. doi: 10.1186/s13045-022-01271-x. PubMed PMID: 35509102; PMCID: PMC9069733.

18.	Xu J, Song F, Lyu H, Kobayashi M, Zhang B, Zhao Z, Hou Y, Wang X, Luan Y, Jia B, Stasiak L, Wong JH, Wang Q, Jin Q, Jin Q, Fu Y, Yang H, Hardison RC, Dovat S, ..., Yue F. Subtype-specific 3D genome alteration in acute myeloid leukaemia. Nature. 2022;611(7935):387-98. Epub 20221026. doi: 10.1038/s41586-022-05365-x. PubMed PMID: 36289338; PMCID: PMC10060167.

19.	Narang S, Ghebrechristos Y, Evensen NA, Murrell N, Jasinski S, Ostrow TH, Teachey DT, Raetz EA, Lionnet T, Witkowski M, Aifantis I, Tsirigos A, Carroll WL. Clonal evolution of the 3D chromatin landscape in patients with relapsed pediatric B-cell acute lymphoblastic leukemia. Nat Commun. 2024;15(1):7425. Epub 20240828. doi: 10.1038/s41467-024-51492-6. PubMed PMID: 39198446; PMCID: PMC11358475.

20.	Mathur R, Wang Q, Schupp PG, Nikolic A, Hilz S, Hong C, Grishanina NR, Kwok D, Stevers NO, Jin Q, Youngblood MW, Stasiak LA, Hou Y, Wang J, Yamaguchi TN, Lafontaine M, Shai A, Smirnov IV, Solomon DA, ..., Costello JF. Glioblastoma evolution and heterogeneity from a 3D whole-tumor perspective. Cell. 2024;187(2):446-63 e16. doi: 10.1016/j.cell.2023.12.013. PubMed PMID: 38242087; PMCID: PMC10832360.

21.	Akdemir KC, Le VT, Kim JM, Killcoyne S, King DA, Lin YP, Tian Y, Inoue A, Amin SB, Robinson FS, Nimmakayalu M, Herrera RE, Lynn EJ, Chan K, Seth S, Klimczak LJ, Gerstung M, Gordenin DA, O'Brien J, ..., Andrew Futreal P. Somatic mutation distributions in cancer genomes vary with three-dimensional chromatin structure. Nat Genet. 2020;52(11):1178-88. Epub 20201005. doi: 10.1038/s41588-020-0708-0. PubMed PMID: 33020667; PMCID: PMC8350746.

22.	Dubois F, Sidiropoulos N, Weischenfeldt J, Beroukhim R. Structural variations in cancer and the 3D genome. Nat Rev Cancer. 2022;22(9):533-46. Epub 20220628. doi: 10.1038/s41568-022-00488-9. PubMed PMID: 35764888; PMCID: PMC10423586.

23.	Breves SL, Di Giammartino DC, Nicholson J, Cirigliano S, Mahmood SR, Lee UJ, Martinez-Fundichely A, Jungverdorben J, Singhania R, Rajkumar S, Kirou R, Studer L, Khurana E, Polyzos A, Fine HA, Apostolou E. Three-dimensional regulatory hubs support oncogenic programs in glioblastoma. Mol Cell. 2025;85(7):1330-48 e6. Epub 20250326. doi: 10.1016/j.molcel.2025.03.007. PubMed PMID: 40147440; PMCID: PMC12009607.

24.	Gambi G, Boccalatte F, Rodriguez Hernaez J, Lin Z, Nadorp B, Polyzos A, Tan J, Avrampou K, Inghirami G, Kentsis A, Apostolou E, Aifantis I, Tsirigos A. 3D chromatin hubs as regulatory units of identity and survival in human acute leukemia. Mol Cell. 2025;85(1):42-60 e7. Epub 20241223. doi: 10.1016/j.molcel.2024.11.040. PubMed PMID: 39719705; PMCID: PMC11934262.

25.	Strom AR, Biggs RJ, Banigan EJ, Wang X, Chiu K, Herman C, Collado J, Yue F, Ritland Politz JC, Tait LJ, Scalzo D, Telling A, Groudine M, Brangwynne CP, Marko JF, Stephens AD. HP1alpha is a chromatin crosslinker that controls nuclear and mitotic chromosome mechanics. Elife. 2021;10. Epub 20210609. doi: 10.7554/eLife.63972. PubMed PMID: 34106828; PMCID: PMC8233041.

26.	Heo SJ, Driscoll TP, Thorpe SD, Nerurkar NL, Baker BM, Yang MT, Chen CS, Lee DA, Mauck RL. Differentiation alters stem cell nuclear architecture, mechanics, and mechano-sensitivity. Elife. 2016;5. Epub 20161130. doi: 10.7554/eLife.18207. PubMed PMID: 27901466; PMCID: PMC5148611.

27.	Patta I, Zand M, Lee L, Mishra S, Bortnick A, Lu H, Prusty A, McArdle S, Mikulski Z, Wang HY, Cheng CS, Fisch KM, Hu M, Murre C. Nuclear morphology is shaped by loop-extrusion programs. Nature. 2024;627(8002):196-203. Epub 20240214. doi: 10.1038/s41586-024-07086-9. PubMed PMID: 38355805; PMCID: PMC11052650.

28.   Scott AK, Rafuse M, Neu CP. Mechanically induced alterations in chromatin architecture
guide the balance between cell plasticity and mechanical memory. Front Cell Dev Biol.
2023;11:1084759. Epub 20230418. doi: 10.3389/fcell.2023.1084759. PubMed PMID: 37143893;
PMCID: PMC10151697.
29.   Shivashankar GV. Mechanical forces and the 3D genome. Curr Opin Struct Biol.
2023;83:102728. Epub 20231109. doi: 10.1016/j.sbi.2023.102728. PubMed PMID: 37948897.
30.   Denais CM, Gilbert RM, Isermann P, McGregor AL, te Lindert M, Weigelin B, Davidson
PM, Friedl P, Wolf K, Lammerding J. Nuclear envelope rupture and repair during cancer cell
migration. Science. 2016;352(6283):353-8. Epub 20160324. doi: 10.1126/science.aad7297.
PubMed PMID: 27013428; PMCID: PMC4833568.
31.   Shroff NP, Xu P, Kim S, Shelton ER, Gross BJ, Liu Y, Gomez CO, Ye Q, Drennon TY, Hu
JK, Green JBA, Campas O, Klein OD. Proliferation-driven mechanical compression induces
signalling centre formation during mammalian organ development. Nat Cell Biol. 2024;26(4):519-
29. Epub 20240403. doi: 10.1038/s41556-024-01380-4. PubMed PMID: 38570617.
32.   Tse JM, Cheng G, Tyrrell JA, Wilcox-Adelman SA, Boucher Y, Jain RK, Munn LL.
Mechanical compression drives cancer cells toward invasive phenotype. Proc Natl Acad Sci U S
A. 2012;109(3):911-6. Epub 20111227. doi: 10.1073/pnas.1118910109. PubMed PMID:
22203958; PMCID: PMC3271885.
33.   Dupont S, Wickstrom SA. Mechanical regulation of chromatin and transcription. Nat Rev
Genet. 2022;23(10):624-43. Epub 20220523. doi: 10.1038/s41576-022-00493-6. PubMed PMID:
35606569.
34.   Pujadas Liwag EM, Acosta N, Almassalha LM, Su YP, Gong R, Kanemaki MT, Stephens
AD, Backman V. Nuclear blebs are associated with destabilized chromatin-packing domains. J
Cell Sci. 2025;138(3). Epub 20250211. doi: 10.1242/jcs.262161. PubMed PMID: 39878045;
PMCID: PMC11883274.
35.   Gupta PB, Onder TT, Jiang G, Tao K, Kuperwasser C, Weinberg RA, Lander ES.
Identification of selective inhibitors of cancer stem cells by high-throughput screening. Cell.
2009;138(4):645-59. Epub 20090813. doi: 10.1016/j.cell.2009.06.034. PubMed PMID: 19682730;
PMCID: PMC4892125.
36.   Rhim AD, Mirek ET, Aiello NM, Maitra A, Bailey JM, McAllister F, Reichert M, Beatty GL,
Rustgi AK, Vonderheide RH, Leach SD, Stanger BZ. EMT and dissemination precede pancreatic
tumor formation. Cell. 2012;148(1-2):349-61. doi: 10.1016/j.cell.2011.11.025. PubMed PMID:
22265420; PMCID: PMC3266542.
37.   Yu M, Bardia A, Wittner BS, Stott SL, Smas ME, Ting DT, Isakoff SJ, Ciciliano JC, Wells
MN, Shah AM, Concannon KF, Donaldson MC, Sequist LV, Brachtel E, Sgroi D, Baselga J,
Ramaswamy S, Toner M, Haber DA, Maheswaran S. Circulating breast tumor cells exhibit
dynamic changes in epithelial and mesenchymal composition. Science. 2013;339(6119):580-4.
doi: 10.1126/science.1228522. PubMed PMID: 23372014; PMCID: PMC3760262.
38.   Pastushenko I, Brisebarre A, Sifrim A, Fioramonti M, Revenco T, Boumahdi S, Van
Keymeulen A, Brown D, Moers V, Lemaire S, De Clercq S, Minguijon E, Balsat C, Sokolow Y,
Dubois C, De Cock F, Scozzaro S, Sopena F, Lanas A, D'Haene N, Salmon I, Marine JC, Voet T,
Sotiropoulou PA, Blanpain C. Identification of the tumour transition states occurring during EMT.
Nature. 2018;556(7702):463-8. Epub 20180418. doi: 10.1038/s41586-018-0040-3. PubMed
PMID: 29670281.
39.   Karras P, Bordeu I, Pozniak J, Nowosad A, Pazzi C, Van Raemdonck N, Landeloos E, Van
Herck Y, Pedri D, Bervoets G, Makhzami S, Khoo JH, Pavie B, Lamote J, Marin-Bejar O, Dewaele
M, Liang H, Zhang X, Hua Y, Wouters J, Browaeys R, Bergers G, Saeys Y, Bosisio F, van den
Oord J, Lambrechts D, Rustgi AK, Bechter O, Blanpain C, Simons BD, Rambow F, Marine JC. A
cellular hierarchy in melanoma uncouples growth and metastasis. Nature. 2022;610(7930):190-
8. Epub 20220921. doi: 10.1038/s41586-022-05242-7. PubMed PMID: 36131018; PMCID:
PMC10439739.

40.   Bajaj S, Donnelly D, Call M, Johannet P, Moran U, Polsky D, Shapiro R, Berman R, Pavlick
A, Weber J, Zhong J, Osman I. Melanoma Prognosis: Accuracy of the American Joint Committee
on Cancer Staging Manual Eighth Edition. J Natl Cancer Inst. 2020;112(9):921-8. doi:
10.1093/jnci/djaa008. PubMed PMID: 31977051; PMCID: PMC7492758.
41.   Sun Q, Lee W, Mohri Y, Takeo M, Lim CH, Xu X, Myung P, Atit RP, Taketo MM, Moubarak
RS, Schober M, Osman I, Gay DL, Saur D, Nishimura EK, Ito M. A novel mouse model
demonstrates that oncogenic melanocyte stem cells engender melanoma resembling human
disease. Nat Commun. 2019;10(1):5023. Epub 20191104. doi: 10.1038/s41467-019-12733-1.
PubMed PMID: 31685822; PMCID: PMC6828673.
42.   Vallius T, Shi Y, Novikov E, Pant SM, Pelletier R, Chen YA, Tefft JB, Johnson AN, Maliga
Z, Wan G, Murphy G, Santagata S, Semenov YR, Liu D, Lian CG, Sorger PK. Spatial determinants
of tumor cell dedifferentiation and plasticity in primary cutaneous melanoma. bioRxiv. 2025. Epub
20250624. doi: 10.1101/2025.06.21.660851. PubMed PMID: 40667360; PMCID: PMC12262277.
43.   Rambow F, Marine JC, Goding CR. Melanoma plasticity and phenotypic diversity:
therapeutic barriers and opportunities. Genes Dev. 2019;33(19-20):1295-318. doi:
10.1101/gad.329771.119. PubMed PMID: 31575676; PMCID: PMC6771388.
44.   Hoek KS, Goding CR. Cancer stem cells versus phenotype-switching in melanoma.
Pigment Cell Melanoma Res. 2010;23(6):746-59. Epub 20100820. doi: 10.1111/j.1755-
148X.2010.00757.x. PubMed PMID: 20726948.
45.   Wouters J, Kalender-Atak Z, Minnoye L, Spanier KI, De Waegeneer M, Bravo Gonzalez-
Blas C, Mauduit D, Davie K, Hulselmans G, Najem A, Dewaele M, Pedri D, Rambow F, Makhzami
S, Christiaens V, Ceyssens F, Ghanem G, Marine JC, Poovathingal S, Aerts S. Robust gene
expression programs underlie recurrent cell states and phenotype switching in melanoma. Nat
Cell Biol. 2020;22(8):986-98. Epub 20200803. doi: 10.1038/s41556-020-0547-3. PubMed PMID:
32753671.
46.   Laurette P, Strub T, Koludrovic D, Keime C, Le Gras S, Seberg H, Van Otterloo E,
Imrichova H, Siddaway R, Aerts S, Cornell RA, Mengus G, Davidson I. Transcription factor MITF
and remodeller BRG1 define chromatin organisation at regulatory elements in melanoma cells.
Elife. 2015;4. Epub 20150324. doi: 10.7554/eLife.06857. PubMed PMID: 25803486; PMCID:
PMC4407272.
47.   Verfaillie A, Imrichova H, Atak ZK, Dewaele M, Rambow F, Hulselmans G, Christiaens V,
Svetlichnyy D, Luciani F, Van den Mooter L, Claerhout S, Fiers M, Journe F, Ghanem GE,
Herrmann C, Halder G, Marine JC, Aerts S. Decoding the regulatory landscape of melanoma
reveals TEADS as regulators of the invasive cell state. Nat Commun. 2015;6:6683. Epub
20150409. doi: 10.1038/ncomms7683. PubMed PMID: 25865119; PMCID: PMC4403341.
48.   Tirosh I, Izar B, Prakadan SM, Wadsworth MH, 2nd, Treacy D, Trombetta JJ, Rotem A,
Rodman C, Lian C, Murphy G, Fallahi-Sichani M, Dutton-Regester K, Lin JR, Cohen O, Shah P,
Lu D, Genshaft AS, Hughes TK, Ziegler CG, ..., Garraway LA. Dissecting the multicellular
ecosystem of metastatic melanoma by single-cell RNA-seq. Science. 2016;352(6282):189-96.
doi: 10.1126/science.aad0501. PubMed PMID: 27124452; PMCID: PMC4944528.
49.   Rambow F, Rogiers A, Marin-Bejar O, Aibar S, Femel J, Dewaele M, Karras P, Brown D,
Chang YH, Debiec-Rychter M, Adriaens C, Radaelli E, Wolter P, Bechter O, Dummer R, Levesque
M, Piris A, Frederick DT, Boland G, ..., Marine JC. Toward Minimal Residual Disease-Directed
Therapy in Melanoma. Cell. 2018;174(4):843-55 e19. Epub 20180712. doi:
10.1016/j.cell.2018.06.025. PubMed PMID: 30017245.
50.   Marin-Bejar O, Rogiers A, Dewaele M, Femel J, Karras P, Pozniak J, Bervoets G, Van
Raemdonck N, Pedri D, Swings T, Demeulemeester J, Borght SV, Lehnert S, Bosisio F, van den
Oord JJ, Bempt IV, Lambrechts D, Voet T, Bechter O, ..., Marine JC. Evolutionary predictability of
genetic versus nongenetic resistance to anticancer drugs in melanoma. Cancer Cell.
2021;39(8):1135-49 e8. Epub 20210617. doi: 10.1016/j.ccell.2021.05.015. PubMed PMID:
34143978.

51.   Tsoi J, Robert L, Paraiso K, Galvan C, Sheu KM, Lay J, Wong DJL, Atefi M, Shirazi R,
Wang X, Braas D, Grasso CS, Palaskas N, Ribas A, Graeber TG. Multi-stage Differentiation
Defines Melanoma Subtypes with Differential Vulnerability to Drug-Induced Iron-Dependent
Oxidative Stress. Cancer Cell. 2018;33(5):890-904 e5. Epub 20180412. doi:
10.1016/j.ccell.2018.03.017. PubMed PMID: 29657129; PMCID: PMC5953834.
52.   Hoek KS, Eichhoff OM, Schlegel NC, Dobbeling U, Kobert N, Schaerer L, Hemmi S,
Dummer R. In vivo switching of human melanoma cells between proliferative and invasive states.
Cancer Res. 2008;68(3):650-6. doi: 10.1158/0008-5472.CAN-07-2491. PubMed PMID:
18245463.
53.   Widmer DS, Cheng PF, Eichhoff OM, Belloni BC, Zipser MC, Schlegel NC, Javelaud D,
Mauviel A, Dummer R, Hoek KS. Systematic classification of melanoma cells by phenotype-
specific gene expression mapping. Pigment Cell Melanoma Res. 2012;25(3):343-53. Epub
20120302. doi: 10.1111/j.1755-148X.2012.00986.x. PubMed PMID: 22336146.
54.   Schmidl C, Rendeiro AF, Sheffield NC, Bock C. ChIPmentation: fast, robust, low-input
ChIP-seq for histones and transcription factors. Nat Methods. 2015;12(10):963-5. Epub
20150817. doi: 10.1038/nmeth.3542. PubMed PMID: 26280331; PMCID: PMC4589892.
55.   Conway K, Edmiston SN, Vondras A, Reiner A, Corcoran DL, Shen R, Parrish EA, Hao H,
Lin L, Kenney JM, Ilelaboye G, Kostrzewa CE, Kuan PF, Busam KJ, Lezcano C, Lee TK,
Hernando E, Googe PB, Ollila DW, Moschos S, Gorlov I, Amos CI, Ernstoff MS, Cust AE, Wilmott
JS, Scolyer RA, Mann GJ, Vergara IA, Ko J, Rees JR, Yan S, Nagore E, Bosenberg M, Rothberg
BG, Osman I, Lee JE, Saenger Y, Bogner P, Thompson CL, Gerstenblith M, Holmen SL, Funchain
P, Brunsgaard E, Depcik-Smith ND, Luo L, Boyce T, Orlow I, Begg CB, Berwick M, Thomas NE,
Inter MELSG. DNA Methylation Classes of Stage II and III Primary Melanomas and Their Clinical
and Prognostic Significance. JCO Precis Oncol. 2024;8:e2400375. Epub 20241107. doi:
10.1200/PO-24-00375. PubMed PMID: 39509669; PMCID: PMC11737429.
56.   Chang L, Xie Y, Taylor B, Wang Z, Sun J, Armand EJ, Mishra S, Xu J, Tastemel M, Lie A,
Gibbs ZA, Indralingam HS, Tan TM, Bejar R, Chen CC, Furnari FB, Hu M, Ren B. Droplet Hi-
C enables scalable, single-cell profiling of chromatin architecture in heterogeneous tissues. Nat
Biotechnol. 2025;43(10):1694-707. Epub 20241018. doi: 10.1038/s41587-024-02447-1. PubMed
PMID: 39424717; PMCID: PMC12520981.
57.   Sakellaropoulos T, Do C, Jiang G, Cova G, Meyn P, Dimartino D, Ramaswami S, Heguy
A, Tsirigos A, Skok JA. MethNet: a robust approach to identify regulatory hubs and their distal
targets from cancer data. Nat Commun. 2024;15(1):6027. Epub 20240717. doi: 10.1038/s41467-
024-50380-3. PubMed PMID: 39025865; PMCID: PMC11258126.
58.   Doench JG, Fusi N, Sullender M, Hegde M, Vaimberg EW, Donovan KF, Smith I, Tothova
Z, Wilen C, Orchard R, Virgin HW, Listgarten J, Root DE. Optimized sgRNA design to maximize
activity and minimize off-target effects of CRISPR-Cas9. Nat Biotechnol. 2016;34(2):184-91.
Epub 20160118. doi: 10.1038/nbt.3437. PubMed PMID: 26780180; PMCID: PMC4744125.
59.   Gladstein S, Stawarz A, Almassalha LM, Cherkezyan L, Chandler JE, Zhou X,
Subramanian H, Backman V. Measuring Nanoscale Chromatin Heterogeneity with Partial Wave
Spectroscopic Microscopy. Methods Mol Biol. 2018;1745:337-60. doi: 10.1007/978-1-4939-7680-
5_19. PubMed PMID: 29476478; PMCID: PMC5983358.
60.   Almassalha LM, Bauer GM, Chandler JE, Gladstein S, Cherkezyan L, Stypula-Cyrus Y,
Weinberg S, Zhang D, Thusgaard Ruhoff P, Roy HK, Subramanian H, Chandel NS, Szleifer I,
Backman V. Label-free imaging of the native, living cellular nanoarchitecture using partial-wave
spectroscopic microscopy. Proc Natl Acad Sci U S A. 2016;113(42):E6372-E81. Epub 20161004.
doi: 10.1073/pnas.1608198113. PubMed PMID: 27702891; PMCID: PMC5081614.
61.   Gladstein S, Almassalha LM, Cherkezyan L, Chandler JE, Eshein A, Eid A, Zhang D, Wu
W, Bauer GM, Stephens AD, Morochnik S, Subramanian H, Marko JF, Ameer GA, Szleifer I,
Backman V. Multimodal interference-based imaging of nanoscale structure and macromolecular

motion uncovers UV induced cellular paroxysm. Nat Commun. 2019;10(1):1652. Epub 20190410. doi: 10.1038/s41467-019-09717-6. PubMed PMID: 30971691; PMCID: PMC6458150.

62.   Virk RKA, Wu W, Almassalha LM, Bauer GM, Li Y, VanDerway D, Frederick J, Zhang D, Eshein A, Roy HK, Szleifer I, Backman V. Disordered chromatin packing regulates phenotypic plasticity. Sci Adv. 2020;6(2):eaax6232. Epub 20200108. doi: 10.1126/sciadv.aax6232. PubMed PMID: 31934628; PMCID: PMC6949045.

63.   Eid A, Eshein A, Li Y, Virk R, Van Derway D, Zhang D, Taflove A, Backman V. Characterizing chromatin packing scaling in whole nuclei using interferometric microscopy. Opt Lett. 2020;45(17):4810-3. doi: 10.1364/OL.400231. PubMed PMID: 32870863; PMCID: PMC7951997.

64.   Pujadas Liwag EM, Wei X, Acosta N, Carter LM, Yang J, Almassalha LM, Jain S, Daneshkhah A, Rao SSP, Seker-Polat F, MacQuarrie KL, Ibarra J, Agrawal V, Aiden EL, Kanemaki MT, Backman V, Adli M. Depletion of lamins B1 and B2 promotes chromatin mobility and induces differential gene expression by a mesoscale-motion-dependent mechanism. Genome Biol. 2024;25(1):77. Epub 20240322. doi: 10.1186/s13059-024-03212-y. PubMed PMID: 38519987; PMCID: PMC10958841.

65.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9(4):357-9. Epub 20120304. doi: 10.1038/nmeth.1923. PubMed PMID: 22388286; PMCID: PMC3322381.

66.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project Data Processing S. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. Epub 20090608. doi: 10.1093/bioinformatics/btp352. PubMed PMID: 19505943; PMCID: PMC2723002.

67.   Ramirez F, Dundar F, Diehl S, Gruning BA, Manke T. deepTools: a flexible platform for exploring deep-sequencing data. Nucleic Acids Res. 2014;42(Web Server issue):W187-91. Epub 20140505. doi: 10.1093/nar/gku365. PubMed PMID: 24799436; PMCID: PMC4086134.

68.   Liu T. Use model-based Analysis of ChIP-Seq (MACS) to analyze short reads generated by sequencing protein-DNA interactions in embryonic stem cells. Methods Mol Biol. 2014;1150:81-95. doi: 10.1007/978-1-4939-0512-6_4. PubMed PMID: 24743991.

69.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics. 2010;26(6):841-2. Epub 20100128. doi: 10.1093/bioinformatics/btq033. PubMed PMID: 20110278; PMCID: PMC2832824.

70.   Sandelin A, Alkema W, Engstrom P, Wasserman WW, Lenhard B. JASPAR: an open-access database for eukaryotic transcription factor binding profiles. Nucleic Acids Res. 2004;32(Database issue):D91-4. doi: 10.1093/nar/gkh012. PubMed PMID: 14681366; PMCID: PMC308747.

71.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK. Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. Mol Cell. 2010;38(4):576-89. doi: 10.1016/j.molcel.2010.05.004. PubMed PMID: 20513432; PMCID: PMC2898526.

72.   Machanick P, Bailey TL. MEME-ChIP: motif analysis of large DNA datasets. Bioinformatics. 2011;27(12):1696-7. Epub 20110412. doi: 10.1093/bioinformatics/btr189. PubMed PMID: 21486936; PMCID: PMC3106185.

73.   Ross-Innes CS, Stark R, Teschendorff AE, Holmes KA, Ali HR, Dunning MJ, Brown GD, Gojis O, Ellis IO, Green AR, Ali S, Chin SF, Palmieri C, Caldas C, Carroll JS. Differential oestrogen receptor binding is associated with clinical outcome in breast cancer. Nature. 2012;481(7381):389-93. Epub 20120104. doi: 10.1038/nature10730. PubMed PMID: 22217937; PMCID: PMC3272464.

74.   Servant N, Varoquaux N, Lajoie BR, Viara E, Chen CJ, Vert JP, Heard E, Dekker J, Barillot E. HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. Genome Biol.

2015;16:259. Epub 20151201. doi: 10.1186/s13059-015-0831-x. PubMed PMID: 26619908; PMCID: PMC4665391.

75.   Berico P, Flores Yanke AF, Vand Rajabpour F, Do C, Simonin Wilmer I, Sakellaropoulos T, Delclaux Obieta I, Stagnitta R, Vazquez Cruz E, Osman I, Skok J, Robles-Espinoza CD, Lund AW, Schober M, Hernando E. A targetable developmental program co-regulates angiogenesis and immune evasion. bioRxiv. 2024. Epub 20241221. doi: 10.1101/2024.12.21.628442. PubMed PMID: 39764043; PMCID: PMC11702693.

76.   Open2C, Abdennur N, Abraham S, Fudenberg G, Flyamer IM, Galitsyna AA, Goloborodko A, Imakaev M, Oksuz BA, Venev SV, Xiao Y. Cooltools: Enabling high-resolution Hi-C analysis in Python. PLoS Comput Biol. 2024;20(5):e1012067. Epub 20240506. doi: 10.1371/journal.pcbi.1012067. PubMed PMID: 38709825; PMCID: PMC11098495.

77.   Durand NC, Shamim MS, Machol I, Rao SS, Huntley MH, Lander ES, Aiden EL. Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments. Cell Syst. 2016;3(1):95-8. doi: 10.1016/j.cels.2016.07.002. PubMed PMID: 27467249; PMCID: PMC5846465.

78.   Wang S, Lee S, Chu C, Jain D, Kerpedjiev P, Nelson GM, Walsh JM, Alver BH, Park PJ. HiNT: a computational method for detecting copy number variations and translocations from Hi-C data. Genome Biol. 2020;21(1):73. Epub 20200323. doi: 10.1186/s13059-020-01986-5. PubMed PMID: 32293513; PMCID: PMC7087379.

79.   Forcato M, Nicoletti C, Pal K, Livi CM, Ferrari F, Bicciato S. Comparison of computational methods for Hi-C data analysis. Nat Methods. 2017;14(7):679-85. Epub 20170612. doi: 10.1038/nmeth.4325. PubMed PMID: 28604721; PMCID: PMC5493985.

80.   Salameh TJ, Wang X, Song F, Zhang B, Wright SM, Khunsriraksakul C, Ruan Y, Yue F. A supervised learning framework for chromatin loop detection in genome-wide contact maps. Nat Commun. 2020;11(1):3428. Epub 20200709. doi: 10.1038/s41467-020-17239-9. PubMed PMID: 32647330; PMCID: PMC7347923.

81.   van der Weide RH, van den Brand T, Haarhuis JHI, Teunissen H, Rowland BD, de Wit E. Hi-C analyses with GENOVA: a case study with cohesin variants. NAR Genom Bioinform. 2021;3(2):lqab040. Epub 20210522. doi: 10.1093/nargab/lqab040. PubMed PMID: 34046591; PMCID: PMC8140737.

82.   Fulco CP, Nasser J, Jones TR, Munson G, Bergman DT, Subramanian V, Grossman SR, Anyoha R, Doughty BR, Patwardhan TA, Nguyen TH, Kane M, Perez EM, Durand NC, Lareau CA, Stamenova EK, Aiden EL, Lander ES, Engreitz JM. Activity-by-contact model of enhancer-promoter regulation from thousands of CRISPR perturbations. Nat Genet. 2019;51(12):1664-9. Epub 20191129. doi: 10.1038/s41588-019-0538-0. PubMed PMID: 31784727; PMCID: PMC6886585.

83.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550. doi: 10.1186/s13059-014-0550-8. PubMed PMID: 25516281; PMCID: PMC4302049.

84.   Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, Srivastava A, Molla G, Madad S, Fernandez-Granda C, Satija R. Dictionary learning for integrative, multimodal and scalable single-cell analysis. Nat Biotechnol. 2024;42(2):293-304. Epub 20230525. doi: 10.1038/s41587-023-01767-y. PubMed PMID: 37231261; PMCID: PMC10928517.

85.   Pozniak J, Pedri D, Landeloos E, Van Herck Y, Antoranz A, Vanwynsberghe L, Nowosad A, Roda N, Makhzami S, Bervoets G, Maciel LF, Pulido-Vicuna CA, Pollaris L, Seurinck R, Zhao F, Flem-Karlsen K, Damsky W, Chen L, Karagianni D, Cinque S, Kint S, Vandereyken K, Rombaut B, Voet T, Vernaillen F, Annaert W, Lambrechts D, Boecxstaens V, Saeys Y, van den Oord J, Bosisio F, Karras P, Shain AH, Bosenberg M, Leucci E, Paschen A, Rambow F, Bechter O, Marine JC. A TCF4-dependent gene regulatory network confers resistance to immunotherapy in melanoma. Cell. 2024;187(1):166-83 e25. doi: 10.1016/j.cell.2023.11.037. PubMed PMID: 38181739.

86.   Aibar S, Gonzalez-Blas CB, Moerman T, Huynh-Thu VA, Imrichova H, Hulselmans G,
Rambow F, Marine JC, Geurts P, Aerts J, van den Oord J, Atak ZK, Wouters J, Aerts S. SCENIC:
single-cell regulatory network inference and clustering. Nat Methods. 2017;14(11):1083-6. Epub
20171009. doi: 10.1038/nmeth.4463. PubMed PMID: 28991892; PMCID: PMC5937676.
87.   Stuart T, Srivastava A, Madad S, Lareau CA, Satija R. Author Correction: Single-cell
chromatin state analysis with Signac. Nat Methods. 2022;19(2):257. doi: 10.1038/s41592-022-
01393-7. PubMed PMID: 34997233.
88.   Li W, Xu H, Xiao T, Cong L, Love MI, Zhang F, Irizarry RA, Liu JS, Brown M, Liu XS.
MAGeCK enables robust identification of essential genes from genome-scale CRISPR/Cas9
knockout screens. Genome Biol. 2014;15(12):554. doi: 10.1186/s13059-014-0554-4. PubMed
PMID: 25476604; PMCID: PMC4290824.
89.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21. Epub
20121025. doi: 10.1093/bioinformatics/bts635. PubMed PMID: 23104886; PMCID: PMC3530905.
90.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics. 2010;26(1):139-40. Epub
20091111. doi: 10.1093/bioinformatics/btp616. PubMed PMID: 19910308; PMCID: PMC2796818.
91.   Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. BMC Bioinformatics. 2011;12:323. Epub 20110804. doi:
10.1186/1471-2105-12-323. PubMed PMID: 21816040; PMCID: PMC3163565.
92.   Zhou W, Triche TJ, Jr., Laird PW, Shen H. SeSAMe: reducing artifactual detection of DNA
methylation by Infinium BeadChips in genomic deletions. Nucleic Acids Res. 2018;46(20):e123.
doi: 10.1093/nar/gky691. PubMed PMID: 30085201; PMCID: PMC6237738.
93.   Colaprico A, Silva TC, Olsen C, Garofano L, Cava C, Garolini D, Sabedot TS, Malta TM,
Pagnotta SM, Castiglioni I, Ceccarelli M, Bontempi G, Noushmehr H. TCGAbiolinks: an
R/Bioconductor package for integrative analysis of TCGA data. Nucleic Acids Res.
2016;44(8):e71. Epub 20151223. doi: 10.1093/nar/gkv1507. PubMed PMID: 26704973; PMCID:
PMC4856967.
94.   Lawrence M, Huber W, Pages H, Aboyoun P, Carlson M, Gentleman R, Morgan MT, Carey
VJ. Software for computing and annotating genomic ranges. PLoS Comput Biol.
2013;9(8):e1003118. Epub 20130808. doi: 10.1371/journal.pcbi.1003118. PubMed PMID:
23950696; PMCID: PMC3738458.
95.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ,
Heuer ML, Larsson E, Antipin Y, Reva B, Goldberg AP, Sander C, Schultz N. The cBio cancer
genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer
Discov. 2012;2(5):401-4. doi: 10.1158/2159-8290.CD-12-0095. PubMed PMID: 22588877;
PMCID: PMC3956037.
96.   Arafeh R, Shibue T, Dempster JM, Hahn WC, Vazquez F. The present and future of the
Cancer Dependency Map. Nat Rev Cancer. 2025;25(1):59-73. Epub 20241028. doi:
10.1038/s41568-024-00763-x. PubMed PMID: 39468210.
97.   Goldman MJ, Craft B, Hastie M, Repecka K, McDade F, Kamath A, Banerjee A, Luo Y,
Rogers D, Brooks AN, Zhu J, Haussler D. Visualizing and interpreting cancer genomics data via
the Xena platform. Nat Biotechnol. 2020;38(6):675-8. doi: 10.1038/s41587-020-0546-8. PubMed
PMID: 32444850; PMCID: PMC7386072.
98.   Yost KE, Zhao Y, Hung KL, Zhu K, Xu D, Corces MR, Shams S, Louie BH, Sarmashghi S,
Sundaram L, Luebeck J, Clarke S, Doane AS, Granja JM, Choudhry H, Imielinski M, Cherniack
AD, Khurana E, Bafna V, Felau I, Zenklusen JC, Laird PW, Curtis C, Cancer Genome Atlas
Analysis N, Greenleaf WJ, Chang HY. Three-dimensional genome landscape of primary human
cancers. Nat Genet. 2025;57(5):1189-200. Epub 20250512. doi: 10.1038/s41588-025-02188-0.
PubMed PMID: 40355593; PMCID: PMC12081301.

99. Almassalha LM, Bauer GM, Wu W, Cherkezyan L, Zhang D, Kendra A, Gladstein S,
Chandler JE, VanDerway D, Seagle BL, Ugolkov A, Billadeau DD, O'Halloran TV, Mazar AP, Roy
HK, Szleifer I, Shahabi S, Backman V. Macrogenomic engineering via modulation of the scaling
of chromatin packing density. Nat Biomed Eng. 2017;1(11):902-13. Epub 20171106. doi:
10.1038/s41551-017-0153-2. PubMed PMID: 29450107; PMCID: PMC5809134.
100. Stephens AD, Liu PZ, Banigan EJ, Almassalha LM, Backman V, Adam SA, Goldman RD,
Marko JF. Chromatin histone modifications and rigidity affect nuclear morphology independent of
lamins. Mol Biol Cell. 2018;29(2):220-33. Epub 20171115. doi: 10.1091/mbc.E17-06-0410.
PubMed PMID: 29142071; PMCID: PMC5909933.
101. Almassalha LM, Carignano M, Liwag EP, Li WS, Gong R, Acosta N, Dunton CL, Gonzalez
PC, Carter LM, Kakkaramadam R, Kroger M, MacQuarrie KL, Frederick J, Ye IC, Su P, Kuo T,
Medina KI, Pritchard JA, Skol A, Nap R, Kanemaki M, Dravid V, Szleifer I, Backman V. Chromatin
conformation, gene transcription, and nucleosome remodeling as an emergent system. Sci Adv.
2025;11(2):eadq6652. Epub 20250110. doi: 10.1126/sciadv.adq6652. PubMed PMID: 39792661;
PMCID: PMC11721585.
102. Lieberman-Aiden E, van Berkum NL, Williams L, Imakaev M, Ragoczy T, Telling A, Amit I,
Lajoie BR, Sabo PJ, Dorschner MO, Sandstrom R, Bernstein B, Bender MA, Groudine M, Gnirke
A, Stamatoyannopoulos J, Mirny LA, Lander ES, Dekker J. Comprehensive mapping of long-
range interactions reveals folding principles of the human genome. Science. 2009;326(5950):289-
93. doi: 10.1126/science.1181369. PubMed PMID: 19815776; PMCID: PMC2858594.
103. Simonis M, Klous P, Splinter E, Moshkin Y, Willemsen R, de Wit E, van Steensel B, de
Laat W. Nuclear organization of active and inactive chromatin domains uncovered by
chromosome conformation capture-on-chip (4C). Nat Genet. 2006;38(11):1348-54. Epub
20061008. doi: 10.1038/ng1896. PubMed PMID: 17033623.
104. Kleffman K, Levinson G, Rose IVL, Blumenberg LM, Shadaloey SAA, Dhabaria A, Wong
E, Galan-Echevarria F, Karz A, Argibay D, Von Itter R, Floristan A, Baptiste G, Eskow NM, Tranos
JA, Chen J, Vega YSdMEC, Call M, Rogers R, Jour G, Wadghiri YZ, Osman I, Li YM, Mathews P,
DeMattos RB, Ueberheide B, Ruggles KV, Liddelow SA, Schneider RJ, Hernando E. Melanoma-
Secreted Amyloid Beta Suppresses Neuroinflammation and Promotes Brain Metastasis. Cancer
Discov. 2022;12(5):1314-35. doi: 10.1158/2159-8290.CD-21-1006. PubMed PMID: 35262173;
PMCID: PMC9069488.
105. Ibrahim M, Illa-Bochaca I, Jour G, Vega-Saenz de Miera E, Fracasso J, Ruggles K, Osman
I, Schober M. NF1 Loss Promotes EGFR Activation and Confers Sensitivity to EGFR Inhibition in
NF1 Mutant Melanoma. Cancer Res. 2025. Epub 20250610. doi: 10.1158/0008-5472.CAN-24-
3904. PubMed PMID: 40494652; PMCID: PMC12221223.
106. Liu L, Luo Q, Sun J, Ju Y, Morita Y, Song G. Chromatin organization regulated by EZH2-
mediated H3K27me3 is required for OPN-induced migration of bone marrow-derived
mesenchymal stem cells. Int J Biochem Cell Biol. 2018;96:29-39. Epub 20180111. doi:
10.1016/j.biocel.2018.01.006. PubMed PMID: 29337251.
107. Stephens AD, Banigan EJ, Adam SA, Goldman RD, Marko JF. Chromatin and lamin A
determine two different mechanical response regimes of the cell nucleus. Mol Biol Cell.
2017;28(14):1984-96. Epub 20170105. doi: 10.1091/mbc.E16-09-0653. PubMed PMID:
28057760; PMCID: PMC5541848.
108. Stephens AD, Banigan EJ, Marko JF. Chromatin's physical properties shape the nucleus
and its functions. Curr Opin Cell Biol. 2019;58:76-84. Epub 20190316. doi:
10.1016/j.ceb.2019.02.006. PubMed PMID: 30889417; PMCID: PMC6692209.
109. Wang X, Luan Y, Yue F. EagleC: A deep-learning framework for detecting a full range of
structural variations from bulk and single-cell contact maps. Sci Adv. 2022;8(24):eabn9215. Epub
20220615. doi: 10.1126/sciadv.abn9215. PubMed PMID: 35704579; PMCID: PMC9200291.
110. Corces MR, Granja JM, Shams S, Louie BH, Seoane JA, Zhou W, Silva TC, Groeneveld
C, Wong CK, Cho SW, Satpathy AT, Mumbach MR, Hoadley KA, Robertson AG, Sheffield NC,

Felau I, Castro MAA, Berman BP, Staudt LM, Zenklusen JC, Laird PW, Curtis C, Cancer Genome Atlas Analysis N, Greenleaf WJ, Chang HY. The chromatin accessibility landscape of primary human cancers. Science. 2018;362(6413). doi: 10.1126/science.aav1898. PubMed PMID: 30361341; PMCID: PMC6408149.

111.   Puram SV, Tirosh I, Parikh AS, Patel AP, Yizhak K, Gillespie S, Rodman C, Luo CL, Mroz EA, Emerick KS, Deschler DG, Varvares MA, Mylvaganam R, Rozenblatt-Rosen O, Rocco JW, Faquin WC, Lin DT, Regev A, Bernstein BE. Single-Cell Transcriptomic Analysis of Primary and Metastatic Tumor Ecosystems in Head and Neck Cancer. Cell. 2017;171(7):1611-24 e24. Epub 20171130. doi: 10.1016/j.cell.2017.10.044. PubMed PMID: 29198524; PMCID: PMC5878932.

112.   Chung W, Eum HH, Lee HO, Lee KM, Lee HB, Kim KT, Ryu HS, Kim S, Lee JE, Park YH, Kan Z, Han W, Park WY. Single-cell RNA-seq enables comprehensive tumour and immune cell profiling in primary breast cancer. Nat Commun. 2017;8:15081. Epub 20170505. doi: 10.1038/ncomms15081. PubMed PMID: 28474673; PMCID: PMC5424158.

113.   Patel AP, Tirosh I, Trombetta JJ, Shalek AK, Gillespie SM, Wakimoto H, Cahill DP, Nahed BV, Curry WT, Martuza RL, Louis DN, Rozenblatt-Rosen O, Suva ML, Regev A, Bernstein BE. Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma. Science. 2014;344(6190):1396-401. Epub 20140612. doi: 10.1126/science.1254257. PubMed PMID: 24925914; PMCID: PMC4123637.

114.   Rambow F, Rogiers A, Marin-Bejar O, Aibar S, Femel J, Dewaele M, Karras P, Brown D, Chang YH, Debiec-Rychter M, Adriaens C, Radaelli E, Wolter P, Bechter O, Dummer R, Levesque M, Piris A, Frederick DT, Boland G, Flaherty KT, van den Oord J, Voet T, Aerts S, Lund AW, Marine JC. Toward Minimal Residual Disease-Directed Therapy in Melanoma. Cell. 2018;174(4):843-55 e19. Epub 20180712. doi: 10.1016/j.cell.2018.06.025. PubMed PMID: 30017245.

115.   Polyak K, Weinberg RA. Transitions between epithelial and mesenchymal states: acquisition of malignant and stem cell traits. Nat Rev Cancer. 2009;9(4):265-73. Epub 20090305. doi: 10.1038/nrc2620. PubMed PMID: 19262571.

116.   Barkley D, Moncada R, Pour M, Liberman DA, Dryg I, Werba G, Wang W, Baron M, Rao A, Xia B, Franca GS, Weil A, Delair DF, Hajdu C, Lund AW, Osman I, Yanai I. Cancer cell states recur across tumor types and form specific interactions with the tumor microenvironment. Nat Genet. 2022;54(8):1192-201. Epub 20220805. doi: 10.1038/s41588-022-01141-9. PubMed PMID: 35931863; PMCID: PMC9886402.

117.   Shaffer SM, Dunagin MC, Torborg SR, Torre EA, Emert B, Krepler C, Beqiri M, Sproesser K, Brafford PA, Xiao M, Eggan E, Anastopoulos IN, Vargas-Garcia CA, Singh A, Nathanson KL, Herlyn M, Raj A. Rare cell variability and drug-induced reprogramming as a mode of cancer drug resistance. Nature. 2017;546(7658):431-+. doi: 10.1038/nature22794 http://www.nature.com/nature/journal/vaop/ncurrent/abs/nature22794.html#supplementary-information.

118.   Tan YY, Li JJ, Zhao GY, Huang KC, Cardenas H, Wang YN, Matei D, Cheng JX. Metabolic reprogramming from glycolysis to fatty acid uptake and beta-oxidation in platinum-resistant cancer cells. Nature Communications. 2022;13(1). doi: 10.1038/s41467-022-32101-w. PubMed PMID: WOS:000836609800006.

119.   Sundaram L, Kumar A, Zatzman M, Salcedo A, Ravindra N, Shams S, Louie BH, Bagdatli ST, Myers MA, Sarmashghi S, Choi HY, Choi WY, Yost KE, Zhao Y, Granja JM, Hinoue T, Hayes DN, Cherniack A, Felau I, Choudhry H, Zenklusen JC, Farh KK, McPherson A, Curtis C, Laird PW, Cancer Genome Atlas Analysis Networkdouble d, Demchok JA, Yang L, Tarnuzzer R, Caesar-Johnson SJ, Wang Z, Doane AS, Khurana E, Castro MAA, Lazar AJ, Broom BM, Weinstein JN, Akbani R, Kumar SV, Raphael BJ, Wong CK, Stuart JM, Safavi R, Benz CC, Johnson BK, Kyi C, Shen H, Corces MR, Chang HY, Greenleaf WJ. Single-cell chromatin accessibility reveals malignant regulatory programs in primary human cancers. Science. 2024;385(6713):eadk9217. Epub 20240906. doi: 10.1126/science.adk9217. PubMed PMID: 39236169; PMCID: PMC12289346.

120.   Do C, Jiang G, Cova G, Zappile P, Heguy A, Skok JA. The coordination between CTCF,
cohesin and TFs impacts nucleosome repositioning and chromatin insulation to define state
specific 3D chromatin folding. bioRxiv. 2025. Epub 20250521. doi: 10.1101/2024.11.02.620823.
PubMed PMID: 39605491; PMCID: PMC11601386.
121.   Rao SSP, Huang SC, Glenn St Hilaire B, Engreitz JM, Perez EM, Kieffer-Kwon KR,
Sanborn AL, Johnstone SE, Bascom GD, Bochkov ID, Huang X, Shamim MS, Shin J, Turner D,
Ye Z, Omer AD, Robinson JT, Schlick T, Bernstein BE, Casellas R, Lander ES, Aiden EL. Cohesin
Loss Eliminates All Loop Domains. Cell. 2017;171(2):305-20 e24. doi: 10.1016/j.cell.2017.09.026.
PubMed PMID: 28985562; PMCID: PMC5846482.
122.   Phillips JE, Corces VG. CTCF: master weaver of the genome. Cell. 2009;137(7):1194-
211. doi: 10.1016/j.cell.2009.06.001. PubMed PMID: 19563753; PMCID: PMC3040116.
123.   Narita T, Kilic S, Higashijima Y, Scherer NM, Pappas G, Maskey E, Choudhary C.
Disentangling the architectural and non-architectural functions of CTCF and cohesin in gene
regulation. Nat Genet. 2025. Epub 20251118. doi: 10.1038/s41588-025-02404-x. PubMed PMID:
41254162.
124.   Zhan P, Xi GM, Zhang B, Wu Y, Liu HB, Liu YF, Xu WJ, Zhu Q, Cai F, Zhou ZJ, Miu YY,
Wang XX, Jin JJ, Li Q, Lv TF, Song Y. NCAPG2 promotes tumour proliferation by regulating G2/M
phase and associates with poor prognosis in lung adenocarcinoma. J Cell Mol Med.
2017;21(4):665-76. Epub 20161115. doi: 10.1111/jcmm.13010. PubMed PMID: 27862966;
PMCID: PMC5345611.
125.   Hu WF, Krieger KL, Lagundzin D, Li X, Cheung RS, Taniguchi T, Johnson KR, Bessho T,
Monteiro ANA, Woods NT. CTDP1 regulates breast cancer survival and DNA repair through
BRCT-specific interactions with FANCI. Cell Death Discov. 2019;5:105. Epub 20190619. doi:
10.1038/s41420-019-0185-3. PubMed PMID: 31240132; PMCID: PMC6584691.
126.   Marques E, Englund JI, Tervonen TA, Virkunen E, Laakso M, Myllynen M, Makela A,
Ahvenainen M, Lepikhova T, Monni O, Hautaniemi S, Klefstrom J. Par6G suppresses cell
proliferation and is targeted by loss-of-function mutations in multiple cancers. Oncogene.
2016;35(11):1386-98. Epub 20150615. doi: 10.1038/onc.2015.196. PubMed PMID: 26073086;
PMCID: PMC4800288.
127.   Shoshan E, Braeuer RR, Kamiya T, Mobley AK, Huang L, Vasquez ME, Velazquez-Torres
G, Chakravarti N, Ivan C, Prieto V, Villares GJ, Bar-Eli M. NFAT1 Directly Regulates IL8 and
MMP3 to Promote Melanoma Tumor Growth and Metastasis. Cancer Res. 2016;76(11):3145-55.
Epub 20160324. doi: 10.1158/0008-5472.CAN-15-2511. PubMed PMID: 27013197; PMCID:
PMC4891299.
128.   Lee YC, Tsai KW, Liao JB, Kuo WT, Chang YC, Yang YF. High expression of tight junction
protein 1 as a predictive biomarker for bladder cancer grade and staging. Sci Rep.
2022;12(1):1496. Epub 20220127. doi: 10.1038/s41598-022-05631-y. PubMed PMID: 35087173;
PMCID: PMC8795112.
129.   Martinez-Noel G, Luck K, Kuhnle S, Desbuleux A, Szajner P, Galligan JT, Rodriguez D,
Zheng L, Boyland K, Leclere F, Zhong Q, Hill DE, Vidal M, Howley PM. Network Analysis of
UBE3A/E6AP-Associated Proteins Provides Connections to Several Distinct Cellular Processes.
J Mol Biol. 2018;430(7):1024-50. Epub 20180206. doi: 10.1016/j.jmb.2018.01.021. PubMed
PMID: 29426014; PMCID: PMC5866790.
130.   Janssen SM, Moscona R, Elchebly M, Papadakis AI, Redpath M, Wang H, Rubin E, van
Kempen LC, Spatz A. BORIS/CTCFL promotes a switch from a proliferative towards an invasive
phenotype in melanoma cells. Cell Death Discov. 2020;6:1. Epub 20200102. doi:
10.1038/s41420-019-0235-x. PubMed PMID: 32123577; PMCID: PMC7026120.
131.   Moscona R, Janssen SM, Elchebly M, Papadakis AI, Rubin E, Spatz A. BORIS/CTCFL-
mediated chromatin accessibility alterations promote a pro-invasive transcriptional signature in
melanoma cells. Pigment Cell Melanoma Res. 2023;36(3-4):299-313. Epub 20230421. doi:
10.1111/pcmr.13089. PubMed PMID: 37082838.

132. Xu X, Jarajapu V, Chadourne M, Soto-Vargas Z, Ochoa-Rios SL, Jasani N, Wang K,
Mecozzi N, Yao J, Yu X, Karreth FA. FRA1 drives melanoma metastasis through an actionable
transcriptional network. Oncogene. 2025. Epub 20251124. doi: 10.1038/s41388-025-03632-5.
PubMed PMID: 41286309.
133. Acosta N, Gong R, Su Y, Frederick J, Medina KI, Li WS, Mohammadian K, Almassalha L,
Wang G, Backman V. Three-color single-molecule localization microscopy in chromatin. Light:
Science & Applications. 2025;14(1):123. doi: 10.1038/s41377-025-01786-1.
134. Nicetto D, Donahue G, Jain T, Peng T, Sidoli S, Sheng L, Montavon T, Becker JS,
Grindheim JM, Blahnik K, Garcia BA, Tan K, Bonasio R, Jenuwein T, Zaret KS. H3K9me3-
heterochromatin loss at protein-coding genes enables developmental lineage specification.
Science. 2019;363(6424):294-7. Epub 20190103. doi: 10.1126/science.aau0583. PubMed PMID:
30606806; PMCID: PMC6664818.
135. Kim YJ, Sheu KM, Tsoi J, Abril-Rodriguez G, Medina E, Grasso CS, Torrejon DY,
Champhekar AS, Litchfield K, Swanton C, Speiser DE, Scumpia PO, Hoffmann A, Graeber TG,
Puig-Saus C, Ribas A. Melanoma dedifferentiation induced by IFN-gamma epigenetic remodeling
in response to anti-PD-1 therapy. J Clin Invest. 2021;131(12). doi: 10.1172/JCI145859. PubMed
PMID: 33914706; PMCID: PMC8203459.
136. Fontanals-Cirera B, Hasson D, Vardabasso C, Di Micco R, Agrawal P, Chowdhury A,
Gantz M, de Pablos-Aragoneses A, Morgenstern A, Wu P, Filipescu D, Valle-Garcia D, Darvishian
F, Roe JS, Davies MA, Vakoc CR, Hernando E, Bernstein E. Harnessing BET Inhibitor Sensitivity
Reveals AMIGO2 as a Melanoma Survival Gene. Mol Cell. 2017;68(4):731-44 e9. doi:
10.1016/j.molcel.2017.11.004. PubMed PMID: 29149598; PMCID: PMC5993436.
1965

Figure 1

Figure 2

Figure 3

Figure 4

a 12-273BM (CDX)

scRNAseq   loHiC

IFN−active
Melanocytic
Mesenchymal
Mitotic
Neural Crest−like

b *cis*

E1 values (20% quantile)

AA
AB:BA
BB

MEL MES   MEL MES   MEL MES

c 12-273BM TADs

#TADs

TADs size (bp)

MEL   MES

d 12-273BM TADs

Insulation score

MEL
MES

-0.6   boundary   0.6Mb

e FFPE TADs

0.1182

#TADs

MEL   MES

0.0775

TADs size (bp)

MEL   MES

Figure 5

Figure 6

Figure 7