De novo origin and evolution of an antimicrobial protein function

Titas Sil[1,2], Caitlin H. Kowalski[1,3], Sierra Scamfer[2,4], Matthew F. Barber[1,2]

[1]Institute of Ecology and Evolution, University of Oregon; Eugene, OR 97403, USA.

[2]Department of Biology, University of Oregon; Eugene, OR 97403, USA.

[3]Department of Dermatology, Dartmouth Hitchcock Medical Center; Lebanon, NH 03756, USA.

[4]Institute of Molecular Biology, University of Oregon; Eugene, OR 97403, USA.

<u>Corresponding author</u>

Matthew F. Barber

Email: mfbarber@uoregon.edu

Keywords: protein evolution, antimicrobial peptides, innate immunity, pathogens, lactoferrin

Abstract

Antimicrobial peptides (AMPs) constitute key components of innate immunity across the tree of
life. Canonical AMPs are typically translated as small proteins and secreted from host cells to act
against microbes. However, cryptic AMP-like domains are also embedded within diverse proteins
not classically associated with antimicrobial function. How such embedded AMPs first emerge
and diversify remains unclear. Here we retrace the origin and evolution of the abundant
mammalian protein lactoferrin and its embedded AMP, lactoferricin. By resurrecting extinct
lactoferrin ancestors dating back to the earliest mammals, we identify a gradual enrichment of
cationic and hydrophobic amino acids in the lactoferricin domain over time. These changes
enabled ancient lactoferricin to first rupture bacterial membranes, an activity that was later
enhanced in extant mammals conferring potent bactericidal activity against diverse bacteria. In
addition, we find that recent natural selection within the lactoferricin domain has continued to
modulate antimicrobial activity on recent evolutionary timescales. In particular, we pinpoint a
single rapidly evolving site in lactoferricin among great apes that significantly enhances
antimicrobial potency against major pathogenic bacteria. Together our study illustrates how novel
immune protein functions can arise, evolve, and diversify to strengthen host defense against
diverse microbial pathogens.

Significance Statement
Cryptic antimicrobial peptide (AMP) domains have been increasingly identified within larger
proteins, yet how these embedded AMPs first emerge and diversify remains unclear. Here we
retrace the origin and evolution of the abundant mammalian immunity protein lactoferrin and its
embedded AMP, lactoferricin. By resurrecting extinct mammalian lactoferrin ancestors, we
identify molecular alterations that first enabled lactoferricin to rupture bacterial membranes and
later acquire bacterial-killing activity. Our findings illustrate how new protein functions can quickly
arise and evolve to confer immune defense against deadly pathogens. More broadly, this work

highlights the proteome as a reservoir of latent antimicrobial defenses, suggesting new strategies
for discovering AMP-based therapeutics to combat the escalating threat of antibiotic-resistance.

Main Text

Introduction

The evolution of immune function is vital for species survival in response to pathogen antagonism
(*1–3*). Antimicrobial peptides (AMPs), also known as host defense peptides, are major
contributors to innate immunity across all domains of life, displaying a range of antimicrobial and
immunomodulatory activities (*4–7*). Well-characterized AMPs are typically small (less than 50
amino acids) and positively charged, promoting interactions with negatively charged microbial cell
envelopes (*4*, *5*, *8*). These interactions can lead to membrane disruption, cell lysis, and
impairment of essential cellular processes, ultimately inhibiting microbial growth or triggering cell
death (*8*). Beyond canonical AMPs, an increasing number of studies have identified the presence
of AMP-like domains embedded within larger proteins (*9–13*). These domains can in some cases
be liberated through proteolytic cleavage, suggesting a mechanism by which diverse proteins
may acquire immune functionality. At present, little is known regarding the origins of these
embedded AMP domains or the molecular changes that enabled the acquisition of antimicrobial
activity.

The mammalian protein lactoferrin provides an exemplar for the evolution of antimicrobial
functions. Lactoferrin is an abundant secreted iron-binding protein that arose by duplication of the
transferrin gene in the ancestor of placental mammals approximately 160 million years ago (Fig.
1A) (*14*, *15*). Transferrin proteins are typically composed of two homologous domains, termed the
N- and C-lobes, each of which binds a single ferric iron ion with high affinity (*15–18*). Serum
transferrin mediates iron transport in the bloodstream, delivering this essential metal nutrient to
cells via receptor-mediated endocytosis (*15*). Transferrin family proteins can also provide a
defensive benefit by sequestering iron from invasive pathogens, a process termed "nutritional
immunity"(*19–21*). Lactoferrin, in contrast to transferrin, is abundantly expressed in mammalian
milk, colostrum, tears, secondary granules of neutrophils, and other body fluids (*22*, *23*). Since its
birth and divergence, lactoferrin has also acquired novel antimicrobial functions which are absent
in extant transferrin (*23*). The N-terminal region of the lactoferrin N-lobe can be proteolytically
cleaved by abundant host proteases such as pepsin, trypsin, and chymotrypsin to generate
cationic AMPs, including lactoferricin and lactoferrampin (*24–29*). These peptides are capable of
killing diverse bacterial and fungal pathogens as well as inhibiting microbial biofilm formation (*26*,
28, *30–32*). Lactoferrin-derived AMPs share key features with conventional AMPs, including a
high proportion of positively charged and hydrophobic amino acids that confer amphipathic
properties (Fig. 1B, S1). Despite its functional importance, the evolutionary origin of antimicrobial
activity within the lactoferricin domain remains poorly understood. Previously we found that
lactoferricin exhibits evidence of repeated positive selection in simian primates, with codons
possessing an elevated rate of non-synonymous to synonymous substitutions (dN/dS) (*33*). This
suggests that lactoferricin has continued to diversify within mammals since the acquisition of
antimicrobial activity. Determining how this novel function arose and subsequently evolved in an
otherwise well-conserved iron-transport protein could offer broader insights into the origin of
antimicrobial proteins and the mechanisms by which new biological functions arise. Here we
retrace the stepwise emergence of lactoferricin antimicrobial activity in ancestral mammals, as
well as demonstrate how recent natural selection has enhanced the function of this abundant host
defense protein.

Results

Ancestral lactoferrin accumulated positive charges after gene birth
To investigate the origin and evolutionary history of the lactoferricin domain, we first compared
lactoferricin sequences across diverse mammals (Fig. S1). Given that lactoferricin peptides have

been reported of different lengths in different species, we focused our studies on the core 25 amino
acids conserved across lactoferricin orthologs (amino acids 17-42 in mature human lactoferrin) (*24*,
25). Most homologs possessed elevated isoelectric points indicating enrichment of positively
charged amino acids such as arginine (R), lysine (K), and histidine (H). They also contained
interspersed hydrophobic residues characteristic of AMPs. Despite these conserved
physiochemical properties, lactoferricin amino acid sequences were highly diverse across
orthologs, making it difficult to pinpoint amino acids that may have been responsible for
antimicrobial function (Fig. S1B).
105
To retrace the mutations that led to the emergence of a novel antimicrobial function, we
reconstructed ancestral lactoferrin and transferrin sequences across diverse mammals (Fig. 1C).
Ancestral sequence reconstruction provides a powerful approach to characterize the function of
ancient proteins (*34–36*). Using the Topiary software pipeline (*37*) and publicly available transferrin
and mammalian lactoferrin sequences from NCBI, we predicted ancestral amino acid sequences
of transferrin and lactoferrin proteins (Fig. S2A). We focused on ancestral proteins (prefixed "Anc")
at key nodes of the phylogenetic tree where the lactoferricin domain underwent significant
divergence (Fig. 1C, Fig. S2B). Each predicted sequence was associated with posterior probability
scores, which identified the most likely residue at each site as well as the most probable alternative
(Fig. S3). These data were used to generate corresponding alternative (Altall_Anc) peptide
sequences, in which alternative amino acid states with a posterior probability greater than 0.2
replaced the most probable ancestral amino acid. To assess the robustness of our findings, we
included both the primary ancestral reconstructions and their alternate versions in subsequent
functional analyses (Fig. S4).
120
To compare the antimicrobial function of lactoferricin between ancestral and extant variants, we
chose human and bovine lactoferrin (hLF and bLF, respectively) as extant references, given the
extensive previous work on these orthologs in the field (*22*, *24*, *25*, *27*). As a negative control we
selected human serum transferrin (hTF) as it did not exhibit antimicrobial function in the N-terminal
region (Fig. S1A). We referred to the last common ancestor of mammalian transferrin prior to the
gene duplication event as AncTF. Gene duplication produced the first ancestral lactoferrin, which
we designated AncLF1 (Fig. 1C). We also considered two intermediate ancestors, denoted AncLF2
and AncLF3. AncLF2 represents the last common ancestor of hLF and bLF in the gene phylogeny,
and AncLF3 represents a bovine-specific ancestor that accumulated multiple substitutions in the
lactoferricin region relative to AncLF2 (Fig. S2B). We then predicted the structures of these full-
length ancestral proteins using AlphaFold2 (Fig. 1C). When the surface electrostatic potentials
were compared, we observed a progressive increase in positive charge density (blue surface) in
the N-terminal region of lactoferrin following the duplication event (Fig. 1C). This suggests that the
emergence of positively charged surfaces in the N-terminus coincided with the origin of lactoferrin
itself in early placental mammals.
136
We next examined the lactoferricin domains of these reconstructed lactoferrin ancestors, which we
refer to as AncLFcin1, AncLFcin2, and AncLFcin3 (Fig. S2B). As the corresponding region in
transferrin (residues 17-42) does not form a known antimicrobial peptide, we referred to these
homologous sequences as AncTF17-42 in ancestral transferrin and hTF17-42 in human serum
transferrin. We observed that cationic amino acids became more abundant in lactoferricin post-
duplication, with the emergence of a K5 substitution in AncLFcin1, R8 in AncLFcin2, and R9 in
AncLFcin3 (Fig. S2B). To quantify this trend, we calculated peptide isoelectric points, which
increased from AncTF17-42 to bLFcin (Fig. S2B). Furthermore, we noted the emergence of
hydrophobic residues adjacent to positively charged amino acids in the lactoferricin domain after
duplication (e.g., alanine 1 in AncLFcin1 and AncLFcin2). Based on these observations, we
hypothesized that the gradual increase in positive charge and hydrophobicity enabled lactoferricin
to acquire antimicrobial function by mediating association with bacterial cell envelopes.
149
Early origin of antimicrobial activity in lactoferricin
To characterize the antimicrobial potency of ancestral and extant lactoferricin domains, we
synthesized these peptides and tested their activity against several bacterial pathogens (Fig. 2).

Among Gram-negative bacteria we included *Pseudomonas aeruginosa*, a pathogen frequently
associated with nosocomial infections, chronic wounds, and cystic fibrosis, as well as a reference
strain of the enteric bacterium *Escherichia coli* (*38–40*). We also tested representative Gram-
positive bacteria including *Staphylococcus aureus*, a major cause of invasive hospital and
community-acquired infections, as well as *Streptococcus agalactiae*, a frequent agent of severe
infections in neonates (*41–43*).
159
To compare the antimicrobial potency of ancestral and extant lactoferricin homologs, we measured
bacterial growth in presence of the peptides at various concentrations over 24 hours. Human
lactoferrin concentrations range widely *in vivo* from 7-8 mg/mL in colostrum, 1-3 mg/mL in breast
milk, 0.1-2 mg/mL in tears and saliva, and even lower levels (0.001-0.1 mg/mL) in other secretory
body fluids (*44*). We therefore used physiologically relevant concentrations of each peptide, ranging
from 0.1-1mg/mL, to assess their dose-dependent antimicrobial activities. Bacterial growth at each
concentration was measured by normalized area under the curve (AUC) calculations, where higher
AUC corresponds to increased bacterial growth. Relative to the untreated control, peptides
exhibiting <50% bacterial growth were classified as having high antimicrobial potency, those with
50–75% growth as moderate, and those with >75% growth as weak. We observed that all bacteria
grew to high levels in the presence of hTF17-42, indicating it has little to no antimicrobial activity
as expected (Fig. 2). Similarly, the ancestral-derived peptide AncTF17-42 exhibited negligible
antimicrobial activity across all doses and pathogens (Fig. 2). AncLFcin1, in contrast, reduced
growth by >50% at higher doses against a subset of pathogens, for example at 400 µg/mL against
P. aeruginosa PAO1 and at 800 µg/mL against *E. coli* (Fig. 2A, B). These findings suggest that
antimicrobial activity in this domain arose after, rather than before, the birth of the lactoferrin gene
itself.
177
AncLFcin2, representing the last common ancestor of hLF and bLF, exhibited higher potency at
lower concentrations across all pathogens tested relative to AncLFcin1 (Fig. 2). For example, it was
highly potent at 100 µg/mL against *P. aeruginosa* PAO1, and at 50 µg/mL against *E. coli* and *S.*
aureus MN8 (Fig. 2A, 2C, 2E). Against other pathogens, AncLFcin2 also reduced bacterial growth
by >50% in the 100-400 µg/mL range. These results suggest that antimicrobial activity of the
lactoferricin domain was substantially enhanced between AncLFcin1 and AncLFcin2. AncLFcin3
exhibited broad antimicrobial activity in the 100-800 µg/mL range but overall was less effective than
bLFcin.
186
To validate the robustness of our ancestral sequence reconstruction, we also tested alternate
versions of each ancestor against *P. aeruginosa* and *S. aureus* (Fig. S4). Overall trends in
antimicrobial activity were largely consistent with those observed for the primary reconstructed
ancestors. The only exception was Altall_AncTF17-42, which exhibited weak activity against *S.*
aureus at the highest concentrations (Fig. S4). bLFcin was consistently more potent than hLFcin
against all pathogens tested, significantly reducing bacterial growth at concentrations as low as 50
µg/mL (Fig. 2). Notably, hLFcin was more effective against Gram-negative bacteria where it
restricted growth at 100-200 µg/mL. Against Gram-positive bacteria, the antimicrobial potency of
hLFcin only became evident at concentrations above 200 µg/mL. Interestingly, AncLFcin2 was
more effective than hLFcin against Gram-positive bacteria. Overall these data indicate that modest
antimicrobial activity first emerged in AncLFcin1, immediately after the birth of lactoferrin in ancient
mammals, which then increased in subsequent ancestors. However, given that hLFcin exhibited
weaker antimicrobial activity than some earlier ancestors, our findings also suggest that the
evolution of lactoferricin antimicrobial function did not increase uniformly during mammalian
evolution.
202
Stepwise enhancement of bactericidal activity during lactoferricin evolution
To further investigate the evolution of antimicrobial function in lactoferricin, we assessed growth
kinetics and viability of representative Gram-negative (*P. aeruginosa*) and Gram-positive (*S.*
aureus) bacteria in the presence of ancestral and extant lactoferricin peptides (Fig. 3). In addition
to analysing growth curves and calculating AUCs, we quantified bacterial survival by measuring
colony-forming units (CFUs) over time. Based on the differences in dose-dependent activity

observed previously, we used 100 µg/mL of the peptides against *P. aeruginosa* and 800 µg/mL
against *S. aureus* in subsequent experiments.
211
Consistent with our previous results, extant and ancestral transferrin-derived peptides, hTF17-42
and AncTF17-42 respectively, exhibited no detectable effects on *P. aeruginosa* growth and survival
(Fig. 3A–C). AncLFcin1 displayed an intermediate phenotype, with a modest but significant
reduction in overall *P. aeruginosa* growth (Fig. 3A, B). We also observed a rapid decrease in *P.*
aeruginosa viability within 2 hours of treatment, followed by a partial recovery (Fig. 3C). In contrast,
AncLFcin2 consistently suppressed *P. aeruginosa* growth throughout the experiment, resulting in
a significant 1000-fold reduction in bacterial viability after 24 hours of treatment (Fig. 3A-C).
AncLFcin3, the ancestor of bLFcin, inhibited *P. aeruginosa* growth and caused 100-1000 fold
reduction in viability during the first 15 hours of treatment, after which bacterial growth resumed.
hLFcin also consistently suppressed *P. aeruginosa* growth, causing a 10,000 fold reduction in
viability at the end of the treatment. Finally, bLFcin exhibited the most potent bactericidal effect,
reflected by minimal AUC values and no detectable *P. aeruginosa* colonies within 2 hours of
treatment (Fig. 3A-C).
225
Lactoferricin activity profiles against *S. aureus* differed from those of *P. aeruginosa* in several ways
(Fig. 3D–F). hTF17-42 and AncTF17-42 exhibited modest inhibitory effects on *S. aureus* later in
the treatment period (Fig. 3D, E), but these differences did not translate into reductions in bacterial
viability as measured by CFU counts (Fig. 3F). AncLFcin1 exerted partial inhibition against *S.*
aureus, with little reduction in bacterial viability. AncLFcin2 strongly suppressed bacterial growth,
resulting in an approximately 100,000 fold reduction in *S. aureus* viability by the end of treatment.
AncLFcin3 also significantly inhibited *S. aureus* growth and reduced its viability over the course of
the experiment. In contrast, hLFcin initially inhibited bacterial growth and caused a rapid 100-fold
reduction in viability. Subsequently, *S. aureus* resumed growth, although this was not reflected in
the optical density, possibly due to a reduction in cell size. Again, bLFcin remained the most potent
peptide, eliminating detectable colonies within two hours (Fig. 3F). Collectively these findings
indicate that weak antimicrobial activity emerged in the earliest lactoferricin ancestor, with
bactericidal activity substantially enhanced in subsequent orthologs.
239
Lactoferricin membrane permeabilizing activity arose early after gene birth
While both ancestral and extant lactoferricin peptides exhibited some degree of bactericidal activity,
they varied dramatically in their potency (Fig. 3). This led us to further examine the molecular
mechanisms underlying differences in their activities. A common feature of many cationic AMPs is
their ability to disrupt and permeabilize bacterial membranes (*4*, *6*). To investigate this function, we
measured the membrane permeabilizing ability of ancestral and extant lactoferricin peptides
against *P. aeruginosa*. Membrane permeabilization was measured by staining with propidium
iodide (PI) which enters the bacterial cell and binds DNA only when cell membranes are
compromised (*45*). We found that all ancestral and extant lactoferricins, but not transferrin-derived
peptides, were capable of permeabilizing bacterial membranes within 30 minutes of incubation as
detected by PI staining (Fig. 4A). This finding indicates that bacterial membrane permeabilization
was an early trait that emerged in the lactoferricin domain shortly after gene birth.
252
Membrane permeabilization by AMPs can induce bacterial stress responses as well as alter
membrane polarization and ion flux, which in turn impacts cell physiology and morphology (*46*, *47*).
To determine the extent to which ancestral and extant lactoferricins affect target cell physiology,
we measured bacterial membrane polarization. *P. aeruginosa* cells were treated with the cationic
dye 3,3'-dipropylthiadicarbocyanine iodide (DiSC$_3$(5)) that accumulates in polarized membranes
quenching its fluorescence (*48*). Upon membrane depolarization, DiSC$_3$(5) is released, and
fluorescence increases. We found that all ancestral and extant lactoferricin-treated cells were
hyperpolarized, as demonstrated by a reduction in fluorescence, compared to untreated, hTF17-
42, or AncTF17-42-treated cells (Fig. 4B).
262
Given that all lactoferricin peptides exhibited both membrane-permeabilizing and polarization-
altering activities, we sought to understand how these peptides nonetheless differed in their overall

antimicrobial potency. To address this, we shifted from population-level assays to single-cell
morphological characterization. We hypothesized that lactoferricin variants with increased
antimicrobial activity would exert greater impact on bacterial cell envelope integrity. To visualize
this, we treated *P. aeruginosa* with the lipophilic dye Nile red, which homogenously binds to
membranes and produces brightly stained foci in regions of increased fluidity, indicative of
membrane perturbation(*45*). Using spinning-disc confocal microscopy, hTF17-42 and AncTF17-42
exhibited no visible effects on bacterial morphology (Fig. 4C). In contrast, cells treated with
AncLFcin1, AncLFcin2, or AncLFcin3 displayed puncta-like structures on their membranes,
suggesting localized membrane disturbance. These cells also appeared significantly smaller than
untreated controls. This effect was most pronounced with hLFcin and bLFcin, where cells became
markedly reduced in size, rounded, and membranes were barely distinguishable from the cytosol
(Fig. 4C). Quantitative measurements of cell size confirmed that ancestral lactoferricins induced
moderate but significant cell shrinkage, whereas extant variants amplified this effect (Fig. 4D).
278
Membrane disruption is a common mechanism of AMP function, which can cause leaking of
cytoplasmic content and shrinkage of cells (*45*, *49*, *50*). Such perturbations can be transient and
resealable or irreversible, depending on the extent of the damage (*50*, *51*). Our results suggest that
bacteria are able to recover from ancestral lactoferricin-induced membrane damage, whereas
extant variants, particularly bLFcin, caused irreversible cell envelope collapse. Together, these
findings indicate that the lactoferricin domain initially possessed membrane-permeabilizing activity,
which intensified during evolution to produce potent bactericidal effects.
286
Rapid evolution of lactoferricin antimicrobial activity in primates
In addition to investigating the molecular origins of the lactoferricin domain, we also sought to
determine how lactoferricin antimicrobial activity has changed over recent evolutionary timescales.
Previous work by our group found evidence that lactoferrin has been subject to repeated positive
selection within human populations as well as between simian primate species, suggesting
antimicrobial activity may have been modulated via recurrent adaptation (*18*, *33*). We identified
several sites in the N-lobe of lactoferrin with elevated rates of nonsynonymous to synonymous
substitutions (dN/dS), including two within the lactoferricin region (Fig. 5A, S5) (*33*). For example,
position 5 varies between primates, with humans and many monkeys encoding a glutamine (Q),
whereas all other great apes encode an arginine (R). In addition, position 12 is variable among
primates as well as polymorphic within human populations, generally toggling between cationic
arginine and lysine (K12 or R12) residues (*33*).
299
To assess the functional consequences of natural selection at sites 5 and 12, we generated two
point mutants in human lactoferricin: hLFcinK12R and hLFcinQ5R. hLFcinK12R exhibited
antimicrobial activity comparable to wild type hLFcin against both *P. aeruginosa* and *S. aureus,*
indicating that the K12R substitution does not substantially alter activity against these pathogens
(Fig. 5B, C). In contrast, the hLFcinQ5R mutant displayed significant enhancement in antimicrobial
potency against *S. aureus* relative to hLFcin (Fig. 5B, C). Beyond the addition of a single positive
charge, this effect may stem from the positioning of R5 adjacent to the aromatic hydrophobic
residue tryptophan (W6). We noted a similar pattern in AncLFcin2, which exhibited greater potency
against *S. aureus* than hLFcin and contains a lysine (K5) adjacent to phenylalanine, another
aromatic hydrophobic residue (Fig. S2B, 2C-D, 3D-F). Notably, the most potent variant, bLFcin,
contains both combinations, which may contribute to its heightened antimicrobial activity. Taken
together, these findings indicate that antimicrobial function of the lactoferricin domain has been
repeatedly modulated by the forces of natural selection, leading to enhanced activity across
mammalian lineages.
314
315 **Discussion**
316
AMPs have long been appreciated as deeply conserved effectors of innate immunity. Recent
large-scale analyses further suggest that AMP-like domains are far more widespread than
previously appreciated (*10–13*). Computational studies have detected AMP-like domains
embedded within diverse proteins not previously associated with direct antimicrobial function,

including interferon gamma, IL-13, histone H1, gasdermin E, and dynactin, among others (*10–*
13). The existence of such cryptic antimicrobial functions suggests a previously unappreciated
layer of host defense, though the evolutionary origins of such embedded AMPs remain largely
uncharacterized. Leveraging mammalian lactoferrin and its embedded AMP lactoferricin as a
model, we observed that even the earliest lactoferricin ancestor possessed the ability to
permeabilize bacterial membranes and alter membrane potential, a property that was further
enhanced in later ancestors (Fig. 4). This increased activity was accompanied by a marked
perturbation of bacterial cells, likely due to leakage of cytoplasmic content (*49–51*). Membrane
permeabilization by AMPs is often reversible depending on the severity of damage (*50*, *51*).
Hence, it is reasonable to speculate that early lactoferricin domains induced perturbations that
bacteria could repair, allowing eventual recovery. In contrast, more potent lactoferricin variants
such as bLFcin induced irreversible damage at much lower concentrations, leading to complete
lysis or severe morphological damage that prevented recovery (Fig. 4). These findings
demonstrate that membrane permeabilization and hyperpolarization were early antimicrobial traits
of lactoferricin, and that bactericidal activity intensified over time.
336
We noted that some ancestral lactoferricins exerted higher antimicrobial activity than their extant
counterparts. For example, AncLFcin2 and AncLFcin3 were consistently more bactericidal
against *S. aureus* than human lactoferricin (Fig. 2, 3). Thus, the evolution of antimicrobial function
in lactoferrin does not reflect a simple linear increase in potency but has rather fluctuated along
different lineages depending on the target microbe. We considered whether a reduction in
lactoferricin antimicrobial activity could indicate a trade-off to minimize toxicity against host cells.
However, treatment of bovine erythrocytes revealed no significant hemolysis in the presence of
any lactoferricin peptides, inconsistent with this hypothesis (Fig. S6). An alternative explanation is
that some bacteria may have evolved host-specific resistance to these AMPs. *S. aureus*, for
example, is human associated with many of its virulence factors and toxins exhibiting narrow
tropism for humans (*52–54*). Together these results underscore how ancestral sequence
reconstruction can uncover unique variants of extant AMPs that could serve as safe and effective
pathogen treatment options, particularly in the era of growing antibiotic resistance (*5*, *55*).
350
AMPs have frequently been identified among the most rapidly evolving genes in host genomes,
likely reflecting repeated adaptation to counteract diverse microbes (*56*, *57*). In contrast, far less
is known about how AMP domains embedded within larger proteins emerge and evolve in
response to selection. We previously identified two sites exhibiting signatures of positive selection
in the lactoferricin domain among primates (*33*). Notably, one of these substitutions (Q5R) was
sufficient to significantly enhance the antimicrobial activity of human lactoferricin, particularly
against *S. aureus* (Fig. 5). This observation suggests the importance of this region for
lactoferricin's activity. Notably, the adjacent sixth residue is invariably an aromatic hydrophobic
amino acid, either phenylalanine (F) or tryptophan (W), across all lactoferricin sequences (Fig.
S2B). AncLFcin2, which is effective against all pathogens tested, also possesses cationic and
aromatic hydrophobic residues at these sites (K5F6). The most potent variant tested, bovine
lactoferricin, contains three such pairs of cationic and aromatic residues (F1K2, R5W6, and
W7R8). Although hTF and AncTF both contain a F6R7 amino acid pair, the effect of the positively
charged arginine is likely diminished by the neighboring aspartate (D8). Given that cationic nature
and hydrophobicity are key features of many AMPs, the juxtaposition of a cationic residue with an
aromatic hydrophobic residue within lactoferrin may reflect positive epistasis between these sites.
Indeed, epistasis between nearby R and W residues has been reported previously in antiviral
proteins (*58*). It is well established that cationic amino acids mediate electrostatic interactions with
the negatively charged bacterial envelopes, whereas hydrophobic residues insert into
membranes and contribute to disruption of the lipid bilayer (*59*). Future studies will determine how
selection and epistasis together contribute to antimicrobial potency in these and other AMPs.
372
In addition to providing protection against invasive pathogens, the expression of AMPs at barrier
tissues positions them to broadly shape host-associated microbiotas. Work in *Drosophila* has
demonstrated how natural selection and polymorphism in AMP genes can impact diverse resident
microbes (*56*, *60*, *61*), as well as confer defense against microbes within specific ecological

niches (*62*). Given that lactoferrin is among the most abundant secreted proteins in many
mammalian barrier tissues, genetic variation may similarly shape the diversity and composition of
host-associated microbial communities. By retracing the origin and evolution of antimicrobial
activity in mammalian lactoferrin, our study illustrates how new protein functions can emerge to
promote host defense against major microbial pathogens.
382
383
384 **Materials and Methods**
385
Ancestral sequence reconstruction
Vertebrate transferrin and mammalian lactoferrin homologs were retrieved from the NCBI and
UniProt databases for ancestral sequence reconstruction. Multiple sequence alignments and
maximum likelihood phylogenetic trees were generated using the Topiary pipeline (*1*).
Alignments were manually curated to remove insertions and deletions. Using Topiary, ancestral
sequences were inferred at each node, with posterior probabilities calculated for each site and
node to assess reconstruction confidence.
393
394 **Peptide synthesis**
395 All peptides were commercially synthesized by GenScript. Lyophilized peptides were
396 resuspended in phosphate-buffered saline (PBS, pH 7.0) to a concentration of 10 mg/mL and
397 stored at –80°C.
398
Structural prediction of ancestral sequences
Protein structure predictions for the reconstructed ancestral sequences were performed using
ColabFold, a streamlined implementation of AlphaFold2 (*2*). Prior to structure prediction, the N-
terminal region corresponding to the human lactoferrin signal peptide (hLF1–19) was removed
from all ancestral sequences. Predicted protein structures were visualized and analyzed using
UCSF ChimeraX (version 1.8, released 2024-06-10) (*3*). The theoretical isoelectric point (pI) of
each peptide was calculated using the ExPASy Compute pI/Mw tool (*4*).
406
Bacterial growth assays
Overnight cultures of all bacterial strains were prepared by inoculating a single colony into low-
nutrient 5% tryptic soy broth (TSB, v/v) to model the low-nutrient conditions of the host
environment (*5*) and incubated at 250 rpm overnight, unless otherwise specified. The following
day, cultures were diluted to an optical density at 600 nm (OD_{600}) of 0.02 in fresh 5% TSB. For
growth assays, 150 μL of the diluted culture was added to each well of a 96-well microtiter plate,
followed by the addition of peptides at the desired concentrations. Plates were incubated for 24
hours with continuous shaking in a Synergy H1 microplate reader (BioTek), and bacterial growth
was monitored by measuring OD_{600} at 10-minute intervals. Area under the curve (AUC) was
calculated using GraphPad Prism software.
417
Bacterial survival assay
Overnight cultures were diluted to an OD_{600} of 0.02 in fresh medium, and antimicrobial peptides
were added at the indicated concentrations in 1.5 mL microcentrifuge tubes. To assess bacterial
viability over time, samples were taken at defined time points during the treatment period. At each
time point, cultures were serially diluted 10-fold (up to 10^{-7}) and plated on tryptic soy agar (TSA)
to enumerate colony-forming units (CFUs). Plates were incubated at 37°C overnight, and CFUs
were counted the following day to evaluate the time-course of bacterial survival.
425
Membrane permeabilization measurements
Overnight cultures of *Pseudomonas aeruginosa* PAO1 were subcultured by diluting to an OD_{600}
of 0.3 in fresh 5% TSB. Aliquots of 100 μL were transferred into 1.5 mL microcentrifuge tubes and
treated with the peptides for 30 minutes. Cells were pelleted by centrifugation at 5,000 × g for 1
minute. Pellets were resuspended in 300 μL of 20 μM propidium iodide (Sigma-Aldrich P4170-
10MG) and incubated in the dark for 15 minutes to prevent photobleaching. Excess dye was
removed by centrifugation at 5,000 × g for 1 minute, the supernatant was discarded, and pellets

were resuspended in 300 μL of phosphate-buffered saline (PBS, pH 7.0). For fluorescence
measurement, 100 μL aliquots from each sample were transferred in triplicate into black-walled,
clear-bottom 96-well plates (without lids). Fluorescence intensity was measured from the bottom
of the plate using a microplate reader with excitation at 535 nm and emission at 617 nm.
437
Membrane potential measurements
The membrane depolarizing activity of peptides was assessed using the voltage-sensitive dye
DiSC$_3$(5) (MedChem Express #HY-D0085-25mg) as previously described(6). *Pseudomonas*
aeruginosa PAO1 overnight cultures were diluted in fresh 5% TSB to OD$_{600}$ of 0.3. DiSC$_3$(5) (6),
prepared as a stock solution in DMSO, was added to the bacterial suspension at a final
concentration of 5 μM. Following a brief equilibration period to allow dye accumulation across
polarized membranes, peptides were added at a final concentration of 100 μg/mL in a black,
clear-bottom 96-well plate. Fluorescence was monitored kinetically over a 1-hour period at 37 °C
using a Synergy H1 microplate reader (BioTek), with excitation and emission wavelengths set to
610 nm and 660 nm, respectively. A decrease in fluorescence intensity was interpreted as
membrane hyperpolarization, reflecting increased accumulation of the cationic, lipophilic dye
within the bacterial cytoplasmic membrane, where self-quenching reduces the overall
fluorescence signal.
451
Nile red staining, microscopy and cell size measurements
The Nile red staining protocol was adapted and modified from Dombach *et al.*, 2021 (7).
Overnight bacterial cultures were diluted in fresh 5% tryptic soy broth (TSB, v/v) to an optical
density at 600 nm (OD$_{600}$) of 0.3. From this, 100 μL aliquots were transferred to 1.5 mL
microcentrifuge tubes for peptide treatment. Samples were incubated at 37 °C for 40 minutes with
shaking at 250 rpm. Following peptide treatment, Nile red (Sigma-Aldrich 72485-100MG) was
added to each tube at a final concentration of 10 μg/mL, followed by a 5-minute incubation at
37 °C. Cells were then fixed by adding paraformaldehyde to a final concentration of 4%, and
incubated at room temperature for 10 minutes. After fixation, cells were pelleted by centrifugation
at 10,000 × g for 1 minute and resuspended in 10 μL of PBS, pH 7.0. For microscopy, 7 μL of the
resuspended cells was applied to a coverslip, overlaid with 0.5% agarose, and gently
compressed with a glass slide to immobilize the sample. Imaging was performed using a SoRA
spinning disk confocal microscope (Nikon CSU-W1) equipped with a 60X water-immersion
objective. Bacterial cell size was measured using the Fiji package of ImageJ software (8).
466
Hemolysis assay
The hemolysis assay was adapted from Greco *et al.*, 2020 (9). Bovine red blood cells (RBCs)
were isolated by centrifuging 1 mL of whole blood at 3000 × g for 2 min. The RBC pellet was
washed five times with phosphate-buffered saline (PBS) by resuspension and centrifugation until
the supernatant was clear, then diluted 1:10 in PBS. In a 96-well plate, 100 μL of the diluted RBC
suspension was mixed with each peptide solution (final concentration: 1 mg/mL) and incubated at
37 °C for 1 h. Absorbance was measured at 405 nm using a microplate reader, and the
percentage of hemolysis was calculated relative to a positive control (1% Triton X-100).
475
476 **Author Contributions:** T.S. and M.F.B. designed research; T.S. performed research; T.S.,
477 C.H.K., and S.S. contributed to data collection; T.S. and M.F.B. wrote the paper; T.S., C.H.K.,
478 S.S., and M.F.B edited the paper.
479

Competing Interest Statement: Authors declare that they have no competing interests.

Acknowledgements: We thank Mike Harms and Jose Sanchez-Borbon for assistance with
ancestral sequence reconstruction. We are thankful to the members of Barber lab for their
feedback and support, as well as Karen Guillemin, Jarrod Smith, and Melanie Spero for helpful
insights and discussion. We thank the Genomics and Cell Characterization Core Facility (GC3F)
at UO and Adam Fries for assistance with microscopy.

485
486 **References**

1. M. D. Daugherty, H. S. Malik, Rules of engagement: molecular insights from host-virus arms races. *Annu. Rev. Genet.* **46**, 677–700 (2012).

2. M. F. Barber, J. R. Fitzgerald, Mechanisms of host adaptation by bacterial pathogens. *FEMS Microbiol. Rev.* **48** (2024).

3. Z. Pancer, M. D. Cooper, The evolution of adaptive immunity. *Annu. Rev. Immunol.* **24**, 497–518 (2006).

4. N. G. Oliveira Júnior, C. M. Souza, D. F. Buccini, M. H. Cardoso, O. L. Franco, Antimicrobial peptides: structure, functions and translational applications. *Nat. Rev. Microbiol.*, doi: 10.1038/s41579-025-01200-y (2025).

5. M. Magana, M. Pushpanathan, A. L. Santos, L. Leanse, M. Fernandez, A. Ioannidis, M. A. Giulianotti, Y. Apidianakis, S. Bradfute, A. L. Ferguson, A. Cherkasov, M. N. Seleem, C. Pinilla, C. de la Fuente-Nunez, T. Lazaridis, T. Dai, R. A. Houghten, R. E. W. Hancock, G. P. Tegos, The value of antimicrobial peptides in the age of resistance. *Lancet Infect. Dis.* **20**, e216–e230 (2020).

6. B. P. Lazzaro, M. Zasloff, J. Rolff, Antimicrobial peptides: Application informed by evolution. *Science* **368**, eaau5480 (2020).

7. R. E. Hancock, M. G. Scott, The role of antimicrobial peptides in animal defenses. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 8856–8861 (2000).

8. M. Zasloff, Antimicrobial peptides in health and disease, *The New England journal of medicine*. **347** (2002)pp. 1199–1200.

9. H. S. Kim, H. Yoon, I. Minn, C. B. Park, W. T. Lee, M. Zasloff, S. C. Kim, Pepsin-mediated processing of the cytoplasmic histone H2A to strong antimicrobial peptide buforin I. *J. Immunol.* **165**, 3268–3274 (2000).

10. N. Y. Yount, D. C. Weaver, E. Y. Lee, M. W. Lee, H. Wang, L. C. Chan, G. C. L. Wong, M. R. Yeaman, Unifying structural signature of eukaryotic α-helical host defense peptides. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 6944–6953 (2019).

11. K. Goldberg, A. Lobov, P. Antonello, M. D. Shmueli, I. Yakir, T. Weizman, A. Ulman, D. Sheban, E. Laser, M. P. Kramer, R. Shteinvil, G. Chen, A. Ibraheem, V. Sysoeva, V. Fishbain-Yoskovitz, G. Mohapatra, A. Abramov, S. Shimshi, K. Ogneva, M. Nandy, S. Amidror, H. Bootz-Maoz, S. H. Kuo, N. Dezorella, A. Kacen, A. Javitt, G. W. Lau, N. Yissachar, Z. Hayouka, Y. Merbl, Cell-autonomous innate immunity by proteasome-derived defence peptides. *Nature*, doi: 10.1038/s41586-025-08615-w (2025).

12. G. Marsman, X. Zheng, D. Čerina, K. A. Lacey, M. Liu, D. Humme, C. Goosmann, V. Brinkmann, C. J. Harbort, V. J. Torres, A. Zychlinsky, Histone H1 kills MRSA. *Cell Rep.* **43**, 114969 (2024).

13. M. D. T. Torres, A. Cesaro, C. de la Fuente-Nunez, Peptides from non-immune proteins target infections through antimicrobial and immunomodulatory properties. *Trends Biotechnol.* **43**, 184–205 (2025).

14. A. L. Hughes, R. Friedman, Evolutionary diversification of the vertebrate transferrin multi-gene family. *Immunogenetics* **66**, 651–661 (2014).

15. L. A. Lambert, Molecular evolution of the transferrin family and associated receptors. *Biochim. Biophys. Acta* **1820**, 244–255 (2012).

16. M. Haridas, B. F. Anderson, E. N. Baker, Structure of human diferric lactoferrin refined at 2.2 Å resolution. *Acta Crystallogr. D Biol. Crystallogr.* **51**, 629–646 (1995).

17. N. Yang, H. Zhang, M. Wang, Q. Hao, H. Sun, Iron and bismuth bound human serum transferrin reveals a partially-opened conformation in the N-lobe. *Sci. Rep.* **2**, 999 (2012).

18. M. F. Barber, N. C. Elde, Escape from bacterial iron piracy through rapid evolution of transferrin. *Science* **346**, 1362–1366 (2014).

19. C. C. Murdoch, E. P. Skaar, Nutritional immunity: the battle for nutrient metals at the host-pathogen interface. *Nat. Rev. Microbiol.*, doi: 10.1038/s41579-022-00745-6 (2022).

20. I. Iatsenko, A. Marra, J.-P. Boquete, J. Peña, B. Lemaitre, Iron sequestration by transferrin 1 mediates nutritional immunity in Drosophila melanogaster. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 7317–7325 (2020).

21. M. F. Barber, N. C. Elde, Buried Treasure: Evolutionary Perspectives on Microbial Iron Piracy. *Trends Genet.* **31**, 627–636 (2015).

22. I. A. García-Montoya, T. S. Cendón, S. Arévalo-Gallegos, Q. Rascón-Cruz, Lactoferrin a
multiple bioactive protein: an overview. *Biochim. Biophys. Acta* **1820**, 226–236 (2012).
23. P. Kowalczyk, K. Kaczyńska, P. Kleczkowska, I. Bukowska-Ośko, K. Kramkowski, D.
Sulejczak, The lactoferrin phenomenon-A miracle molecule. *Molecules* **27**, 2941 (2022).
24. W. Bellamy, M. Takase, K. Yamauchi, H. Wakabayashi, K. Kawase, M. Tomita,
Identification of the bactericidal domain of lactoferrin. *Biochim. Biophys. Acta* **1121**, 130–
136 (1992).
25. M. Sinha, S. Kaushik, P. Kaur, S. Sharma, T. P. Singh, Antimicrobial lactoferrin peptides:
the hidden players in the protective function of a multifunctional protein. *Int. J. Pept.*
2013, 390230 (2013).
26. G. Ramamourthy, H. J. Vogel, Antibiofilm activity of lactoferrin-derived synthetic peptides
against Pseudomonas aeruginosa PAO1. *Biochem. Cell Biol.* **99**, 138–148 (2021).
27. A. Ohradanova-Repic, R. Praženicová, L. Gebetsberger, T. Moskalets, R. Skrabana, O.
Cehlar, G. Tajti, H. Stockinger, V. Leksa, Time to kill and time to heal: The multifaceted
role of lactoferrin and lactoferricin in host defense. *Pharmaceutics* **15** (2023).
28. M. A. León-Calvijo, A. L. Leal-Castro, G. A. Almanzar-Reina, J. E. Rosas-Pérez, J. E.
García-Castañeda, Z. J. Rivera-Monroy, Antibacterial activity of synthetic peptides
derived from lactoferricin against Escherichia coli ATCC 25922 and Enterococcus
faecalis ATCC 29212. *Biomed Res. Int.* **2015**, 453826 (2015).
29. M. Nakamura, N. Tsuda, T. Miyata, M. Ikenaga, Antimicrobial effect and mechanism of
bovine lactoferrin against the potato common scab pathogen Streptomyces scabiei. *PLoS*
One **17**, e0264094 (2022).
30. K. E. Fernandes, D. A. Carter, The antifungal activity of lactoferrin and its derived
peptides: Mechanisms of action and synergy with drugs against fungal pathogens. *Front.*
Microbiol. **8**, 2 (2017).
31. H. Liao, S. Liu, H. Wang, H. Su, Z. Liu, Enhanced antifungal activity of bovine lactoferrin-
producing probiotic Lactobacillus casei in the murine model of vulvovaginal candidiasis.
BMC Microbiol. **19**, 7 (2019).
32. P. K. Singh, M. R. Parsek, E. P. Greenberg, M. J. Welsh, A component of innate
immunity prevents bacterial biofilm development. *Nature* **417**, 552–555 (2002).
33. M. F. Barber, Z. Kronenberg, M. Yandell, N. C. Elde, Antimicrobial Functions of
Lactoferrin Promote Genetic Conflicts in Ancient Primates and Modern Humans. *PLoS*
Genet. **12**, e1006063 (2016).
34. J. T. Bridgham, S. M. Carroll, J. W. Thornton, Evolution of hormone-receptor complexity
by molecular exploitation. *Science* **312**, 97–101 (2006).
35. M. J. Harms, J. W. Thornton, Analyzing protein structure and function using ancestral
gene reconstruction. *Curr. Opin. Struct. Biol.* **20**, 360–366 (2010).
36. T. N. Starr, L. K. Picton, J. W. Thornton, Alternative evolutionary histories in the
sequence space of an ancient protein. *Nature* **549**, 409–413 (2017).
37. K. N. Orlandi, S. R. Phillips, Z. R. Sailer, J. L. Harman, M. J. Harms, Topiary: Pruning the
manual labor from ancestral sequence reconstruction. *Protein Sci.* **32**, e4551 (2023).
38. M. Letizia, S. P. Diggle, M. Whiteley, Pseudomonas aeruginosa: ecology, evolution,
pathogenesis and antimicrobial susceptibility. *Nat. Rev. Microbiol.* **23**, 701–717 (2025).
39. A. Grace, R. Sahu, D. R. Owen, V. A. Dennis, Pseudomonas aeruginosa reference
strains PAO1 and PA14: A genomic, phenotypic, and therapeutic review. *Front. Microbiol.*
13, 1023523 (2022).
40. T. D. Minogue, H. A. Daligault, K. W. Davenport, K. A. Bishop-Lilly, S. M. Broomall, D. C.
Bruce, P. S. Chain, O. Chertkov, S. R. Coyne, T. Freitas, K. G. Frey, H. S. Gibbons, J.
Jaissle, C. L. Redden, C. N. Rosenzweig, Y. Xu, S. L. Johnson, Complete genome
assembly of Escherichia coli ATCC 25922, a serotype O6 reference strain. *Genome*
Announc. **2** (2014).
41. G. Y. C. Cheung, J. S. Bae, M. Otto, Pathogenicity and virulence of Staphylococcus
aureus. *Virulence* **12**, 547–569 (2021).
42. J. B. Parsons, A. Mourad, B. P. Conlon, T. Kielian, V. G. Fowler Jr, Methicillin-resistant
and susceptible Staphylococcus aureus: tolerance, immune evasion and treatment. *Nat.*
Rev. Microbiol., doi: 10.1038/s41579-025-01226-2 (2025).

43. H. C. Maisey, K. S. Doran, V. Nizet, Recent advances in understanding the molecular
basis of group B Streptococcus virulence. *Expert Rev. Mol. Med.* **10**, e27 (2008).
44. X. Cao, Y. Ren, Q. Lu, K. Wang, Y. Wu, Y. Wang, Y. Zhang, X.-S. Cui, Z. Yang, Z. Chen,
Lactoferrin: A glycoprotein that plays an active role in human health. *Front. Nutr.* **9**,
1018336 (2022).
45. N. Malanovic, J. A. Buttress, D. Vejzovic, A. Ön, P. Piller, D. Kolb, K. Lohner, H. Strahl,
Disruption of the cytoplasmic membrane structure and barrier function underlies the
potent antiseptic activity of octenidine in Gram-positive bacteria. *Appl. Environ. Microbiol.*
88, e0018022 (2022).
46. K. N. Bhaumik, R. Spohn, A. Dunai, L. Daruka, G. Olajos, F. Zákány, A. Hetényi, C. Pál,
T. A. Martinek, Chemically diverse antimicrobial peptides induce hyperpolarization of the
E. coli membrane. *Commun. Biol.* **7**, 1264 (2024).
47. D.-Y. D. Lee, L. Galera-Laporta, M. Bialecka-Fornal, E. C. Moon, Z. Shen, S. P. Briggs, J.
Garcia-Ojalvo, G. M. Süel, Magnesium flux modulates ribosomes to increase bacterial
survival. *Cell* **177**, 352-360.e13 (2019).
48. J. D. Te Winkel, D. A. Gray, K. H. Seistrup, L. W. Hamoen, H. Strahl, Analysis of
antimicrobial-triggered membrane depolarization using voltage sensitive dyes. *Front. Cell
Dev. Biol.* **4**, 29 (2016).
49. E. H.-L. Chen, C.-H. Wang, Y.-T. Liao, F.-Y. Chan, Y. Kanaoka, T. Uchihashi, K. Kato, L.
Lai, Y.-W. Chang, M.-C. Ho, R. P.-Y. Chen, Visualizing the membrane disruption action
of antimicrobial peptides by cryo-electron tomography. *Nat. Commun.* **14**, 5464 (2023).
50. Z. Yang, H. Choi, J. C. Weisshaar, Melittin-induced permeabilization, re-sealing, and re-
permeabilization of E. coli membranes. *Biophys. J.* **114**, 368–379 (2018).
51. A. Rice, A. C. Zourou, M. L. Cotten, R. W. Pastor, A unified model of transient poration
induced by antimicrobial peptides. *Proc. Natl. Acad. Sci. U. S. A.* **122**, e2510294122
(2025).
52. K. Tam, V. J. Torres, Staphylococcus aureus Secreted Toxins and Extracellular
Enzymes. *Microbiol Spectr* **7** (2019).
53. K. J. Koymans, M. Vrieling, R. D. Gorham Jr, J. A. G. van Strijp, Staphylococcal immune
evasion proteins: Structure, function, and host adaptation. *Curr. Top. Microbiol. Immunol.*
409, 441–489 (2017).
54. A. F. Haag, J. R. Fitzgerald, J. R. Penadés, Staphylococcus aureus in animals. *Microbiol.
Spectr.* **7** (2019).
55. B. Antunes, C. Zanchi, P. R. Johnston, B. Maron, C. Witzany, R. R. Regoes, Z. Hayouka,
J. Rolff, The evolution of antimicrobial peptide resistance in Pseudomonas aeruginosa is
severely constrained by random peptide mixtures. *PLoS Biol.* **22**, e3002692 (2024).
56. R. L. Unckless, V. M. Howick, B. P. Lazzaro, Convergent Balancing Selection on an
Antimicrobial Peptide in Drosophila. *CURBIO* **26**, 257–262 (2016).
57. A. van Dijk, R. Guabiraba, G. Bailleul, C. Schouler, H. P. Haagsman, A.-C. Lalmanach,
Evolutionary diversification of defensins and cathelicidins in birds and primates. *Mol.
Immunol.* **157**, 53–69 (2023).
58. R. A. Geiger, D. Khera, J. L. Tenthorey, G. Kochs, L. Graf, M. Emerman, H. S. Malik,
Heterozygous and generalist MxA super-restrictors overcome breadth-specificity trade-
offs in antiviral restriction. *Sci. Adv.* **11**, eadu0062 (2025).
59. K. J. Cheng, S. Shastry, J. D. Campolargo, M. J. Hallock, T. V. Pogorelov, Charge,
hydrophobicity, and lipid type drive antimicrobial peptides' unique perturbation
ensembles. *Biochemistry* **64**, 1484–1500 (2025).
60. M. A. Hanson, P. T. Hamilton, S. J. Perlman, Immune genes and divergent antimicrobial
peptides in flies of the subgenus Drosophila. *BMC Evol. Biol.* **16**, 228 (2016).
61. J. R. Chapman, T. Hill, R. L. Unckless, Balancing selection drives the maintenance of
genetic variation in Drosophila antimicrobial peptides. *Genome Biol. Evol.* **11**, 2691–2701
(2019).
62. M. A. Hanson, L. Grollmus, B. Lemaitre, Ecology-relevant bacteria drive the evolution of
host antimicrobial peptides in Drosophila. *Science* **381**, eadg5725 (2023).
63. M. Mirdita, K. Schütze, Y. Moriwaki, L. Heo, S. Ovchinnikov, M. Steinegger, ColabFold:
making protein folding accessible to all. *Nat. Methods* **19**, 679–682 (2022).

64. E. F. Pettersen, T. D. Goddard, C. C. Huang, G. S. Couch, D. M. Greenblatt, E. C. Meng,
T. E. Ferrin, UCSF Chimera--a visualization system for exploratory research and
analysis. *J. Comput. Chem.* **25**, 1605–1612 (2004).
65. E. Gasteiger, C. Hoogland, A. Gattiker, S. Duvaud, M. R. Wilkins, R. D. Appel, A.
Bairoch, "Protein identification and analysis tools on the ExPASy server" in *The*
Proteomics Protocols Handbook (Humana Press, Totowa, NJ, 2005), pp. 571–607.
66. J. L. Dombach, J. L. J. Quintana, C. S. Detweiler, Staphylococcal bacterial persister cells,
biofilms, and intracellular infection are disrupted by JD1, a membrane-damaging small
molecule. *MBio* **12**, e0180121 (2021).
67. J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S.
Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D. J. White, V. Hartenstein,
K. Eliceiri, P. Tomancak, A. Cardona, Fiji: an open-source platform for biological-image
analysis. *Nat. Methods* **9**, 676–682 (2012).
68. I. Greco, N. Molchanova, E. Holmedal, H. Jenssen, B. D. Hummel, J. L. Watts, J.
Håkansson, P. R. Hansen, J. Svenson, Correlation between hemolytic activity,
cytotoxicity and systemic in vivo toxicity of synthetic antimicrobial peptides. *Sci. Rep.* **10**,
13206 (2020).
671

673
Figure 1. Retracing the emergence of antimicrobial function in mammalian lactoferrin. (**A**)
The extant lactoferrin (PDB: 1lfg) and serum transferrin (PDB: 3qyt) genes arose via duplication

of transferrin in the ancestor of placental mammals. Surface electrostatic potential is shown.
While transferrin maintains key roles in iron binding and transport, lactoferrin has acquired new
functions including direct bactericidal activity. (**B**) Structure of the N-terminal lactoferricin region of
lactoferrin, enriched for positively charged amino acids. (**C**) Phylogenetic tree of reconstructed
ancestral lactoferrin homologs (AncTF and AncLF) with extant human transferrin (hTF), human
lactoferrin (hLF), and bovine lactoferrin (bLF) illustrated with surface electrostatic potentials.
Structures of ancestral proteins were predicted using AlphaFold2. AncTF represents the last
common ancestor of mammalian transferrin and lactoferrin, after which the first lactoferrin
ancestor (AncLF1) arose by gene duplication. AncLF2 is the common ancestor of hLF and bLF,
while AncLF3 represents a bLF ancestor.

Figure 2. origin of antimicrobial activity in lactoferricin. Heatmaps depict area under the
curve (AUC) values normalized against the untreated control from bacterial growth curves
performed in the presence of indicated peptides at specified concentrations. Gram-negative
strains are shown in pink, Gram-positive strains in purple.
691

Figure 3. Stepwise evolution of bactericidal activity in ancestral lactoferricin. (**A-C**)
Pseudomonas aeruginosa strain PAO1 treated with 100 μg/mL of indicated peptides, (**D-F**)
Staphylococcus aureus strain JE2 treated with 800 μg/mL of indicated peptides. (**A, D**) Growth
curves measured by optical density at 600 nm over time. (**B, E**) Corresponding area under the
curve (AUC) measurements normalized against untreated control based on bacterial growth
curves. (**C, F**) Bacterial survival curves measured as colony forming units (CFUs). All data
represent mean and standard error based on three biological replicates. Statistical significance
was determined using one-way ANOVA.

702
Figure 4. Lactoferricin membrane permeabilizing activity emerged early after gene
duplication. *Pseudomonas aeruginosa* strain PAO1 was treated with 100 µg/mL of indicated
peptides for 30-60 minutes to assess membrane disruption and cell morphology. (**A**) Membrane
permeabilization measured by propidium iodide staining after 30 minutes of treatment. (**B**)
Membrane polarization measured by DiSC3(5) dye after 1 hour of treatment. Peptides were
added 5 minutes after addition of the dye (shown with the dotted line). Lower fluorescence
indicates hyperpolarization. Data represent mean and standard error based on three biological
replicates. (**C**) Confocal microscopy images of peptide-treated cells at 30 minutes (scale bar,
1µM). (**D**) Quantification of bacterial cell size from microscopy images (n=100).
712

713
Figure 5. Natural selection enhanced lactoferricin antimicrobial potency in great apes. (A)
Alignment of lactoferricin orthologs from humans, chimpanzees, and rhesus macaques. Two sites
(positions 5 and 12) exhibit elevated dN/dS across primates. (B) Antimicrobial activity of hLFcin
and mutant peptides against *Pseudomonas aeruginosa* PAO1 at 100 μg/mL. (C) Antimicrobial
activity of hLFcin and mutant peptides against *Staphylococcus aureus* strain JE2 at 800 μg/mL.
Data represent mean and standard error based on three biological replicates.
720

 Supplementary figures

Fig. S1. Diversity and evolution of lactoferricin sequences. (A) Antimicrobial activity of extant
human and bovine lactoferricin (hLFcin and bLFcin, respectively) against *Pseudomonas*
aeruginosa PAO1 compared with the equivalent region in human transferrin (hTF17-42). (B)
Amino acid alignment of the 25-amino acid stretch of lactoferricin region across placental
mammals. Blue, red and green letters denote positively charged, negatively charged, and
hydrophobic aromatic residues, respectively.
729

Fig. S2. Ancestral sequence reconstruction (ASR) of vertebrate transferrin family proteins.
(A) Representative taxa included in the ASR. A total of 376 sequences of transferrin, lactoferrin, and melanotransferrin from diverse vertebrate species were used to reconstruct the ancestral sequences. (B) Alignment of the lactoferricin region from hTF, AncTF, selected AncLFs, hLF, and bLF, with their corresponding isoelectric points. Higher isoelectric points indicate higher incidence of positive charges.

Fig. S3. Reconstructed ancestral sequences and alternate variants. Posterior probabilities of
each site are shown for (A) AncTF, (B) AncLF1, (C) AncLF2 and (D) AncLF3. Blue indicates the
most probable amino acid at each position, while red represents the *alt_all* sequence containing
the second-most probable amino acid at each site. The lactoferricin regions are highlighted with
cyan rectangles in the full-length sequences. (E) Sequences of the lactoferricin regions in the
reconstructed ancestors. Yellow highlight denotes residues that differ between the most probable
(blue) and *alt_all* (red) sequences.

747
748 **Fig. S4. Antimicrobial activities of alternate ancestral peptides.** Heatmaps show area under
749 the curve (AUC) values normalized against the untreated control from bacterial growth curves in
750 the presence of the indicated peptides at specified concentrations. Gram-negative strains are
751 shown in pink, Gram-positive strains in purple.
752

753
754 **Fig. S5. Evolution of lactoferricin across simian primates.** Lactoferricin positions 5 and 12
755 exhibited elevated dN/dS consistent with repeated positive selection. In Great Apes, glutamine
756 (Q) at position 5 is replaced by arginine (R), adjacent to a tryptophan (W). At position 12, residues
757 vary between arginine (R) and lysine (K) across primate species.
758
759

760
761 **Fig. S6. Hemolysis of ancestral and extant peptides.** Bovine red blood cells (RBCs) were
762 treated with peptides at 1 mg/mL. Complete lysis (100% hemolysis) was achieved with 1% Triton
763 X-100 as a positive control. Hemolysis by the peptides is shown as a percentage relative to this
764 control. Data represent three biological replicates with the standard error.
765