Multimodal Profiling of Repair-associated Immune Dynamics in a Mouse Model of

Menstruation

Rebecca J. Ainslie[1,2], Phoebe M. Kirkwood[1], Hanning Li[1], Symeon Gerasimou[1], Philippa T.K.

Saunders[1], Ioannis Simitsidellis[1], and Douglas A. Gibson[1,2*]

[1]*Centre for Reproductive Health and [2]Centre for Inflammation Research, Institute for*

Regeneration and Repair, University of Edinburgh, Edinburgh BioQuarter, 4-5 Little France

Crescent, Edinburgh EH16 4UU.

[*]Corresponding Author

Douglas Gibson, Principal Investigator & Sir Henry Dale Fellow

Institute for Regeneration and Repair

The University of Edinburgh

Edinburgh BioQuarter

4-5 Little France Drive

Edinburgh

EH16 4UU

E-mail: D.A.Gibson@ed.ac.uk

Abstract

Despite the importance of inflammation to menstruation, we lack detailed understanding of

how immune dynamics contribute to endometrial repair. We performed detailed phenotypic

characterisation of uterine immune cells using single cell RNA sequencing, flow cytometry and

multiplex immunohistochemistry in a mouse model of simulated menstruation. Uterine tissues

were collected from age-matched controls or from key phases of menstruation, including

tissue breakdown, repair and remodelling. Our findings reveal distinct compositional changes

across different phases of menstruation and highlight predominant roles for monocytes,

macrophages, and neutrophils in endometrial repair. Immunohistochemistry revealed the

spatial association of these myeloid cell subsets with areas of tissue repair and remodelling.

Bioinformatic analysis highlighted key roles for monocyte, macrophage and neutrophil

signalling during endometrial repair with thrombospondin 1 and secreted phosphoprotein 1

emerging as key signalling pathways. These data significantly advance our understanding of

menstrual physiology and identifies potential therapeutic targets for menstrual disorders.

40-word summary:

This study investigates immune cell dynamics in a mouse model of menstruation, highlighting

roles for monocytes, macrophages, and neutrophils in non-fibrotic endometrial repair, and

identifies key signalling pathways as potential therapeutic targets for menstrual disorders.

<u>Introduction</u>

Menstruation is characterised by an inflammatory cascade resulting in the breakdown and shedding of the functional layer of the endometrium followed by phases of endometrial repair and remodelling [1-4].This extensive inflammatory process must be tightly regulated to maintain reproductive function, with deficits in endometrial repair associated with reproductive health disorders [5]. Despite the importance of inflammation to menstruation and reproductive health, our current understanding of how inflammatory cells contribute to endometrial repair is limited.

Studies investigating endometrial immune cell composition during human menstruation have relied on immunohistochemistry analysis of fixed tissues with limited direct profiling of immune cell subsets. These studies have reported that the abundance of monocytes and macrophages dramatically increase prior to menstruation [6] whilst neutrophils are increased during endometrial breakdown and repair [7, 8]. Profiling of lymphoid cells is more limited. A study in the 1990's suggested CD8+ T cells are increased post-menstruation during the early-mid proliferative phase of the cycle [9]. B cells are sparse in the endometrium throughout the cycle and therefore their role during menstruation has not been investigated [10]. In contrast, while uterine natural killer cells (uNKs) are abundant in the late secretory phase and in association with fertility regulation, their numbers are reportedly low during menstruation [11, 12] suggesting they may not play a significant role in endometrial repair.

The generation of human scRNAseq datasets have increased our understanding of the composition of endometrial immune cells in the proliferative and secretory phases of the menstrual cycle [13, 14]. However, studies profiling immune cells in tissue samples from the menstrual phase are lacking. To overcome these limitations we, and others, have developed and validated a refined mouse model of simulated menstruation [15] to gain insights into tissue dynamics, immune cell composition and mechanisms of scarless repair. Although mice don't naturally menstruate, the physiological changes that occur in the human endometrium can be modelled using hormonal priming and artificial stimulation of implantation. Following hormone

withdrawal, uterine tissues undergo reproducible breakdown and repair, recapitulating key

features of human menstruation including rapid re-epithelialisation, stromal cell remodelling

and endothelial cell proliferation [15].

We have previously used this mouse model of menstruation to investigate the role of uterine

macrophages during endometrial repair which identified distinct monocyte and macrophage

populations with spatially discrete distributions[16]. Specifically, immunohistochemistry

analysis revealed that monocyte-derived macrophages were located in areas of the

endometrium actively undergoing repair whilst tissue-resident macrophages were restricted to

areas that have previously undergone repair [16]. Macrophages have been shown to

contribute to endometrial shedding through the production of matrix metalloproteinases that

degrade the extracellular matrix and through the production of vascular endothelial growth

factor [17, 18]. Neutrophils are abundant during menstruation and are associated with

apoptosis and cytokine production but whether they contribute to endometrial repair has not

been explored [7, 8]. Cytokines and growth factors that support tissue breakdown and repair

include pro-inflammatory cytokines (IL-1β, TNF-α, IL-6)[19, 20], chemokines (CCL2, CCL5,

CXCL8) [8, 21] and regulators of matrix degradation and remodelling (MIF, MMPs,

prostaglandins, VEGF) [1, 22, 23]. However, expression of these mediators has mostly been

reported in whole tissues, limiting interpretation of cell-specific factors. Although immune cell

types and mediators in menstruation have been profiled to some extent, deeper phenotyping

of endometrial repair processes is needed to advance our understanding of menstrual

physiology.

Menstruation is a complex and dynamic inflammatory process, but the contribution of distinct

immune cell subsets and mediators remains poorly understood. To address this, we performed

detailed phenotypic characterisation of uterine immune cells in a mouse model of simulated

menstruation scRNAseq, flow cytometry and multiplex immunohistochemistry. We describe

temporal and spatial association of myeloid cells with regions of tissue repair and highlight

putative repair-associated signalling pathways associated with these cell types. Collectively,

these data provide new insights into inflammatory regulation of menstruation and identify key

pathways could be future therapeutic targets in menstrual disorders.

Materials and Methods

Mouse model of simulated menstruation

All animal work was carried out according to best practise in line with the Animal Welfare and

Ethical Review Body (AWERB) and under licensed approval from the UK Home Office. The

mouse model of simulated menstruation was conducted in accordance with standard protocols

that have been described in detail in previous studies[8, 15, 16, 24, 25]. Briefly, female mice

between eight to ten weeks of age underwent dorsal bilateral ovariectomy (d0) and were given

daily injections of 17β-oestradiol (E2) in sesame oil (100ng/100µl, d7-9). A progesterone (P4)-

secreting pellet was inserted subcutaneously on d13; mice also received daily injections of E2

(5ng/100µl; d13-15). On d15, decidualisation was induced by trans-cervical introduction of

sesame oil into the uterine lumen (40µl, d15). Removal of the P4 pellet (d19) results in a rapid

decline in progesterone, stimulating the breakdown and shedding of the decidual mass which

simulates human menstruation in the mouse. Uterine tissues were collected at various time

points following progesterone withdrawal (0-, 12-, 24-, and 48 hr) as well as from age-matched

mice (~3-month-old, oestrus cycling). A schematic of the mouse model of simulated

menstruation and gross morphological changes that occur in the endometrium during phases

of tissue breakdown, repair and remodelling can be seen in figure S1.

Bioinformatics analysis of single cell RNA sequencing data:

ScRNAseq analysis was performed on total leukocytes (CD45$^+$ live singlets) isolated from

uterine tissues 24- (n=4) and 48 hrs (n=4) following progesterone withdrawal, representing

time points of endometrial tissue repair and remodelling respectively, and compared to age-

matched cycling/control tissues (n=8). CD45$^+$ cells were recovered using fluorescence-

activated cell sorting (FACS), pooling 25,000 cells per sample to give a total of 100,000 cells

for input to the 24/48hr dataset and 200,000 cells for input to the cycling dataset. Isolated cell

suspensions were counted, and viability confirmed to be >85% using a TC20 Automated Cell

Counter (BioRad, Cat. No. 1450102). Cell suspensions were processed using the 10x

Chromium Controller (10x Genomics, USA) to partition single cells into Gel bead-in-Emulsions

(GEMs) containing unique 10x barcodes and cDNA libraries generated following standard

protocol

ls as previously described[24, 25]. Sequencing was carried out on the Illumina NovaSeq

platform using bespoke 10x parameters, according to standard protocols (Edinburgh

Genomics: http://genomics.ed.ac.uk/, Edinburgh).

Raw data files were processed using 10x Cell Ranger (version 2.0.1,

https://www.10xgenomics.com) and downstream QC, clustering and gene expression analysis

performed in *Seurat [26]* (Version 3) following analysis pipelines described previously[24, 25].

Data were filtered based on standard QC metrics: number of unique genes detected in each

cell (nFeature_RNA >200 & <5000); the percentage of mitochondrial genes (<10%); and the

percentage ribosomal genes (<10%). The Scrublet python package was used to further

identify and remove putative doublets from scRNAseq data[27]. Individual datasets were

combined using Seurat's 'anchor-based' integration workflow to facilitate accurate

comparative analysis between experimental conditions. Following integration, data were

normalised and scaled prior to unsupervised clustering based on PCA analysis of the most

variably expressed genes using a graph-based approach ('FindNeighbours', 'FindClusters). A

quantitative method was used alongside the 'ElbowPlot' function to determine the number of

principle components required to capture the majority of the variance in the data[28]. Resultant

clusters were visualised using the uniform manifold approximation and projection (UMAP)

method and differential gene expression analysis was performed ('FindAllMarkers') to identify

cluster specific gene signatures using the non-parametric Wilcoxon rank sum test and p-value

threshold of <0.05. Over-represented functional annotations associated with the differentially

expressed genes (DEGs) were detected using the clusterProfiler package[29] using core

functions to interpret data in the context of biological function, pathways, and networks.

The R package CellChat [30] (version 2) was utilised to investigate cellular interactions

between immune cell types. CellChat estimates the probability of cell-cell communication via

the integration of single cell gene expression data with a database (CellChatDB) of known

signalling ligands and receptors generated from the literature.

The data discussed in this study will be deposited in NCBI's Gene Expression Omnibus on

publication [31].

Flow cytometry:

Uterine tissue samples were dissected and transferred to PBS kept at 4°C prior to the tissue

digestion protocol. Individual uterine horns were analysed as distinct samples given the

potential for each to exhibit varying degrees of decidualisation in response to the

stimulus. Tissue processing for flow cytometry analysis was performed as described

previously [24, 25]. Briefly, uterine tissues were manually dissociated and incubated with

collagenase (1mg/ml) and DNase (0.1mg/ml) for 30 min at 37°C. Tissues were further

dispersed and washed in FACS buffer (PBS Ca2-Mg2-: 5% charcoal stripped foetal calf serum

(CSFCS – Cat. No. 12676029), 2mM EDTA) and subsequently strained through 70μm and

40μm cell strainers. Cell suspensions were centrifuged at 400rcf for 5 min and cell pellets re-

suspended in 1ml ACK lysing buffer (Gibco, Cat. No. A10492-01) for 1 min then washed.

Uterine cell suspensions were incubated with LIVE/DEAD Red (Life Technologies Ltd,

L34971) for 30 min on ice followed by incubation with Mouse TruStain FcXTM (Biolegend

101319; 1:1000) for 10 min on ice. Cell suspensions were subsequently incubated with

optimised dilutions of fluorescently conjugated extracellular antibodies for 30 min on ice, as

detailed in Table 1 –- Figure 2.

Following antibody incubation, single cell suspensions were washed and re-suspended in

FACS buffer at 4°C and analysed using a Cytek Aurora (Cytek Biosciences) and SpectroFlo

software. Data analysis was performed using FlowJo analysis software (FlowJo LLC, Oregon,

USA).

Tissue recovery, processing and Haematoxylin and Eosin staining

Mouse uteri were fixed in 10% neutral buffered formalin for 24 hr at 4°C before transfer into

70% ethanol, at which point they could be stored long term. Tissues were then processed

according to standard protocols for paraffin embedding by the University of Edinburgh

Histology Facility. Sections from formalin-fixed paraffin-embedded (FFPE) blocks were cut at

5um thickness using a microtome and the sections were placed on Xtra adhesive pre-cleaned

micro slides (Surgipath, Leica Biosystems) and placed in at 60°C for 15 min prior to staining.

For visualisation of tissue morphology, staining was performed with haematoxylin and eosin.

Immunohistochemistry and Immunofluorescence

For immunohistochemistry and immunofluorescence, FFPE uterine sections of 5um thickness

were deparaffinised and rehydrated using a series of xylene and alcohols followed by antigen

retrieval in Citrate retrieval buffer (pH6) for 5 min in an Instant Pot. Following washes with tap

water, the sections were incubated with 3% (v/v) hydrogen peroxide solution in PBS for 30

min at room temperature, followed by several blocking steps with serum from the species the

secondary antibody was raised in, to avoid nonspecific binding of antibodies. Sections were

incubated with the primary antibody diluted in serum blocking solution overnight at 4°C.

Negative controls were included, in which the primary antibody was omitted to identify any

nonspecific binding by the secondary antibody. Primary antibodies against CD3 (1:250

dilution; NB600-1441, Novus Biologicals), B220 (1:250 dilution; 14-0452-82, Invitrogen), CD14

(1:6000 dilution; ab221678, Abcam), CD64 (1:6000 dilution; 50086-R008, Sino Biological),

Ly6G (1:10000 dilution; 127649, Biolegend) and DBA-lectin (1:1000 dilution; B-1035, Vector)

were used in this study.

The following day, sections were washed with 0.05% Tween in PBS followed by an incubation

with an HRP-conjugated secondary antibody (1:500 dilution) in serum blocking solution for 30

min at room temperature followed by an incubation with either DAB substrate for 5 min at room

temperature (ab64238, Abcam) or Opal Polaris solution (FP1487001KT [Opal 520],

FP1488001KT [Opal 570], FP1496001KT [Opal 650], Akoya Biosciences) for 10 min at room

temperature. After staining, sections were washed and counterstained either with

Haematoxylin or DAPI (1:3000 dilution; 10236276001, Sigma-Aldrich). For chromogenic

immunohistochemistry, the sections were dehydrated in a series of alcohols and mounted on

coverslips with Pertex mounting medium (SEA-0100-00A, CellPath), while for

immunofluorescence, the sections were mounted on coverslips with Fluoromount (0100-01,

Cambridge Bioscience). Slides were imaged with a slide scanner (Axioscan 7, Zeiss) or an

inverted microscope (Axio Observer, Zeiss) and images were exported using the Zen 3.9

software (Zeiss).

Statistical analysis

Statistical analysis was performed using Graphpad prism software (version 10). Brown-

Forsythe and Welch tests were used to determine significance between experimental

conditions in data that were normally distributed. Non-parametric testing was utilised where

sample sizes were insufficient to confirm normality of data distribution or where the data was

not normally distributed; Kruskal-Wallis tests were used to assess differences between

experimental conditions. All data are presented as mean ± SD and criteria for significance is

$p < 0.05$.

<u>Results</u>

Single-cell RNA sequencing identifies distinct immune cell populations in the mouse uterus

associated with endometrial tissue repair and remodelling.

To investigate the phenotype and function of immune cells during endometrial repair, single

cell RNA sequencing (scRNAseq) analysis was performed on CD45$^+$ immune cells isolated

from uterine tissues recovered at key time points of an established mouse model of

menstruation: 24- and 48 hr post progesterone withdrawal and in age-matched controls

('cycling') (figure S1). These timepoints are representative of active tissue repair (24 hr,

'repair') and tissue remodelling (48 hr, 'remodelling'). Following quality control and filtering of

sequencing data, 13,498 CD45$^+$ immune cells were detected across 3 experimental conditions

(cycling: 5,379 cells; repair: 3,761 cells; and remodelling: 4,358 cells) and integrated for

downstream analysis. Unsupervised clustering resulted in the identification of 8

transcriptionally distinct clusters of immune cells in uterine tissues: neutrophils, eosinophils,

monocytes, macrophages, dendritic cells, NK cells, T cells and B cells (Figure 1A). Cell

clusters were annotated using both the expression of established immune cell gene markers

(Figure 1B) and analysis of the top 10 most differentially expressed genes (DEGs) per cluster

(Figure 1C & 1D).

Analysis of immune cell clusters relative to experimental condition identified distinct changes

in immune cell composition during endometrial repair and remodelling. Innate immune

populations including neutrophils, monocytes and to a lesser degree macrophages, dendritic

cells and eosinophils were predominant in the 'repair' dataset, while adaptive immune

populations such as T, B and NK cells, were largely only present in the 'cycling' and

'remodelling' data (Figure 1E & 1F). Relative abundance analysis within each experimental

condition showed changes in uterine immune cell composition in response to endometrial

repair (Figure 1G). Most notable was an increase in the proportion of neutrophils and

monocytes during active endometrial repair (24-hour) which remained elevated during the

remodelling phase (48-hour).

Unbiased gene ontology (GO) analysis was used to identify biological processes associated

with the transcriptome of each immune cell cluster in the combined dataset. Top processes

associated with subsets were consistent with their phenotypic annotation, further validating

their identification as transcriptionally and functionally discrete immune cells (Figure S2A).

Monocytes were associated with 'wound healing' (Figure S2A(v)) macrophages with

phagocytosis (Figure S2A(vi)) and neutrophils with cell chemotaxis/migration (Figure

S2A(vii)).

To explore the functional role of monocytes, macrophages and neutrophils in endometrial

repair and remodelling, the expression of gene sets associated with essential tissue repair

processes, generated from literature searches and MsigDB [32], were interrogated in

scRNAseq data (Figure S2B). Neutrophils were found to have the highest expression of genes

associated with NETosis, chemoattraction and certain remodelling enzymes while

macrophages and monocytes expressed genes associated with phagocytosis. Notably, when

assessing the expression of genes associated with 'wound healing' each cell type showed

expression of a distinct subset of genes (monocytes: *Anxa5*, *Cav1*, *F10*, *Fn1*, *Pdpn*, *Sdc1*,

Sdc4, Thbd; macrophages: *Apoe*, *Axl*, *C3*, *Ccr2*, *Gnas*; neutrophils: *Slc11a1, Cd9, Lrg1, Plaur,*

S100a9) (Figure S3B). Taken together, GO analyses suggest that myeloid cells are

predominant during endometrial tissue repair and may have key role in wound healing

following menstruation.

Flow cytometry analysis confirms dynamic changes in immune cell composition and the

predominance of myeloid cells during endometrial repair.

To validate putative repair-associated changes in uterine immune cell composition identified

by *in silico* analyses, flow cytometry was performed on uterine tissues collected over a

comprehensive time course of endometrial repair using the mouse model of menstruation

(figure S1). A flow cytometry panel was designed to interrogate the immune cell populations

identified by scRNAseq using phenotypic markers as outlined in Supplementary Table 1.

Using the gating strategy outlined in figure 2A, flow cytometry analysis corroborated the

immune cell composition identified by scRNAseq analysis in cycling, repair (24 hr) and

remodelling (48 hr) tissues. This was complemented by analysis of additional time points

which provided resolution on immune cell dynamics prior to and during tissue breakdown

(Figure 2B & 2C). At 0 hr (prior to breakdown), there was an increase in the proportion of

monocytes within the uterus and a decrease in the proportion of NK cells and dendritic cells

(DCs) when compared to cycling control. At 12 hr during tissue breakdown there was a large

compositional shift characterised by a significant increase in the proportion of neutrophils in

the uterus, accounting for 87.82% ± 14.06 of all CD45$^+$ immune cells at this timepoint. The

proportion of neutrophils peaked at 24 hr (91.86% ± 2.60 of CD45$^+$ cells) during repair and

neutrophils were also the most abundant cell type at 48 hr (55.7% ± 12.72 of CD45$^+$ cells),

with the numbers of T cells, NK cells and macrophages also increasing at 48 hr (Figure 2B &

2C).

Lymphocyte populations are infrequent during endometrial repair.

T cells (CD45$^+$ Ly6G$^-$ NK1.1$^-$ CD19$^-$ CD3$^+$) were quantified in uterine tissue digests using flow

cytometry (Figure 3A). Both the relative abundance (percentage of CD45$^+$ cells) and absolute

numbers of T cells were significantly increased during tissue remodelling (48 hr; 8.52% ± 4.71

and 47.87 count/mg ± 27.80) when compared to earlier phases of breakdown (12 hr; 1.72% ±

3.84 and 8.06 count/mg ± 9.67) and repair (24 hr; 0.19% ± 0.13 and 11.77 count/mg ± 13.23)

(Figure 3B & 3C). IHC was performed on tissue samples from matching time points/conditions.

CD3$^+$ T cells were detected in the endometrium and myometrium of both 24- and 48-hour

tissues (Figure 3D; black arrows & Figure S4A). T cells were more readily detected in 48-hour

tissues but there was no spatially distinct distribution associated with different time-points

(Figure 3D; black arrows & Figure S4A).

NK cells (CD45$^+$ Ly6G$^-$ CD3$^-$ CD19$^-$ NK1.1$^+$) were more readily detected (Figure 3E) and

abundant in cycling mice (62.35 count/mg ± 120.30), numbers were lowest during breakdown

(12 hr; 9.069 count/mg ± 23.66) and repair (24 hr; 26.44 count/mg ± 30.64) but more abundant

in remodelling tissues (48 hr; 43.35 count/mg ± 32.52) (Figure 3G). Detection of NK cells using

DBA-lectin in uterine tissue sections located NK cells to the decidualised tissue prior to

breakdown (0-hour), in close proximity to the luminal epithelium and adjacent to the decidual-

endometrial boundary. In 24-hour tissues NK cells were infrequent, located predominantly in

the shed decidual tissue (Figure 3H) and in 48-hour tissues very few NK cells could be

detected (Figure S4B).

Quantitative analysis of B cells (CD45$^+$ Ly6G$^-$ CD3$^-$ NK1.1$^-$ CD19$^+$) demonstrated that B cells

are abundant in the uterus prior to breakdown (0 hr; 38.66 count/mg ± 24.97) but significantly

decreased thereafter (Figure 3J & 3K). Detection of B220$^+$ B cells in uterine tissue sections by

IHC showed B cells located in both the endometrium and myometrium prior to breakdown

(Figure 3L; 0 hr, black arrows) but they were infrequent during repair and remodelling (Figure

3L; black arrows & Figure S4C).

Myeloid cell populations are dynamically regulated during endometrial breakdown, repair, and

remodelling.

Flow cytometry and immunohistochemistry analyses demonstrated changes in the abundance

and location of key myeloid cell types in the uterus during endometrial repair. Monocytes

(Ly6C$^+$ MHCII$^-$; Figure 4A) were the most abundant immune cell present in the uterus prior to

breakdown (0-hour; 103.8 ± 50.46) (Figure 4B & 4C). The number of monocytes in the uterus

was then significantly decreased during breakdown when compared to prior to breakdown (0

hr; 103.80 count/mg ± 50.46 vs 12 hr; 25.40 count/mg ± 54.57, $p < 0.0001$). Monocytes were

significantly increased during repair (12 hr; 25.40 count/mg ± 54.57 vs 24 hr; 101.1 count/mg

± 49.67, $p < 0.0001$) and significantly decreased during remodelling (24 hr; 101.1 count/mg ±

49.67 vs 48 hr; 10.44 count/mg ± 8.27, $p = 0.0018$) (Figure 4C). Immunofluorescent staining

was performed using an anti-CD14 antibody to investigate spatial distribution of monocytes in

uterine tissues. CD14$^+$ cells were predominantly localised to the boundary between the

underlying endometrium and the detached decidual tissue during breakdown (12 hr; Figure

S6A) and subsequent repair (24 hr; Figure 4J and Figure S6A).

Macrophages counts were significantly decreased during tissue breakdown when compared

to tissue repair (12 hr; 4.71 count/mg ± 8.17 vs 24 hr; 21.78 count/mg ± 26.47, $p = 0.0013$)

and remodelling time points (12 hr; 4.71 count/mg ± 8.17 vs 48 hr; 16.61 count/mg ± 10.69, p

< 0.0001) (Figure 4F). Macrophages (F4/80$^+$CD64$^+$; Figure 4D) were decreased during

endometrial repair but increased in the remodelling timepoint (24 hr; 0.36% ± 0.28 vs 48 hr;

2.80% ± 1.22, $p = 0.004$) (Figure 4E). CD64$^+$ macrophages were detected by

immunohistochemistry in all uterine tissue compartments throughout breakdown and repair

(0-24 hr; Figure S6B), in both the decidualised and non-decidualised endometrium, as well as

the myometrium. In 48-hour tissues, macrophages were most abundant in the stroma of the

remodelled endometrium (Figure 4K and Figure S6B).

Neutrophils (Ly6G$^+$SiglecF$^-$, Figure 4G) were the most abundant immune cell type present in

the uterus during breakdown (12 hr; 87.82% ± 14.06), repair (24 hr; 91.86% ± 2.60), and

remodelling (48 hr; 55.71% ± 12.72) (Figure 4H). Neutrophils counts in the uterus were

significantly increased during breakdown compared to cycling controls (12 hr; 1374 count/mg

± 2398 vs cycling; 56.25 count/mg ± 175.80, p < 0.0001). Neutrophil abundance peaked during

endometrial repair (24 hr; 4946 count/mg ± 2344) and although neutrophil numbers were

relatively decreased during remodelling they were still significantly increased compared to

cycling controls (Figure 4I; cycling; 56.25 count/mg ± 175.80 vs 48 hr; 504.5 count/mg ± 570.6,

p = 0.016). Immunofluorescent staining identified abundant Ly6G$^+$ neutrophils in uterine

tissues during breakdown (12 hr) and repair (24 hr) specifically localised to the boundary

between the detached decidual tissue and the underlying endometrium (Figure S6C). In 48-

hour tissues, Ly6G$^+$ neutrophils were readily detected throughout the remodelled

endometrium (Figure S6C).

Dendritic cells (DCs; CD11c$^+$MHCII$^+$; Figure S5A) were infrequent during tissue breakdown

(12 hr; 5.63 count/mg ± 11.91) but were modestly increased during repair (24 hr; 15.80

count/mg ± 13.51) and remodelling (48 hr; 6.59 count/mg ± 6.019) (Figure S5B & S5C).

Eosinophils (Ly6G$^-$ SiglecF$^+$; Figure S5D) were infrequent in cycling tissues (3.26 count/mg ±

6.22) and during phases of tissue breakdown (12 hr; 2.14 count/mg ± 5.08) and repair (24 hr;

2.43 count/mg ± 4.20). Notably, eosinophils were significantly increased during remodelling

(48 hr; 17.60 count/mg ± 20.43) but were infrequent overall (Figure S5F). Due to their low

abundance, DCs and eosinophils were not analysed by IHC.

Monocytes, neutrophils, and macrophages co-localise to areas of breakdown and tissue

repair.

To understand the distribution of key myeloid cell types relative to each other in uterine tissue

sections, multiplex IHC for the detection of CD14 (monocytes), CD64 (macrophages) and

Ly6G (neutrophils) was performed using uterine tissues collected from across the time course

(Figure 4J-K, Figure S6D). Prior to tissue breakdown (0 hr), relatively few CD14$^+$ or Ly6G$^+$

cells could be detected but CD64$^+$ macrophages were abundant throughout the tissue (Figure S6D). At 12 hr, during tissue breakdown, CD14$^+$ monocytes and Ly6G$^+$ neutrophils were detected clustered within the same regions of the tissue and were localised to regions of shedding, where decidualised tissue was detaching, as well as in the underlying residual endometrium (Figure S5D). CD64$^+$ macrophages were located predominantly in the outer myometrial layers at 12 hr and only diffusely detected within regions of tissue breakdown (Figure S6D).

At 24 hr, CD14$^+$ monocytes and Ly6G$^+$ neutrophils were identified in the detaching decidual tissue but were also detected in regions of underlying endometrium undergoing tissue repair (Figure 4J & S6D). CD64$^+$ macrophages were also detected throughout the repairing endometrium as well as in the decidual tissue, in association with regions where detachment had not yet taken place (Figure 4J & S6D). At this time point, monocytes, neutrophils and macrophages were detected within regions of active tissue repair but were less frequent in areas of intact tissue. At 48 hr the decidual tissue is almost completely expelled, the luminal epithelium is largely restored, and the underlying stroma is undergoing remodelling. CD14$^+$ monocytes were less frequent at 48 hr and localised only to regions of decidual detachment on the endometrial surface, while Ly6G$^+$ neutrophils were abundant and dispersed throughout the remodelling endometrium (Figure 4K & S6D). Notably, CD64$^+$ macrophages were more abundant at 48 hr with an apparent expansion throughout the remodelling endometrium in regions both proximal and distal to sites of recent tissue repair (Figure 4K & S6D).

CellChat analysis of scRNAseq data identifies monocyte, macrophage and neutrophil signalling pathways associated with endometrial repair.

Given the association of monocytes, macrophages and neutrophils with tissue repair and remodelling in the endometrium, further *in silico* analyses were performed to infer putative signalling pathways and interactions associated with these cell types. CellChat analysis of ligand-receptor expression was performed on scRNAseq data, and immune cell interactions were assessed. Monocytes, macrophages and neutrophils were identified as both dominant

senders (high ligand expression) and dominant receivers (high receptor expression) in the

communication network, scoring highly in both outgoing and incoming interaction strength

(Figure 5A). The top signalling pathways found to contribute to outgoing/incoming signalling

across all immune cell types were CCL, CXCL, MIF, FN1, SPP1 and THBS (Figure 5B).

Analysis of communication patterns between cell types for these signalling pathways

highlighted the predominance of monocytes (CCL, CXCL, MIF, FN1, SPP1 and THBS),

macrophages (CCL, MIF and FN1) and neutrophils (CCL, CXCL, and THBS) as key

contributors to cell communication of repair-associated signalling pathways (Figure 5C).

The relationship between specific ligand-receptor pairs and immune cell subsets is

summarised in the chord diagrams for each signalling pathway, where the width of each chord

corresponds to the strength of the interaction (Figure 5D). The relationship between different

CCL chemokine ligands and corresponding receptors showed a predominance of ligand

expression associated with monocytes (*Ccl2*, *Ccl3*, *Ccl4*, *Ccl6* and *Ccl9*), macrophages (*Ccl2*,

Ccl3, *Ccl4*, *Ccl6 and Ccl9*) and neutrophils (*Ccl3*, *Ccl6* and *Ccl9*) (Figure 5Di). Crosstalk in

CXCL pathway was predominantly from monocytes and neutrophils signalling *Cxcl2* and *Cxcl3*

to *Cxcr2* on neutrophils and eosinophils (Figure 5Dii). *Mif* and *Fn1* signalled from monocytes

and macrophages to corresponding receptors on multiple cell types (Figure 5Diii and iv). *Spp1*

ligand was predominantly associated with monocytes signalling to distinct reciprocal receptors

including *Cd44* on neutrophils and integrins on monocytes and macrophages(Figure 5Dv).

Similarly, *Thbs1* signalling was predominantly from monocytes and neutrophils which

signalled via *Cd47* on all immune cell types in the dataset (Figure 5Dv and vi).

To examine the association of these signalling pathways more discretely, the top overall

signalling pathways were assessed for each experimental condition (Figure 5E). While similar

signalling pathways ranked highly, their association with specific immune cell subsets shifted

according to experimental conditions. The relative strength of THBS signalling was strongest

for neutrophils in cycling, but more strongly associated with monocytes in repair and

remodelling tissues. MIF signalling was strongly associated with DCs in cycling, but more

strongly associated with monocytes in repair, and with macrophages in remodelling tissues.

Notably, FN1 and SPP1 signalling was strongly associated with monocytes in all conditions.

Due to their strength of association with top signalling pathways, monocytes, macrophages

and neutrophils were extracted for further communication analyses (Figure 5F). Pairwise

comparisons between experimental conditions (cycling vs repair, cycling vs remodelling,

repair vs remodelling) revealed changes in THBS1, SPP1, MIF, CCL and CXCL signalling

pathways in monocytes, macrophages and neutrophils relative to phases of endometrial repair

and remodelling (Figure 5F). The greatest interaction strength for incoming and outgoing

signals in monocytes was associated with THBS1 in repair and SPP1 in remodelling.

Macrophage signalling also differed by condition with greatest signalling strength for THBS1

associated with repair while MIF and SPP1 were associated with remodelling. In each instance

these signalling pathways were associated with a large differential signal strength for both

incoming and outgoing interactions. Neutrophil signalling was also associated with THBS1

and SPP1 in repair and remodelling respectively. CCL signalling strength was reduced in

monocytes and macrophages but increased in neutrophils in remodelling. CXCL signalling

was strongly associated with neutrophils and both outgoing and incoming signalling strength

was upregulated during repair (24 hr) and remodelling (48 hr). Monocytes, macrophages and

neutrophils were strongly associated with THBS1 signalling during endometrial repair and

SPP1 and MIF signalling during remodelling. Chemokine signalling was distinctly associated

with neutrophils during remodelling consistent with roles in immune cell crosstalk and

recruitment at this time point.

Discussion

In this study, we combined a mouse model of menstruation with comprehensive immune cell profiling to characterise the contribution of distinct immune subsets to endometrial repair. We found that uterine immune cell composition was dynamic, with pronounced temporal and spatial changes in monocytes, macrophages and neutrophils which were associated with regions of tissue repair and remodelling. These data suggest that monocyte and neutrophil infiltration is a key feature of endometrial repair, typical of early inflammation responses observed in other tissue contexts[33, 34]. Neutrophils, typically associated with acute inflammatory responses, were abundant during tissue repair as well as remodelling, while macrophages, which regulate tissue repair processes, were relatively decreased during active repair. Myeloid cells were enriched for both pro- and anti-inflammatory signalling pathways, highlighting the discrete balance of immune cell functions that support this unique repair process. Utilising scRNAseq data we were able to identify key signalling pathways associated with myeloid cells during menstruation which included THBS1 and SPP1 as important mediators of endometrial repair. Collectively, these data provide new insights into the immunology of menstruation and highlight specific spatiotemporal roles for myeloid cells during endometrial repair.

We found that monocytes and macrophages were transcriptionally similar with overlapping expression of markers such as *Csf1r*, *Cd68*, *Msr1* and *Fn1*. However, *Inha* and *Cxl3* expression was distinct to monocytes consistent with potential roles in regulating migration and adhesion of inflammatory cells to sites of tissue repair [35]. GOBP analysis highlighted enrichment for genes associated with wound healing and endoplasmic reticulum stress. Monocytes expressed apoptosis- and phagocytosis-associated genes as well as a subset of wound healing genes, such as *Pdpn* and *Cav1*, that were distinct from those expressed by macrophages and neutrophils. Specifically, *Cav1* expression has been shown to increase during monocyte adhesion and drives monocyte-to-macrophage differentiation [36].

CD14+ monocytes were located in both the decidualised tissue and underlying endometrium

and were detected in close association with macrophages and neutrophils. Cell chat analysis

identified monocytes as major drivers of intercellular communication in the uterus with the

highest incoming and outgoing signalling strength amongst immune cells, followed by

macrophages and neutrophils. Monocytes were associated with signalling via CCL, MIF,

SPP1 and THBS and were the predominant outgoing signal source for these signalling

pathways. Signalling via THBS and SPP1 was strongly associated with monocytes from repair

and remodelling tissues, respectively.

SPP1 (Secreted Phosphoprotein 1, also known as Osteopontin) plays multiple roles in tissue

repair and is reported to modulate inflammation and tissue remodelling. SPP1 mediates cell-

to-cell and cell-to-matrix interactions, via integrins and CD44, which can promote macrophage

migration and effector functions[37]. SPP1 also increases expression of S100A9, IL-1β, and

chemokines like CCL2, CCL3 and CCL4 which skews monocytes toward a pro-inflammatory

phenotype[38]. Notably, these genes were enriched in the transcriptome of uterine monocytes

during menstruation in the current study. Thrombospondin 1 (TSP-1) promotes cell migration

and tissue remodelling via TGF-β and integrin signalling which may support tissue repair in

the endometrium [39] [40]. However, it can also inhibit repair by suppressing angiogenesis, a

crucial component of endometrial repair [41]. TSP-1 signalling was enriched in monocytes

from repair tissues, but the signal strength was relatively decreased in remodelling tissues,

when vascular remodelling is more prominent, highlighting the discrete and dynamic

contributions of different inflammatory cells and mediators to endometrial repair during

menstruation.

We have previously shown that distinct subsets of macrophages associate with regions

endometrial repair during menstruation [16] but the function/phenotype of uterine

macrophages has not been elucidated. In the current study, we found that macrophages were

enriched for signalling pathways important for tissue repair and remodelling. This included

complement pathway which has roles in clearing cellular debris, recruiting immune cells to

sites of injury, releasing growth factors and regulating blood clotting in tissue repair processes.

Differential gene expression analysis highlighted that uterine macrophages distinctly

expressed key components of the complement pathway, including *C1qa*, which is a marker of

tissue resident macrophages in other contexts [42]. Macrophages were also enriched for

genes involved in galectin signalling indicative of a pro-resolving phenotype [43]. Similar to

monocytes, macrophages were also associated with MIF and FN1 signalling networks which

were key pathways of cell-cell communication. The transcriptomic similarities between

macrophages and monocytes are likely representative of linked differentiation trajectories

within uterine tissues. Although the ontogeny and kinetics of monocytes and macrophages in

the uterus are yet to be fully defined, studies using tissue explants, or organ transplants,

suggest the uterine macrophage niche is maintained via monocyte-driven replenishment [44,

45]. This is consistent with the compositional changes in macrophages and monocytes

described in the current study, where monocytes are predominant during repair, but

macrophages, likely monocyte-derived, are more abundant than monocytes during

remodelling.

ScRNAseq data analysis confirmed that neutrophils were a major source of chemokines as

they were enriched for both CCL and CXCL-mediated signalling pathways, indicating

regulation of immune cell recruitment within the repairing endometrium. Neutrophils were

associated with pro-inflammatory signalling via TNF pathways, which, interestingly, was

prominent during remodelling rather than repair. While TNF is a classic pro-inflammatory

mediator, Annexin1 signalling was also enriched in neutrophils at this timepoint. Annexin1 has

anti-inflammatory functions by downregulating neutrophil activation [46, 47]. The neutrophil

transcriptome profile was consistently associated with both pro- and anti-inflammatory

pathways which may suggest neutrophil phenotype is temporally altered in the uterine tissue

environment across repair and remodelling, with a switch to anti-inflammatory functions during

the latter phase. Assessing differentially expressed genes in neutrophils highlighted that in

addition to classic neutrophils markers such as *S100a8*, *S100a9* and *Cxcl2*, uterine

neutrophils also expressed *Dusp1* and *Acod1* indicating an anti-inflammatory phenotype.

Notably, Dusp1 (Dual Specificity Phosphatase 1) has key roles in resolution of inflammation

by deactivating pro-inflammatory mitogen-activated protein kinase (MAPK)-dependent

pathways and influencing production of neutrophil chemoattractants like Cxcl1[48]. This may

reflect a functional requirement for restraining classic effector functions of neutrophils within

the uterus in the context of endometrial repair. Expression of Acod1 (Aconitate Decarboxylase

1) in neutrophils regulates inflammation via production of itaconate. This shift in

immunometabolism is associated with anti-inflammatory protein production, protecting cells

from oxidative stress as well as regulation of NET formation and neutrophil migration [49, 50].

Thus, although neutrophils are abundant during menstruation, their transcriptomic phenotype

supports roles in coordinating immune responses via chemokine secretion and promoting

resolution of inflammation through suppression of pro-inflammatory signalling pathways.

Transcriptome profiling in the current study supports reparative functions for neutrophils during

menstruation consistent with emerging roles for neutrophils in contributing to tissue repair in

other contexts [51-53]. Indeed, Kaitu'u-Lino et al. previously demonstrated the importance of

monocytes and macrophages to endometrial breakdown and repair using antibody-based

depletion (RB6-8C5) of Gr-1 positive cells in a mouse model of menstruation [54].

Our findings using a mouse model of simulated menstruation extend and complement those

from analysis of human endometrial tissues. Armstrong et al., investigated mRNA expression

levels of inflammatory cytokines and chemokines during menstruation in the human

endometrium and found increased expression of CCL2, CXCL8, IL6 and TNF during the

menstrual phase of the cycle[8]. Importantly, the current study provides much needed

resolution to the specific cellular sources of these mediators which may be beneficial to

translational studies aiming to understand the pathophysiology of menstrual dysfunction in

reproductive health disorders. Dysregulated inflammatory responses are a common feature of

reproductive health disorders such as heavy menstrual bleeding (HMB) and endometriosis,

with changes to macrophage and neutrophil function reported[55]. While studies

characterising specific phenotypes of immune cell subsets during menstruation are lacking,

changes in expression to key mediators described in the current study have been reported.

For example, TSP-1 expression is reported to be decreased in endometrial tissues from

women with HMB [41] while overexpression of SPP1 has been reported in macrophages from

women with endometriosis[56]. Further research will be needed to better characterise immune

dysfunction in human reproductive disorders, but the comprehensive phenotyping performed

in the current study provides tractable targets for further translational investigation.

<u>Conclusion</u>

Here, we have comprehensively profiled immune cells using a combination of scRNAseq, flow

cytometry and multiplex immunohistochemistry and identified that monocytes, macrophage

and neutrophils are spatially and temporally regulated to support endometrial repair. These

data provide a more complete understanding of how different immune cell populations

contribute to endometrial tissue repair and identify key cell-specific mediators and pathways.

Future studies should assess the importance of myeloid cells to endometrial repair through

lineage tracing, cell ablation and comparison to human datasets. Improved understanding of

endometrial repair will, in turn, support identification of potential pathways that are

dysregulated in reproductive health disorders in women.

<u>References</u>

1.   Critchley, H.O., et al., *The endocrinology of menstruation--a role for the immune*
556   *system.* Clin Endocrinol (Oxf), 2001. **55**(6): p. 701-10.
557   2.   Critchley, H.O.D., et al., *Physiology of the Endometrium and Regulation of*
558   *Menstruation.* Physiol Rev, 2020. **100**(3): p. 1149-1179.
559   3.   Kirkwood, P.M., I.W. Shaw, and P.T.K. Saunders, *Mechanisms of Scarless Repair at*
560   *Time of Menstruation: Insights From Mouse Models.* Front Reprod Health, 2021. **3**: p.
561   801843.
562   4.   Maybin, J.A. and H.O. Critchley, *Steroid regulation of menstrual bleeding and*
563   *endometrial repair.* Rev Endocr Metab Disord, 2012. **13**(4): p. 253-63.
564   5.   Hu, X., et al., *Cyclical endometrial repair and regeneration: Molecular mechanisms,*
565   *diseases, and therapeutic interventions.* MedComm (2020), 2023. **4**(6): p. e425.
566   6.   Lee, S.K., et al., *Immune cells in the female reproductive tract.* Immune Netw, 2015.
567   **15**(1): p. 16-26.
568   7.   Lathbury, L.J. and L.A. Salamonsen, *In-vitro studies of the potential role of neutrophils*
569   *in the process of menstruation.* Mol Hum Reprod, 2000. **6**(10): p. 899-906.
570   8.   Armstrong, G.M., et al., *Endometrial apoptosis and neutrophil infiltration during*
571   *menstruation exhibits spatial and temporal dynamics that are recapitulated in a mouse*
572   *model.* Sci Rep, 2017. **7**(1): p. 17416.
573   9.   White, H.D., et al., *CD3+ CD8+ CTL activity within the human female reproductive*
574   *tract: influence of stage of the menstrual cycle and menopause.* J Immunol, 1997.
575   **158**(6): p. 3017-27.
576   10.   Chen, C.K., et al., *Increased expressions of CD69 and HLA-DR but not of CD25 or*
577   *CD71 on endometrial T lymphocytes of nonpregnant women.* Hum Immunol, 1995.
578   **42**(3): p. 227-32.
579   11.   Flynn, L., et al., *Menstrual cycle dependent fluctuations in NK and T-lymphocyte*
580   *subsets from non-pregnant human endometrium.* Am J Reprod Immunol, 2000. **43**(4):
581   p. 209-17.
582   12.   Starkey, P.M., L.M. Clover, and M.C. Rees, *Variation during the menstrual cycle of*
583   *immune cell populations in human endometrium.* Eur J Obstet Gynecol Reprod Biol,
584   1991. **39**(3): p. 203-7.
585   13.   Chen, K., et al., *Single-cell transcriptomic analysis of immune cell dynamics in the*
586   *healthy human endometrium.* Biochem Biophys Rep, 2024. **39**: p. 101802.
587   14.   Wang, W., et al., *Single-cell transcriptomic atlas of the human endometrium during the*
588   *menstrual cycle.* Nat Med, 2020. **26**(10): p. 1644-1653.
589   15.   Cousins, F.L., et al., *Evidence from a mouse model that epithelial cell migration and*
590   *mesenchymal-epithelial transition contribute to rapid restoration of uterine tissue*
591   *integrity during menstruation.* PLoS One, 2014. **9**(1): p. e86378.
592   16.   Cousins, F.L., et al., *Evidence for a dynamic role for mononuclear phagocytes during*
593   *endometrial repair and remodelling.* Sci Rep, 2016. **6**(1): p. 36748.
594   17.   Salamonsen, L.A., et al., *Regulation of matrix metalloproteinases in human*
595   *endometrium.* Hum Reprod, 2000. **15 Suppl 3**: p. 112-9.
596   18.   Thiruchelvam, U., et al., *The importance of the macrophage within the human*
597   *endometrium.* J Leukoc Biol, 2013. **93**(2): p. 217-25.
598   19.   Tabibzadeh, S. and X.Z. Sun, *Cytokine expression in human endometrium throughout*
599   *the menstrual cycle.* Hum Reprod, 1992. **7**(9): p. 1214-21.
600   20.   Jones, R.L., et al., *Identification of chemokines important for leukocyte recruitment to*
601   *the human endometrium at the times of embryo implantation and menstruation.* J Clin
602   Endocrinol Metab, 2004. **89**(12): p. 6155-67.

21. Crona Guterstam, Y., et al., *The cytokine profile of menstrual blood.* Acta Obstet
Gynecol Scand, 2021. **100**(2): p. 339-346.
22. Hampton, A.L. and L.A. Salamonsen, *Expression of messenger ribonucleic acid*
encoding matrix metalloproteinases and their tissue inhibitors is related to
menstruation. J Endocrinol, 1994. **141**(1): p. R1-3.
23. Sharkey, A.M., et al., *Vascular endothelial growth factor expression in human*
endometrium is regulated by hypoxia. J Clin Endocrinol Metab, 2000. **85**(1): p. 402-9.
24. Kirkwood, P.M., et al., *Single-cell RNA sequencing redefines the mesenchymal cell*
landscape of mouse endometrium. FASEB J, 2021. **35**(4): p. e21285.
25. Kirkwood, P.M., et al., *Single-cell RNA sequencing and lineage tracing confirm*
mesenchyme to epithelial transformation (MET) contributes to repair of the
endometrium at menstruation. Elife, 2022. **11**.
26. Stuart, T. and R. Satija, *Integrative single-cell analysis.* Nat Rev Genet, 2019. **20**(5): p.
257-272.
27. Wolock, S.L., R. Lopez, and A.M. Klein, *Scrublet: Computational Identification of Cell*
Doublets in Single-Cell Transcriptomic Data. Cell Syst, 2019. **8**(4): p. 281-291 e9.
28. Core, H.C.B. *Elbow plot: Quantitative approach*. Introduction to Single-cell RNA-seq
- ARCHIVED. ; Available from: https://hbctraining.github.io/scRNA-
seq/lessons/elbow_plot_metric.html.
29. Yu, G., et al., *clusterProfiler: an R package for comparing biological themes among*
gene clusters. OMICS, 2012. **16**(5): p. 284-7.
30. Jin, S., et al., *Inference and analysis of cell-cell communication using CellChat.* Nat
Commun, 2021. **12**(1): p. 1088.
31. Edgar, R., M. Domrachev, and A.E. Lash, *Gene Expression Omnibus: NCBI gene*
expression and hybridization array data repository. Nucleic Acids Res, 2002. **30**(1): p.
207-10.
32. Liberzon, A., et al., *The Molecular Signatures Database (MSigDB) hallmark gene set*
collection. Cell Syst, 2015. **1**(6): p. 417-425.
33. Kolaczkowska, E. and P. Kubes, *Neutrophil recruitment and function in health and*
inflammation. Nat Rev Immunol, 2013. **13**(3): p. 159-75.
34. Serbina, N.V., et al., *Monocyte-mediated defense against microbial pathogens.* Annu
Rev Immunol, 2008. **26**(Volume 26, 2008): p. 421-52.
35. Petraglia, F., et al., *Inhibin and activin modulate human monocyte chemotaxis and*
human lymphocyte interferon-gamma production. J Clin Endocrinol Metab, 1991.
72(2): p. 496-502.
36. Fu, Y., et al., *Caveolin-1 plays a critical role in the differentiation of monocytes into*
macrophages. Arterioscler Thromb Vasc Biol, 2012. **32**(9): p. e117-25.
37. Kitano, T., et al., *Transcriptome Analysis Identified SPP1-Positive Monocytes as a Key*
in Extracellular Matrix Formation in Thrombi. J Am Heart Assoc, 2025. **14**(19): p.
e044299.
38. MacDonald, L., et al., *COVID-19 and RA share an SPP1 myeloid pathway that drives*
PD-L1+ neutrophils and CD14+ monocytes. JCI Insight, 2021. **6**(13).
39. Wilson, Z.S., et al., *Critical role of thrombospondin-1 in promoting intestinal mucosal*
wound repair. JCI Insight, 2024. **9**(17).
40. Streit, M., et al., *Thrombospondin-1 suppresses wound healing and granulation tissue*
formation in the skin of transgenic mice. EMBO J, 2000. **19**(13): p. 3272-82.
41. Rae, M., et al., *Cortisol inactivation by 11beta-hydroxysteroid dehydrogenase-2 may*
enhance endometrial angiogenesis via reduced thrombospondin-1 in heavy
menstruation. J Clin Endocrinol Metab, 2009. **94**(4): p. 1443-50.

42. Horowitz, A., et al., *C1QA is an invariant biomarker for tissue macrophages.* bioRxiv, 2024.

43. Yaseen, H., et al., *Galectin-1 Facilitates Macrophage Reprogramming and Resolution of Inflammation Through IFN-beta.* Front Pharmacol, 2020. **11**: p. 901.

44. Bister, J., et al., *Tissue-specific nonheritable influences drive endometrial immune system variation.* Sci Immunol, 2024. **9**(94): p. eadj7168.

45. Guo, Y., et al., *Comprehensive analysis of leukocytes, vascularization and matrix metalloproteinases in human menstrual xenograft model.* PLoS One, 2011. **6**(2): p. e16840.

46. Sugimoto, M.A., et al., *Annexin A1 and the Resolution of Inflammation: Modulation of Neutrophil Recruitment, Apoptosis, and Clearance.* J Immunol Res, 2016. **2016**(1): p. 8239258.

47. Guido, B.C., et al., *Annexin-A1 peptide down-regulates the leukocyte recruitment and up-regulates interleukin-10 release into lung after intestinal ischemia-reperfusion in mice.* J Inflamm (Lond), 2013. **10**(1): p. 10.

48. Hoppstadter, J. and A.J. Ammit, *Role of Dual-Specificity Phosphatase 1 in Glucocorticoid-Driven Anti-inflammatory Responses.* Front Immunol, 2019. **10**: p. 1446.

49. Ji, J., et al., *IRG1/ACOD1 promotes neutrophil reverse migration and alleviates local inflammation.* J Leukoc Biol, 2024. **116**(4): p. 854-863.

50. Zhao, N., et al., *Neutrophil Irg1/itaconate axis protects against experimental colitis by suppressing local inflammation and maintaining hematopoietic homeostasis.* Mol Biomed, 2025. **6**(1): p. 143.

51. Rizo-Tellez, S.A. and J.G. Filep, *Beyond host defense and tissue injury: the emerging role of neutrophils in tissue repair.* Am J Physiol Cell Physiol, 2024. **326**(3): p. C661-C683.

52. Peiseler, M. and P. Kubes, *More friend than foe: the emerging role of neutrophils in tissue repair.* J Clin Invest, 2019. **129**(7): p. 2629-2639.

53. Loh, W. and S. Vermeren, *Anti-Inflammatory Neutrophil Functions in the Resolution of Inflammation and Tissue Repair.* Cells, 2022. **11**(24).

54. Kaitu'u-Lino, T.J., N.B. Morison, and L.A. Salamonsen, *Neutrophil depletion retards endometrial repair in a mouse model.* Cell Tissue Res, 2007. **328**(1): p. 197-206.

55. Wilson, T.R., S. Kasper, and K.A. Burns, *An emerging role for neutrophils in the pathogenesis of endometriosis.* 2025. **3**(1): p. 9.

56. Zhang, S., et al., *SPP1 as a key modulator of M2 macrophage polarization promotes endometriosis progression via activation of the FAK/PI3K/AKT pathway: A bioinformatics and experimental study.* Int Immunopharmacol, 2025. **166**: p. 115563.

Acknowledgments

General

We are sincerely grateful to the members of the SURF imaging suite at the University of

Edinburgh and the Veterinary and other staff in Biological Services for their assistance. Flow

cytometry data were generated with support from the IRR Flow Cytometry and cell sorting

facility, University of Edinburgh.

For the purpose of open access, the author has applied a Creative Commons Attribution (CC

BY) licence to any Author Accepted Manuscript version arising from this submission.

Funding

This work was supported by the Wellcome Trust (Fellowship 220656/Z/20/Z to DAG), and the

Medical Research Council studentship (MR/W006804/1; RJA/DAG).

Author contributions

Conceptualization: DAG

Methodology: RJA, PMK, HL, SG, IS, DAG

Analysis: RJA, PMK, HL, SG, IS, DAG

Funding acquisition: DAG

Writing – original draft: RJA, PMK, IS, DAG

Writing – review & editing: RJA, PMK, PTKS, IS, DAG

Data and materials

All data are available in the main text or the supplementary materials

Figures

Figure 1: Single cell RNA sequencing analysis identifies 8 uterine immune cell

populations in a mouse model of endometrial repair.

(A) Uniform manifold approximation and projection (UMAP) visualisation: population restricted

CD45$^+$ immune cells isolated from mouse uterine tissue samples taken from steady state

control mice (cycling, n=8) and menstruating mice 24- (endometrial repair, n=4) and 48 hr

(endometrial remodelling, n=4) after progesterone withdrawal. **(B)** Dot plot: Expression of

canonical immune cell markers: neutrophils (*S100a8*, *S100a9*, *Cxcr2*), eosinophils (*SiglecF*,

Hic1), monocytes (*Cd14*), macrophages (*Csfr1, Cd68, Adgre1*, *Fcgr1, Mertk*), dendritic cells

(*H2-Ab1, Tmem123, Xcr1, Cd209a, Clec9a, Clec10a*), natural killer cells (*Gzmb, Klrb1c,*

Ncr1), T cells (*Cd3e, Gata3*), and B cells (*Cd19, Ighd*). Dot size: percentage of cells in each

cluster expressing the gene; Dot colour: average expression per cluster from low (lavender)

to high (dark blue). **(C)** Violin plot: Expression of the top 3 differentially expressed genes in

each cell cluster displayed as log (TPM$^+$1). TPM= transcript count per million. **(D)** Heatmap

(yellow, high; purple, low): Top 10 differentially expressed genes in each cell cluster (logFC >

0.5, p-value <0.05, Wilcoxon rank-sum test). Top axis is colour coded and named by immune

cell type. **(E)** Feature plot: Distribution of cells from each experimental condition. **(F)** Bar plot:

Summary of immune cell origin across experimental conditions. Each colour represents each

experimental condition, and the y-axis denotes the contribution of each experimental condition

to each immune cell cluster. **(G)** Stacked bar plot: Single cell RNA sequencing analysis of the

relative abundance of uterine immune cell populations during endometrial repair. Each colour

represents a distinct immune cell type, and the y-axis denotes the percentage of total CD45$^+$

immune cells.

B) **Flow cytometry – Relative abundance of CD45+ cells**

C) **Flow cytometry – Relative abundance of CD45+ cells**

Cycling (Steady state)	0hrs (Prior to breakdown)	12hrs (Breakdown)	24hrs (Repair)	48hrs (Remodel)	Cluster
8.627	8.458	1.310	0.107	0.161	B cells
5.477	3.266	1.410	0.189	8.519	T cells
16.326	5.626	0.316	0.415	6.468	NK cells
7.235	0.648	0.277	0.269	0.996	Dendritic cells
4.114	2.074	0.351	0.361	2.801	Macrophages
6.477	24.100	1.248	1.976	1.477	Monocytes
0.680	0.075	0.198	0.034	1.151	Eosinophils
15.788	15.776	87.818	91.856	55.710	Neutrophils

Figure 2: Flow cytometry analysis reveals dynamics changes in uterine immune cell composition during a mouse model of endometrial repair and remodelling. (A) Flow cytometry gating strategy employed to identify and quantify distinct immune cell populations in uterine tissue digests. **(B)** Stacked bar plot: relative abundance of uterine immune cell populations across phases of endometrial repair and remodelling (0-,12-, 24-, 48 hr following progesterone withdrawal) compared to cycling/steady state control tissue: colours represent distinct immune cell types, and the y-axis denotes the percentage of total CD45+ immune cells in each dataset. **(C)** Table: Quantification of flow cytometry analysis of the relative abundance

746   of uterine immune cell populations across phases of endometrial repair and remodelling: 0 hr

747   (n=17), 12 hr (n=22), 24 hr (n=16) and 48 hr (n=10) following progesterone withdrawal

748   compared to cycling/steady state control tissue (n=15).

Figure 3: Flow cytometry and immunohistochemistry reveal dynamic changes in uterine lymphocyte populations during endometrial repair using a mouse model of

simulated menstruation. (A) Population restricted FC analysis to identify CD45$^+$ Ly6G$^-$

NK1.1$^-$ CD19$^-$ CD3$^+$ T cells in uterine tissue digests. **(B)** Bar plot, FC quantification: Relative

abundance of T cells calculated as a percentage of total CD45$^+$ cells per uterine horn. **(C)** Bar

plot, FC quantification: Absolute counts of T cells per uterine horn. **(D)** Immunohistochemistry:

detection of CD3$^+$ T cells (black arrows) in uterine tissues 24- and 48 hr after progesterone

withdrawal (representative images: 24 hr n=5, 48 hr n=6, scale bar= 50μm). **(E)** Population

restricted FC analysis to identify CD45$^+$ Ly6G$^-$ CD3$^-$ CD19$^-$ NK1.1$^+$ NK cells in uterine tissue

digests. **(F)** Bar plot, FC quantification: Relative abundance of NK cells calculated as a

percentage of total CD45$^+$ cells per uterine horn. **(G)** Bar plot, FC quantification: Absolute

counts of NK cells per uterine horn. **(H)** Immunohistochemistry: detection of DBA-bound NK

cells in uterine tissues 24- and 48 hr after progesterone withdrawal (representative images:

24 hr n=4, 48 hr n=4, scale bar= 50μm). **(I)** Population restricted FC analysis to identify CD45$^+$

Ly6G$^-$ CD3$^-$ NK1.1$^-$ CD19$^+$ B cells in uterine tissue digests **(J)** Bar plot, FC quantification:

Relative abundance of B cells calculated as a percentage of total CD45$^+$ cells per uterine horn.

(K) Bar plot, FC quantification: Absolute counts of B cells per uterine horn. **(L)**

Immunohistochemistry: detection of B220$^+$ B cells (black arrows) in uterine tissues 24- and 48

hr after progesterone withdrawal (representative images: 24 hr n=4, 48 hr n=4, scale bar=

50μm). (FC analysis groups: control (n=15), 0 hr (prior to breakdown, n=17), 12 hr (tissue

breakdown, n=22), 24 hr (tissue repair, n=16) and 48 hr (tissue remodelling, n=10) after

progesterone withdrawal; statistical comparisons were made using Kruskal-Wallis tests with

multiple comparisons. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$).

Figure 4: Flow cytometry and immunohistochemistry reveal dynamic changes in uterine myeloid cell populations during endometrial repair using a mouse model of

simulated menstruation. **(A)** Population restricted FC analysis to identify CD45$^+$ CD3$^-$ CD19$^-$ NK1.1$^-$ CD11b$^+$ Ly6G$^-$ SiglecF$^-$ CD64$^-$ Ly6C$^+$ MHCII$^-$ monocytes in uterine tissue digests. **(B)** Bar plot, FC quantification: Relative abundance of monocytes calculated as a percentage of total CD45$^+$ cells per uterine horn. **(C)** Bar plot, FC quantification: Absolute counts of monocytes per uterine horn. **(D)** Population restricted FC analysis of CD45$^+$ CD3$^-$ CD19$^-$ NK1.1$^-$ CD11b$^+$ Ly6G$^-$ SiglecF$^-$ CD64$^+$ Ly6C$^-$ F4/80$^+$ MHCII$^+$ macrophages in uterine tissue digests. **(E)** Bar plot, FC quantification: Relative abundance of macrophages calculated as a percentage of total CD45$^+$ cells per uterine horn. **(F)** Bar plot, FC quantification: Absolute counts of macrophages per uterine horn . **(G)** Population restricted FC analysis of CD45$^+$ CD3$^-$ CD19$^-$ NK1.1$^-$ CD11b$^+$ Ly6G$^+$ SiglecF$^-$ neutrophils in uterine tissue digests. **(H)** Bar plot, FC quantification: Relative abundance of neutrophils calculated as a percentage of total CD45$^+$ cells per uterine horn. **(I)** Bar plot, FC quantification: Absolute counts of neutrophils per uterine horn. (FC analysis groups: control (n=15), 0 hr (prior to breakdown, n=17), 12 hr (tissue breakdown, n=22), 24 hr (tissue repair, n=16) and 48 hr (tissue remodelling, n=10) after progesterone withdrawal; statistical comparisons were made using Kruskal-Wallis tests with multiple comparisons. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$). **(J-K)** Immunohistochemical detection of CD14$^+$ monocytes (cyan), CD64$^+$ macrophages (yellow) and Ly6G$^+$ neutrophils (magenta) (merged and split channel) in uterine tissue cross sections at **(J)** 24 hr (repair, n=5) and **(K)** 48 hr (remodelling, n=3) following progesterone withdrawal (merged and split channel representative images, scale bar = 1000µm).

Figure 5 – CellChat analysis of scRNAseq immune cell data infers intercellular communication networks associated with endometrial repair and remodelling. (A) Scatterplot: visualisation of overall signalling role of each cell group as determined by total outgoing and incoming communication probability (dot size = total number of inferred links, dot colour = different cell groups). **(B)** Heatmap: Outgoing and incoming signalling patterns relative to cell type identifying top pathways in scRANseq data. **(C)** Circle plot: inter- and

intracellular communication networks between immune cell types for **(i)** CCL, **(ii)** CXCL, **(iii)** MIF, **(iv)** FN1, **(v)** SPP1, **(vii)** THBS signalling pathways (circle size = number of cells in group, connection width = probability of interaction). **(D)** Chord diagram: ligand-receptor pairs and their weighted interactions between immune cells for **(i)** CCL, **(ii)** CXCL, **(iii)** MIF, **(iv)** FN1, **(v)** SPP1, **(vii)** THBS signalling pathways (chord width = strength of interactions). **(E)** Heatmap: Overall signalling patterns of immune cells relative to experimental condition, top 8 signalling pathways highlighted with arrows (intensity of colour= relative signalling strength). **(F)** Scatterplot: pairwise comparison of relative strength of signalling pathways in monocytes, macrophages and neutrophils between experimental data set: cycling vs repair; cycling vs remodelling; repair vs remodelling.

<u>Supplementary data</u>

Supplementary figure 1: Mouse model of simulated menstruation characterised by

endometrial tissue breakdown and repair. (A) Schematic detailing the timeline of surgical

procedures and hormonal stimulation used to induce a menstruation-like event in mice. **(B)**

Haematoxylin and eosin (H&E) staining of uterine tissue sections demonstrate macro- and

microscopic tissue architecture in **(i)** steady state mouse uterus and mouse uterus at various

time points of tissue breakdown and repair: **(ii)** 0hrs, prior to breakdown; **(iii)** 12hrs, tissue

breakdown; **(iv)** 24hrs, endometrial repair; **(v)** 48hrs, endometrial remodelling (time post

progesterone withdrawal; scale bar=1000µm).

Fluorophore	Antigen	Clone	Catalogue number	Dilution
BV421	CD64	X54-7/7.1	139309	1/100
BV510	CD45	30-F11	103138	1/200
BV570	CD3	17A2	100225	1/40
BV785	CD11c	N418	117335	1/100
PE-Fire 640	Ly6G	1A8	127675	1/400
PE/Dazzle 594	Siglec F	E50-2440	562757	1/400
PE/Cy7	F4/80	BM8	123114	1/200
PerCP/Cy5.5	Ly6C	HK1.4	128012	1/100
AF700	MHC II	M5/114.15.2	107622	1/200
APC/Cy7	CD11b	M1/70	101226	1/200
Pacific blue	NK1.1	PK136	108722	1/200
Spark blue 550	CD19	6D5	115565	1/200
LIVE/DEAD Red			L34971	1 test per sample

828 **Supplementary Table 1: Flow cytometry antibody panel utilised to characterise uterine**

829 **immune cell dynamics.**

A)

B)

Supplementary figure 2: Gene Ontology (GO) analysis reveals biological processes

(BP) associated with the transcriptome of each immune cell cluster. (A) Bar plot:

Unbiased analysis of GOBP terms associated with the genetic signature of each immune cell

cluster: **(i)** T cells **(ii)** B cells **(iii)** NK cells **(iv)** Dendritic cells **(v)** Monocytes **(vi)** Macrophages

(vii) Neutrophils **(viii)** Eosinophils. Bar size= qscore: -log(p.adjust) ; bar colour= P.adjust

value. **(B)** Heatmap (yellow, high; purple, low): Biased analysis of gene signatures (GSEA

gene sets) associated with known wound tissue repair GO terms by neutrophil, monocyte and

macrophage clusters: apoptosis, NETosis, wound healing, phagocytosis, chemoattraction,

remodelling enzymes. Top axis is colour coded and named by immune cell type.

Supplementary figure 3: Average gene expression analysis for each immune cell

cluster between the 3 experimental conditions. Scatter plot: Average gene expression

analysis for each immune cell cluster **(A)** T cells **(B)** B cells **(C)** NK cells **(D)** Dendritic cells **(E)**

Monocytes **(F)** Macrophages **(G)** Neutrophils **(H)** Eosinophils between different experimental

conditions: **(i)** Cycling vs 24 hr **(ii)** Cycling vs 48 hr **(iii)** 24 hr vs 48 hr.

Supplementary figure 4: Immunohistochemistry analysis of uterine lymphocyte dynamics during endometrial repair using a mouse model of simulated menstruation.
Immunohistochemistry: detection of **(A)** CD3+ T cells, **(B)** DBA-lectin+ NK cells and **(C)** B220+ B cells in uterine tissue sections at 0hr (prior to breakdown, n=4), 12hr (tissue breakdown, n=4-6), 24hr (tissue repair, n=4-5) and 48hr (tissue remodelling, n=4-6) hr after progesterone withdrawal (representative images, scale bar=500µm).

Supplementary figure 5: Flow cytometry analysis of dendritic cells and eosinophils in uterine tissues endometrial repair using a mouse model of simulated menstruation. (A) Population restricted FC analysis to identify CD45+ CD3- CD19- NK1.1- Ly6G- CD64- CD11c+ MHCII+ dendritic cells (DCs) in uterine tissue digests. **(B)** Bar plot, FC quantification: Relative abundance of DCs calculated as a percentage of total CD45+ cells per uterine horn. **(C)** Bar plot, FC quantification: Absolute counts of DCs per uterine horn. **(D)** Population restricted FC analysis to CD45+ CD3- CD19- NK1.1- CD11b+ Ly6G- SiglecF+ eosinophils in uterine tissue digests. **(E)** Bar plot, FC quantification: Relative abundance of eosinophils calculated as a percentage of total CD45+ cells per uterine horn. **(F)** Bar plot, FC quantification: Absolute counts of eosinophils per uterine horn. (FC analysis groups: control (n=15), 0hr (prior to breakdown, n=17), 12hr (tissue breakdown, n=22), 24hr (tissue repair, n=16) and 48hr (tissue remodelling, n=10) after progesterone withdrawal; statistical comparisons were made using Kruskal-Wallis tests with multiple comparisons. and * p< 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001).

Supplementary figure 6: Immunohistochemistry analysis of myeloid cell dynamics during endometrial repair using a mouse model of simulated menstruation. Immunohistochemical detection of **(A)** CD14+ monocytes (cyan), **(B)** CD64+ macrophages (yellow) and **(C)** Ly6G+ neutrophils (magenta) in uterine tissue sections at 0hr (prior to breakdown, n=4), 12hr (tissue breakdown, n=5), 24hr (tissue repair, n=5) and 48hr (tissue

879   remodelling, n=3) following progesterone withdrawal (representative images, scale

880   bar=1000µm). (D) Merged channel representative images (scale bar=1000µm).

882 **<u>Appendix</u>**

883 <u>Abbreviations</u>

Abbreviation	**Full term**
Acod1	Aconitate decarboxylase 1
Adgre1	Adhesion G protein–coupled receptor E1 (F4/80)
Anxa5	Annexin A5
Apoe	Apolipoprotein E
APP	Amyloid precursor protein
AWERB	Animal Welfare and Ethical Review Body
Axl	AXL receptor tyrosine kinase
Cav1	Caveolin 1
C3	Complement component 3
Ccl2	C–C motif chemokine ligand 2
Ccl3	C–C motif chemokine ligand 3
Ccl4	C–C motif chemokine ligand 4
Ccl5	C–C motif chemokine ligand 5
Ccl6	C–C motif chemokine ligand 6
Ccl9	C–C motif chemokine ligand 9
CCL	C–C motif chemokine ligand
CCR2	C–C motif chemokine receptor 2
Cd3e	CD3 epsilon chain
CD3	Cluster of differentiation 3
CD9	Cluster of differentiation 9
CD11b	Integrin alpha M
CD11c	Integrin alpha X
CD14	Cluster of differentiation 14
CD19	Cluster of differentiation 19
CD45	Protein tyrosine phosphatase receptor type C
CD64	Fc gamma receptor I
Cd68	Cluster of differentiation 68
Cdc1	Conventional dendritic cell 1
cDNA	Complementary DNA
Clec9a	C-type lectin domain family 9 member A
Clec10	C-type lectin domain family 10
Csf1r	Colony stimulating factor 1 receptor
CXCL	C-X-C motif chemokine ligand
Cxcl1	C-X-C motif chemokine ligand 1
Cxcl3	C-X-C motif chemokine ligand 3
Cxcr2	C-X-C motif chemokine receptor 2
CXCL8	C-X-C motif chemokine ligand 8
DCs	Dendritic cells
DEGs	Differentially expressed genes
Dusp1	Dual specificity phosphatase 1
E2	Estradiol
EDTA	Ethylenediaminetetraacetic acid

Abbreviation	Full term
F10	Coagulation factor X
F4/80	Macrophage marker (Adgre1)
FACS	Fluorescence-activated cell sorting
FC	Flow cytometry
FFPE	Formalin-fixed paraffin-embedded
FN1	Fibronectin 1
Fn1	Fibronectin 1
FSC-A	Forward scatter area
FSC-H	Forward scatter height
GEMs	Gel bead-in-Emulsions
GO	Gene ontology
GOBP	Gene ontology biological processes
Gata3	GATA binding protein 3
Gnas	GNAS complex locus
Gzmb	Granzyme B
H2-Ab1	MHC class II antigen A, beta 1
Hic1	Hypermethylated in cancer 1
HMB	Human menopausal blood
HRP	Horseradish peroxidase
Ighd	Immunoglobulin heavy constant delta
IL-1β	Interleukin 1 beta
IL-6	Interleukin 6
Inha	Inhibin alpha
ISG15	Interferon-stimulated gene 15
IFIT1	Interferon-induced protein with tetratricopeptide repeats 1
IFIT3	Interferon-induced protein with tetratricopeptide repeats 3
Klrb1c	Killer cell lectin-like receptor B1C
Lrg1	Leucine-rich alpha-2-glycoprotein 1
Ly6C	Lymphocyte antigen 6 complex, locus C
Ly6G	Lymphocyte antigen 6 complex, locus G
MAPK	Mitogen-activated protein kinase
Mertk	MER proto-oncogene tyrosine kinase
MHC II	Major histocompatibility complex class II
MIF	Macrophage migration inhibitory factor
MMP	Matrix metalloproteinase
Msr1	Macrophage scavenger receptor 1
NET	Neutrophil extracellular trap
NK1.1	Natural killer cell marker
Ncr1	Natural cytotoxicity triggering receptor 1
P4	Progesterone
PCA	Principal component analysis
Pdpn	Podoplanin
PBS	Phosphate-buffered saline
Plaur	Plasminogen activator, urokinase receptor
QC	Quality control

Abbreviation Full term

Abbreviation	Full term
RNA	Ribonucleic acid
Rsad2	Radical S-adenosyl methionine domain containing 2
S100a8	S100 calcium-binding protein A8
S100a9	S100 calcium-binding protein A9
ScRNAseq	Single-cell RNA sequencing
Sdc1	Syndecan 1
Sdc4	Syndecan 4
Sell	L-selectin
SiglecF	Sialic acid-binding Ig-like lectin F
Slc11a1	Solute carrier family 11 member 1
SPP1	Secreted phosphoprotein 1
SSC-A	Side scatter area
SSC-H	Side scatter height
TGF-β	Transforming growth factor beta
TGFBI	Transforming growth factor beta-induced
THBS1	Thrombospondin 1
Thbd	Thrombomodulin
Tmem123	Transmembrane protein 123
TNF-α	Tumour necrosis factor alpha
TPM	Transcripts per million
TSP-1	Thrombospondin-1
UMAP	Uniform Manifold Approximation and Projection
uNKs	Uterine natural killer cells
VEGF	Vascular endothelial growth factor
Xcr1	X-C motif chemokine receptor 1