# 2-Deoxyglucose dendrimer-enabled niclosamide delivery to FRβ-expressing macrophages alleviates endometriosis progression and associated hyperalgesia

Anubhav Dhull[1#], Mingxin Shi[2#], Madeleine E. Harvey[2#], Taylor M. Page[2], Rishi Sharma[1], Kenneth John Goody[1], Aqib Iqbal Dar[1], Nash S. Moawad[3], Vijay Kumar Sirohi[4], Paul S. Cooke[4], Kanako Hayashi[2]*, Anjali Sharma[1]*

[1]*Department of Chemistry, College of Arts and Sciences, Washington State University, Pullman, WA 99164, USA.*

[2]*School of Molecular Biosciences, Center for Reproductive Biology, Washington State University, Pullman, WA 99164, USA.*

[3]*Department of Obstetrics & Gynecology, University of Florida College of Medicine, P.O. Box 100294, Gainesville, FL 32610, USA.*

[4]*Department of Physiological Sciences, University of Florida, P.O. Box 100144, Gainesville, FL 32611, USA.*

[#]*These authors contributed equally to this work*

*Correspondence:

Anjali Sharma, Department of Chemistry, College of Arts and Sciences, Washington State University, Pullman, WA 99164, USA.

Kanako Hayashi, School of Molecular Biosciences, Center for Reproductive Biology, Washington State University, Pullman, WA 99164, USA.

E-mail: anjali.sharma@wsu.edu, k.hayashi@wsu.edu

*Abstract*

Endometriosis is a chronic, incurable disease. Due to limited efficacy, high recurrence rates, and

serious side effects of current treatments, development of new, targeted, non-hormonal therapies

is urgently needed. We previously reported that niclosamide, an FDA-approved anthelmintic

drug, attenuates endometriotic lesion growth. We further identified folate receptor-$\beta$ (FR$\beta$)-

positive macrophages as contributors to disease progression. Significantly, niclosamide inhibits

FR$\beta^+$ macrophages and reduces inflammation, innervation, and angiogenesis. To develop

niclosamide as a non-hormonal and selective immune cell-targeted therapy for endometriosis, we

engineered a folic acid-conjugated 2-deoxyglucose dendrimer (FA-2DG-D) using click chemistry

to enable selective FR$\beta$-mediated uptake. Conjugation of niclosamide to FA-2DG-D yielded a

targeted nanotherapeutic (FA-2DG-D-Niclo) with enhanced aqueous solubility, controlled

intracellular release, and excellent batch-to-batch reproducibility. In a mouse model of

endometriosis, FA-2DG-D demonstrated lesion-specific accumulation and selective

internalization by FR$\beta^+$ macrophages with minimal off-target organ retention. A single

intraperitoneal dose of FA-2DG-D-Niclo (25 or 50 mg/kg/bw of niclosamide) significantly

reduced FR$\beta^+$ macrophage burden, suppressed lesion number and volume, and markedly

improved endometriosis-associated hyperalgesia at two weeks post-treatment. Together, these

findings establish FR$\beta^+$ macrophages as a potential target in endometriosis and present FA-2DG-

D-Niclo as a non-hormonal, macrophage-focused nanomedicine for precise and effective

endometriosis treatment.

**Keywords**: endometriosis, niclosamide, dendrimer, macrophages, non-hormonal therapy

**Graphic Abstract**

*Created via Biorender.com*

## 1. Introduction

Endometriosis affects an estimated 10% of reproductive-aged women, roughly 190 million individuals worldwide, and remains one of the most debilitating gynecological disorders [1]. The disease is characterized by chronic pelvic pain, dysmenorrhea, dyspareunia, and infertility, all of which severely diminish the quality of life for patients and their families [2-4]. In the United States alone, the annual economic burden is estimated at $78-119 billion, encompassing healthcare costs, productivity losses, and reduced workforce participation [5]. Despite its prevalence and impact, endometriosis remains incurable, and the therapeutic landscape has changed little over the past several decades. Current medical management relies on hormonal suppression of estrogen production or action [6, 7], such as oral contraceptives, progestins, and/or gonadotropin-releasing hormone (GnRH) agonists/antagonists [6, 8], as well as surgical excision of lesions. However, hormonal treatments are limited in efficacy, have high recurrence rates, and cause significant side effects. Surgical excision offers temporary relief of symptoms, yet nearly half of surgically treated women experience recurrence within five years, and ~12% fail to achieve meaningful pain reduction after surgery [9, 10]. Thus, there is an urgent need to identify new therapeutic targets to develop novel non-hormonal treatments in endometriosis.

A growing body of evidence implicates immune dysfunction, particularly macrophage dysregulation, in driving lesion establishment, chronic inflammation, angiogenesis, and neurogenesis in endometriosis [11-13]. Recently, our single-cell transcriptomic analysis identified a unique FRβ-expressing macrophage population, differentiated from monocyte-derived proinflammatory macrophages and characterized by residential macrophage traits in mouse endometriosis [14]. The population of macrophages is also identified in the lesions of endometriosis patients [15]. The selective upregulation of FRβ$^+$ on these macrophages can be used as an accessible and disease-restricted cellular target for ligand-guided targeted drug delivery.

Our prior studies demonstrated that niclosamide significantly suppressed endometriotic lesion growth, alleviated aberrant peritoneal inflammation, and decreased vascularization and innervation in lesions using two distinct mouse models of endometriosis without compromising fertility [13, 14, 16]. Niclosamide (5-chloro-*N*-(2-chloro-4-nitrophenyl)-2-hydroxybenzamide) is a chlorinated salicylanilide derivative that is FDA-approved as an anthelmintic drug. Currently, multiple clinical trials are ongoing to repurpose it for the treatment of other diseases, including cancer and metabolic diseases [17]. However, the clinical use of niclosamide is significantly

limited by its poor aqueous solubility and correspondingly low oral bioavailability. To address these challenges, researchers are actively developing innovative formulation strategies such as water-soluble derivatives, solid dispersions, and nanocarrier-based delivery systems to enhance its solubility, pharmacokinetics and overall therapeutic potential [18]. Additionally, systemic niclosamide treatment inhibits several pathways, including STAT3, NFκB, and WNT signaling, across all cell types [11, 12, 16], including other macrophages [14]. Disrupting these essential pathways across all cell types would lead to unwanted side effects and this poses a significant barrier to therapeutic use of niclosamide. Importantly, niclosamide can also reduce elevated FRβ$^+$ macrophages and restore the perturbed microenvironment in endometriosis by finely regulating macrophage dynamics [14], suggesting that FRβ$^+$ macrophages can serve as a viable therapeutic target in endometriosis. Therefore, targeted delivery of niclosamide to FRβ$^+$ macrophages offers a strategy to maximize therapeutic efficacy while minimizing systemic toxicity.

In this context, dendrimer nanotechnology has emerged as a powerful tool for targeted therapeutic delivery, enabling selective engagement of pathogenic cells while minimizing off-target toxicity [19-24]. Dendrimers are monodisperse, highly branched macromolecules whose multivalent surface architecture and tunable chemistry allow precise conjugation of targeting ligands, therapeutic agents, and imaging probes [25-27]. Building on our previously synthesized 2-deoxy-glucose (2DG) functionalized dendrimer (2DG-D) [28], which incorporates 2DG moieties to enhance solubility and macrophage interaction, we engineered a folic acid (FA), a ligand of FRβ, conjugated dendrimer (FA-2DG-D) by covalently coupling FA to the dendrimer surface through orthogonal click chemistry. This design exploits ligand-FRβ binding to selectively direct dendrimer uptake into FRβ$^+$ macrophages within endometriotic lesions and peritoneum [29, 30]. To achieve targeted delivery, we further developed the niclosamide conjugated FA-2DG-D (FA-2DG-D-Niclo). Here, we report the design, synthesis, and in vivo evaluation of FA-2DG-D and FA-2DG-D-Niclo in a mouse model of endometriosis. Our results demonstrate that FA-2DG-D has FRβ$^+$ macrophage-specific targeting in the lesions and peritoneum and delivers niclosamide into FRβ$^+$ macrophages with ligand-receptor internalization. This targeted dendrimer platform offers a promising strategy for therapeutic delivery in endometriosis, addressing a critical unmet need for more precise and effective interventions.

**2. Methods**

### 2.1. Human Sample Collection

All procedures involving human subjects were performed in accordance with ethical approval by the Institutional Review Board (IRB) guidelines for the University of Florida under IRB code NO. IRB202003131. All patients were recruited from the Division of Minimally Invasive Gynecologic Surgery of the University of Florida Health Shands Hospital, Gainesville, Florida. Participants were either previously diagnosed with endometriosis or suspected of having endometriosis, with symptoms including ovarian cysts, chronic pelvic pain, dysmenorrhea, dyschezia, dyspareunia, uterine bleeding, and myofascial pain syndrome. Endometriosis is further confirmed by pathological analysis of the excised lesions obtained during surgery. Informed written consent was obtained from each participant before the procedure. In this study, peritoneal washings from 4 patients aged 19 to 37 years (**Supplementary Table S1**) were collected at the beginning of the procedure. Briefly, the pelvic cavity was irrigated using approximately 100 mL of sterile saline, and approximately 80 mL of peritoneal washings containing pelvic exudate cells were collected from each patient and placed on ice immediately. Red blood cells were removed by resuspending cells with 500 µL Red Blood Cell Lysis Buffer (BioLegend, Cat.#420301). Cells were then counted and cryopreserved with freezing media containing 90% fetal bovine serum (FBS) and 10% dimethyl sulfoxide, and stored at −80 °C for further analysis.

### 2.2. Single-cell RNA-Sequencing and Analysis

### 2.2.1. Library Preparation and Sequencing

To perform single-cell RNA sequencing (scRNA-seq), frozen pelvic exudate cells were thawed in a 37 °C water bath and prepared following the 10x Genomics protocol named "Fresh frozen human peripheral blood mononuclear cells for single-cell RNA sequencing" (document #CG00039 Rev E). Briefly, thawed cells were slowly transferred into a 50 mL conical tube and recovered by sequential 1:1 dilution with warmed FBS for a total of 5 times. Cell viability was assessed to ensure samples with viability greater than 80%. After centrifuging at 300g for 5 min at room temperature, the cells were resuspended in phosphate-buffered saline (PBS) with 0.04% bovine serum albumin (BSA) at a concentration of 900-1,200 cells/µl. Cells were then placed on ice and immediately proceeded to library construction using a 10x Genomics Chromium Next GEM Single Cell 3' Kit v3.1, following the standard kit user guide (document #CG000204) for a target of 6000 cells. Libraries were sequenced on Illumina NovaSeq 6000. FASTQ files were

aligned to the human reference genome GRCh38 via Cell Ranger v7.1.0 [31], and an average of

107,903 reads per cell was achieved for each RNA library (**Supplementary Table S2**).

*2.2.2. External Human Endometrial and Pelvic Exudate Cell scRNA-seq Datasets*

We integrated publicly available human scRNA-seq datasets from eutopic endometrium

with or without endometriosis, ectopic endometrium, and pelvic exudate cells. Processed Cell

Ranger output files or raw FASTQ files were downloaded from the Gene Expression Omnibus

(GEO), ArrayExpress, or the Sequence Read Archive (SRA). These datasets included: (1) Wang et

al. (GSE111976) [32], (2) Lai et al. (GSE183837) [33], (3) Garcia-Alonso et al. (E-MTAB-10287)

[34], (4) Tan et al. (GSE179640) [15], (5) Fonseca et al. (GSE213216) [35], (6) Huang et al.

(GSE214411) [36], (7) Marečková et al. (E-MTAB-14039) [37], (8) Shin et al. (PRJNA932195)

[38], (9) Han et al. (GSE228030) [39], and (10) Zou et al. (PRJNA713993) [40], as shown in

**Figure 1A and B**. For all datasets, only samples collected from women of reproductive age were

included. For endometrial scRNA-seq datasets from (1) to (8), patients labeled as undergoing

hormonal treatments during sample collection were also excluded. Samples used in this study are

summarized in **Supplementary Table S3**.

*2.2.3. Single-cell Data Pre-processing and Clustering*

RNA matrices from both newly generated and existing public datasets were individually

imported to Scanpy v.1.8.1 for further analysis [41]. Low-quality cells were excluded using the

following criteria: (1) genes that were detected in fewer than 3 cells, (2) cells with fewer than 300

genes, (3) a maximum of 100,000 UMIs, and (4) a maximum of 20% mitochondrial content.

Doublets were detected and removed using Scrublet [42]. Filtered matrices from endometrial or

pelvic cell scRNA-seq datasets were combined separately and analyzed independently. For each

combined endometrial or pelvic cell scRNA-seq matrix, gene expression values were normalized

with a global scaling factor of 10, 000 and log-transformed. Top 3,000, or 2,000 highly variable

genes for the endometrial, or pelvic cell datasets, respectively, were identified by Scanpy. The

mitochondrial genes, haemoglobin genes, and ribosomal genes were excluded from the highly

variable gene set [43]. Principal component analysis (PCA) was performed using the highly

variable gene set. After Harmony (v.0.0.10) [44] batch effect correction, harmony-corrected

components were used to generate the neighborhood graph and perform dimensionality reduction

using uniform manifold approximation and projection (UMAP). The Leiden algorithm was then used for clustering analysis [45]. Marker genes of each cluster were identified using the Wilcoxon rank-sum test [46] with the default setting (**Supplementary Table S4**). Major cell types were determined by manually annotating cell clusters with previously established marker genes from the literature [15, 39, 47, 48].

### 2.2.4. Identification of Enriched Genes and Gene Ontology Analysis in FOLR2$^+$ Endometrial and Pelvic Cells

To understand the transcriptional profiles of folate receptor beta-positive (FOLR2$^+$) macrophages, cells with specific FOLR2 expression (FOLR2>0) were labeled and extracted from endometrial and pelvic cell matrices. Enriched genes in FOLR2$^+$ cells were identified by the Wilcoxon rank-sum test in a one-versus-rest fashion, with a minimum log$_2$(fold change) of 2 (**Supplementary Table S5**). Gene Ontology (GO) was performed on the identified enriched genes in FOLR2$^+$ endometrial or pelvic cells using clusterProfiler (v.4.2.2) [49]. The enriched GO list was further filtered with the *simplify* function (cutoff = 0.7) to remove redundant terms (**Supplementary Table S6**). Differentially expressed genes (DEGs) among eutopic endometrium without (Control; Cont) or with endometriosis (EuE), and ectopic endometrium (EcE) in the endometrial FOLR2$^+$ macrophages, and DEGs between control (Cont) and endometriotic (Endo) pelvic exudate cells, were identified by Wilcoxon rank-sum test, with a minimum log$_2$(fold change) of 0.5 (**Supplementary Table S7**).

### 2.3. Chemistry Experimental Section

Experimental methods related to materials, instrumentation, synthesis protocols, characterization of compounds (**Supplementary Figures S1-S15 and S17-S34**), in vitro drug release studies, and FA-2DG-D-Niclo formulation stability studies are presented in the **Supplementary Information**.

### 2.4. Animals

C57BL/6 mice were purchased from Inotiv and housed in an environment-controlled animal facility (20-22 °C, ~40% relative humidity, and 12:12 light-dark cycle) with *ad libitum*

206 access to food and water. All animal experiments were conducted at Washington State University
207 in accordance with NIH guidelines for care and use of laboratory animals (protocol #6751).
208

*2.5. Mouse Model of Endometriosis*

An experimental mouse model of endometriosis was employed, using procedures
previously described and adopted by us, with minor modifications [50]. Briefly, to induce
endometriosis-like lesions (ELL), female donor mice were injected subcutaneously with pregnant
mare serum gonadotropin (PMSG; 5 IU; Sigma, St. Louis, MO) to stimulate an estrogenic response
within the uterus. Uteri were harvested from these donors 41 hours after PMSG injection. The
endometrium was then scraped from myometrium and dissected into fragments (1-2 mm per side),
and introduced as the source of syngeneic mouse endometrium (50 mg of fragments) via injection
(in 0.2 mL of PBS) into the peritoneal cavity in the ovary-intact recipient females under anesthesia
via inhaled isoflurane.
219

*2.6. Animal Study Design*
*2.6.1. In Vivo Dendrimer Biodistribution*

Adult female mice (n=36, ~3 months old) were randomly divided into three groups
(n=3/group/timepoint): 1) sham controls administered with Cy5-labeled dendrimer FA-2DG-D-
Cy5; 2) ELL-induced mice administered with 2DG-D-Cy5; 3) ELL-induced mice administered
with FA-2DG-D-Cy5. The animals in the ELL groups received donor endometrium to induce
endometriotic lesions, and the sham control mice received PBS. One week following lesion
induction, all mice received FA-2DG-D-Cy5 or 2DG-D-Cy5 injected into the tail vein, as listed in
the above group description, at a dose of 50 mg/kg (300 μL). Mice were then euthanized at 1, 6,
24, and 72 hours after injection. The whole skin was gently peeled off to acquire the whole-body
Cy5 fluorescence distribution using an in vivo imaging system (IVIS; Revvity). Lesions and/or
major organs, including the brain, heart, spleen, lungs, liver and kidneys, were dissected to obtain
ex vivo fluorescence images and stored at -80 °C. The fluorescence intensity of the region of
interest (ROI) was calculated with the Living Image software (4.7.4) accompanying IVIS.
Amounts of dendrimer in major organs and lesions were further examined ex vivo using
fluorescence intensity-based quantification.
236

### 2.6.2. Ex Vivo Dendrimer Quantification

Frozen organs (lesions, liver, spleen, heart, kidneys, brain and lungs) were slowly thawed on ice and weighed. Tissue samples were then dissected to obtain known masses from each organ. Samples were homogenized in methanol (1 mL per 100 mg tissue) with stainless steel beads using a tissue homogenizer. The homogenized samples were centrifuged at 4°C, and the clear supernatants were transferred to protein LoBind Eppendorf tubes and stored at -80°C. For fluorescence quantification, thawed supernatants were centrifuged again before analysis. Fluorescence intensity was measured on a Varioskan Lux Multimode Microplate Reader (ThermoFisher), using $\lambda\_ex = 650$ nm and $\lambda\_em = 670$ nm for Cy5. Background fluorescence from control solvent was subtracted, and corrected fluorescence values were converted to 2DG-D-Cy5 and FA-2DG-D-Cy5 concentrations using calibration curves generated at 5-10 slit widths.

### 2.6.3. In Vivo Cellular Uptake of Dendrimer

To evaluate the specificity of cellular uptake of the dendrimer, endometriotic lesions were induced in 6 female mice. On Day 7 following lesion induction, mice received an intraperitoneal (i.p.) injection of FA-2DG-D-Cy5 or 2DG-D-Cy5 (50 mg/kg/bw, n=3/group). Peritoneal cells, lesions, kidney, liver and spleen were collected at 2 weeks after dendrimer injection. Specific cellular uptake of the dendrimer was then assessed by immunostaining.

### 2.6.4. In Vivo Efficacy of Dendrimer-Niclosamide

The efficacy of targeted delivery of niclosamide with FA-conjugated dendrimer (FA-2DG-D-Niclo) was examined using the mouse model of endometriosis described above. On Day 7, ELL-induced mice were administered a single i.p. injection of control (FA-2DG-D), or 25 mg or 50 mg/kg body weight (bw) of FA-2DG-D-Niclo. Behavioral tests were performed on Day -1 (a day before lesion induction), Day 7 (before dosing), and Day 14 or 21 (1 or 2 weeks after dosing, n=16 at Day -1 and Day 7, n=8 at Day 14 and Day 21). Mice were then euthanized on Day 14 or 21 after the last behavioral test. Peritoneal fluid was recovered by lavage (4 mL x 2 of ice-cold PBS with 3% FBS) for immune profiling. The number and volume of lesions were recorded. Lesions were fixed in 4% paraformaldehyde for further analysis.

### 2.7. von Frey Test

A standard behavioral (mechanical sensitivity) test was performed before sample

collection, as described by our laboratory previously [51, 52]. Mice (n=8/group/time point) were

allowed to acclimate quietly in the testing room for 30 minutes under red light. Then, von Frey

filaments (BIO-VF-M, Bioseb) were applied 10 times to the skin perpendicular to the lower

abdomen and bilateral hind paws to evaluate local or systemic sensitivity, respectively. The force

in grams (g) of the filament evoking a withdrawal response (50% response count as sensitive) was

recorded. Three behaviors were considered positive responses to filament stimulation: 1) sharp

retraction of the abdomen, 2) immediate licking and/or scratching of the area of filament

stimulation, or 3) jumping. All behavioral tests were performed with investigators blinded to

treatment groups. These data were then subsequently analyzed by another blinded investigator.

***2.8. Flow Cytometry***

Single-cell suspensions of peritoneal exudate cells were used for immune cell profiling by

flow cytometry as described previously [51, 52]. Briefly, peritoneal exudate cells were lysed using

Red Blood Cell Lysis Buffer (BioLegend) and incubated at room temperature for 20 minutes with

Zombie Aqua™ Fixable Viability dye (BioLegend). Cells were blocked on ice for 20 minutes with

Fc Block anti-CD16/CD32 (ThermoFisher) and stained with fluorochrome-conjugated

monoclonal antibodies for 1 hour (**Supplementary Table S8**). Samples (n=8/group) were acquired

with the Attune NxT Acoustic Focusing Cytometer using Attune NxT software (ThermoFisher),

and data were analyzed with FlowJo v10.4 software (FLOWJO).

***2.9. Immunohistochemistry and Confocal Imaging***

The peritoneal exudate cell suspension was first fixed with 4% paraformaldehyde at RT for

15 min, and blocked with 10% donkey serum at RT for 30 min. Cells were then incubated with

FRβ primary antibody (1:200, Abcam, Cat.#ab302532) at 4°C overnight and AlexaFluor 488-

conjugated F(ab') secondary antibody (Molecular Probe) at RT for 1 hour. Following DAPI

staining, the cells were resuspended with Fluoromount-G (SouthernBiotech) and mounted on

slides. The mounted slides were imaged using a Leica SP8-X Confocal Microscope.

Immunohistochemistry of FRβ on cross-sections (5 μm) of paraffin-embedded tissues,

including lesion, kidney, and liver, was performed using the FRβ primary antibody (1:400) and

AlexaFluor 488-conjugated F(ab') secondary antibody, or VECTASTAIN ABC kit (Vector Lab).

Immunostaining images were acquired by Leica SP8-X Confocal Microscope or Leica DM4 B

microscopy. FRβ-positive cells in lesions were counted by Image J in an area of 0.07244 $mm^2$, and

the number of $FRβ^+$ cells per 1 $mm^2$ was shown.

### 2.10. Statistical Analysis

Statistical analyses were performed using GraphPad Prism (version 9.5). To compare

differences in dendrimer organ uptake among groups, one-way ANOVA followed by the Tukey

multiple comparison tests was used. To analyze behavioral test results, time-dependent differences

within a single group were compared using one-way ANOVA followed by the Dunnett's multiple-

comparison test. To compare effects of different niclosamide dosages of niclosamide at each time

point, one-way ANOVA was used with Tukey multiple comparison tests. Differences in lesion

number, volume, and $FRβ^+$ or $CD45^+$ cell counts between groups at each time point were analyzed

using one-way ANOVA followed by the Tukey multiple-comparison test. In all cases, data were

tested for normal distribution using the Shapiro-Wilk normality test. If data were normally

distributed, one-way ANOVA was used to analyze differences among groups. If data were not

normally distributed, the Kruskal-Wallis test was performed, as stated in the figure caption.

Significance was defined as $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***), or $p < 0.0001$ (****).

## 3. Results and Discussion

### 3.1. FRβ-expressing Macrophages in Eutopic and Ectopic Endometrium, and Pelvic Exudate
### Cells in Endometriosis

We performed an integrated analysis with our newly generated pelvic exudate cell scRNA-

seq dataset from women with endometriosis (**Supplementary Tables S1 and S2**), along with

publicly available human datasets, including eutopic endometrium without or with endometriosis,

ectopic endometrium (lesion), and pelvic exudate cells (**Figure 1A-B, Supplementary Table S3**).

After data merging and quality control filtering, 748,509, and 76,187 total high-quality endometrial

and pelvic exudate cells were retained, respectively. Clustering analysis identified 7 major cell

types in the endometrium (EM), including B lymphocytes ($CD79A^+$), endothelial cells

($PECAM1^+$), epithelial cells ($EPCAM^+$), myeloid cells ($LYZ^+$), perivascular cells ($RGS5^+$),

stromal cells ($DCN^+$), and T lymphocytes and natural killer (NK) cells ($CD3E^+$ and $CD2^+$) (**Figure
1C and E, Supplementary Table S4**). In the pelvic cavity (P), B lymphocytes ($CD79A^+$),

dendritic cells (CD1C[+]), macrophages (Mac, CD14[+]), mast cells (TPSAB1[+]), NK cells (GNLY[+]), and T lymphocytes (CD3D[+]) were assigned as overarching cell types (**Figure 1D and F, Supplementary Table S4**). As expected, UMAP projection confirmed that FOLR2[+] cells are localized within the myeloid cell clusters of endometrial and pelvic exudate cells (**Figure 1G-H**). Gene Ontology (GO) enrichment analysis indicated that both endometrial and pelvic FOLR2[+] macrophages were primarily associated with myeloid migration and activation and immune regulation (**Figure 1I-J, Supplementary Table S6**). In particular, biological processes associated with cytokine production, leukocyte cell-cell adhesion, and interleukin-6 (IL-6) production were predominantly enriched in the FOLR2[+] macrophages of the endometrium. Stronger association with phagocytosis, cholesterol regulation, and response to bacteria was detected in the pelvic FOLR2[+] macrophages, consistent with the primary roles of residential FOLR2[+] macrophages in host defense and metabolic processes [53-55]. Comparative analysis showed a significantly higher abundance of FOLR2[+] macrophages in ectopic endometrium (EcE) than in eutopic endometrium with (EuE) and without (Cont) endometriosis (**Figure 1K**). Similarly, in pelvic exudate cells, significantly elevated numbers of FOLR2[+] macrophages were observed in patients with endometriosis (**Figure 1L**). Furthermore, DEG analysis of endometrial and pelvic FOLR2[+] macrophages demonstrated that significant upregulation of proinflammatory (CCL2, IL-6R, and NFKB1), angiogenesis (FGFR1, HIF1A, and MMP2), and adhesion-related genes (ITGAV, TGFBR1, and TGFBR2) in endometriosis-associated endometrium (EuE and/or EcE) and pelvic cells (Endo) (**Figure 1M-N, Supplementary Table S7**). Collectively, these results indicate an expansion of a pro-disease FOLR2[+] macrophage subpopulation in the ectopic endometrium and pelvic cavity in endometriosis patients with pro-disease functions.

### *3.2. Synthesis and Characterization of FA-2DG-D Using Click Chemistry*

A major challenge in developing nanomaterial-based drug delivery systems lies in the complexity and limited reproducibility of synthetic procedures, scalability constraints, and poor batch-to-batch consistency. To overcome these limitations, we used efficient copper(I) catalyzed alkyne-azide cycloaddition (CuAAC) click reaction to design and synthesize the FA-2DG-D conjugate capable of selectively targeting FRβ[+] macrophages for treating endometriosis. This click chemistry-based approach offers a robust and versatile strategy for achieving precise, reproducible,

and site-specific conjugation of targeting ligands or therapeutic molecules to polymeric

nanocarriers, including dendrimers [28, 56-58].

Folic acid (**1**) was functionalized with an azide group to enable its participation in the

CuAAC reaction. The folate receptor recognizes folic acid through its pterin and *p*-aminobenzoate

moieties via multiple hydrogen bonds and electrostatic interactions within the receptor pocket [59].

To preserve these recognition elements, conjugation with the azide linker was performed through

the α-carboxylic acid group of the glutamic acid residue, forming a stable amide linkage with the

terminal amine of the linker. The carboxylic acid in the ligand was activated using

dicyclohexylcarbodiimide (DCC) and *N*-Hydroxysuccinimide (NHS), and the resulting NHS-

activated intermediate was reacted with azido-PEG5-amine (**2**) in the presence of DIPEA to yield

the desired FA-Azide (**3**). Successful formation of the azide-functionalized folic acid was

confirmed by $^1$H NMR spectroscopy, which showed characteristic PEG proton signals from the

azide linker at δ 3.0 (2H), 3.4 (2H), and 3.5-3.7 (20H) ppm, along with a new amide proton signal

at δ 7.3 ppm (**Figures 2B and Supplementary Figure S1**). HPLC analysis confirmed the purity

of the product, exhibiting a single peak with a retention time of 13.3 minutes with more than 98%

purity (**Supplementary Figure S3**). To enable the subsequent CuAAC coupling, terminal alkyne

groups were introduced on the surface of 2DG-D. The primary hydroxyl groups of the surface-

exposed 2DG moieties were utilized for this modification. 2DG-D (**4**) was reacted with 5-hexynoic

acid (**5**) in the presence of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC)

and 4-(dimethylamino)pyridine (DMAP) to afford 2DG-D-Hexyne (**6**) (**Figure 2A**). $^1$H NMR

spectrum showed new methylene proton signals from the hexynoic linker in the aliphatic region at

δ 1.75-1.68 (8H), 2.46-2.34, and 2.74-2.58 (8H) ppm (**Figures 2B and Supplementary Figure**

**S5**), confirming the attachment of 4 hexyne arms. HPLC analysis showed a shift in retention time

from 14.0 minutes for 2DG-D (**4**) to 19.7 minutes for 2DG-D-Hexyne (**6**), confirming successful

surface modification (**Supplementary Figures S7 and S15**). Finally, the CuAAC reaction

between FA-Azide (**3**) and 2DG-D-Hexyne (**6**) yielded the FRβ$^+$ macrophage-targeted platform

FA-2DG-D (**7**). Previous studies have demonstrated that dendrimers bearing a single folic acid

ligand exhibit monovalent, reversible binding with folate-binding protein (FBP), whereas

dendrimers conjugated with two or more folic acid molecules engage in multivalent interactions,

resulting in stronger receptor binding [60-62]. Given the high hydrophobicity of folic acid,

conjugating more than 2 ligand molecules could potentially reduce the dendrimer's aqueous

solubility. Our CuAAC click chemistry approach ensured desired loading of folic acid on the

dendrimer. The number of attached molecules was confirmed through the comparative integration

of aromatic protons from the dendrimer at $\delta$7.2 ppm (16H), and the aromatic and amide protons of

folic acid at $\delta$8.7 ppm (2H), 7.7 ppm (4H), 7.0 ppm (4H) and at 6.7 ppm (4H). The HPLC

chromatogram showed a retention time shift from 19.7 minutes for 2DG-D-Hexyne (**6**) to 16.1

minutes for FA-2DG-D (**7**) (**Supplementary Figures S7** and **S10**). The purity of FA-2DG-D (**7**)

was determined to be more than 96% by HPLC (**Supplementary Figure S10**). Dynamic light

scattering (DLS) analysis revealed a hydrodynamic diameter of $6.6 \pm 0.4$ nm and a zeta potential

of $-5.8 \pm 0.3$ mV (**Supplementary Figures S11 and S12**).

To enable visualization and assessment of uptake of the dendrimer in FR$\beta^+$ macrophages

in endometriosis, the dendrimer was further functionalized with a near-infrared dye, cyanine 5

(Cy5). The FA-2DG-D (**7**) containing approximately two terminal alkynes on its surface was

conjugated with Cy5-azide (**8**) via a CuAAC reaction, yielding the fluorescently labeled FA-2DG-

D-Cy5 (**9**). Successful conjugation was confirmed by the appearance of characteristic Cy5 proton

signals in the $^1$H NMR spectrum (**Figure 2B**). The number of Cy5 molecules attached to the

dendrimer surface was estimated by proton integration. Comparison of the dendrimer aromatic

protons at $\delta$ 7.2 ppm (16H) with the Cy5 aromatic protons at $\delta$ 8.4 (4H), 8.2 (4H), 7.4 (4H), 6.6

(2H), and 6.4 ppm (2H) indicated the attachment of approximately two Cy5 molecules (**Figures**

**2B and Supplementary Figure S13**). The HPLC chromatogram of FA-2DG-D-Cy5 (**9**) exhibited

a retention time shift from 16.1 to 17.7 minutes following Cy5 conjugation, with a purity exceeding

97% (**Supplementary Figures S14** and **S15**). In addition, the Cy5-labeled dendrimer displayed a

characteristic UV absorbance at 650 nm, corresponding to Cy5, which further confirmed

successful conjugation. A Cy5-labeled 2DG-D dendrimer (2DG-D-Cy5), lacking the folic acid

targeting ligand, was prepared as a non-targeted control for uptake studies following our previously

reported procedure [27].

*3.3. Fluorescence Imaging and Quantitative Analysis of the Biodistribution of Systemically*

*Administered 2DG-D-Cy5 and FA-2DG-D-Cy5 in a Mouse Model of Endometriosis*

We first evaluated the selective FR$\beta^+$ macrophages targeting capability of FA-2DG-D-Cy5

in ELL-induced mice at 1, 6, 24, and 72 hours (n=3) and compared it with sham controls (n=3).

To evaluate the contribution of folic acid (FA) toward lesion targeting, biodistribution studies of

non-targeted 2DG-D-Cy5 in ELL-induced mice (n=3) were also performed under identical

conditions (**Figure 3**). One week after lesion induction, 50 mg/kg bw of fluorescently labeled (FA-

2DG-D-Cy5) or non-targeted (2DG-D-Cy5) was injected into the tail vein. Cy5 fluorescent signals

were calculated via the IVIS system at 1, 6, 24, and 72 hours. $FR\beta^+$ macrophages in the lesions

showed a pronounced elevation in fluorescent (Cy5) signal starting at 1 hour post-injection (green

arrows), which persisted above that of the 2DG-D-Cy5 group over the 72-hour observation period

(**Figure 3A**). The ex vivo fluorescent results at 72 hours showed specific uptake of FA-2DG-D-

Cy5 in the lesion, although some nonspecific uptake of FA-2DG-D-Cy5 and 2DG-D-Cy5 was

observed primarily in the liver and kidneys (**Figure 3B-C**). For more precise quantification, frozen

tissue samples from the 1, 6, 24, and 72-hour groups were homogenized, and fluorescence intensity

was measured to assess quantitative dendrimer uptake in lesions and major organs. Both FA-2DG-

D-Cy5 and 2DG-D-Cy5 showed strong, persistent accumulation within endometriosis lesions

beginning at 1-hour post-injection and sustained through 72 hours **(Figures 3D and S16)**. By 72

hours, lesion uptake of FA-2DG-D-Cy5 was significantly higher than that of 2DG-D-Cy5,

suggesting its targeted accumulation in $FR\beta^+$ macrophages that are abundant in endometriotic

lesions (**Figure 3D-E and Supplementary Figure S16**). Moreover, FA-2DG-D-Cy5

demonstrated minimal accumulation (<5% Injected Dose (% I.D.)) in other major organs,

indicating rapid clearance from off-target tissues **(Figure 3E)**. Importantly, unlike many

nanoparticle systems that exhibit up to ~80% hepatic accumulation, FA-2DG-D-Cy5 showed only

~4% liver accumulation, comparable to the previously characterized 2DG-D-Cy5 platform,

demonstrating its favorable pharmacokinetic profile [27].

**3.4. Confocal Imaging of Cellular Uptake of Intraperitoneally Administered FA-2DG-D-Cy5**

**and 2DG-D-Cy5 in a Mouse Model of Endometriosis**

We next examined the targeting ability, including selective cellular uptake, localization,

and retention of FA-2DG-D-Cy5 or 2DG-D-Cy5 in ELL-induced mice 2 weeks after dendrimer

administration (**Figure 4**). Confocal imaging showed that FA-2DGD-Cy5 (red) was selectively

targeted and taken up by $FR\beta^+$ macrophages (green), not only in lesions but also in peritoneal

exudate cells (**Figure 4A-B, white arrows**), where $FR\beta^+$ macrophages were elevated in ELL-

induced mice. In contrast, cellular uptake of 2DG-D-Cy5 without FA ligand was limited in the

lesion and peritoneal cells (**Figure 4A-B**). Given the potential uptake of non-targeted 2DG-D-Cy5
by activated macrophages [28], we further quantified nonspecific FA-2DG-D-Cy5 internalization
in peritoneal exudate cells. The uptake was an average of 0.59%, indicating minimal nonspecific
internalization. No dendrimer retention was observed in the kidneys, liver, or spleen in ELL-
induced mice 2 weeks following dendrimer administration **(Figure 4C)**, indicating that FA-2DG-
D-Cy5 underwent complete clearance, and off-target accumulation in major organs was negligible.
These results suggest that FA-2DGD-Cy5 dendrimer is internalized only by ligand-receptor
endocytosis in FRβ$^+$ macrophages in the lesions and peritoneal cells. As it remains stable within
cells for at least 2 weeks, it functions as an excellent nanocarrier for drug delivery.

*3.5. Synthesis and Characterization of FA-2DG-D-Niclo Conjugate Using Click Chemistry*

We developed a niclosamide-labeled dendrimer platform, FA-2DG-D-Niclo, by
conjugating niclosamide to the FA-2DG-D, to enable targeted delivery of niclosamide to disease-
driving FRβ$^+$ macrophages as a therapy for endometriosis. To achieve high conjugation efficiency
and precise drug loading, CuAAC was employed. A niclosamide bearing an azide linker (Niclo-
Azide) was first synthesized by functionalizing the phenolic -OH group of niclosamide (**Figure
5A**). Several linker architectures were explored to identify a construct that combined physiological
stability with controlled intracellular release. Initially, aliphatic and PEG-based carboxylic acids
were coupled to niclosamide through EDC/DMAP-mediated esterification, aiming to generate
ester linkages that are labile in the intracellular conditions and stable at physiological pH.
However, due to the proximity of the ester group to the aromatic ring, the resulting phenolic esters
exhibited instability and underwent rapid cleavage during subsequent CuAAC reactions [63]. To
circumvent this issue, a previous report on a disulfide-linker-based niclosamide prodrug was
followed with some modifications to design and synthesize niclosamide-azide (niclo-azide)
containing a disulfide linker [64]. It was reported that the disulfide linker remains stable under
extracellular conditions but undergoes selective cleavage in the intracellular environment, thereby
releasing free niclosamide [64].

The synthesis of niclo-azide was carried out through a multistep route (**Figure 5A**). The
Boc-protected disulfide derivative **(10)** was synthesized using a published procedure [64]. It was
then deprotected using 4 N HCl in dioxane to afford compound **11** as the amine hydrochloride salt,
which was directly converted to the corresponding carbamoyl chloride intermediate **12** using

triphosgene, following coupling with niclosamide **(13)** under mild basic conditions to afford the desired Niclo-Azide **(14) (Figure 5A)**. Successful formation of the product was confirmed by comparative integration in ¹H NMR spectroscopy, which showed characteristic aromatic protons from the niclosamide scaffold δ 8.4-7.3 ppm (6H) and the aliphatic protons from the disulfide linker between δ 3.7-2.7 ppm (11H) **(Figures 5B and Supplementary Figure S19)**. The product purity (~ 95.9%), was verified by HPLC, which showed a chromatogram with a retention time of 14.7 minutes **(Figures 6A and Supplementary Figure S21).** To enable subsequent CuAAC coupling, 2DG-D **(4)** was partially modified with 5-hexynoic acid. Approximately 12 hydroxyl groups per dendrimer were modified to yield 2DG-D-Hexyne' **(15) (Figure 5A)**. The formation of the required product was confirmed by the appearance of characteristic methylene proton signals from the hexynoic linker in the aliphatic region of the ¹H NMR spectrum at δ 1.7 (24H), 2.4 (24H) and at 2.2 ppm **(Figures 5B and Supplementary Figure S23)** which confirmed the presence of 12 hexyne arms. HPLC analysis of 2DG-D-Hexyne' **(15)** showed the chromatogram at 10.4 minutes with a purity exceeding 97% **(Figure 6A)**. Subsequently, a CuAAC reaction between FA-Azide **(3)** and 2DG-D-Hexyne' **(15)** yielded FA-2DG-D' **(16)**. The ¹H NMR spectrum of FA-2DG-D' **(16)** showed characteristic aromatic resonances corresponding to folic acid, confirming successful conjugation of 2 molecules per dendrimer **(Supplementary Figure S26).** The resulting construct retained approximately ten free terminal alkyne groups per dendrimer. The HPLC chromatogram exhibited a retention time shift from 10.4 minutes for 2DG-D-Hexyne' to 9.5 minutes for FA-2DG-D' **(16)**, with a purity of ~99% **(Figures 6A and Supplementary Figure S28)**. Lastly, FA-2DG-D' **(16)** was conjugated with Niclo-Azide **(14)** through CuAAC to obtain FA-2DG-D-Niclo **(17)**. Consistent with our previous studies [56, 58], we aimed to maintain the drug loading between 10 and 15% to ensure aqueous solubility of the dendrimer conjugate. Drug loading was calculated by ¹H NMR spectroscopy through comparison of the dendrimer aromatic protons at δ 7.2 ppm (16H) with the newly aromatic signals at δ 8.4 (10H), 8.3 (10H), and 8.2-7.8 ppm, along with distinct amide protons of niclosamide at δ 10.43 ppm (10H), confirmed the conjugation of approximately ten niclosamide molecules per dendrimer, corresponding to ~12% drug loading by weight **(Figures 5B and Supplementary Figure S29)**. HPLC analysis revealed a shift in retention time from 9.5 minutes for FA-2DG-D' **(17)** to 11.2 minutes for FA-2DG-D-Niclo **(17)**, with a final purity of more than 98% **(Supplementary Figure S31)**. DLS analysis indicated

a hydrodynamic diameter of 7.6 ± 0.6 nm and a zeta potential of -7.7 ± 0.6 mV **(Figure 6C and Supplementary Figures S32 and S33)**.

The physicochemical properties of FA-2DG-D-Niclo **(23)** are presented in **Figure 7A.** While niclosamide is practically insoluble in water (~5 µg/mL) [65], the dendrimer-drug conjugate exhibited excellent aqueous solubility (~150 mg/mL). At a drug loading of 12%, this corresponds to an effective niclosamide solubility of approximately 18 mg/mL in water, representing a ~3,600 fold increase in solubility compared to free niclosamide **(Figure 7A)**.

The shelf stability of FA-2DG-D-Niclo **(17)** (1 mg/mL) was assessed in PBS at 4 °C, room temperature (25 °C), and 40 °C over a 30-day period. The formulation remained stable throughout the study, retaining more than 96% purity by HPLC at all tested temperatures **(Supplementary Figure S34)**. No shifts in retention time were observed over 30 days, confirming the chemical stability of the formulation under these conditions.

### 3.6. Scalable and Reproducible Synthesis of FA-2DG-D-Niclo

Ensuring synthetic reproducibility and scalability remains a central barrier in advancing nanoparticle-based therapeutics toward clinical applications. To overcome these challenges, we established a streamlined and scalable synthetic workflow for FA-2DG-D-Niclo **(17)** that enables rigorous characterization of intermediates at each stage. To assess batch-to-batch consistency, three independent 400 mg-scale batches were synthesized and evaluated by $^1$H NMR and HPLC. The $^1$H NMR spectra of all three batches **(Figure 7B, left)** showed identical chemical profiles and consistent drug and ligand loading. Similarly, HPLC analysis revealed overlapping chromatogram peaks with retention times of ~11.2 minutes and purities exceeding 98%, confirming robust batch-to-batch reproducibility **(Figure 7B, right)**.

### 3.7. In vitro drug release studies for FA-2DG-D-Niclo

We further assessed the in vitro drug-release profile of the conjugate under conditions mimicking both the extracellular environment (PBS + 0.1 mM GSH and 100% human plasma) and the intracellular microenvironment (pH 5.5 citrate buffer + 10mM GSH) **(Figure 7C)**. Since niclosamide is attached to the dendrimer through a disulfide linkage, it enables glutathione (GSH) responsive cleavage [66]. Upon cleavage, the generated free thiol is known to undergo an intramolecular nucleophilic attack on the tertiary carbamate, ultimately yielding free niclosamide

[64]. When conjugated to the targeted FA-2DG-D, this design facilitates controlled release within

the intracellular environments while limiting premature systemic exposure outside the lesion sites.

In PBS and 0.1mM GSH, the conjugate was found to be relatively stable. Less than 40% of the

drug was observed to be released in around 2 weeks, with <10% drug release detected for the first

5 days. Notably less than 5% of the drug was released within 8 hours. When incubated in human

plasma, less than 35% of the drug was released over 5 days, indicating stability in physiological

conditions **(Figure 7C).** Under intracellular conditions (pH 5.5 citrate buffer + 10mM GSH), the

conjugate showed a sustained release profile, with ~45% released in the first 24 hours, ~80% by

two days, and reaching ~95% over 12 days **(Figure 7C)**. The targeted and sustained intracellular

drug release profile is important to maintain the efficacy of the drug for a long time and to avoid

off-target effects.

***3.8. FA-2DG-D-Niclo attenuates lesion progression and endometriosis-associated hyperalgesia.***

Lastly, we assessed the efficacy of FA-2DG-D-Niclo in endometriosis disease progression

and endometriosis-associated hyperalgesia. After the lesion induction, a single dose of FA-2DG-

D-Niclo dendrimer (25 or 50 mg/kg bw of niclosamide) or FA-2DG-D without niclosamide control

was administered i.p. in ELL-induced mice at Day 7, as shown in **Figure 8A**. Then, the efficacy

of dendrimers was examined 7 (Day 14) and 14 (Day 21) days after dendrimer administration. We

first confirmed the suppressing capability of FRβ$^+$ macrophages in the lesions and peritoneal cavity

by FA-2DG-D-Niclo dendrimer (**Figure 8B-C**). Single dosing of FA-2DG-D-Niclo (50 mg/kg bw)

significantly reduced FRβ$^+$ macrophages in the lesions at Day 21, although the reduction of FRβ$^+$

macrophages was not observed at Day 14 (**Figure 8B**). Peritoneal FRβ$^+$ macrophage population

was reduced by both doses (25 and 50 mg/kg bw) of FA-2DG-D-Niclo (**Figure 8C**). The results

indicate that 7 days after dendrimer administration may not be sufficient to reduce FRβ$^+$

macrophages in both lesions and the peritoneum. Indeed, lesion number and volume were also

significantly reduced at Day 21 by FA-2DG-D-Niclo at both doses, but there were no significant

differences in lesion number or volume at Day 14 (**Figure 9A**). However, we realized that three

mice treated with both doses and examined at Day 14 did not have lesions. Furthermore, four and

five mice, at doses of 25 and 50 mg/kg bw of niclosamide, respectively, did not have lesions by

Day 21, either. Thus, a single dose of FA-2DG-D-Niclo administration completely inhibited lesion

growth and progression in 37.5-62.5% of mice, whereas all control mice developed lesions on

Days 14 and 21. While reduced FRβ$^+$ macrophages in the remaining lesions were observed only
at Day 21, a few mice at Day 14 lacked lesions for examination, suggesting that FA-2DG-D-Niclo
was likely effective for lesion growth 7 days after administration. In support of this implication,
abdominal and hindpaw hypersensitivities were already improved at Day 14 (**Figure 9B**). The
abdominal and hindpaw sensitivity 7 days after lesion induction was high compared to before
lesion induction (Day -1). At Day 14, FA-2DG-D-Niclo at a dose of 50 mg/kg bw of niclosamide
significantly reduced the abdominal sensitivity, which is considered indicative of peripheral
visceral pain. Both doses of FA-2DG-D-Niclo improved hindpaw sensitivity, which is likely be
affected by peripheral and central sensitization. By Day 21, FA-2DG-D-Niclo at both doses further
improved abdominal and hindpaw sensitivities. The results clearly showed that a single dose of
FA-2DG-D-Niclo inhibited disease progression and endometriosis-associated hyperalgesia.
Notably, a single dose of FA-2DG-D-Niclo remained effective for 2 weeks after administration.
The current FDA-approved niclosamide treatment requires a high dose (2g/day) due to its low
water solubility and is limited to oral administration for 7 days. Niclosamide treatment for patients
who have endometriosis would need to be longer due to the chronic nature of the disease. Long-
term systemic treatment with niclosamide could cause off-target actions. Therefore, the 2DG-D-
based delivery of niclosamide and targeting of disease-specific FRβ$^+$ macrophages is highly
translational, as FA-2DG-D can deliver niclosamide intracellularly into FRβ$^+$ macrophages,
thereby enhancing efficacy, reducing side effects, and treating chronic diseases such as
endometriosis.
594

**4. Conclusions**

With integrated single-cell transcriptomic analysis, we further identified a significant
recruitment of pro-disease FRβ$^+$ macrophages within human endometriotic lesions and the pelvic
cavity, mirroring our observations in the mouse model. Importantly, overexpression of FRβ on
these macrophages provides a ligand-guided entry point, enabling receptor-specific targeting for
both imaging and therapeutic interventions. Leveraging this receptor overexpression, we
engineered a folic acid-conjugated dendrimer (FA-2DG-D) to exploit FRβ-mediated uptake for
targeted imaging and drug delivery in endometriotic lesions. FA-2DG-D offers several critical
advantages for translational development. The dendrimer architecture enables precise,
reproducible conjugation of targeting ligands and therapeutic payloads via click chemistry,

resulting in well-defined nanostructures, excellent aqueous solubility, and robust batch-to-batch consistency. Conjugation of niclosamide to generate FA-2DG-D-Niclo markedly enhanced the solubility of this poorly water-soluble drug, enabled controlled, intracellularly triggered release via a disulfide linker, and minimized premature systemic exposure. Importantly, the formulation demonstrated high stability under physiological and storage conditions. In a mouse model of endometriosis, the Cy5-labelled platform demonstrated lesion and peritoneal cell-specific accumulation and intracellular internalization along with a favorable biodistribution with minimal off-target organ accumulation. FA-2DG-D-Niclo enabled targeted delivery to disease-driving macrophages. Administration of FA-2DG-D-Niclo led to significant reductions in lesion number and volume, and improved hyperalgesia, highlighting both therapeutic efficacy and immunomodulatory activity. This work demonstrates the potential of FA-2DG-D platform to target FRβ⁺ macrophages for the treatment of endometriosis. By addressing key limitations of current hormonal and systemic treatments, this approach establishes a non-hormonal, FRβ⁺ macrophage-focused strategy for the treatment of endometriosis.

**Supporting Information**

The chemistry experimental details and supplementary figures related to characterization, stability studies, and bioditribution are provided in supporting information.

**Data Availability Statement**

The raw data of scRNA sequencing have been deposited to NCBI/SRA (PRJNA1418167). Additional data or materials related to the findings of this study are available from the corresponding author upon reasonable request.

**Funding Information**

This work was supported by NIH grant R01HD104619 (to Kanako Hayashi) and a Washington Research Foundation Technology Commercialization Phase 1 award (to Kanako Hayashi and Anjali Sharma).

**Declaration of Interest Statement:**

635     The authors A.S, K.H, R.S, A.D, M.S, M.E.H, and A.I.D have pending patents and
636 invention disclosures on 2DG and/or FA-2DG dendrimers and their conjugates.
637

**Authorship Contribution Statement**
A.S. and K.H. designed the research; A.D., A.S., and R.S. designed and synthesized the
dendrimers; M.S., M.H., and T.P. performed the animal experiments and/or analyzed the data;
A.D., K.J.G., and A.I.D. performed quantitative biodistribution analysis; N.S.M., V.K.S., and
P.S.C. collected/or prepared the human samples; M.S. performed single-cell RNA sequencing
and analysis; A.D., M.S., R.S., A.S., and K.H. wrote the paper; all authors read, reviewed, edited,
and approved the manuscript.
645
646

## References

1.    Zondervan, K. T.; Becker, C. M.; Missmer, S. A.  **Endometriosis**. *The New England journal of medicine* 2020, **382**, (13), 1244-1256.

2.    Burney, R. O.; Giudice, L. C.  **Pathogenesis and pathophysiology of endometriosis**. *Fertility and sterility* 2012, **98**, (3), 511-9.

3.    As-Sanie, S.; Mackenzie, S. C.; Morrison, L.; Schrepf, A.; Zondervan, K. T.; Horne, A. W.; Missmer, S. A.  **Endometriosis: A Review**. *JAMA* 2025, **334**, (1), 64-78.

4.    Giudice, L. C.; Kao, L. C.  **Endometriosis**. *Lancet (London, England)* 2004, **364**, (9447), 1789-99.

5.    Ellis, K.; Munro, D.; Clarke, J.  **Endometriosis Is Undervalued: A Call to Action**. *Frontiers in global women's health* 2022, **3**, 902371.

6.    Marla, S.; Mortlock, S.; Heinosalo, T.; Poutanen, M.; Montgomery, G. W.; McKinnon, B. D.  **Gene expression profiles separate endometriosis lesion subtypes and indicate a sensitivity of endometrioma to estrogen suppressive treatments through elevated ESR2 expression**. *BMC Medicine* 2023, **21**, (1), 460.

7.    Edi, R.; Cheng, T.  **Endometriosis: Evaluation and Treatment**. *American family physician* 2022, **106**, (4), 397-404.

8.    Alonso, A.; Gunther, K.; Maheux-Lacroix, S.; Abbott, J.  **Medical management of endometriosis**. *Current opinion in obstetrics & gynecology* 2024, **36**, (5), 353-361.

9.    Guo, S.-W.  **Recurrence of endometriosis and its control**. *Human Reproduction Update* 2009, **15**, (4), 441-461.

10.    Singh, S. S.; Gude, K.; Perdeaux, E.; Gattrell, W. T.; Becker, C. M.  **Surgical Outcomes in Patients With Endometriosis: A Systematic Review**. *Journal of obstetrics and gynaecology Canada : JOGC = Journal d'obstetrique et gynecologie du Canada : JOGC* 2020, **42**, (7), 881-888.e11.

11.    Sekulovski, N.; Whorton, A. E.; Shi, M.; MacLean, J. A., II; Hayashi, K.  **Endometriotic inflammatory microenvironment induced by macrophages can be targeted by niclosamide†**. *Biology of reproduction* 2019, **100**, (2), 398-408.

12.    Sekulovski, N.; Whorton, A. E.; Tanaka, T.; Hirota, Y.; Shi, M.; MacLean, J. A.; de Mola, J. R. L.; Groesch, K.; Diaz-Sylvester, P.; Wilson, T.; Hayashi, K.  **Niclosamide suppresses macrophage-induced inflammation in endometriosis†**. *Biology of reproduction* 2020, **102**, (5), 1011-1019.

13.    Shi, M.; Sekulovski, N.; Whorton, A. E.; MacLean, J. A., 2nd; Greaves, E.; Hayashi, K.  **Efficacy of niclosamide on the intra-abdominal inflammatory environment in endometriosis**. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* 2021, **35**, (5), e21584.

14.    Zhao, L.; Shi, M.; Winuthayanon, S.; MacLean, J. A., 2nd; Hayashi, K.  **Niclosamide targets the dynamic progression of macrophages for the resolution of endometriosis in a mouse model**. *Communications biology* 2022, **5**, (1), 1225.

15.    Tan, Y.; Flynn, W. F.; Sivajothi, S.; Luo, D.; Bozal, S. B.; Davé, M.; Luciano, A. A.; Robson, P.; Luciano, D. E.; Courtois, E. T.  **Single-cell analysis of endometriosis reveals a coordinated transcriptional programme driving immunotolerance and angiogenesis across eutopic and ectopic tissues**. *Nature cell biology* 2022, **24**, (8), 1306-1318.

16.    Prather, G. R.; MacLean, J. A., 2nd; Shi, M.; Boadu, D. K.; Paquet, M.; Hayashi, K.  **Niclosamide As a Potential Nonsteroidal Therapy for Endometriosis That Preserves**

**Reproductive Function in an Experimental Mouse Model**. *Biology of reproduction* 2016, **95**, (4), 76.

17.      Chen, W.; Mook, R. A., Jr.; Premont, R. T.; Wang, J.   **Niclosamide: Beyond an antihelminthic drug**. *Cellular signalling* 2018, **41**, 89-96.

18.      Jug, M.; Laffleur, F.; Millotti, G.   **Revisiting Niclosamide Formulation Approaches - a Pathway Toward Drug Repositioning**. *Drug design, development and therapy* 2024, **18**, 4153-4182.

19.      Rani, A.; Bisen, S.; Sharma, R.; Dar, A. I.; Sharma, A.; Dhull, A.; Palmer, N.; Goody, K. J.; Singh, N. K.; Sharma, A.   **Systemic targeting of aberrant neovascular tufts using trehalose-dendrimer nanocarriers for the treatment of proliferative retinopathies**. *Theranostics* 2026, **16**, (3), 1504-1527.

20.      Pulukuri, A. J.; Dhull, A.; Dar, A. I.; Rani, A.; Sharma, R.; Berkman, C. E.; Sharma, A.   **Dendrimer-Mediated Delivery Enhances Therapeutic Efficacy in Triple-Negative Breast Cancer**. *Biomacromolecules* 2025, **26**, (9), 5979-6000.

21.      Sharma, R.; Porterfield, J. E.; An, H.-T.; Jimenez, A. S.; Lee, S.; Kannan, S.; Sharma, A.; Kannan, R. M.   **Rationally Designed Galactose Dendrimer for Hepatocyte-Specific Targeting and Intracellular Drug Delivery for the Treatment of Liver Disorders**. *Biomacromolecules* 2021, **22**, (8), 3574-3589.

22.      Monge, P.; Søgaard, A. B.; Andersen, D. G.; Chandrawati, R.; Zelikin, A. N.   **Synthetic chemical ligands and cognate antibodies for biorthogonal drug targeting and cell engineering**. *Advanced Drug Delivery Reviews* 2021, **170**, 281-293.

23.      Nam, H. Y.; Nam, K.; Hahn, H. J.; Kim, B. H.; Lim, H. J.; Kim, H. J.; Choi, J. S.; Park, J.-S.   **Biodegradable PAMAM ester for enhanced transfection efficiency with low cytotoxicity**. *Biomaterials* 2009, **30**, (4), 665-673.

24.      Huang, W.-C.; Lu, I. L.; Chiang, W.-H.; Lin, Y.-W.; Tsai, Y.-C.; Chen, H.-H.; Chang, C.-W.; Chiang, C.-S.; Chiu, H.-C.   **Tumortropic adipose-derived stem cells carrying smart nanotherapeutics for targeted delivery and dual-modality therapy of orthotopic glioblastoma**. *Journal of Controlled Release* 2017, **254**, 119-130.

25.      Dhull, A.; Yu, C.; Wilmoth, A. H.; Chen, M.; Sharma, A.; Yiu, S.   **Dendrimers in Corneal Drug Delivery: Recent Developments and Translational Opportunities**. *Pharmaceutics* 2023, **15**, (6).

26.      Dhull, A.; Wei, J.; Pulukuri, A. J.; Rani, A.; Sharma, R.; Mesbahi, N.; Yoon, H.; Savoy, E. A.; Xaivong Vi, S.; Goody, K. J.; Berkman, C. E.; Wu, B. J.; Sharma, A.   **PSMA-targeted dendrimer as an efficient anticancer drug delivery vehicle for prostate cancer**. *Nanoscale* 2024, **16**, (11), 5634-5652.

27.      Dhull, A.; Zhang, Z.; Sharma, R.; Dar, A. I.; Rani, A.; Wei, J.; Gopalakrishnan, S.; Ghannam, A.; Hahn, V.; Pulukuri, A. J.; Tasevski, S.; Moughni, S.; Wu, B. J.; Sharma, A.   **Discovery of 2-deoxy glucose surfaced mixed layer dendrimer: a smart neuron targeted systemic drug delivery system for brain diseases**. *Theranostics* 2024, **14**, (8), 3221-3245.

28.      Dhull, A.; Park, K. W.; Dar, A. I.; Wang, A.; Rani, A.; Sharma, R.; Valdez, T. A.; Sharma, A.   **Mixed-Layered Glycodendrimer Probe for Imaging Inflammation at Surgical Site Infections**. *ACS Sensors* 2025, **10**, (3), 2234-2243.

29.      Puligujja, P.; McMillan, J.; Kendrick, L.; Li, T.; Balkundi, S.; Smith, N.; Veerubhotla, R. S.; Edagwa, B. J.; Kabanov, A. V.; Bronich, T.; Gendelman, H. E.; Liu, X. M.   **Macrophage folate receptor-targeted antiretroviral therapy facilitates drug entry, retention, antiretroviral**

activities and biodistribution for reduction of human immunodeficiency virus infections. *Nanomedicine : nanotechnology, biology, and medicine* 2013, **9**, (8), 1263-73.

30.	Chandrupatla, D.; Molthoff, C. F. M.; Lammertsma, A. A.; van der Laken, C. J.; Jansen, G. **The folate receptor β as a macrophage-mediated imaging and therapeutic target in rheumatoid arthritis**. *Drug delivery and translational research* 2019, **9**, (1), 366-378.

31.	Zheng, G. X.; Terry, J. M.; Belgrader, P.; Ryvkin, P.; Bent, Z. W.; Wilson, R.; Ziraldo, S. B.; Wheeler, T. D.; McDermott, G. P.; Zhu, J.; Gregory, M. T.; Shuga, J.; Montesclaros, L.; Underwood, J. G.; Masquelier, D. A.; Nishimura, S. Y.; Schnall-Levin, M.; Wyatt, P. W.; Hindson, C. M.; Bharadwaj, R.; Wong, A.; Ness, K. D.; Beppu, L. W.; Deeg, H. J.; McFarland, C.; Loeb, K. R.; Valente, W. J.; Ericson, N. G.; Stevens, E. A.; Radich, J. P.; Mikkelsen, T. S.; Hindson, B. J.; Bielas, J. H. **Massively parallel digital transcriptional profiling of single cells**. *Nature communications* 2017, **8**, 14049.

32.	Wang, W.; Vilella, F.; Alama, P.; Moreno, I.; Mignardi, M.; Isakova, A.; Pan, W.; Simon, C.; Quake, S. R. **Single-cell transcriptomic atlas of the human endometrium during the menstrual cycle**. *Nature medicine* 2020, **26**, (10), 1644-1653.

33.	Lai, Z. Z.; Wang, Y.; Zhou, W. J.; Liang, Z.; Shi, J. W.; Yang, H. L.; Xie, F.; Chen, W. D.; Zhu, R.; Zhang, C.; Mei, J.; Zhao, J. Y.; Ye, J. F.; Zhang, T.; Li, M. Q. **Single-cell transcriptome profiling of the human endometrium of patients with recurrent implantation failure**. *Theranostics* 2022, **12**, (15), 6527-6547.

34.	Garcia-Alonso, L.; Handfield, L. F.; Roberts, K.; Nikolakopoulou, K.; Fernando, R. C.; Gardner, L.; Woodhams, B.; Arutyunyan, A.; Polanski, K.; Hoo, R.; Sancho-Serra, C.; Li, T.; Kwakwa, K.; Tuck, E.; Lorenzi, V.; Massalha, H.; Prete, M.; Kleshchevnikov, V.; Tarkowska, A.; Porter, T.; Mazzeo, C. I.; van Dongen, S.; Dabrowska, M.; Vaskivskyi, V.; Mahbubani, K. T.; Park, J. E.; Jimenez-Linan, M.; Campos, L.; Kiselev, V. Y.; Lindskog, C.; Ayuk, P.; Prigmore, E.; Stratton, M. R.; Saeb-Parsy, K.; Moffett, A.; Moore, L.; Bayraktar, O. A.; Teichmann, S. A.; Turco, M. Y.; Vento-Tormo, R. **Mapping the temporal and spatial dynamics of the human endometrium in vivo and in vitro**. *Nature genetics* 2021, **53**, (12), 1698-1711.

35.	Fonseca, M. A. S.; Haro, M.; Wright, K. N.; Lin, X.; Abbasi, F.; Sun, J.; Hernandez, L.; Orr, N. L.; Hong, J.; Choi-Kuaea, Y.; Maluf, H. M.; Balzer, B. L.; Fishburn, A.; Hickey, R.; Cass, I.; Goodridge, H. S.; Truong, M.; Wang, Y.; Pisarska, M. D.; Dinh, H. Q.; El-Naggar, A.; Huntsman, D. G.; Anglesio, M. S.; Goodman, M. T.; Medeiros, F.; Siedhoff, M.; Lawrenson, K. **Single-cell transcriptomic analysis of endometriosis**. *Nature genetics* 2023, **55**, (2), 255-267.

36.	Huang, X.; Wu, L.; Pei, T.; Liu, D.; Liu, C.; Luo, B.; Xiao, L.; Li, Y.; Wang, R.; Ouyang, Y.; Zhu, H.; Huang, W. **Single-cell transcriptome analysis reveals endometrial immune microenvironment in minimal/mild endometriosis**. *Clinical and experimental immunology* 2023, **212**, (3), 285-295.

37.	Marečková, M.; Garcia-Alonso, L.; Moullet, M.; Lorenzi, V.; Petryszak, R.; Sancho-Serra, C.; Oszlanczi, A.; Icoresi Mazzeo, C.; Wong, F. C. K.; Kelava, I.; Hoffman, S.; Krassowski, M.; Garbutt, K.; Gaitskell, K.; Yancheva, S.; Woon, E. V.; Male, V.; Granne, I.; Hellner, K.; Mahbubani, K. T.; Saeb-Parsy, K.; Lotfollahi, M.; Prigmore, E.; Southcombe, J.; Dragovic, R. A.; Becker, C. M.; Zondervan, K. T.; Vento-Tormo, R. **An integrated single-cell reference atlas of the human endometrium**. *Nature genetics* 2024, **56**, (9), 1925-1937.

38.	Shin, S.; Chung, Y. J.; Moon, S. W.; Choi, E. J.; Kim, M. R.; Chung, Y. J.; Lee, S. H. **Single-cell profiling identifies distinct hormonal, immunologic, and inflammatory signatures of endometriosis-constituting cells**. *The Journal of pathology* 2023, **261**, (3), 323-334.

39. Han, J.; Gallerand, A.; Erlich, E. C.; Helmink, B. A.; Mair, I.; Li, X.; Eckhouse, S. R.; Dimou, F. M.; Shakhsheer, B. A.; Phelps, H. M.; Chan, M. M.; Mintz, R. L.; Lee, D. D.; Schilling, J. D.; Finlay, C. M.; Allen, J. E.; Jakubzick, C. V.; Else, K. J.; Onufer, E. J.; Zhang, N.; Randolph, G. J. **Human serous cavity macrophages and dendritic cells possess counterparts in the mouse with a distinct distribution between species**. *Nature immunology* 2024, **25**, (1), 155-165.

40. Zou, G.; Wang, J.; Xu, X.; Xu, P.; Zhu, L.; Yu, Q.; Peng, Y.; Guo, X.; Li, T.; Zhang, X. **Cell subtypes and immune dysfunction in peritoneal fluid of endometriosis revealed by single-cell RNA-sequencing**. *Cell & bioscience* 2021, **11**, (1), 98.

41. Wolf, F. A.; Angerer, P.; Theis, F. J. **SCANPY: large-scale single-cell gene expression data analysis**. *Genome biology* 2018, **19**, (1), 15.

42. Wolock, S. L.; Lopez, R.; Klein, A. M. **Scrublet: Computational Identification of Cell Doublets in Single-Cell Transcriptomic Data**. *Cell systems* 2019, **8**, (4), 281-291.e9.

43. Giotti, B.; Chen, S. H.; Barnett, M. W.; Regan, T.; Ly, T.; Wiemann, S.; Hume, D. A.; Freeman, T. C. **Assembly of a parts list of the human mitotic cell cycle machinery**. *Journal of molecular cell biology* 2019, **11**, (8), 703-718.

44. Korsunsky, I.; Millard, N.; Fan, J.; Slowikowski, K.; Zhang, F.; Wei, K.; Baglaenko, Y.; Brenner, M.; Loh, P. R.; Raychaudhuri, S. **Fast, sensitive and accurate integration of single-cell data with Harmony**. *Nature methods* 2019, **16**, (12), 1289-1296.

45. Traag, V. A.; Waltman, L.; van Eck, N. J. **From Louvain to Leiden: guaranteeing well-connected communities**. *Scientific reports* 2019, **9**, (1), 5233.

46. Soneson, C.; Robinson, M. D. **Bias, robustness and scalability in single-cell differential expression analysis**. *Nature methods* 2018, **15**, (4), 255-261.

47. Wu, C.; Boey, D.; Bril, O.; Grootens, J.; Vijayabaskar, M. S.; Sorini, C.; Ekoff, M.; Wilson, N. K.; Ungerstedt, J. S.; Nilsson, G.; Dahlin, J. S. **Single-cell transcriptomics reveals the identity and regulators of human mast cell progenitors**. *Blood advances* 2022, **6**, (15), 4439-4449.

48. Panjwani, M. K.; Grassmann, S.; Sottile, R.; Le Luduec, J. B.; Kontopoulos, T.; van der Ploeg, K.; Sun, J. C.; Hsu, K. C. **Single-cell profiling aligns CD56(bright) and cytomegalovirus-induced adaptive natural killer cells to a naïve-memory relationship**. *Frontiers in immunology* 2024, **15**, 1499492.

49. Yu, G.; Wang, L. G.; Han, Y.; He, Q. Y. **clusterProfiler: an R package for comparing biological themes among gene clusters**. *Omics : a journal of integrative biology* 2012, **16**, (5), 284-7.

50. Nothnick, W. B.; Colvin, A.; Cheng, K. F.; Al-Abed, Y. **Inhibition of macrophage migration inhibitory factor reduces endometriotic implant size in mice with experimentally induced disease**. *Journal of endometriosis* 2011, **3**, (3), 135-142.

51. Herup-Wheeler, T.; Shi, M.; Harvey, M. E.; Talwar, C.; Kommagani, R.; MacLean, J. A., 2nd; Hayashi, K. **High-fat diets promote peritoneal inflammation and augment endometriosis-associated abdominal hyperalgesia**. *Frontiers in endocrinology* 2024, **15**, 1336496.

52. Shi, M.; MacLean, J. A., 2nd; Hayashi, K. **The involvement of peritoneal GATA6(+) macrophages in the pathogenesis of endometriosis**. *Frontiers in immunology* 2024, **15**, 1396000.

53. Davies, L. C.; Rice, C. M.; Palmieri, E. M.; Taylor, P. R.; Kuhns, D. B.; McVicar, D. W. **Peritoneal tissue-resident macrophages are metabolically poised to engage microbes using tissue-niche fuels**. *Nature communications* 2017, **8**, (1), 2074.

54.     Vega-Pérez, A.; Villarrubia, L. H.; Godio, C.; Gutiérrez-González, A.; Feo-Lucas, L.; Ferriz, M.; Martínez-Puente, N.; Alcaín, J.; Mora, A.; Sabio, G.; López-Bravo, M.; Ardavín, C. **Resident macrophage-dependent immune cell scaffolds drive anti-bacterial defense in the peritoneal cavity**. *Immunity* 2021, **54**, (11), 2578-2594.e5.

55.     Jiang, D.; Xiao, R.; Bai, J.; Wang, X.; Fang, Y.; Liu, J.; Wang, X.; Cui, M.; You, L.; Zhao, Y. **FOLR2(+) macrophages in cancer: allies or enemies**. *Cell communication and signaling : CCS* 2025, **23**, (1), 261.

56.     Rani, A.; Pulukuri, A. J.; Wei, J.; Dhull, A.; Dar, A. I.; Sharma, R.; Mesbahi, N.; Savoy, E. A.; Yoon, H.; Wu, B. J.; Berkman, C. E.; Sharma, A. **PSMA-Targeted 2-Deoxyglucose-Based Dendrimer Nanomedicine for the Treatment of Prostate Cancer**. *Biomacromolecules* 2024, **25**, (9), 6164-6180.

57.     Dar, A. I.; Jain, V.; Rani, A.; Pulukuri, A. J.; Gonzalez, J. C.; Dhull, A.; Sharma, R.; Sharma, A. **Silibinin-Conjugated Galactose Dendrimers for Targeted Treatment of Hepatocellular Carcinoma**. *ACS Applied Materials & Interfaces* 2025, **17**, (14), 20980-21000.

58.     Dar, A. I.; Zhang, Z.; Gopalakrishnan, S.; Sharma, R.; Pulukuri, A. J.; Rani, A.; Dhull, A.; Gonzalez, J. C.; Atoui, T.; Mashal, Y.; Naseer, Z.; Calmi, J.; Sharma, A. **Neuron-targeted 2-deoxyglucose-dendrimer-rosiglitazone nanotherapy mitigates neuroinflammation and cognitive deficits in pediatric traumatic brain injury**. *Bioengineering & Translational Medicine* **n/a**, (n/a), e70053.

59.     Chen, C.; Ke, J.; Zhou, X. E.; Yi, W.; Brunzelle, J. S.; Li, J.; Yong, E. L.; Xu, H. E.; Melcher, K. **Structural basis for molecular recognition of folic acid by folate receptors**. *Nature* 2013, **500**, (7463), 486-9.

60.     van Dongen, M. A.; Silpe, J. E.; Dougherty, C. A.; Kanduluru, A. K.; Choi, S. K.; Orr, B. G.; Low, P. S.; Banaszak Holl, M. M. **Avidity mechanism of dendrimer-folic acid conjugates**. *Molecular pharmaceutics* 2014, **11**, (5), 1696-706.

61.     Hong, S.; Leroueil, P. R.; Majoros, I. J.; Orr, B. G.; Baker, J. R., Jr.; Banaszak Holl, M. M. **The binding avidity of a nanoparticle-based multivalent targeted drug delivery platform**. *Chemistry & biology* 2007, **14**, (1), 107-15.

62.     Thomas, T. P.; Huang, B.; Choi, S. K.; Silpe, J. E.; Kotlyar, A.; Desai, A. M.; Zong, H.; Gam, J.; Joice, M.; Baker, J. R., Jr. **Polyvalent dendrimer-methotrexate as a folate receptor-targeted cancer therapeutic**. *Molecular pharmaceutics* 2012, **9**, (9), 2669-2676.

63.     de Souza, H. M. R.; Guedes, J. S.; Freitas, R.; Gelves, L. G. V.; Fokoue, H. H.; Sant'Anna, C. M. R.; Barreiro, E. J.; Lima, L. M. **Comparative chemical and biological hydrolytic stability of homologous esters and isosteres**. *Journal of enzyme inhibition and medicinal chemistry* 2022, **37**, (1), 718-727.

64.     Kim, J. H.; Park, S.; Jung, E.; Shin, J.; Kim, Y. J.; Kim, J. Y.; Sessler, J. L.; Seo, J. H.; Kim, J. S. **A dual-action niclosamide-based prodrug that targets cancer stem cells and inhibits TNBC metastasis**. *Proceedings of the National Academy of Sciences of the United States of America* 2023, **120**, (21), e2304081120.

65.     Li, J.; Chang, Y.; Zou, Z. Z. **Enhancing the in vivo bioavailability and absorption of Niclosamide with amorphous solid dispersion via solvent method**. *RPS Pharmacy and Pharmacology Reports* 2024, **3**, (4).

66.     Navath, R. S.; Kurtoglu, Y. E.; Wang, B.; Kannan, S.; Romero, R.; Kannan, R. M. **Dendrimer-drug conjugates for tailored intracellular drug release based on glutathione levels**. *Bioconjugate chemistry* 2008, **19**, (12), 2446-55.

**Figure and Legends**

*Figure 1. scRNA-seq analysis of FOLR2[+] macrophages in endometrium and pelvic exudate cells*

*in endometriosis. (A, B) Lists of human scRNA-seq datasets analyzed and sample size per dataset*

*in the groups of (A) eutopic endometrium with (EuE) and without (Cont) endometriosis, ectopic*

*endometrium (lesion, EcE), and (B) pelvic exudate cells from subjects with (Endo) and without*

*(Cont) endometriosis. (C, D) UMAP plots showing major cell types identified in the endometrium*

*(EM) and pelvic cells (P). (E, F) Violin plot representing marker gene expression for each major*

*cell type identified in endometrial and pelvic cell datasets. (G, H) UMAP projections showing*

*expression patterns of FOLR2 in endometrial and pelvic cells. (I, J) Bar plots showing enriched*

*gene ontology (GO) biological processes in FOLR2[+] macrophages from endometrium and pelvic*

*cells. (K, L) Proportions of endometrial and pelvic FOLR2-expressing cells to total cells within*

*each tissue type. Differences among groups were analyzed with one-way ANOVA followed by the*

*Tukey multiple comparisons test. \*\*P<0.01, \*\*\*P<0.001. (M, N) Dot plot showing representative*

*significantly differentially expressed genes (DEGs) involved in proinflammatory, angiogenesis,*

*and adhesion processes among groups of endometrium and pelvic cells (adjusted p-value <0.05).*

*Figure 2. Preparation and structural elucidation of FA-2DG-D-Cy5 conjugate. (A) Schematic*

*representation for the synthesis of FA-2DG-D-Cy5 using CuAAC reaction. Reagents and*

*conditions: (i) DMSO, DCC, NHS, DIPEA, RT, 12 hours, 70% (ii) DMF, EDC.HCl, DMAP, RT, 12*

*hours, 80% (iii) CuSO$_4$.5H$_2$O, Sodium ascorbate, DMF, DI Water, 60 ºC, 10 hours, 84% (iv)*

*CuSO$_4$.5H$_2$O, Sodium ascorbate, DMF, DI Water, 40 ºC, 6 hours, 88% (B) [1]H NMR spectra*

*showing characteristic proton signals of 2DG-D, 2DG-D-Hexyne, Folic Acid, FA-Azide, FA-2DG-*

*D and FA-2DG-D-Cy5 at each step of the synthesis.*

*Figure 3. In vivo and ex vivo fluorescence imaging and whole-body biodistribution of FA-2DG-D-Cy5. (A) Whole-body in vivo fluorescence images of mice following intravenous (tail vein) administration of FA-2DG-D-Cy5 in sham controls (left) and ELL-induced mice (right), or 2DG-D-Cy5 administered to ELL-induced mice (center) at 1, 6, 24, and 72 hours (n = 3/group/timepoint). (B) Ex vivo fluorescence images of major organs collected at 72 hours post-injection. Organs are arranged from top to bottom as: brain, heart, spleen, lungs, liver, kidneys, and lesion. (C) Quantitative analysis of in vivo fluorescence signals in major organs at 72 hours (n = 3). (D) Heat map showing the mean percentage of injected dose (I.D.) in lesions and major organs at 1, 6, 24, and 72 hours after FA-2DG-D-Cy5 administration in sham controls and ELL-induced mice, or ELL-induced mice treated with 2DG-D-Cy5 (n=3/group). Heatmap was generated with the ComplexHeatmap package of R. (E) Quantitative biodistribution of FA-2DG-D-Cy5 and 2DG-D-Cy5 in lesions and major organs at 72 hours in sham and ELL-induced mice (n = 3). Data for (D) and (E) were obtained by fluorescence spectroscopy of homogenized tissues and are shown as the percentage of the injected dose per organ. Differences in each tissue among groups for data in (C) and (E) were analyzed by one-way ANOVA followed by the Tukey multiple-comparison test. $p < 0.05$ (\*), $p < 0.01$ (\*\*). ELL, endometriosis-like lesion.*

*Figure 4. Cellular uptake of FA-2DG-D-Cy5 and 2DG-D-Cy5 at 2 weeks following intraperitoneal administration. Representative images showing selective uptake of FA-2DG-D-Cy5 or 2DG-D-Cy5 (A) in peritoneal cells, (B) in endometriosis-like lesions (ELL), and (C) kidney, liver, and spleen from ELL-induced mice (n=3). Peritoneal cells were co-stained with FRβ (green) and DAPI (blue). The Cy5 signal (red) was directly acquired using a confocal microscope. White arrows denote Cy5-positive signal in FRβ[+] cells.*

*Figure 5. Preparation and structural elucidation of FA-2DG-D-Niclo conjugate. (A) Schematic representation for the synthesis of (i) Niclo-Azide and (ii) FA-2DG-D-Niclo using CuAAC reaction. Reagents and conditions: (i) DCM, HCl, Dioxane, RT, 3 hours (ii) DCM, TEA, Triphosgene, -25 °C, 1 hour (iii) DCM, TEA, DMAP, RT, 16 hours, 49% (iv) DMF, EDC.HCl, DMAP, RT, 12 hours, 82% (v) CuSO$_4$.5H$_2$O, Sodium ascorbate, DMF, DI Water, 60 °C, 10 hours, 85% (vi) CuSO$_4$.5H$_2$O, Sodium ascorbate, DMF, DI Water, 40 °C, 10 hours, 75% (B) $^1$H NMR spectra showing*

*characteristic proton signals of FA-2DG-D', Compound 10, Niclosamide, Niclo-Azide and FA-*
*2DG-D-Niclo at each step of the synthesis.*

*__Figure 6__. __Characterization of FA-2DG-D-Niclo conjugate and intermediates:__ (__A__) HPLC*
*chromatograms of intermediates at various synthesis steps and percentage purity of FA-2DG-D-*
*Niclo. (__B__) (i) Hydrodynamic diameter, and (ii) Zeta Potential of FA-2DG-D analyzed by dynamic*
*light scattering (DLS) in triplicate. (__C__) (i) Hydrodynamic diameter, and (ii) zeta potential of FA-*
*2DG-D-Niclo analyzed by DLS in triplicate.*

*__Figure 7__. __Characterization of FA-2DG-D-Niclo conjugate and intermediates:__ (__A__) Table showing*
*the physicochemical properties of FA-2DG-D-Niclo and image showing the increase in solubility*
*of niclosamide after conjugation with FA-2DG-D scaffold (FA-2DG-D-Niclo). (__B__) The figure*
*shows the reproducibility in the synthesis of multiple 400 mg-scale batches of FA-2DG-D-Niclo*
*via $^1$H NMR and HPLC. The $^1$H NMR spectra of 3 batches (400 mg scale) show characteristic*
*protons from FA-2DG-D-Niclo. The HPLC chromatographs show reproducibility in the purity of*
*these batches with all these batches showing >99% purity. (__C__) In vitro drug release profile of FA-*
*2DG-D-Niclo conjugate in PBS and 0.1mM GSH, intracellular conditions at 37ºC after 15 days,*
*and in (100%) human plasma after 120 hours.*

*__Figure 8__. __Quantitative analysis of FRβ$^+$ cells in the lesion and peritoneum following dendrimer__*
*__treatments.__ (__A__) Study design as described in Methods. (__B__) Representative immunohistochemical*
*images (left) and quantification of FRβ$^+$ cells (right) in lesions at 14 or 21 days after lesion*
*induction (n=3-8, as dendrimer-Niclo inhibited lesion development). One-way ANOVA followed*
*by the Tukey multiple-comparison test was used to assess differences among groups. (__C__)*
*Representative flow plots (left) and quantification of peritoneal FRβ$^+$ cells (right). Differences in*
*peritoneal FRβ$^+$ cells among groups at each time point were assessed by the Kruskal-Wallis test.*
*Data in all figures are shown as the mean ± SEM (n=8). p < 0.05 (*), or p < 0.01 (**).*

*Figure 9. **Efficacy of FA-2DG-D-Niclo on treating endometriosis.** (A) Quantification of lesion*

*numbers and volumes in mice at Days 14 and 21 after lesion induction. Lesion volumes were*

*calculated as length $\times$ width$^2$. Differences in lesion number and volume among groups at each*

*time point were assessed by the Kruskal-Wallis test. (**B**) Evaluation of endometriosis-associated*

*hyperalgesia. Abdominal (left) and hindpaw (right) withdrawal thresholds were assessed by the*

*von Frey test. Data represent the group mean threshold at the corresponding time point (n=16 at*

*Day -1 and Day 7, n=8 at Day 14 and Day 21). Red arrows show the time of i.p. administration*

*of FA-2DG-D or FA-2DG-D-Niclo. The time course differences within each treatment group*

*were compared using a Kruskal-Wallis test followed by the Dunnett multiple-comparison test.*

*The colored stars or ns (non-significance) represent the differences at the corresponding time*

*point relative to Day -1 within the treatment group. To compare dosing effects of niclosamide at*

*each time point among groups, the Kruskal-Wallis test was used. Data are shown as mean $\pm$*

*SEM (n=8). $p < 0.05$ (\*), $p < 0.01$ (\*\*), $p < 0.001$ (\*\*\*), or $p < 0.0001$ (\*\*\*\*).*

**A. Synthesis of FA-2DG-D-Cy5**

**B. ¹H NMR Characterization of FA-2DG-D-Cy5**

# A. Peritoneal cells

# B. Lesions

# C. Major organs

**A. Synthesis of FA-2DG-D-Niclo**

**(i) Synthesis of Niclo-Azide**

**(ii) Synthesis of FA-2DG-D-Niclo**

**B. ¹H NMR Characterization of FA-2DG-D-Niclo**

**A. HPLC Characterization of FA-2DG-D-Niclo**

2DG-D-Hexyne
@254 nm

FA-2DG-D'
@254 nm

Niclosamide
@331 nm

Niclo-Azide
@331 nm

FA-2DG-D-Niclo
@331 nm
>98% Purity

**B. Size and Zeta Potential of FA-2DG-D**

$6.6 \pm 0.4$ nm

$-5.8 \pm 0.3$ mV

**C. Size and Zeta Potential of FA-2DG-D-Niclo**

$7.6 \pm 0.6$ nm

$-7.7 \pm 0.6$ mV

## A. Physicochemical Properties and Solubility of FA-2DG-D-Niclo

| Physicochemical Properties of FA-2DG-D-Niclo | |
|---|---|
| Hydrodynamic Diameter | 7.6 ± 0.6 nm |
| Zeta Potential | -7.7 ± 0.6 mV |
| HPLC Purity | >98% |
| Drug Loading | ~12% |
| Solubility | 150 mg/mL |

1. FA-2DG-D-Niclo at 18 mg/mL (Niclosamide equivalent concentration)
2. Niclosamide at 1 mg/mL

## B. Reproducible Synthesis of FA-2DG-D-Niclo

$^1$H NMR Spectra (*DMSO-$d_6$*)

HPLC Chromatogram

## C. *In Vitro* Drug Release Profile of FA-2DG-D-Niclo

pH 5.5 + 10mM GSH
pH 7.4 + 0.1mM GSH

Human Plasma Release (Until 5 Days)

**A**

**Study Design**

Von Frey test

Treatment groups (n=16/group):
① FA-2DG-D (Control)
② FA-2DG-D-Niclo 25 mg/kg/b.w.
③ FA-2DG-D-Niclo 50 mg/kg/b.w.

i.p. single dose

Behavioral tests and tissue collection

Von Frey test + Collect: Peritoneal fluid Lesions

Lesion Induction    Treatment    n=8/group/time point

Day -1    Day 0    Day 7    Day 14    Day 21

Age: 12 weeks old    Von Frey test before treatment

**B**

Control    Niclo 25 mg/kg    Niclo 50 mg/kg

Day14    IgG    50 µm

Day21

FRβ+ cells/mm²

○ Control
○ Niclo 25 mg/kg
● Niclo 50 mg/kg

Day 14    Day 21

Days after lesion induction

**C**

Live/CD45+

Control    Niclo 25 mg/kg    Niclo 50 mg/kg

Day14    FRβ+

Day21

FSC-H    FRβ-PE

FRβ+ (cells/uL)

○ Control
○ Niclo 25 mg/kg
● Niclo 50 mg/kg

Day 14    Day 21

Days after lesion induction

**A**

**B**