Structural determinants of broadly neutralizing human antibodies binding to

morphological dengue virus variants.

Ananya Chatterjee[1†], Ankush Roy[1†], Srividhya Srinivasan[1†], Sylvia Charles[1π], Jay

Lubow[3Ψ], Leslie Goo[3τ] and Vidya Mangala Prasad[1,2*]

† equal contribution

Affiliations:

1. Molecular Biophysics Unit, Indian Institute of Science, Karnataka 560012, India.

2. Centre for Infectious Diseases Research, Indian Institute of Science, Karnataka

560012, India.

3. Vaccine and Infectious Disease Division, Fred Hutchinson Cancer Center, Seattle

98109, Washington, USA.

[π]Present address: Department of Chemical Engineering, University of Bath, Bath,

UK.

[Ψ] Present address: ImmunoVec, 570 Westwood Plaza Building 114 ,90095, Los

Angeles, California, USA.

[τ] Present address: Vaccine Company Inc., South San Francisco, California, USA.

*Correspondence to: vmprasad@iisc.ac.in

Abstract:

Dengue virus (DENV) is a global disease threat with tropical regions bearing the bulk

of disease burden. Presence of four DENV serotypes with varying antigenicity has

complicated development of effective, pan-serotype strategies for prevention and

treatment of dengue disease. Here, we report single-particle cryo-electron microscopy

(cryo-EM) structures of DENV in complex with three human-derived broadly

neutralizing antibodies (bnAbs), revealing an antibody class which preferentially uses

its heavy chain for potency and antigen recognition. These heavy chain-driven E-dimer

recognizing (HEDR) bnAbs potently neutralize all DENV serotypes by binding to a

quaternary epitope encompassing the critical fusion loop and conserved glycans on

DENV's surface E-glycoprotein dimer. We also demonstrate the presence of capsule-

shaped DENV, with HEDR antibodies binding such tubular morphologies in addition

to spherical virions. Cryo-EM helical reconstructions of Fab-bound tubular virions from

different DENV strains establishes that the pattern of E-glycoprotein's helical

arrangement is dependent solely on the virus strain. The structural data also

demonstrate that binding of highly potent HEDR bnAb D14.F25.S02 causes distortion

of DENV particles. Collectively, these results elucidate key features of HEDR bnAbs

while illustrating the importance of eliciting immune responses that can neutralize

diverse DENV morphologies.

Introduction:

DENV is an enveloped RNA virus which affects nearly 400 million individuals worldwide giving rise to diverse clinical outcomes, including asymptomatic dengue fever, joint pain, rashes, dengue shock syndrome and severe dengue haemorrhagic fever[1]. The canonical mature and infectious DENV is an icosahedral particle, 500 Å in diameter, with its outer membrane surface coated by its main antigenic protein, the envelope (E) glycoprotein (Fig.1a,b). Each E-protein monomer (~50kDa molecular weight) consists of three domains—DI, DII, and DIII—with N-linked glycosylation at Asn67 and Asn153 amino acid residues[2,3] (Fig.1c). The E-protein's DII domain contains the highly conserved fusion loop (~14 amino acids long) at its distal end, which is crucial for viral membrane fusion within host cells during infection[3–5] (Fig.1c). On the mature virion surface, the E-proteins are arranged into 90 head-tail homodimers with three such E-dimers forming an E-raft that comprises two asymmetric units of the virus's icosahedral surface arrangement[2] (Fig.1b,c). Despite global icosahedral symmetry, the local environment for each of the E-proteins in the asymmetric unit is different, and consequently, the local environment experienced by the E-proteins are distinct at the icosahedral 5-fold (5f), 3-fold (3f) and 2-fold (2f) symmetry positions[2] (Fig.1b,c).

DENV presents almost identical surface epitopes across its four serotypes [6–8], giving rise to cross-reactive antibodies that are, often, poorly-neutralizing against heterologous DENV strains[9,10]. Secondary infections of DENV can, thus, cause severe disease due to antibody-dependent enhancement (ADE) of viral infection [11,12]. There are few reported examples of broadly neutralizing antibodies (bnAbs) that neutralize all DENV serotypes[10–12]. All these bnAbs bind on or near the conserved fusion loop region of DENV's E-protein, highlighting the significance of this region as an

immunogenic target. Notably, largest breadth in neutralization activity across DENV

serotypes has been observed in antibodies that recognize a combination of the fusion

loop along with adjacent regions on a E-protein dimer[13–21]. From previously reported

literature, few of the best characterized bnAbs which bind a quaternary epitope

comprising the fusion loop and adjacent regions on the DENV E-protein are the E-

dimer epitope (EDE) bnAbs, which are further sub-classified as EDE1 and EDE2

bnAbs[19,20]. EDE1 bnAbs strongly neutralize both DENV and Zika viruses with bnAb

binding being independent of N153 glycosylation on DENV E-protein. In contrast, the

EDE2 bnAbs potently neutralize DENV, with lower activity against ZIKV, and require

interaction with E-protein's N153 glycan.

DENV populations in nature are a mixture of its mature and immature forms,

including virions that are partially mature to various proportions[6,22,23]. The mature and

immature DENV structures are markedly different, with the mature DENV surface

displaying 90 non-covalently bound, antiparallel E-dimers along with buried membrane

(M) proteins on its surface, whereas the immature DENV displays 60 trimeric spikes

of E-protein and precursor membrane (prM) protein heterodimers. Thus, incomplete

virus maturation adds a layer of structural heterogeneity to circulating DENV

populations. Additionally, in the past decade, DENV virions have been reported to

deviate from their icosahedral structures to exhibit other morphologies such as the

expanded/bumpy virus[24,25] and club-shaped tubular virions[26] at temperatures of 37° C

and above. These heterogeneities in DENV structural forms exhibit variable

proportions of distinct quaternary surface arrangements that are not yet fully

understood, with little data available on the ability of bnAbs to recognize these various

morphologies.

We describe here, cryo-EM structures of three potent, broadly neutralizing

human antibodies, J9, D14.F25.S02 and D14.F05.S03, in complex with the complete,

infectious DENV virion. All three bnAbs were isolated from patients that recovered

from secondary DENV infection[16,18]. J9 and D14.F05.S03 neutralize only DENV

whereas D14.F25.S02 neutralizes DENV as well as its relative, Zika virus (ZIKV).

These select bnAbs show strong neutralization activity against DENV which is

comparable, and in some instances, better than other known bnAbs[16,18]. The

structures described here demonstrate that these bnAbs form a unique class of potent

antibodies that bind to the quaternary E-dimer epitope encompassing the fusion loop

and adjacent regions of the E-proteins, with distinct features that differentiate these

from previously described bnAbs. We also report helical cryo-EM structures of tubular

DENV morphologies bound to these bnAbs, showing the capability of these bnAbs to

bind variant DENV morphologies. Furthermore, using different DENV strains, we

demonstrate that E-protein arrangement on the helical DENV-bnAb structures is

dictated by the virus strain and not the bound antibody.

Results:

Cryo-EM structures of spherical DENV particles bound to bnAbs J9,

D14.F25.S02 and D14.F05.S03:

In this study, we have used two DENV-2 serotype strains from different geographical

origins: THSTI/TRC/01 (India) and US/BID-V594/2006 (Puerto Rico). Both these

strains have been isolated from humans, and the respective E-glycoproteins share

98% identity (99% similarity) between each other and ~98% identity (~99.6%

similarity) with the E-protein of well-studied DENV-2/16881 strain (Supplementary Fig.

1). Using these strains, cryo-EM structures of spherical DENV complexed with Fabs

from three potent bnAbs, namely, J9, D14.F25.S02 and D14.F05.S03 were calculated

with icosahedral symmetry applied (Supplementary Fig.2-4). The icosahedral cryo-EM

structures of the DENV+J9 Fab, DENV+ D14.F25.S02 Fab and DENV+D14.F05.S03

Fab were determined to highest global resolutions of 3.85 Å, 4.1 Å and 4.5 Å

respectively (Fig1c-e, Supplementary Fig.5 and Supplementary Table 1-3). Local

resolution of the cryo-EM density maps, calculated from the viral membrane outwards,

ranges between 2.4 Å to 6.4 Å for DENV+J9 Fab complex, 2.5 Å to 7.5 Å for

DENV+D14.F25.S02 Fab complex, and 3.5 Å to 7.0 Å for DENV+D14.F05.S03 Fab

complex respectively (Supplementary Fig.5). Localised reconstruction of sub-

icosahedral symmetry-related positions further improved local map features and

glycan densities for both the DENV+J9 as well as DENV+D14.F05.S03 complex maps

(Supplementary Fig.6, 7).

In the above described cryo-EM maps of icosahedral DENV complexes,

densities for the E and M glycoproteins as well as the antibody-virus interface were

well-resolved with amino acid side-chains distinguishable in most regions of the virus

surface, permitting real-space refinement of atomic models of DENV E-protein and the

respective Fab variable regions (Fig.2a,d,g, Supplementary Fig.8, Supplementary

Tables 1-3). Though one Fab (~50kDa molecular weight) can bind to each E-protein

for all three bnAbs, none of the bnAbs show full occupancy of all the possible 180 E-

protein binding sites on the DENV virion, with each bnAb showing a different pattern

of binding across the virion (Fig.1d-i) suggesting that complete occupancy of all

available epitopes may not be necessary for neutralization activity of these bnAbs.

Figure 1. Cryo-EM structures of HEDR bnAbs complexed with DENV.

a. Surface and cross-sectional view of dengue virus with black arrows indicating the different virus parts: E (envelope glycoprotein layer), OM (Outer lipid membrane), TM (Transmembrane space), IM (inner lipid membrane) and G (genome core).

b. Surface representation of DENV virion showing icosahedral arrangement of E-
proteins in grey. Black triangles indicate the icosahedral asymmetric unit. A single
E- raft comprising of three E-dimers is coloured in beige and marked by a rhombus.
Each asymmetric unit repeats itself 60 times to form the spherical virus surface.
Despite the global icosahedral symmetry, the local environment around each E-
protein in the asymmetric unit is distinct and consequently, the local environment
experienced by the E-proteins at the 5-fold, 3-fold and 2-fold vertices are different,
which introduces an element of non-symmetry in virus-antibody interactions at
these vertices.

c. Cartoon representation of an E-dimer raft, also referred to as an E-raft, consisting
of three adjacent E-dimers. Parts of the E-protein are colored as: Domain-1 (DI) in
mustard yellow, Domain-2 (DII) in brown with the fusion loop (FL) in green and
Domain-3 (DIII) in black.

d-f. Radially coloured, high-resolution cryo-EM reconstructions of DENV complexed
with J9 (d), D14.F25.S02 (e) and D14.F05.S03 (f) Fab fragments.

g-i. Surface model representation of the cryo-EM reconstructions shown in d-f panels.
The E-proteins are colored as grey surface similar to panel b, The variable region of
the Fab fragments is represented also as surface models. **g.** DENV complexed with
J9 Fab (Orange- heavy chain, Yellow- light chain). **h**. DENV complexed with D14.F25.S02
Fab (Dark blue- heavy chain, Light blue- light chain). **i.** DENV complexed with
D14.F05.S03 fab (Dark pink- heavy chain, Light pink- light chain).

In all panels, the 5-fold, 3-fold and 2-fold icosahedral symmetry vertices are indicated
by a hexagon, triangle and ellipse, respectively.

Epitope features of bnAbs J9, D14.F25.S02 and D14.F05.S03 on icosahedral

DENV:

All three bnAbs bind to the fusion loop region and interact with the N153 glycan, with the Fab footprints covering parts of DI and DIII domains from the adjacent E-protein in an E-dimer (Fig.2, Fig.3a and Supplementary Fig.9, 10). The bnAbs require interactions that span the E-dimer surface (Fig.2b,e,h, Fig.3a, Supplementary Tables 4-6), which explains why these were previously reported as not binding E-monomers in solution[16,18]. All three bnAbs show overlapping epitope footprints, despite differences in their angle of approach to the E-protein (Fig.3, Supplementary Fig.10). All three bnAbs bind to the E-protein's fusion loop using its heavy chain CDR3 (HCDR3) loop (Fig.2a,d,g, Fig.3 a,b and Supplementary Figure 9). The tip of J9's HCDR3 is positioned towards the intra-E-dimer cleft, whereas the D14.F05.S03's HCDR3 loop is positioned on top of the fusion loop (Fig.3b). Notably, in D14.F25.S02, its HCDR3 loop interacts in a sideway fashion with the E-protein's fusion loop, such that the HCDR3 is inserted within the inter-dimer space between adjacent E-dimers (Fig.3b). Furthermore, the HDCR3 from all three bnAbs also make contacts with the conserved N153 glycan on E-protein (Fig.2c,f,i).

The overall epitope footprint of the three bnAbs (Fig.2b,e,h and Fig.3a) overlaps broadly with previously described EDE bnAbs[20]. However, the clusters of E-protein amino acid residues that are critical for bnAb binding and neutralization, as determined from these structural data and corroborating functional assays[16,18], are distinct between the different bnAbs, indicating local variations in binding determinants (Supplementary Fig.9). Furthermore, all three bnAbs make majority of interactions with

the DENV E-protein using its heavy chain (Supplementary Tables 4-7), with the bnAb

heavy chain involved in interactions with both the fusion loop (Fig.2a,d,g and 3b) and

conserved N153 glycan (Fig.2c,f,i), grouping them as a separate class of their own,

hereon referred to as heavy chain-driven E-dimer recognizing (HEDR) antibodies.

Interactions with E-protein glycans by the HEDR bnAbs, J9, D14.F25.S02 and

D14.F05.S03:

The HEDR bnAbs can be classified into two categories, similar to the EDE

bnAbs[19,20], based on their dependence on E-protein's N153 glycan for neutralization

activity. D14.F25.S02 forms HEDR category-1 which is independent of N153

glycosylation for its neutralization activity. Nevertheless, D14.F25.S02 bnAb makes

multiple contacts with the N153 glycan (Fig2f, Supplementary Table 5), in contrast to

the previously described EDE1 bnAbs which induce disordering of the N150 loop

containing the N153 glycan[15,20]. In further contrast to EDE1 bnAbs[20], HEDR category-

1 bnAb, D14.F25.S02, does not show any interactions with the E-protein's N67 glycan

(Fig. 2f). On the other hand, HEDR category-2 bnAbs include J9 and D14.F05.S03,

show many similarities with the EDE2 bnAbs, including the requirement of N153

glycosylation for its neutralization activity[16,19]. J9 and D14.F05.S03 make close

contacts to the N153 glycan making it an integral part of the bnAb epitope (Fig.2c,i,

Supplementary Table 4, 6). J9 uses its heavy chain to interact with the lower part of

the N153 glycan via hydrophobic and electrostatic interactions, whereas the light chain

contacts the terminal glycan residues via primarily electrostatic interactions (Fig.2c,

Supplementary Table 4). Similar pattern of engagement is seen with D14.F05.S03

also with the Fab heavy chain making multiple contacts to the lower part of the N153

glycan (Fig.2i, Supplementary Table 6). J9 and D14.F05.S03 also make close contacts

with the E-protein's other conserved glycan at N67 position (Fig2c,i, Supplementary

Tables 4, 6), though this glycan is not necessary for its neutralization activity. As the

spatial position of N153 glycan in DENV's E-protein is different with respect to the

equivalent N154 glycan position in ZIKV's E-protein, the structural data suggests that

close interactions with N153 glycan may influence differential recognition of DENV by

the HEDR category-2 bnAbs, J9 and D14.F05.S03.

Figure 2. Quaternary binding epitopes of HEDR bnAbs on DENV E-dimer.

a,d,g. Density map cutouts of DENV E-dimer and Fab complexes from the

icosahedral cryo-EM structures shown in Figure 1d-f. The maps cutouts also include

the built-in atomic models for each of the complexes (DENV-E+M protein and Fab

variable regions) showing the accuracy of the protein models. In all panels of this

figure, the E-dimer ribbon models are shown in dark grey. The ribbon models of each Fab variable region (Fv) are also shown, with J9 Fv colored in orange (heavy chain) and yellow (light chain), D14.F25.S02 colored in dark blue (heavy chain) and cyan (light chain), D14.F05.S03 colored in dark pink (heavy chain) and light pink (light chain) and fusion loop region in green.

b,e,h. Top and side view of atomic models of E-dimer ectodomain complexed with Fab variable regions of J9 (b), D14.F25.S02 (c) and D14.F05.S03 (e) respectively. In all three atomic model representations, the glycan positions are marked as N153 and N67.

c,f,i. Interactions of bnAbs J9 (h), D14.F25.S02 (i) and D14.F05.S03 (j) with conserved glycans, N153 (left) and N67 (right) on E-protein. Black dotted lines indicate the closest interacting residues. Glycan units are numbered 1 to 5 starting from the closest unit to the Asn side chain.

a Overlap of Fab variable region map densities of J9, D14.F25.S02 and D14.F05.S03 on E-protein dimer

Fab axes

10 Å

Fab footprints on E-dimer

10Å

N67 glycan   N153 glycan

J9
D14.F25.S02
D14.F05.S03
E protein
Fusion Loop

10 Å

b Overlap of atomic models of J9, D14.F25.S02 and D14.F05.S03 Fab variable regions on E-protein monomer

10 Å

2Å

Figure 3. Overlap of bnAb binding locations and epitopes.

a. Overlapping location of the map density volumes of all three Fab variable regions

(J9, D14.F25.S02 and D14.F05.S03) on E+M-dimer. Zoomed-in panel shows the top

view of bnAb footprints on one of the E-dimer binding positions. The bnAb footprint

outlines are colored as J9: yellow, D14.F25.S02: blue and D14.F05.S03: magenta.

b. Location of overlapping binding positions of all three Fabs on E-monomer. Top

panels show the Fabs as density volumes and E-monomer as a ribbon model built-in

to its electron density. Bottom panels show the Fabs and E-monomer as ribbon

models. Left panels show the side view and the right panels show the top view. Color

scheme is the same as panel a. Zoomed-in top view shows the HCDR3 loop

positions of J9 (yellow), D14.F25.S02 fab (blue) and D14.F05.S03 fab (magenta)

with respect to the E-protein's fusion loop (green).

Morphological features of tubular and capsule-shaped DENV present in virus

preparations:

Virus preparations used in this study were produced by infection of mosquito

cells and incubated with Fabs at room temperature without any treatment at higher

temperatures. Nonetheless, DENV particles with tube-like morphologies and bound

Fab were observed as a small percentage (<15%) in all the cryo-EM datasets collected

for both the US/BID-V594/2006 as well as THSTI/TRC/01 strains. Individual virions

were captured from both DENV strains with different lengths of tube-like extensions

connected to a spherical head (Supplementary Fig.13-14), reminiscent of the club-

shaped DENV morphology described previously[26]. Furthermore, in case of the

US/BID-V594/2006 strain, completely tubular virions similar to cylindrical capsules

were also observed (Fig.4a,b). The capsule-shaped DENV particles have similar

widths as the tube-like extensions observed developing from spherical virions of the

same strain, suggesting that capsule-shaped virions form via alterations to the

spherical virus (Supplementary Fig. 11). The capsule-shaped virions have an inner

diameter of 110 Å and 120 Å in the D14.F25.S02- and J9-bound datasets respectively,

with average length of particles being 79nm and 74nm respectively (Fig.4a,b). Thus,

the capsule-shaped virions enclose a volume large enough to contain 1.9 – 2.2 copies

of the viral genome (see Methods for details), indicating that these morphological

variants can accommodate the viral genome in its entirety.

Cryo-EM structures of tubular DENV2-US/BID-V594/2006 virions bound to

D14.F25.S02 and J9 Fab fragments:

Cryo-EM reconstruction of the tubular regions of DENV2-US/BID-

V594/2006+D14.F25.S02 Fab complex using helical symmetry parameters achieved

a global resolution of 14.7 Å (Fig.4c; Supplementary Fig.11a-f, 12a). In this structure,

E-dimers are arranged one next to another with a gentle rise, wrapping tightly around

the membrane tube (Fig.4c-e, Supplementary Fig.11 a-e), with two Fabs bound per E-

dimer, in similar positions as in the icosahedral structures. Overlapping three adjacent

E-dimer atomic models from the helical structure with the icosahedral E-raft structure

shows good agreement with an r.m.s.d value of 1.04 Å, confirming that the E-raft

contacts seen in spherical virions are preserved in the helical structure of DENV2-

US/BID-V594/2006+D14.F25.S02 Fab complex (Fig.4e,f). This helical arrangement of

E-proteins resembles the E-protein arrangement previously reported in the helical

structure of ZIKV, a close relative of DENV, complexed with an EDE1 antibody[26].

Similar helical patterns were observed in the J9+US/BID-V594/2006 dataset also, but

only a very low resolution helical cryo-EM reconstruction was determined due to few

tubular particle numbers (Fig.4g; Supplementary Fig.11g-i, 12b). Rigid body fitting of

the E-dimer and Fab complex model into the J9+DENV2-US/BID-V594/2006 map

(Fig.4h, Supplementary Fig. 11i) showed that the E-dimer arrangement is similar to

that observed with the D14.F25.S02+DENV2-US/BID-V594/2006 helical

reconstruction (Fig.4e), albeit with a rotation of 26º along an axis perpendicular to the

plane of the E-dimer.

Figure 4. Capsule-shaped DENV show helical symmetry arrangement in Fab-bound state.

a-b. Representative raw cryo-EM images of capsule-shaped DENV2-US/BID-V594/2006 virions bound with D14.F25-S02 Fab (a) and J9 Fab (b). Spiky Fab projections perpendicular to the smooth virus surface is observed with presence of ordered arrangement of proteins visible.

c. Map representation of surface (left) and longitudinal cut-out (centre) and top view (right) of the helical reconstruction of D14.F25.S02 Fab complexed with DENV2-

US/BID-V594/2006. M, E and F represent the membrane, E-protein and Fab density

regions, respectively, in the density map.

d. Surface representation of fitted models of the E-dimer and D14.F25.S02 Fab

variable region into the Fab-bound DENV2-US/BID-V594/2006 helical structure.

e. Ribbon diagram of adjacent E-dimer positions in the D14.F25.S02 + DENV2-

US/BID-V594/2006 helical structure.

f. Overlap of three adjacent E-dimer atomic models from the D14.F25.S02 + DENV2-

US/BID-V594/2006 helical structure (purple) to an E-raft from the DENV2-US/BID-

V594/2006 icosahedral structure (green), showing similarity in E-dimer arrangement

and an r.m.s.d value of 1.04Å.

g. Map representation of surface (left) and longitudinal cut-out (centre) and top view

(right) of the helical reconstruction of J9 Fab complexed with DENV2-US/BID-

V594/2006. M, E and F represent the membrane, E-protein and Fab density regions,

respectively, in the density map.

h. Surface representation of fitted of the E-dimer and J9 Fab variable region into the

Fab-bound DENV2-US/BID-V594/2006 helical structure.

i. Ribbon diagram of adjacent E-dimer positions in the J9 + DENV2-US/BID-

V594/2006 helical structure. This E-dimer arrangement is similar to that seen in the

D14.F25.S02+DENV2-US/BID-V594/2006 helical structure after rotation of 26º along

an axis perpendicular to the plane of the E-dimer. R.m.s.d between the two E-dimer

arrangements in panel e and i (after superposition with rotation) is 0.875Å.

Cryo-EM structures of tubular DENV2-THSTI/TRC/01 virions bound to J9, D14.F05.S03 and D14.F25.S02 Fab fragments:

For the DENV2-THSTI/TRC/01 strain, Fab-bound particles with a spherical head and narrow tube-like extension were observed (Supplementary Fig.13b,f, 14a). These tubular extensions were selected and subjected to unsupervised 2D classification, which also showed clear helical patterns (Supplementary Fig.13b,g 14b). Helical cryo-EM reconstructions of THSTI/TRC/01 virions were achieved to global resolutions of 14.41Å for J9-bound, 17.3Å for D14.F05.S03-bound and 17.37 Å for the D14.F25.S02-bound particles respectively (Fig.5, Supplementary Fig. 13, 14, 15). The outer and inner diameters of THSTI/TRC/01 tubular particles were 312 Å and 20 Å for the J9-virus complex, 280 Å and 55 Å for D14.F05.S03-virus complex, and 320 Å and 20 Å for the D14.F25.S02-virus complex respectively (Fig.5). The narrow inner diameters of the tubes in the THSTI/TRC/01 strains cannot accommodate the viral genome, indicating that the viral RNA is contained in the spherical head regions only. The narrow inner tube diameter could, thus, restrict these virions from transitioning to complete cylinders or capsule-shaped particles.

Despite the differences in tube diameters observed in all three helical THSTI/TRC/01 structures, the E-dimers adopt a similar steep step-like arrangement of E-dimer around the membrane tube (Fig.5), which differs remarkably from the E-dimer arrangement observed in the helical US/BID-V594/2006 structures (Fig.3). In both the J9- and D14.F05.S03-bound helical structures from THSTI/TRC/01, two Fabs are bound per E-dimer with the external tips of the Fab densities joined to each other. Rigid body fitting of the Fab atomic models into the map density indicates that the

constant regions of the Fabs would clash with each other in this conformation (Fig.5a,b,

Supplementary Figs. 13d,e, 14e-g). However, no such clashes were observed in the

respective icosahedral structural models (Fig.1d,f, Supplementary Fig. 13e,14g). In

spherical DENV virions, the E-protein conformation has a slight curve to match the

curvature of the virus. Modelling the Fab structures to a flat conformation of the E-

proteins, generally observed in purified E-protein structures[20,27], shows clashing of Fab

constant domains similar to that observed in the helical structures of the DENV-

THSTI/TRC/01+ Fab complexes (Supplementary Fig.13d,e, 14e-g). These analyses

suggest that the E-dimer adopts a flatter conformation in the tubular virions when

compared to its spherical counterpart. In case of D14.F25.S02, only one Fab density

is observed per E-dimer in the DENV-THSTI/TRC/01 tubular reconstruction. Fitting the

atomic models of the E-dimers and D14.F25.S02 Fab variable region in the cryo-EM

map suggests that the close juxtaposition of the tips of adjacent E-dimers in the helical

structure restricts binding of a second Fab due to probable clashes (Fig.5c,

Supplementary Fig.13f-k).

Figure 5: Helical reconstruction of DENV2/THSTI/TRC/01 complexed with J9,

D14.F05.S03 and D14.F25.S02 Fab fragments. Side view, central cut-out, top view

and surface rendered helical density map of DENV2/THSTI/TRC/01 + J9 Fab **(a)**,

DENV2/THSTI/TRC/01 + D14.F05.S03 Fab **(b)**, and DENV2/THSTI/TRC/01 +

D14.F25.S02 Fab **(c)** complexes. Each helical density map is colored based on its

radius from the central helical axis. In the surface rendered view (right panels of a-c),

only the Fab variable region is shown. (d) Comparison of adjacent E-dimer positions

in all three Fab-bound THSTI/TRC/01 helical structures showing similar E-dimer

arrangements. This similarity in adjacent E-dimer arrangement is reflected in the

similar helical parameters obtained for the three helical structures of

DENV2/THSTI/TRC/01 complexed with J9 (Rise=15.19 Å, Twist=80.62°),

D14.F05.S03 (Rise=15.01 Å, Twist=79.94°) and D14.F25.S02 (Rise=15.13 Å,

Twist=80.09°).

Structural features of D14.F25.S02 Fab and IgG binding on spherical DENV and

its implications:

Three-dimensional (3D) classification of spherical virions from the DENV2-US/BID-

V594/2006 + D14.F25.S02 Fab complex dataset resulted in two distinct classes of

bound virions. The 4.1Å resolution cryo-EM structure was achieved with the larger

class of particles which contained Fab bound to only the icosahedral 5-fold (5f) and 3-

fold (3f) symmetry positions with negligible density observed at the icosahedral 2-fold

axes (2f) (hereafter referred to as 5f-3f map) (Fig.1e). The Fab occupancy is very low

in the cryo-EM map, with 5f and 3f Fab occupancy estimated to be approximately 25

% and 18 % respectively (calculated as percentage of the average map value of

Fab/E-protein[28]). Moreover, though the Fab positions are similar at both 5f and 3f

positions, only the Fab density at 5f position was defined enough to carry out real

space refinement of the Fab variable chain region.

The smaller class contained particles where the D14.F25.S02 Fab was bound

only at the icosahedral 3-fold (3f) and 2-fold positions (2f) (hereafter referred to as 2f-

3f map) (Fig.6a). Refinement of the 2f-3f class, with icosahedral symmetry imposed,

resulted in a cryo-EM map of 8.45 Å global resolution (Fig5a, Supplementary

Fig.16a,b). The Fab occupancy in the 2f-3f map was 39%, with both 3f and 2f positions

bound equivalently. However, only 20% of particles formed the 2f-3f class compared

to the 5f-3f class, indicating that the Fab binds to the 5f position robustly. The Fab

footprint and angle of approach are equivalent in both 2f-3f and 5f-3f maps (Fig.1e,

2c, Fig.7a, Supplementary Fig. 16c). Though there is no evidence of potential clashes,

comparing the Fab structural models bound to symmetry-related positions on the virion

suggest that binding of D14.F25.S02 to 5f position affects Fab binding to the 2f

positions, as is seen in the negligible Fab occupancy at 2f positions in the 5f-3f map.

Attempts to perform localized reconstruction of the Fab-bound E-dimers at the 5f and

3f symmetry positions did not yield any 3D classes with better Fab densities, which

could be due to the low Fab occupancy. Analysis of 2D classes shows that saturation

level binding of D14.F25.S02 bnAb causes distortion of the spherical virions

(Supplementary Fig.16d). Spherical virions are only observed in 2D classes that

contain feeble Fab projections suggesting lower binding ratio of Fab molecules to

virions (Supplementary Fig.16d).

Cryo-EM analyses of DENV2-US/BID-V594/2006 complexed to D14.F25.S02

IgG was also attempted to get a broader sense of bnAb epitope binding

(Supplementary Fig.17a). Massive aggregation of virions was observed in the dataset

with few isolated virions (Supplementary Fig.17b). Single-particle cryo-EM

reconstruction was calculated from a small subset of non-aggregated and loosely

aggregated virions to a global resolution of 7.7 Å with a local resolution range of 4.5

to 10 Å (Fig.6b, Supplementary Fig.17a,c,d). The IgG binding positions on the virion

are identical to that observed with the Fab (Figure 1,2 and Supplementary Fig.16), and

the bnAb binding occupancy at all symmetry-related epitope sites was approximately

26%. 3D classification did not yield any differentially bound IgG subsets as observed

with the Fab-bound dataset, which could be due to the lower number of isolated

particles in the IgG dataset. The IgG map density could be resolved only till the Fab

region, and no density was observed for the Fc portion of the IgG, even at very low

thresholds (Supplementary Fig.17f). This observation along with massive aggregation

seen in the dataset suggests that the two arms of the IgG cross-link adjacent virions.

The overall low occupancy of Fab arms in the averaged IgG+DENV complex

reconstruction further reinforces that the D14.F25.S02 binding is relatively weak

across different epitope positions on the virus.

Despite several attempts, it was not possible to achieve a high resolution cryo-

EM structure of D14.F25.S02 Fab complexed to the DENV2-THSTI/TRC/01 strain.

The best possible 3D reconstructed map of D14.F25.S02 Fab+DENV2-THSTI/TRC/01

complex achieved a global resolution of ~19 Å showing Fabs bound equivalently at all

symmetry related positions in the cryo-EM map (Fig.6c). However, the Fab and E-

protein features lack definition, suggesting higher variability in the constituent virions

(Fig.6c, Supplementary Fig.18). Analysis of 2D classes from the DENV2-

THSTI/TRC/01+ D14.F25.S02 Fab dataset showed clear distortion of Fab-bound

virion particles, indicating that the virus morphology was dramatically affected by

D14.F25.S02 Fab binding (Supplementary Fig.18b).

446   Analysing the binding epitope of D14.F25.S02 shows that the bnAb contacts

span both intra-E-dimer and inter-E-dimer regions (Fig.2c, 5d,e and Supplementary

Fig.9). Majority of the Fab interactions within an E-dimer is carried out by the Fab

heavy chain through contacts to DI and DII domains, with the Fab light chain making

minimal contacts with DI and DIII domains (Supplementary Fig.9 and Supplementary

Table 5). However, the Fab light chain is solely involved in inter-E-dimer contacts

between adjacent E-rafts and also within an E-raft (Fig.6d,e). At the icosahedral 5f

position, the Fab light chain has a different local environment when compared to the

2f and 3f positions and makes numerous contacts to the DI and DIII domains of E-

proteins in adjacent E-dimer rafts (Fig. 5d, Supplemental Table 2). E-protein residues,

Tyr 178 and Met 297, which have been implicated in neutralization activity of

D14.F25.S02[16] are located within bnAb footprint at the 5f position, indicating that these

inter-raft contacts are essential for the bnAb activity (Fig.6d). In contrast,

D14.F25.S02's light chain interactions with adjacent E-dimers within an E-raft are

minimal (Fig.6e, Supplementary Table 5) and not necessary for its binding or

neutralization activity[16].

Figure 6. **Cryo-EM reconstructions of D14.F25.S02 Fab and IgG bound to DENV strains.**

a. Surface rendered 2f-3f cryo-EM density map of D14.F25.S02 Fab + DENV2-US/BID-V594/2006 complex.

b. Surface rendered cryo-EM density map of D14.F25.S02 IgG + DENV2-US/BID-V594/2006 complex.

c. Surface rendered cryo-EM density map of D14.F25.S02 Fab + DENV2-THSTI/TRC/01 complex. Panels a-c are colored radially from the centre and icosahedral asymmetric unit labelled as a dotted triangle.

d. Top view of icosahedral 5f position from D14.F25.S02 Fab + DENV2-US/BID-

V594/2006 complex showing inter-E-raft interactions made by D14.F25.S02 light

chain. E-proteins are shown as ribbons and Fabs are shown as surface

representations. Zoomed-in panels show top view and side view of inter-E-raft

interaction regions. Amino acid residues important for neutralization have been

depicted as green-colored balls. In the zoomed-in top view, cyan outline indicates the

Fab footprint. In the side-view, contacts between the Fab light chain and E-protein

residues have been labelled as dotted lines, (i) S96->P384 (ii) S26->E174 (iii) G31->

E174 (iv) G30-> E174 (v) K33-> G179 (vi) S69-> T176 (vii) G70->T176 respectively.

Residue contacts are calculated as residues with <8Å distance between Cα-Cα

backbone of Fab and E-protein respectively.

e. Top view of inter-E-dimer interactions within an E-raft from the D14.F25.S02 Fab +

DENV2-US/BID-V594/2006 complex structure showing inter-E-dimer interactions

made by Fab light chain at 3f and 2f positions.

Structural features of J9 and D14.F05.S03 Fab bound to spherical DENV strains

and its implications:

The highest resolution, 3.85 Å cryo-EM structure of J9 Fab+DENV was

achieved in complex with the DENV2-US/BID-V594/2006 strain (Fig.1d,g, 2b). In case

of D14.F05.S03, structure determination to a global resolution of 4.5 Å was achieved

in complex with the DENV2-THSTI/TRC/01 strain (Fig.1f,i, 2d). Additionally, a 7.2 Å

cryo-EM structure of J9 Fab complexed with DENV2-THSTI/TRC/01 was also

determined, which showed clear density for the J9 Fab and identical binding features

as seen with the DENV2-US/BID-V594/2006 strain (Supplementary Fig.19). These

results demonstrate that the DENV2-THSTI/TRC/01 strain consists of homogenous

mature particles suitable for single-particle cryo-EM analyses. These results also

highlight that it is the binding of D14.F25.S02 Fab to DENV2 and not DENV2-strain

dependent properties which impeded higher resolution cryo-EM structure

determination of D14.F25.S02 Fab-bound DENV2-THSTI/TRC/01 sample.

In comparison to the D14.F25.S02 bound spherical DENV structures, the

occupancy of bound Fab was much higher in the case of D14.F05.S03 and J9.

D14.F05.S03 showed high occupancy of Fabs at its 3f and 2f positions (82%) with

relatively lower occupancy (50%) at the 5f positions (Fig.7a) (calculated as percentage

of the average map value of Fab/E-protein[28]). In case of J9, the fab bound equivalently

to the 5f and 3f positions (~60%) with lower occupancy at the 2f position (~35%)

(Fig.1d, Fig.7b). Analyses of fab interactions using the real-space refined atomic

models show that both D14.F05.S03 and J9 make minimal inter-E-dimer contacts. The

Fab light chains of both bnAbs making potential electrostatic interactions with Thr226

and Gln227 residues from an adjacent E-dimer within an E-raft (Fig.7a,b). However,

as in the case of D14.D25.S02, these intra-E-raft, inter-E-dimer contact residues are

not essential for Fab binding[16]. At the 5f position, the inter-E-dimer contacts change

for both J9 and D14.F05.S03. For D14.F05.S03, the orientation of the Fab atomic

model at 5f results in clashes with the atomic model of the E-protein from an adjacent

E-raft due to close juxtaposition of the protein chains (Fig.7a, Supplementary Table

6). These potential clashes of D14.F05.S03's light chain at the 5f position combined

with lack of inter-Fab stacking as seen at the 2f and 3f positions, could cause higher

variability in the Fab binding at the 5f vertices, which in turn may account for the lower

occupancy of D14.F05.S03 at 5f vertices as seen in the icosahedral-averaged cryo-

EM map (Fig.1f). For J9, the Fab light chain at 5f interacts with E-protein from an

adjacent E-dimer raft maintaining electrostatic interactions (Fig.7b). The lower J9

occupancy at 2f position could be due steric hindrance caused by J9 binding to

adjacent 5f and 3f positions in an E- raft (Fig.7b).

Figure 7. D14.F05.S03 and J9 Fab arrangement on DENV E-raft. **a.** Low threshold,

surface map rendering of D14.F05.S03 + DENV2-US/BID-V594/2006 map showing

higher Fab occupancy near 2-fold and 3-fold (3f) positions compared to the 5-fold (5f).

Based on the atomic models of the Fab and E-protein built into the electron density,

potential clashes seem likely between the Fab light chain (L1-S26) and E-protein (E2'-

P384) at the 5f position, with unrealistic inter-atomic distances of 2.01 Å between the

protein backbones. In contrast, at the 3f position, the closest distance between Fab

light chain (S26) and E-protein (E2-T266) is 7.51 Å. **b.** Low threshold, surface map

rendering of J9 + DENV2-US/BID-V594/2006 map showing higher Fab occupancy

near 5-fold (5f) and 3-fold (3f) positions compared to the 2-fold. Zoomed-in insets in

both panels show interaction of Fab light chain CDR1 (L1) with E-protein residues.

Based on the atomic model built in the electron density map, at the 5f position, the

closest distance between atoms in the J9 fab light chain (L1-S26) and E protein (E2'-

P384) from adjacent raft is 7.79 Å. At the 3f position, J9 fab light chain (L1 S26) residue

and E protein (T266) residue from adjacent raft is 8.86 Å. Residue contacts are

calculated as residues with distance between Ca-Ca backbone of Fab and E-protein

respectively.

Discussion:

HEDR bnAbs bind mature as well as tubular DENV as shown in this work

(Fig.8), along with incompletely mature virions as reported previously[16,18].

Expanded/bumpy DENV particles have been described to be present only at 37°C or

above [24,25] and all three bnAbs described here potently neutralize DENV at 37C[16,18].

The different DENV morphologies, ranging from spherical (mature, partially mature,

immature), to expanded and tubular virions, present varying quaternary surface

epitope arrangements and topology. Our current work along with previous reports[29]

shows that bnAb binding is differential to different DENV morphologies, suggesting

that these variation in DENV morphologies might serve as an adaptive mechanism for

escaping antibody-mediated neutralization and immune evasion by the virus. The

common surface feature across these virion forms is the presence of the E-dimer

configuration, which further emphasizes the importance of the E-dimer quaternary unit

as a critical antigenic target.

Figure 8. Structural features of HEDR bnAb-bound DENV complexes and bnAb

binding epitopes.

On comparing the five tubular DENV structures reported here, the helical arrangements of E-dimers were similar within each DENV strain type, irrespective of the Fab bound. The binding of an antibody, that locks the E-dimers together, seems to be a prerequisite to form a globally regular arrangement of E-dimers on tubular DENV, but the specific arrangement achieved by the E-dimers is influenced by the virus strain type. In case of DENV2-US/BID-V594/2006 strain, the helical arrangement resembles the E-raft sequentially wrapping around the tubular virion's length, suggesting tighter inter-E-dimer and inter-E-raft interactions. However, in the THSTI/TRC/01 strain, the relative strength of inter-E-dimer and E-raft interactions appears to be weaker as the helical arrangement is drastically different from the icosahedral E-raft arrangement.

The bnAb complex structures described here, show that these bnAbs directly interact with the fusion loop on E-protein and occupy similar epitopes that bridge the E-dimer, strongly suggesting that the bnAbs lock the E-dimers in the pre-fusion state. The global bnAb footprints overlap with prM binding region on E-protein, which is similar to that observed with EDE bnAbs[15,20], highlighting the conserved nature of this epitope. Fusion loop exposure on the E-protein is necessary for attachment to endosomal membranes and initiate viral membrane fusion in dengue virus cellular entry[5,6]. Similarly, dissociation of E-dimers is necessary for conformational changes in the E-proteins from pre-fusion to post-fusion structure, which has again been shown to be essential for successful virus infection in cells[30,31]. Thus, the bnAbs likely inhibit virus infection in cells by blocking exposure of the fusion loop and impeding E-dimer dissociation, both steps that are essential for viral entry into host cells. Indeed,

previous report on the J9 bnAb has shown that the bnAb obstructs early entry steps in

virus infection[18], which further corroborates our results.

The heavy chain of the HEDR bnAbs interacts with the fusion loop and make

majority of the contacts within an E-dimer. The heavy chains of J9 and D4.F05.S03

are involved in interactions with both the conserved glycans and makes robust

contacts than the respective Fab's light chains. The light chain of the bnAbs is involved

in inter-E-dimer contacts both within and between E-rafts, likely stabilizing the bnAb

binding on the virions. As seen in case of D14.F25.S02, the inter-E-raft light chain

interactions are specific and necessary for neutralization activity of the bnAb.

However, the bnAbs also bind the tubular forms of DENV with good occupancy despite

the disruption in bnAb light-chain contacts as evident in the helical DENV2-

THSTI/TRC/01 complex structures. Thus, the bnAb heavy chain drives the overall

epitope recognition and potency of these HEDR antibodies.

From our results, it is evident that binding of HEDR category 1 bnAb,

D14.F25.S02, distorts spherical DENV. In case of the DENV2-THST/TRC/01 strain,

the effect of D14.F25.S02 binding at saturating concentrations had a dramatic effect

on virion structure and symmetry, impeding higher resolution structure determination

of the virion+Fab complex. However, the Fab binding did not destroy virion

organization but rather appears to have introduced significant local movements in the

E-proteins, as the virions' icosahedral symmetry was still discernible from the low-

resolution 3D density map. Remarkably, a 4.1 Å cryo-EM structure was achievable for

D14.F25.S02 to DENV2-US/BID-V594/2006 strain, indicating that this strain was

relatively resistant to distortion due to bnAb binding which could be a further reflection

of stronger interactions between the E-dimer and E-rafts in the US/BID-V594/2006

strain when compared to THSTI/TRC/01 strain. Nevertheless, analyses of 2D classes

and cryo-EM density maps show that binding of D14.F25.S02 to the US/BID-

V594/2006 virions does affect the topology of the E-protein surface, leading to many

virion particles being discarded during the 3D reconstruction process.

The major class of particles obtained from the D14.F25.S02+DENV2-US/BID-

V594/2006 dataset had the highest Fab occupancy at the icosahedral 5f position. The

local E-dimer arrangement at the 5f position likely facilitates D14.F25.S02 binding as

the larger gap between E-dimers surrounding the 5f vertices would allow easier

insertion of the bnAb HCDR3 loop between adjacent E-dimers. The bnAb also makes

inter-raft contacts that are unique to the 5f position which have been reported to be

crucial for its neutralization activity. These observations indicate that D14.F25.S02

binds preferentially to the 5f position on the icosahedral virus. However, the binding at

5f vertices effectively inhibits bnAb binding to the icosahedral 2f position, suggesting

that the bnAb binding at 5f causes local structural variations which limits bnAb binding

at the other sites. Notably, when D14.F25.S02 binds the 2f and 3f positions on the

virus, there is no binding observed at the 5f position, suggesting that once

D14.F25.S02 binds at 2f position, binding at 5f position is no longer possible for the

bnAb. These observations imply that binding of D14.F25.S02 at the 5f position causes

perturbations in the adjacent E-raft which impedes bnAb binding to the 2f position,

whereas binding of bnAb first to the 2f and 3f positions stabilizes the E-raft such that

E-dimer movement near the 5f position is restricted and thus, could inhibit bnAb

binding. Similar reports of local distortion to DENV structure have been reported for

EDE1 bnAb C10 which also strongly neutralizes DENV and ZIKV[28]. Binding of C10 to

purified DENV E-dimers has been shown to cause hinge-like motions in the E-

protein[15], along with preferred binding of bnAb to 5f and 3f positions and increased

dynamics of E-proteins with increasing Fab concentrations[28]. Thus, despite

differences in binding determinants and specific interactions to the E-proteins, HEDR

category-1 bnAbs and EDE1 bnAbs show similarities in disrupting E-protein

organization on spherical DENV surface.

HEDR category-2 bnAbs, J9 as well as D14.F05.S03, neutralize only DENV

and require interactions with the E-protein's conserved N153 glycan for their

neutralization activity. The HCDR3 loop of these bnAbs interacts with both the N153

glycan as well as the fusion loop on the E-protein. J9 and D14.F05.S03 also interact

with N67 glycan on E-protein which is not observed for D14.F25.S02, suggesting a

role for N67 glycan in epitope recognition. Notably, no distortion of the virion structure

is observed on binding with HEDR category-2 bnAbs.

Structural elucidation of these pan-DENV bnAb epitopes, as described here,

emphasizes the need for vaccine immunogen design that most closely imitate the E-

protein quaternary epitopes, including glycosylation, as presented on mature DENV

surface to elicit broad neutralization responses. For both, D14.F25.S02 and

D14.F05.S03, bivalent engagement as IgGs is necessary for potent neutralization

activity against DENV[16]. Based on the structural data obtained from D14.F25.S02 IgG

bound to DENV2 and previous published reports[13,32,33], it seems possible that virion

aggregation along with locking of E-dimers maybe essential for the neutralization

activity of these bnAbs. Despite the distortion caused by D14.F25.S02 binding to

DENV, the bnAb still has one of the highest reported neutralization potencies for a

pan-DENV and ZIKV bnAb[16]. When comparing neutralization of DENV serotypes

alone, J9 has one of the highest reported neutralization potencies reported till-date[18].

J9 has a relatively narrow footprint on the E-dimer surface, with most residues

important for its neutralization potency located near the fusion loop and the N153-

glycan loop. J9 also makes minimal inter-E-dimer and inter-E-raft contacts, with

primary determinants of bnAb binding localized to individual E-dimers. Furthermore,

J9 is capable of neutralizing DENV as IgG as well as Fab fragment. The potent activity

of J9 as Fab fragment increases its applicability as a therapeutic antibody while

avoiding any ADE effects caused by Fc regions present on IgGs. Similarly, despite the

need for bivalent engagement, antibody engineering[34,35] of D14.F25.S02 could be

leveraged for developing the bnAb as a pan-DENV+ZIKV therapeutic antibody. Thus,

engineered versions of these bnAbs may provide as much needed therapeutics

against severe dengue infections.

Materials and Methods:

<u>Virus sample preparation:</u>

Dengue virus serotype 2 strains THSTI/TRC/01 (GenBank: UNZ79673.1) was

obtained as a kind gift from Dr. Guruprasad Medigishi, THSTI, Faridabad and US/BID-

V594/2006 (GenBank: ACA48992.1) was obtained through BEI Resources, NIAID,

NIH: Dengue Virus Type2, DENV-2/US/BID-V594/2006, NR-43280. Both virus strains

were propagated with minimal passages in mosquito cells (P-6) to 1L culture volumes

and purified in a similar manner as reported previously[2]. Mosquito cells (C6/36) were

cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal

bovine serum (FBS) and 1% nonessential amino acids (NEAA). The cells were

maintained at 28°C with 5% CO_2 in a 10-stack cell factory system (Nunc). Upon

reaching 70-75% confluency, cells were infected with DENV-2 using a multiplicity of

infection (MOI) of 0.1. The infection was allowed to proceed at 30°C with 5% CO2.

Post 2 hours of infection, the media was changed to complete media supplemented

with 5% FBS. At 20 hours post-infection, media was again changed to complete media

supplemented with 2% FBS. Supernatant medium containing the virus was collected

on 4th and 6th day post-infection. Virus-containing media was centrifuged at 9500xg

for 30 minutes and filtered through a 0.22-micron filter to remove all cell debris. The

filtrate was precipitated with 8% polyethylene glycol (PEG) overnight at 4°C, followed

by centrifugation at 9500xg for one hour to collect the PEG pellet. The pellet was then

resuspended in NTE buffer (120 mM NaCl, 20 mM Tris, 1 mM EDTA, pH 8.0). The

resuspended PEG pellet was sedimented through a 24% sucrose cushion. The

dissolved sucrose pellet applied to a potassium tartrate gradient (35%-10%) and

centrifugated at 32000 rpm for 2 hours at 4°C in a SW41-Ti swinging bucket rotor

(Beckman Coulter). Bands corresponding to the virus was extracted from the 20%

fraction of the discontinuous gradient, buffer exchanged and concentrated in NTE

buffer using a 100 kDa centrifugal filter. The purity and concentration of the virus

sample was estimated by visualising under negative staining transmission electron

microscopy (TEM) analysis.

<u>Fab preparation:</u>

Purified IgG of D14.F25.S02, J9, and D14.F05.S03 bnAbs were prepared as

previously reported[16,18]. BnAbs were digested with papain individually to generate Fab

fragments (~50kDa molecular weight) using a commercially available Fab

Fragmentation kit (G Biosciences) (Cat. # 786-273). Fabs were then separated in

flowthrough by passing over a Protein-A column. Concentrations of Fabs obtained for

F25.S02, J9, and F05.S03 were 0.9 mg/ml, 0.6 mg/ml and 1.01 mg/ml, respectively.

Concentration of F25.S02 IgG sample was 1mg/ml.

Cryo-EM sample preparation:

Purified DENV samples at an approximate concentration of 0.25 mg/ml (E-

glycoprotein concentration) were incubated with the respective Fabs and F25.S02

IgG at room temperature for 45 minutes. In case of D14.F25.S02, molar ratio of Fab

to virus was 4:1 and molar ratio of IgG to virus was 1:1. For J9 and D14.F05.S03,

molar ratio of Fab to virus was maintained at 2:1 and 2.5:1 respectively. For

estimating the concentration of dengue virus needed for the experiments, the

concentration of E-glycoprotein estimated from SDS-PAGE was used. Ultra-thin

lacey carbon grids (300 or 400 mesh) were glow discharged using 25 mA current for

30s with a PELCO EasiGlow machine. Aliquots of 3ul virus-Ab mixture were applied

on glow discharged grids and plunge-frozen with FEI Vitrobot mark IV for DENV-2-

THST1/TRC/01 + Fab complexes (Blot time – 8s, Temperature - 4° C, Humidity –

100%) and Leica EM GP2 for DENV2-US/BID-V594/2006 + Ab complexes (Blot time

– 3.5s, Temperature - 4° C, Humidity – 100%).

Cryo-EM data collection:

All DENV-2-THST1/TRC/01 and Fab complex samples were imaged in Talos Arctica

(FEI, ThermoFisher Inc.), operating at 200 keV, with a K2 summit electron detector

(Gatan Inc.). Micrographs were collected with a pixel size of 1.16 Å/pixel, at a total

dose of 38 - 40 e$^-$/Å^2 fractionated into 20 frames per image, and with defocus range of

-1.5 to -2.5 µm.

DENV2 US/BID-V594/2006 samples complexed with J9 Fab, F25.S02 Fab and IgG

were imaged using Titan Krios G3I, operating at 300 keV with a K3 detector and an

energy filter (Bioquantum imaging) operating at a slit width 20eV. Micrographs for each

dataset were collected with a total dose of 40 – 42 e$^-$/Å^2 fractionated into 38 frames

per image at a defocus range of -1 µm to -2.5µm and with a pixel size of 1.1 Å/pixel.

<u>Cryo-EM data processing:</u>

A combination of RELION-4.0[36] and cryoSPARC version 4.7.0 software package[37]

was used for data processing following standard single particle analysis (SPA) steps.

Frame alignment was carried out using Relion's own implementation of motion

correction or Cryosparc's Patch motion correction. CTF estimation was performed

using CTFFIND4[38].

<u>D14.F25.S02-virus complex datasets:</u> For the D14.F25.S02 Fab + DENV2 US/BID-

V594/2006 complex, 7405 micrographs were collected at a magnification of 81000x

with corresponding pixel size of 1.1Å/Pixel. Dataset of D14.F25.S02 IgG + DENV2

US/BID-V594/2006 was also collected at the same magnification and microscope

settings as the Fab datasets with a total of 4172 images. Virion particles were manually

picked from both datasets, with 22000 particles picked from the Fab dataset and 8176

particles from the IgG dataset. Subsequent rounds of 2D classification and 3D

classification were performed on the individual particle datasets. For the Fab complex,

a 3D class with 2305 particles was selected and used for non-uniform refinement[39]

and CTF refinement[40] which resulted in a 4.09 Å density map showing Fab density

near 5f and 3f symmetry vertices. A smaller 3D class containing 905 particles was

selected for non-uniform refinement which resulted in an 8.45 Å density map of the

Fab bound near the 2f-3f symmetry vertices. For the IgG dataset, 2D classification, 3D

refinement and post-processing was carried out in RELION-4.0. The final 3D

reconstruction of 7.65 Å global resolution was calculated from 1045 particles selected

from 3D classification.

For the D14.F25.S02 Fab + DENV-2-THST1/TRC/01 complex, 4000 micrographs

were collected at a magnification of 36000x with corresponding pixel size of

1.16Å/Pixel. From these images, 26782 particles were manually picked and subjected

to unsupervised 2D classification in RELION-4.0. The best classes which showed

apparent Fab densities were selected for reference-based 3D classification in

RELION-4.0. Different starting models, including a sphere, mature DENV structure

and the determined DENV2-US/BID/V594/2006+D14.F25.S02 Fab structure were

used for 3D classification. All the classification routines resulted in similar results.

Finally, a total of 4715 particles from the best classes of 3D classification from

RELION4 were utilised for 3D non-uniform refinement in cryoSPARC-4.7.0 to generate

a 3D density map at 19 Å global resolution.

<u>J9-virus complex datasets</u>: A total of 4756 micrographs were collected for the J9 +

DENV2 US/BID-V594/2006 complex at a magnification of 81000x with corresponding

pixel size of 1.1Å/Pixel. Around 15000 particles were manually picked in RELION-4.0.

Further, the particle coordinates were imported to cryoSPARC-4.7.0. After 2D and 3D

classification, 2900 best particles were used for iterative non-uniform refinement and

CTF refinement, resulting in an electron density map of 3.85 Å global resolution.

For the J9 + DENV-2-THST1/TRC/01 complex, 1100 images were collected at a

magnification of 36000x with corresponding pixel size of 1.16Å/Pixel. Of the 10108

manually picked particles, only 668 particles showed clear Fab densities after 2D

classification in RELION-4.0. These particles were then used non-uniform refinement

in cryoSPARC-4.7.0. The 3D refinement resulted in an electron density map of 7.21 Å

global resolution.

<u>D14.F05.S03-virus complex datasets</u>: A total of 3176 micrographs were collected at a

magnification of 36000x with corresponding pixel size of 1.16Å/pixel for D14.F05.S03

+ DENV-2-THST1/TRC/01 complex. From these images, 32650 particles were

manually picked using RELION-4.0 software. The manually picked particle

coordinates were further imported to cryoSPARC-4.7.0 for further processing. A final

set of 4299 particles were selected for 3D refinement following non-reference 2D

classification and reference-based 3D classification. Subsequent iterative non-uniform

refinement and CTF refinement resulted in a 3D density map with global resolution of

4.51 Å.

<u>Real space refinement and model fitting in icosahedral maps:</u>

The complete IgG sequence of D14.F25.S02 and the Fab portion sequence of

D14.F25.S02, J9, and D14.F05.S03 were used to generate atomic models in

AlphaFold3 web server[41]. The sequences of DENV-2-THST1/TRC/01 and DENV2-

US/BID-V594/2006 were used as queries for homology modelling in the SWISS-

MODEL web server[42] using mature DENV-2 structures (PDB IDs:7KV8 and 7V3F) as

a template. The asymmetric unit of the virus and Fab models were initially rigid body

fitted separately in respective maps. For the different high-resolution maps, different

approaches were used for map sharpening. For D14.F25.S02 and D14.F05.S03

maps, map sharpening was done using the Phenix software package[43]. In case of the

J9 map, map sharpening was carried out using DeepEMhancer[44]. Further, the

generated atomic models were real-space refined using the Phenix software[45,46].

Geometry restraints and Ramachandran clashes were adjusted as needed using

COOT[47]. Glycan components were added to the E-protein atomic models using higher

resolution E-protein structures (PDB IDs: 4UT6, 4UTA) as reference. Glycan residues

(N-acetyl glucosamine (NAG), α-D-mannopyranose (MAN) and β-D-mannopyranose

(BMA)) were manually added to N67 and N153 residues of the E protein using COOT

according to the map densities and were included in the atomic models for real-space

refinement.

For the highest resolution J9 Fab-virus complex map, the model used for real-space

refinement included the complete asymmetric unit containing three E-protein

monomers and Fabs at 5f and 3f position. For the highest resolution D14.F25.S02

Fab-virus complex map, model used for real-space refinement included the E-dimer

with Fab at 5f position. For the highest resolution D14.F05.S03 Fab-virus complex

map, model used for real-space refinement included the complete asymmetric unit

containing 3 E-protein monomers and Fabs at 3f position. Maps and models were

analysed and figures prepared using UCSF Chimera[48] and UCSF ChimeraX[49].

<u>Localized reconstruction at sub-icosahedral symmetry positions:</u>

Particles from the initial 3D classification of icosahedral reconstruction were imported

to SCIPION-3[50]. Vector coordinates of 5 and 3fold vertices were applied and the sub-

particles were defined[51]. The defined sub-particles were extracted, and 3D

classification and refinement were performed in RELION-4.0 according to standard

procedures. Circular masks which covered the Fabs and half of the E-dimer length at

a given vertex, were applied during classification and initial refinement steps.

The localized reconstruction from 5-fold and 3-fold symmetry positions of the J9 +

DENV2/US/HBID-V594/2006 complex was performed with an initial set of 42886 and

71487 defined and filtered sub-particles, respectively. After classification, 20077 and

26545 best-aligned sub-particles were used for 3D refinement resulting in 4.55 Å (5-

fold) and 4.4 Å (3-fold) resolution maps. For the D14.F05.S03 + DENV-2

THSTI/TRC/01 complex, localized reconstruction for the 3f vertices was performed

with 193844 defined sub-particles. After 3D classification, 75433 sub-particles were

used in the final refinement to produce a map of 4.84 Å global resolution. In case of

D14.F25.S02, 3D classification of sub-particles extracted at 5-fold and 3-fold positions

failed to give any class that showed clear Fab densities and thus were not refined

further. The 3D density maps were sharpened in DeepEMhancer[44]. Glycans where

manually added to improved density regions using COOT[47] based on higher resolution

E-protein structures (PDB IDs: 4UT6 and 4UTA) as references. The atomic models

were fitted as rigid bodies using UCSF Chimera[48] followed by real-space refined in

Phenix[45].

<u>Difference map calculations</u>:

The atomic models of each E-dimer + Fab complex were aligned using the E-dimer as

reference in UCSF Chimera. Next, the atomic models of the three Fabs, J9,

D14.F25.S02, and D14.F05.S03, from the complex structures were used to generate

volumes in UCSF Chimera, maintaining equal grid spacing and scaling. The difference

between each pair of Fab volumes was calculated using the RELION-4.0 command

interface (relion_image_handler). The computed difference volumes show the

common footprint shared between Fabs, which lies over the E-protein's fusion loop

region.

 <u>Helical reconstructions of tubular DENV regions:</u>

For the helical reconstruction of tubular regions from DENV2 US/BID-V594/2006

virions, helical segments were manually picked in segment picking of RELION-4.0. For

both D14.F25.S02- and J9-bound virions, the initial helical rise and twist for DENV2

US/BID-V594/2006 tubular particles were estimated manually in UCSF Chimera by

arranging the Fab+E-dimer models around a cylinder of same dimensions as the virion

segments with minimal clashes. A 2D projection of the manually arranged helical

geometries was then used to match with the arrangement observed in 2D class

averages of tubular particles with no symmetry applied. An initial rise and twist for

D14.F25.S02-bound segments (rise=10 Å and twist-=30°) and for J9-bound segments

(rise=13Å and twist=36°) was used for reference-free helical 2D classification[52] in

RELION-4.0. Best 2D classes containing a total of 2188 segments for D14.F25.S02-

bound dataset and 1361 segments for J9-bound dataset were subjected to 3D

classification with initial estimates using a 3D-cylinder as starting model. In case of

D14.F25.SO2 dataset, 405 segments were imported for final helical 3D reconstruction

in cryoSPARC-4.7.0 using non-uniform refinement option with rise and twist of 11.25Å

and 29.48° which yielded a density map of global resolution 14.7 Å. For the J9 dataset,

a best class with 253 segments was used for helical 3D reconstruction in RELION-4.0

with rise and twist of 12.83Å and 35.83° that resulted in a 3D map with estimated global

resolution of 44Å.

For bnAb bound tubular-shaped particles found in DENV-2 THSTI/TRC/01 datasets,

the tubular-shaped regions of the virions were manually picked following the helical

particle picking protocol of RELION-4.0 and unsupervised 2D classification was

performed. Class averages with distinct Fourier amplitude patterns were imported to

PyHi Fourier Bessel indexing software[53] and helical rise and twist values were

calculated for the antibody-virus complexes. From the calculated rise and twist,

another round of unsupervised helical 2D classification was performed, and the

particles were extracted with the same parameters for reference-based 3D

classification. For initial 3D classifications 3D-cylinder was used as startup model. The

best 3D classes were selected, and the particle co-ordinates were transferred to

cryoSPARC-4.7.0 for final helical reconstruction using non-uniform refinement option.

For the J9 + DENV-2 THSTI/TRC/01 complex, 19792 segments were manually

picked, and after classification routines, 5769 segments were used for the final 3D

refinement, which resulted in a helical density map of 14.41 Å. Similarly, for the

D14.F25.S02 and D14.F05.S03 complexes, 8264 and 8045 segments were

respectively picked in manual mode, and final sets of 1355 (for D14.F25.S02) and

1311 segments (for D14.F05.S03) were used to generate 3D maps at global

resolutions of 17.37 Å and 17.29 Å respectively.

In all helical reconstructions, a hollow cylindrical mask was used which focused on the

E-protein and membrane layer of DENV tubular particles during all the refinement

steps.

<u>Inner volume calculations for capsule-shaped DENV2-US/BID-V4594/2006:</u>

For D14.F25.S02-bound virions:

Average length of cylindrical portion in capsule-shaped DENV = 79 nm

Inner radius of capsule = 5.5 nm.

Thus, internal volume of cylindrical portion in capsule-shaped DENV = 7.503×10^{-24}

m^3

Assuming, volume of one copy of viral genome[26,54] = 3.8×10^{-24} m^3

Number of genome copies that can be possibly accommodated in capsule-shaped

DENV = 1.9

For J9-bound virions:

Average length of cylindrical portion in capsule-shaped DENV = 74.149nm

Inner radius of capsule= 6nm

Thus, internal volume of cylindrical portion of capsule-shaped DENV = 8.38×10^{-24}

m3

Assuming, volume of one copy of viral genome[26,54] = $3.8 \times 10\text{-}24$ m^3

Number of genome copies that can be possibly accommodated in capsule-shaped

DENV = 2.2

References:

1.   Bhatt, S. *et al.* The global distribution and burden of dengue. *Nature* **496**, 504–507
(2013).
2.   Kuhn, R. J. *et al.* Structure of Dengue Virus. *Cell* **108**, 717–725 (2002).
3.   Modis, Y., Ogata, S., Clements, D. & Harrison, S. C. A ligand-binding pocket in the
dengue virus envelope glycoprotein. *Proceedings of the National Academy of*
Sciences **100**, 6986–6991 (2003).
4.   Zhao, R. *et al.* Flavivirus: From Structure to Therapeutics Development. *Life* **11**, 615
(2021).
5.   Huang, C. Y.-H. *et al.* The dengue virus type 2 envelope protein fusion peptide is
essential for membrane fusion. *Virology* **396**, 305–315 (2010).
6.   Nicholls, C. M. R., Sevvana, M. & Kuhn, R. J. Structure-guided paradigm shifts in
flavivirus assembly and maturation mechanisms. in 33–83 (2020).
doi:10.1016/bs.aivir.2020.08.003.
7.   Chong, L. C. & Khan, A. M. Identification of highly conserved, serotype-specific
dengue virus sequences: implications for vaccine design. *BMC Genomics* **20**, 921
(2019).
8.   Dejnirattisai, W. *et al.* Cross-Reacting Antibodies Enhance Dengue Virus Infection in
Humans. *Science (1979)* **328**, 745–748 (2010).
9.   Rey, F. A., Stiasny, K., Vaney, M.-C., Dellarole, M. & Heinz, F. X. The bright and the
dark side of human antibody responses to flaviviruses: lessons for vaccine design.
EMBO Rep **19**, 206–224 (2018).

10. Morrone, S. R. & Lok, S.-M. Structural perspectives of antibody-dependent
enhancement of infection of dengue virus. *Curr Opin Virol* **36**, 1–8 (2019).
11. Katzelnick, L. C. *et al.* Antibody-dependent enhancement of severe dengue disease in
humans. *Science* **358**, 929–932 (2017).
12. Pierson, T. C. & Diamond, M. S. The continued threat of emerging flaviviruses. *Nat*
Microbiol **5**, 796–812 (2020).
13. Zhang, S. *et al.* A Human Antibody Neutralizes Different Flaviviruses by Using
Different Mechanisms. *Cell Rep* **31**, 107584 (2020).
14. Dejnirattisai, W. *et al.* Dengue virus sero-cross-reactivity drives antibody-dependent
enhancement of infection with zika virus. *Nat Immunol* **17**, 1102–1108 (2016).
15. Sharma, A. *et al.* The epitope arrangement on flavivirus particles contributes to Mab
C10's extraordinary neutralization breadth across Zika and dengue viruses. *Cell* **184**,
6052-6066.e18 (2021).
16. Lubow, J. *et al.* Single B cell transcriptomics identifies multiple isotypes of broadly
neutralizing antibodies against flaviviruses. *PLoS Pathog* **19**, e1011722 (2023).
17. Barba-Spaeth, G. *et al.* Structural basis of potent Zika–dengue virus antibody cross-
neutralization. *Nature* **536**, 48–53 (2016).
18. Durham, N. D. *et al.* Broadly neutralizing human antibodies against dengue virus
identified by single B cell transcriptomics. *Elife* **8**, (2019).
19. Dejnirattisai, W. *et al.* A new class of highly potent, broadly neutralizing antibodies
isolated from viremic patients infected with dengue virus. *Nat Immunol* **16**, 170–177
(2015).
20. Rouvinski, A. *et al.* Recognition determinants of broadly neutralizing human
antibodies against dengue viruses. *Nature* **520**, 109–113 (2015).
21. Rey, F. A., Stiasny, K., Vaney, M.-C., Dellarole, M. & Heinz, F. X. The bright and the
dark side of human antibody responses to flaviviruses: lessons for vaccine design.
EMBO Rep **19**, 206–224 (2018).
22. Plevka, P. *et al.* Maturation of flaviviruses starts from one or more icosahedrally
independent nucleation centres. *EMBO Rep* **12**, 602–606 (2011).
23. Junjhon, J. *et al.* Influence of pr-M Cleavage on the Heterogeneity of Extracellular
Dengue Virus Particles. *J Virol* **84**, 8353–8358 (2010).
24. Zhang, X. *et al.* Dengue structure differs at the temperatures of its human and
mosquito hosts. *Proc Natl Acad Sci U S A* **110**, 6795–9 (2013).
25. Fibriansah, G. *et al.* Structural changes in dengue virus when exposed to a
temperature of 37°C. *J Virol* **87**, 7585–92 (2013).
26. Morrone, S. R. *et al.* High flavivirus structural plasticity demonstrated by a non-
spherical morphological variant. *Nat Commun* **11**, 3112 (2020).
27. Kudlacek, S. T. *et al.* Designed, highly expressing, thermostable dengue virus 2
envelope protein dimers elicit quaternary epitope antibodies. *Sci Adv* **7**, eabg4084
(2021).
28. Lim, X.-X. *et al.* Human antibody C10 neutralizes by diminishing Zika but enhancing
dengue virus dynamics. *Cell* **184**, 6067-6080.e13 (2021).
29. Fibriansah, G., Lim, X.-N. & Lok, S.-M. Morphological Diversity and Dynamics of
Dengue Virus Affecting Antigenicity. *Viruses* **13**, 1446 (2021).
30. Chao, L. H. *et al.* How small-molecule inhibitors of dengue-virus infection interfere
with viral membrane fusion. *Elife* **7**, (2018).
31. Stiasny, K., Fritz, R., Pangerl, K. & Heinz, F. X. Molecular mechanisms of flavivirus
membrane fusion. *Amino Acids* **41**, 1159–1163 (2011).

32. Zhou, Q. F. *et al.* Structural basis of Chikungunya virus inhibition by monoclonal antibodies. *Proceedings of the National Academy of Sciences* **117**, 27637–27645 (2020).

33. Wang, J. *et al.* A Human Bi-specific Antibody against Zika Virus with High Therapeutic Potential. *Cell* **171**, 229-241.e15 (2017).

34. Hale, G. Living in LALA land? Forty years of attenuating Fc effector functions. *Immunol Rev* **328**, 422–437 (2024).

35. Rosenstein, S. *et al.* Production of F(ab')2 from Monoclonal and Polyclonal Antibodies. *Curr Protoc Mol Biol* **131**, e119 (2020).

36. Kimanius, D., Dong, L., Sharov, G., Nakane, T. & Scheres, S. H. W. New tools for automated cryo-EM single-particle analysis in RELION-4.0. *Biochem J* **478**, 4169–4185 (2021).

37. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290–296 (2017).

38. Rohou, A. & Grigorieff, N. CTFFIND4: Fast and accurate defocus estimation from electron micrographs. *J Struct Biol* **192**, 216–21 (2015).

39. Punjani, A., Zhang, H. & Fleet, D. J. Non-uniform refinement: adaptive regularization improves single-particle cryo-EM reconstruction. *Nat Methods* **17**, 1214–1221 (2020).

40. Zivanov, J., Nakane, T. & Scheres, S. H. W. Estimation of high-order aberrations and anisotropic magnification from cryo-EM data sets in RELION-3.1. *IUCrJ* **7**, 253–267 (2020).

41. Abramson, J. *et al.* Addendum: Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **636**, E4 (2024).

42. Waterhouse, A. *et al.* SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Res* **46**, W296–W303 (2018).

43. Terwilliger, T. C., Sobolev, O. V, Afonine, P. V & Adams, P. D. Automated map sharpening by maximization of detail and connectivity. *Acta Crystallogr D Struct Biol* **74**, 545–559 (2018).

44. Sanchez-Garcia, R. *et al.* DeepEMhancer: a deep learning solution for cryo-EM volume post-processing. *Commun Biol* **4**, 874 (2021).

45. Afonine, P. V *et al.* Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta Crystallogr D Struct Biol* **74**, 531–544 (2018).

46. Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861–877 (2019).

47. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486–501 (2010).

48. Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem* **25**, 1605–12 (2004).

49. Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci* **32**, e4792 (2023).

50. de la Rosa-Trevín, J. M. *et al.* Scipion: A software framework toward integration, reproducibility and validation in 3D electron microscopy. *J Struct Biol* **195**, 93–9 (2016).

51. Ilca, S. L. *et al.* Localized reconstruction of subunits from electron cryomicroscopy images of macromolecular complexes. *Nat Commun* **6**, 8843 (2015).

52. He, S. & Scheres, S. H. W. Helical reconstruction in RELION. *J Struct Biol* **198**, 163–
176 (2017).
53. Zhang, X. Python-based Helix Indexer: A graphical user interface program for finding
symmetry of helical assembly through Fourier-Bessel indexing of electron microscopic
data. *Protein Sci* **31**, 107–117 (2022).
54. Yoffe, A. M. *et al.* Predicting the sizes of large RNA molecules. *Proc Natl Acad Sci U*
S A **105**, 16153–8 (2008).

1046

1047

1048 **Data availability**

1049 Cryo-electron microscopy density maps and corresponding atomic models presented

1050 in this study has been submitted to the Electron Microscopy Data Bank (EMDB) and

1051 RCSB Protein Data Bank (PDB). Accession codes for the same are as follows:

S.no.	Sample description	EMDB ID	PDB ID
1.	J9 Fab + DENV2-US/BID-V594/2006 icosahedral reconstruction	EMD-63759	9MAV
2.	5-fold Localized reconstruction of J9 Fab + DENV2-US/BID-V594/2006 icosahedral reconstruction	EMD-63760	9MAX
3.	3-fold Localized reconstruction of J9 Fab + DENV2-US/BID-V594/2006 icosahedral reconstruction	EMD-63761	9MAY
4.	J9 Fab + DENV2-US/BID-V594/2006 helical reconstruction	EMD-63762	9MAZ
5.	J9 Fab + DENV2-THSTI/TRC/01 icosahedral reconstruction	EMD-63763	9MB0
6.	J9 Fab + DENV2-THSTI/TRC/01 helical reconstruction	EMD-63764	9MB1

7.	D14.F25.S02 Fab + DENV2-US/BID-V594/2006 icosahedral reconstruction (5f-3f map)	EMD-63643	9M5H, 9U33
8.	D14.F25.S02 Fab + DENV2-US/BID-V594/2006 icosahedral reconstruction (2f-3f map)	EMD-63813	9U38
9.	D14.F25.S02 IgG + DENV2-US/BID-V594/2006 icosahedral reconstruction	EMD-63812	9U37
10.	D14.F25.S02 Fab + DENV2-US/BID-V594/2006 helical reconstruction	EMD-63809	9U34
11.	D14.F25.S02 Fab + DENV2-THSTI/TRC/01 helical reconstruction	EMD-63810	9U35
12.	D14.F25.S02 Fab + DENV2-THSTI/TRC/01 icosahedral reconstruction	EMD-63811	9U36
13.	D14.F05.S03 Fab + DENV2-THSTI/TRC/01 icosahedral reconstruction	EMD-63801	9MCF
14.	3-fold Localized reconstruction of D14.F05.S03 Fab + DENV2-THSTI/TRC/01 icosahedral reconstruction	EMD-63800	9MCE
15.	D14.F05.S03 Fab + DENV2-THSTI/TRC/01 helical reconstruction	EMD-63815	9U3J

1052

1053

Competing interests

1055 The authors declare no competing interests.

1056

Supplementary Information:

1058 Supplementary figures and tables including extended data information are included

1059 in separate supplementary information document.

1060

1061 Acknowledgements

We thank the Advanced Center for Cryo-Electron Microscopy (ACCEM) facility at

Indian Institute of Science, Bangalore (IISc) and the DST-SATHI Cryo-EM facility at

IIT-Delhi for data collection time and support. We also thank the Bioimaging facility at

IISc which houses the 120kV room temperature TEM for time and support. We thank

Dr. Kelly Lee, University of Washington for his support during project initiation and

scientific discussions. This study was supported by SERB Power grant

(SPG/2021/002433) to V.M.P. This study was also partially supported through the

India Alliance DBT Wellcome Intermediate fellowship (IA/I/22/1/506233) and start-up

grant fund (IISc) to V.M.P; IoE fellowship to S.S and A.R, and Prime Minister's

Research Fellowship (PMRF), Govt. of India, to A.C.

1072

1073 Author contributions:

1074 Conceptualization L.G and V.M.P; IgG purification, J.L; virus purification, A.C and

1075 S.C; Fab production, virus-complex preparations, cryo-EM data collection and data

1076 processing, A.R, A.C and S.S; EM data analysis, A.C, A.R, S.S and V.M.P; Writing,

1077 V.M.P, A.C, A.R and S.S; resources, V.M.P.; funding acquisition, V.M.P.