# Multi-compartment spatiotemporal metabolic modeling of the chicken gut guides dietary intervention design

Irina Utkina[1], Mohammadali Alizadeh[2], Shayan Sharif[2], John Parkinson[1,3,*]

[1] Program in Molecular Medicine, Hospital for Sick Children, Toronto, Ontario, Canada

[2] Department of Pathobiology, Ontario Veterinary College, University of Guelph, Guelph, Ontario, Canada

[3] Department of Biochemistry, University of Toronto, Toronto ON, Canada

* Contact information for the Corresponding Author:

Program in Molecular Medicine, The Hospital for Sick Children, Toronto, ON, Canada M5G 0A4

jparkin@sickkids.ca

**Keywords:** Microbiome; Metabolic Modeling; Poultry Gut; Dietary Interventions; Metabolic Interactions

## ABSTRACT

Understanding the interactions between diet and the gut microbiome is critical for identifying dietary interventions that support gut health. This is of particular importance for poultry where the elimination of antibiotic growth promoters has resulted in an alarming rise in enteric infections with significant economic consequences. While the application of computational models capable of dissecting the metabolic interactions supporting gut communities has shown promise, they remain limited, largely ignoring the physiological and geographical considerations of the poultry gastrointestinal tract. To address these limitations, we developed the first multi-compartment, spatiotemporally-resolved metabolic model of the chicken gastrointestinal tract. Our six-compartment framework integrates avian-specific physiological features including bidirectional flow, feeding-fasting cycles, and compartment-specific environmental parameters. The model captured distinct metabolic specialization along the gut, with upper compartments enriched for biosynthetic pathways and lower compartments specialized for fermentation. *In silico* screening of 34 dietary supplements revealed context-dependent metabolic responses and predicted cellulose, starch, and L-threonine as robust enhancers of short-chain fatty acid production. A controlled feeding trial validated key predictions, particularly for butyrate, with prediction accuracy further improved through integration of trial-specific microbial community data. Our findings demonstrate that community composition is a major driver of metabolic outcomes and underscore the need for context-specific modeling. Our framework provides a mechanistic platform for rational dietary intervention design and is broadly adaptable to other animal or human gastrointestinal systems.

# INTRODUCTION

Over the past eight decades, antibiotic growth promoters (AGPs) have played an important role in livestock production, promoting gut health and reducing the burden of infectious disease[1]. However, the rise of antimicrobial resistance (AMR) associated with the use of AGPs (with AMR hotspots persisting across major poultry-producing regions in Asia, South America, and North America[2]) has created an urgent need for efficacious alternatives[3,45,62]. Research efforts have identified several promising alternatives, including phytochemicals, organic acids, probiotics, and prebiotics[7,8]. However, these interventions produce inconsistent outcomes across production systems, with substantial variability in efficacy between studies and farms[7–9]. This inconsistency reflects a critical challenge: intervention efficacy depends on baseline conditions, including diet composition, microbiota structure, and host physiology[9]. Consequently, identification of effective alternatives requires a more complete understanding of how potential interventions interact in the context with these additional factors.

At the core of these relationships are the metabolic interactions between gut microbiota, responsible for the degradation of dietary components and provision of nutrients, both to the microbial community as well as the host. Helping dissect these relationships are frameworks such as constraint-based metabolic models capable of explicitly simulating biochemical reaction networks. Recent advances have scaled these models from single species to community-scale frameworks capable of capturing interspecies interactions and metabolite exchange[10,11]. Tools such as MICOM and BacArena now enable quantitative simulations of microbiome metabolism under dietary perturbations[12–14]. In humans, these approaches have accurately predicted individual short-chain fatty acid (SCFA; metabolites associated with gut health[15]) profiles and revealed mechanistic links between microbial metabolism, host physiology, and diet[16,17]. Multi-compartment and agent-based frameworks have been established for mammalian systems, capturing pH gradients, oxygen diffusion, motility, and absorption along the gastrointestinal tract (GIT)[13,19–23]. Further, studies involving these platforms have shown that modeling the spatial organization of microbial communities improves predictive accuracy for metabolite fluxes and dietary intervention outcomes[19,21,22].

Despite the limited application of community metabolic modeling to livestock, initial studies have shown much promise, revealing the impact of keystone taxa, as well as AGPs and their potential alternatives[24–26]. A significant limitation in these models, however, has been the

focus on a single compartment, typically the ceca. Such models neglect the potential

contributions of other compartments that feature distinct physical and chemical

environments. Consequently, they do not capture the progressive processing of dietary

components as they transit the GIT, or the functional contributions of individual taxa that

may vary across sites – a limitation that is particularly relevant in the search for health-

promoting probiotics that may colonize different regions[27,28]. The omission of these features

impacts predictive accuracy for interventions affecting multiple gut sites and underscore the

need for physiologically grounded, multi-compartment frameworks.

To address these limitations, we developed the first multi-compartment, spatiotemporally

resolved metabolic model of the chicken GIT. Our six-compartment framework (gizzard,

duodenum, jejunum, ileum, cecum, colon) integrates avian-specific physiological features,

including bidirectional flow through peristalsis and reverse peristalsis, feeding-fasting cycles

with diurnal shifts in gut motility, compartment-specific environmental parameters (pH,

oxygen gradients, transit times), and nutrient absorption by the host. To reflect realistic

variation in poultry production systems, the model was parameterized using 16S rRNA

sequence survey data from chickens fed corn- or wheat-based diets, two nutritionally distinct

formulations known to drive markedly different gut community structures and fermentative

outputs[29–31]. The framework captured compartment-specific metabolic profiles, taxon-

specific SCFA production, and temporal metabolite dynamics across feeding cycles.

Systematic screening of 34 dietary supplements revealed context-dependent metabolic

responses shaped by microbial community composition and gut region and identified

cellulose, starch, and L-threonine as robust enhancers of SCFA production in the lower GIT.

A controlled feeding trial provided direct experimental validation, with the accurate

prediction of butyrate production. We further show that prediction accuracy increases with

the use of community compositions representative of the birds being tested, that we suggest

should be a prerequisite for developing precision nutrition strategies. Overall, our framework

provides a mechanistic platform for rational design of dietary interventions to replace AGPs

and advance strategies for sustainable poultry production.

## RESULTS

**Multi-compartment framework enables spatially resolved metabolic modeling**

Previous metabolic models of chicken gut microbiomes have focused exclusively on cecal metabolism, neglecting how community composition varies across gut regions and how dietary nutrients are sequentially transformed during transit. To address these limitations, we developed a six-compartment model predicting both taxon-specific metabolic contributions and the fate of dietary inputs along the gastrointestinal tract. This framework connects adjacent gut sections, spanning from gizzard to colon, allowing the flow of both microbes and metabolites, and incorporates host physiological features including pH, oxygen gradients, motility, enzymatic digestion, and nutrient absorption.

Microbial community models for each compartment were built using 16S rRNA profiles from 10-day-old chickens fed either corn- (n=5) or wheat-based (n=5) diets. For each compartment, we selected representative taxa by identifying the most abundant bacterial families (comprising >1% relative abundance) and choosing the dominant taxa (represented as distinct Amplicon Sequence Variants (ASVs)) within those families that were mapped to high-quality chicken gut-derived MAGs[34–36]. Taxon selection was further informed by two additional criteria: documented prevalence in poultry microbiome literature and transcriptional activity in metatranscriptomic data from the same trial[9]. Genome-scale metabolic models for the resulting 29 bacterial taxa representing the core microbiota across compartments were reconstructed with gapseq[38]. To ensure the inclusion of key fiber-degrading pathways, models were additionally curated with carbohydrate-active enzymes derived from CAZyme predictions.

To account for physiological differences between gut sections, compartment-specific parameters such as pH, oxygen levels, and bacterial population sizes were set to reflect previously reported gradients along the tract (**Figure 1B**; see Methods). Nutritional inputs were calibrated to average daily feed intake and bacterial density, with feeding intervals matching poultry husbandry regimes (see Supplementary Methods). Transit of material (metabolites and taxa) between compartments was defined with reference to previous studies[89,100]. Nighttime dynamics, including reverse peristalsis, were simulated to mimic the fasted state and the reintroduction of digesta into the upper compartments. Finally, host-driven digestion and absorption of fatty acids, amino acids, and simple sugars were

incorporated, with fractions removed from the relevant sections to reflect host uptake.
Together, the developed comprehensive *in silico* framework provides a spatially and
temporally resolved representation of microbial community dynamics in the chicken GIT.

**Compartment-specific metabolic capabilities reveal distinct functional**
**specialization along the chicken GIT**

The 29 selected taxa span diverse phylogenetic groups and metabolic capabilities, with
compartment-specific abundances reflecting the selective pressures of each gut region
(**Figure 1C**). Metabolic capacities varied substantially across taxa: Enterobacteriaceae and
other metabolic generalists (taxa with large, broadly distributed repertoires) encoded the
highest numbers of reactions and pathways, while obligate anaerobes such as
Oscillospiraceae displayed more specialized repertoires, reflecting adaptation to fiber
fermentation in the lower gut. PCoA of enzyme content revealed clear metabolic stratification
among the reconstructed models (**Figure 1D**). Metabolic generalists (*Escherichia, Klebsiella,*
*Pseudomonas*) were separated from fermentation specialists (members of Lachnospiraceae
and Oscillospiraceae families), with *Lactobacillus* species forming an isolated cluster likely
reflecting their specialized fermentative metabolism. Notably, *Streptococcus* and
*Romboutsia*, despite showing diet-dependent prevalence patterns in upper GIT compartments,
clustered closely together, indicating metabolic similarity in the oxygenated, nutrient-rich
upper gut. This suggests that taxonomic shifts driven by diet do not necessarily translate to
functional divergence in the upper GIT communities. This contrasts with lower GIT
communities, where taxonomic differences between corn- and wheat-fed birds
(Lachnospiraceae/Clostridiaceae versus Enterobacteriaceae dominance) corresponded to
distinct metabolic profiles.

Enzyme-level analysis further highlighted taxon-specific metabolic niches (**Figure S2**).
Obligate anaerobes restricted to lower GIT compartments (members of Ruminococcaceae,
Oscillospiraceae and Erysipelotrichaceae) encoded limited enzyme repertoires ($510\pm62$ ECs)
concentrated in glycan metabolism pathways ($331\pm44$ pathways), consistent with
specialization in cecal and colonic polysaccharide fermentation. By contrast, facultative
anaerobes present throughout the GIT including cecum and colon (Enterobacteriaceae,
*Pseudomonas*, and *Acinetobacter*) encoded substantially larger enzyme repertoires ($913\pm145$
ECs) spanning diverse metabolic pathways ($600\pm81$ pathways), reflecting their capacity as
metabolic generalists. Their presence in the lower GIT likely supports obligate anaerobes

through oxygen consumption and diverse metabolite production. Upper GIT-associated facultative anaerobes (*Streptococcus*, *Romboutsia*, *Lactobacillus*) encoded intermediate metabolic repertoires (505±114 ECs, 357±80 pathways), reflecting adaptation to rapidly available nutrients in the upper gut. Certain taxa, including *Sphingomonas*, *Acinetobacter*, and *Klebsiella*, uniquely possessed enzymes linked to xenobiotic degradation, suggesting ecological roles in processing dietary or environmental compounds that other members of the gut microbiome are unable to metabolize.

At the community level, PCoA based on reaction abundances revealed distinct clustering patterns that highlight metabolic specialization along the chicken gastrointestinal tract (**Figure 1E**). Although the metabolic profiles of the microbial communities differed significantly across gut compartments (PERMANOVA p=0.001), the pairwise differences between compartments were less pronounced (TukeyHSD p>0.05). The primary separation (PC1) distinguished upper from lower GIT communities based on contrasting metabolic strategies (**Figure S3A**). Upper GIT compartments were specifically enriched for bile salt hydrolases (glycocholate, glycochenodeoxycholate, taurodeoxycholate amidohydrolases), membrane lipid synthesis pathways (phosphatidate phosphatase, glycerol-3-phosphate cytidylyltransferase), and fatty acid biosynthesis (acetyl-CoA carboxylase), consistent with rapid biosynthetic activity in nutrient-rich, oxygen-replete conditions (**Figure S3B**). In contrast, cecal and colonic communities formed distinct clusters driven by enrichment in anaerobic fermentation pathways. Key metabolic specializations included SCFA synthesis (butyryl-CoA dehydrogenase, 3-hydroxybutanoyl-CoA oxidoreductase, butanoyl-CoA phosphotransferase for butyrate and propionate production), polysaccharide degradation (galacturonan hydrolases, arabinofuranosidases), and amino acid fermentation. The colon's metabolic profiles partially overlapped with upper GIT profiles, reflecting the direct anatomical linkage between the ileum and colon, enabling the gradual transition in microbial activity and community composition as the digesta moves from the small to the large intestine (**Figure 1E**).

**Temporal dynamics of metabolite concentrations reveal compartment-specific signatures**

Model simulations predicted taxon-specific SCFA production patterns that align with known metabolic capabilities (**Figure 2A**). *Escherichia flexneri* and *Klebsiella pneumoniae*, along with representatives of Lachnospiraceae and Erysipelotrichaceae families, were predicted to

be among the major contributors to acetate production in the ileum, cecum, and colon.
Acetate consumption by *Faecalibacterium* and *Clostridium* spp. fueled butyrate
production[102,103], while *Streptococci,* particularly abundant in wheat-fed birds, displayed
metabolic flexibility, producing acetate in the more oxygenated ileal environment but
consuming it under anaerobic, glucose-scarce conditions in the cecum and colon. Such shifts
align with experimental evidence for environment-dependent regulation of Streptococcal
acetate metabolism[104–107]. *Bacteroides* species emerged as the primary propionate producers
in the lower GIT, corroborating the well-documented capacities of this genus[108](**Figure 2A**).
Together, these findings demonstrate that the six-compartment model captures both taxon-
level metabolic repertoires and community-level metabolic specialization, including the
adaptation of microbial metabolism to the varying physiological conditions of different gut
sections and reliable prediction of SCFA production, consistent with previous reports.

Metabolite profiles across 96-hour simulations revealed dynamic compartment-specific
signatures that shifted predictably with feeding-fasting cycles (**Figure 2B**). At the beginning
of the feeding period, metabolite profiles clustered tightly within their respective
compartments, reflecting the compartment-specific metabolic activity following fasting, with
overlapping cecum and colon profiles, consistent with their similar environments and
functionality. Notably, the unprocessed diet formed a distinct cluster, that remained separated
from all gut compartments throughout simulations**,** demonstrating that microbial communities
extensively transform dietary inputs into modified metabolite profiles rather than simply
passing through dietary compounds unchanged. As feeding progressed, jejunal and ileal
profiles showed increased dispersion, with the ileum shifting away from the cecum and colon
and aligning more with the duodenum and jejunum, likely reflecting intensified nutrient
processing as digesta transit accelerates. Cecal and colonic profiles remained stable and
distinct, reinforcing their role as fermentation sites. During fasting, distal compartments
maintained their clustering but shifted modestly toward the small intestine, while jejunal and
ileal profiles converged, consistent with retrograde flow (reflux) of contents during periods of
reduced peristaltic activity and nutrient scarcity, which redistributes metabolites backward
along the GIT. The reflux mechanism is likewise responsible for the observed similarity in
profiles between the duodenum and the gizzard.

Analysis of specific metabolite classes further illustrated these compartmental and temporal
dynamics (**Figure 2C,D**). Simple sugars such as glucose, fructose, and sucrose accumulated

in the duodenum and jejunum during feeding periods but were largely depleted in distal compartments. Residual glucose observed in the cecum and colon originated from the progressive breakdown of starch-derived oligosaccharides (maltoundecaose, maltohexaose, amylotriose). During fasting, glucose levels increased in the small intestine as a consequence of reflux and continued digestion of residual oligosaccharides. During the feeding period, amino acids including methionine, threonine, and alanine peaked in the duodenum and jejunum and declined downstream, reflecting rapid absorption in the small intestine; during fasting, reflux redistributed amino acids in upstream compartments, sustaining microbial growth in the absence of dietary inputs. Fermentation products (acetate, propionate, butyrate, ethanol, lactate and formate) reached the highest concentrations in the cecum and colon, particularly during fasting when the community relies on residual substrates from starch digestion and dietary fiber components.

Beyond accurately recapitulating concentration changes in primary substrates and fermentation products, the model also captured expected spatial and temporal dynamics of bile acids, fatty acids, and neuroactive metabolites. For example, concentrations of the secondary bile acid glycochenodeoxycholate peaked in the duodenum and jejunum during feeding and sharply declined in downstream compartments, while fasting caused modest increases in the upper compartments due to reflux (**Figure 2C**). Hexanoate concentrations peaked in the upper GIT (gizzard, duodenum, jejunum) during fasting, likely due to reflux-mediated redistribution, consistent with previous reports of higher medium-chain fatty acids in the chicken upper GIT relative to the lower GIT[109]. Interestingly, gamma-aminobutyric acid (GABA), a neurotransmitter produced by the gut microbiota and known for its neuroactive roles[110], showed distinct temporal shifts across the GIT (**Figure 2C**). After fasting, GABA concentrations were highest in the cecum and colon (0.0025 ± 0.0007 and 0.0023 ± 0.0005 mM, respectively), reflecting sustained microbial production overnight. During feeding, concentrations increased in the duodenum (22-fold, to 0.0016 ± 0.0007 mM) and ileum (to 0.0023 ± 0.0011 mM). The jejunum showed the greatest fluctuation of GABA across the feeding-fasting cycle (Δ0.0025 mM), while cecal and colonic concentrations remained relatively stable. Such findings may have implications for the dynamics of GABA availability and its role in the gut-brain axis.

The time-dependent changes in metabolite concentrations observed *in silico* reveal how gut microbial communities respond to substrate availability and gut motility. Additionally, these

270 temporal patterns highlight a critical methodological consideration: metabolite concentrations
271 vary dramatically with sampling time. For instance, during fasting, SCFA levels peak in the
272 cecum and colon, as fermentation intensifies and transit slows, allowing accumulation. Such
273 findings underscore the importance of the timing of sample collection in experimental studies
274 of gut metabolism[111].

## Supplementation with L-threonine or starch is predicted to enhance SCFA production by diverse microbial communities

Establishing a model that recapitulates the spatial dynamics of metabolism in the chicken
GIT, provides a valuable framework for examining the impact of dietary supplements. Of
particular interest is understanding how differences in gut microbial communities may impact
results and which supplements may be least susceptible to variations in these communities.
With established roles in gut health[15], we focused investigations on the production of SCFAs
and screened 34 candidate supplements including established prebiotics (e.g., inulin[112],
XOS[113], GOS[114] and MOS[115]), non-digestible carbohydrates with documented beneficial
effects on chicken gut health and performance (e.g., beta-glucans[116],
arabinooligosaccharides[117], resistant starches[118,119]), essential nutrients commonly
supplemented in poultry feed (e.g., thiamin[119], riboflavin[120] and zinc[121]), and metabolites
identified through flux balance analysis as potential growth-limiting factors for beneficial
bacteria (e.g., raffinose, L-leucine and pantothenic acid). Simulations involving
supplementation of each metabolite were performed for 96-hours for each of the ten
communities sampled (5 wheat-based and 5 corn-based with 5 technical *in silico* replicates
performed for each community, see Methods).

Overall, we found that the metabolic response of each supplement varied according to both
the supplement itself and the type of community (i.e., wheat- vs corn-based; **Figure 3A;**
**Figure S4**). Further analysis of SCFA production fluxes demonstrated that multiple
supplements could substantially alter SCFA production rates (**Figure 3B,C; Figure S4**). For
instance, high doses of amino acid supplementation showed consistent results: L-threonine
and glycine were predicted to significantly increase acetate production in both ileum and
cecum irrespective of baseline community composition. Selected complex carbohydrates
demonstrated similarly robust impacts: high doses of starch and cellulose were predicted to
enhance production of all three SCFAs in both the ileum and cecum relative to controls.
Several compounds displayed dose-dependent effects, with opposing outcomes at different

concentrations (**Figure S4**). Most notably for propionate production: maltoheptaose

supplementation at lower doses had minimal impact, while higher doses significantly

decreased production, particularly in the cecum. Pantothenic acid showed similar

concentration-dependent divergence, with low doses predicted to enhance SCFA production

while higher doses reduced production. These opposing responses likely reflect shifts in

community dynamics where higher substrate concentrations may favor fast-growing species

that outcompete SCFA producers, reducing community-level yields. Conversely, certain

commonly used additives in poultry production (e.g., riboflavin[120], several

mannooligosaccharides[115,122]) were unexpectedly predicted to decrease SCFA production

across samples. Polysaccharide chain length also influenced SCFA production, with longer

lengths of starch and mannooligosaccharides decreasing SCFA production. This is likely a

consequence of longer chains requiring more specialized or complete sets of degradation

enzymes, which are encoded by fewer bacterial genomes in the community.

To identify promising candidates for potential application in the poultry industry, we focused

on supplements that consistently increased SCFA production. Across compartments and

diets, three supplements, namely, cellulose, starch, and L-threonine, were predicted to

consistently enhance SCFA production (**Figures 3A,C; Figure S4B**). Cellulose, a complex

polysaccharide, serves as a dietary fiber in poultry nutrition[39,40] and is primarily fermented to

produce SCFAs by cellulolytic bacteria such as Ruminococcaceae, Lachnospiraceae, and

other Clostridiales[123]. Starch, a readily fermentable carbohydrate, is a major energy source for

both the host and its gut microbiota. The digestion of starch starts in the small intestine, and a

portion reaches the lower GIT, where bacterial enzymes degrade it into fermentable

oligosaccharides and monosaccharides, subsequently converted to SCFAs through microbial

fermentation pathways. L-threonine is an essential amino acid for chickens[124]. While

primarily absorbed in the small intestine, a fraction of dietary L-threonine can reach the

cecum and colon, where it serves as a substrate for bacterial species encoding threonine

degradation pathways, yielding acetate and propionate as byproducts[125–127]. Based on their

reproducible *in silico* effects across diverse microbial communities within the reference

dataset, we selected these three candidates for validation in chicken feeding trials, testing two

dietary doses of each to determine whether predicted metabolic benefits translated *in vivo*.

**Context-specific community models improved prediction accuracy**

To evaluate the predictive capacity of our modeling framework, we performed a trial
investigating the impact of two doses each of cellulose, starch, and L-threonine
supplementation in broilers fed a corn-based diet. Ileal and cecal samples were collected at
day 14 for 16S rRNA sequencing and targeted metabolomics (see Methods). Comparisons
between predicted and experimentally derived metabolite concentrations revealed a spectrum
of correlations (**Figure 4A**, upper panel). Among the metabolites tested, butyrate and
fumarate concentration fold changes showed the highest agreement between predictions and
experimental measurements (Spearman's $\rho$=0.49 for both; **Figure 4B**). In contrast, acetate
(Spearman's $\rho$=-0.14), succinate (Spearman's $\rho$=0.09), and $\alpha$-ketoglutarate (Spearman's $\rho$=-
0.2) did not correlate with predictions, while lactate predictions were inversely correlated
with trial measurements (Spearman's $\rho$=-1.0, q=0.039). A potential confounder in these
correlations is the microbial community composition, which differed between the validation
trial (conducted in Guelph, Ontario) and the original trial used to construct the six-
compartment models (conducted in Edmonton, Alberta). PCoA of cecal 16S rRNA profiles
revealed distinct clustering of the trial-derived communities compared to both the original
full-resolution dataset and the simplified composition used in the 6-compartment model
(**Figure 4B**). PERMANOVA analysis confirmed statistically significant differences in
community composition ($p < 0.05$) across both treatment groups and data sources, indicating
substantial shifts in community structure between cohorts.
354
These compositional differences reflected distinct taxonomic distributions between trials
(**Figures S1, S6**). For example, Lactobacillaceae were more abundant in the original
reference dataset (average 5.3 ± 3.9%, **Figure S1**) compared to trial communities (average
0.51 ± 1.88%, **Figure S6**) in cecal samples, contributing to predicted lactate accumulation in
the original model and likely explaining the negative correlation for lactate. Notably,
Enterobacteriaceae dominated the cecum in the reference dataset (average 21.3 ± 7.8%) but
were nearly absent in trial communities (average 0.85 ± 1.8%). As facultative anaerobes,
Enterobacteriaceae, mainly represented by *E. coli* in our models, produce acetate, succinate,
and formate as major end products (**Figure 2A;Figure S8**) and are known to mediate cross-
feeding networks in cecal communities[25]. The combined depletion of these families could
alter both metabolite production and metabolic interactions, potentially explaining poor
predictions for acetate and succinate. Conversely, Ruminococcaceae expanded from 4.2 ±
3.0% (reference) to 29.4 ± 8.4% (trial) in the cecal communities. This family comprises
obligate anaerobic fiber degraders known to produce butyrate, including *Faecalibacterium*, a

key butyrate producer in our models (**Figure 2A;Figure S8**). This may explain why butyrate

predictions maintained reasonable correlation (Spearman's $\rho=0.49$) despite the community

restructuring. These findings demonstrated that while metabolic predictions were robust to

variation within a reference cohort (**Figure 3**), community composition differences between

facilities impacted prediction accuracy for certain metabolites, highlighting the need for

context-specific modeling.

To address this limitation, we constructed a two-compartment model (reflecting the two

compartments sampled in the validation trial – ileum and cecum) based on communities from

the trial 16S profiles (see Methods). Trial-informed models were found to increase prediction

accuracy across multiple metabolites (**Figure 4A**, lower panels). For instance, propionate

correlation increased substantially from Spearman's $\rho=0.31$ to $\rho=0.94$ (**Figure 4A**), aligning

with elevated concentrations in the low-starch treatment group where *Bacteroides* were

uniquely detected (**Figure S6**). Similarly, predictions for succinate improved from

Spearman's $\rho=0.09$ to $\rho=0.71$, and α-ketoglutarate from Spearman's $\rho=-0.2$ to $\rho=0.6$.

Butyrate correlation showed modest improvement (Spearman's $\rho=0.49$ to $\rho=0.54$), while

lactate predictions shifted from significant inverse correlation (Spearman's $\rho=-1.0$, $q=0.03$) to

a more moderate mismatch (Spearman's $\rho=-0.31$), consistent with reduced Lactobacillaceae

abundance in trial-derived inputs. However, even with trial-specific community data, the

models failed to accurately predict certain metabolites, revealing limitations beyond

community composition effects. Acetate, which is broadly produced and absorbed throughout

the gut, remained poorly predicted (Spearman's $\rho=-0.14$ to $\rho=-0.09$), a pattern consistent with

acetate prediction challenges reported in previous community modeling studies[128]. Fumarate

correlation declined (Spearman's $\rho=0.49$ to $\rho=-0.60$), with measured concentrations showing

consistent depletion across all dietary treatments (ileum Kruskal-Wallis (KW) $p=0.004$,

cecum KW $p=0.002$, significant post-hoc Dunn and Wilcoxon rank-sum tests, **Figure S7AB**),

despite predicted net microbial production suggesting rapid microbial consumption or host

absorption not captured in the model. In contrast, butyrate increases under low-dose cellulose

and starch supplementation aligned well with predicted production by members of

Ruminococcaceae, Butyricicoccaceae, and Lachnospiraceae families (cecum KW $p=0.029$),

demonstrating accurate predictions when key producer taxa were present (**Figure S8**).

We note that high inter-individual variability and small sample sizes (n=6 per treatment)

limited detection of statistically significant patterns for several metabolites, though trial-

informed models consistently improved rank-order agreement with experimental trends.
Together, these results demonstrate that community-matched metabolic models can
reproduce *in vivo* fermentation patterns when grounded in appropriate taxonomic inputs,
though host metabolic processes and biological variability remain important sources of
prediction uncertainty.
408

## DISCUSSION

Optimizing poultry nutrition to promote gut health requires an understanding of how dietary inputs are transformed into metabolic outputs across the gastrointestinal tract. We developed a spatiotemporally resolved computational framework that simulates microbial community metabolism throughout the chicken GIT. This framework represents the most comprehensive spatial characterizations of gut metabolism in any livestock system performed to date. Our multi-compartment model captures three critical features: 1) spatial organization across the entire GIT through the simulation of six connected compartments; 2) temporal dynamics through inclusion of feeding-fasting cycles; and 3) the bidirectional flow of microbes and metabolites between neighbouring compartments through peristalsis and reverse peristalsis. Together these features represent a significant advance over previous metabolic modeling efforts in poultry that focused specifically on cecal metabolism[24,25]. By incorporating five additional compartments (gizzard, duodenum, jejunum, ileum, and colon), our framework treats the cecum as part of an interconnected system where upstream metabolic interconversions reshape substrate availability in the lower GIT and downstream metabolism. Our simulations revealed that metabolite landscapes and production rates vary substantially across compartments and over time – patterns that single-compartment models cannot capture. Upper gut compartments displayed distinct metabolic profiles enriched for bile salt hydrolases, membrane lipid synthesis, and fatty acid biosynthesis, while cecal and colonic communities specialized in SCFA synthesis pathways and polysaccharide degradation. The unprocessed diet remained distinct from metabolic profiles of all gut compartments throughout simulations, demonstrating that microbial communities extensively transform dietary inputs rather than simply passing compounds unchanged. Importantly, overnight reflux redistributes metabolites and helps explain convergences in metabolite profiles between adjacent regions.

Recent advances in mammalian gut modeling, including multi-compartment frameworks in mice[17,129], human community-level FBA approaches[12,16,130], and the CODY model of the human colon[21], have demonstrated the value of spatially explicit and physiologically informed models for understanding host-microbiome metabolism. Our work builds on these concepts and extends them to poultry, where avian-specific physiological features, such as rapid upper GIT transit, steep oxygen gradients, and reverse peristalsis, play a defining role in nutrient processing and microbial ecology. Incorporating these traits allowed us to capture

metabolite redistribution and temporal oscillations characteristic of the chicken gut. By

modeling taxa individually rather than as a single metabolic entity, we predicted taxon-

specific contributions to SCFA production: *Bacteroides* as the primary propionate producers,

*Faecalibacterium* and *Clostridium* spp. as key butyrate producers, and Enterobacteriaceae

and Lachnospiraceae as major acetate producers. Our study showed that the upper GIT

(gizzard, duodenum and jejunum) communities were dominated by facultative anaerobes

encoding fewer metabolic enzymes and enriched for bile salt hydrolases and fatty acid

biosynthesis, reflecting adaptation to readily available nutrients under aerobic conditions. The

ileum harbored mixed aerobic and anaerobic taxa, reflecting its transitional position between

upper and lower GIT, while cecal and colonic communities were enriched in Clostridiales

families with the capacity to ferment complex polysaccharides. Through the inclusion of

spatial and temporal dynamics, our model captured metabolite accumulation associated with

fasting and reflux. For example, fasting was associated with an increase in the production of

SCFAs in the lower GIT (cecum and colon) as well as signaling molecules such as GABA in

the upper GIT. These observations indicate that the timing of sample collection has the

potential to substantially influence metabolite levels, consistent with recent studies showing

that feeding cycles drive dramatic differences in metabolic measurements[111].

Having established a spatiotemporally resolved framework, we leveraged its predictive

capabilities to systematically screen dietary interventions. Simulations involving 34 candidate

metabolite supplements revealed that their impacts were largely context-dependent, with

metabolic responses varying by substrate type, concentration, polymer chain length,

microbial community composition, and gut compartment. Some additives commonly used in

poultry production (riboflavin, mannooligosaccharides) were unexpectedly predicted to

decrease SCFA production, while others showed opposite dose-dependent effects (e.g.,

maltoheptaose, pantothenic acid). However, our model predicted three compounds –

cellulose, starch, and L-threonine – to consistently enhance SCFA production across diets and

compartments. The robustness of these predicted effects despite high inter-individual

microbiome variation were tested in a poultry feed trial, enabling direct assessment of our

model's prediction accuracy.

Our experimental validation through a controlled feeding trial yielded critical insights into the

determinants of predictive accuracy in community metabolic modeling. Butyrate responses to

cellulose and starch supplementation aligned with model predictions, while other metabolites

showed variable agreement with experimental measurements. For example, lactate

accumulated in simulations but not *in vivo*, which reflected the much lower abundance of

Lactobacillaceae in the trial cohort compared to the reference community used for model

construction. Similarly, Enterobacteriaceae abundances were substantially lower in trial birds,

while Ruminococcaceae increased 7-fold relative to the reference dataset, which could

disrupt prediction for metabolites predominantly produced by specific taxa belonging to these

families (e.g., acetate, butyrate). Since one of our main aims was to identify supplements that

led to robust changes in SCFA production irrespective of community composition, this

finding suggests that predictions may have limited relevance for poultry raised under

different conditions and/or locales. By incorporating 16S rDNA survey data collected from

the ileum and ceca during the trial within a more simplified two-compartment model,

prediction accuracy improved, particularly for propionate, succinate, and α-ketoglutarate

(Spearman's ρ=0.31→0.94; ρ=0.09→0.71; and ρ=-0.20→0.60, respectively). This

improvement underscores taxonomic structure as a primary determinant of metabolic

function, aligning with recent observations that community composition drives differences in

SCFA production[16]. At the same time, we note that even trial-informed model could not

predict acetate (Spearman's ρ=-0.09), consistent with challenges reported in other community

modeling studies, or fumarate (Spearman's ρ=-0.60), which showed consistent experimental

depletion despite predicted microbial production, likely reflecting rapid host absorption not

captured by our simplified fractional removal rates. This suggests future models could be

improved through an improved understanding of host-metabolite absorption. These findings

highlight that while metabolic predictions were robust to dietary and inter-individual

variation within a reference cohort, cross-facility differences in community composition

significantly impact accuracy, and host metabolic processes remain important sources of

uncertainty.

Implementation-specific limitations are more readily addressable as data availability

improves. The model's reliance on automated genome-scale reconstructions means that

enzymatic capabilities may be incomplete for underrepresented taxa. Our framework

currently models only bacteria, excluding fungi and other microorganisms that may

contribute to gut metabolism. The model assumes uniform distribution and mixing within the

compartments, which may not accurately reflect the spatial microenvironments where

localized metabolite gradients shape community interactions. Estimated rates of nutrient

absorption and gut transit show inter-individual variation that our population-averaged

parameters cannot capture. Additionally, the absence of host metabolic networks means we
cannot represent how host processes (e.g., immune responses) affect microbial metabolism.
Nevertheless, this modeling framework is adaptable and can readily accommodate future
advances in microbial genomics and metabolic modeling. As new high-quality poultry MAGs
and physiological measurements become available, they can be readily incorporated in future
models. Incorporating gene and protein expression (metatranscriptomic and proteomic data)
will further improve the accuracy of individual reconstructions, while coupling microbial
models with host metabolic networks would improve representation of host absorption and
host-microbiome metabolic exchange.

While our six-compartment model represents a significant step forward, it serves as a proof-
of-concept for systematic modeling of the poultry GIT. By bridging *in silico* predictions with
controlled feeding trials, we demonstrate both the promise and the limitations of current
metabolic modeling approaches. This work lays the foundation for evaluating dietary
interventions in poultry, guiding the design of targeted supplements that enhance SCFA
production and gut health. More broadly, it illustrates how multi-compartment community
models can accelerate hypothesis generation, reduce reliance on large-scale animal trials, and
extend to other host-associated microbiomes where spatial structure and host physiology play
decisive roles.

# METHODS

## Development of a six-compartment spatiotemporal model
### Processing of 16S rRNA gene data
Reads were processed using the QIIME2[32] package (version 2023.2), which involved
trimming, filtering, and removing low-quality and chimeric sequences. The resulting high-
quality data was then processed following the deblur pipeline. Sequences were clustered into
amplicon sequence variant (ASV) groups and singleton sequences removed. Taxonomic
classification was performed using the classify-hybrid-vsearch-sklearn function against the
SILVA database (release 138.1)[32,33]. ASVs with an abundance less than 0.01% were removed
to reduce the potential for observing bleed-through ASVs. ASVs identified as chloroplast or
mitochondrial contaminants were also removed, and ASV tables were subsequently

normalized and rarefied to a depth of 10,000 reads per sample to ensure comparability across

datasets.

**Selection of representative species and model reconstruction**

To represent the most abundant taxa within each gut compartment, the top five ASVs per

bacterial family were used as blastn queries against several collections of high-quality

chicken gut-derived MAGs: a set of 12,339 MAGs from integrated 799 public chicken gut

microbiome samples[34], 469 MAGs from 24 chicken cecum samples[35], and 461 MAG from

five chicken gut compartments of 30 chickens[36]. Sequences meeting thresholds of 97%

identity, e-value threshold of 1e-70, as well as passing assembly quality thresholds of ≥90%

completeness, and ≤5% contamination were retained for further analysis. For ASVs that did

not yield any MAG matches meeting these criteria, the 16S rRNA sequence was subjected to

a blastn search against the NCBI RefSeq 16S database[37]. The highest scoring sequence

alignments were then evaluated based on three criteria: (i) presence in chicken gut-derived

MAG collections, (ii) prior documentation in poultry microbiome studies, and (iii)

transcriptional activity confirmed by metatranscriptomic data from the same reference trial[9].

MAGs for taxa meeting these criteria and representing the most prevalent members of their

genus in the chicken gut were selected for downstream metabolic model reconstruction.

Complete details of genome sources, genome quality metrics, and taxonomic classifications

for all selected MAGs are provided in Supplementary **Table S1**. Gapseq version 1.2[38] with

default settings was used to reconstruct genome-scale metabolic models. Draft metabolic

models were curated manually to improve representation of carbohydrate metabolism (see

Supplementary Methods).

**Construction of sample-specific gut microbiota community models**

Compartment-specific models were constructed for 60 samples (2 diets × 5 birds × 6

compartments) using the list of mapped taxa and their family-level abundances from 16S

data. Representative species models were combined according to compartment-specific

abundance profiles. In cases where sequencing data were unavailable for one compartment

within a sample, the missing values were imputed using average relative abundances from the

same dietary group and compartment. Samples missing data for more than one compartment

were excluded.

**Simulation framework and gastrointestinal configuration**

All microbial community simulations were conducted using BacArena, an R-based platform

for community modeling[13]. Bacteria were represented as individual agents occupying a 100 ×

100 grid (10,000 cells total), where each cell could grow, move, and exchange metabolites

with the surrounding environment. Growth optimization was performed for each individual at

every time step using the microbial biomass reaction as an objective. Simulations were run

for 96 hours (six-compartment model) or 48 hours (two-compartment model), each consisting

of a 16-hour daytime phase (representing daylight exposure and feeding) followed by an

eight-hour nighttime phase (mimicking sleeping and fasting conditions). To account for

stochasticity in bacterial placement, movement, and replication, each bird-specific model was

simulated with 5 replicates using different random seeds. The simulations were conducted

with a time step of 1 hour, representing an hour of bacterial growth per iteration. To

incorporate physiological factors, including absorption, peristalsis, reflux (reverse peristalsis),

feeding, excretion, pH and oxygen homeostasis, the simulation environment was updated

every hour (see more details in the **Supplementary Methods**). After each hour of simulation,

these processes were executed to adjust the environment accordingly before proceeding with

the simulation. The relevant parameters used during simulation initialization were defined for

each compartment based on literature sources (**Table S2A**).

**Comparative analysis of the functional capabilities of community models**

To compare the metabolic capabilities across different gut compartments and samples, I first

extracted the complete set of reactions from their respective community metabolic models.

To quantify the metabolic potential within each sample, accounting for the varying

contributions of different species, I summarized the reaction counts within each sample,

where each reaction count was weighted by the relative abundance of the species contributing

to that reaction. The resulting incidence matrices contained information on the reaction

abundance in each community. Bray-Curtis dissimilarity matrices were constructed from the

incidence matrices described using *vegdist* function from the vegan R-package, and Principal

Coordinate Analysis was performed using *cmdscale* function from stats R-package.

**Comparison of control simulations and simulations with supplementation**

To evaluate the effects of different compounds on community fluxes and metabolite

concentrations, fold changes were calculated for each combination of gut compartment,

metabolite, and supplemented compound by dividing the metabolite concentration or

community flux in the experimental group by the mean concentration or flux of 5 replicates

from the control group. Additionally, given the experimental design, where validation is

performed through chicken trials using 10 chickens per treatment, I subsampled 10 replicates

from the original 25 by randomly selecting 2 out of 5 replicates from each of the 5 samples

per diet. This subsampling approach resulting in 10 subsampled replicates per condition was
designed to match the number of biological replicates to be used in the experimental
validation and maintain the integrity of the statistical comparisons. Significant differences
were determined by Wilcoxon signed-rank tests, that were conducted on the fold changes
under the null hypothesis that the median fold change is equal to 1 (indicating no effect of the
treatment). P-values were calculated using a two-sided test, and to account for the multiple
comparisons inherent in this analysis, the Benjamini-Hochberg procedure was applied to
control the False Discovery Rate (FDR). Results were considered statistically significant at
an FDR-adjusted p-value threshold of 0.05.

## Experimental validation and trial-informed model refinement

*In vivo* **validation through chicken trials**

While *in silico* simulations demonstrated significant effects of supplementation over 96
hours, the translation between *in silico* time units and actual community dynamics *in vivo* is
not direct. Microbial growth rates, metabolic activities, and community interactions in the
living gut are influenced by multiple physiological factors not fully captured in the model.
Therefore, to ensure sufficient time for supplement-induced changes to manifest in the gut
microbiome, a longer supplementation period was implemented in the validation
experiments.

Seventy one-day-old broiler chickens were raised on standard corn-based starter diet for 7
days before random assignment to seven treatment groups (n=10 per group): control (G1),
cellulose at 2g/100g (G2) and 8g/100g (G3), L-threonine at 1g/100g (G4) and 5g/100g (G5),
and starch at 5g/100g (G6) and 10g/100g (G7). Supplementation lasted 7 days, birds were
euthanized on day 14 and samples were collected from ileum and cecum, stored on dry ice,
and then kept at −80 °C for microbiome analysis. All animal experiments were approved by
the Animal Care Committee of the University of Guelph and adhered to the guidelines for the
use of animals (Animal Utilization Protocol #4815).

### DNA extraction

DNA was extracted from 10 mg of chicken gut samples (ceca and ileum) using the
ZymoBIOMICS DNA miniprep kit with the addition of the ZR BashingBead (2.0 mm) tubes
to increase homogenization and lysis of the samples. The extraction protocol was followed as
indicated in the manual with a final elution volume of 50uL. 5 µL of the ZymoBIOMICS
Spike-In Control I (D6320) was added to ileum samples and 10 µL to caeca. Positive (Zymo

Microbial Community Standard) and negative (no input) extraction controls were run
alongside the sample extractions.

**16S rRNA gene sequencing and processing**

The V4 hypervariable region of the 16S rRNA gene was amplified using uniquely barcoded
515F (forward) and 806R (reverse) sequencing primers to allow for multiplexing[41].
Amplification reactions were performed using 12.5 µL of KAPA2G Robust HotStart
ReadyMix (KAPA Biosystems), 1.5 µL of 10 µM forward and reverse primers, 7.5 µL of
sterile water and 2 µL of DNA. The V4 region was amplified by cycling the reaction at 95°C
for 3 minutes, 18x cycles of 95°C for 15 seconds, 50°C for 15 seconds and 72°C for 15
seconds, followed by a 5-minute 72°C extension. All amplification reactions were done in
duplicate to reduce amplification bias, pooled, and checked on a 1% agarose TBE gel. Pooled
duplicates were quantified using PicoGreen and combined by even concentrations. The
library was then purified using Ampure XP beads and loaded on to the Illumina MiSeq for
sequencing, according to manufacturer instructions (Illumina, San Diego, CA). Sequencing is
performed using the V2 (150bp x 2) chemistry. A single-species (*Pseudomonas aeruginosa*
DNA), a mock community (Zymo Microbial Community DNA Standard) and a template-free
negative control were included in the sequencing runs. Sequencing data were processed using
the identical QIIME2 pipeline detailed in section "Processing of 16S rRNA gene data" above,
ensuring direct comparability between reference and trial datasets.

**Targeted quantitative metabolomics data processing and analysis**

A targeted quantitative metabolomics workflow was performed using a custom reverse-phase
LC-MS/MS assay on an AB Sciex 5500 QTRAP® (Applied Biosystems/MDS Sciex)
equipped with an Agilent 1290 UHPLC. Metabolites were identified and quantified by
multiple reaction monitoring using isotope-labeled and other internal standards. The assay
was prepared in a 96-deep-well plate format with an attached filter plate; the first 14 wells
were allocated to one blank, three zero samples, seven calibration standards, and three quality
control samples.

For all metabolites except organic acids, samples were thawed on ice and extracted. Extracts
were loaded onto the filter plate, dried under nitrogen, derivatized with phenyl-
isothiocyanate, incubated, and dried again. Metabolites were then extracted with 300 µL
extraction solvent and collected by centrifugation into the lower deep-well plate, followed by
dilution with MS running solvent. For organic acids, 150 µL ice-cold methanol and 10 µL
isotope-labeled internal standard mix were added to extracted samples for overnight protein

precipitation. Samples were centrifuged at 13,000×g for 20 min, and 50 μL supernatant was

loaded into a 96-deep-well plate. 3-nitrophenylhydrazine reagent was added, followed by 2h

incubation; BHT stabilizer and water were then added prior to LC-MS injection. Samples

were analyzed by an LC method followed by a direct injection method. Data were acquired

and processed in Analyst v1.6.3.

Dry-weight normalized concentrations were processed using MetaboAnalyst web platform[42].

Quality control filtering removed low-variance features (interquartile range < 5th percentile)

and low-abundance features (median intensity < 5th percentile). The filtered dataset was

normalized using probabilistic quotient normalization (PQN) with the control group (G1) as

reference to account for dilution effects and technical variation. Fold changes for each

metabolite were calculated relative to the control group median within each compartment.

Statistical comparisons between treatment and control groups employed two-sample

Wilcoxon tests with Benjamini-Hochberg FDR correction. To evaluate agreement between

model predictions and experimental results, Spearman's rank correlation coefficients were

computed across all metabolites and treatment conditions, comparing predicted and observed

fold changes within each compartment.

**Construction of trial-informed 2-compartment models**

From 16S rRNA data of 112 samples, 694 ASVs were filtered to retain those with ≥0.5%

relative abundance, reducing sequencing noise. Functional profiles were predicted using

PICRUSt2[43], and ASVs were clustered by enzymatic similarity (Jaccard distance, 0.85

threshold), yielding 267 functionally distinct representatives. These were matched to high-

quality MAGs through blastn-search against four chicken MAG catalogues: 461 MAG from

five chicken gut compartments of 30 chickens[36], MGnify chicken-gut v1.0 collection (13,386

genomes)[44], a set of 12,339 MAGs from integrated 799 public chicken gut microbiome

samples[34], and 461 MAG from five chicken gut compartments of 30 chickens[36]. Matching

was performed in three sequential passes on unmatched ASVs: (i) 97% sequence identity

threshold with ≥80% query coverage; (ii) 95% identity with ≥70% coverage; (iii) 90%

identity with ≥70% coverage. For each BLAST hit, a combined quality score was computed

incorporating both sequence similarity (60% weight) and assembly quality (40% weight),

where: sequence similarity = 0.4*identity + 0.3*coverage + 0.3*min[100,100×bitscore/500]),

assembly quality = completeness - 5*contamination. Final selection included top-scoring hits

per ASV, resulting in 170 unique MAGs. Complete details of genome sources, genome

quality metrics, and taxonomic classifications for all selected MAGs are provided in

Supplementary **Table S3**. Metabolic models were reconstructed using gapseq version 1.4.0[38]

with default settings for 44 bird pairs with sufficient data quality.

Bird-specific models paired ileal and cecal communities based on sample-derived ASV

abundances, creating 88 compartment-specific models (44 birds $\times$ 2 compartments). Model

assembly followed the same procedure as for the multi-compartment system, ensuring direct

comparability.

**Definition of *in silico* dietary input**

Nutritional information from the corn-based grower diet was adjusted to account for upstream

digestion and absorption to create an *in silico* representation of dietary metabolites reaching

the distal small intestine. The base diet composition was derived from the same poultry feed

tables used in the 6-compartment model, with modifications to simulate the metabolic

environment of the distal jejunum/proximal ileum. Metabolite concentrations entering the

ileum were reduced based on published ileal digestibility coefficients for corn-based poultry

diets. Mineral retention rates were set according to jejunal and ileal digestibility studies[45–51].

Similarly, carbohydrate retention reflected measurements on upper small intestine

digestion[52–54], while amino acids were retained at 25% based on typical ileal digestibility[52,55].

Fatty acid retention varied by saturation level[56], and cellulose was retained at 50% to reflect

partial fermentation in the upper gut. Essential metabolites (host-supplied metabolites, such

as mucins, urea, and metabolites supporting growth of all community members) were added

at baseline concentrations. The composition of the *in silico* diets used is available in the

**Table S4**.

**Design and simulation of a trial-informed 2-compartment (ileum-cecum) model**

The design and configuration were identical to the six-compartment GIT simulation

framework (described in the **Supplementary Methods**) with all the processes applicable to

ileum and cecum implemented in the same way, except the reflux happening overnight. In the

simplified 2-compartment model, reverse peristalsis was not considered due to the lack of

upstream compartments and colon, involved in the process.

To maintain realistic metabolic growth rates, we implemented cooperative trade-off FBA

(ctFBA) with L2 regularization, adapted from the MICOM framework[12]. This approach

balances individual species growth rates against community-level biomass production

through a two-step optimization. First, the maximum total community biomass production

rate is determined, then the individual species growth rates are optimized while constraining

total community growth to a fraction ($\tau$, trade-off parameter) of the maximum, using L2

regularization to minimize growth rate variance. The trade-off parameter ($\tau$) was calibrated

independently for each compartment to ensure at least 90% of species achieved positive

growth rates while maintaining physiologically realistic maximum growth rates (<3 h⁻¹ in

the ileum, <5 h⁻¹ in the cecum). Calibration was performed at simulation initialization using

the seeded microbial community and initial medium conditions, similar to MICOM's

approach.

The relevant parameters used during simulation initialization were defined for each

compartment based on literature sources (**Table S2B**).

**Simulation of dietary supplementation for validation experiments**

To simulate dietary supplementation treatments, supplement concentrations entering the

ileum were adjusted to account for upstream absorption and saturation effects. Cellulose

supplementation at 2g/100g (G2) and 8g/100g (G3) was modeled as 1g/100g and 4g/100g

reaching the ileum, maintaining an approximately 50% passage rate. L-Threonine

supplementation at 1g/100g (G4) and 5g/100g (G5) was modeled as 0.4 g/100g and 2.5

g/100g reaching the ileum. While 40-60% of dietary threonine undergoes first-pass gut

extraction, high doses saturate intestinal absorption, allowing greater proportions to reach the

distal ileum. Starch supplementation at 5g/100g (G6) and 10g/100g (G7) in feed was modeled

as 1.25g/100g and 2.0g/100g reaching the ileum, respectively, reflecting saturation of glucose

absorption in the proximal small intestine at higher doses. These adjustments ensured

physiologically realistic ileal concentrations while maintaining experimental dose-response

relationships.

**DATA AVAILABILITY**

For the 6-compartment model, 16S rDNA sequence datasets from a previous trial examining the impact of AGPs and diet on the poultry gut microbiome were downloaded from the SRA (project PRJNA614900). The previous study[9] details the animal trial conditions, sequencing data retrieval, and 16S rDNA sequencing. The 16S rDNA sequence datasets generated in this study from the feeding trial are deposited under the BioProject PRJNA1403483.

**CODE AVAILABILITY**

Scripts, source data and metabolic models required to reproduce the analyses of this study can be accessed through GitHub at https://github.com/ParkinsonLab/ChickenGIT_modeling

# ETHICS DECLARATIONS

**COMPETING INTERESTS**

The authors declare that they have no competing interests.

**FUNDING**

This work was funded by grants from the Natural Sciences and Engineering Research Council of Canada to J.P. (RGPIN-2019-06852/RGPIN-2025-06006), and Ontario Research Fund-Research Excellence to J.P. and S.S. High-performance computing was provided by the SciNet HPC Consortium; SciNet is funded by the following: the Canada Foundation for Innovation under the auspices of Digital Alliance Canada, the Government of Ontario, Ontario Research Fund–Research Excellence, and the University of Toronto. The funders had no role in the design of the study, collection of data and analysis, preparation of the manuscript, and decision to publish.

**AKNOWLEDGEMENTS**

We thank the Centre for the Analysis of Genome Evolution & Function (CAGEF) at the University of Toronto for 16S rRNA amplicon sequencing services and The Metabolomics Innovation Centre (TMIC) for metabolomics profiling services.

**AUTHORS' CONTRIBUTIONS**

J.P. and I.U. conceived and designed the study. I.U., J.P., M.A. and S.S. designed the validation chicken trials; M.A. and S.S. conducted the chicken feeding trials, obtained regulatory approvals, and collected intestinal samples. I.U. performed metabolic model

792    reconstruction and curation, simulations, data analyses, and visualization. J.P. and I.U wrote

793    the manuscript. All authors reviewed and/or edited the manuscript.

# REFERENCES

1. Dibner, J. J. & Richards, J. D. Antibiotic growth promoters in agriculture: History and mode of action. *Poult. Sci.* **84**, 634–643 (2005).

2. Zhao, C., Wang, Y., Mulchandani, R. & Van Boeckel, T. P. Global surveillance of antimicrobial resistance in food animals using priority drugs maps. *Nat. Commun.* **15**, (2024).

3. European Union. Ban on antibiotics as growth promoters in animal feed enters into effect. *Regulation* (2006).

4. Antimicrobial consumption and resistance in bacteria from humans and food-producing animals: Fourth joint inter-agency report on integrated analysis of antimicrobial agent consumption and occurrence of antimicrobial resistance in bacteria from humans and food-producing animals in the EU/EEA JIACRA IV – 2019–2021. *EFSA Journal* **22**, (2024).

5. Singer, R. S. *et al.* Raising Animals Without Antibiotics: U.S. Producer and Veterinarian Experiences and Opinions. *Front. Vet. Sci.* **6**, (2019).

6. Acosta, A. *et al.* The future of antibiotic use in livestock. *Nature Communications* **16**, (2025).

7. Rahman, M. R. T., Fliss, I. & Biron, E. Insights in the Development and Uses of Alternatives to Antibiotic Growth Promoters in Poultry and Swine Production. *Antibiotics* vol. 11 Preprint at https://doi.org/10.3390/antibiotics11060766 (2022).

8. Patience, J. F. & Ramirez, A. Invited review: strategic adoption of antibiotic-free pork production: the importance of a holistic approach. *Translational Animal Science* vol. 6 Preprint at https://doi.org/10.1093/tas/txac063 (2022).

9. Zou, A. *et al.* Systematic profiling of the chicken gut microbiome reveals dietary supplementation with antibiotics alters expression of multiple microbial pathways with minimal impact on community structure. *Microbiome* **10**, (2022).

10. Orth, J. D., Thiele, I. & Palsson, B. Ø. What is flux balance analysis? *Nat. Biotechnol.* **28**, 245–248 (2010).

11. O'Brien, E. J., Monk, J. M. & Palsson, B. O. Using genome-scale models to predict biological capabilities. *Cell* vol. 161 Preprint at https://doi.org/10.1016/j.cell.2015.05.019 (2015).

12. Diener, C., Gibbons, S. M. & Resendis-Antonio, O. MICOM: Metagenome-Scale Modeling To Infer Metabolic Interactions in the Gut Microbiota. *mSystems* **5**, (2020).

828    13.    Bauer, E., Zimmermann, J., Baldini, F., Thiele, I. & Kaleta, C. BacArena: Individual-
829           based metabolic modeling of heterogeneous microbes in complex communities. *PLoS*
830           *Comput. Biol.* **13**, (2017).

831    14.    Bauer, E. & Thiele, I. From metagenomic data to personalized in silico microbiotas：
832           predicting dietary supplements for Crohn ' s disease. *NPJ Syst. Biol. Appl.* **4**, (2018).

833    15.    Mukhopadhya, I. & Louis, P. Gut microbiota-derived short-chain fatty acids and their
834           role in human health and disease. *Nature Reviews Microbiology* vol. 23 Preprint at
835           https://doi.org/10.1038/s41579-025-01183-w (2025).

836    16.    Quinn-Bohmann, N. *et al.* Microbial community-scale metabolic modelling predicts
837           personalized short-chain fatty acid production profiles in the human gut. *Nat.*
838           *Microbiol.* **9**, 1700–1712 (2024).

839    17.    Best, L. *et al.* Metabolic modelling reveals the aging-associated decline of host–
840           microbiome metabolic interactions in mice. *Nat. Microbiol.* **10**, 973–991 (2025).

841    18.    Heinken, A. *et al.* A genome-scale metabolic reconstruction resource of 247,092
842           diverse human microbes spanning multiple continents, age groups, and body sites. *Cell*
843           *Syst.* **16**, (2025).

844    19.    Marinos, G., Zimmermann, J., Taubenheim, J. & Kaleta, C. Virtual Colon:
845           Spatiotemporal modelling of metabolic interactions in a computational colonic
846           environment. Preprint at https://doi.org/10.1101/2024.06.11.598488 (2024).

847    20.    Versluis, D. M. *et al.* A Multiscale Spatiotemporal Model Including a Switch from
848           Aerobic to Anaerobic Metabolism Reproduces Succession in the Early Infant Gut
849           Microbiota. *mSystems* **7**, (2022).

850    21.    Geng, J., Boyang, J., Li, G., López-Isunza, F. & Nielsen, J. CODY enables
851           quantitatively spatiotemporal predictions on in vivo gut microbial variability induced
852           by diet intervention. *Syst. Biol. (Stevenage).* **118**, (2021).

853    22.    Javad, A.-D., Andrew, D., Amin, V. & K, M. M. R. MetaBiome: a multiscale model
854           integrating agent-based and metabolic networks to reveal spatial regulation in gut
855           mucosal microbial communities. *mSystems* **10**, e01652-24 (2025).

856    23.    Chan, Friedman, Wu & Maranas. Predicting the Longitudinally and Radially Varying
857           Gut Microbiota Composition Using Multi-Scale Microbial Metabolic Modeling.
858           *Processes* **7**, 394 (2019).

859    24.    Plata, G. *et al.* Growth promotion and antibiotic induced metabolic shifts in the
860           chicken gut microbiome. *Commun. Biol.* **5**, (2022).

861    25.    Utkina, I., Fan, Y., Willing, B. P. & Parkinson, J. Metabolic modeling of microbial
862            communities in the chicken ceca reveals a landscape of competition and co-operation.
863            *Microbiome* **13**, 248 (2025).

864    26.    Al-Nijir, M., Chuck, C. J., Bedford, M. R. & Henk, D. A. Metabolic modelling
865            uncovers the complex interplay between fungal probiotics, poultry microbiomes, and
866            diet. *Microbiome* **12**, (2024).

867    27.    Taverniti, V. *et al.* Probiotics Modulate Mouse Gut Microbiota and Influence Intestinal
868            Immune and Serotonergic Gene Expression in a Site-Specific Fashion. *Front.*
869            *Microbiol.* **12**, (2021).

870    28.    Jensen, B. A. H. *et al.* Small intestine vs. colon ecology and physiology: Why it
871            matters in probiotic administration. *Cell Reports Medicine* vol. 4 Preprint at
872            https://doi.org/10.1016/j.xcrm.2023.101190 (2023).

873    29.    Borda-Molina, D. *et al.* Caeca Microbial Variation in Broiler Chickens as a Result of
874            Dietary Combinations Using Two Cereal Types, Supplementation of Crude Protein
875            and Sodium Butyrate. *Front. Microbiol.* **11**, (2021).

876    30.    Kim, E. *et al.* The flow of non-starch polysaccharides along the gastrointestinal tract of
877            broiler chickens fed either a wheat- or maize-based diet. *Animal Nutrition* **9**, (2022).

878    31.    Nguyen, H. T., Bedford, M. R., Wu, S. B. & Morgan, N. K. Soluble non-starch
879            polysaccharide modulates broiler gastrointestinal tract environment. *Poult. Sci.* **100**,
880            (2021).

881    32.    Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data
882            science using QIIME 2. *Nature Biotechnology* vol. 37 Preprint at
883            https://doi.org/10.1038/s41587-019-0209-9 (2019).

884    33.    Kaehler, B. D. *et al.* Species abundance information improves sequence taxonomy
885            classification accuracy. *Nat. Commun.* **10**, (2019).

886    34.    Feng, Y. *et al.* Metagenome-assembled genomes and gene catalog from the chicken
887            gut microbiome aid in deciphering antibiotic resistomes. *Commun. Biol.* **4**, (2021).

888    35.    Glendinning, L., Stewart, R. D., Pallen, M. J., Watson, K. A. & Watson, M. Assembly
889            of hundreds of novel bacterial genomes from the chicken caecum. *Genome Biol.* **21**, 1–
890            16 (2020).

891    36.    Zhang, Y. *et al.* Improved microbial genomes and gene catalog of the chicken gut from
892            metagenomic sequencing of high-fidelity long reads. *Gigascience* **11**, (2022).

893    37.    O'Leary, N. A. *et al.* Reference sequence (RefSeq) database at NCBI: Current status,
894            taxonomic expansion, and functional annotation. *Nucleic Acids Res.* **44**, (2016).

895    38.    Zimmermann, J., Kaleta, C. & Waschina, S. gapseq: informed prediction of bacterial
896            metabolic pathways and reconstruction of accurate metabolic models. *Genome Biol.*
897            **22**, (2021).

898    39.    Tejeda, O. J. & Kim, W. K. Role of dietary fiber in poultry nutrition. *Animals* vol. 11
899            Preprint at https://doi.org/10.3390/ani11020461 (2021).

900    40.    Hetland, H., Choct, M. & Svihus, B. Role of insoluble non-starch polysaccharides in
901            poultry nutrition. in *World's Poultry Science Journal* vol. 60 (2004).

902    41.    Caporaso, J. G. *et al.* QIIME allows analysis of high-throughput community
903            sequencing data. *Nature Methods* vol. 7 Preprint at https://doi.org/10.1038/nmeth.f.303
904            (2010).

905    42.    Pang, Z. *et al.* MetaboAnalyst 5.0: Narrowing the gap between raw spectra and
906            functional insights. *Nucleic Acids Res.* **49**, (2021).

907    43.    Douglas, G. M. *et al.* PICRUSt2 for prediction of metagenome functions. *Nature
908            Biotechnology* vol. 38 Preprint at https://doi.org/10.1038/s41587-020-0548-6 (2020).

909    44.    Gurbich, T. A. *et al.* MGnify Genomes: A Resource for Biome-specific Microbial
910            Genome Catalogues: MGnify biome-specific genome catalogues. *J. Mol. Biol.* **435**,
911            (2023).

912    45.    Guenter, W. & Sell, J. L. Magnesium absorption and secretion along the
913            gastrointestinal tract of the chicken. *J. Nutr.* **103**, (1973).

914    46.    David, L. S., Abdollahi, M. R., Ravindran, G., Walk, C. L. & Ravindran, V. Studies on
915            the measurement of ileal calcium digestibility of calcium sources in broiler chickens.
916            *Poult. Sci.* **98**, (2019).

917    47.    Haetinger, V. S. *et al.* Ileal phosphorus digestibility of soybean meal for broiler
918            chickens remains consistent across institutions in a collaborative study regardless of
919            non-phytate phosphorus concentration in the pre-experimental starter diet. *Poult. Sci.*
920            **103**, (2024).

921    48.    Mutucumarana, R. K., Ravindran, V., Ravindran, G. & Cowieson, A. J. Measurement
922            of true ileal digestibility and total tract retention of phosphorus in corn and canola meal
923            for broiler chickens. *Poult. Sci.* **93**, (2014).

924    49.    Chang'a, E. P. *et al.* Energy utilization, nutrient digestibility and bone quality of
925            broiler chickens fed Tanzania-type diets in different forms with enzymes. *J. Anim. Sci.
926            Technol.* **61**, (2019).

927    50.    Thomas, D. V. & Ravindran, V. Mineral retention in young broiler chicks fed diets
928            based on wheat, Sorghum or maize. *Asian-Australas. J. Anim. Sci.* **23**, (2010).

51.  Ward, T. L., Watkins, K. L. & Southern, L. L. Interactive effects of dietary copper,
water copper, and Eimeria spp. infection on growth, water intake, and plasma and liver
copper concentrations of poults. *Poult. Sci.* **74**, (1995).

52.  Macelline, S. P., Chrystal, P. V., Selle, P. H. & Liu, S. Y. Protein sources and starch-
protein digestive dynamics manipulate growth performance in broiler chickens defined
by an equilateral-triangle response surface design. *Animal Nutrition* **9**, (2022).

53.  Stefanello, C. *et al.* Starch digestibility, energy utilization, and growth performance of
broilers fed corn-soybean basal diets supplemented with enzymes. *Poult. Sci.* **94**,
(2015).

54.  Khalil, M. M., Abdollahi, M. R., Zaefarian, F. & Ravindran, V. Measurement of ileal
endogenous energy losses and true ileal digestible energy of cereal grains for broiler
chickens. *Poult. Sci.* **99**, (2020).

55.  Truong, H. H., Chrystal, P. V., Moss, A. F., Selle, P. H. & Liu, S. Y. Rapid protein
disappearance rates along the small intestine advantage poultry performance and
influence the post-enteral availability of amino acids. *British Journal of Nutrition* **118**,
(2017).

56.  Tancharoenrat, P., Ravindran, V., Zaefarian, F. & Ravindran, G. Digestion of fat and
fatty acids along the gastrointestinal tract of broiler chickens. *Poult. Sci.* **93**, 371–379
(2014).

57.  Seaver, S. M. D. *et al.* The ModelSEED Biochemistry Database for the integration of
metabolic annotations and the reconstruction, comparison and analysis of metabolic
models for plants, fungi and microbes. *Nucleic Acids Res.* **49**, (2021).

58.  Zheng, J. *et al.* DbCAN3: Automated carbohydrate-Active enzyme and substrate
annotation. *Nucleic Acids Res.* **51**, (2023).

59.  Cantarel, B. I. *et al.* The Carbohydrate-Active EnZymes database (CAZy): An expert
resource for glycogenomics. *Nucleic Acids Res.* **37**, (2009).

60.  Bansal, P. *et al.* Rhea, the reaction knowledgebase in 2022. *Nucleic Acids Res.* **50**,
(2022).

61.  Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: New
perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res.* **45**, D353–
D361 (2017).

62.  Schomburg, I., Chang, A. & Schomburg, D. BRENDA, enzyme data and metabolic
information. *Nucleic Acids Res.* **30**, (2002).

63.    Karp, P. D. *et al.* The BioCyc collection of microbial genomes and metabolic
pathways. *Brief. Bioinform.* **20**, (2018).

64.    Kanehisa, M. & Goto, S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic*
*Acids Research* vol. 28 Preprint at https://doi.org/10.1093/nar/28.1.27 (2000).

65.    Karp, P. D., Riley, M., Paley, S. M. & Pellegrini-Toole, A. The MetaCyc database.
*Nucleic Acids Res.* **30**, (2002).

66.    King, Z. A. *et al.* BiGG Models: A platform for integrating, standardizing and sharing
genome-scale models. *Nucleic Acids Res.* **44**, (2016).

67.    Ravindran, V. Feed enzymes: The science, practice, and metabolic realities. *Journal of*
*Applied Poultry Research* **22**, 628–636 (2013).

68.    Shang, Y., Kumar, S., Oakley, B. & Kim, W. K. Chicken Gut Microbiota￼:
Importance and Detection Technology. *Front. Vet. Sci.* **5**, (2018).

69.    Aruwa, C. E., Pillay, C., Nyaga, M. M. & Sabiu, S. Poultry gut health – microbiome
functions, environmental impacts, microbiome engineering and advancements in
characterization technologies. *Journal of Animal Science and Biotechnology* vol. 12
Preprint at https://doi.org/10.1186/s40104-021-00640-9 (2021).

70.    Bindari, Y. R. & Gerber, P. F. Centennial Review: Factors affecting the chicken
gastrointestinal microbial composition and their association with gut health and
productive performance. *Poultry Science* vol. 101 Preprint at
https://doi.org/10.1016/j.psj.2021.101612 (2022).

71.    Yadav, S. & Jha, R. Strategies to modulate the intestinal microbiota and their effects
on nutrient utilization, performance, and health of poultry. *J. Anim. Sci. Biotechnol.* **10**,
1–11 (2019).

72.    AFZ. INRA-CIRAD-AFZ feed tables: composition and nutritive values of feeds for
cattle, sheep, goats, pigs, poultry, rabbits, horses, and salmonids. Feedtables.com,
INRAE, CIRAD, AFZ. (2017).

73.    Noronha, A. *et al.* The Virtual Metabolic Human database￼: integrating human and
gut microbiome metabolism with nutrition and disease. *Nucleic Acids Res.* **47**, 614–
624 (2018).

74.    Grieshop, C. M. *et al.* Chemical and Nutritional Characteristics of United States
Soybeans and Soybean Meals. *J. Agric. Food Chem.* **51**, (2003).

75.    Wu, Y. V. Determination of Neutral Sugars in Corn Distillers' Dried Grains, Corn
Distillers' Dried Solubles, and Corn Distillers' Dried Grains with Solubles. *J. Agric.*
*Food Chem.* **42**, (1994).

76. Zilic, S., Milasinovic, M., Terzic, D., Barac, M. & Ignjatovic-Micic, D. Grain
characteristics and composition of maize specialty hybrids. *Spanish Journal of*
*Agricultural Research* **9**, (1970).

77. Miya, S. P., Modi, A. T., Tesfay, S. Z. & Mabhaudhi, T. Maize grain soluble sugar and
protein contents in response to simulated hail damage. *South African Journal of Plant*
*and Soil* **35**, (2018).

78. Schmidt, M. & Sciurba, E. Determination of FODMAP contents of common wheat and
rye breads and the effects of processing on the final contents. *European Food*
*Research and Technology* **247**, (2021).

79. Svihus, B. Function of the digestive system. *Journal of Applied Poultry Research* **23**,
306–314 (2014).

80. Barash, I., Nitsan, Z. & NIR, Z. Metabolic and behavioural adaptation of light-bodied
chicks to meal feeding. *Br. Poult. Sci.* **33**, (1992).

81. Taylor, J., Sakkas, P. & Kyriazakis, I. What are the limits to feed intake of broilers on
bulky feeds? *Poult. Sci.* **100**, (2021).

82. Smith, D. P. & Berrang, M. E. Prevalence and numbers of bacteria in broiler crop and
gizzard contents. *Poult. Sci.* **85**, (2006).

83. Hassanin, F., Hassan, M., Shaltout, F., Shawqy, N. & Abd-Elhameed, G.
Bacteriological criteria of chicken giblets. *Benha Vet. Med. J.* **33**, (2017).

84. Dänicke, S., Vahjen, W., Simon, O. & Jeroch, H. Effects of dietary fat type and
xylanase supplementation to rye-based broiler diets on selected bacterial groups
adhering to the intestinal epithelium, on transit time of feed, and on nutrient
digestibility. *Poult. Sci.* **78**, (1999).

85. Shires, A., Thompson, J. R., Turner, B. V., Kennedy, P. M. & Goh, Y. K. Rate of
passage of corn-canola meal and corn-soybean meal diets through the gastrointestinal
tract of broiler and White Leghorn chickens. *Poult. Sci.* **66**, (1987).

86. Chee, S. H., Iji, P. A., Choct, M., Mikkelsen, L. L. & Kocher, A. Functional
interactions of manno-oligosaccharides with dietary threonine in chicken
gastrointestinal tract. III. Feed passage rate. *Br. Poult. Sci.* **51**, (2010).

87. Duke, G. E., Eccleston, E., Louis, C. F. & Bedbury, H. P. Cellulose digestion by
domestic turkeys fed low or high fiber diets. *Journal of Nutrition* **114**, 95–102 (1984).

88. Duke, G. E. Relationship of cecal and colonic motility to diet, habitat, and cecal
anatomy in several avian species. *Journal of Experimental Zoology* **252**, 38–47 (1989).

89.  Duve, L. R., Steenfeldt, S., Thodberg, K. & Nielsen, B. L. Splitting the scotoperiod:
Effects on feeding behaviour, intestinal fill and digestive transit time in broiler
chickens. *Br. Poult. Sci.* **52**, (2011).

90.  Adelsohn, D. S., Scanes, C. G. & Decuypere, E. Diurnal—nocturnal changes in food
intake gut storage of ingesta, food transit time and metabolism in growing broiler
chickens: A model for temporal control of energy balance. *Br. Poult. Sci.* **34**, (1993).

91.  Sacranie, A., Iji, P., L. Mikkelsen, L. & Choct, M. Occurrence of reverse peristalsis in
broiler chickens. in *Conference: Australian Poultry Science Symposium, Sydney, New*
*South Wales, Australia Volume: 19* (Sydney, New South Wales, Australia, 2007).

92.  Wise, M. G. & Siragusa, G. R. Quantitative analysis of the intestinal bacterial
community in one- to three-week-old commercially reared broiler chickens fed
conventional or antibiotic-free vegetable-based diets. *J. Appl. Microbiol.* **102**, (2007).

93.  Riesenfeld, G., Sklan, D., Bar, A., Eisner, U. & Hurwitz, S. Glucose absorption and
starch digestion in the intestine of the chicken. *Journal of Nutrition* **110**, 117–121
(1980).

94.  Bielorai, R. & Iosif, B. Amino acid absorption and endogenous amino acids in the
lower ileum and excreta of chicks. *Journal of Nutrition* **117**, 1459–1462 (1987).

95.  Bielorai, R., Tamir, M. & Hurwitz, S. Amino acid absorption along the intestinal tract
of chicks fed heated and raw soybean meal. *Journal of Nutrition* **107**, 1775–1778
(1977).

96.  Tancharoenrat, P., Ravindran, V., Zaefarian, F. & Ravindran, G. Digestion of fat and
fatty acids along the gastrointestinal tract of broiler chickens. *Poult. Sci.* **93**, 371–379
(2014).

97.  Rodriguez-Sanchez, R., Tres, A., Sala, R., Guardiola, F. & Barroeta, A. C. Evolution
of lipid classes and fatty acid digestibility along the gastrointestinal tract of broiler
chickens fed different fat sources at different ages. *Poult. Sci.* **98**, 1341–1353 (2019).

98.  He, W., Li, P. & Wu, G. Amino Acid Nutrition and Metabolism in Chickens. in
*Advances in Experimental Medicine and Biology* vol. 1285 (2021).

99.  Scanes, C. G. & Pierzchala-Koziec, K. Biology of the gastro-intestinal tract in poultry.
*Avian Biol. Res.* **7**, 193–222 (2014).

100. Rodrigues, I. & Choct, M. The foregut and its manipulation via feeding practices in the
chicken. *Poult. Sci.* **97**, 3188–3206 (2018).

101. Sergeant, M. J. *et al.* Extensive microbial and functional diversity within the chicken
cecal microbiome. *PLoS One* **9**, (2014).

102.    Rios-Covian, D., Gueimonde, M., Duncan, S. H., Flint, H. J. & De Los Reyes-Gavilan,
1064            C. G. Enhanced butyrate formation by cross-feeding between Faecalibacterium
1065            prausnitzii and Bifidobacterium adolescentis. *FEMS Microbiol. Lett.* **362**, (2015).

1066    103.    Louis, P. & Flint, H. J. Diversity, metabolism and microbial ecology of butyrate-
1067            producing bacteria from the human large intestine. *FEMS Microbiol. Lett.* **294**, (2009).

1068    104.    Iwami, Y., Abbe, K., Takahashi☐Abbe, S. & Yamada, T. Acid production by
1069            streptococci growing at low pH in a chemostat under anaerobic conditions. *Oral*
1070            *Microbiol. Immunol.* **7**, (1992).

1071    105.    Carlsson, J. & Griffith, C. J. Fermentation products and bacterial yields in glucose-
1072            limited and nitrogen-limited cultures of streptococci. *Arch. Oral Biol.* **19**, (1974).

1073    106.    SHOCKMAN, G. D. The acetate requirement of Streptococcus faecalis. *J. Bacteriol.*
1074            **72**, (1956).

1075    107.    Beelen, P., Hoeven, J. S., Jong, M. H. & Hoogendoorn, H. The effect of oxygen on the
1076            growth and acid production of Streptococcus mutans and Streptococcus sanguis .
1077            *FEMS Microbiol. Lett.* **38**, (1986).

1078    108.    Shin, J. H. *et al.* Bacteroides and related species: The keystone taxa of the human gut
1079            microbiota.           *Anaerobe*          vol.           85          Preprint          at
1080            https://doi.org/10.1016/j.anaerobe.2024.102819 (2024).

1081    109.    Lyte, J. M. *et al.* Inclusion of trans-cinnamaldehyde and caprylic acid in feed results in
1082            detectable concentrations in the chicken gut and reduces foodborne pathogen carriage.
1083            *Poult. Sci.* **103**, (2024).

1084    110.    Ge, S. *et al.* GABA regulates synaptic integration of newly generated neurons in the
1085            adult brain. *Nature* **439**, (2006).

1086    111.    Allaband, C. *et al.* Time of sample collection is critical for the replicability of
1087            microbiome    analyses.    *Nature    Metabolism*    vol.    6    1282–1293    Preprint    at
1088            https://doi.org/10.1038/s42255-024-01064-1 (2024).

1089    112.    Xia, Y. *et al.* Dietary inulin supplementation modulates the composition and activities
1090            of carbohydratemetabolizing organisms in the cecal microbiota of broiler chickens.
1091            *PLoS One* **16**, (2021).

1092    113.    De Maesschalck, C. *et al.* Effects of Xylo-oligosaccharides on broiler chicken
1093            performance and microbiota. *Appl. Environ. Microbiol.* **81**, (2015).

1094    114.    Richards, P. J. *et al.* Galacto-Oligosaccharides Modulate the Juvenile Gut Microbiome
1095            and Innate Immunity To Improve Broiler Chicken Performance. *mSystems* **5**, (2020).

115.    Jahan, A. A., González Ortiz, G., Moss, A. F., Bhuiyan, M. M. & Morgan, N. K. Role
of supplemental oligosaccharides in poultry diets. *World's Poultry Science Journal*
vol. 78 Preprint at https://doi.org/10.1080/00439339.2022.2067805 (2022).

116.    Schwartz, B. & Vetvicka, V. Review: β☐glucans as effective antibiotic alternatives in
poultry. *Molecules* vol. 26 Preprint at https://doi.org/10.3390/molecules26123560
(2021).

117.    Morgan, N. K. Advances in prebiotics for poultry: role of the caeca and
oligosaccharides. *Animal Production Science* vol. 63 Preprint at
https://doi.org/10.1071/AN23011 (2023).

118.    Zhang, Y. *et al.* Dietary resistant starch modifies the composition and function of
caecal microbiota of broilers. *J. Sci. Food Agric.* **100**, (2020).

119.    Suckeveris, D. *et al.* Supra-nutritional levels of selected B vitamins in animal or
vegetable diets for broiler chicken. *Revista Brasileira de Ciencia Avicola / Brazilian*
*Journal of Poultry Science* **22**, (2020).

120.    Leiber, F. *et al.* Effects of riboflavin supplementation level on health, performance,
and fertility of organic broiler parent stock and their chicks. *Animal* **16**, (2022).

121.    Santos, T. S. dos *et al.* Effects of Inorganic Zn and Cu Supplementation on Gut Health
in Broiler Chickens Challenged With Eimeria spp. *Front. Vet. Sci.* **7**, (2020).

122.    Biswas, A., Mohan, N., Dev, K., Mir, N. A. & Tiwari, A. K. Effect of dietary mannan
oligosaccharides and fructo-oligosaccharides on physico-chemical indices, antioxidant
and oxidative stability of broiler chicken meat. *Sci. Rep.* **11**, (2021).

123.    Froidurot, A. & Julliand, V. Cellulolytic bacteria in the large intestine of mammals.
*Gut Microbes* vol. 14 Preprint at https://doi.org/10.1080/19490976.2022.2031694
(2022).

124.    Faure, M. *et al.* Specific amino acids increase mucin synthesis and microbiota in
dextran sulfate sodium-treated rats. *Journal of Nutrition* **136**, (2006).

125.    Bell, S. C. & Turner, J. M. Bacterial catabolism of threonine. Threonine degradation
initiated by L-threonine hydrolyase (deaminating) in a species of Corynebacterium.
*Biochemical Journal* **164**, (1977).

126.    Bell, S. C. & Turner, J. M. Bacterial catabolism of threonine. Threonine degradation
initiated by L-threonine acetaldehyde-lyase (aldolase) in species of Pseudomonas.
*Biochemical Journal* **166**, (1977).

127.    Simanshu, D. K., Chittori, S., Savithri, H. S. & Murthy, M. R. N. Structure and
function of enzymes involved in the anaerobic degradation of L-threonine to

propionate. *Journal of Biosciences* vol. 32 Preprint at https://doi.org/10.1007/s12038-
007-0121-1 (2007).

128.  Quinn-Bohmann, N. *et al.* Microbial community-scale metabolic modeling predicts
personalized short chain fatty acid production profiles in the human gut. *bioRxiv*
Preprint at https://doi.org/10.1101/2023.02.28.530516 (2023).

129.  Zimmermann-Kogadeeva, M., Bencivenga-Barry, N. A., Zimmermann, M., Bork, P. &
Goodman, A. L. Multiomics and quantitative modelling disentangle diet, host, and
microbiota contributions to the host metabolome. *bioRxiv* 2022.08.15.503927 (2022)
doi:10.1101/2022.08.15.503927.

130.  Carr, A. V., Baliga, N. S., Diener, C. & Gibbons, S. M. Personalized Clostridioides
difficile colonization risk prediction and probiotic therapy assessment in the human
gut. *Cell Syst.* **16**, 101367 (2025).

131.  Gutierrez del alamo, A., Verstegen, M. W. A., Den hartog, L. A., Perez de ayala, P. &
Villamide, M. J. Wheat starch digestion rate affects broiler performance. *Poult. Sci.*
**88**, (2009).

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

**FIGURE LEGENDS**

**Figure 1. A spatiotemporal, multi-compartment metabolic modeling framework for the chicken gastrointestinal tract microbiome. A.** Overview of the modeling workflow. Community composition in each gut section (gizzard, duodenum, jejunum, ileum, cecum, colon) was parameterized using 16S rRNA profiles from corn- and wheat-fed chickens. Representative taxa were mapped to high-quality chicken gut-derived MAGs and reconstructed into genome-scale metabolic models with gapseq. Models were curated with CAZyme predictions (dbCAN3) to ensure representation of fiber-degrading pathways and integrated into sample- and compartment-specific community metabolic models. **B.** The six compartments are connected to simulate the flow of microbes and metabolites along the tract, with host physiological parameters including pH, oxygen, motility, digestion, and absorption incorporated to reflect site-specific conditions. **C.** Representative taxa selected to comprise core microbiota, shown in a phylogenetic tree with compartment-specific relative abundances (averaged across 10 samples) and the metabolic breadth of each corresponding GEM (number of EC numbers, reactions, pathways). **D.** PCoA of model reaction content using Bray-Curtis dissimilarity, illustrating clustering of taxa by their predicted metabolic repertoires. Each dot corresponds to a model of the taxon labeled. **E.** PCoA of reaction abundance profiles at the sample level, weighted by relative taxon abundance, showing clear separation of communities by gut compartment and partial separation by diet.

**Figure 2. Temporal dynamics of metabolite concentrations across the chicken gastrointestinal tract during feeding and fasting cycles. A.** Taxon-specific flux predictions for SCFAs over 96-hour simulations. **B.** Principal Coordinate Analysis (PCoA) of metabolite profiles across six gut compartments (gizzard, duodenum, jejunum, ileum, cecum, colon) at three time points: beginning of the feeding period after fasting, middle of the feeding period, and middle of the fasting period. Points are color-coded by compartment (corn- and wheat-fed samples combined to emphasize conserved compartment-specific signatures), with distinct clusters reflecting compartment-specific metabolic activities and shifts over time across both diets. **C.** Concentration profiles of selected metabolites across gut compartments at three time points: beginning of the feeding period (pink), middle of the feeding period (light blue), and middle of the fasting period (dark brown). The plots indicate the mean concentrations and standard deviations of metabolite concentrations from samples corresponding to chickens fed corn-based diet (n=5). **D.** Dynamics of metabolite concentrations across gut compartments during 96 hours of simulation for a selection of metabolites. Rows represent specific metabolites, with color intensity indicating concentration levels. The results shown were retrieved from simulation of one of the samples from corn-fed chicken.

**Figure 3**. **Impact of dietary supplementation on metabolite profiles and SCFA production across gut compartments. A.** PCoA of metabolite concentrations after 96-hour simulations. Representative samples from corn-fed (Corn5-2) and wheat-fed (Wheat11-3) chickens are shown, comparing control conditions (black dots) with various dietary supplements across the ileum, cecum, and colon compartments. Key compounds discussed in the Results section (L-threonine, starch, pantothenic acid, cellulose, high-dose amino acids) are emphasized with thicker outlines. Arrows indicate L-threonine, cellulose, and starch - the three compounds selected for validation trials. Top panels: PCoA based on all metabolite concentrations. Bottom panels: PCoA based solely on SCFA concentrations. Supplement types are indicated by symbols and colors, with dosing shown after underscores (e.g., Inulin_2 = 2g per 100g diet). Complete dataset including all samples and compartments is presented in Supplementary Figure 4. **B.** Distribution of SCFA production responses to

supplementation. Stacked bar plots show the proportion of tested compounds that significantly increased (green), decreased (red), or did not affect (gray) community-level SCFA production in each compartment. **C.** Scatter plots showing fold changes in butyrate (x-axis) versus propionate (y-axis) production for all tested supplements. Bubble size represents the magnitude of acetate fold change. Colors indicate which SCFA changes were statistically significant: red = all three SCFAs; yellow = acetate + butyrate; orange = acetate + propionate; green = butyrate + propionate; brown = acetate only; purple = butyrate only; blue-gray = propionate only; transparent gray = no significant changes. Dashed lines at fold change = 1.0 indicate average production levels in control simulations.

**Figure 4. Trial-specific community composition data substantially improves metabolic predictions for cecal fermentation. A.** Correlation between predicted and experimental log☐(fold change) values for seven fermentation metabolites in the cecum. Top row: Predictions from the 6-compartment model after 96-hour simulations compared to experimental measurements from the validation trial. Bottom row: Predictions from the trial-informed 2-compartment model after 48-hour simulations compared to experimental measurements. Each panel shows one metabolite with treatment groups indicated by color. Dashed lines represent linear regression fits. Spearman correlation coefficient (rho), uncorrected p-value (p), and FDR-adjusted q-value (q) are displayed for each metabolite. Fold changes were calculated relative to control group medians. **B**. Community composition differences. PCoA plots comparing taxonomic profiles between original reference data (Initial data), simplified model communities (Simulation), and trial communities (Validation trial) for cecum. PERMANOVA confirms significant separation between data sources (p < 0.05), highlighting taxonomic basis for model refinement.

## SUPPLEMENTAL MATERIAL

**Supplementary Figure 1. Taxonomic composition of gut microbiota in different gut compartments.** The relative abundance of bacterial families derived from 16S rRNA survey data is shown for samples from the gizzard, duodenum, jejunum, ileum, cecum, and colon, separated by corn and wheat diets. Color bars represent different taxonomic families, with relative proportions depicted for each gut compartment.

**Supplementary Figure 2. Distribution of enzyme commission numbers (ECs) across metabolic models.** The columns of the heatmap represent individual ECs, while the rows correspond to metabolic models representing different taxa. ECs are annotated using KEGG pathway designations, indicated in two layers to reflect primary and secondary pathway associations. For ECs lacking KEGG pathway annotation, MetaCyc pathway annotation was utilized when available. For ECs mapped to 3 or more major pathway categories, "Multiple superpathways" category was assigned. Each row represents a different taxon, annotated by taxonomic family.

**Supplementary Figure 3. Reaction-level drivers of compartmental separation observed in Figure 1E. A.** Distribution of PC1 (left) and PC2 (right) scores from the principal coordinates analysis (PCoA shown in **Figure 1E**) across gut compartments. PCoA was performed on Bray-Curtis dissimilarities of community metabolic reaction profiles weighted by species relative abundances. PC scores reflect the relative contribution of reaction sets associated with each axis rather than distances to compartment centroids. Dashed red lines

indicate zero for each axis. PC1 primarily separates upper gastrointestinal tract compartments (gizzard, duodenum, jejunum, ileum) from lower gut compartments (cecum, colon), whereas PC2 captures additional metabolic stratification along the proximal-distal axis. **B.** Heatmap showing mean compartment-level abundances of metabolic reactions most strongly associated with PC1 and PC2 in Fig. 1E. Reactions were selected based on the highest positive and negative Spearman correlations between individual reaction abundances and PCoA axis scores (top loadings). Values are column-scaled (z-scores per reaction) to emphasize relative enrichment across compartments. Reactions associated with positive PC1 scores (upper GIT) include bile salt hydrolases, membrane lipid synthesis pathways, and fatty acid biosynthesis, consistent with biosynthetically active, oxygen-replete environments. In contrast, reactions associated with negative PC1 scores (lower GIT) are enriched for anaerobic fermentation processes, including short-chain fatty acid biosynthesis, polysaccharide degradation, and amino acid fermentation.

**Supplementary Figure 4. PCoA of metabolite concentrations after 96-hour simulations.** All 10 samples from corn-fed and wheat-fed chickens are shown, comparing control conditions (black dots) with various dietary supplements across all six gut compartments. Top panels: PCoA based on all metabolite concentrations. Bottom panels: PCoA based solely on SCFA concentrations. Supplement types are indicated by symbols and colors, with dosing shown after underscores (e.g., Inulin_2 = 2g per 100g diet).

**Supplementary Figure 5. SCFA production rates in simulations with supplementation.** Fold changes in total SCFA production by the microbial community over 96 hours of simulation when supplemented with a specific compound (y-axis), compared to the control, in ileum and cecum. The y-axis lists the various dietary supplements tested, with their concentrations indicated after the underscore (e.g., Inulin_2 denotes supplementation with 2g per 100g of diet). Each dot represents a fold change in community flux in a replicate (n=5) of a sample (n=10). Significance based on two-sided Wilcoxon signed-rank test with Benjamini–Hochberg correction for multiple testing: * – pvalue- adjusted < 0.05, ** – pvalue-adjusted < 0.01, *** – pvalue-adjusted < 0.001.

**Supplementary Figure 6. Taxonomic composition of gut microbiota in trial-derived samples from ileum and cecum gut compartments.** The relative abundances of bacterial families are derived from 16S rRNA survey data. Color bars represent different taxonomic families, with relative proportions depicted for each gut compartment.

**Supplementary Figure 7. Experimental validation of metabolite concentrations in response to dietary supplementation.** Targeted metabolomics measurements from chicken ileum and cecum samples following 7-day dietary supplementation. **A**. Box plots showing metabolite concentrations (μmol/g dry weight) across treatment groups for both ileum and cecum compartments. Each metabolite panel displays individual data points overlaid on box plots, with Kruskal-Wallis (KW) p-values shown in the top-left of each facet. Significant pairwise comparisons between treatment groups and control (determined by Dunn's post-hoc test with Benjamini-Hochberg correction) are indicated by brackets with asterisks. **B**. Bar plots showing log2 fold changes relative to control group for each treatment in ileum and cecum. Asterisks indicate statistical significance from two-sample Wilcoxon rank-sum tests with Benjamini-Hochberg correction (* - $p < 0.05$, ** - $p < 0.01$, *** - $p < 0.001$). Fumarate showed consistent depletion across all treatments (KW p=0.002), contrasting with model predictions of net microbial production, suggesting rapid consumption or host absorption. Butyrate exhibited significant treatment effects (KW p=0.029), with increases under low-dose

cellulose and starch aligning with predicted production by Ruminococcaceae, Butyricicoccaceae, and Lachnospiraceae. Lactate showed significant treatment effects (KW p=0.034) with high inter-individual variability, consistent with its role as both a fermentation product and rapidly consumed substrate. Succinate showed predominantly negative fold changes despite model predictions of increased production under starch (Bacteroidaceae) and cellulose (Lachnospiraceae), indicating additional metabolic sinks not captured in models. Acetate production increased in ileum, especially under cellulose and L-threonine treatments, however no consistent treatment effects were observed in cecum, reflecting acetate broad production across multiple families.

**Supplementary Figure 8. Predicted taxonomic contributions to metabolite production in the trial-informed 2-compartment model.** Bacterial flux predictions from 48-hour simulations of the trial-informed 2-compartment (ileum-cecum) model showing which taxonomic families contribute to production of seven fermentation metabolites. Horizontal stacked bar plots display cumulative metabolic flux (fmol/gdW/hour) for acetate, propionate, butyrate, lactate, fumarate, succinate, and alpha-ketoglutarate across all seven treatment groups (G1 (control), G2 (cellulose 2g/100g), G3 (cellulose 8g/100g), G4 (L-threonine 1g/100g), G5 (L-threonine 5g/100g), G6 (starch 5g/100g), and G7 (starch 10g/100g)) in ileum and cecum compartments. Each color represents a taxonomic family. Flux values represent total community production summed across all 48 hours of simulation and normalized by community size. The figure illustrates predicted treatment-specific modulation of bacterial metabolism, with cellulose and starch supplementation enhancing butyrate production capacity in cellulose- and starch-fermenting families, as referenced in the main text analyses.

**Supplementary Table S1. MAG selections for 29 bacterial species in multi-compartment GIT metabolic models.** Genome assignments for 29 bacterial species selected for genome-scale metabolic modeling of chicken gastrointestinal tract compartments. Species represent prevalent taxa from day 10 post-hatch samples across six GIT sites. Table includes species names, MAG identifiers, genome source databases (Intestinal Microbial Genomes ANI99 collection, HoloFood chicken gut MAG catalog), match quality categories, CheckM quality metrics (completeness, contamination), originating GIT site, assembly type, complete taxonomic classification, and selection notes. NCBI reference genomes were used when high-quality matches were unavailable in the three chicken gut MAG databases used as a reference.

**Supplementary Table S2. A. Simulation parameters for 6-compartment model.** Parameters for spatial metabolic simulations using BacArena. Table S2A provides parameters for the six-compartment gastrointestinal tract model representing gizzard, duodenum, jejunum, ileum, cecum, and colon. Parameters include initial bacterial community sizes per compartment, spatial grid dimensions (arena size), basal nutrient availability (essential metabolite concentrations), physicochemical conditions (pH, oxygen concentration), and dietary input schedules (proportion and timing of feed addition to gizzard). Additional parameters specify differential removal rates for day versus night periods and substrate-level removal percentages, reflecting circadian feeding patterns and intestinal motility in broiler chickens. **B. Simulation parameters for ileum-cecum model.** Table S2B provides parameters for the simplified two-compartment ileum-cecum model, including bacterial removal rates and frequencies that simulate digesta flow and GIT transit. Additional parameters specify differential removal rates for day versus night periods and substrate-level

removal percentages, reflecting circadian feeding patterns and intestinal motility in broiler chickens.

**Supplementary Table S3.MAG assignements for metabolic models in ileum and cecum microbiomes.** Representative MAG assignments for 267 functionally clustered ASVs from paired chicken ileum and cecum samples (validation trial). ASVs were clustered by 75% functional similarity (Jaccard index) of PICRUSt2-predicted enzyme profiles, with representatives selected to maximize MAG coverage. MAGs were selected from three chicken gut databases: ANI99 collection (Huang et al. 2022), MGnify chicken gut catalog (13,389 genomes), and HoloFood 12k MAG catalog. Selection prioritized combined BLAST quality (identity, bit score) and genome quality (completeness, contamination). Table includes: sample identifiers, treatment groups (control vs. prebiotic), body site (ileum/cecum), cumulative relative abundance per functional cluster, MAG identifiers, genome source databases, BLAST statistics (identity, bit score), MAG quality metrics (completeness, contamination), match quality categories (excellent/good/acceptable), and full taxonomic classifications.

**Supplementary Table S4. Diet compositions for metabolic simulations.** Complete metabolic compositions of corn-based and wheat-based grower diets used in simulations. Feed compositions from the original trial were translated into corresponding metabolites using animal feed tables mapping nutrient content to feed items. Oil compositions were sourced from vmh.life. Sugar and fiber categories were resolved into specific carbohydrates and polysaccharides based on published compositional studies, with cellulose used as the primary fiber source.

**Supplementary Table S5. Reactions added to and removed from metabolic models based on CAZyme gene annotations.** Manual curation of metabolic reactions to improve representation of complex carbohydrate metabolism beyond gapseq's default modelSEED database. Sheet 1 documents 55 reactions added based on CAZymes detected by dbCAN3 analysis. Reactions were incorporated directly from modelSEED when exact matches existed, or manually defined using multiple biochemical databases for enzymes not in modelSEED. Includes polysaccharide degradation reactions (arabinan, xyloglucan, xylan, etc.) and associated transport reactions, with detailed information on reaction definitions, EC numbers, stoichiometry, reversibility, affected species models, and newly introduced metabolites with chemical formulas. Sheet 2 documents 14 reactions removed from models when both: (1) the corresponding CAZyme was not detected by dbCAN3 in the genome, and (2) no significant sequence similarity was found during gapseq reaction search. Removed reactions include incorrectly predicted fructan biosynthesis and starch degradation activities.

**A** Cecum - 6-compartment model

Cecum - validation trial-informed 2-compartment model

Treatment
- G1 - Control
- G2 - Cellulose Low
- G3 - Cellulose High
- G4 - L−Threonine Low
- G5 - L−Threonine High
- G6 - Starch Low
- G7 - Starch High

**B** Comparison of microbial compositions in the cecum across data sources

Treatment
- G1 - Control
- G2 - Cellulose Low
- G3 - Cellulose High
- G4 - L−Threonine Low
- G5 - L−Threonine High
- G6 - Starch Low
- G7 - Starch High

Data source: ▲ Initial 16S data  ■ Simulation  ● Validation trial