Tissue-scale mechanics controls differentiation strategy and dynamics of epithelial

multilayering

Clémentine Villeneuve[1,2]*, Somiealo Azote Epse Hassikpezi [3]*, Marga Albu[2], Matthias Rübsam[4],

Leah C. Biggs[1,2], Sabrina Vinzens[2], Kai Kruse[5], Anubhav Prakash[2], Peter Zentis[6], Elizabeth Lawson-

Keister[3], Gautier Follain[7,8,9], Johanna Ivaska[7,10,11], Carien M. Niessen[4§], M. Lisa Manning[3§], and Sara

A. Wickström[1,2,12§]

[1]Stem Cells and Metabolism Research Program, Faculty of Medicine, University of Helsinki, 00290 Helsinki Finland

[2]Department of Cell and Tissue Dynamics, Max Planck Institute for Molecular Biomedicine, 48149 Münster, Germany

[3]Department of Physics and BioInspired Institute, Syracuse University, Syracuse, NY, USA.

[4]Department Cell Biology of the Skin, Cologne Excellence Cluster on Cellular Stress Responses in Aging Associated
Diseases (CECAD), Center for Molecular Medicine Cologne, University Hospital Cologne, University of Cologne,
Joseph-Stelzmann-Str. 26, 50931 Cologne, Germany.

[5]Max Planck Institute for Molecular Biomedicine, 48149 Münster, Germany

[6]Cluster of Excellence for Aging-Associated Diseases, CECAD Imaging Facility Cologne, University of Cologne,
Cologne, Germany.

[7]Turku Bioscience Centre, University of Turku and Åbo Akademi University, Turku, 20520, Finland
[8]Faculty of Science and Engineering, Cell Biology, Åbo Akademi University, Turku, 20520, Finland

[9]Turku Collegium for Science, Medicine and Technology (TCSMT), Turku, 20520, Finland

[10]Department of Life Technologies, University of Turku, FI-20520 Turku, Finland.

[11]InFLAMES Research Flagship Center, University of Turku, Turku, Finland.

[12]Helsinki Institute of Life Science, Biomedicum Helsinki, University of Helsinki, FI-00290 Helsinki, Finland

*These authors contributed equally

§These authors jointly supervised this work

carien.niessen@uni-koeln.de

mmanning@syr.edu

sara.wickstrom@mpi-muenster.mpg.de

Abstract

Generating and maintaining multilayered epithelia requires coordinated cell division, differentiation,

and tissue architecture, yet the precise mechanisms of multilayering remain unclear. Using the

developing mouse epidermis, we show that basal stem cells adopt distinct multilayering strategies

depending on tissue mechanics. Combining quantitative morphometry, embryo live imaging and

physical modeling, we observe that early in development, the epidermis is fluid-like, allowing

undifferentiated cells to move suprabasally through perpendicular divisions or basal detachment

before differentiating. As the tissue matures and rigidifies, a mechanical barrier is established that

only allows upward movement of basal cells that have committed to differentiation. The final step of

this commitment is delamination that requires Notch signaling, triggered by increased tissue stiffness

and jamming. This mechanical regulation orchestrates a feedback loop that induces cell upward

motion precisely when the basal layer becomes crowded. Together, our findings identify tissue

mechanics as the key determinant of how tissues drive multilayering and reveal mechanically

regulated Notch signaling as a driver of epidermal delamination.

Introduction

Stratified epithelia such as the skin epidermis are multilayered tissues where the basal layer consists

of undifferentiated epithelial stem cells adhering to the basement membrane, while the suprabasal

layers contain post-mitotic, differentiated cells [1-3]. Unlike other dynamically self-renewing epithelia

such as the intestine, in which clear tissue-scale geometric cues such as curvature compartmentalize

the stem cell away from differentiated progeny [4], it remains unclear how stratified epithelia establish

and organize this spatial organization and differentiation dynamics during development and into

adulthood.

To initiate or maintain epidermal stratification, the basal stem cells generate suprabasal cells via two

distinct strategies: perpendicular divisions or detachment from the basal layer, termed delamination

[2,3,5]. Early in development, proliferation rates in the epidermis are high - to match the rapid growth of

the organism it covers, and to generate sufficient suprabasal cells for efficient establishment of a

functional life-essential barrier [6]. During this period, perpendicular cell divisions generate suprabasal

daughter cells [7,8], while also some delaminations are observed [9,10]. In contrast, during adult

homeostasis differentiation does not depend on cell divisions, with cells moving suprabasally

exclusively through delamination to replenish the most superficial, desquamating cornified layers,

followed by parallel divisions to maintain basal cell density [11,12]. Key open questions remain: Why

does this switch in strategies occur? And how is upward movement robustly regulated in the absence

of obvious niche inputs like tissue curvature?

Here we demonstrate that multilayering strategy is controlled by a mechanical barrier that is

established during development and hinders basal cells from moving upward. We establish a new

approach to quantitively measure this mechanical barrier by combining a 3D biophysical model for

cell and tissue mechanics with imaging of cell and tissue geometries in the basal layer and at the

basal-suprabasal interface. We find that at early developmental stages, embryonic (E) day 13.5/14.5,

there is little mechanical distinction between basal and suprabasal layers allowing basal cells to easily

place cells suprabasally either through perpendicular division or through rapid detachment assisted

by fluctuations from cell divisions. Around E15.5, a mechanical boundary between the basal and

suprabasal layers emerges, and basal cells must now undergo profound transcriptional

reprogramming– e.g., commit to delamination – in order to cross this boundary. How is this decision

to commit regulated? Our data suggests an elegant feedback mechanism involving Notch signaling.

Tissue crowding and stiffening trigger a transient Notch signal in a small subset of cells, causing

these cells to execute a transcriptional switch to complete delamination. Collectively, our study

shows that changes in tissue mechanics that drive intercompartment boundary formation are essential

determinants for a switch from early embryonic to a mature differentiation and multilayering

strategy. Moreover, we identify Notch signaling as an integrative feedback relay to precisely time the

transition between the mechanically separated basal and suprabasal compartments within the mature

stratified epithelium.

Results

Basal cells employ distinct embryonic stage-specific strategies to generate and

compartmentalize suprabasal layers

To understand which mechanisms are used for epidermal multilayering, we employed a whole

embryo live imaging system [9,13] to visualize long term (over 8 h) cell dynamics at E14.5 and E15.5

(Fig. 1A). While perpendicular divisions were prevalent at E14.5 as previously reported [8], parallel

oriented cell divisions became predominant at E15.5 with the overall frequency of cell divisions

decreasing (Fig. 1B). As reported [6], cell divisions also occurred suprabasally at E14.5 but much less

at E15.5 (Fig. 1B, Supplementary Fig. 1A). Imaging of fixed tissues showed that significantly more

undifferentiated cells were found in the first suprabasal layer at E14.5 than at E15.5 (Supplementary

Fig. 1B), as detected by the absence of the differentiation marker Keratin (K)10 [14] in suprabasal cells.

A fraction of these suprabasal undifferentiated cells were still proliferating at E14.5 but not at E15.5

(Fig. 1C). This suggested that differentiation is not associated with upward movement at early stages.

We then asked whether cell delamination dynamics also changed as a function of developmental

stage. At E14.5, detachment of cells from the basement membrane was regularly observed, with

upward movement lasting around 120 minutes. This detachment was frequently associated with a cell

division, either occurring after a perpendicular division or followed by a mitosis in the suprabasal

layer (Fig. 1D; Supplementary Movie 1). In contrast, at E15.5 delaminations were substantially

slower, lasting at least 300 minutes, and were not associated with divisions (Fig. 1E; Supplementary

Movie 2). The geometry of delaminating cells also changed; at E14.5 cells rounded up for

delamination, while at E15.5 cells transitioned instead into axially elongated and basally constricted,

wedge-like shapes prior to delamination (Fig. 1D-F).

To better understand how these changes in division and delamination strategies relate to cell

differentiation, and to identify the mechanisms involved, we turned to fixed tissue imaging. Previous

work has implicated the desmosomal cadherin Desmoglein1 (Dsg1) in delamination [15]. At E14.5 no

Dsg1 or K10-positive differentiating cells were detected in the basal layer (Fig. 1G). In contrast, at

E15.5 Dsg1+ K10+ cells were now observed basally, indicating that these cells had already initiated

commitment towards differentiation. These double-positive basal cells exhibited enhanced

localization of Dsg1 (Fig. 1H) and E-cadherin (Supplementary Fig. 1C) particularly at apical

interfaces, displaying wedge-like shapes similar to delaminating cells seen in live imaging (Fig. 1H).

To precisely quantify the shape changes of these K10+-basal cells at E15.5, we performed super-

resolution imaging of skin whole mounts. K10+ basal cells were still attached to the basement

membrane as shown by $\alpha6\beta4$ integrin staining, (Fig. 1I), and displayed a wedge-like shape

characterized by decreased basal adhesion length and increased apical adhesion surface (Fig. 1I).

Interestingly, integrins were enriched at these narrow basal footprints, indicating that differentiation

is initiated prior to complete cell detachment from the basement membrane (Fig. 1J).

Collectively, these live imaging experiments are consistent with earlier work on fixed samples

showing that at early developmental stages embryos engage both perpendicular cell divisions and

rapid delaminations associated with suprabasal divisions to generate suprabasal layers [6-8,10]. Our data

further indicates that at E15.5 and E16.5 embryos employ a substantially slower, division-

independent delamination process associated with remodeling of cell adhesion and changes in cell

shape to position cells suprabasally.

A biophysical model predicts that in the presence of a mechanical boundary, delamination depends on the mechanical properties of the delaminating cell

To understand the mechanistic basis for this switch in multilayering strategy, we turned to quantitative mechanical modelling. For this we utilized open-source 3D vertex model code [16] and extended a previously published 3D vertex model of the epidermis [13] (see Supplementary Methods). Similar models were recently used to study clonal dynamics of tumor cells in stratified epithelia [17] and initiation of multilayering in single-layered epithelia [18]. The model consisted of suprabasal cells (brown), basal cells (light blue) and a basement membrane (dark blue; Fig. 2A). A key model feature was basal layer rigidity, governed by cell shape. We therefore parameterized this by Δs, the distance in shape space to the fluid-solid transition [16,19,20] (Fig. 2A, inset). The model also included heterotypic interfacial tensions between basal and adjacent suprabasal cells (σ_a; Fig. 2A, magenta), and a potentially wetting interfacial tension between basal cells and the basement membrane (σ_b; Fig. 2A green). Previous work [13] had shown that these three "tissue background" parameters (σ_a, σ_b, and Δs) generate a robust mechanical model that is capable of capturing key features of the multilayered tissue architecture observed *in vivo*. A central feature of this mechanical model is generation of a mechanical boundary between the basal and suprabasal layer that ensures that small active mechanical fluctuations do not induce mixing between basal and suprabasal cells, and the layers are robustly separated, as observed at E15.5 onwards.

We first sought to predict whether the presence of a mechanical boundary could explain the embryonic stage-specific differences in delamination speed and shape of the delaminating cell (see Fig, 1), similar to what has been theoretically predicted for cell extrusion in simple epithelia [21]. To this end we specifically altered interfacial tensions of the delaminating cell – lateral σ_{dl}, basal σ_{db}, and apical σ_{da} (Fig. 2B). Past experimental observations [9,22] suggested that basal layer rigidity Δs is also relevant for delamination. We therefore also allowed it to vary while keeping the other "tissue background" parameters σ_b and σ_a fixed. We found that for a cell to move upward into the suprabasal layer in the presence of small active fluctuations, it indeed needed only to change its interfacial tensions (Fig. 2C). We proceeded to precisely predict how changes in delaminating cell mechanics and the mechanical boundary itself alter delamination rates. To quantify these rates, we simulated a large ensemble of conditions (N>85) for each set of model parameters (3240 sets), and measured the fraction of cells that had delaminated in a fixed time window ($\Delta t = 2000$ simulation time units, estimated to be 20 – 200 min [23,24], see Supplemental Methods). We found that delamination rates are a complex function of the four variables (σ_{da}, σ_{dl}, σ_{db}, Δs; Supplementary Fig. 2A). To parse the resulting large parameter space, we searched for a data collapse in

delamination rates [25], and further investigated whether the rates were consistent with an Arrhenius

process governed by an energy barrier, i.e. where the rate is proportional to $e^{-E/kT_{eff}}$, where $k\,T_{eff}$

sets the magnitude of active fluctuations [26-28]. We identified a specific combination of model

parameters where all delamination rate curves collapsed on top of each other (Fig. 2D; non-collapsed

data in Supplementary Fig. 2), indicating that a delaminating cell experiences a mechanical energy

barrier to move upwards. Further, this barrier can be modulated by the mechanical properties of the

delaminating cell, in the following form:

165
$$E \;=\; F(\,\sigma_{da} \;-\; \sigma_{db} \;+\; c\,\varDelta s\,)$$

Here F is a monotonically increasing function of its argument (see Supplemental Methods). The first

two terms indicate that the energy barrier depends on the difference between the basal and apical

interface tensions of the delaminating cell. To decrease the energy barrier and delaminate more

efficiently, a delaminating cell needs to increase the tension (i.e. decrease the wetting) with the

basement membrane, while also decreasing its apical interfacial tension with the suprabasal cells.

This is consistent with the experimentally observed emergence of a wedge-like cell shape and apical

Dsg1 enrichment from E15.5 onwards (see Fig. 1G-I). The third term indicated that the energy

barrier increases with increasing surrounding basal layer rigidity $\varDelta s$. Interestingly, altering the

interfacial tension σ_{dl} of the delaminating cell did not change the delamination rate (Fig. 2D).

Overall, these results indicated that the non-mixing of basal and suprabasal cells observed

experimentally at E15.5 and E16.5 (see Fig. 1) could be explained by a mechanical boundary

between these cell layers, and that cells would be able to cross this barrier by changing their own

mechanical properties.

A mechanical boundary emerges at E15.5 impacting delamination.

Next, we investigated whether such a mechanical intratissue boundary emerges during development.

To this end we performed detailed quantitative morphometry of basal cells across developmental

stages. 3D segmentation revealed substantial transformations of cell shapes: whereas basal cells at

E14.5 showed considerable shape heterogeneity and a high degree of shape deformation, at E15.5

cells shortened and became more cuboidal, and finally transitioned into homogenous, prolate cell

shapes at E16.5 (Fig. 3A-C). To infer changes in interfacial tensions from these morphological

transitions, we built on previous work suggesting that an apical curvature angle γ and cell height h

reflect a force balance along each interface [29,30]. Further, interfacial properties – orientation angle of

the lateral interface (β) and the roughness of the apical interface (R) - depend on heterotypic

interfacial tensions [31,32]. We therefore measured these observables *in vivo* in the basal layer (Fig. 3D-

H). These measurements revealed a decrease in cell height between E15.5 and E16.5, and non-

monotonic behaviors for the lateral interface orientation angle from E14.5 to E16.5 (Fig. 3E),

consistent with the observed stage-specific cell shape transitions (Fig 3A-C). As in previous work, R

quantifies the tissue-scale roughness – variations between the heights of the lateral junctions across

the basal layer (Fig. 3F) – which are distinct from the cell-scale curvatures defined by γ (Fig. 3G).

This can be seen by linking the images in Fig 3G to the data in Fig 3E and H - at E15.5 the cell-scale

curvature γ is high but over larger length scales the apical surface is relatively flat (low R), while at

E14.5 the cell-scale curvature γ is low but with a higher tissue-scale roughness R. At E16.5, γ is even

higher than at E15.5, while R is lower that E15.5. Collectively these shape changes were indicative of

dynamic, stage-specific changes in apical heterotypic interfacial tension, wetting at the basement

membrane, and overall tissue stiffness.

We then proceeded to integrate the experimental quantifications with the 3D vertex model by

converting experimental measurements to simulation units, followed by model simulations across a

wide range of parameter conditions (Fig. 3D, I, J). For three observables (R, β, h), we discovered that

the data could be collapsed as a simple function of the model parameters (Fig. 3I; Supplementary Fig.

3.A-C). As a result, the average value of each *in vivo* observable (Fig. 3E, H) specified a point on the

y-axis of the respective simulation plot (Fig. 3I), and therefore generated a unique constraint on the

model parameters. Although the curved interfaces described by γ are not explicitly embedded in 3D

vertex models, we can use the assumption of force balance[29,30] to write γ directly as function of σ_b

and Δs (Fig. 3J), and thereby directly extract a fourth constraint on vertex model parameters.

Collectively this benchmarking process of *in vivo* measurements to simulations resulted in a set of

four constraint equations for the three unknowns $(\sigma_b, \sigma_a, \Delta s)$ at each developmental stage (Fig. 3D);

we solved this system of equations for the best-fit model parameters at each stage using a standard

over-constrained solver[33] (Fig. 3K; see Supplementary Methods).

These data demonstrated that at E14.5, the basal layer is nearly fluid-like, with very low tissue

stiffness parameterized by Δs (Fig. 3N). At this stage the interfacial tension between basal and

suprabasal cells is low, coupled to a positive interfacial tension at the cell-basement membrane

interface, suggesting only low levels of integrin-based wetting. At E15.5, the tension at the basement

membrane interface became nearly zero, suggesting wetting behavior (Fig. 3K). This was

accompanied by increased interfacial tension at the basal-suprabasal interface, as well as increased

basal layer stiffness (Fig. 3K). At E16.5, similar trends remained, while tissue stiffness increased

further (Fig. 3K). Combined, these changes generated a four-fold increase in the magnitude of the

mechanical energy barrier for a basal cell to delaminate from E14.5 to E15.5 (Supplementary Fig.

3F-H).

Building on our understanding from the data collapse in Fig. 2, we were then able to collapse the

rates of delamination as a function of the differences in the interfacial tensions between the

delaminating cell and the background basal cells at the apical ($\Delta\sigma_{da,a}$) and basal ($\Delta\sigma_{db,b}$) interfaces

(Fig. 3L; Supplementary Methods Fig. 3C, see Supplemental Methods). This analysis showed that at

E14.5, the probability of a standard basal cell (0.0 on the x-axis of Fig 3L) delaminating during the

time window Δt was less than 20%, but only a small change in basal and apical tensions (~15%, 0.15

on the x-axis in Fig 3L) was sufficient to generate a massive increase in delamination probability,

increasing it to nearly 100% in the same time window. In contrast, at E15.5 and E16.5, a basal cell

must change its mechanical properties substantially (by appr. 40%, 0.4 on the x-axis of Fig. 3L) in

order to generate a similar increase in delamination probability.

Collectively, these data suggest that a mechanical boundary forms between the basal and suprabasal

layers at around E15.5, thus requiring cells to have a robust mechanism to commit to delamination

and overcome this barrier, while such a mechanism is not necessary at E14.5.

239   A committed basal population characterized by Notch activation emerges at E15.5

To identify the molecular driver for this predicted mechanical state change required for delamination

at E15.5 and later, we investigated the evolution of epidermal cell states using single cell RNA

sequencing (scRNAseq; Fig. 4A). We obtained single-cell suspensions of skin from E13.5, E14.5,

E15.5. and E16.5 embryos, isolated and sequenced in parallel to avoid batch effects (Fig. 4A). The

quality-controlled and filtered dataset, obtained from 2 biological replicates/stage, comprised of

>15,000 cells/stage that were annotated based on known marker genes (Supplementary Fig. 4A, B).

Focusing on the epidermal (non-hair follicle) clusters, we identified cycling and non-cycling basal

cells as well as suprabasal cell states (Supplementary Fig. 4C, D). Leiden clustering revealed that cell

states were clustered separately according to their developmental stage, indicating that key cell states

(basal and suprabasal cells) underwent substantial transcriptional changes in transcriptomes while

progressing through development (Fig. 4B).

Closer inspection of key differentiation genes such as *Krt1*, *Krtdap* and *Notch1* revealed that at E14.5

their expression was restricted to the suprabasal cell populations. In contrast, at E15.5 and E16.5 a

subset of basal cells also showed expression of differentiation genes (Fig. 4C), in agreement with

some basal cells showing K10 and Dsg1 expression at this stage (Fig.1G, H) . To conclusively

identify a committed basal cell population, we examined a Harmony integrated dataset of E14.5-

E16.5 epidermal cells. Leiden clustering of this integrated dataset revealed a distinct population of

cells characterized by expression of both basal stem cell (e.g. *Trp63* and *Itga6*) and terminal

differentiation genes (Supplementary Fig. 4E, F), indicating that only from E15.5 onward, a subset of

basal cells commit to differentiation. Next to this hybrid transcriptional signature of differentiation

and stemness, this subset of committed basal cells also showed enhanced expression of the suprabasal

adhesion proteins E-cadherin and desmoglein (Fig. 4D), consistent with a downregulation in

heterotypic interfacial tension between the committed cell and suprabasal cells predicted by the

model (Fig. 2D and 3O).

As Notch activity has been shown to be critical for terminal differentiation in keratinocytes [34], we

examined Notch pathway activity specifically in the basal layer. Receptor-ligand enrichment analyses

(CellPhoneDB; [35] predicted increased basal activity of especially Notch1-Jagged2 signaling from

E15.5 onward (Fig. 4E). Indeed, immunofluorescence analysis revealed that Notch activity (cleaved

intracellular domain; N-ICD), was detected in single cells only in the basal layer of E15.5 and E16.5

epidermis, but not at E14.5 (Fig. 4F, G). Intriguingly, these Notch1-active cells were also positive for

the differentiation marker K10 and displayed a wedge-shaped morphology (Fig. 4F), indicating that

these cells are committed to differentiation and undergoing delamination as observed by the

K10/Dsg-1 positive cells (Fig.1G, H)

Collectively these data indicated the emergence of a committed basal cell state characterized by a

hybrid profile of stemness and differentiation, a change in cell shape, a switch to a suprabasal cell

adhesion profile and Notch activity.

Generation of the mechanical boundary at E15.5 is concomitant with stiffening of the basement

membrane and rigidification of the basal layer

We hypothesized that maturation of the tissue and specifically the basement membrane could drive

the emergence of the mechanical barrier at E15.5. Combined analysis of the scRNAseq data (Fig.

5A) and quantitative multiplex tissue imaging using Imaging Mass Cytometry [36], revealed an

increase in basement membrane maturation and cell-matrix adhesion (Fig.5A-C; Supplementary Fig.

5A). Importantly, collagen IV, a key component of the basement membrane that is essential for its

rigidification and stabilization, was increased from E14.5 to E16.5 (Fig. 5B-C), suggesting

maturation of the BM from E14.5-16.5. In agreement, the hemidesmosomal α6β4 integrin was also

increased during this transition, and became more and more confined to the epidermal-dermal

junction (Fig. 5B, C; Supplementary Fig. 5A). Consistently, atomic force microscopy-mediated force

spectroscopy confirmed a significant stiffening of the basement membrane from E14.5 to E16.5 (Fig.

5D).

Based on observations in other systems that link changes in tension to mechanical boundary

formation [9,16,19,20,22,37,38] we speculated that this change in tissue mechanics promoted establishment

of a mechanical boundary between basal and suprabasal cells. Quantification myosin phosphorylation

(pMLC2) indicated increased actomyosin tension at apical cell-cell junctions and specifically at the

interface between basal and suprabasal cells at E15.5 compared to E14.5 (Fig. 5E), providing further

support for the geometry-based tension inference indicating emergence of a mechanical boundary

(Fig. 3N). Furthermore, analyses of cell shapes and tissue dynamics from live imaging showed that

the cells in the basal layer of E14.5 embryos were larger and more irregular shaped compared to cells

at E15.5, as already seen in fixed tissue imaging (Fig. 5F, G; Fig. 3 A-C; Supplementary Movie 3). In

agreement, at E15.5 cell density was increased, (Fig. 5F, H), with decreased mobility of the basal

layer cells as quantified by particle image velocimetry (Fig. 5F, I; Supplementary Movie 3), in line

with previous *in vitro* work implicating a jamming transition in initiation of delamination [9]. Together,

these changes were consistent with a tissue-scale rigidity transition in the developing epidermis that

occurs from E14.5-16.5, from an initially floppier basement membrane and a fluid-like basal layer to

a stiff basement membrane and a jammed basal layer with enhanced cell-matrix adhesion and apical

actomyosin tension.

Tissue rigidity and cell shape changes drive Notch-dependent delamination and control of

tissue crowding

We next investigated whether the changes to tissue rigidity are sufficient to induce basal layer

jamming and the emergence of a mechanical barrier. For this we mimicked tissue maturation *in vitro*

by modulating cell density and substrate rigidity by engineering micropatterned adhesive islands on

polyacrylamide hydrogels of defined stiffness (Fig. 6A). Culturing primary epidermal progenitors on

micropatterned islands revealed that increasing substrate stiffness under confinement not only

substantially enhanced differentiation (Fig. 6B), but, importantly, promoted efficient suprabasal

separation of these differentiated cell populations from the undifferentiated cells that remained

basally positioned (Fig. 6C), indicating that increasing substrate stiffness is sufficient to induce

boundary formation.

As the emergence of basal committed cells was associated with Notch activity (Fig. 4D), we then

asked whether basal layer crowding was sufficient to induce Notch signaling. For this, we performed

live imaging of primary epidermal progenitors isolated from Notch1 activity reporter mice [39] to

identify whether and when Notch signaling is activated in basal cells triggered to differentiate and

form a suprabasal layer (Fig. 6D). As expected, all progenitors were initially Notch-reporter negative,

but upon application of calcium to trigger differentiation and multilayering, individual cells gradually

switched on the reporter followed by delamination (Fig. 6D; Supplementary Movie 4). Intriguingly,

prior to turning on Notch1 signaling these cells were more elongated and resided in denser tissue

regions than reporter negative-cells (Fig. 6D), thus indicating that crowding and increased cell-cell to

cell-matrix contact area promoted Notch activation.

We then asked whether Notch1 transcriptional activity is necessary for cell fate commitment and

delamination by deleting the Notch transcription factor, Rbpj, in the epidermis (Rbpj-KO; [40,41]). At

E14.5, the epidermis of Rbpj-KO mice appeared largely comparable to control littermates (Fig. 6E).

At E15.5, and more prominently at E16.5, the epidermis became structurally more abnormal,

characterized by a specific thinning of the suprabasal differentiated layers (Fig. 6F). Moreover, basal

cells were increased in height and became more columnar in E15.5 and E16.5 Rbpj-KO mice

compared to controls (Fig. 6G), accompanied by higher cell density (Fig. 6D), indicating increased

crowding of the basal layer. To examine if this increased crowding was due to reduced commitment

to differentiation or decreased delamination, we quantified differentiating, delaminating cells. We

observed emergence of the committed basal cell population in both control and Rbpj-KO epidermis

(Fig. 6H), indicating that the initial commitment to differentiate does not depend on Notch

transcriptional activity. Instead, we observed an increase in the number of wedge-shaped committed

cells undergoing delamination at this stage (Fig. 6H), suggesting an inability to overcome the

mechanical energy barrier and move suprabasally.

To identify the molecular mechanism by which Notch transcriptional activity regulates efficient

delamination, we performed scRNAseq analysis across E14.5-E16.5 from control and Rbpj-KO

mouse skin. Overall, basal cell populations were transcriptionally similar between controls and Rbpj-

KO mice (Fig. 6I). However, as described previously [34], Rbpj-KO cells of all developmental stages

displayed impaired terminal differentiation (Fig. 6I; Supplementary Fig. 6A). Intriguingly, at E15.5 a

distinct Leiden cluster representing committed basal cell population emerged, mainly composed of

Rbpj-KO cells (Fig. 6I, J; Supplementary Fig. 6B). Consistently, at E16.5 the Leiden cluster

consisting of committed cells was dominated by Rbpj-KO cells (Fig. 6I, J; Supplementary Fig. 6B),

validating the immunofluorescence observations that Rbpj-KO still commit to initial differentiation

but cannot efficiently delaminate, leading to accumulation of committed basal cells.

To identify master transcriptional switches downstream of Notch that drive the final step of

delamination, we performed differential gene expression analyses of committed cells between control

and Rbpj-KO at E15.5 and E16.5. This analysis identified two interesting significantly altered

regulators in Rbpj-KO committed cells: the zinc finger protein Zpf800, a master regulator of

intestinal stem cell differentiation [42], was not upregulated (Fig. 6K left panel; Supplementary Fig. 6C,

D). At the same time Rbpj-KO basal cells failed to downregulate the Ep300 Inhibitor of

Differentiation (Eid1) (Fig. 6K, right panel). Eid1 acts as a transcriptional repressor through

inhibition of p300-mediated H3K27 acetylation (H3K27ac), thus preventing the activation of

enhancers that promote differentiation [43,44]. Indeed, RNA *in situ* hybridization and quantitative

immunofluorescence confirmed that while control embryos downregulated Eid1 in the suprabasal

layers and, as a consequence, showed substantially reduced H3K27ac levels in delaminating cells and

suprabasally, Rbpj-KO embryos maintained both high Eid1 and H3K27ac in delaminating cells and

suprabasal layers, in line with reduced differentiation (Fig. 6L).

Collectively, these data demonstrate that stiffening of the substrate is sufficient to drive basal layer

jamming and the formation of a mechanical barrier between basal and suprabasal compartments. This

basal layer jamming and crowding then triggers Notch signaling required for completion of

delamination and control of basal layer density.

Discussion

Here we show that the developing epidermis employs different strategies for multilayering that are

driven by changes in tissue mechanics. Early during development of this stratified epithelium, there

is no substantial mechanical energy barrier between basal and suprabasal cell layers. This allows fast

upward movement of basal cells independent of basal fate commitment by either dividing

perpendicularly or detaching quickly in order to rapidly establish a stratified tissue. Similar

mechanical-instability-driven mechanisms have been proposed by previous modeling [18,21]. At later

stages (E15.5 and beyond), maturation and stiffening of the basement membrane enhance basal cell-

matrix adhesion and crowding to rigidify the tissue, generating a substantial mechanical boundary

between basal and suprabasal cells. In order to cross this boundary, cells from this stage on must now

commit to differentiation and undergo subsequent transcriptional rewiring, controlled by Notch

activation, to complete delamination. Interestingly, this delamination process in the matured

epidermis is substantially slower than the rapid detachment observed by us and others in early

embryonic development[9,10]. As Notch activation itself is regulated by tissue rigidification and

crowding, as also found in other systems[45,46], it suggests an elegant control mechanism for

homeostasis in an open niche not characterised by clear geometrical cues: tissue crowding itself is

sufficient in a rigidified tissue to trigger Notch activation necessary to control the rate and timing of

delamination. This feedback loop thus allows accurate control over the densities of both basal and

suprabasal layers in coordination with compartmentalization of stem cells and differentiated progeny.

Many open questions remain. A key open question is how the system decides which subset of cells in

the basal layer becomes committed and Notch active. Previous work has suggested that since Notch

is a surface receptor, its activity depends on tissue crowding together with associated cell shape and

surface area changes that regulate the density of such receptors [46-48]. This could explain the

emergence of Notch active cells in elongated cells within dense regions – these cells would have

transiently maximized surface area to receive Notch ligand signal from their neighbors. Previous

work has investigated a 2D vertex model where cell fate decisions are coupled to cell areas [49], which

may be a useful starting point for predicting features we observe here. Future modeling work should

focus on understanding how changes to tissue stiffness and cell shape induce Notch signaling in a

'correct' subset of basal cells (e.g. they are spaced out and the right fraction depending on the degree

of crowding), and on adding the effects of differently oriented cell divisions in basal layer — in

different developmental stages – to the 3D model.

Downstream of Notch signaling, it will be intriguing to understand how the transcriptional

reprogramming, involving Eid1and Zfp800 associated with global changes in H3K27ac translates to

control of delamination timing. It is interesting to note that the direct downstream targets of Eid1- the

histone acetyltransferases P300 and CBP - are evolutionarily conserved maintenance factors of

lineage-specifying core-transcriptional networks. They function as transcriptional rheostats that

control global H3K27ac levels, where the duration of their activity determines the long-term impact

on chromatin state [50,51]. Upon their inhibition, >8h of hypoacetylation is required to trigger a

permanent switch and "point of no return" to H3K27me3-mediated chromatin silencing [50].

Intriguingly, this is the approximate duration of the slower delamination process after E15.5. Thus,

Notch may function as a mechanical sensor to ensure that terminal differentiation is not initiated

prematurely in areas of low basal cell density. Precise coordination of onset of terminal

differentiation with basal stem cell layer density is likely to be a critical feature in dynamically self-

renewing tissues with an "open stem cell niche". Collectively, we propose that in such tissues

mechanical boundaries are required to separate biological activities of cycling stem cells and post-

mitotic terminally differentiated cells. Here, precisely controlled cell state transitions control the

transit between the compartments to robustly maintain tissue homeostasis and barrier integrity.

Methods

Mice

Membrane-targeted Tomato reporter mice (R26R$^{mT/mG}$), nucleus-targeted Tomato reporter mice (R26R$^{nT/nG}$) and high-activity Notch1 activation-dependent reporter mice (N1IP: CreHI) were from were from JAX laboratories (stock #007676, #:023035 and #027234 respectively). Rbpj floxed mice were obtained from JAX (stock # 034200) and crossed with the K14-Cre line [40] to obtain epidermis-specific depletion. Embryos were staged based on vaginal plug (taken as embryonic day 0) and limb morphology. All mouse studies were approved and carried out in accordance with the guidelines of the Ministry for Environment, Agriculture, Conservation and Consumer Protection of the State of North Rhine-Westphalia (LANUW; 81-02.04.2022.A065; and 2024-155), Germany.

Ex vivo embryo imaging

Imaging was performed essentially as described earlier [9,13]. Briefly, embryos were dissected at embryonic stages E14.5 or E15.5 and immobilized on custom-built imaging chambers engineered on top of Lumox-teflon imaging plates to allow gas exchange (Sarstedt). Whole embryos were immersed in epidermal growth medium (CnT-PR, CELLnTEC Advanced Cell Systems) supplemented with calcium-depleted 10% fetal bovine serum (HyClone/Thermo Fisher), 20 U/ml penicillin-streptomycin (Thermo Fisher), 20 mM Hepes, 1.8 mM CaCl2, and 10 ng/ml mouse recombinant EGF (Novus; NBP2-34952). Imaging was performed using an Andor Dragonfly 505 high-speed spinning disk confocal microscope (Oxford Instruments) with an inverted Nikon Eclipse Ti2 microscope (Nikon) equipped with 488 nm and 546 nm lasers, an Andor Zyla 4.2 sCMOS camera and an environmental chamber set at 37°C, 5% CO2 using a 40X dry objective. Images were acquired using Fusion 2.0 software (Andor). After acquisition, the image series were 3D drift corrected using the ImageJ plug-in Fast4Dreg [52] combined with manual registration, if required.

Immunofluorescence and confocal microscopy

For skin whole mounts, embryos were dissected from pregnant mice and directly fixed in pre-warmed (37°C) 4% paraformaldehyde (PFA) at room temperature for 3 h. After multiple PBS washes, the skin was carefully peeled using forceps and placed in blocking solution (0.5% Triton-X, 5% bovine serum albumin (BSA), 3% normal goat serum (NGS)) for 1 h at room temperature. For paraffin sections tissue biopsies were fixed in 4% PFA, embedded in paraffin and sectioned. Sections were de-paraffinized using a graded alcohol series and antigen retrieval was carried out using target

retrieval solution (Dako) at pH 6 or pH 9 in a pressure cooker. For cryosections, tissue samples were

placed unfixed into cryomolds, embedded in OCT (Tissue Tek) and allowed to harden on dry ice. 8

μm thick sections were cut, air-dried and fixed with 4% PFA. For both paraffin and cryofixed

sections, samples were blocked in 5% BSA, 3% NGS. For paraffin sections antibodies were diluted

in Dako antibody diluent (Agilent). For cryosections antibodies were diluted in blocking solution.

Cultured epidermal cells were fixed in 4% PFA for 15 min at room temperature After PBS washes,

samples were placed in 0.3% Triton-X, 5% BSA for one hour at room temperature after which

primary antibodies were diluted in the same solution. In all cases antibodies were incubated overnight

in moist chamber at 4°C. Following PBS washes, secondary antibodies were diluted in 1% BSA in

PBS and incubated for 1h at room temperature.

For super-resolution imaging of Actin (Phalloidin), K10, Dsg1, E-cadherin, and Integrin α6, the

embryos were fixed in pre-warmed 4% PFA at 37°C for 1 hour and rinsed with PBS. The back skin

was peeled using forceps and incubated in blocking solution (2% Triton-x, 5% BSA in PBS) for 24

hours at 37°C. Primary antibodies were diluted in the blocking solution and incubated 48 hours at

37°C on shaker followed by several washes with 2% Triton-X with a final overnight wash.

Secondary antibodies were diluted in blocking solution and incubated 24 hours at 4°C followed by

several washes with 2% Triton-X.

For pMLC2 staining embryos were fixed in 4% PFA dissolved in HBSS for 1 hour on a shaker at

room temperature. Back skins samples are then peeled and permeabilized with 0.02% PBS-Tween 20

for 30mins and blocked with blocking solution (5% NGS,1% BSA) for 2 hours before incubating

with primary antibody overnight in 4°C. Samples are then thoroughly washed with 0.02% PBS-

Tween 20, incubated with secondary antibodies at 1:400 dilution for 1 hour at room temperature and

washed. All tissue samples were mounted in Elvanol.

The following primary antibodies were used: guinea-pig anti-Keratin 10 (Progen, #GP-10; 1/300);

rabbit anti-p-MLC2 (p-Ser19; Cell Signaling, 3675; 1:200), rabbit anti-cleaved-Notch1 (Cell

signaling; #4147S; 1/50), mouse anti-Desmoglein 1/2 (Progen, #61002; 1/200), rabbit anti-TGM1

(360; Proteintech, #12912-3-AP; 1/300), rabbit anti P63 (Genetex #GTX102425; 1/100), guinea-pig

anti-Keratin 14 (Progen, GP-CK14; 1:300), mouse anti E-cadherin (BD Biosciences, #610181;

1/200), rabbit anti Cyclin B1 (Cell Signaling, #4138S; 1/200), mouse anti H3K27ac (Active Motif,

#39685, 1/200), rat anti integrin α6 (Biorad #MCA699; 1/200) . F-actin was detected using Alexa-

Fluor-conjugated phalloidin (1:500). Bound primary antibody was detected by incubation with

Alexa-Fluor 488-, 568- or 647 -conjugated antibodies (Invitrogen). Nuclei were counterstained with

4′, 6-diamidino-2-phenylindole (DAPI, Invitrogen). Slides were mounted in Elvanol.

Fluorescence images were collected by laser scanning confocal microscopy (LSM980; Zeiss) with

with Zeiss ZEN Microscopy software (Zeiss ZEN version 3.7) using a 40X water immersion

objective. Images were acquired at room temperature using sequential scanning of frames of 1 μm

thick confocal planes (pinhole 1). Images were collected with the same settings for all samples within

an experiment. For super-resolution imaging a 63X oil objective and the Airyscan 2 SR (super-

resolution) mode with default resolution settings were used. Laser power and image size were kept at

a minimum to avoid photo-bleaching. All airyscan images were 3D processed using the Zeiss

Airyscan Processing mode with auto filter and standard strength.

Embryonic epidermal vertex analysis

Membrane-tomato positive E14.5, E15.5 and E16.5 embryos were isolated, fixed immediately in

4%PFA over night at 4°C and washed in PBS. The back skin was then removed using micro-scissors

and forceps and permeabilized in PBS/0.5% Triton-X for 1h at RT. Nuclei were stained by

incubating the skin in DAPI diluted in DAKO antibody diluent for 1h at RT. The skin was then

mounted in Mowiol.

Confocal stacks were recorded at a voxel size of X: 0.361, Y: 0361, Z:0.299 using a Leica SP8

equipped with an APO CS2 63x/1.40 oil objective. To determine basal cell vertices, confocal stacks

were processed in Imaris and minimal XZ or YZ projection were exported using the extended section

view. Using the segmented line tool in ImageJ a line was drawn across the apical and basal vertices

of basal cells. Coordinates were extracted and "Julia" was used in Visual Code Studio to determine

cell height (h), apical and basal vertex distances (a) and the lateral interface orientation (angle beta).

The lines connecting the basal and apical vertices, respectively, were used to determine the apical

layer roughness (R) (See also Supplementary Methods). The apical curvature angle gamma was

determined manually using the angle tool in ImageJ.

Image analyses

For tissue-scale analysis, 2D segmentation were performed from live imaging data on

whole embryos or cultured epidermal progenitor monolayers using Cellpose4SAM (v4.0.8) [53]. 2D

cell shape parameters were measured using custom Python script and cell densities were measured

using ImageJ. For 3D analyses of cell shapes of delaminating cells from embryo live imaging, Imaris

11 was used to manually create 3D surfaces of delaminating and corresponding neighboring basal

cells. For 3D segmentation of basal cells from super resolution images, K10 negative basal cells were

manually segmented in 3D using Imaris. Meshes were exported from Imaris surfaces after which

shapes were analyzed by quantifying Spherical Harmonics [54] using a custom Python script.

In fixed tissue samples, basal committed cells and delaminations were identified based on the

expression of specific markers (K10 and/or Desmoglein) and their contact with the basement

membrane, and subsequently manually counted. Active-Notch positive cells were quantified

manually. The method was applied across all stages and conditions. Integrin α6, E-cadherin and

Desmoglein mean intensities were manually measured at the basal, apical and lateral membranes on

Image J. To quantify epidermis thickness, the distance between epidermal basement membrane and

most superficial point of the epidermis was measured. To quantify specifically the basal layer

thickness versus the differentiated layer thickness, the expression of specific markers (K10 and

desmoglein) has been used to distinguish the differentiated layers (K10+) from the basal layer.

To measure boundary straightness between the bottom and upper layers in micropatterns of soft and

stiff polyacrylamide (PAA) hydrogels, both a segmented line following the top of each nucleus in the

bottom layer and a straight line were drawn. The lengths of these lines were measured using ImageJ,

and the ratio of the straight line to the segmented line was calculated.

Eid1 RNA signal was quantified by segmenting suprabasal cells based on Keratin-DAP–positive

cells. Mean fluorescence intensity of Eid1 mRNA was measured using Fiji.

3D vertex model

A dynamic three-dimensional model of stratified epithelial tissues was developed using the 3D vertex

framework (Zhang and Schwarz, 2022). In the 3D vertex model, cells are represented as polyhedra,

each characterized by a homeostatic volume V_0 and surface area S_0. The geometry of each cell is

defined by its vertices, which serve as the degrees of freedom of the system. The modeled epithelium

consists of two main cell compartments. The upper compartment represents the suprabasal

differentiated cell layer, while the middle one-cell-thick layer corresponds to the basal stem cell layer.

Because the vertex model treats cells as space-filling polyhedra, we modeled the basement membrane

as an additional layer of interconnected "cellular network" directly attached beneath the basal layer. In

this framework, cells exert forces on one another through shape-based interactions governed by the

following energy functional :

$E = \sum_{\alpha} K_S (S_\alpha - S_0)^2 + \sum_{\alpha} K_V (V_\alpha - V_0)^2 + \sum_{\alpha\beta} \boldsymbol{\sigma}_{\alpha\beta} S_\alpha$ (1)

The first term of the energy functional describes the competition between contractile forces generated

by cell actomyosin networks (which enable cells to minimize their shape or surface area to resist their

environments), and adhesive forces mediated by cadherin-based, which promote the elongation of cell–

cell contacts during neighbors' exchange (T1s). The second term accounts for the mechanical

resistance of cells to volume fluctuations, as most cells are fluid-filled and resist changes in volume.

The third term incorporates heterotypic interfacial tension σ_a between the basal cells and suprabasal

cells, and the tension σ_b associated with integrin-based wetting behaviors at basal-basement membrane

interface.

Previous studies have shown that differential expression of adhesion molecules and cytoskeletal

components associated with cell fate specification (through variations in protein or gene expression)

can modify cell–cell interactions, leading to differences in adhesion strength and heterotypic interfacial

tension [9,13,55-63]. Such tension differences are known to play a central role in establishing and

maintaining sharp tissue boundaries [61]. Moreover, basal cells adhere to the basement membrane via

integrins, which introduce a wetting interaction characterized by σ_b [64]. The parameters K_S and K_V

denote the elastic moduli associated with the surface area and volume of the cells, respectively. Upon

non-dimensionalization, this energy functional allows the definition of a key order parameter known

as the target shape index, $s_0 = S_0 V_0^{-\frac{2}{3}}$, which controls the mechanical rigidity of the tissue. By tuning

s_0, the epithelial confluent tissue exhibits a rigidity transition from a solid-like to a fluid-like state, with

a critical value at $s_0 = 5.4$ [16,20].

To quantitatively predict the mechanical parameters and cellular mechanisms underlying mechanical

barrier formation between the basal and suprabasal layers, we first simulated the epithelium in the

absence of cell delamination. The system evolves according to overdamped Langevin dynamics,

where vertex positions are continuously updated to minimize the total mechanical energy:

$$\partial_t r_i = -\mu \nabla_i E_i + \sqrt{2D_t}\, \boldsymbol{\eta}(t). \qquad (2)$$

The left-hand side represents viscous drag forces, the first term on the right-hand side corresponds to

interaction forces derived from the energy potential, and the second term represents thermal

fluctuations in vertex positions r_i . Here $\boldsymbol{\eta}$ is an uncorrelated stochastic Gaussian white noise with

unit amplitude, $D_t = \mu k_B T_{\text{eff}}$, where μ is the mobility (inverse frictional drag), k_B Boltzmann

constant, and T_{eff} is an effective temperature parameter that sets the scale of the fluctuations. Unless

otherwise noted, for our simulations we choose $\mu = 1$, $k_B = 1$ and $T_{\text{eff}} = 0.002$. This choice of T_{eff}

allows fluctuations of interface lengths but is not sufficient to generate diffusion or fluidity in the
basal layer.

From the simulated model, we can compute the effective tensions governing different interfaces: a)
The homotypic tension between basal cells $\sigma_{bb} = 4K_S\Delta s_b$ where $\Delta s = s_\alpha - s_0$ measures the basal
layer stiffness (and $\Delta s = 0$ at the transition point); b) The effective heterotypic tension at the basal-
suprabasal interface $\Gamma_a = 2\sigma_a + 2K_s(\Delta s_s + \Delta s_b)$, where Δs_s the stiffness of the suprabasal layer; c)
The effective wetting tension $\Gamma_{bm} = 2\sigma_b + 2K_s(\Delta s_b + \Delta s_m)$ between basal cells and the basement
membrane (constituted of a cellular network with a small shape parameter). Unless otherwise
noted, $\Delta s_m = 0.26$, which ensures the basement membrane remains very stiff and solid-like.
589
590 For more detailed description see Supplementary Methods.
591

Single-cell RNA sequencing (scRNAseq)
E13.5, E14.5, E15.5 and E16.5 embryos from wild type, control or K14Cre-RBP-J floxed/floxed
mice (two embryos/genotype/stages) were collected simultaneously and back skins were peeled off.
Single-cell suspensions were generated by enzymatic digestion by incubating back skin tissues in a
mixture of DNAse I (40 µg/ml), Liberase (0.125 mg/ml; Roche Cat. NO. 05401119001), and papain
(30 U/ml; Worthington) in Minimum Essential Medium, Spinner modification (Gibco 11380) for 15
min (E13.5), 20 min (E14.5), 30 min (E15.5) or 40min (E16.5) at 37°C followed by mechanical
dissociation. The reaction was stopped with 22% FBS and single cells were washed with FACS
buffer (0.2% chelated FBS, 2 mM EDTA in 1x DPBS (Sigma Aldrich, D8537)). For wild type
sequencing, cells were labeled with BD Mouse Single-Cell Multiplexing Kit (Cat # 633793): E13.5
rep 1: Sample Tag (ST) 1; E13.5 rep 2: ST 2; E14.5 rep1: ST 3; E14.5 rep2: ST 4; E15.5 rep1: ST 5;
E15.5 rep2: ST 6; E16.5 rep1: ST 7, E16.5 rep2: ST 8. For K14Cre-Rbpj floxed and control
sequencing, surface biotinylation was used [65]. Briefly, cells were labeled with 50 µg/mL EZ-Link
sulfo-NHS-biotin (Thermo Fisher Scientific, 21217) on ice for 10 min, followed by incubation in
0.6 µg/ml TotalSeq-A PE streptavidin hashtag antibodies (BioLegend) for 30 min on ice as follows:
E14.5 Rbpj-KO 1: TotalSeq-A0951 PE Streptavidin 405251 ; E14.5 Rbpj-KO 2: TotalSeq™-A0952
PE Streptavidin 405253; E14.5 control 1: TotalSeq-A0953 PE Streptavidin 405255; E14.5 control 2:
TotalSeq-A0954 PE Streptavidin 405257; E15.5 Rbpj-KO 1: TotalSeq-A0951 PE Streptavidin
405251 ; E15.5 control 1: TotalSeq-A0953 PE Streptavidin 405255; E15.5 control 2: TotalSeq-
A0954 PE Streptavidin 405257; E16.5 Rbpj-KO 1: TotalSeq™-A0951 PE Streptavidin 405251 ;
E16.5 Rbpj-KO 2: TotalSeq-A0952 PE Streptavidin 405253; E16.5 control 1: TotalSeq-A0953 PE

Streptavidin 405255; E16.5 control 2: TotalSeq-A0954 PE Streptavidin 405257. Labeled cells were

loaded on a microwell cartridge of the BD Rhapsody Express system (BD) following the

manufacturer's instructions. Single-cell whole transcriptome analysis libraries were prepared

according to the manufacturer's instructions using BD Rhapsody WTA Reagent kit (633802; BD),

except for the Rbpj-KO and control sample tag libraries in which totalseq-ADT-oligo1 (5'-

TGCTCTTCCGATCTTGGCACCCGAGAATTCCA-3') primer in place of Sample Tag PCR1 and

Sample Tag PCR2 primers during the Sample Tag library amplification [65]. Libraries were sequenced

on the Illumina NextSeq 500 using High Output Kit v2.5 (150 cycles; Illumina) for 2×75-bp paired-

end reads with 8-bp single index aiming sequencing depth of >20,000 reads per cell for each sample.

scRNAseq analyses

Read Data QC and Mapping

Illumina sequencing results were demultiplexed and converted to FASTQ format using Illumina's

bcl2fastq software (version 2.20.0.422). For K14Cre-RBP-J floxed/floxed data a custom

demultiplexing script was used to assign barcodes to sample tags (rhapsody-demultiplex with

parameters '-w BD_CLS1_v2.txt BD_CLS2_v2.txt BD_CLS3_v2.txt -a

TGGCACCCGAGAATTCCA <fastq files> sample_tags.txt', and sample tags as described above).

Preprocessed read data was aligned to the mouse reference genome (GRCm39) and counted with

STAR [66] (version 2.7.11b) through its single-cell functionality STARsolo, using the following

parameters:

WT: `--runThreadN 16 --soloType CB_UMI_Complex --soloCellFilter None --outSAMtype BAM

SortedByCoordinate --soloFeatures Gene GeneFull GeneFull_Ex50pAS --soloAdapterSequence

NNNNNNNNNGTGANNNNNNNNNNGACA --soloCBmatchWLtype 1MM --soloCBposition

2_0_2_8 2_13_2_21 3_1_3_9 --soloUMIposition 3_10_3_17 --soloCBwhitelist BD_CLS1_v2.txt

BD_CLS2_v2.txt BD_CLS3_v2.txt --runRNGseed 1 --soloMultiMappers EM --readFilesCommand

zcat --outSAMattributes NH HI AS nM NM MD jM jI MC ch CB UB GX GN`

K14Cre-RBP-J floxed/floxed: `--soloType CB_UMI_Complex --soloAdapterSequence

NNNNNNNNNGTGANNNNNNNNNNGACA --soloCBposition 2_0_2_8 2_13_2_21 3_1_3_9 --

soloUMIposition 3_10_3_17 --soloBarcodeReadLength 0 --soloCellFilter None --soloCBwhitelist

BD_CLS1_v2.txt BD_CLS2_v2.txt BD_CLS3_v2.txt --soloCBmatchWLtype 1MM --

soloUMIfiltering MultiGeneUMI_CR --soloUMIdedup 1MM_CR --outSAMtype BAM

SortedByCoordinate --soloFeatures Gene GeneFull GeneFull_Ex50pAS --soloMultiMappers EM --

outSAMattributes NH HI AS nM NM MD jM jI MC ch CB UB GX GN --outSAMmultNmax 1 --

outMultimapperOrder Random --runRNGseed 1 --outFilterScoreMin 30 --outFilterMatchNmin 15`

Data Filtering, Normalisation, and Visualization

Preprocessing of the feature-count-matrix output by STARsolo was largely performed within the

scanpy [67]ecosystem, with the exception of scran normalization [68], which was performed in R (R Core

Team, 2017). All visualization was done using [69] functions. Default parameters were used unless

detailed otherwise.

For wild type:

Barcodes with less than 3000 UMI were discarded.

Only barcodes with less than 15% of counts stemming from mitotic genes were retained.

For K14Cre-RBP-J floxed/floxed and control:

Barcodes with less than 2000 UMI were discarded.

Only barcodes with less than 20% of counts stemming from mitotic genes were retained.

Only barcodes with less than 70% of counts stemming from hemoglobin genes were retained.

And barcodes were filtered for a complexity value above 0.75, where complexity is defined as the

log10-transformed number of expressed genes divided by the log10-transformed number of total

counts per barcode. Scran was used for normalization.

The separate samples were merged into a single object, G2M and S phase scores were assigned to

each cell using gene lists from [70] and the scanpy scanpy.tl.score_genes_cell_cycle function, and

regressed out of the gene expression matrix using sc.pp.regress_out. 3000 highly variable genes were

calculated from the normalized matrix (sc.pp.highly_variable_genes, flavor "seurat"), and the first

100 principal components were calculated with scanpy. Neighbourhood graph calculation (30

neighbors), UMAP dimensionality reduction [71], and Leiden clustering [72] were performed. scrublet [73]

and/or scDblFinder were then used for doublet prediction.

Contaminant or undesired cell populations such as clusters of cells that, either manually or by

scrublet/scDblFinder, could be classified as doublets based on their expression patterns, were

removed. For this, cells were clustered at resolution 0.05, and the cluster corresponding to the

epidermal cells were isolated. These were then reclustered at resolution 0.1 (wild type) and 0.2

(K14Cre-RBP-J floxed/floxed) as described above, and undesired clusters were removed. Finally,

PCA, nearest neighbor calculation, and UMAP dimensionality reduction were repeated for the clean

epidermis data. For WT, stages E14.5, E15.5, and E16.5 were additionally analysed separately with

batch-correction using Harmony [74] after cell-cycle regression.

Differential expression analysis

Differentially expressed genes were calculated using a pseudobulk approach [75], comparing aggregate

counts with 2 pseudoreplicates for each condition (pyDeSEQ2 0.4.8).

Ligand-receptor interaction analysis

We used CellphoneDB (v2.1.7 [35] with default parameters to analyse ligand-receptor interactions.

Mouse gene names were converted to human gene names prior to the analysis.

Gene Set Enrichment Analysis

We ranked differential expression results by pyDESeq2 shrunken log2-fold changes and used

GSEApy (v1.0.6 [76]) "prerank" to calculate the enrichment statistics.

Imaging Mass Cytometry

Primary antibodies from bovine serum albumin (BSA)- and azide-free formulations were conjugated

using the Maxpar X8 Labeling Kit (Standard BioTools) according to the manufacturer's protocol.

Stainings were performed on frozen cryosections according to the manufacturer's protocol

(Fluidigm). Briefly, 5 μm cryosections were prepared and stored at −80 °C until use. On the day of

the experiment, slides were equilibrated at −20 °C for 1 h and then fixed in 4% paraformaldehyde

(PFA) for 30 min at 4 °C. After washing in PBS, sections were incubated in blocking solution (3%

BSA) for 45 min at room temperature, followed by incubation with the antibody cocktail prepared in

0.5% BSA for 1 h in a hydration chamber at room temperature. Slides were then rinsed in 0.2%

Triton X-100/PBS, stained with Cell-ID Intercalator-IR (Fluidigm #201192A) for 30 min at room

temperature, washed in Milli-Q water, air-dried for 30 min, and stored until imaging. The samples

were laser-ablated at 200 Hz using a Hyperion imaging mass cytometry system (Fluidigm) coupled to

a Helios time-of-flight mass cytometer with a resolution of 1 μm. Raw MiniCAD Design (MCD) files

generated by the Hyperion Imaging System were first inspected using the MCD Viewer (Standard

BioTools) to verify signal intensity and overall acquisition quality, and then exported as TIFF images

for downstream analysis on Image J. ROIs delineating the basement membrane were drawn, and

signal intensity at the basement membrane was measured using ImageJ software.

Particle Imaging Velocimetry (PIV)

PIV analysis of whole-embryo live imaging was performed as previously described, with minor

adjustments [13]. Briefly, prior to analysis, each movie was processed as follows: a 3D drift correction

was applied using the Fast4DReg plug-in in Fiji and sagittal cross-sections containing the basal

layers were selected. A median filter of 2 pixels was applied in Fiji, followed by background

subtraction using a rolling ball radius of 50 pixels.The movies were then imported into PIVLab

version 2.62. Tissue flow tracking was restricted to the interfollicular epidermis. To exclude the hair

follicle placodes from the analysis, ROIs covering the placodes were manually drawn on the raw

images and imported into PIVLab. These masks were overlaid onto the flow field to exclude the

placode regions. To prevent tracking errors due to intensity variations, image contrast was enhanced

using CLAHE (Contrast Limited Adaptive Histogram Equalization) with a box size of 64 pixels.

Denoising was performed using a Wiener Noise Filter with a maximum pixel count of 3. Tissue flow

tracking was carried out using a multi-pass, windowed FFT-based approach, with window sizes of 64

and 32 pixels for the first and second passes, respectively, and a 50% overlap. The resulting vector

fields were visually inspected, and any erroneous vectors were manually removed from the analysis.

Culture of primary epidermal progenitor cells

Primary epidermal cells were isolated from newborn mice essentially as described [13]. Briefly, after

dissection, skins were treated with antibiotic/antimycotic solution (Sigma A5955) for 5 min, and

epidermal single cell suspensions were generated by incubating skin pieces in 0.8% trypsin for

30 min at 37 °C. Cells were suspended in Keratinocyte Growth Medium (MEM Spinner's

modification (Sigma), 5 µg/ml insulin (Sigma), 10 ng/ml EGF (Sigma), 10 µg/ml transferrin (Sigma),

10 µM phosphoethanolamine (Sigma), 10 µM ethanolamine (Sigma), 0.36 µg ml/ml hydrocortisone

(Calbiochem), 2 mM glutamine (Gibco), 100 U ml/ml penicillin, 100 µg ml/ml streptomycin (Gibco),

10% chelated fetal calf serum (Gibco), 5 µM Y27632, 20 ng ml/ml mouse recombinant vascular

endothelial growth factor (VEGF) and 20 ng ml/ml human recombinant fibroblast growth factor-2

(FGF-2) (all from Miltenyi Biotec) and seeded on cell culture plates coated with 30 µg ml^{-1} collagen

I (Millipore) and 10 µg ml^{-1} fibronectin (Millipore) for 1 h at 37 °C.

Micropatterned hydrogels

Circular micropatterns (100 μm diameter) were generated on glass coverslips activated by oxygen

plasma treatment for 3 min (PDC-002-CE Expanded Plasma Cleaner), as described previously [9].

Activated coverslips were incubated with 1× PLL (Surface Solutions) for 30 min at room

temperature, rinsed with 10 mM HEPES (pH 7.8), and subsequently incubated with PEG-SVA (50

mg/ml in 10 mM HEPES, pH 7.8) for 1 h at room temperature. Coverslips were rinsed with Milli-Q

water and air-dried before patterning by deep-UV (Primo; Alveole). PEG-cleaved regions were

rinsed with Milli-Q water and incubated with 50 μg/ml fibronectin for 30 min at room temperature.

In parallel, glass-bottom plates (Ibidi, #81218-200) were silanized. Polyacrylamide gels of defined

stiffness were prepared by placing a drop of acrylamide/bis-acrylamide solution between the

micropatterned coverslip and the silanized plate. After 45 min of polymerization, the coverslip was

removed, transferring the ECM micropatterns onto the gel surface. Primary epidermal progenitors

were then seeded onto the patterned hydrogels.

Live imaging of epidermal progenitor monolayers

Freshly isolated epidermal progenitors isolated from N1IP: CreHI reporter mice crossed with

R26R$^{mT/mG}$ mice to label the plasma membrane and detect Cre activity were grown to confluency in

Keratinocyte Growth Medium and plated on 35 mm glass bottom dish with 14 mm micro-well plates

(Cellvis; D35-14-1.5-N). Upon confluency, 200 μM Ca^{2+} was added to induce differentiation.

Imaging was performed using an Andor Dragonfly 505 high-speed spinning disk confocal

microscope (Oxford Instruments) with an inverted Nikon Eclipse Ti2 microscope (Nikon) equipped

with 488 nm and 546 nm lasers, an Andor Zyla 4.2 sCMOS camera and an environmental chamber

set at 37°C, 5% CO2 using a 40X dry objective. Images were acquired using Fusion 2.0 software

(Andor) with multiple positions and drift stabilizer.

Atomic Force Microscopy

For Atomic Force Microscopy (AFM), embryonic skin was dissected, immediately nitrogen frozen

and then embedded in OCT blocks. AFM measurements of the basement membrane were performed

on freshly cut 20 μm-thick cryosections using JPK Nano Wizard IV (Bruker Nano) atomic force

microscope mounted on an Olympus IX73 inverted fluorescent microscopy (Olympus) and operated

via JPK SPM Control Software v.5. Cryosections were equilibrated in PBS supplemented with

protease inhibitors and measurements were performed within 20 minutes of thawing the samples.

Triangular non-conductive Silicon Nitride cantilevers (MLCT, Bruker Daltonics) with a nominal

spring constant of 0,07 Nm^{-1} were used for the nanoindentation experiments. For all indentation

experiments, forces of up to 3 nN were applied, and the velocities of cantilever approach and

retraction were kept constant at 10 μms^{-1} ensuring an indentation depth of 500 nm. All analysis were

performed with JPK Data Processing Software (Bruker Nano). Prior to fitting the Hertz model

corrected by the tip geometry to obtain Young's Modulus (Poisson's ratio of 0.5), the offset was

removed from the baseline, contact point was identified, and the cantilever bending was subtracted

from all force curves. Force curves were filtered for quality and all valid forces curves were

analyzed.

RNA in situ hybridization

RNA in situ hybridization was performed using RNAscope technology according to the

manufacturer's instructions (ACD, RNAscope Multiplex Fluorescent v2 Assay, kit: RNAscope

Multiplex Fluorescent v2 Assay ACD, #323100). Briefly, paraffin-embedded tissue sections (5 μm

thick) were deparaffinized in xylene, rehydrated through an ethanol series, and incubated in hydrogen

peroxide for 10 min at RT. Sections were then subjected to antigen retrieval in RNAscope Target

Retrieval solution for 15 min at 100°C using a pressure cooker. RNAscope Protease Plus (ACD;

#322331) was applied to each section and incubated at 40°C for 30 min. The Eid1 RNAscope probe

(ACD; #1713581-C3) and Keratin-DAP RNAscope probe (ACD; #500671-C1) were hybridized

according to the manufacturer's protocol using the RNAscope Multiplex Fluorescent v2 Assay

(ACD; #323100), followed by signal detection with TSA Plus Fluorescein (Akoya,

#NEL741001KT), TSA Plus Cyanine 3 System (Akoya, #NEL744001KT). All hybridization and

amplification steps were performed in a HybEZ II oven (ACD). DAPI solution was incubated for 30

seconds at room temperature. Sections were mounted in Elvanol and imaged using a Zeiss LSM 980

confocal microscope equipped with a 40× water-immersion objective, using Zeiss ZEN software

(version 3.5).

Statistics and reproducibility

Statistical analyses were performed using GraphPad Prism software (GraphPad, version 9) or in R

(version 4.2.2). Statistical significance was determined by the specific tests indicated in the

corresponding figure legends. Only 2-tailed tests were used. In all cases where a test for normally

distributed data was used, normal distribution was confirmed with the Kolmogorov–Smirnov test (α

= 0.05). All experiments presented in the manuscript were repeated at least in 3 independent

replicates.

Data availability

Sequencing datasets are available at GEO (GSE315748 and GSE315746). All analysis scripts,

custom code, and data that support the conclusions are available from the authors on request.

Acknowledgements

We thank Claudia Ortmeier, Manuela Haustein, Hermann vom Bruch, and Sandra Heising for expert

technical assistance, Silvia Fre (Institute Curie, France) for advice on the project and general support,

Fabien Bertillot for advice on image analyses, Yekaterina Miroshnikova for advice on

micropatterning, and the Max Planck Institute BioOptics and Sequencing core facilities for support

with experiments. Imaging Mass Cytometry was performed at the Cell Imaging and Cytometry Core,

Turku Bioscience Centre (Turku, Finland), with the support of Biocenter Finland. This work was

supported by European Union's Horizon 2020 research and innovation programme under Marie

Skłodowska-Curie grant agreement no. 101032331 (to CV), Research Council of Finland (332402)

and The Turku Collegium for Science, Medicine, and Technology (to GF), Academy of Finland

Center of Excellence BarrierForce (346131) and R'Life Programme consortium NucleoMech

(330033) (to JI and SAW) the Sigrid Juselius Foundation, Helsinki Institute of Life Science, and the

Max Planck Society (all to SAW), NSF-CMMI-1334611 (to SA and MLM), grant number 2023-

329572 from the Chan Zuckerberg Initiative DAF, an advised fund of the Silicon Valley Community

Foundation (to MLM), and the Syracuse University Office of Research Future Faculty Fellow

Program (to SA). CMN is supported by the Deutsche Forschungsgemeinschaft (DFG, German

Research Foundation: Germany´s Excellence Strategy – EXC 2030 – 390661388; SPP 1782 NI

1234/6-2, Project-ID 388932620; FOR 2743 NI 1234/7-1 Project-ID 678823; ANR/DFG NI 1234/9

(Project-ID 505673300) and the European Union NETSKINMODELS, COST Action no. CA21108.

Author contributions

CV, MA, and MR designed and performed experiments and analyzed data. SHAE and MLM

developed vertex model and performed all simulations and analyses. KK analyzed sequencing data.

SV and LCB supported live imaging and single cell sequencing experiments. AP performed and

analyzed immunofluorescence experiments, PZ assisted with imaging. GF and JI performed mass

cytometry experiment. CMN, MLM and SAW conceived and supervised the study, designed

experiments, and analyzed data. MLM and SAW wrote the paper. All authors commented and edited

the manuscript.

Competing interests statement

The authors declare no competing interests.

References:

1.   Kulukian, A., and Fuchs, E. (2013). Spindle orientation and epidermal morphogenesis. Philos
Trans R Soc Lond B Biol Sci *368*, 20130016. 10.1098/rstb.2013.0016.
2.   Niessen, C.M., Manning, M.L., and Wickstrom, S.A. (2025). Mechanochemical Principles of
Epidermal Tissue Dynamics. Cold Spring Harb Perspect Biol *17*.
10.1101/cshperspect.a041518.
3.   Tai, K., Cockburn, K., and Greco, V. (2019). Flexibility sustains epithelial tissue homeostasis.
Curr Opin Cell Biol *60*, 84–91. 10.1016/j.ceb.2019.04.009.
4.   Beumer, J., and Clevers, H. (2024). Hallmarks of stemness in mammalian tissues. Cell Stem
Cell *31*, 7–24. 10.1016/j.stem.2023.12.006.
5.   Williams, S.E., and Fuchs, E. (2013). Oriented divisions, fate decisions. Curr Opin Cell Biol
25, 749–758. 10.1016/j.ceb.2013.08.003.
6.   Damen, M., Wirtz, L., Soroka, E., Khatif, H., Kukat, C., Simons, B.D., and Bazzi, H. (2021).
High proliferation and delamination during skin epidermal stratification. Nat Commun *12*,
3227. 10.1038/s41467-021-23386-4.
7.   Box, K., Joyce, B.W., and Devenport, D. (2019). Epithelial geometry regulates spindle
orientation and progenitor fate during formation of the mammalian epidermis. Elife *8*.
10.7554/eLife.47102.
8.   Lechler, T., and Fuchs, E. (2005). Asymmetric cell divisions promote stratification and
differentiation of mammalian skin. Nature *437*, 275–280. 10.1038/nature03922.
9.   Miroshnikova, Y.A., Le, H.Q., Schneider, D., Thalheim, T., Rubsam, M., Bremicker, N.,
Polleux, J., Kamprad, N., Tarantola, M., Wang, I., et al. (2018). Adhesion forces and cortical
tension couple cell proliferation and differentiation to drive epidermal stratification. Nat Cell
Biol *20*, 69–80. 10.1038/s41556-017-0005-z.
10.   Williams, S.E., Ratliff, L.A., Postiglione, M.P., Knoblich, J.A., and Fuchs, E. (2014). Par3-
mInsc and Galphai3 cooperate to promote oriented epidermal cell divisions through LGN. Nat
Cell Biol *16*, 758–769. 10.1038/ncb3001.
11.   Cockburn, K., Annusver, K., Gonzalez, D.G., Ganesan, S., May, D.P., Mesa, K.R.,
Kawaguchi, K., Kasper, M., and Greco, V. (2022). Gradual differentiation uncoupled from
cell cycle exit generates heterogeneity in the epidermal stem cell layer. Nat Cell Biol *24*,
1692–1700. 10.1038/s41556-022-01021-8.
12.   Mesa, K.R., Kawaguchi, K., Cockburn, K., Gonzalez, D., Boucher, J., Xin, T., Klein, A.M.,
and Greco, V. (2018). Homeostatic Epidermal Stem Cell Self-Renewal Is Driven by Local
Differentiation. Cell Stem Cell *23*, 677–686 e674. 10.1016/j.stem.2018.09.005.

13. Villeneuve, C., Hashmi, A., Ylivinkka, I., Lawson-Keister, E., Miroshnikova, Y.A., Perez-
Gonzalez, C., Myllymaki, S.M., Bertillot, F., Yadav, B., Zhang, T., et al. (2024). Mechanical
forces across compartments coordinate cell shape and fate transitions to generate tissue
architecture. Nat Cell Biol *26*, 207–218. 10.1038/s41556-023-01332-4.
14. Moll, R., Franke, W.W., Schiller, D.L., Geiger, B., and Krepler, R. (1982). The catalog of
human cytokeratins: patterns of expression in normal epithelia, tumors and cultured cells. Cell
31, 11–24. 10.1016/0092-8674(82)90400-7.
15. Nekrasova, O., Harmon, R.M., Broussard, J.A., Koetsier, J.L., Godsel, L.M., Fitz, G.N.,
Gardel, M.L., and Green, K.J. (2018). Desmosomal cadherin association with Tctex-1 and
cortactin-Arp2/3 drives perijunctional actin polymerization to promote keratinocyte
delamination. Nat Commun *9*, 1053. 10.1038/s41467-018-03414-6.
16. Zhang, T., and Schwarz, J.M. (2022). Topologically-protected interior for three-dimensional
confluent cellular collectives. Physical Review Research *4*, 043148.
10.1103/PhysRevResearch.4.043148.
17. Sahu, P., Monteiro-Ferreira, S., Canato, S., Soares, R.M., Sanchez-Danes, A., and Hannezo,
E. (2025). Mechanical control of cell fate decisions in the skin epidermis. Nat Commun *16*,
8440. 10.1038/s41467-025-62882-9.
18. Fukamachi, S., Datta, R., Wuergezhen, D., Ichikawa, T., Yagasaki, R., Horiguchi, S.A.,
Leeaw, P., Omachi, K., Sada, A., Imayoshi, I., et al. (2024). Epithelium Stratifies via
Nucleation and Growth Induced by Foam-Geometric Instability. bioRxiv,
2024.2012.2026.630435. 10.1101/2024.12.26.630435.
19. Bi, D., Lopez, J.H., Schwarz, J.M., and Manning, M.L. (2015). A density-independent rigidity
transition in biological tissues. Nature Physics *11*, 1074–1079. 10.1038/nphys3471.
20. Merkel, M., and Manning, M.L. (2018). A geometrically controlled rigidity transition in a
model for confluent 3D tissues. New Journal of Physics *20*, 022002. 10.1088/1367-
2630/aaaa13.
21. Okuda, S., and Fujimoto, K. (2020). A Mechanical Instability in Planar Epithelial Monolayers
Leads to Cell Extrusion. Biophysical Journal *118*, 2549–2560.
https://doi.org/10.1016/j.bpj.2020.03.028.
22. Devany, J., Sussman, D.M., Yamamoto, T., Manning, M.L., and Gardel, M.L. (2021). Cell
cycle-dependent active stress drives epithelia remodeling. Proc Natl Acad Sci U S A *118*.
10.1073/pnas.1917853118.
23. Erdemci-Tandogan, G., Clark, M.J., Amack, J.D., and Manning, M.L. (2018). Tissue Flow
Induces Cell Shape Changes During Organogenesis. Biophys J *115*, 2259–2270.
10.1016/j.bpj.2018.10.028.
24. Serwane, F., Mongera, A., Rowghanian, P., Kealhofer, D.A., Lucio, A.A., Hockenbery, Z.M.,
and Campas, O. (2017). In vivo quantification of spatially varying mechanical properties in
developing tissues. Nat Methods *14*, 181–186. 10.1038/nmeth.4101.
25. Somendra, M.B., and Flavio, S. (2001). A measure of data collapse for scaling. Journal of
Physics A: Mathematical and General *34*, 6375. 10.1088/0305-4470/34/33/302.
26. Bi, D., Lopez, J.H., Schwarz, J.M., and Manning, M.L. (2014). Energy barriers and cell
migration in densely packed tissues. Soft Matter *10*, 1885–1890. 10.1039/c3sm52893f.
27. McQuarrie, D.A. (1976). Statistical Mechanics (Harper & Row).
28. Mombach, J.C.M., Robert D. , Graner, F., Gillet, G., Thomas, G.L., Idiart, M., and Rieu, J.P.
(2005). Rounding of aggregates of biological cells: Experiments and simulations. Physica A:
Statistical Mechanics and its Applications *352* (*525*). 10.1016/j.physa.2005.02.008.
29. Maitre, J.L., Berthoumieux, H., Krens, S.F., Salbreux, G., Julicher, F., Paluch, E., and
Heisenberg, C.P. (2012). Adhesion functions in cell sorting by mechanically coupling the
cortices of adhering cells. Science *338*, 253–256. 10.1126/science.1225399.

30. Pawlizak, S., Fritsch, A.W., Grosser, S., Ahrens, D., Thalheim, T., Riedel, S., Kießling, T.R.,
Oswald, L., Zink, M., Manning, M.L., and Käs, J.A. (2015). Testing the differential adhesion
hypothesis across the epithelial−mesenchymal transition. New Journal of Physics *17*, 083049.
10.1088/1367-2630/17/8/083049.
31. Sahu, P., Schwarz, J.M., and Manning, M.L. (2021). Geometric signatures of tissue surface
tension in a three-dimensional model of confluent tissue. New Journal of Physics *23*, 093043.
10.1088/1367-2630/ac23f1.
32. Sahu, P., Sussman, D.M., Rübsam, M., Mertz, A.F., Horsley, V., Dufresne, E.R., Niessen,
C.M., Marchetti, M.C., Manning, M.L., and Schwarz, J.M. (2020). Small-scale demixing in
confluent biological tissues. Soft Matter *16*, 3325–3337. 10.1039/C9SM01084J.
33. Press, W.H., Teukolsky, S.A., Vetterling, W.T., and Flannery, B.P. (2020). Numerical
Recipes: The Art of Scientific Computing (Cambridge University Press).
34. Blanpain, C., Lowry, W.E., Pasolli, H.A., and Fuchs, E. (2006). Canonical notch signaling
functions as a commitment switch in the epidermal lineage. Genes Dev *20*, 3022–3035.
10.1101/gad.1477606.
35. Efremova, M., Vento-Tormo, M., Teichmann, S.A., and Vento-Tormo, R. (2020).
CellPhoneDB: inferring cell-cell communication from combined expression of multi-subunit
ligand-receptor complexes. Nat Protoc *15*, 1484–1506. 10.1038/s41596-020-0292-x.
36. Giesen, C., Wang, H.A., Schapiro, D., Zivanovic, N., Jacobs, A., Hattendorf, B., Schuffler,
P.J., Grolimund, D., Buhmann, J.M., Brandt, S., et al. (2014). Highly multiplexed imaging of
tumor tissues with subcellular resolution by mass cytometry. Nat Methods *11*, 417–422.
10.1038/nmeth.2869.
37. Röper, K. (2012). Anisotropy of Crumbs and aPKC Drives Myosin Cable Assembly during
Tube Formation. Developmental Cell *23*, 939–953. 10.1016/j.devcel.2012.09.013.
38. Rustarazo-Calvo, L., Pallares-Cartes, C., Aguirre-Tamaral, A., Floris, E., Hingerl, M.,
Autorino, C., Maula Khan, A.U., Corominas-Murtra, B., and Petridou, N.I. (2025). Adhesion-
driven tissue rigidification triggers epithelial cell polarity. bioRxiv, 2025.2003.2018.644006.
10.1101/2025.03.18.644006.
39. Liu, Z., Turkoz, A., Jackson, E.N., Corbo, J.C., Engelbach, J.A., Garbow, J.R., Piwnica-
Worms, D.R., and Kopan, R. (2011). Notch1 loss of heterozygosity causes vascular tumors
and lethal hemorrhage in mice. J Clin Invest *121*, 800–808. 10.1172/JCI43114.
40. Hafner, M., Wenk, J., Nenci, A., Pasparakis, M., Scharffetter-Kochanek, K., Smyth, N.,
Peters, T., Kess, D., Holtkotter, O., Shephard, P., et al. (2004). Keratin 14 Cre transgenic mice
authenticate keratin 14 as an oocyte-expressed protein. Genesis *38*, 176–181.
10.1002/gene.20016.
41. Tanigaki, K., Han, H., Yamamoto, N., Tashiro, K., Ikegawa, M., Kuroda, K., Suzuki, A.,
Nakano, T., and Honjo, T. (2002). Notch-RBP-J signaling is involved in cell fate
determination of marginal zone B cells. Nat Immunol *3*, 443–450. 10.1038/ni793.
42. Lin, L., DeMartino, J., Wang, D., van Son, G.J.F., van der Linden, R., Begthel, H., Korving,
J., Andersson-Rolf, A., van den Brink, S., Lopez-Iglesias, C., et al. (2023). Unbiased
transcription factor CRISPR screen identifies ZNF800 as master repressor of enteroendocrine
differentiation. Science *382*, 451–458. 10.1126/science.adi2246.
43. Miyake, S., Sellers, W.R., Safran, M., Li, X., Zhao, W., Grossman, S.R., Gan, J., DeCaprio,
J.A., Adams, P.D., and Kaelin, W.G., Jr. (2000). Cells degrade a novel inhibitor of
differentiation with E1A-like properties upon exiting the cell cycle. Mol Cell Biol *20*, 8889–
8902. 10.1128/MCB.20.23.8889-8902.2000.
44. Witte, S., Bradley, A., Enright, A.J., and Muljo, S.A. (2015). High-density P300 enhancers
control cell state transitions. BMC Genomics *16*, 903. 10.1186/s12864-015-1905-6.
45. Andrews, T.G.R., Cornwall-Scoones, J., Ramel, M.-C., Gupta, K., Briscoe, J., and Priya, R.
(2025). Mechanochemical coupling of cell shape and organ function optimizes heart size and

contractile efficiency in zebrafish. Developmental Cell *60*, 3167–3184.e3168.
10.1016/j.devcel.2025.07.011.

46. 984 Priya, R., Allanki, S., Gentile, A., Mansingh, S., Uribe, V., Maischein, H.M., and Stainier,
985 D.Y.R. (2020). Tension heterogeneity directs form and fate to pattern the myocardial wall.
986 Nature *588*, 130–134. 10.1038/s41586-020-2946-9.

47. 987 Khait, I., Orsher, Y., Golan, O., Binshtok, U., Gordon-Bar, N., Amir-Zilberstein, L., and
988 Sprinzak, D. (2016). Quantitative Analysis of Delta-like 1 Membrane Dynamics Elucidates
989 the Role of Contact Geometry on Notch Signaling. Cell Rep *14*, 225–233.
990 10.1016/j.celrep.2015.12.040.

48. 991 Shaya, O., Binshtok, U., Hersch, M., Rivkin, D., Weinreb, S., Amir-Zilberstein, L., Khamaisi,
992 B., Oppenheim, O., Desai, R.A., Goodyear, R.J., et al. (2017). Cell-Cell Contact Area Affects
993 Notch Signaling and Notch-Dependent Patterning. Dev Cell *40*, 505–511 e506.
994 10.1016/j.devcel.2017.02.009.

49. 995 Sarate, R.M., Hochstetter, J., Valet, M., Hallou, A., Song, Y., Bansaccal, N., Ligare, M.,
996 Aragona, M., Engelman, D., Bauduin, A., et al. (2024). Dynamic regulation of tissue fluidity
997 controls skin repair during wound healing. Cell *187*, 5298–5315.e5219.
998 10.1016/j.cell.2024.07.031.

50. 999 Hogg, S.J., Motorna, O., Cluse, L.A., Johanson, T.M., Coughlan, H.D., Raviram, R., Myers,
1000 R.M., Costacurta, M., Todorovski, I., Pijpers, L., et al. (2021). Targeting histone acetylation
1001 dynamics and oncogenic transcription by catalytic P300/CBP inhibition. Mol Cell *81*, 2183–
1002 2200 e2113. 10.1016/j.molcel.2021.04.015.

51. 1003 Weinert, B.T., Narita, T., Satpathy, S., Srinivasan, B., Hansen, B.K., Scholz, C., Hamilton,
1004 W.B., Zucconi, B.E., Wang, W.W., Liu, W.R., et al. (2018). Time-Resolved Analysis Reveals
1005 Rapid Dynamics and Broad Scope of the CBP/p300 Acetylome. Cell *174*, 231–244 e212.
1006 10.1016/j.cell.2018.04.033.

52. 1007 Pylvanainen, J.W., Laine, R.F., Saraiva, B.M.S., Ghimire, S., Follain, G., Henriques, R., and
1008 Jacquemet, G. (2023). Fast4DReg - fast registration of 4D microscopy datasets. J Cell Sci
1009 *136*. 10.1242/jcs.260728.

53. 1010 Pachitariu, M., Rariden, M., and Stringer, C. (2025). Cellpose-SAM: superhuman
1011 generalization for cellular segmentation. bioRxiv, 2025.2004.2028.651001.
1012 10.1101/2025.04.28.651001.

54. 1013 Shen, L., Farid, H., and McPeek, M.A. (2009). Modeling three-dimensional morphological
1014 structures using spherical harmonics. Evolution *63*, 1003–1016. 10.1111/j.1558-
1015 5646.2008.00557.x.

55. 1016 Biggs, L.C., Kim, C.S., Miroshnikova, Y.A., and Wickstrom, S.A. (2020). Mechanical Forces
1017 in the Skin: Roles in Tissue Architecture, Stability, and Function. J Invest Dermatol *140*,
1018 284–290. 10.1016/j.jid.2019.06.137.

56. 1019 Dahmann, C., Oates, A.C., and Brand, M. (2011). Boundary formation and maintenance in
1020 tissue development. Nat Rev Genet *12*, 43–55. 10.1038/nrg2902.

57. 1021 Foty, R.A., and Steinberg, M.S. (2005). The differential adhesion hypothesis: a direct
1022 evaluation. Dev Biol *278*, 255–263. 10.1016/j.ydbio.2004.11.012.

58. 1023 Harris, T.J., and Tepass, U. (2010). Adherens junctions: from molecules to morphogenesis.
1024 Nat Rev Mol Cell Biol *11*, 502–514. 10.1038/nrm2927.

59. 1025 Joost, S., Zeisel, A., Jacob, T., Sun, X., La Manno, G., Lonnerberg, P., Linnarsson, S., and
1026 Kasper, M. (2016). Single-Cell Transcriptomics Reveals that Differentiation and Spatial
1027 Signatures Shape Epidermal and Hair Follicle Heterogeneity. Cell Syst *3*, 221–237 e229.
1028 10.1016/j.cels.2016.08.010.

60. 1029 Krieg, M., Arboleda-Estudillo, Y., Puech, P.H., Kafer, J., Graner, F., Muller, D.J., and
Heisenberg, C.P. (2008). Tensile forces govern germ-layer organization in zebrafish. Nat Cell
Biol *10*, 429–436. 10.1038/ncb1705.

61. Monier, B., Pelissier-Monier, A., Brand, A.H., and Sanson, B. (2010). An actomyosin-based
barrier inhibits cell mixing at compartmental boundaries in Drosophila embryos. Nat Cell
Biol *12*, 60–69. 10.1038/ncb2005.
62. Rubsam, M., Broussard, J.A., Wickstrom, S.A., Nekrasova, O., Green, K.J., and Niessen,
C.M. (2018). Adherens Junctions and Desmosomes Coordinate Mechanics and Signaling to
Orchestrate Tissue Morphogenesis and Function: An Evolutionary Perspective. Cold Spring
Harb Perspect Biol *10*. 10.1101/cshperspect.a029207.
63. Wickstrom, S.A., and Niessen, C.M. (2018). Cell adhesion and mechanics as drivers of tissue
organization and differentiation: local cues for large scale organization. Curr Opin Cell Biol
54, 89–97. 10.1016/j.ceb.2018.05.003.
64. Watt, F.M. (2002). Role of integrins in regulating epidermal adhesion, growth and
differentiation. The EMBO Journal *21*, 3919–3926. 10.1093/emboj/cdf399.
65. Sugimoto, M., Tada, Y., Shichino, S., Koyamatsu, S., Tsumaki, N., and Abe, K. (2022).
Universal Surface Biotinylation: a simple, versatile and cost-effective sample multiplexing
method for single-cell RNA-seq analysis. DNA Res *29*. 10.1093/dnares/dsac017.
66. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics
29, 15–21. 10.1093/bioinformatics/bts635.
67. Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell gene
expression data analysis. Genome Biol *19*, 15. 10.1186/s13059-017-1382-0.
68. Lun, A.T., Bach, K., and Marioni, J.C. (2016). Pooling across cells to normalize single-cell
RNA sequencing data with many zero counts. Genome Biol *17*, 75. 10.1186/s13059-016-
0947-7.
69. Hunter, J.D. (2007). Matplotlib: A 2D Graphics Environment. Computing in Science &
Engineering *9*, 90–95. 10.1109/MCSE.2007.55.
70. Tirosh, I., Izar, B., Prakadan, S.M., Wadsworth, M.H., 2nd, Treacy, D., Trombetta, J.J.,
Rotem, A., Rodman, C., Lian, C., Murphy, G., et al. (2016). Dissecting the multicellular
ecosystem of metastatic melanoma by single-cell RNA-seq. Science *352*, 189–196.
10.1126/science.aad0501.
71. McInnes, L.H., J.; Melville, J. (2020). UMAP: Uniform Manifold Approximation and
Projection for Dimension Reduction. https://arxiv.org/abs/1802.03426}, .
72. Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden: guaranteeing
well-connected communities. Sci Rep *9*, 5233. 10.1038/s41598-019-41695-z.
73. Wolock, S.L., Lopez, R., and Klein, A.M. (2019). Scrublet: Computational Identification of
Cell Doublets in Single-Cell Transcriptomic Data. Cell Systems *8*, 281–291.e289.
https://doi.org/10.1016/j.cels.2018.11.005.
74. Korsunsky, I., Millard, N., Fan, J., Slowikowski, K., Zhang, F., Wei, K., Baglaenko, Y.,
Brenner, M., Loh, P.R., and Raychaudhuri, S. (2019). Fast, sensitive and accurate integration
of single-cell data with Harmony. Nat Methods *16*, 1289–1296. 10.1038/s41592-019-0619-0.
75. Squair, J.W., Gautier, M., Kathe, C., Anderson, M.A., James, N.D., Hutson, T.H., Hudelle,
R., Qaiser, T., Matson, K.J.E., Barraud, Q., et al. (2021). Confronting false discoveries in
single-cell differential expression. Nat Commun *12*, 5692. 10.1038/s41467-021-25960-2.
76. Fang, Z., Liu, X., and Peltz, G. (2023). GSEApy: a comprehensive package for performing
gene set enrichment analysis in Python. Bioinformatics *39*. 10.1093/bioinformatics/btac757.
1076
1077

Main Figures

1079

Figure 1. Basal cells employ distinct embryonic stage-specific strategies to generate suprabasal

layers

A. Schematics of the *ex vivo* embryo imaging setup. **B.** Quantification of parallel (left panel), basal

(middle panel) and suprabasal (right panel) cell divisions in E14.5 and E15.5 embryos (mean ±SD;

n=4 (E14.5) /5 (E15.5) embryos; Student's t-test (left and middle panel), Mann-Whitney (right

panel)). **C.** Representative images and quantification of CyclinB-positive undifferentiated cells (K10-neg) in the suprabasal layers of E14.5 and E15.5 embryos (30 μm; mean ±SD; n=4 embryos/stage; Mann-Whitney). **D.** Representative snapshots and 3D rendering (lower panel) from live imaging movies of E14.5 epidermis from nTomato reporter mice. Red asterisk marks cell departing from basal layer; double red asterisk marks daughter of perpendicular division that subsequently moves suprabasally, blue asterisks marks appearance of cells in the suprabasal layer, red dotted circle marks completed delamination. Images representative of 3 embryos, scale bars 20 μm. **E.** Representative snapshots and 3D rendering (lower panel) from live imaging movies of E15.5 epidermis from nTomato reporter mice. Red asterisk marks cell departing from basal layer; blue asterisk marks appearance of this cell in the suprabasal layer, red dotted circle marks completed delamination. Images representative of 3 embryos, scale bars 20 μm. **F.** Quantification of nuclear oblateness (roundness; upper panel) and asymmetry of deformation (wedge-shape; lower panel) of delaminating cells (Del.) and their immediate neighbors (Neigh.) from 3D renderings. Note that while at E14.5 delaminating cells round up, at E15.5 they elongate in an asymmetric manner (min-to-max box plots; n=3 embryos/stage). **G.** Representative images and quantification of tissue sections from E14.5 - 16.5 embryos stained against K10, Dsg1 and DAPI. Note emergence of K10-Dsg1- positive cells at 15.5 (red asterisks). Dotted line marks basement membrane (scale bars 20 μm; mean ±SD; n=4 embryos; Two-way ANOVA/Tukey's). **H.** Representative images and quantification of tissue sections from E15.5 embryos stained against K10, Dsg1 and DAPI. Note correlation of wedge-like shape and apical Dsg1 intensity. White asterisks mark wedge-like cells with high apical Dsg1, blue asterisks mark cuboidal neighboring cell (scale bars 10 μm; mean ±SD; n=87 cells pooled across 3 embryos; Simple linear regression; Pearson's correlation). **I, J.** Representative images and quantification of delaminating cells in E15.5 embryo 3D whole mounts stained against integrin α6 (Itga6), K10 and DAPI. Note wedge-like shape and integrin α6-positive basal footprint of the delaminating cell (white asterisks; scale bars 5 μm; min-max box and whiskers; n=37 cells pooled across 3 embryos; Mann-Whitney (left, middle panel) Friedman/Dunn's).

1112

1113

Figure 2. A biophysical model predicts that in the presence of a mechanical barrier, delamination depends on the mechanical properties of the delaminating cell.

A. Snapshot from 3D vertex model for stratified epithelium geometry, illustrating suprabasal cells (brown), basal cells (light blue) and basement membrane (dark blue), interfacial tensions between the apical surface of basal cells and the adjacent suprabasal cell (, magenta), basal cells and basement membrane (σ_b, green), and cell shapes (inset) with different 3D shape parameters. **B.** Simulation snapshot including a delaminating cell type, with potentially different delaminating cell-basal cell lateral interfaces σ_{dl}, delaminating cell-basement membrane interfaces (σ_b, yellow), and delaminating cell-suprabasal cell interfaces (σ_{da}, red). **C.** Snapshot of simulation showing cell in process of delaminating (cell integrates fully into the suprabasal layer by end of simulation). **D.** Fraction of cells that delaminate after a time t = X, as a function of σ_{db}, for many different values of the other parameters. Data averaged over an ensemble of n=20 simulations for each simulation condition. **E.** Delamination fraction collapses as a specific function of vertex model parameters.

Figure 3. A mechanical boundary emerges at E15.5 impacting delamination.

A. Representative super resolution images of tissue cross sections across developmental stages. Scale bars 10 μm. **B.** 3D rendering of cell shapes across developmental stages from data in (A). Scale bars 5μm. **C.** Quantification of 3D cell shapes from (B) using spherical harmonics reveals a decrease in deformed shapes and an increase in prolateness from E14.5 to E16.5 (n=19 (E14.5, E15.5)/ 30 (E16.5) cells pooled across 3 embryos; Kruskal-Wallis/Dunn's). **D.** Schematic representation of cell-scale geometric parameters and their integration into the vertex model. **E.** Quantification of lateral interface height, variability of orientation angle and apical surface angle across developmental stages

(n>400 cells (height); >25 fields of view (apical angle) >740 cells (lateral angle) per timepoint pooled across 3 mice (E14.5, E15.4), 5 mice (E15.5); Kruskal-Wallis/Dunn's). **F.** Schematic representation of tissue-scale geometric parameter of surface roughness in the 3D vertex model. **G, H.** Representative experimental whole mount (B) and cross section images (C) of surface roughness measurements (scale bars 10 μm). **I.** Collapsed simulation data measured in the 3D model simulations across mechanical parameter space ($\sigma_a, \sigma_b, \Delta s$). Corresponding averages of *in vivo* observables converted to simulation units specified as green, orange and blue symbols (n>85 simulations for each parameter set). **J.** Apical angle simulation data as a function of vertex model parameters required by force balance (n>85 simulations for each parameter set). **K.** Best-fit vertex model parameters σ_b (green), σ_a (pink), Δs (blue) to the cell and tissue geometry observed at each stage; error bars represent uncertainty in the fit (see Supplemental Methods). **L.** Simulation-predicted delamination rates when delaminating cells alter their mechanical properties from the background basal cells by $\Delta\sigma_{da,a}$ (apical) and $\Delta\sigma_{db,b}$ (basal). Rate is the fraction that delaminates in a fixed time window t = 2000 timesteps, corresponding to $\tau \sim 20$ simulation time units (n>85 simulations per parameter set).

Figure 4. A committed basal population characterized by Notch activation emerges at E15.5

A. Schematics of the scRNAseq experiment. **B.** UMAP representation of the Leiden clusters of
filtered epidermal cells from scRNAseq. Embryonic stages of cells are color coded. **C.** UMAP
projections of basal identity and differentiation gene expression across E14.5-E16.5. Arrowheads
mark transition population between basal and suprabasal cells. **D.** Dotplot of selected genes across

E14.5-E16.5 basal cells. Note increased expression of differentiation genes in basal cells of E15.5

and E16.5 embryos. **E.** Dot plot of receptor-ligand (r+l) analyses between basal stem cells across

E14.5-E16.5. Note increased predicted Jag2-Notch signaling interaction at E15.5 and E16.5. **F-H.**

Representative images (F) and quantification (G, H) of tissue sections from E14.5- 16.5 embryos

stained against Notch intracellular domain (N-ICD) and K10. Note emergence of N-ICD-K10-

positive delaminating basal cells at 15.5. Dotted line marks basement membrane (mean ±SD; scale

bars 20 μm; n=3 embryos/stage; Two-way ANOVA/Tukey's (G); Mann-Whitney (H)).

1164

1165   **Figure 5. Generation of the mechanical boundary at E15.5 is concomitant with stiffening of the**

1166   **basement membrane and rigidification of the basal layer**

A. Dot plot of Gene set enrichment analyses (GSEA) from differentially expressed genes in basal cells of E14.5 versus E15.5/E16.5 embryos from scRNAseq data. **B, C.** Representative images (B) and quantification (C) of tissue sections from E14.5- 16.5 embryos analyzed by mass cytometry. Note increased expression of basement membrane Collagen IV (counterstained with K10) and integrins α6 and β1 (mean ±SD; scale bars 50 μm; n=3 embryos/stage; Two-way ANOVA/Tukey's (Left and middle panel), Kruskal-Wallis/Dunn's (right panel)). **D.** Basement membrane Young's moduli extracted from AFM force indentation experiments (Tukey's box and whiskers; n>200 force curves pooled across 3embryos/stage; Kolmogorov-Smirnov). **E.** Representative images and quantification of skin wholemounts from E14.5 and E15.5 embryos stained for pMLC2 and phalloidin. Note increased pMLC2 intensity at cell junctions and interface between basal and suprabasal layers (arrows) (min-max box and whiskers; scale bars 10 μm; n=5 embryos/stage; Mann-Whitney). **F.** Representative snapshots from live imaging movies and subsequent PIV analyses of velocity magnitudes of E14.5 and E15.5 epidermis from membrane-Tomato (mTomato) reporter mice. Red masks mark excluded hair follicle areas (scale bars 50 µm). **G.** Quantification of 2D cell perimeter, area and circularity from data in (A) (Tukey's box and whiskers; n>2300 cells pooled across 3 embryos/stage; Mann-Whitney). **H.** Quantification of cell densities from data in (A) (mean ±SD; n=4 embryos/stage; Mann-Whitney). **I.** Quantification of velocity magnitudes from data in (A). Note attenuated velocity at E15.5 (mean ±SD; n=4 embryos/stage; Linear Mixed-Effects model.

1185

Figure 6. Tissue rigidity and cell shape changes drive Notch- delamination and control of tissue crowding

A. Schematic illustration of micropatterning of soft and stiff polyacrylamide (PAA) hydrogels. **B.** Representative images of primary epidermal progenitors stained with K10 (green) and phalloidin (magenta) cultured on 100-µm-diameter micropatterns on 2.5 kPa (soft) or 8 kPa (stiff) PAA gels. K10 mean intensity was quantified as a function of the distance from the bottom of the micropatterns. Note the increased K10 mean intensity in the suprabasal region on stiff substrates. **C.** Quantification of boundary straightness between the basal layer (bottom) and the first suprabasal layer (upper layer). (Tukey's box and whiskers, n= 10 (soft); 15 (stiff) patterns pooled across 3 independent experiments; Mann-Whitney). **D.** Representative snapshots and quantification of live imaging movies from epidermal progenitor cell monolayers from mTomato/mGFP-Cre reporter mice crossed with Notch1-ICD-Cre to detect cells that activate Notch (mGFP+). Note increased elongation and local cell density of cells (white asterisk) that subsequently activate Notch signaling (red asterisk). Dotted circle marks delamination of Notch-Reporter-positive (NotchRep+) cell (min-max box and whiskers; n=38 Notch+ cells pooled across 3 independent experiments; Mann-Whitney; scale bars 10 µm). **E.** Representative images of tissue sections from E14.5-16.5 embryos of control and Rbpj-KO mice stained against K10 (magenta) and Desmoglein 1 (Dsg1; white) (images representative of >3 embryos/stage; scale bars 20 µm). **F.** Quantification of epidermal thickness from data in (A). Note reduced total and differentiated layer thickness of Rbpj-KO epidermis at E16.5 (mean ±SD; n=3 embryos/E14.5 and E15.5, 4/E16.5; Mann-Whitney). **G.** Quantification of basal layer thickness and cell height from data in (E). Note increase basal layer thickness and cell height of Rbpj-KO epidermis emerging at E15.5 (mean ±SD; n=3 embryos/E14.5 and E15.5, 4/E16.5; Mann-Whitney). **H.** Quantification of delaminating cells from data in (E). Note increase in delaminating cells in Rbpj-KO epidermis at E15.5 (mean ±SD; n=3 (E14.5), 5 (E15.5), 4 (E16.5) embryos; Mann-Whitney). **I.** UMAP representation of the Leiden clusters (left panel) and biological samples (right panel) of filtered epidermal cells from scRNAseq. **J.** Dotplot of selected genes across E14.5-E16.5 epidermal cells. Clusters 3 and 6 (in red) represent committed basal cell populations enriched in Rbpj-KO cells of E15.5 and E16.5 embryos. **K.** Violin plots of Zfp800 of Eid1 expression across E14.5-E16.5 in control and Rbpj-KO epidermal cells. **L.** Representative images and H3K27ac nuclear intensity quantification from tissue sections from E15.5 embryos of control and Rbpj-KO mice stained for H3K27ac (grey), K10 (magenta) and DAPI (cyan) (n= 176 (CNL), 234 (Rbpj-KO) cells pooled across 3 (CNL) /4 (Rbpj-KO) embryos; Student's t-test; Scale bars 20 µm).

Supplementary Figures

1220

Supplementary Figure 1.

A. Quantification of differentiated (K10-positive) CyclinB1+ cells in the suprabasal layer (mean

±SD; n=4 embryos /stage; Mann-Whitney) **B.** Representative images and quantification of

delaminating cells in E15.5 embryo from 3D whole mounts stained against E-cadherin, K10 and

DAPI. Delaminating cell is marked with a white asterisk; scale bars 10 µm; min-max box and

whiskers; n=38 cells pooled across 3 embryos; Friedman/Dunn's).

1227

Supplementary Figure 2. Systematic quantification of delamination rates as a function of the

properties of the delaminating and stiffness of the surrounding tissue Δs

A. Uncollapsed delamination rates. **B.** Delamination rates plotted as a function of $\sigma_{db} - \sigma_{da}$, which

do not fully collapse, indicating that an additional energetic contribution is required. Figure 2D in the

main text shows that adding the term $c \, \Delta s$ leads to a collapse of the curves onto a single master curve

as a function of $\sigma_{db} - \sigma_{da} + c \, \Delta s$.

1234

Supplementary Figure 3. Geometric observables for mechanical barrier and quantification of delamination rates and associated energy barriers as function of delaminating cell properties.
A-C. Uncollapsed plots of basal-suprabasal interface roughness, basal cell lateral interface height, and lateral interface orientation, respectively, measured in the 3D model simulations across mechanical parameter space $(\sigma_a, \sigma_b, \Delta s)$. Corresponding collapsed plots are shown in main Fig. 3G, I, K. **D.** Quantification of delamination rates data as function of the delaminating cell tensions $(\sigma_{db}, \sigma_{da})$ parameter space. **E.** Data from (F) collapses as a function of $\sigma_{db} - \sigma_{da}$. **F.** Curve fits to the collapsed delamination rate data shown in panel (B) across the developmental stages. **G.** Quantification of the differences in energy barriers between developmental stages as a function of the differences in delaminating cell properties $(\sigma_{da}, \sigma_{db})$. **H.** Normalized averages (normalized by the energy barrier difference between stages E16 and E15) of the differences in energy barriers between stages E15.5 and E14.5, and between E16.5 and E14.5, plotted as a function of $\sigma_{db} - \sigma_{da}$.

1247

Supplementary Figure 4.

A. Bar plot showing number of sequenced cells/genotype. **B.** Dot plot of selected key cell type-specific marker genes across annotated cell types. **C.** UMAP visualization of Leiden clusters of

1251 epidermal cells with main cell states annotated. **D.** Heatmap of marker genes for epidermal Leiden

1252 clusters. **E.** UMAP visualization of Harmony integrated epidermal cells. Red arrow indicates

1253 committed cell population. **F.** Dot plot of selected key cell type-specific marker genes across

1254 annotated cell types from Harmony integrated epidermal cells. Red dotted rectangle indicates

1255 committed cell population.

1256

1257 **Supplementary Figure 5.**

1258 **A.** Representative images and quantifications of tissue sections from E14.5- 16.5 embryos analyzed

1259 by mass cytometry. Note increased expression of integrin β4 (mean ±SD; scale bars 50 µm; n=3

1260 embryos/stage; Two-way ANOVA/Tukey's.

Supplementary Figure 6.

A. Volcano plot showing differentially expressed genes between Rbpj-KO and CNL epidermis from differentiated cell clusters 5 and 8. **B.** Bar plot showing number of sequenced cells/embryonic stage and genotype across Leiden clusters. **C.** Volcano plot showing differentially expressed genes between E15.5 Rbpj-KO and CNL epidermis from clusters 6 and 0. **D.** Volcano plot showing differentially expressed genes between E16.5 Rbpj-KO and CNL epidermis from cluster 3. **E.** Representative images and quantification of tissue sections from E15.5 embryos of control and Rbpj-KO mice of the expression of Eid1 (magenta) and Keratin-dap (grey) detected by RNA in situ hybridization. Dotted lines mark basal layer (n=4 embryos/condition; Mann-Whitney). Scale bars 20 μm.