**Tissue-scale mechanics controls differentiation strategy and dynamics of epithelial multilayering**

Clémentine Villeneuve[1, 2]*, Somiealo Azote Epse Hassikpezi [3]*, Marga Albu[2], Matthias Rübsam[4], Leah C. Biggs[1,2], Sabrina Vinzens[2], Kai Kruse[5], Anubhav Prakash[2], Peter Zentis[6], Elizabeth Lawson-Keister[3], Gautier Follain[7,8,9], Johanna Ivaska[7,10,11], Carien M. Niessen[4§], M. Lisa Manning[3§], and Sara A. Wickström[1,2,12§]

[1]Stem Cells and Metabolism Research Program, Faculty of Medicine, University of Helsinki, 00290 Helsinki Finland

[2]Department of Cell and Tissue Dynamics, Max Planck Institute for Molecular Biomedicine, 48149 Münster, Germany

[3]Department of Physics and BioInspired Institute, Syracuse University, Syracuse, NY, USA.

[4]Department Cell Biology of the Skin, Cologne Excellence Cluster on Cellular Stress Responses in Aging Associated Diseases (CECAD), Center for Molecular Medicine Cologne, University Hospital Cologne, University of Cologne, Joseph-Stelzmann-Str. 26, 50931 Cologne, Germany.

[5]Max Planck Institute for Molecular Biomedicine, 48149 Münster, Germany

[6]Cluster of Excellence for Aging-Associated Diseases, CECAD Imaging Facility Cologne, University of Cologne, Cologne, Germany.

[7]Turku Bioscience Centre, University of Turku and Åbo Akademi University, Turku, 20520, Finland

[8]Faculty of Science and Engineering, Cell Biology, Åbo Akademi University, Turku, 20520, Finland

[9]Turku Collegium for Science, Medicine and Technology (TCSMT), Turku, 20520, Finland

[10]Department of Life Technologies, University of Turku, FI-20520 Turku, Finland.

[11]InFLAMES Research Flagship Center, University of Turku, Turku, Finland.

[12]Helsinki Institute of Life Science, Biomedicum Helsinki, University of Helsinki, FI-00290 Helsinki, Finland

*These authors contributed equally

§These authors jointly supervised this work

carien.niessen@uni-koeln.de

mmanning@syr.edu

sara.wickstrom@mpi-muenster.mpg.de

**Abstract**

Generating and maintaining multilayered epithelia requires coordinated cell division,

differentiation, and tissue architecture, yet how multilayering arises remains unclear. Using

the developing mouse epidermis, we show that basal stem cells adopt distinct multilayering

strategies depending on tissue mechanics. Early in development, the epidermis is fluid-like,

allowing undifferentiated cells to move suprabasally through perpendicular divisions or basal

detachment before differentiating. As the tissue matures and rigidifies, a mechanical barrier is

established that only allows upward movement of basal cells that have committed to

differentiation. The final step of this commitment requires Notch signaling that is triggered

by increased tissue stiffness and jamming, orchestrating a feedback loop that induces cell

upward motion precisely when the bottom layer becomes crowded. Together, our findings

identify tissue mechanics as the key determinant of how tissues drive multilayering and

reveal mechanically regulated Notch signaling as a driver of epithelial delamination.

**Introduction**

Stratified epithelia such as the skin epidermis are multilayered tissues where the basal layer

consists of undifferentiated stem cells adhering to the basement membrane, while the

suprabasal layers contain post-mitotic, differentiated cells [1-3]. Unlike other dynamically

self-renewing epithelia such as the intestine, where clear tissue-scale geometric cues such as

curvature compartmentalize the stem cell niche and differentiated progeny [4], it remains

unclear how stratified epithelia establish and organize this spatial organization and

differentiation dynamics during development and into adulthood.

To initiate or maintain epidermal stratification, stem cells that are restricted to the basal layer

generate suprabasal cells via two distinct strategies: perpendicular divisions or detachment

from the basal layer, termed delamination [2, 3, 5]. Early in development, proliferation rates

in the epidermis are high - to match the rapid growth of the organism it covers, and to

generate sufficient suprabasal cells for efficient establishment of a functional life-essential

barrier [6]. During this period, perpendicular cell divisions generate suprabasal daughter cells

[7]. In contrast, during adult homeostasis differentiation is independent of cell divisions, with

cells moving suprabasally to replenish the most superficial, desquamating cornified layers,

followed by parallel divisions of to maintain basal cell density [8, 9]. Key open questions

remain: When and how does this switch in strategies occur? And how is upward movement

robustly regulated, so that stem cells are retained in the basal layer while differentiating move

at the correct rates, maintaining basal layer density and epithelial thickness in the absence of

obvious signals like morphogens or tissue curvature?

Here we demonstrate that multilayering strategy is controlled by a mechanical barrier that is

established during development and hinders basal cells from moving upward. We establish a

new approach to quantitively measure this mechanical barrier by combining a 3D biophysical

model for cell and tissue mechanics with imaging of cell and tissue geometries in the basal

layer and at the basal-suprabasal interface. We find that at early developmental stages,

embryonic (E) day 13.5/14.5, there is little mechanical distinction between basal and

suprabasal layers, allowing basal cells to easily place cells suprabasally either through

perpendicular division or through rapid detachment assisted by fluctuations from cell

divisions. Starting around E15.5, a mechanical energy barrier between the basal and

suprabasal layers emerges, with cell divisions becoming mostly parallel and daughters not

crossing the barrier. Instead, basal cells must now undergo profound transcriptional

reprogramming– e.g. commit to delamination – in order to cross the barrier. How is this

decision to commit regulated? Our data suggests an elegant feedback mechanism involving

Notch signaling and tissue crowding. We show that Notch activation is sensitive to cell

shapes and tissue rigidity. Tissue crowding-induced changes trigger a transient Notch signal

in a small subset of cells, causing these cells to execute a transcriptional switch to complete

delamination. Collectively, our study shows that changes in tissue mechanics that drive

intercompartment boundary formation are essential determinants for a switch from early

embryonic to a mature differentiation and multilayering strategy. Moreover, we identify

Notch signaling as an integrative feedback relay to coordinate tissue density and cell fate

dynamics to correctly time the transition between the mechanically separated basal and

suprabasal compartments within a mature stratified epithelium.

**Results**

**Basal cells employ distinct embryonic stage-specific strategies to generate and**
**compartmentalize suprabasal layers**

To understand which mechanisms are used for epidermal multilayering, we employed a

whole embryo live imaging system [10, 11] to visualize long term (over 8 h) cell dynamics at

E14.5 and E15.5 (Fig. 1A). While perpendicular divisions were prevalent at E14.5, parallel

oriented cell divisions became predominant at E15.5 with the overall frequency of cell

divisions decreasing (Fig. 1B). As reported [12], cell divisions also occurred suprabasally at

E14.5 but much less at E15.5 (Fig. 1B, Supplementary Fig. 1A). Imaging of fixed tissues

showed that significantly more undifferentiated cells were found in the first suprabasal layer at E14.5 than at E15.5 (Supplementary Fig. 1B), as detected by Keratin (K)-14 expression that marks undifferentiated stem cells and K10 that marks differentiated suprabasal cells [13]. A fraction of these suprabasal undifferentiated cells were still proliferating at E14.5 but not at E15.5 (Fig. 1C). This suggested that differentiation is not associated with upward movement at early stages.

We then asked whether cell delamination itself also changed as a function of developmental stage. At E14.5, detachment of cells from the basement membrane was regularly observed, with upward movement lasting around 120 minutes. This detachment was frequently associated with a cell division, either occurring after a perpendicular division or followed by a mitosis in the suprabasal layer (Fig. 1D; Supplementary Movie 1). In contrast, at E15.5 delaminations were substantially slower, lasting at least 300 minutes, and were not associated with divisions (Fig. 1E; Supplementary Movie 2). The geometry of delaminating cells also changed; at E14.4 cells rounded up for delamination, while at E15.5 cells transitioned instead into elongated and basally constricted, wedge-like shapes prior to delamination (Fig. 1F), indicating a switch in mechanical state.

To better understand how these changes in division and delamination strategies relate to cell differentiation, and to identify the mechanisms involved, we turned to fixed tissue imaging. Previous work has implicated desmosomal cadherins/Desmoglein1 (Dsg1) in delamination [14]. At E14.5 no Dsg1 or K10-positive differentiating cells were detected in the basal layer (Fig. 1G). In contrast, at E15.5 Dsg1+ K10+ basal cells were now observed, indicating that these cells had already initiated commitment towards differentiation. These double-positive basal cells exhibited upregulation of Dsg1 with enhanced localization of Dsg1 (Fig. 1H) and E-cadherin (Supplementary Fig. 1C) at apical interfaces, displaying wedge-like shapes and nuclear deformation similar to delaminating cells seen in live imaging (Fig. 1H).

To precisely quantify the shape changes of these K10+-basal cells at E15.5, we performed super-resolution imaging of skin whole mounts. K10+ basal cells were still attached to the basement membrane as shown by staining for α6β4 integrins, (Fig. 1I), and displayed a wedge-like shape characterized by decreased basal adhesion length and increased apical adhesion surface (Fig. 1I). Interestingly, integrins were enriched at these narrow basal footprints, indicating that differentiation is initiated prior to complete cell detachment from the basement membrane (Fig. 1J).

Collectively, these experiments show that at early developmental stages embryos engage both perpendicular cell divisions and rapid delaminations associated with suprabasal divisions to

generate suprabasal layers. Instead, E15.5 and E16.5 embryos employ a substantially slower,

division-independent delamination process associated with remodeling of cell adhesion and

changes in cell shape to position cells suprabasally.

**A biophysical model predicts that in the presence of a mechanical boundary,**

**delamination depends on the mechanical properties of the delaminating cell**

To understand the mechanistic basis for this switch in multilayering strategy, we turned to

quantitative mechanical modelling. For this we developed 3D vertex model of the epidermis

[11, 15] (see Supplementary Methods). The model consisted of suprabasal cells (brown),

basal cells (light blue) and a basement membrane (dark blue; Fig. 2A). A key model feature

was basal layer rigidity, governed by cell shape. We therefore parameterized this by $\Delta s$, the

distance in shape space to the fluid-solid transition [15-17] (Fig. 2A, inset). The model also

included heterotypic interfacial tensions between basal and adjacent suprabasal cells ($\sigma_a$; Fig.

2A, magenta), and a potentially wetting interfacial tension between basal cells and the

basement membrane ($\sigma_b$ ; Fig. 2A green). A similar Voronoi-based model was recently used

to study clonal dynamics of tumor cells in stratified epithelia {Sahu, 2025 #80}.

Previous work [11, 15] had shown that these three "tissue background" parameters ($\sigma_a$, $\sigma_b$,

and $\Delta s$) generate a robust mechanical model that is capable of capturing key features of the

multilayered tissue architecture observed *in vivo* by generating a mechanical boundary

between the basal and suprabasal layers. This boundary ensures that small active mechanical

fluctuations do not induce mixing between basal and suprabasal cells, and the layers are

robustly separated, as observed at E15.5 onwards.

We sought to predict whether the presence of a mechanical boundary could explain the need

for a mechanical change of the delaminating cell, observed at E15.5/E16.5 (see Fig, 1). We

thus studied the behavior of a potential delaminating basal cell by altering specific interfacial

tensions of this cell – lateral $\sigma_{dl}$, basal $\sigma_{db}$, and apical $\sigma_{da}$ (Fig. 2B). Past experimental

observations [10, 18], suggested that basal layer rigidity $\Delta s$ is also relevant for delamination.

We therefore also allowed it to vary while keeping the other "tissue background" parameters

$\sigma_b$ and $\sigma_a$ fixed. We found that indeed, for a cell to move upward into the suprabasal layer in

the presence of small active fluctuations, it needed only to change its interfacial tensions (Fig.

2C). We proceeded to precisely predict how changes to the mechanics of the delaminating

cell alter the energy barrier and therefore the rates of delamination. To quantify these rates,

we simulated a large ensemble of conditions (N>85) for each set of model parameters (3240

sets), and measured the fraction of cells that had delaminated in a fixed time window ($\Delta t$ =

2000 simulation time units, estimated to be 20 – 200 min [19, 20], see Supplemental

Methods). We found that delamination rates are a complex function of the four variables

($\sigma_{da}$, $\sigma_{dl}$, $\sigma_{db}$, $\Delta s$; Supplementary Fig. 2A). In order to parse the resulting large parameter

space, we adopted physics methodology [21] to search for a data collapse in delamination

rates. We further tested if those rates were consistent with an Arrhenius process governed by

an energy barrier, i.e. where the rate is proportional to $e^{-E/kT_{eff}}$, where $k\,T_{eff}$ sets the

magnitude of active fluctuations [22-24]. We identified a specific combination of model

parameters that generated a data collapse – i.e. where all delamination rate curves fell on top

of one another (Fig. 2D; uncollapsed data in Supplementary Fig. 2). This indicated that a

delaminating cell experiences a mechanical energy barrier to move upwards, and that this

barrier can be modulated by the mechanical properties of the delaminating cell, in the

following form:

180
$$E \;=\; F(\,\sigma_{da} - \sigma_{db} + c\,\Delta s)$$

Here $F$ is a monotonically increasing function of its argument (see Supplemental Methods).

The first two terms indicate that the energy barrier depends on the difference between the

basal and apical interface tensions of the delaminating cell. To decrease the energy barrier

and delaminate more efficiently, a delaminating cell needs to increase the tension (i.e.

decrease the wetting) with the basement membrane, while also decreasing its apical

heterotypic interfacial tension with the suprabasal cells. This is consistent with the observed

emergence of a wedge-like cell shape and apical Dsg1 enrichment observed in the

experiments from E15.5 onwards (see Fig. 1G-I). The third term indicated that the energy

barrier increases with increasing surrounding basal layer stiffness $\Delta s$. Interestingly, the

energy barrier did not depend on $\sigma_{dl}$ - changes to the delaminating cell's lateral interfacial

tension did not change the delamination rate (Fig. 2D). Overall, these results indicated that

the non-mixing of basal and suprabasal cells observed at E15.5 and E16.5 is consistent with

the existence of a mechanical energy barrier, and that cells can cross this barrier by changing

their own mechanical properties.

**A substantial mechanical energy barrier emerges at E15.5 impacting delamination.**

Next, we sought to determine precisely if such a barrier emerges during development, and to

quantify its impact on delamination. Previous work suggests that the apical curvature angle $\gamma$

and the cell height $h$ reflect a force balance along each interface [25, 26], and that interfacial

properties – lateral orientation $\beta$ and apical interface roughness $R$ - depend on heterotypic

interfacial tensions [27, 28]. We therefore measured these observables and their variation *in*

*vivo* in the basal layer (Fig. 3A-E) and integrated them with the 3D vertex model by

converting the measurements to simulation units. We then simulated the model across a wide

range of parameter conditions (Fig. 3F-M). For three observables ($R, \beta, h$), we discovered

that the data could be collapsed as a simple function of the model parameters (Fig. 3F-K;

Supplementary Fig. 3.A-C). As a result, the average value of each *in vivo* observable (Fig.

3F, H, J) specified a point on the y-axis of the respective simulation plot (Fig. 3G, I, K), and

therefore generated a unique constraint on the model parameters. For $\gamma$ (Fig. 3L), force

balance allowed us to quantify this directly as function of $\sigma_b$ and $\Delta s$ (Fig. 3M). This

benchmarking process resulted in a set of four constraint equations for the three unknowns

($\sigma_b$, $\sigma_a$, $\Delta s$) at each developmental stage; we solved this system of equations for the best-fit

model parameters at each stage using a standard over-constrained solver [29] (Fig. 3N; see

Supplementary Methods).

These data demonstrated that at E14.5, the basal layer is nearly fluidlike, with very low tissue

stiffness parameterized by $\Delta s$ (Fig. 3N). At this stage the interfacial tension between basal

and suprabasal cells is low, coupled to a small, positive interfacial tension at the basement

membrane interface, suggesting only low levels of integrin-based wetting. At E15.5, the

tension at the basement membrane interface became negative, suggesting wetting behavior,

accompanied by increased interfacial tension at the basal-suprabasal interface, as well as

increased basal layer stiffness. At E16.5, similar trends remained, while tissue stiffness

increased further (Fig. 3N). Combined, these changes generated a four-fold increase in the

magnitude of the mechanical energy barrier for a basal cell to delaminate from E14.5 to

E15.5 (Supplementary Fig. 3F-H).

This substantial increase in the mechanical barrier between E14.5 and E15.5 has important

implications for strategies for delamination. Building upon our understanding from the data

collapse in Fig. 2, we were then able to collapse the rates of delamination as a function of the

differences in the interfacial tensions between the delaminating cell and the background basal

cells at the apical ($\Delta\sigma_{da,a}$) and basal ($\Delta\sigma_{db,b}$) interfaces (Fig. 3O; Supplementary Methods

Fig. 3C, see Supplemental Methods). This analysis showed that at E14.5, the probability of a

standard basal cell (0.0 on the x-axis of Fig 3O) delaminating during the time window $\Delta t$ was

less than 20%, but only a small change in basal and apical tensions (~15%, 0.15 on the x-axis

in Fig 3O) was sufficient to generate a massive increase in the delamination probability,

increasing it to nearly 100% in the same time window. In contrast, at E15.5 and E16.5, a

basal cell must change its mechanical properties substantially (by more than 50%, 0.5 on the

x-axis of Fig. 3O) in order to generate a similar increase in delamination probability.

Collectively, these data indicate that a mechanical boundary forms between the basal and

suprabasal layers at around E15.5, thus requiring cells to have a robust mechanism to commit

to delamination and overcome this barrier, while such a mechanism is not necessary at E14.5.

**240 A committed basal population characterized by Notch activation emerges at E15.5**

To identify the molecular driver for this predicted mechanical state change required for

delamination at E15.5 and later, we investigated the evolution of epidermal cell states using

single cell RNA sequencing (scRNAseq; Fig. 4A). We obtained single-cell suspensions of

skin from E13.5, E14.5, E15.5. and E16.5 embryos, isolated and sequenced in parallel to

avoid batch effects. The quality-controlled and filtered dataset, obtained from 2 biological

replicates/stage, comprised of >15,000 cells/stage that were annotated based on known

marker genes (Fig. 4A; Supplementary Fig. 4A, B). Focusing on the epidermal (non-hair

follicle) clusters, we identified cycling and non-cycling basal cells as well as suprabasal cell

states (Supplementary Fig. 4C, D). Leiden clustering revealed that cell states were clustered

separately according to their developmental stage, indicating that key cell states (basal and

suprabasal cells) underwent substantial transcriptional changes in transcriptomes while

progressing through development (Fig. 4B).

Closer inspection of key differentiation genes such as *Krt1*, *Krtdap* and *Notch1* revealed that

at E14.5 their expression was restricted to the suprabasal cell populations. In contrast, at

E15.5 and E16.5 a subset of basal cells also showed expression of differentiation genes (Fig.

4C). To conclusively identify a committed basal cell population, we examined a Harmony

integrated dataset of E14.5-E16.5 epidermal cells. Leiden clustering of this integrated dataset

revealed a distinct population of cells characterized by expression of both basal stem cell

(e.g. *Trp63* and *Itga6*) and terminal differentiation genes (Supplementary Fig. 4E, F),

indicating that only from E15.5 onward, a subset of basal cells commit to differentiation.

Next to this hybrid transcriptional signature of differentiation and stemness, this subset of

committed basal cells also showed enhanced expression of the suprabasal adhesion proteins

E-cadherin and desmoglein (Fig. 4D), consistent with a downregulation in heterotypic

interfacial tension between the committed cell and suprabasal cells predicted by the model

(Fig. 2D and 3O).

As Notch activity has been shown to be critical for terminal differentiation in keratinocytes
[30], we examined Notch pathway activity specifically in the basal layer. Receptor-ligand
enrichment analyses (CellPhoneDB; [31] predicted increased basal activity of especially
Notch1-Jagged2 signaling from E15.5 onward (Fig. 4E). Indeed, immunofluorescence
analysis revealed that Notch activity (cleaved intracellular domain; N-ICD), was detected in
single cells only in the basal layer of E15.5 and E16.5 epidermis, but not at E14.5 (Fig. 4F,
G). Intriguingly, these Notch1-active cells were also positive for the differentiation marker
K10 and displayed a wedge-shaped morphology (Fig. 4F), indicating that these cells are
committed to differentiation and undergoing delamination.

Collectively these data indicated the emergence of a committed basal cell state characterized
by a hybrid profile of stemness and differentiation, a change in cell shape, a switch to a
suprabasal cell adhesion profile and Notch activity.

**Generation of the mechanical boundary at E15.5 is concomitant with stiffening of the**
**basement membrane and rigidification of the basal layer**

We hypothesized that maturation of the tissue and specifically the basement membrane could
drive the emergence of the mechanical barrier at E15.5. Combined analysis of the scRNAseq
data (Fig. 5A) and quantitative multiplex tissue imaging using Imaging Mass Cytometry [32],
revealed an increase in basement membrane maturation and cell-matrix adhesion (Fig.5A-C;
Supplementary Fig. 5A). Importantly, collagen IV, a key component of the basement
membrane that is essential for its rigidification and stabilization, was increased from E14.5 to
E16.5 (Fig. 5B-C), suggesting maturation of the BM from E14.5-16.5. In agreement, the
hemidesmosomal $\alpha6\beta4$ integrin was also increased during this transition, and became more
and more confined to the epidermal-dermal junction (Fig. 5B, C; Supplementary Fig. 5A).
Consistently, atomic force microscopy-mediated force spectroscopy confirmed a significant
stiffening of the basement membrane from E14.5 to E16.5 (Fig. 5D).

Based on observations in other systems [10, 15-18, 33] we speculated that this change in
basement membrane mechanics promoted establishment of a mechanical barrier between
basal and suprabasal cells through rigidification of the basal layer. Analyses of actomyosin
tension by quantifying myosin phosphorylation (pMLC2) showed increased tension at apical
cell-cell junctions and specifically at the interface between basal and suprabasal cells at
E15.5 compared to E14.5 (Fig. 5E), indicating emergence of a mechanical barrier, providing
further support for the geometry-based apical tension inference (Fig. 3N). Furthermore,
analyses of cell shapes and tissue dynamics from live imaging showed that the cells in the
basal layer of E14.5 embryos were larger and more irregular shaped compared to cells at

E15.5 (Fig. 5F, G; Supplementary Movie 3). In agreement, at E15.5 cell density was

increased, (Fig. 5F, H), with decreased mobility of the basal layer cells as quantified by

particle image velocimetry (Fig. 5F, I; Supplementary Movie 3), in line with the geometry-

based basal tissue stiffness inference (Fig. 3N). Together, these changes were consistent with

a tissue-scale rigidity transition in the developing epidermis that occurs from E14.5-16.5,

from an initially floppier basement membrane and a fluid-like basal layer to a stiff basement

membrane and a jammed basal layer with enhanced cell-matrix adhesion and apical

actomyosin tension.

**Tissue rigidity and cell shape changes drive Notch-dependent delamination and control**

**of tissue crowding**

We next investigated whether the changes to tissue rigidity are sufficient to induce basal

layer jamming and the emergence of a mechanical barrier. For this we mimicked tissue

maturation *in vitro* by modulating cell density and substrate rigidity by engineering

micropatterned adhesive islands on polyacrylamide hydrogels of defined stiffness (Fig. 6A).

Culturing primary epidermal progenitors on micropatterned islands revealed that increasing

substrate stiffness under confinement not only substantially enhanced differentiation (Fig.

6B), but, importantly, promoted efficient separation of the undifferentiated basal and

differentiated suprabasal cell populations (Fig. 6C), indicative of boundary formation.

As the scRNAseq associated the emergence of committed cells with Notch activity (Fig. 4D),

we then asked whether basal layer crowding was sufficient to induce Notch signaling. For

this, we performed live imaging of primary epidermal progenitors isolated from Notch1

activity reporter mice [34] to identify emergence of Notch-active basal cells in cells triggered

to differentiate and form a suprabasal layer (Fig. 6D). As expected, all progenitors were

initially Notch-reporter negative, but upon application of calcium to trigger differentiation

and multilayering, individual cells gradually switched on the reporter followed by

delamination (Fig. 6D; Supplementary Movie 4). Intriguingly, prior to turning on Notch1

signaling these cells were more elongated and resided in denser tissue regions than reporter

negative-cells (Fig. 6D), thus indicating that crowding and increased cell-cell to cell-matrix

contact area promoted Notch activation.

We then asked whether Notch1 transcriptional activity is necessary for cell fate commitment

and delamination by deleting the Notch transcription factor, Rbpj, in the epidermis (Rbpj-

KO; [35, 36]). At E14.5, the epidermis of Rbpj-KO mice appeared largely comparable to

control littermates (Fig. 6E). At E15.5, and more prominently at E16.5, the epidermis became

structurally more abnormal, characterized by a specific thinning of the suprabasal

differentiated layers (Fig. 6F). Moreover, basal cells were increased in height and became

more columnar in E15.5 and E16.5 Rbpj-KO mice compared to controls (Fig. 6G),

accompanied by higher cell density (Fig. 6D), indicating increased crowding of the basal

layer. To examine if this increased crowding was due to reduced commitment to

differentiation or decreased delamination, we quantified differentiating, delaminating cells.

We observed emergence of the committed basal cell population in both control and Rbpj-KO

epidermis (Fig. 6H), indicating that the initial commitment to differentiate does not depend

on Notch activity. Instead, we observed an increase in the number of wedge-shaped

committed cells undergoing delamination at this stage (Fig. 6H), suggesting an inability to

overcome the mechanical energy barrier and move suprabasally.

To identify the molecular mechanism by which Notch transcriptional activity regulates

efficient delamination, we performed scRNAseq analysis across E14.5-E16.5 from control

and Rbpj-KO mouse skin. Overall, basal cell populations were transcriptionally similar

between controls and Rbpj-KO mice (Fig. 6I). However, as described previously [30], Rbpj-

KO cells of all developmental stages displayed impaired terminal differentiation (Fig. 6I;

Supplementary Fig. 6A). Intriguingly, at E15.5 a distinct Leiden cluster representing

committed basal cell population emerged, mainly composed of Rbpj-KO cells (Fig. 6I, J;

Supplementary Fig. 6B). Consistently, at E16.5 the Leiden cluster consisting of committed

cells was dominated by Rbpj-KO cells (Fig. 6I, J; Supplementary Fig. 6B), validating the

immunofluorescence observations that Rbpj-KO still commit to initial differentiation but

cannot efficiently delaminate, leading to accumulation of committed basal cells.

To identify master transcriptional switches downstream of Notch that drive the final step of

delamination, we performed differential gene expression analyses of committed cells between

control and Rbpj-KO at E15.5 and E16.5. This analysis identified two interesting

significantly altered regulators in Rbpj-KO committed cells: the zinc finger protein Zpf800, a

master regulator of intestinal stem cell differentiation [37], was not upregulated (Fig. 6K left

panel; Supplementary Fig. 6C, D). At the same time Rbpj-KO basal cells failed to

downregulate the Ep300 Inhibitor of Differentiation (Eid1) (Fig. 6K, right panel). Eid1 acts

as a transcriptional repressor through inhibition of p300-mediated H3K27 acetylation

(H3K27ac), thus preventing the activation of enhancers that promote differentiation [38, 39].

Indeed, RNA in situ hybridization and quantitative immunofluorescence confirmed that while

control embryos downregulated Eid1 in the suprabasal layers and, as a consequence, showed

substantially reduced H3K27ac levels in delaminating cells and suprabasally, Rbpj-KO

embryos maintained both high Eid1 and H3K27ac in delaminating cells and suprabasal

layers, in line with reduced differentiation (Fig. 6L).

Collectively, these data demonstrate that stiffening of the substrate is sufficient to drive basal

layer jamming and the formation of a mechanical barrier between basal and suprabasal

compartments. This basal layer jamming and crowding then triggers Notch signaling required

for completion of delamination and control of basal layer density.

**Discussion**

Here we show that the developing epidermis employs different strategies for multilayering

that are driven by changes in tissue mechanics. Early during development of this stratified

epithelium, there is no substantial energy barrier between basal and suprabasal cell layers.

This allows fast upward movement of basal cells independent of basal fate commitment by

either dividing perpendicularly or detaching quickly in order to rapidly establish a stratified

tissue. At later stages (E15.5 and beyond), maturation and stiffening of the basement

membrane enhance basal cell-matrix adhesion and crowding to rigidify the tissue. This

results in the generation of a substantial mechanical energy barrier between basal and

suprabasal cells. In order to cross this barrier, cells must commit to differentiation and

undergo subsequent transcriptional rewiring, controlled by Notch activation, to complete

delamination. Interestingly, this delamination process in the matured epidermis is

substantially slower than the rapid detachment observed by us and others in early embryonic

development [40]. As Notch activation itself is regulated by tissue rigidification and

crowding, it suggests an elegant control mechanism for homeostasis in an open niche, where

tissue crowding in a rigidified tissue triggers Notch activation to control the rate and timing

of delamination. This allows accurate control over the densities of both basal and suprabasal

layers.

Many open questions remain. A key open question is how the system decides which subset of

cells in the basal layer becomes committed and Notch active. Previous work has suggested

that since Notch is a surface receptor, its activity depends the cell shapes and surface areas

that regulate the density of such receptors [41, 42]. This could explain the emergence of

Notch active cells in elongated cells within dense regions – these cells would have transiently

maximized surface area to receive Notch ligand signal from their neighbors. Previous work

has investigated a 2D vertex model where cell fate decisions are coupled to cell areas [43],

which may be a useful starting point for predicting features we observe here. Future modeling

work should focus on understanding how changes to tissue stiffness and cell shape induce

Notch signaling in a 'correct' subset of basal cells (e.g. they are spaced out and the right fraction depending on the degree of crowding), and on adding the effects of differently oriented cell divisions in basal layer — in different developmental stages – to the 3D model.

Downstream of Notch signaling, it will be intriguing to understand how the transcriptional reprogramming, involving Eid1and Zfp800 associated with global changes in H3K27ac translates to control of delamination timing. It is interesting to note that the direct downstream targets of Eid1- the histone acetyltransferases P300 and CBP - are evolutionarily conserved maintenance factors of lineage-specifying core-transcriptional networks. They function as transcriptional rheostats that control global H3K27ac levels, where the duration of their activity determines the long-term impact on chromatin state [44, 45]. Upon their inhibition, >8h of hypoacetylation is required to trigger a permanent switch and "point of no return" to H3K27me3-mediated chromatin silencing [44]. Intriguingly, this is the approximate duration of the slower delamination process after E15.5. Thus, Notch may function as a mechanical sensor to ensure that terminal differentiation is not initiated prematurely in areas of low basal cell density. Precise coordination of onset of terminal differentiation with basal stem cell layer density is likely to be a critical feature in dynamically self-renewing tissues with an "open stem cell niche". Collectively, we propose that in such tissues mechanical boundaries are required to separate biological activities of cycling stem cells and post-mitotic terminally differentiated cells. Here, precisely controlled cell state transitions control the transit between the compartments to robustly maintain tissue homeostasis and barrier integrity.

## Methods

### Mice

Membrane-targeted Tomato reporter mice (R26R$^{mT/mG}$), nucleus-targeted Tomato reporter mice (R26R$^{nT/nG}$) and high-activity Notch1 activation-dependent reporter mice (N1IP: CreHI) were from were from JAX laboratories (stock #007676, #:023035 and #027234 respectively). Rbpj floxed mice were obtained from JAX (stock # 034200) and crossed with the K14-Cre line [35] to obtain epidermis-specific depletion. Embryos were staged based on vaginal plug (taken as embryonic day 0) and limb morphology. All mouse studies were approved and carried out in accordance with the guidelines of the Ministry for Environment, Agriculture, Conservation and Consumer Protection of the State of North Rhine-Westphalia (LANUW; 81-02.04.2022.A065; and 2024-155), Germany.

**Ex vivo embryo imaging**

Imaging was performed essentially as described earlier [10, 11]. Briefly, embryos were dissected at embryonic stages E14.5 or E15.5 and immobilized on custom-built imaging chambers engineered on top of Lumox-teflon imaging plates to allow gas exchange (Sarstedt). Whole embryos were immersed in epidermal growth medium (CnT-PR, CELLnTEC Advanced Cell Systems) supplemented with calcium-depleted 10% fetal bovine serum (HyClone/Thermo Fisher), 20 U/ml penicillin-streptomycin (Thermo Fisher), 20 mM Hepes, 1.8 mM $CaCl_2$, and 10 ng/ml mouse recombinant EGF (Novus; NBP2-34952). Imaging was performed using an Andor Dragonfly 505 high-speed spinning disk confocal microscope (Oxford Instruments) with an inverted Nikon Eclipse Ti2 microscope (Nikon) equipped with 488 nm and 546 nm lasers, an Andor Zyla 4.2 sCMOS camera and an environmental chamber set at 37°C, 5% $CO_2$ using a 40X dry objective. Images were acquired using Fusion 2.0 software (Andor). After acquisition, the image series were 3D drift corrected using the ImageJ plug-in Fast4Dreg [46] combined with manual registration, if required.

**Immunofluorescence and confocal microscopy**

For skin whole mounts, embryos were dissected from pregnant mice and directly fixed in pre-warmed (37°C) 4% paraformaldehyde (PFA) at room temperature for 3 h. After multiple PBS washes, the skin was carefully peeled using forceps and placed in blocking solution (0.5% Triton-X, 5% bovine serum albumin (BSA), 3% normal goat serum (NGS)) for 1 h at room temperature. For paraffin sections tissue biopsies were fixed in 4% PFA, embedded in paraffin and sectioned. Sections were de-paraffinized using a graded alcohol series and antigen retrieval was carried out using target retrieval solution (Dako) at pH 6 or pH 9 in a pressure cooker. For cryosections, tissue samples were placed unfixed into cryomolds, embedded in OCT (Tissue Tek) and allowed to harden on dry ice. 8 μm thick sections were cut, air-dried and fixed with 4% PFA. For both paraffin and cryofixed sections, samples were blocked in 5% BSA, 3% NGS. For paraffin sections antibodies were diluted in Dako antibody diluent (Agilent). For cryosections antibodies were diluted in blocking solution. Cultured epidermal cells were fixed in 4% PFA for 15 min at room temperature After PBS washes, samples were placed in 0.3% Triton-X, 5% BSA for one hour at room temperature after which primary antibodies were diluted in the same solution. In all cases antibodies were incubated overnight in moist chamber at 4°C. Following PBS washes, secondary antibodies were diluted in 1% BSA in PBS and incubated for 1h at room temperature.

For super-resolution imaging of Dsg1, K10, E-cadherin, and Integrin α6, the embryos were

fixed in pre-warmed 4% PFA at 37°C for 1 hour and rinsed with PBS. The back skin was

peeled using forceps and incubated in blocking solution for 24 hours at 37°C. Primary

antibodies were diluted in the blocking solution and incubated 48 hours at 37°C on shaker

followed by several washes with 2% Triton-X with a final overnight wash. Secondary

antibodies were diluted in blocking solution and incubated 24 hours at 4°C followed by

several washes with 2% Triton-X.

For pMLC2 staining embryos were fixed in 4% PFA dissolved in HBSS for 1 hour on a

shaker at room temperature. Back skins samples are then peeled and permeabilized with

0.02% PBS-Tween 20 for 30mins and blocked with blocking solution (5% NGS,1% BSA) for

2 hours before incubating with primary antibody overnight in 4ºC. Samples are then

thoroughly washed with 0.02% PBS-Tween 20, incubated with secondary antibodies at 1:400

dilution for 1 hour at room temperature and washed. All tissue samples were mounted in

Elvanol.

The following primary antibodies were used: guinea-pig anti-Keratin 10 (Progen, #GP-10;

1/300);  rabbit anti-p-MLC2 (p-Ser19; Cell Signaling, 3675; 1:200), rabbit anti-cleaved-

Notch1 (Cell signaling; #4147S; 1/50), mouse anti-Desmoglein 1/2 (Progen, #61002; 1/200),

rabbit anti-TGM1 (360; Proteintech, #12912-3-AP; 1/300), rabbit anti P63 (Genetex

#GTX102425; 1/100), guinea-pig anti-Keratin 14 (Progen, GP-CK14; 1:300). Bound primary

antibody was detected by incubation with Alexa-Fluor 488-, 568- or 647 -conjugated

antibodies (Invitrogen). Nuclei were counterstained with 4′, 6-diamidino-2-phenylindole

(DAPI, Invitrogen). Slides were mounted in Elvanol.

Fluorescence images were collected by laser scanning confocal microscopy (LSM980; Zeiss)

with with Zeiss ZEN Microscopy software (Zeiss ZEN version 3.7) using a 40X water

immersion objective. Images were acquired at room temperature using sequential scanning of

frames of 1 μm thick confocal planes (pinhole 1). Images were collected with the same

settings for all samples within an experiment. For super-resolution imaging a 63X oil

objective and the Airyscan 2 SR (super-resolution) mode with default resolution settings

were used. Laser power and image size were kept at a minimum to avoid photo-bleaching.

All airyscan images were 3D processed using the Zeiss Airyscan Processing mode with auto

filter and standard strength.

**Embryonic epidermal vertex analysis**

Membrane-tomato positive E14.5, E15.5 and E16.5 embryos were isolated, fixed

immediately in 4%PFA over night at 4°C and washed in PBS. The back skin was then

removed using micro-scissors and forceps and permeabilized in PBS/0.5% Triton-X for 1h at RT. Nuclei were stained by incubating the skin in DAPI diluted in DAKO antibody diluent for 1h at RT. The skin was then mounted in Mowiol.

Confocal stacks were recorded at a voxel size of X: 0.361, Y: 0361, Z:0.299 using a Leica SP8 equipped with an APO CS2 63x/1.40 oil objective. To determine basal cell vertices, confocal stacks were processed in Imaris and minimal XZ or YZ projection were exported using the extended section view. Using the segmented line tool in ImageJ a line was drawn across the apical and basal vertices of basal cells. Coordinates were extracted and "Julia" was used in Visual Code Studio to determine cell height (h), apical and basal vertex distances (a) and the lateral interface orientation (angle beta). The lines connecting the basal and apical vertices, respectively, were used to determine the apical layer roughness (R) (See also Supplementary Methods). The apical curvature angle gamma was determined manually using the angle tool in ImageJ.

**Image analyses**

For tissue-scale analysis, 2D segmentation were performed from live imaging data on whole embryos or cultured epidermal progenitor monolayers using Cellpose4SAM (v4.0.8) [47]. 2D cell shape parameters were measured using custom Python script and cell densities were measured using ImageJ. For analyses of cell shapes of delaminating cells from embryo live imaging, Imaris 11 was used to manually create 3D surfaces of delaminating and corresponding neighboring basal cells. Meshes were exported from Imaris surfaces after which shapes were analyzed by quantifying Spherical Harmonics [48] using a custom Python script.

In fixed tissue samples, basal committed cells and delaminations were identified based on the expression of specific markers (K10 and/or Desmoglein) and their contact with the basement membrane, and subsequently manually counted. Active-Notch positive cells were quantified manually. The method was applied across all stages and conditions. Integrin $\alpha6$, E-cadherin and Desmoglein mean intensities were manually measured at the basal, apical and lateral membranes on Image J. To quantify epidermis thickness, the distance between epidermal basement membrane and most superficial point of the epidermis was measured. To quantify specifically the basal layer thickness versus the differentiated layer thickness, the expression of specific markers (K10 and desmoglein) has been used to distinguish the differentiated layers (K10+) from the basal layer.

To measure boundary straightness between the bottom and upper layers in micropatterns of

soft and stiff polyacrylamide (PAA) hydrogels, both a segmented line following the top of

each nucleus in the bottom layer and a straight line were drawn. The lengths of these lines

were measured using ImageJ, and the ratio of the straight line to the segmented line was

calculated.

Eid1 RNA signal was quantified by segmenting suprabasal cells based on Keratin-DAP–

positive cells. Mean fluorescence intensity of Eid1 mRNA was measured using Fiji.

**3D vertex model**

A dynamic three-dimensional model of stratified epithelial tissues was developed using the 3D

vertex framework (Zhang and Schwarz, 2022). In the 3D vertex model, cells are represented

as polyhedra, each characterized by a homeostatic volume $V_0$ and surface area $S_0$. The

geometry of each cell is defined by its vertices, which serve as the degrees of freedom of the

system. The modeled epithelium consists of two main cell compartments. The upper

compartment represents the suprabasal differentiated cell layer, while the middle one-cell-thick

layer corresponds to the basal stem cell layer. Because the vertex model treats cells as space-

filling polyhedra, we modeled the basement membrane as an additional layer of interconnected

"cellular network" directly attached beneath the basal layer. In this framework, cells exert

forces on one another through shape-based interactions governed by the following energy

functional :

$E = \sum_\alpha K_S (S_\alpha - S_0)^2 + \sum_\alpha K_V (V_\alpha - V_0)^2 + \sum_{\alpha\beta} \boldsymbol{\sigma}_{\alpha\beta} S_\alpha$   (1)

The first term of the energy functional describes the competition between contractile forces

generated by cell actomyosin networks (which enable cells to minimize their shape or surface

area to resist their environments), and adhesive forces mediated by cadherin-based, which

promote the elongation of cell–cell contacts during neighbors' exchange (T1s). The second

term accounts for the mechanical resistance of cells to volume fluctuations, as most cells are

fluid-filled and resist changes in volume. The third term incorporates heterotypic interfacial

tension $\sigma_a$ between the basal cells and suprabasal cells, and the tension $\sigma_b$ associated with

integrin-based wetting behaviors at basal-basement membrane interface.

Previous studies have shown that differential expression of adhesion molecules and

cytoskeletal components associated with cell fate specification (through variations in protein

or gene expression) can modify cell–cell interactions, leading to differences in adhesion

strength and heterotypic interfacial tension [10, 11, 49-57]. Such tension differences are known

to play a central role in establishing and maintaining sharp tissue boundaries [55]. Moreover,
basal cells adhere to the basement membrane via integrins, which introduce a wetting
interaction characterized by $\sigma_b$ [58]. The parameters $K_S$ and $K_V$ denote the elastic moduli
associated with the surface area and volume of the cells, respectively. Upon non-
dimensionalization, this energy functional allows the definition of a key order parameter
known as the target shape index, $s_0 = S_0 V_0^{-\frac{2}{3}}$, which controls the mechanical rigidity of the
tissue. By tuning $s_0$, the epithelial confluent tissue exhibits a rigidity transition from a solid-
like to a fluid-like state, with a critical value at $s_0 = 5.4$ [15, 17].
To quantitatively predict the mechanical parameters and cellular mechanisms underlying
mechanical barrier formation between the basal and suprabasal layers, we first simulated the
epithelium in the absence of cell delamination. The system evolves according to overdamped
Langevin dynamics, where vertex positions are continuously updated to minimize the total
mechanical energy:

$$\partial_t r_i = -\mu \nabla_i E_i + \sqrt{2D_t}\, \boldsymbol{\eta}(t). \qquad (2)$$

The left-hand side represents viscous drag forces, the first term on the right-hand side
corresponds to interaction forces derived from the energy potential, and the second term
represents thermal fluctuations in vertex positions $r_i$ . Here $\boldsymbol{\eta}$ is an uncorrelated stochastic
Gaussian white noise with unit amplitude, $D_t = \mu k_B T_{\text{eff}}$ , where $\mu$ is the mobility (inverse
frictional drag), $k_B$ Boltzmann constant, and $T_{\text{eff}}$ is an effective temperature parameter that
sets the scale of the fluctuations. Unless otherwise noted, for our simulations we choose $\mu = $
1, $k_B = 1$ and $T_{\text{eff}} = 0.002$. This choice of $T_{\text{eff}}$ allows fluctuations of interface lengths but
is not sufficient to generate diffusion or fluidity in the basal layer.

From the simulated model, we can compute the effective tensions governing different
interfaces: a) The homotypic tension between basal cells $\sigma_{bb} = 4K_S \Delta s_b$ where $\Delta s = s_\alpha - s_0$
measures the basal layer stiffness (and $\Delta s = 0$ at the transition point); b) The effective
heterotypic tension at the basal-suprabasal interface $\Gamma_a = 2\sigma_a + 2K_s(\Delta s_s + \Delta s_b)$, where $\Delta s_s$
the stiffness of the suprabasal layer; c) The effective wetting tension $\Gamma_{bm} = 2\,\sigma_b + 2K_s(\Delta s_b +$
$\Delta s_m)$ between basal cells and the basement membrane (constituted of a cellular network with

a small shape parameter). Unless otherwise noted, $\Delta s_m = 0.26,$ which ensures the basement

membrane remains very stiff and solid-like.

For more detailed description see Supplementary Methods.

**Single-cell RNA sequencing (scRNAseq)**

E13.5, E14.5, E15.5 and E16.5 embryos from wild type, control or K14Cre-RBP-J

floxed/floxed mice (two embryos/genotype/stages) were collected simultaneously and back

skins were peeled off. Single-cell suspensions were generated by enzymatic digestion by

incubating back skin tissues in a mixture of DNAse I (40 µg/ml), Liberase (0.125 mg/ml;

Roche Cat. NO. 05401119001), and papain (30 U/ml; Worthington) in Minimum Essential

Medium, Spinner modification (Gibco 11380) for 15 min (E13.5), 20 min (E14.5), 30 min

(E15.5) or 40min (E16.5) at 37°C followed by mechanical dissociation. The reaction was

stopped with 22% FBS and single cells were washed with FACS buffer (0.2% chelated FBS,

2 mM EDTA in 1x DPBS (Sigma Aldrich, D8537)). For wild type sequencing, cells were

labeled with BD Mouse Single-Cell Multiplexing Kit (Cat # 633793): E13.5 rep 1: Sample

Tag (ST) 1; E13.5 rep 2: ST 2; E14.5 rep1: ST 3; E14.5 rep2: ST 4; E15.5 rep1: ST 5; E15.5

rep2: ST 6; E16.5 rep1: ST 7, E16.5 rep2: ST 8. For K14Cre-Rbpj floxed and control

sequencing, surface biotinylation was used [59]. Briefly, cells were labeled with 50 µg/mL

EZ-Link sulfo-NHS-biotin (Thermo Fisher Scientific, 21217) on ice for 10 min, followed by

incubation in 0.6 µg/ml TotalSeq-A PE streptavidin hashtag antibodies (BioLegend) for 30

min on ice as follows: E14.5 Rbpj-KO 1: TotalSeq™-A0951 PE Streptavidin 405251 ; E14.5

Rbpj-KO 2: TotalSeq™-A0952 PE Streptavidin 405253; E14.5 control 1: TotalSeq™-A0953

PE Streptavidin 405255; E14.5 control 2: TotalSeq™-A0954 PE Streptavidin 405257; E15.5

Rbpj-KO 1: TotalSeq™-A0951 PE Streptavidin 405251 ; E15.5 control 1: TotalSeq™-

A0953 PE Streptavidin 405255; E15.5 control 2: TotalSeq™-A0954 PE Streptavidin 405257;

E16.5 Rbpj-KO 1: TotalSeq™-A0951 PE Streptavidin 405251 ; E16.5 Rbpj-KO 2:

TotalSeq™-A0952 PE Streptavidin 405253; E16.5 control 1: TotalSeq™-A0953 PE

Streptavidin 405255; E16.5 control 2: TotalSeq™-A0954 PE Streptavidin 405257. Labeled

cells were loaded on a microwell cartridge of the BD Rhapsody Express system (BD)

following the manufacturer's instructions. Single-cell whole transcriptome analysis libraries

were prepared according to the manufacturer's instructions using BD Rhapsody WTA

Reagent kit (633802; BD), except for the Rbpj-KO and control sample tag libraries in which

totalseq-ADT-oligo1 (5'-TGCTCTTCCGATCTTGGCACCCGAGAATTCCA-3') primer in

place of Sample Tag PCR1 and Sample Tag PCR2 primers during the Sample Tag library

amplification [59]. Libraries were sequenced on the Illumina NextSeq 500 using High Output

Kit v2.5 (150 cycles; Illumina) for 2 × 75-bp paired-end reads with 8-bp single index aiming

sequencing depth of >20,000 reads per cell for each sample.

**scRNAseq analyses**

*Read Data QC and Mapping*

Illumina sequencing results were demultiplexed and converted to FASTQ format using

Illumina's bcl2fastq software (version 2.20.0.422). For K14Cre-RBP-J floxed/floxed data a

custom demultiplexing script was used to assign barcodes to sample tags (rhapsody-

demultiplex with parameters '-w BD_CLS1_v2.txt BD_CLS2_v2.txt BD_CLS3_v2.txt -a

TGGCACCCGAGAATTCCA <fastq files> sample_tags.txt', and sample tags as described

above).

Preprocessed read data was aligned to the mouse reference genome (GRCm39) and counted

with STAR [60] (version 2.7.11b) through its single-cell functionality STARsolo, using the

following parameters:

WT: `--runThreadN 16 --soloType CB_UMI_Complex --soloCellFilter None --outSAMtype

BAM SortedByCoordinate --soloFeatures Gene GeneFull GeneFull_Ex50pAS --

soloAdapterSequence NNNNNNNNNGTGANNNNNNNNNGACA --soloCBmatchWLtype

1MM --soloCBposition 2_0_2_8 2_13_2_21 3_1_3_9 --soloUMIposition 3_10_3_17 --

soloCBwhitelist BD_CLS1_v2.txt  BD_CLS2_v2.txt BD_CLS3_v2.txt --runRNGseed 1 --

soloMultiMappers EM --readFilesCommand zcat --outSAMattributes NH HI AS nM NM

MD jM jI MC ch CB UB GX GN`

K14Cre-RBP-J floxed/floxed: `--soloType CB_UMI_Complex --soloAdapterSequence

NNNNNNNNNGTGANNNNNNNNNGACA --soloCBposition 2_0_2_8 2_13_2_21

3_1_3_9 --soloUMIposition 3_10_3_17 --soloBarcodeReadLength 0 --soloCellFilter None --

soloCBwhitelist BD_CLS1_v2.txt BD_CLS2_v2.txt BD_CLS3_v2.txt --

soloCBmatchWLtype 1MM --soloUMIfiltering MultiGeneUMI_CR --soloUMIdedup

1MM_CR --outSAMtype BAM SortedByCoordinate --soloFeatures Gene GeneFull

GeneFull_Ex50pAS --soloMultiMappers EM --outSAMattributes NH HI AS nM NM MD jM

jI MC ch CB UB GX GN --outSAMmultNmax 1 --outMultimapperOrder Random --

runRNGseed 1 --outFilterScoreMin 30 --outFilterMatchNmin 15`

*Data Filtering, Normalisation, and Visualization*

Preprocessing of the feature-count-matrix output by STARsolo was largely performed within the scanpy [61]ecosystem, with the exception of scran normalization [62], which was performed in R (R Core Team, 2017). All visualization was done using [63] functions. Default parameters were used unless detailed otherwise.

For wild type:

Barcodes with less than 3000 UMI were discarded.

Only barcodes with less than 15% of counts stemming from mitotic genes were retained.

For K14Cre-RBP-J floxed/floxed and control:

Barcodes with less than 2000 UMI were discarded.

Only barcodes with less than 20% of counts stemming from mitotic genes were retained.

Only barcodes with less than 70% of counts stemming from hemoglobin genes were retained.

And barcodes were filtered for a complexity value above 0.75, where complexity is defined as the log10-transformed number of expressed genes divided by the log10-transformed number of total counts per barcode. Scran was used for normalization.

The separate samples were merged into a single object, G2M and S phase scores were assigned to each cell using gene lists from [64] and the scanpy scanpy.tl.score_genes_cell_cycle function, and regressed out of the gene expression matrix using sc.pp.regress_out. 3000 highly variable genes were calculated from the normalized matrix (sc.pp.highly_variable_genes, flavor "seurat"), and the first 100 principal components were calculated with scanpy. Neighbourhood graph calculation (30 neighbors), UMAP dimensionality reduction [65], and Leiden clustering [66] were performed. scrublet [67] and/or scDblFinder were then used for doublet prediction.

Contaminant or undesired cell populations such as clusters of cells that, either manually or by scrublet/scDblFinder, could be classified as doublets based on their expression patterns, were removed. For this, cells were clustered at resolution 0.05, and the cluster corresponding to the epidermal cells were isolated. These were then reclustered at resolution 0.1 (wild type) and 0.2 (K14Cre-RBP-J floxed/floxed) as described above, and undesired clusters were removed. Finally, PCA, nearest neighbor calculation, and UMAP dimensionality reduction were repeated for the clean epidermis data. For WT, stages E14.5, E15.5, and E16.5 were

additionally analysed separately with batch-correction using Harmony [68] after cell-cycle

regression.

*Differential expression analysis*

Differentially expressed genes were calculated using a pseudobulk approach [69], comparing

aggregate counts with 2 pseudoreplicates for each condition (pyDeSEQ2 0.4.8).

*Ligand-receptor interaction analysis*

We used CellphoneDB (v2.1.7 [31] with default parameters to analyse ligand-receptor

interactions. Mouse gene names were converted to human gene names prior to the analysis.

*Gene Set Enrichment Analysis*

We ranked differential expression results by pyDESeq2 shrunken log2-fold changes and used

GSEApy (v1.0.6 [70]) "prerank" to calculate the enrichment statistics.

**Imaging Mass Cytometry**

Primary antibodies from bovine serum albumin (BSA)- and azide-free formulations were

conjugated using the Maxpar X8 Labeling Kit (Standard BioTools) according to the

manufacturer's protocol. Stainings were performed on frozen cryosections according to the

manufacturer's protocol (Fluidigm). Briefly, 5 μm cryosections were prepared and stored at

−80 °C until use. On the day of the experiment, slides were equilibrated at −20 °C for 1 h and

then fixed in 4% paraformaldehyde (PFA) for 30 min at 4 °C. After washing in PBS, sections

were incubated in blocking solution (3% BSA) for 45 min at room temperature, followed by

incubation with the antibody cocktail prepared in 0.5% BSA for 1 h in a hydration chamber at

room temperature. Slides were then rinsed in 0.2% Triton X-100/PBS, stained with Cell-ID

Intercalator-IR (Fluidigm #201192A) for 30 min at room temperature, washed in Milli-Q

water, air-dried for 30 min, and stored until imaging. The samples were laser-ablated at 200

Hz using a Hyperion imaging mass cytometry system (Fluidigm) coupled to a Helios time-of-

flight mass cytometer with a resolution of 1 μm. Raw MiniCAD Design (MCD) files

generated by the Hyperion Imaging System were first inspected using the MCD Viewer

(Standard BioTools) to verify signal intensity and overall acquisition quality, and then

exported as TIFF images for downstream analysis on Image J. ROIs delineating the basement

membrane were drawn, and signal intensity at the basement membrane was measured using

ImageJ software.

**Atomic Force Microscopy**

For Atomic Force Microscopy (AFM), embryonic skin was dissected, immediately nitrogen frozen and then embedded in OCT blocks. AFM measurements of the basement membrane were performed on freshly cut 20 μm-thick cryosections using JPK Nano Wizard IV (Bruker Nano) atomic force microscope mounted on an Olympus IX73 inverted fluorescent microscopy (Olympus) and operated via JPK SPM Control Software v.5. Cryosections were equilibrated in PBS supplemented with protease inhibitors and measurements were performed within 20 minutes of thawing the samples. Triangular non-conductive Silicon Nitride cantilevers (MLCT, Bruker Daltonics) with a nominal spring constant of 0,07 Nm$^{-1}$ were used for the nanoindentation experiments. For all indentation experiments, forces of up to 3 nN were applied, and the velocities of cantilever approach and retraction were kept constant at 10 μms$^{-1}$ ensuring an indentation depth of 500 nm. All analysis were performed with JPK Data Processing Software (Bruker Nano). Prior to fitting the Hertz model corrected by the tip geometry to obtain Young's Modulus (Poisson's ratio of 0.5), the offset was removed from the baseline, contact point was identified, and the cantilever bending was subtracted from all force curves. Force curves were filtered for quality and all valid forces curves were analyzed.

**Particle Imaging Velocimetry (PIV)**

PIV analysis of whole-embryo live imaging was performed as previously described, with minor adjustments [11]. Briefly, prior to analysis, each movie was processed as follows: a 3D drift correction was applied using the Fast4DReg plug-in in Fiji and sagittal cross-sections containing the basal layers were selected. A median filter of 2 pixels was applied in Fiji, followed by background subtraction using a rolling ball radius of 50 pixels.The movies were then imported into PIVLab version 2.62. Tissue flow tracking was restricted to the interfollicular epidermis. To exclude the hair follicle placodes from the analysis, ROIs covering the placodes were manually drawn on the raw images and imported into PIVLab. These masks were overlaid onto the flow field to exclude the placode regions. To prevent tracking errors due to intensity variations, image contrast was enhanced using CLAHE (Contrast Limited Adaptive Histogram Equalization) with a box size of 64 pixels. Denoising was performed using a Wiener Noise Filter with a maximum pixel count of 3. Tissue flow tracking was carried out using a multi-pass, windowed FFT-based approach, with window sizes of 64 and 32 pixels for the first and second passes, respectively, and a 50% overlap. The

779    resulting vector fields were visually inspected, and any erroneous vectors were manually

780    removed from the analysis.

**Culture of primary epidermal progenitor cells**

Primary epidermal cells were isolated from newborn mice essentially as described [11].

Briefly, after dissection, skins were treated with antibiotic/antimycotic solution (Sigma

A5955) for 5 min, and epidermal single cell suspensions were generated by incubating skin

pieces in 0.8% trypsin for 30 min at 37 °C. Cells were suspended in Keratinocyte Growth

Medium (MEM Spinner's modification (Sigma), 5 µg/ml insulin (Sigma), 10 ng/ml EGF

(Sigma), 10 µg/ml transferrin (Sigma), 10 µM phosphoethanolamine (Sigma), 10 µM

ethanolamine (Sigma), 0.36 µg ml/ml hydrocortisone (Calbiochem), 2 mM glutamine

(Gibco), 100 U ml/ml penicillin, 100 µg ml/ml streptomycin (Gibco), 10% chelated fetal calf

serum (Gibco), 5 µM Y27632, 20 ng ml/ml mouse recombinant vascular endothelial growth

factor (VEGF) and 20 ng ml/ml human recombinant fibroblast growth factor-2 (FGF-2) (all

from Miltenyi Biotec) and seeded on cell culture plates coated with 30 µg ml$^{-1}$ collagen I

(Millipore) and 10 µg ml$^{-1}$ fibronectin (Millipore) for 1 h at 37 °C.

**Micropatterned hydrogels**

Circular micropatterns (100 µm diameter) were generated on glass coverslips activated by

oxygen plasma treatment for 3 min (PDC-002-CE Expanded Plasma Cleaner), as described

previously [10]. Activated coverslips were incubated with 1× PLL (Surface Solutions) for 30

min at room temperature, rinsed with 10 mM HEPES (pH 7.8), and subsequently incubated

with PEG-SVA (50 mg/ml in 10 mM HEPES, pH 7.8) for 1 h at room temperature.

Coverslips were rinsed with Milli-Q water and air-dried before patterning by deep-UV

(Primo; Alveole). PEG-cleaved regions were rinsed with Milli-Q water and incubated with 50

µg/ml fibronectin for 30 min at room temperature. In parallel, glass-bottom plates (Ibidi,

#81218-200) were silanized. Polyacrylamide gels of defined stiffness were prepared by

placing a drop of acrylamide/bis-acrylamide solution between the micropatterned coverslip

and the silanized plate. After 45 min of polymerization, the coverslip was removed,

transferring the ECM micropatterns onto the gel surface. Primary epidermal progenitors were

then seeded onto the patterned hydrogels.

**Live imaging of epidermal progenitor monolayers**

Freshly isolated epidermal progenitors isolated from  N1IP: CreHI reporter mice crossed with

R26R$^{mT/mG}$ mice to label the plasma membrane and detect Cre activity were grown to

confluency in Keratinocyte Growth Medium and plated on 35 mm glass bottom dish with 14

mm micro-well plates (Cellvis; D35-14-1.5-N). Upon confluency, 200 µM $Ca^{2+}$ was added to

induce differentiation. Imaging was performed using an Andor Dragonfly 505 high-speed

spinning disk confocal microscope (Oxford Instruments) with an inverted Nikon Eclipse Ti2

microscope (Nikon) equipped with 488 nm and 546 nm lasers, an Andor Zyla 4.2 sCMOS

camera and an environmental chamber set at 37°C, 5% CO2 using a 40X dry objective.

Images were acquired using Fusion 2.0 software (Andor) with multiple positions and drift

stabilizer.

**RNA in situ hybridization**

RNA in situ hybridization was performed using RNAscope technology according to the

manufacturer's instructions (ACD, RNAscope Multiplex Fluorescent v2 Assay, kit:

RNAscope Multiplex Fluorescent v2 Assay ACD, #323100). Briefly, paraffin-embedded

tissue sections (5 µm thick) were deparaffinized in xylene, rehydrated through an ethanol

series, and incubated in hydrogen peroxide for 10 min at RT. Sections were then subjected to

antigen retrieval in RNAscope Target Retrieval solution for 15 min at 100°C using a pressure

cooker. RNAscope Protease Plus (ACD; #322331) was applied to each section and incubated

at 40°C for 30 min. The Eid1 RNAscope probe (ACD; #1713581-C3) and Keratin-DAP

RNAscope probe (ACD; #500671-C1) were hybridized according to the manufacturer's

protocol using the RNAscope Multiplex Fluorescent v2 Assay (ACD; #323100), followed by

signal detection with TSA Plus Fluorescein (Akoya, #NEL741001KT), TSA Plus Cyanine 3

System (Akoya, #NEL744001KT). All hybridization and amplification steps were performed

in a HybEZ II oven (ACD). DAPI solution was incubated for 30 seconds at room

temperature. Sections were mounted in Elvanol and imaged using a Zeiss LSM 980 confocal

microscope equipped with a 40× water-immersion objective, using Zeiss ZEN software

(version 3.5).

**Statistics and reproducibility**

Statistical analyses were performed using GraphPad Prism software (GraphPad, version 9) or

in R (version 4.2.2). Statistical significance was determined by the specific tests indicated in

the corresponding figure legends. Only 2-tailed tests were used. In all cases where a test for

normally distributed data was used, normal distribution was confirmed with the

Kolmogorov–Smirnov test ($\alpha = 0.05$). All experiments presented in the manuscript were

repeated at least in 3 independent replicates.

**Data availability**

Sequencing datasets are available at GEO (GSE315748 and GSE315746). All analysis

scripts, custom code, and data that support the conclusions are available from the authors on

request.

**Acknowledgements**

We thank Claudia Ortmeier and Sandra Heising for expert technical assistance, Silvia Fre

(Institute Curie, France) for advice on the project and general support, Fabien Bertillot for

advice on image analyses, Yekaterina Miroshnikova for advice on micropatterning, and the

Max Planck Institute BioOptics and Sequencing core facilities for support with experiments.

Imaging Mass Cytometry was performed at the Cell Imaging and Cytometry Core, Turku

Bioscience Centre (Turku, Finland), with the support of Biocenter Finland. This work was

supported by European Union's Horizon 2020 research and innovation programme under

Marie Skłodowska-Curie grant agreement no. 101032331 (to CV), Research Council of

Finland (332402) and The Turku Collegium for Science, Medicine, and Technology (to GF),

Academy of Finland Center of Excellence BarrierForce (346131) and R'Life Programme

consortium NucleoMech (330033) (to JI and SAW) the Sigrid Juselius Foundation, Helsinki

Institute of Life Science, and the Max Planck Society (all to SAW), NSF-CMMI-1334611 (to

SA and MLM), grant number 2023-329572 from the Chan Zuckerberg Initiative DAF, an

advised fund of the Silicon Valley Community Foundation (to MLM), and the Syracuse

University Office of Research Future Faculty Fellow Program (to SA). CMN is supported by

the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation: Germany´s

Excellence Strategy – EXC 2030 – 390661388; SPP 1782 NI 1234/6-2, Project-ID

388932620; FOR 2743 NI 1234/7-1 Project-ID 678823;  ANR/DFG NI 1234/9 (Project-ID

505673300) and the European Union NETSKINMODELS, COST Action no. CA21108.

**Author contributions**

CV, MA, and MR designed and performed experiments and analyzed data. SHAE and MLM

developed vertex model and performed all simulations and analyses. KK performed

sequencing data analysis. SV and LCB supported live imaging and single cell sequencing

experiments. AP performed and analyzed immunofluorescence experiments, PZ assisted with

imaging. GF and JI performed mass cytometry experiment. CMN, MLM and SAW conceived

and supervised the study, designed experiments, and analyzed data. MLM and SAW wrote

the paper. All authors commented and edited the manuscript.

**Competing interests statement**

885 The authors declare no competing interests.

**References:**

1. Kulukian, A. and E. Fuchs, *Spindle orientation and epidermal morphogenesis.* Philos
Trans R Soc Lond B Biol Sci, 2013. **368**(1629): p. 20130016.
2. Niessen, C.M., M.L. Manning, and S.A. Wickstrom, *Mechanochemical Principles of*
*Epidermal Tissue Dynamics.* Cold Spring Harb Perspect Biol, 2025. **17**(6).
3. Tai, K., K. Cockburn, and V. Greco, *Flexibility sustains epithelial tissue homeostasis.*
Curr Opin Cell Biol, 2019. **60**: p. 84-91.
4. Beumer, J. and H. Clevers, *Hallmarks of stemness in mammalian tissues.* Cell Stem
Cell, 2024. **31**(1): p. 7-24.
5. Williams, S.E. and E. Fuchs, *Oriented divisions, fate decisions.* Curr Opin Cell Biol,
2013. **25**(6): p. 749-58.
6. Damen, M., et al., *High proliferation and delamination during skin epidermal*
*stratification.* Nat Commun, 2021. **12**(1): p. 3227.
7. Lechler, T. and E. Fuchs, *Asymmetric cell divisions promote stratification and*
*differentiation of mammalian skin.* Nature, 2005. **437**(7056): p. 275-80.
8. Cockburn, K., et al., *Gradual differentiation uncoupled from cell cycle exit generates*
*heterogeneity in the epidermal stem cell layer.* Nat Cell Biol, 2022. **24**(12): p. 1692-
1700.
9. Mesa, K.R., et al., *Homeostatic Epidermal Stem Cell Self-Renewal Is Driven by Local*
*Differentiation.* Cell Stem Cell, 2018. **23**(5): p. 677-686 e4.
10. Miroshnikova, Y.A., et al., *Adhesion forces and cortical tension couple cell*
*proliferation and differentiation to drive epidermal stratification.* Nat Cell Biol, 2018.
**20**(1): p. 69-80.
11. Villeneuve, C., et al., *Mechanical forces across compartments coordinate cell shape*
*and fate transitions to generate tissue architecture.* Nat Cell Biol, 2024. **26**(2): p. 207-
218.
12. !!! INVALID CITATION !!! (Damen, Wirtz et al. 2021).
13. Moll, R., et al., *The catalog of human cytokeratins: patterns of expression in normal*
*epithelia, tumors and cultured cells.* Cell, 1982. **31**(1): p. 11-24.
14. Nekrasova, O., et al., *Desmosomal cadherin association with Tctex-1 and cortactin-*
*Arp2/3 drives perijunctional actin polymerization to promote keratinocyte*
*delamination.* Nat Commun, 2018. **9**(1): p. 1053.
15. Zhang, T. and J.M. Schwarz, *Topologically-protected interior for three-dimensional*
*confluent cellular collectives.* Physical Review Research, 2022. **4**(4): p. 043148.
16. Bi, D., et al., *A density-independent rigidity transition in biological tissues.* Nature
Physics, 2015. **11**(12): p. 1074-1079.

17.  Merkel, M. and M.L. Manning, *A geometrically controlled rigidity transition in a model for confluent 3D tissues.* New Journal of Physics, 2018. **20**(2): p. 022002.

18.  Devany, J., et al., *Cell cycle-dependent active stress drives epithelia remodeling.* Proc Natl Acad Sci U S A, 2021. **118**(10).

19.  Erdemci-Tandogan, G., et al., *Tissue Flow Induces Cell Shape Changes During Organogenesis.* Biophys J, 2018. **115**(11): p. 2259-2270.

20.  Serwane, F., et al., *In vivo quantification of spatially varying mechanical properties in developing tissues.* Nat Methods, 2017. **14**(2): p. 181-186.

21.  Somendra, M.B. and S. Flavio, *A measure of data collapse for scaling.* Journal of Physics A: Mathematical and General, 2001. **34**(33): p. 6375.

22.  Bi, D., et al., *Energy barriers and cell migration in densely packed tissues.* Soft Matter, 2014. **10**(12): p. 1885-90.

23.  McQuarrie, D.A., *Statistical Mechanics.* 1976, New York: Harper & Row.

24.  Mombach, J.C.M., et al., *Rounding of aggregates of biological cells: Experiments and simulations.* Physica A: Statistical Mechanics and its Applications, 2005. **352**(534).

25.  Maitre, J.L., et al., *Adhesion functions in cell sorting by mechanically coupling the cortices of adhering cells.* Science, 2012. **338**(6104): p. 253-6.

26.  Pawlizak, S., et al., *Testing the differential adhesion hypothesis across the epithelial−mesenchymal transition.* New Journal of Physics, 2015. **17**(8): p. 083049.

27.  Sahu, P., J.M. Schwarz, and M.L. Manning, *Geometric signatures of tissue surface tension in a three-dimensional model of confluent tissue.* New Journal of Physics, 2021. **23**(9): p. 093043.

28.  Sahu, P., et al., *Small-scale demixing in confluent biological tissues.* Soft Matter, 2020. **16**(13): p. 3325-3337.

29.  Press, W.H., et al., *Numerical Recipes: The Art of Scientific Computing.* 2020: Cambridge University Press.

30.  Blanpain, C., et al., *Canonical notch signaling functions as a commitment switch in the epidermal lineage.* Genes Dev, 2006. **20**(21): p. 3022-35.

31.  Efremova, M., et al., *CellPhoneDB: inferring cell-cell communication from combined expression of multi-subunit ligand-receptor complexes.* Nat Protoc, 2020. **15**(4): p. 1484-1506.

32.  Giesen, C., et al., *Highly multiplexed imaging of tumor tissues with subcellular resolution by mass cytometry.* Nat Methods, 2014. **11**(4): p. 417-22.

33.  Rustarazo-Calvo, L., et al., *Adhesion-driven tissue rigidification triggers epithelial cell polarity.* bioRxiv, 2025: p. 2025.03.18.644006.

34.  Liu, Z., et al., *Notch1 loss of heterozygosity causes vascular tumors and lethal hemorrhage in mice.* J Clin Invest, 2011. **121**(2): p. 800-8.

35.  Hafner, M., et al., *Keratin 14 Cre transgenic mice authenticate keratin 14 as an oocyte-expressed protein.* Genesis, 2004. **38**(4): p. 176-81.

36.  Tanigaki, K., et al., *Notch-RBP-J signaling is involved in cell fate determination of marginal zone B cells.* Nat Immunol, 2002. **3**(5): p. 443-50.

37.  Lin, L., et al., *Unbiased transcription factor CRISPR screen identifies ZNF800 as master repressor of enteroendocrine differentiation.* Science, 2023. **382**(6669): p. 451-458.

38.  Miyake, S., et al., *Cells degrade a novel inhibitor of differentiation with E1A-like properties upon exiting the cell cycle.* Mol Cell Biol, 2000. **20**(23): p. 8889-902.

39.  Witte, S., et al., *High-density P300 enhancers control cell state transitions.* BMC Genomics, 2015. **16**: p. 903.

40.  Williams, S.E., et al., *Par3-mInsc and Galphai3 cooperate to promote oriented epidermal cell divisions through LGN.* Nat Cell Biol, 2014. **16**(8): p. 758-69.

41.  Khait, I., et al., *Quantitative Analysis of Delta-like 1 Membrane Dynamics Elucidates the Role of Contact Geometry on Notch Signaling.* Cell Rep, 2016. **14**(2): p. 225-33.

42.    Shaya, O., et al., *Cell-Cell Contact Area Affects Notch Signaling and Notch-*
*Dependent Patterning.* Dev Cell, 2017. **40**(5): p. 505-511 e6.
43.    Sarate, R.M., et al., *Dynamic regulation of tissue fluidity controls skin repair during*
*wound healing.* Cell, 2024. **187**(19): p. 5298-5315.e19.
44.    Hogg, S.J., et al., *Targeting histone acetylation dynamics and oncogenic transcription*
*by catalytic P300/CBP inhibition.* Mol Cell, 2021. **81**(10): p. 2183-2200 e13.
45.    Weinert, B.T., et al., *Time-Resolved Analysis Reveals Rapid Dynamics and Broad*
*Scope of the CBP/p300 Acetylome.* Cell, 2018. **174**(1): p. 231-244 e12.
46.    Pylvanainen, J.W., et al., *Fast4DReg - fast registration of 4D microscopy datasets.* J
Cell Sci, 2023. **136**(4).
47.    Pachitariu, M., M. Rariden, and C. Stringer, *Cellpose-SAM: superhuman*
*generalization for cellular segmentation.* bioRxiv, 2025: p. 2025.04.28.651001.
48.    Shen, L., H. Farid, and M.A. McPeek, *Modeling three-dimensional morphological*
*structures using spherical harmonics.* Evolution, 2009. **63**(4): p. 1003-16.
49.    Biggs, L.C., et al., *Mechanical Forces in the Skin: Roles in Tissue Architecture,*
*Stability, and Function.* J Invest Dermatol, 2020. **140**(2): p. 284-290.
50.    Dahmann, C., A.C. Oates, and M. Brand, *Boundary formation and maintenance in*
*tissue development.* Nat Rev Genet, 2011. **12**(1): p. 43-55.
51.    Foty, R.A. and M.S. Steinberg, *The differential adhesion hypothesis: a direct*
*evaluation.* Dev Biol, 2005. **278**(1): p. 255-63.
52.    Harris, T.J. and U. Tepass, *Adherens junctions: from molecules to morphogenesis.*
Nat Rev Mol Cell Biol, 2010. **11**(7): p. 502-14.
53.    Joost, S., et al., *Single-Cell Transcriptomics Reveals that Differentiation and Spatial*
*Signatures Shape Epidermal and Hair Follicle Heterogeneity.* Cell Syst, 2016. **3**(3):
p. 221-237 e9.
54.    Krieg, M., et al., *Tensile forces govern germ-layer organization in zebrafish.* Nat Cell
Biol, 2008. **10**(4): p. 429-36.
55.    Monier, B., et al., *An actomyosin-based barrier inhibits cell mixing at compartmental*
*boundaries in Drosophila embryos.* Nat Cell Biol, 2010. **12**(1): p. 60-9.
56.    Rubsam, M., et al., *Adherens Junctions and Desmosomes Coordinate Mechanics and*
*Signaling to Orchestrate Tissue Morphogenesis and Function: An Evolutionary*
*Perspective.* Cold Spring Harb Perspect Biol, 2018. **10**(11).
57.    Wickstrom, S.A. and C.M. Niessen, *Cell adhesion and mechanics as drivers of tissue*
*organization and differentiation: local cues for large scale organization.* Curr Opin
Cell Biol, 2018. **54**: p. 89-97.
58.    Watt, F.M., *Role of integrins in regulating epidermal adhesion, growth and*
*differentiation.* The EMBO Journal, 2002. **21**(15): p. 3919-3926.
59.    Sugimoto, M., et al., *Universal Surface Biotinylation: a simple, versatile and cost-*
*effective sample multiplexing method for single-cell RNA-seq analysis.* DNA Res,
2022. **29**(3).
60.    Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013.
**29**(1): p. 15-21.
61.    Wolf, F.A., P. Angerer, and F.J. Theis, *SCANPY: large-scale single-cell gene*
*expression data analysis.* Genome Biol, 2018. **19**(1): p. 15.
62.    Lun, A.T., K. Bach, and J.C. Marioni, *Pooling across cells to normalize single-cell*
*RNA sequencing data with many zero counts.* Genome Biol, 2016. **17**: p. 75.
63.    Hunter, J.D., *Matplotlib: A 2D Graphics Environment.* Computing in Science &
Engineering, 2007. **9**(3): p. 90-95.
64.    Tirosh, I., et al., *Dissecting the multicellular ecosystem of metastatic melanoma by*
*single-cell RNA-seq.* Science, 2016. **352**(6282): p. 189-96.
65.    McInnes, L.H., J.; Melville, J., *UMAP: Uniform Manifold Approximation and*
*Projection for Dimension Reduction.* https://arxiv.org/abs/1802.03426}, , 2020.

66.    Traag, V.A., L. Waltman, and N.J. van Eck, *From Louvain to Leiden: guaranteeing*
*well-connected communities.* Sci Rep, 2019. **9**(1): p. 5233.
67.    Wolock, S.L., R. Lopez, and A.M. Klein, *Scrublet: Computational Identification of*
*Cell Doublets in Single-Cell Transcriptomic Data.* Cell Systems, 2019. **8**(4): p. 281-
291.e9.
68.    Korsunsky, I., et al., *Fast, sensitive and accurate integration of single-cell data with*
*Harmony.* Nat Methods, 2019. **16**(12): p. 1289-1296.
69.    Squair, J.W., et al., *Confronting false discoveries in single-cell differential expression.*
Nat Commun, 2021. **12**(1): p. 5692.
70.    Fang, Z., X. Liu, and G. Peltz, *GSEApy: a comprehensive package for performing*
*gene set enrichment analysis in Python.* Bioinformatics, 2023. **39**(1).
1038

1039    **Main Figures**

**Figure 1. Basal cells employ distinct embryonic stage-specific strategies to generate suprabasal layers**

**A.** Schematics of the *ex vivo* embryo imaging setup. **B.** Quantification of parallel (left panel), basal (middle panel) and suprabasal (right panel) cell divisions in E14.5 and E15.5 embryos (mean ±SD; n=4 (E14.5) /5 (E15.5) embryos; Student's t-test (left and middle panel), Mann-Whitney (right panel)). **C.** Representative images and quantification of CyclinB-positive undifferentiated cells (K10-neg) in the suprabasal layers of E14.5 and E15.5 embryos (30 μm; mean ±SD; n=4 embryos/stage; Mann-Whitney). **D.** Representative snapshots and 3D rendering (lower panel) from live imaging movies of E14.5 epidermis from nTomato reporter mice. Red asterisk marks cell departing from basal layer; double red asterisk marks daughter of perpendicular division that subsequently moves suprabasally, blue asterisks marks appearance of cells in the suprabasal layer, red dotted circle marks completed delamination. Images representative of 3 embryos, scale bars 20 μm. **E.** Representative snapshots and 3D rendering (lower panel) from live imaging movies of E15.5 epidermis from nTomato reporter mice. Red asterisk marks cell departing from basal layer; blue asterisk marks appearance of this cell in the suprabasal layer, red dotted circle marks completed delamination. Images representative of 3 embryos, scale bars 20 μm. **F.** Quantification of nuclear oblateness (roundness; upper panel) and asymmetry of deformation (wedge-shape; lower panel) of delaminating cells (Del.) and their immediate neighbors (Neigh.) from 3D renderings. Note that while at E14.5 delaminating cells round up, at E15.5 they elongate in an asymmetric manner (min-to-max box plots; n=3 embryos/stage). **G.** Representative images and quantification of tissue sections from E14.5 - 16.5 embryos stained against K10, Dsg1 and DAPI. Note emergence of K10-Dsg1- positive cells at 15.5 (red asterisks). Dotted line marks basement membrane (scale bars 20 μm; mean ±SD; n=4 embryos; Two-way ANOVA/Tukey's). **H.** Representative images and quantification of tissue sections from E15.5 embryos stained against K10, Dsg1 and DAPI. Note correlation of wedge-like shape and apical Dsg1 intensity. White asterisks mark wedge-like cells with high apical Dsg1, blue asterisks mark cuboidal neighboring cell (scale bars 10 μm; mean ±SD; n=87 cells pooled across 3 embryos; Simple linear regression; Pearson's correlation). **I, J.** Representative images and quantification of delaminating cells in E15.5 embryo 3D whole mounts stained against integrin α6 (Itga6), K10 and DAPI. Note wedge-like shape and integrin α6-positive basal footprint of the delaminating cell (white asterisks; scale bars 5 μm; min-max box and whiskers; n=37 cells pooled across 3 embryos; Mann-Whitney (left, middle panel) Friedman/Dunn's).

1076

**Figure 2. A biophysical model predicts that in the presence of a mechanical barrier,**

**delamination depends on the mechanical properties of the delaminating cell.**

**A.** Snapshot from 3D vertex model for stratified epithelium geometry, illustrating suprabasal

cells (brown), basal cells (light blue) and basement membrane (dark blue), interfacial

tensions between the apical surface of basal cells and the adjacent suprabasal cell ($\sigma_a$,

magenta), basal cells and basement membrane ($\sigma_b$, green), and cell shapes (inset) with

different 3D shape parameters. **B.** Simulation snapshot including a delaminating cell type,

with potentially different delaminating cell-basal cell lateral interfaces $\sigma_{dl}$, delaminating cell-

basement membrane interfaces ($\sigma_b$, yellow), and delaminating cell-suprabasal cell interfaces

($\sigma_{da}$, red). **C.** Snapshot of simulation showing cell in process of delaminating (cell integrates

fully into the suprabasal layer by end of simulation). **D.** Fraction of cells that delaminate after

a time t = X, as a function of $\sigma_{db}$, for many different values of the other parameters. Data

averaged over an ensemble of n=20 simulations for each simulation condition. **E.**

Delamination fraction collapses as a specific function of vertex model parameters.

1091

**Figure 3. A substantial mechanical energy barrier emerges at around E15.5 impacting delamination.**

**A.** Schematic representation of tissue-scale geometric parameter of surface roughness in the 3D vertex model. **B, C.** Representative experimental whole mount (B) and cross section images (C) of surface roughness measurements (scale bars 10 µm). **D.** Schematic representation of cell-scale geometric parameters. **E.** Representative image of tissue cross section across developmental stages for apical curvature angle (scale bars 5 µm). **F-K** Experimental observables (left panels) and corresponding collapsed simulation data (right panels) measured in the 3D model simulations across mechanical parameter space ($\sigma_a$, $\sigma_b$, $\Delta s$). Corresponding averages of *in vivo* observables converted to simulation units specified as green, orange and blue symbols (experimental N>400 cells (J/K); >25 fields of view (L) >130 cells (I) timepoint pooled from n=3 mice; Kruskal-Wallis/Dunn's; n>85

simulations for each parameter set). **L, M.** Experimentally measured apical angles and

corresponding simulation data as a function of vertex model parameters required by force

balance (experimental N>400 cells; n>85 simulations for each parameter set). **N.** Best-fit

vertex model parameters $\sigma_b$ (green), $\sigma_a$ (pink), $\Delta s$ (blue) to the cell and tissue geometry

observed at each stage; error bars represent uncertainty in the fit (see Supplemental

Methods). **O.** Simulation-predicted delamination rates when delaminating cells alter their

mechanical properties from the background basal cells by $\Delta\sigma_{da,a}$ (apical) and $\Delta\sigma_{db,b}$ (basal).

Rate is the fraction that delaminates in a fixed time window t = 2000 timesteps,

corresponding to $\tau \sim 20$ simulation time units (n>85 simulations per parameter set).

1113

**Figure 4. A committed basal population characterized by Notch activation emerges at E15.5**

**A.** Schematics of the scRNAseq experiment. **B.** UMAP representation of the Leiden clusters of filtered epidermal cells from scRNAseq. Embryonic stages of cells are color coded. **C.** UMAP projections of basal identity and differentiation gene expression across E14.5-E16.5.

Arrowheads mark transition population between basal and suprabasal cells. **D.** Dotplot of

selected genes across E14.5-E16.5 basal cells. Note increased expression of differentiation

genes in basal cells of E15.5 and E16.5 embryos. **E.** Dot plot of receptor-ligand (r+l) analyses

between basal stem cells across E14.5-E16.5. Note increased predicted Jag2-Notch signaling

interaction at E15.5 and E16.5. **F-H.** Representative images (F) and quantification (G, H) of

tissue sections from E14.5- 16.5 embryos stained against Notch intracellular domain (N-ICD)

and K10. Note emergence of N-ICD-K10- positive delaminating basal cells at 15.5. Dotted

line marks basement membrane (mean ±SD; scale bars 20 μm; n=3 embryos/stage; Two-way

ANOVA/Tukey's (G); Mann-Whitney (H)).

1128

**Figure 5. Generation of the mechanical boundary at E15.5 is concomitant with stiffening of the basement membrane and rigidification of the basal layer**

**A.** Dot plot of Gene set enrichment analyses (GSEA) from differentially expressed genes in basal cells of E14.5 versus E15.5/E16.5 embryos from scRNAseq data. **B, C.** Representative images (B) and quantification (C) of tissue sections from E14.5- 16.5 embryos analyzed by mass cytometry. Note increased expression of basement membrane Collagen IV (counterstained with K10) and integrins α6 and β1 (mean ±SD; scale bars 50 μm; n=3 embryos/stage; Two-way ANOVA/Tukey's (Left and middle panel), Kruskal-Wallis/Dunn's (right panel)). **D.** Basement membrane Young's moduli extracted from AFM force indentation experiments (Tukey's box and whiskers; n>200 force curves pooled across 3embryos/stage; Kolmogorov-Smirnov). **E.** Representative images and quantification of skin wholemounts from E14.5 and E15.5 embryos stained for pMLC2 and phalloidin. Note increased pMLC2 intensity at cell junctions and interface between basal and suprabasal layers (arrows) (min-max box and whiskers; scale bars 10 μm; n=5 embryos/stage; Mann-Whitney). **F.** Representative snapshots from live imaging movies and subsequent PIV analyses of velocity magnitudes of E14.5 and E15.5 epidermis from membrane-Tomato (mTomato) reporter mice. Red masks mark excluded hair follicle areas (scale bars 50 μm). **G.** Quantification of 2D cell perimeter, area and circularity from data in (A) (Tukey's box and whiskers; n>2300 cells pooled across 3 embryos/stage; Mann-Whitney). **H.** Quantification of cell densities from data in (A) (mean ±SD; n=4 embryos/stage; Mann-Whitney). **I.** Quantification of velocity magnitudes from data in (A). Note attenuated velocity at E15.5 (mean ±SD; n=4 embryos/stage; Linear Mixed-Effects model.

1151

**Figure 6. Tissue rigidity and cell shape changes drive Notch- delamination and control of tissue crowding**

**A.** Schematic illustration of micropatterning of soft and stiff polyacrylamide (PAA)

hydrogels. **B.** Representative images of primary epidermal progenitors stained with K10

(green) and phalloidin (magenta) cultured on 100-µm-diameter micropatterns on 2.5 kPa
(soft) or 8 kPa (stiff) PAA gels. K10 mean intensity was quantified as a function of the
distance from the bottom of the micropatterns. Note the increased K10 mean intensity in the
suprabasal region on stiff substrates. **C.** Quantification of boundary straightness between the
basal layer (bottom) and the first suprabasal layer (upper layer). (Tukey's box and whiskers,
n= 10 (soft); 15 (stiff) patterns pooled across 3 independent experiments; Mann-Whitney). **D.**
Representative snapshots and quantification of live imaging movies from epidermal
progenitor cell monolayers from mTomato/mGFP-Cre reporter mice crossed with Notch1-
ICD-Cre to detect cells that activate Notch (mGFP+). Note increased elongation and local
cell density of cells (white asterisk) that subsequently activate Notch signaling (red asterisk).
Dotted circle marks delamination of Notch-Reporter-positive (NotchRep+) cell (min-max
box and whiskers; n=38 Notch+ cells pooled across 3 independent experiments; Mann-
Whitney; scale bars 10 µm). **E.** Representative images of tissue sections from E14.5-16.5
embryos of control and Rbpj-KO mice stained against K10 (magenta) and Desmoglein 1
(Dsg1; white) (images representative of >3 embryos/stage; scale bars 20 µm). **F.**
Quantification of epidermal thickness from data in (A). Note reduced total and differentiated
layer thickness of Rbpj-KO epidermis at E16.5 (mean ±SD; n=3 embryos/E14.5 and E15.5,
4/E16.5; Mann-Whitney). **G.** Quantification of basal layer thickness and cell height from data
in (E). Note increase basal layer thickness and cell height of Rbpj-KO epidermis emerging at
E15.5 (mean ±SD; n=3 embryos/E14.5 and E15.5, 4/E16.5; Mann-Whitney). **H.**
Quantification of delaminating cells from data in (E). Note increase in delaminating cells in
Rbpj-KO epidermis at E15.5 (mean ±SD; n=3 (E14.5), 5 (E15.5), 4 (E16.5) embryos; Mann-
Whitney). **I.** UMAP representation of the Leiden clusters (left panel) and biological samples
(right panel) of filtered epidermal cells from scRNAseq. **J.** Dotplot of selected genes across
E14.5-E16.5 epidermal cells. Clusters 3 and 6 (in red) represent committed basal cell
populations enriched in Rbpj-KO cells of E15.5 and E16.5 embryos. **K.** Violin plots of
Zfp800 of Eid1 expression across E14.5-E16.5 in control and Rbpj-KO epidermal cells. **L.**
Representative images and H3K27ac nuclear intensity quantification from tissue sections
from E15.5 embryos of control and Rbpj-KO mice stained for H3K27ac (grey), K10
(magenta) and DAPI (cyan) (n= 176 (CNL), 234 (Rbpj-KO) cells pooled across 3 (CNL) /4
(Rbpj-KO) embryos; Student's t-test; Scale bars 20 µm.)

**Supplementary Figures**

1189

**Supplementary Figure 1.**

**A.** Quantification of differentiated (K10-positive) CyclinB1+ cells in the suprabasal layer
(mean ±SD; n=4 embryos /stage; Mann-Whitney) **B.** Representative images and
quantification of delaminating cells in E15.5 embryo from 3D whole mounts stained against
E-cadherin, K10 and DAPI. Delaminating cell is marked with a white asterisk; scale bars 10
μm; min-max box and whiskers; n=38 cells pooled across 3 embryos; Friedman/Dunn's).

1196

**Supplementary Figure 2. Systematic quantification of delamination rates as a function**
**of the properties of the delaminating and stiffness of the surrounding tissue Δ$s$**

**A.** Uncollapsed delamination rates. **B.** Delamination rates plotted as a function of $\sigma_{db} - \sigma_{da}$,
which do not fully collapse, indicating that an additional energetic contribution is required.
Figure 2D in the main text shows that adding the term $c\,\Delta s$ leads to a collapse of the curves
onto a single master curve as a function of $\sigma_{db} - \sigma_{da} + c\,\Delta s$.

1203

**Supplementary Figure 3. Geometric observables for mechanical barrier and quantification of delamination rates and associated energy barriers as function of delaminating cell properties.**

**A-C.** Uncollapsed plots of basal-suprabasal interface roughness, basal cell lateral interface height, and lateral interface orientation, respectively, measured in the 3D model simulations across mechanical parameter space $(\sigma_a, \sigma_b, \Delta s)$. Corresponding collapsed plots are shown in main Fig. 3G, I, K. **D.** Quantification of delamination rates data as function of the delaminating cell tensions $(\sigma_{db}, \sigma_{da})$ parameter space. **E.** Data from (F) collapses as a function of $\sigma_{db} - \sigma_{da}$. **F.** Curve fits to the collapsed delamination rate data shown in panel (B) across the developmental stages. **G.** Quantification of the differences in energy barriers between developmental stages as a function of the differences in delaminating cell properties $(\sigma_{da}, \sigma_{db})$. **H.** Normalized averages (normalized by the energy barrier difference between stages E16 and E15) of the differences in energy barriers between stages E15.5 and E14.5, and between E16.5 and E14.5, plotted as a function of $\sigma_{db} - \sigma_{da}$.

Supplementary Fig. 4

**Supplementary Figure 4.**

**A.** Bar plot showing number of sequenced cells/genotype. **B.** Dot plot of selected key cell

type-specific marker genes across annotated cell types. **C.** UMAP visualization of Leiden

clusters of epidermal cells with main cell states annotated. **D.** Heatmap of marker genes for

epidermal Leiden clusters. **E.** UMAP visualization of Harmony integrated epidermal cells.

Red arrow indicates committed cell population. **F.** Dot plot of selected key cell type-specific

marker genes across annotated cell types from Harmony integrated epidermal cells. Red

dotted rectangle indicates committed cell population.

1227

1228   **Supplementary Figure 5.**

1229   **A.** Representative images and quantifications of tissue sections from E14.5- 16.5 embryos

1230   analyzed by mass cytometry. Note increased expression of integrin β4 (mean ±SD; scale bars

1231   50 µm; n=3 embryos/stage; Two-way ANOVA/Tukey's.

1232

**Supplementary Figure 6.**

**A.** Volcano plot showing differentially expressed genes between Rbpj-KO and CNL epidermis from differentiated cell clusters 5 and 8. **B.** Bar plot showing number of sequenced cells/embryonic stage and genotype across Leiden clusters. **C.** Volcano plot showing differentially expressed genes between E15.5 Rbpj-KO and CNL epidermis from clusters 6 and 0. **D.** Volcano plot showing differentially expressed genes between E16.5 Rbpj-KO and CNL epidermis from cluster 3. **E.** Representative images and quantification of tissue sections from E15.5 embryos of control and Rbpj-KO mice of the expression of Eid1 (magenta) and Keratin-dap (grey) detected by RNA in situ hybridization. Dotted lines mark basal layer (n=4 embryos/condition; Mann-Whitney). Scale bars 20 µm.