FRONT MATTER

Title

•   Full title: **HSPB8 regulates CTP synthase filaments to couple nucleotide metabolism**

and autophagy in tumors

Authors

Wei-Cheng Lin[1†], Chun-Yen Wang[2†], Kuang-Jing Huang[3], Archan Chakraborty[2,7], Yu-Tsun

Lin[1], Ya-Ju Hsieh[1], Kun-Yi Chien[1,2], Po-Yuan Ke[2,3], Wei-Han Huang[2], Chih-Yun Cheng[4], Ian

Yi-Feng Chang[1], Hsiang-Yu Tang[5], Cheng-Hung Yang[5], Mei-Ling Cheng[4,5], Ying-Chih

Chang[8,9,10,11], Chau-Ting Yeh[6], Chien-Kuo Lee[12], Jau-Song Yu[1,2], Yu-Sun Chang[1], Jörg

Großhans[13], and Li-Mei Pai[1,2,3,6] *

Affiliations

[1]Molecular Medicine Research Center, College of Medicine,

[2]Graduate Institute of Biomedical Sciences, College of Medicine,

[3]Department of Biochemistry and Molecular Biology, College of Medicine,

[4]Department of Biomedical Sciences, College of Medicine,

[5]Metabolomics Core Laboratory, Healthy Aging Research Center,

Chang Gung University, Taoyuan 33302, Taiwan.

[6]Liver Research Center, Chang Gung Memorial Hospital, Linkou, Taiwan.

[7]Pharmacology and Cancer Biology, Duke University, NC 27705, USA.

[8]AcroCyte Therapeutics, New Taipei City, Taiwan.

[9]Department of Chemical Engineering, Stanford University, Stanford, CA, 94305, USA.

[10]Developmental Biology and Regenerative Medicine Center, National Taiwan University,

Taipei 10672, Taiwan.

[11]Genomics Research Center, Academia Sinica, Taipei, Taiwan.

[12]Graduate Institute of Immunology, National Taiwan University College of Medicine, Taipei

100233, Taiwan.

[13]Department of Biology, Philipps University Marburg, Germany.

[†]co-first author

[*]Corresponding author Li-Mei Pai, pai@mail.cgu.edu.tw.

Abstract

Metabolic enzyme compartmentalization is increasingly recognized as a regulatory

strategy, yet its functional relevance in cancer remains incompletely understood. Here, we

demonstrate that polymerization of cytidine triphosphate synthase (CTPS) into filaments is

essential for tumor growth in vivo and for three-dimensional proliferation. Glutamine

deprivation, a hallmark of the nutrient-poor core of solid tumors, promotes CTPS filament

assembly, suppressing enzymatic activity and limiting phospholipid synthesis and autophagic

flux. Using proximity labeling, we identify the chaperone HSPB8 as a filament-associated

regulator. HSPB8 antagonizes CTPS filament assembly by facilitating clearance of misfolded

proteins, thereby restoring CTPS activity. Filament disassembly enhances CTP production,

elevates autophagy-related phosphatidylinositol and phosphatidylethanolamine synthesis, and

accelerates autophagic flux. Disruption of CTPS filament formation impairs xenograft tumor

growth. Consistently, analysis of patient datasets reveals that high CTPS and low HSPB8

expression correlate with poor survival across multiple cancers. These findings identify CTPS

filament dynamics as a metabolic vulnerability in tumors.

MAIN TEXT

Introduction

Adapting to chronic nutrient scarcity is a fundamental challenge for malignant

progression. In solid tumors, poor vascularization leads to profound glutamine deprivation,

forcing cancer cells to develop sophisticated survival strategies (*1*). To maintain amino acid

homeostasis, cells utilize a coordinated network of membrane transport, de novo biosynthesis,

and proteostasis—the balance between protein synthesis and autophagic breakdown. Among

these adaptive responses, cells increasingly rely on asparagine (Asn) to maintain proliferation

under glutamine-depleted conditions (*2-4*). Consequently, asparaginase treatment—which

depletes extracellular and intracellular asparagine—is a key therapeutic strategy (*5, 6*).

However, the efficacy of this treatment is often limited by adaptive cellular responses,

including the induction of autophagy (*7-9*).

Emerging evidence suggests that cellular metabolism is regulated through the dynamic

compartmentalization of enzymes into membrane-less organelles or biomolecular condensates.

Cytidine triphosphate synthase (CTPS), the rate-limiting enzyme in de novo pyrimidine

synthesis, is a prominent example of this phenomenon. CTPS catalyzes the ATP-dependent

conversion of UTP to CTP using glutamine as a nitrogen donor. Under glutamine-deprived

conditions, mammalian CTPS undergoes a phase transition to form higher-order polymers and

filament bundles, facilitated by specific residues in the linker (R294) and glutaminase (H355)

domains (*10, 11*). Furthermore, we have previously shown that the amino acid, histidine can

trigger these filamentous structures to assemble along with cytokeratin networks and allow

rapid cell growth after stress relief (*12, 13*). While these filaments are known to respond to

nutrient cues, the molecular machinery that coordinates their assembly with broader cellular

survival programs remains poorly defined.

The maintenance of proteostasis is primarily overseen by molecular chaperones, among

which the small heat shock protein B8 (HSPB8) plays a specialized role. Unlike general

chaperones, HSPB8 specifically recognizes unfolded peptides and defective ribosomal

products (DRiPs), targeting them for degradation via the chaperone-assisted selective

autophagy (CASA) complex (*14-16*). By forming a functional unit with the co-chaperone

BAG3 and the scaffold protein HSP70, HSPB8 has been shown to actively promote

autophagic flux to clear proteotoxic aggregates (*17-19*). While HSPB8 is well-characterized in

the context of neurodegenerative protein aggregation, its involvement in regulating the

structural dynamics of metabolic enzymes has not been explored. Given that metabolic stress often coincides with proteostatic imbalance, HSPB8 emerges as a potential link between the cellular protein quality control machinery and the adaptive compartmentalization of enzymes like CTPS.

In this study, we identify the availability of asparagine as a key environmental factor governing CTPS filament dynamics. Through proteomic screening, we discovered that HSPB8 serves as a critical metabolic rheostat that antagonizes CTPS polymerization. We demonstrate that filament assembly is driven by the accumulation of unfolded peptides, which act as structural scaffolds for filament bundling. Our results reveal that the HSPB8-mediated disassembly of CTPS filaments is a requisite step for activating autophagic flux under nutrient stress. By releasing soluble, active CTPS, cells increase CTP production to fuel the phospholipid biosynthesis required for autophagosome membrane expansion. Crucially, we show that disrupting this balance to reduce CTPS filament formation leads to significantly impaired tumor growth, identifying the HSPB8-CTPS-autophagy axis as a vital survival response. These findings define a therapeutic vulnerability in cancers with high CTPS expression and provide a new perspective on how cells coordinate their physical and chemical states to survive starvation.

Results

CTPS filament assembly is essential for xenograft tumor progression

Given that CTPS filaments have been identified in clinical tumor specimens (*20*), we hypothesized that these macromolecular structures serve a functional role in oncogenesis. To investigate whether CTPS polymerization regulates tumor growth *in vivo*, we performed xenograft experiments using HCT116 cells expressing either wild-type (WT) CTPS or the filament-deficient H355A mutant (Fig. 1A-B). The filament-deficient H355A mutant cells exhibited a significant reduction in tumor volume compared to WT controls (Fig. 1C-E). Notably, these phenotypic differences occurred in the absence of systemic toxicity, as indicated by stable murine body weights across cohorts (Fig. 1F).

To further elucidate the mechanisms driving this growth disparity, we characterized the proliferative capacity of these cells under varying environmental conditions. Under glutamine deprivation in three-dimensional (3D) culture (Gln(-)S(+)), HCT116 cells showed negligible growth. However, in nutrient-replete conditions (Gln(+)S(+)), the H355A mutant displayed significantly impaired proliferation specifically in suspension sphere assays (Fig. S1B). This contrast is striking given that the mutant and WT cells exhibited indistinguishable growth rates in standard two-dimensional (2D) monolayer cultures (Fig. S1A). Collectively, these data suggest that CTPS filament assembly is not merely a cellular feature but a critical regulator of 3D proliferative capacity and *in vivo* tumor formation.

Asparaginase inhibits CTP synthase polymerization by depleting intracellular asparagine pools

Asparagine (Asn) is a known survival factor during glutamine deprivation (*3*). Consistent with this role, we observed that CTP synthase (CTPS) filament formation under glutamine-

deprived conditions was inhibited by asparaginase (ASNase) treatment (Fig. 2, A to F) without
altering total CTPS protein levels (Fig. 2G). Conversely, exogenous Asn supplementation
promoted CTPS polymerization (Fig. 2, C to F). To further define the specificity of this effect,
we screened the remaining non-essential amino acids—alanine, asparagine, aspartate, and
proline—for their capacity to induce filamentation. Our results confirm that Asn induces
CTPS filaments, comparable to the histidine positive control (Fig. S2A). Supporting this, the
combination of Asn and ASNase treatment nearly abolished CTPS filamentation in EBSS, a
conditional medium without any amino acid supplementation (Fig. S2B).

Given that Asn is absent from DMEM, we hypothesized that serum serves as the primary
source of Asn for filament assembly. Quantitative metabolic analysis revealed that Asn levels
increase during glutamine deprivation relative to standard DMEM but are depleted upon
serum withdrawal or ASNase treatment (Fig. S2C). Although extracellular nutrients can be
scavenged via macropinocytosis to support proliferation under nutrient stress (*21*), Asn can
also be synthesized de novo by asparagine synthetase (ASNS). However, ASNS knockdown
did not affect CTPS filamentation (Fig. S2, D and E), likely due to substrate limitations of
glutamine required for ASNS enzymatic activity (*22*). Collectively, these data demonstrate that
intracellular Asn levels regulate CTPS filament dynamics, suggesting a metabolic adaptation
where cells modulate amino acid homeostasis to confer nutrition stress during glutamine
deprivation.

143 Identification of CTPS-associated proteins via APEX2 proximity labeling
To elucidate the mechanisms governing CTP synthase (CTPS) filament dynamics, we
employed APEX2-based proximity labeling to identify protein interactors in situ (*23*).
Although CTPS filaments assemble during nutrient starvation, their structural integrity is
highly sensitive to cell lysis, which typically precludes traditional biochemical purification.
We therefore generated HEp-2 cell lines stably expressing a CTPS-APEX2 fusion protein to
biotinylate proximal proteins in living cells (*13*). To ensure high specificity for filament-
associated partners, we utilized two negative controls representing a non-filamentous state:
glutamine-induced disassembly and a polymerization-deficient R294D mutant (*12, 24*) (Fig.
3A). Western blot analysis confirmed that both endogenous CTPS and exogenous CTPS-
APEX2 levels were markedly elevated under glutamine deprivation, consistent with reports
that filamentation protects CTPS from proteasomal degradation (*12*). Furthermore, the
enrichment of the known filament-associated protein IMPDH2 (*25*) in the CTPS-APEX2 pull-
down validated the spatial accuracy and efficacy of our proximity labeling system (Fig. 3B).

157 Discovery of HSPB8 as a CTPS filament-associated protein regulated by asparaginase
To define the CTPS filament proteome with high stringency, we integrated data from two
independent proteomics experiments and applied a fold-change cutoff based on IMPDH2
enrichment (Fig. S3, A to C). This filtering strategy yielded 184 overlapping proteins
significantly enriched in pathways involving de novo purine biosynthesis and protein
ubiquitination (Fig. 3C). IMPDH2 and PFAS in the purine de novo biosynthesis category have
been demonstrated to form distinct compartments, filament, and purinosome, respectively (*26,*

27). Notably, the identification of multiple heat shock proteins (e.g., HSPA4, HSPB8, HSPH1)

and deubiquitinating enzymes in the protein ubiquitination category suggests that protein

quality control (PQC) machinery localizes proximally to CTPS filaments during metabolic

stress. Given that asparagine (Asn) is a key effector of ATF4-mediated amino acid

homeostasis, we hypothesized that Asn availability modulates this PQC-filament association

(*4, 28*). Quantitative proteomics and subsequent western blotting validation revealed that the

small heat shock protein HSPB8 is downregulated during glutamine deprivation but

specifically upregulated upon ASNase treatment (Fig. 3, D and E). We further confirmed the

physical proximity between CTPS and HSPB8 through a proximity ligation assay (PLA) (Fig.

S3D), establishing HSPB8 as a putative regulator of CTPS filament dynamics under

glutamine-deprived stress.

HSPB8 regulates the compartmentalization of CTPS filaments

HSPB8, a member of the small heat shock protein (sHSP) family, acts in concert with

BAG3—and often within the chaperone-assisted selective autophagy (CASA) complex

alongside HSP70 and E3 ligases—to facilitate the clearance of misfolded or aggregation-prone

protein. Given its established role in maintaining proteostasis, we hypothesized that the

ASNase-induced upregulation of HSPB8 promotes the disassembly of CTPS filaments. To test

this, we performed loss- and gain-of-function assays to evaluate the requirement of HSPB8 for

CTPS filament dynamics. Under glutamine-deprived conditions, HSPB8 knockdown

significantly enhanced CTPS filament assembly compared to control groups (Fig. 4, A and B).

Conversely, we observed that the restoration of glutamine, which typically triggers filament

disassembly, was less effective in the absence of HSPB8, resulting in persistent filament

structures (Fig. 4, A and B).

In contrast to the knockdown phenotypes, overexpression of HSPB8 markedly reduced

the prevalence of CTPS filaments assembly, suggesting that HSPB8 facilitates the release of

CTPS from filamentous compartments into the cytosol (Fig. 4, C and D). Crucially, we found

that HSPB8 depletion was sufficient to restore CTPS filament assembly even in the presence

of ASNase treatment, which normally prevents filament assembly (Fig. 4, E and F).

Collectively, these results demonstrate that ASNase disassembles CTPS filaments by

upregulating HSPB8, identifying this chaperone as a key molecular rheostat for CTPS

assembly during glutamine-deprived stress.

Clinical significance of the HSPB8-CTPS axis in cancer prognosis

Although CTPS expression and enzymatic activity have been established as drivers of

malignant progression (*29*), the clinical relevance of its regulatory partnership with HSPB8

remains poorly defined. To investigate this, we analyzed expression profiles across several

cancer types using The Cancer Genome Atlas (TCGA) database. Our analysis revealed that

CTPS expression is significantly elevated, while HSPB8 expression is markedly reduced, in

multiple malignancies compared to adjacent normal tissues, including breast invasive

carcinoma (BRCA), colon adenocarcinoma (COAD), head and neck squamous cell carcinoma

(HNSCC), lung adenocarcinoma (LUAD), lung squamous cell carcinoma (LUSC), rectum

adenocarcinoma (READ), stomach adenocarcinoma (STAD), and thyroid carcinoma (THCA)

(Fig. S4, A and B).

To evaluate the prognostic impact of this inverse correlation, we performed Kaplan-Meier

survival analysis. We found that patients with high CTPS expression combined with low

HSPB8 expression exhibited significantly poorer survival outcomes in adrenocortical

carcinoma (ACC), pancreatic adenocarcinoma (PAAD), sarcoma (SARC), and

pheochromocytoma and paraganglioma (PCPG) (Fig. S4C). These clinical data suggest that an

environment characterized by low HSPB8 levels—which we have shown promotes CTPS

filament assembly—may drive tumor progression and serves as a negative prognostic

signature in several human cancers.

Glutamine deprivation-induced misfolded protein accumulation drive CTPS

filamentation

Since HSPB8 targets misfolded proteins for autophagic clearance and our proteomic data

identified protein quality control (PQC) machinery in the CTPS interactome (Fig. 3C), we

hypothesized that the accumulation of unfolded proteins modulates CTPS filament dynamics.

Proximity ligation assays (PLA) confirmed that the CTPS-HSPB8 association is spatially

restricted to CTPS filaments (Fig. S5A). Furthermore, we observed an increased association

between HSPB8 and its co-chaperone BAG3 under glutamine deprivation (Fig. S5B),

suggesting an active role for the HSPB8-BAG3 complex in managing protein misfolding

during metabolic stress.

To evaluate the direct impact of proteotoxic stress on CTPS polymerization, we treated

cells with puromycin to induce the premature termination of translation and the subsequent

generation of defective ribosomal products (DRiPs) (*30*). Under glutamine-deprived

conditions, puromycin treatment resulted in the formation of significantly thicker CTPS

filaments compared to untreated controls (Fig. 5, A and B). The proximity biotin-labeling with

streptavidin pull-down experiment revealed that puromycylated polypeptides are enriched in

the immediate vicinity of CTPS filaments (Fig. 5C), implying that local proteotoxic stress

exerts a stabilizing effect on filament dynamics.

Consistent with its role in protein clearance, the knockdown of HSPB8 led to a marked

accumulation of puromycylated polypeptides (Fig. 5D and Fig. S5C). Conversely, the intensity

of these misfolded protein signals significantly declined upon ASNase treatment (Fig. 5E).

Taken together, these findings demonstrate that HSPB8 regulates CTPS filament dynamics by

modulating the local concentration of unfolded proteins, positioning the filament as a

metabolic structure sensitive to cellular proteostasis.

CTPS filament disassembly promotes autophagic flux under glutamine-deprived stress

Autophagy is a critical survival mechanism during asparaginase-induced cytotoxicity.

Given that CTP is a requisite substrate for the de novo synthesis of choline phospholipids—

catalyzed by PCYT1A (phosphate cytidylyltransferase 1, choline, alpha) to replenish

autophagosome membranes (*31*)—we investigated whether the disassembly of CTPS

filaments enhances autophagic capacity. We previously established that non-filamentous CTPS

possesses higher enzymatic activity than its polymerized counterpart. Consistent with this, our in vivo enzymatic assays, measuring the $^{13}C^{15}N$-CTP to $^{13}C^{15}N$-UTP ratio, demonstrated that filament-null mutants (R294D and H355A) exhibit significantly higher catalytic activity compared to wild-type CTPS (Fig. S6A).

To determine if increased CTPS activity promotes autophagy, we utilized mRFP-GFP-LC3 reporter cell lines to monitor autophagic flux (*32*). Under glutamine deprivation, CTPS^{R294D} or CTPS^{H355A} cells displayed a higher density of mRFP-positive/GFP-negative puncta per cell than CTPS^{WT} cells, indicating an acceleration of autolysosome formation (Fig. 6, A and B; Fig. S6, B and C). Transmission electron microscopy (TEM) further confirmed a marked accumulation of both early (AVi) and late (AVd) autophagic vacuoles in filament-null mutants, whereas wild-type cells exhibited significantly fewer structures (Fig. 6, C and D; Fig. S6, D and E). To distinguish between increased induction and blocked degradation, we treated cells with Bafilomycin A1 (BafA1) to inhibit lysosomal fusion. Under BafA1 treatment, filament-null cells accumulated significantly higher levels of LC3-II compared to wild-type controls, confirming that CTPS disassembly actively enhances autophagic flux (Fig. 6, E and F). Collectively, these findings suggest that the metabolic regulation of CTPS filament dynamics modulates autophagic activity.

HSPB8 promotes autophagy by antagonizing CTPS filament assembly

Building on findings identifying the chaperone HSPB8 as a negative regulator of CTP synthase (CTPS) polymerization, we investigated the role of CTPS filament dynamics within the HSPB8-autophagy axis during glutamine-deprived nutrient stress. HSPB8, a member of the small heat shock protein family, specifically binds unfolded peptides and has been previously implicated in the induction of autophagy (*33*). To determine if CTPS filament disassembly is a requisite step in this process, we evaluated autophagic flux under glutamine- and serum-deprived conditions (Gln(-)S(-))In cells expressing wild-type CTPS (CTPS^{WT}), overexpression of HSPB8 significantly increased the number of autolysosomes (mRFP-positive/GFP-negative) and total autophagic vesicles compared to vehicle-treated controls (Fig. 7, A, C, and E). In contrast, HSPB8 overexpression failed to further enhance autophagic vesicle accumulation in filament-deficient CTPS^{H355A} cells (Fig. 7, B, D, and E). These results suggest that the pro-autophagic effect of HSPB8 under glutamine deprivation is mediated through the inhibition of CTPS filament assembly, identifying the transition from polymerized to dimeric or tetrameric CTPS as a critical metabolic switch for autophagic activation.

Reduced volume of CTPS^{H355A} tumors correlates with excessive autophagy activity

Given that CTPS^{H355A} cells exhibit both impaired filament assembly and elevated autophagy in vitro, we sought to determine if this relationship holds true in a physiological tumor model. We analyzed the widely-used autophagy marker LC3B in CTPS^{WT} - and CTPS^{H355A} -derived tumors (from Fig. 1) to understand how defective filament assembly interferes with tumor progression (Fig. S7A). Consistently, the LC3B-II/LC3B-I ratio, a hallmark of autophagic flux, was elevated in CTPS^{H355A} tumors (Fig. S7B). These findings suggest that the anti-tumor effect observed when CTPS filament assembly is disrupted is

284 associated with an induction of excessive autophagic activity.

Phospholipid biosynthesis is upregulated in filament-deficient CTPS mutants

Because CTPS facilitates phospholipid biosynthesis through CTP production, we
hypothesized that filament-deficient mutants exhibit elevated autophagy due to an increased
supply of membrane building blocks. Given that lipids and lipid-associated proteins are
indispensable for autophagosome formation and maturation, we performed liquid
chromatography-mass spectrometry (LC-MS) to compare the lipidomic profiles of CTPS^{WT}
and CTPS^{R294D} HEp-2 cells. We identified 75 annotated lipids that met the significance criteria
of $P < 0.05$, fold change > 1.3, and a Variable Importance in Projection (VIP) score > 1 (Fig.
8A and Fig. S8, A to C). A heatmap of 43 high-confidence lipids which was verified by
MS/MS identification revealed a consistent elevation in CTPS^{R294D} cells, correlating with the
increased enzymatic activity of the non-filamentous mutant (Fig. 8B). LIPEA pathway
analysis indicated a significant enrichment of autophagy-related lipids, specifically
phosphatidylethanolamines (PE) and phosphatidylinositols (PI), which were significantly
higher in CTPS^{R294D} cells (Fig. 8, C and D). Since PI and PE are essential for autophagosome
membrane initiation and expansion, these findings suggest that the metabolic output of
disassembled CTPS directly supports the lipid requirements for enhanced autophagic flux.

Discussion

Metabolic enzymes increasingly are recognized for their ability to undergo higher-order
compartmentalization in response to physiological stress, yet the functional consequences of
these transitions remain poorly understood. Our findings demonstrate that the polymerization
of CTP synthase (CTPS) into filamentous structures is a critical regulatory event that dictates
tumor growth and metabolic adaptation. By integrating biochemical analysis, 3D cell culture,
and clinical datasets, we show that CTPS filaments serve as a metabolic switch that balances
enzyme activity with proteostatic demands (**Fig. 1 and Fig. S1**). Specifically, under glutamine
deprivation—a hallmark of the nutrient-poor core of solid tumors (*1*)—the assembly of CTPS
filaments suppresses enzymatic activity, thereby modulating phospholipid availability and
autophagic flux to influence tumor progression.

A key revelation of this study is the identification of the HSPB8-CTPS filament axis as a
novel branch of the cellular protein quality control (PQC) system. Building on our previous
evidence that ubiquitination regulates CTPS polymerization (*34, 35*), we identified a suite of
PQC-related proteins associated with the filament interactome (**Fig. 3**). The chaperone
HSPB8, acting within the HSPB8-BAG3 complex, emerged as a master regulator of these
dynamics. Our data suggest that misfolded proteins, or defective ribosomal products (DRiPs),
localize to and stabilize CTPS filaments. By modulating the clearance of these proteotoxic
species, HSPB8 dictates the architectural state of CTPS. The knockdown of HSPB8 promotes
filament bundling, while its overexpression triggers disassembly (**Fig. 4**). This mirrors the role
of the HSPB8-BAG3-HSP70 complex in regulating the phase transition of stress granules,
suggesting a universal role for HSPB8 in maintaining the fluidity and dynamics of
biomolecular condensates and membrane-less organelles.

The biological significance of this regulation is underscored by our analysis of TCGA datasets, where we observed a pervasive inverse relationship between CTPS and HSPB8 expression across 75% of cancer types (**Fig. S4, A and B**). The "high-CTPS/low-HSPB8" signature—which our model predicts would favor stable filament formation—correlates significantly with reduced patient survival (**Fig. S4C**). This suggests that the stabilization of CTPS filaments may provide a selective advantage in aggressive malignancies, likely by sequestering enzymatic activity until metabolic conditions require its mobilization. Consequently, the molecular machinery governing CTPS assembly represents a promising, yet unexplored, therapeutic vulnerability in cancer.

Mechanistically, we establish that the transition from filamentous to soluble CTPS is a prerequisite for autophagic activation (**Fig. 6 and Fig. S6**). While it is well known that autophagy is essential for surviving starvation, we found that the pro-autophagic effects of HSPB8 are strictly dependent on the filamentous state of CTPS (**Fig. 7**). The disassembly of these structures facilitates CTP production, providing the necessary energy and substrates for phosphatidylcholine biosynthesis, which is required for autophagosome membrane expansion. Our lipidomic profiles confirm that autophagy-essential lipids, such as PI and PE, are upregulated upon filament disassembly (**Fig. 8**). In the context of the tumor microenvironment, the mobilization of CTPS and the resulting surge in autophagy may interfere with tumor growth (**Fig. S7**), as excessive autophagic flux can trigger apoptosis or metabolic exhaustion (*36, 37*).

Finally, our study identifies the structural dynamics of CTPS as a direct downstream target of the cellular PQC response. We propose that the accumulation of unfolded peptides serves as a molecular scaffold that promotes CTPS filament bundling. This process is highly sensitive to the cellular methylation status (*12*), as evidenced by the disassembly of CTPS filaments upon treatment with the methylation inhibitor Adox (**Fig. S2F**). Under metabolic stress and asparaginase treatment, the upregulation of HSPB8 facilitates the clearance of these filament-associated misfolded proteins (**Fig. 2 and Fig. 3E**). This HSPB8-mediated transition, potentially modulated by methylation-dependent signaling (*38*), ensures that cells can reallocate metabolic resources from a sequestered state into active CTP synthesis to meet the demands of autophagic flux. Collectively, these findings position the CTPS filament as a structural bridge between protein quality control and metabolic homeostasis, revealing that targeting the unfolded protein-HSPB8 axis could disrupt the safeguards enabling cancer cell survival (**Fig. 8E**).

Materials and methods

Cells

Human HEp-2 and HeLa cells were cultured in Dulbecco's Modified Eagle's Medium (Gibco) supplemented with 10% fetal bovine serum. Human HCT116 cells were culture in McCoy's 5A Medium (Sigma). All of cells were maintained at 37°C under 5% CO_2, and passage was done every 3~4 days. Cells were wash with 1X PBS twice before any medium change.

3D spheroid culture

HCT116 cells were seeded in the R^{3}CE platform (Acrocyte Therapeutics Inc., New Taipei City, Taiwan) and incubated in McCoy's 5A medium (with 10% FBS), and appropriate amount of fresh medium is replenished every 3-4 days. Spheroid images were captured by Nikon Eclipse-TS100 microscope with LeadView-2000AIO camera and the spheroid sizes were quantified by imageJ. The spheroids at day 21 were fixed by 4% formaldehyde for 1 h 30 mins and immunostained with CTPS, followed with image capturing by Leica Stellaris SP8 confocal microscope.

Xenograft mouse

Fresh HCT116 WT-CTPS and H355A-CTPS are suspended in McCoy's 5A (without FBS) with Matrigel (M5A: Matrigel=1:1) and subcutaneously injected into right dorsal flank of NUDE mice (BALB/cAnN.Cg-Foxn1nu/CrlNarl). From tumor start to be seen, tumor volume is calculated by length and width (tumor volume V = ½ (Length × Width2)) twice a week. Until 28 days after injection, mice are sacrificed and tumors are collected for further experiments.

Statistical Analysis

Experiment data are analyzed by Student's t test or Kaplan-Meier analysis. Results includes at least 3 biological repeats show means and standard deviation (SD) values with P-values: *<0.05, **<0.01, ***<0.001, ****<0.0001. Other statistical details are included in figure legends.

References

1. M. Pan *et al.*, Regional glutamine deficiency in tumours promotes dedifferentiation through inhibition of histone demethylation. *Nat Cell Biol* **18**, 1090-1101 (2016).
2. A. S. Krall, S. Xu, T. G. Graeber, D. Braas, H. R. Christofk, Asparagine promotes cancer cell proliferation through use as an amino acid exchange factor. *Nat Commun* **7**, 11457 (2016).
3. N. N. Pavlova *et al.*, As Extracellular Glutamine Levels Decline, Asparagine Becomes an Essential Amino Acid. *Cell Metab* **27**, 428-438 e425 (2018).
4. J. Ye *et al.*, The GCN2-ATF4 pathway is critical for tumour cell survival and proliferation in response to nutrient deprivation. *EMBO J* **29**, 2082-2096 (2010).
5. P. L. Lorenzi, S. Claerhout, G. B. Mills, J. N. Weinstein, A curated census of autophagy-modulating proteins and small molecules: candidate targets for cancer therapy. *Autophagy* **10**, 1316-1326 (2014).
6. L. Willems *et al.*, Inhibiting glutamine uptake represents an attractive new strategy for treating acute myeloid leukemia. *Blood* **122**, 3521-3532 (2013).
7. H. Takahashi *et al.*, Autophagy is required for cell survival under L-asparaginase-induced metabolic stress in acute lymphoblastic leukemia cells. *Oncogene* **36**, 4267-4276 (2017).
8. P. Song *et al.*, The role of autophagy in asparaginase-induced immune suppression of

macrophages. *Cell Death Dis* **8**, e2721 (2017).

9.   L. Hinze *et al.*, Synthetic Lethality of Wnt Pathway Activation and Asparaginase in

Drug-Resistant Acute Leukemias. *Cancer Cell* **35**, 664-676 e667 (2019).

10.   E. M. Lynch *et al.*, Human CTP synthase filament structure reveals the active enzyme

conformation. *Nat Struct Mol Biol* **24**, 507-514 (2017).

11.   E. M. Lynch, J. M. Kollman, Coupled structural transitions enable highly cooperative

regulation of human CTPS2 filaments. *Nat Struct Mol Biol* **27**, 42-48 (2020).

12.   W. C. Lin *et al.*, Histidine-Dependent Protein Methylation Is Required for

Compartmentalization of CTP Synthase. *Cell Rep* **24**, 2733-2745 e2737 (2018).

13.   A. Chakraborty *et al.*, SNAP29 mediates the assembly of histidine-induced CTP

synthase filaments in proximity to the cytokeratin network. *J Cell Sci* **133**,   (2020).

14.   S. Carra, S. J. Seguin, H. Lambert, J. Landry, HspB8 chaperone activity toward

poly(Q)-containing proteins depends on its association with Bag3, a stimulator of

macroautophagy. *J Biol Chem* **283**, 1437-1444 (2008).

15.   M. Ganassi *et al.*, A Surveillance Function of the HSPB8-BAG3-HSP70 Chaperone

Complex Ensures Stress Granule Integrity and Dynamism. *Mol Cell* **63**, 796-810

(2016).

16.   F. Li, H. Xiao, Z. Hu, F. Zhou, B. Yang, Exploring the multifaceted roles of heat shock

protein B8 (HSPB8) in diseases. *Eur J Cell Biol* **97**, 216-229 (2018).

17.   S. Carra, S. J. Seguin, H. Lambert, J. Landry, HspB8 chaperone activity toward

poly(Q)-containing proteins depends on its association with Bag3, a stimulator of

macroautophagy. *J Biol Chem* **283**, 1437-1444 (2008).

18.   V. Arndt *et al.*, Chaperone-assisted selective autophagy is essential for muscle

maintenance. *Curr Biol* **20**, 143-148 (2010).

19.   R. Cristofani *et al.*, The Regulation of the Small Heat Shock Protein B8 in Misfolding

Protein Diseases Causing Motoneuronal and Muscle Cell Death. *Front Neurosci* **13**,

796 (2019).

20.   C. C. Chang *et al.*, CTP synthase forms the cytoophidium in human hepatocellular

carcinoma. *Exp Cell Res* **361**, 292-299 (2017).

21.   C. Commisso *et al.*, Macropinocytosis of protein is an amino acid supply route in Ras-

transformed cells. *Nature* **497**, 633-637 (2013).

22.   J. Zhang *et al.*, Asparagine plays a critical role in regulating cellular adaptation to

glutamine depletion. *Mol Cell* **56**, 205-218 (2014).

23.   S. S. Lam *et al.*, Directed evolution of APEX2 for electron microscopy and proximity

labeling. *Nat Methods* **12**, 51-54 (2015).

24.   S. J. Calise *et al.*, Glutamine deprivation initiates reversible assembly of mammalian

rods and rings. *Cell Mol Life Sci* **71**, 2963-2973 (2014).

25.   C. C. Chang, G. D. Keppeke, L. Y. Sung, J. L. Liu, Interfilament interaction between

IMPDH and CTPS cytoophidia. *FEBS J* **285**, 3753-3768 (2018).

26.   Y. Ji, J. Gu, A. M. Makhov, J. D. Griffith, B. S. Mitchell, Regulation of the interaction

of inosine monophosphate dehydrogenase with mycophenolic Acid by GTP. *J Biol*

448   *Chem* **281**, 206-212 (2006).

449 27. S. An, R. Kumar, E. D. Sheets, S. J. Benkovic, Reversible compartmentalization of de

450   novo purine biosynthetic complexes in living cells. *Science* **320**, 103-106 (2008).

451 28. A. Nakamura *et al.*, Inhibition of GCN2 sensitizes ASNS-low cancer cells to

452   asparaginase by disrupting the amino acid response. *Proc Natl Acad Sci U S A* **115**,

453   E7776-E7785 (2018).

454 29. J. C. Williams, H. Kizaki, G. Weber, H. P. Morris, Increased CTP synthetase activity in

455   cancer cells. *Nature* **271**, 71-73 (1978).

456 30. R. Aviner, The science of puromycin: From studies of ribosome function to

457   applications in biotechnology. *Comput Struct Biotechnol J* **18**, 1074-1083 (2020).

458 31. G. Andrejeva *et al.*, De novo phosphatidylcholine synthesis is required for

459   autophagosome membrane formation and maintenance during autophagy. *Autophagy*

460   **16**, 1044-1060 (2020).

461 32. P. Y. Ke, C. W. Chang, Y. C. Hsiao, Baicalein Activates Parkin-Dependent Mitophagy

462   through NDP52 and OPTN. *Cells* **11**, (2022).

463 33. P. Rusmini *et al.*, The Role of the Heat Shock Protein B8 (HSPB8) in Motoneuron

464   Diseases. *Front Mol Neurosci* **10**, 176 (2017).

465 34. P. Y. Wang *et al.*, Regulation of CTP Synthase Filament Formation During DNA

466   Endoreplication in Drosophila. *Genetics* **201**, 1511-1523 (2015).

467 35. L. M. Pai *et al.*, Ubiquitination and filamentous structure of cytidine triphosphate

468   synthase. *Fly (Austin)* **10**, 108-114 (2016).

469 36. S. Liu, S. Yao, H. Yang, S. Liu, Y. Wang, Autophagy: Regulator of cell death. *Cell*

470   *Death Dis* **14**, 648 (2023).

471 37. T. Iba, J. Helms, C. L. Maier, R. Ferrer, J. H. Levy, Autophagy and autophagic cell

472   death in sepsis: friend or foe? *J Intensive Care* **12**, 41 (2024).

473 38. H. C. Chang *et al.*, Asparagine deprivation enhances T cell antitumour response in

474   patients via ROS-mediated metabolic and signal adaptations. *Nat Metab* **7**, 918-927

475   (2025).

476 39. J. Klawitter, V. Schmitz, J. Klawitter, D. Leibfritz, U. Christians, Development and

477   validation of an assay for the quantification of 11 nucleotides using LC/LC-

478   electrospray ionization-MS. *Anal Biochem* **365**, 230-239 (2007).

479 40. L. Dai *et al.*, Phosphoinositide-specific phospholipase Cgamma1 inhibition induces

480   autophagy in human colon cancer and hepatocellular carcinoma cells. *Scientific reports*

481   **7**, 13912 (2017).

483 **Acknowledgements**

484 We thank Chang Gung University for the Metabolomics Core Laboratory for assistance with

485 metabolomics analysis, Proteomics Core Laboratory for assistance with proteomic analysis,

486 Microscopy Center for assistance with image capture, TEM and ultramicrotomy,

487 Bioinformatics Core Laboratory for assistance with TCGA database analysis, respectively. We

488 thank the National RNAi Core Facility at Academia Sinica for providing shRNA reagents and

related services. We thank Dr. Hui-Kuan Lin, Anna C-C Jang and Tzyy-Choou Wu for their

comments on the manuscript. This work was funded by the Ministry of Science and

Technology, Taiwan (MOST 111-2311-B-182-001 to L.-M.P. and MOST 108-2321-B-182-

004-MY3 to W.-C.L.); the Chang Gung Memorial Hospital (CMRPD1K0591-2,

CMRPD1M0171); the "Molecular Medicine Research Center, Chang Gung University" from

The Featured Areas Research Center Program within the framework of the Higher Education

Sprout Project by the Ministry of Education (MOE) in Taiwan.

Author Contributions

Conceptualization: W.C.L. and L.M.P.

Methodology: K.Y.C., M.L.C., P.Y.K., and C.K.L.

Investigation: W.C.L., C.Y.W., K.J.H., A.C., W.H.H., C.Y.C., Y.T.L., Y.J.H., and H.Y.T.

Validation: A.C.

Visualization: W.C.L., C.Y.W., K.J.H., and A.C.

Data curation: W.C.L., C.Y.W., K.Y.C., Y.T.L., C.H.Y., and M.L.C.

Formal analysis: Y.S.C. and Y.F.C.

Funding acquisition: W.C.L. and L.M.P.

Project administration: L.M.P.

Supervision: Y.S.C. and J.S.Y.

Resources: P.Y.K., C.T.Y., Y.C.C., and C.K.L.

Writing – original draft: W.C.L., C.Y.W., A.C., and L.M.P.

Writing – review & editing: Y.S.C., C.K.L., and J.G.

Competing interests:

513 All other authors declare they have no competing interests.

Data and materials availability:

516 Under uploading

Figures

Figure 1. Disrupting CTPS filament assembly impairs HCT116 tumor growth in vivo

(A) HCT116 cells stably expressing CTPS^{WT} or CTPS^{H355A} were cultured in Gln-free DMEM for 24 h, and immunostained for CTPS (green); nuclei were counterstained with DAPI (blue). **(B)** Schematic of the xenograft mouse experiment. Thirty mice were used, with 15 injected with HCT116 CTPS^{WT} cells and 15 injected with HCT116 CTPS^{H355A} cells. The experiment was terminated day 28 after injection. **(C)** Tumor volume measured by external caliper throughout the experimental period. **(D)** Representative images of tumors derived from CTPS^{WT} or CTPS^{H355A} cells excised at day 28 after injection; tumor weights are quantified in **(E)**. **(F)** Mouse body weight recorded by scale over the course of the experiment.

Figure 2. Asparagine availability is essential for CTPS filament assembly under glutamine deprivation

(A, B) HCT116 cells were incubated in Gln-free DMEM or Gln-free DMEM supplemented with ASNase (2U/ml) for 24 h, followed by immunostaining for CTPS (green) and DAPI (blue). **(C, D)** HEp-2 cells were incubated in Gln-free DMEM supplemented with ASNase or Asn (0.34mM) for 24 h, followed by immunostaining for CTPS (green) and DAPI (blue). **(E,**

537   **F)** HeLa cells were incubated with Gln-free DMEM supplemented with ASNase or Asn
538   (0.34mM) for 24 h, followed by immunostaining against CTPS (green) and DAPI (blue).
539   **(G)** HEp-2 cells were incubated in indicated conditions for 24 h, harvested and lysed for
540   Western blotting analysis. Scale bars: 10 mm.
541
542

Figure 3. Proximity labeling identifies the protein quality control factor HSPB8 as a condition-specific interactor of CTPS

(A) Schematic of the CTPS-APEX2 proximity labeling strategy. **(B)** HEp-2 cells expressing CTPS^{WT}-APEX2 were cultured in DMEM, Gln-free DMEM, or Gln-free DMEM supplemented

with Gln for 24 h. HEp-2 cells expressing CTPS^{R294D}-APEX2 were cultured in Gln-free DMEM

for 24 h. Biotinylated proteins were affinity purified using streptavidin-conjugated magnetic

beads and analyzed by Western blotting. **(C)** Venn diagram showing 184 proteins common to

the comparisons of CTPS^{WT}-APEX2 cells cultured in Gln-free DMEM versus Gln-free DMEM

supplemented with Gln, and CTPS^{WT}-APEX2 versus CTPS^{R294D}-APEX2 cells cultured in Gln-

free DMEM. Enriched pathways associated with CTPS-interacting proteins under Gln

deprivation were identified by Ingenuity Pathway Analysis (IPA). **(D)** Heatmap showing

proteins involved in the unbiquitination pathway in cells cultured in DMEM, Gln-free DMEM,

or Gln-free DMEM supplemented with ASNase. **(E)** HSPB8 protein levels in cells cultured in

DMEM, Gln-free DMEM, or Gln-free DMEM supplemented with ASNase.

Figure 4. HSPB8 is required for the disassembly of CTPS filaments during nutrient stress

(A and B) HCT116 cells expressing shLacZ or shHSPB8 were cultured in glutamine-free DMEM for 24 h, with glutamine added for the final 30 min where indicated. Cells were immunostained for CTPS (green) and DAPI (blue). CTPS filament intensity was quantified using ImageJ, with representative filament lengths indicated by pixel values (yellow). HSPB8 protein levels were assessed by Western blotting.

(C and D) HCT116 cells transfected with HA-mock or HA-HSPB8 were cultured in glutamine-free DMEM for 24 h and immunostained for CTPS (green) and DAPI (blue).

(E and F) shLacZ- and shHSPB8-expressing HCT116 cells were cultured in glutamine-free DMEM supplemented with asparaginase (ASNase) for 24 h and immunostained for CTPS (green) and DAPI (blue).

570   Scale bars, 10 μm.

Figure 5. HSPB8-mediated protein quality control is associated with CTPS filament formation

(A and B) HCT116 cells were cultured in Gln-free DMEM for 24 h, with puromycin added during the final 3 h, followed by immunostaining for CTPS (green) and DAPI (blue). CTPS filament intensity was quantified using ImageJ.

(C) HCT116 cells expressing CTPS^{WT}-APEX2 were cultured in Gln-free DMEM for 24 h, with puromycin (10 µg/ml) added during the final 3 h. Proximity labeling was performed, and

biotinylated proteins were analyzed by Western blotting.

(D) shLacZ- and shHSPB8-expressing HEp-2 cells and **(E)** HEp-2 cells were cultured in Gln-

free DMEM for 24 h, with puromycin (20 μg/ml) added during the final 10 min. Cells were

harvested and analyzed by Western blotting for puromycin incorporation, HSPB8, and actin.

Scale bars, 10 μm.

Figure 6. CTPS filament modulates autophagic flux under glutamine deprivation

(A and B) HEp-2 cells stably expressing CTPS^{WT} or CTPS^{R294D} and the tandem GFP–RFP–

LC3 reporter were cultured in Gln-free DMEM for 24 h. GFP- and RFP-positive puncta were

quantified using ImageJ.

(C and D) Transmission electron microscopy images of CTPS^{WT} and CTPS^{R294D} HEp-2 cells cultured in Gln-free DMEM for 24 h. Autophagic vesicles are indicated by yellow arrowheads.
(E and F) Quantification of LC3-II levels in CTPS^{WT} and CTPS^{R294D} HEp-2 cells cultured in Gln-free DMEM for 24 h, with or without bafilomycin A1 (BafA1; 100 nM) added during the final 16 h.

Figure 7. HSPB8 overexpression enhances autophagy inversely correlated with CTPS filament formation

(A–D) Representative images of autophagic vesicles in HCT116 cells expressing the tandem RFP–GFP–LC3 reporter. Red-only puncta indicate autolysosomes, while yellow puncta (RFP/GFP colocalized) mark autophagic structures; nuclei were counterstained with DAPI (blue).

(A′–D′) Corresponding images showing CTPS filament formation by immunostaining for CTPS (green) with DAPI (blue).

604   **(E)** Quantification of autophagic puncta per cell, including red-only, yellow, and total puncta.

605   Numbers below each condition (n) indicate the number of cells analyzed per group.

606   Scale bars, 10 μm.

609   **Figure 8. Metabolomic profiling of phospholipids in cells expressing CTPS^WT versus**

filament-null CTPS mutant

(A) Scheme of the experiment design. HEp-2 cells stably expressing $CTPS^{WT}$ and $CTPS^{H355A}$ were cultured in Gln-free DMEM for 24 h, and subjected to lipid extraction, followed by LC/MS-MS analysis. Three independent experiments were performed and included in the analysis. **(B)** Heatmap showing the relative abundance of 43 identified metabolites across conditions. **(C)** Pathway annotation analysis of the 43 identified metabolites using LIPEA. **(D)** Phosphatidylserine (PS) and Phosphatidylethanolamine (PE) contribution to autophagy pathway. **(E)** Schematic illustrating the HSPB8–CTPS axis reveals a metabolic vulnerability in glutamine-stressed tumors. CTPS filament dynamics is suggested as a metabolic switch. Accumulation of unfolded peptides promotes CTPS filament bundling, a process modulated by cellular methylation. Under glutamine deprivation, reduced HSPB8 levels lead to the accumulation of misfolded peptides, thereby facilitating CTPS filament assembly. In contrast, increased HSPB8 expression, induced by asparaginase (ASNase), promotes CTPS filament disassembly and enhances CTPS enzymatic activity. Elevated CTPS activity supports CTP-dependent phospholipid synthesis, including phosphatidylinositol (PI) and phosphatidylethanolamine (PE), thereby driving autophagosome expansion and suppressing tumor growth. This regulatory axis reveals a metabolic vulnerability in tumors characterized by high CTPS and low HSPB8 expression.

Abbreviations: Me, methylation; ASNase, asparaginase.