Identification of Human Gut Microbiome–Derived Peptides Targeting Biofilm-Specific Lectin Proteins of *Pseudomonas aeruginosa*

Ayush Amod[#1], Ananya Anurag Anand[#1], Sreyanki Chandra[1], Sarfraz Anwar[1], Mubashra[2], Sameer Srivastava[2], Sintu Kumar Samanta*[1]

# denotes equal contribution

[1]Department of Applied Sciences, Indian Institute of Information Technology Allahabad, Prayagraj-211015, India

[2]Department of Biotechnology, Motilal Nehru National Institute of Technology, Prayagraj-211004, India

Corresponding author* :

Dr. Sintu Kumar Samanta

Associate Professor

Department of Applied Sciences

Indian Institute of Information Technology, Allahabad

Uttar Pradesh- 211015 (India)

Email ID: samantasintu@iiita.ac.in

Abstract

The carbohydrate-binding proteins (LecA and LecB) present within the extracellular polymeric substance (EPS) matrix of *Pseudomonas aeruginosa* play an essential role in maintaining the structural integrity of the biofilms through interactions with the EPS polysaccharides. Therefore, targeting the above lectins can turn out to be one of the promising strategies for disrupting *P. aeruginosa* biofilms. In the current study, we investigated the potency of antimicrobial peptides (AMPs) produced by the human gut microbiome in targeting LecA and LecB proteins of *P. aeruginosa*. Initially, a comprehensive *in-silico* pipeline was developed to identify and characterize putative antibacterial and antibiofilm AMPs produced by the human gut microbiome. These AMPs were then subsequently studied for their interaction with the lectin proteins through molecular docking, MM-GBSA, residue analysis, and molecular dynamics (MD) simulation. Among the studied peptides, amp21 and

amp24 exhibited the strongest interactions with the lectin protein, occupying binding sites overlapping with key active-site residues previously reported for raffinose binding. amp6, amp21, and amp24 were selected for *in vitro* validation based on the MD simulation results of both LecA and LecB proteins. The above selected peptides exhibited minimal hemolytic activity across the tested concentration range. amp21 and amp6 were non-toxic to mammalian cells while amp24 demonstrated cytotoxicity only at higher doses. amp21 was found to be the most potent AMP and inhibited the growth of *P. aeruginosa* by ~60% at 50 µg mL□¹. amp6 and amp21 resulted in a significant disruption of *P. aeruginosa* biofilms. Membrane permeabilization assays and scanning electron microscopy revealed that amp6, amp21, and amp24 damaged the bacterial cell membranes apart from compromising the integrity of the biofilm EPS matrix. These findings establish human-gut microbiome-derived AMPs as promising molecules to target *P. aeruginosa* biofilms via inhibition of lectin proteins.

Keywords: Antimicrobial peptides; biofilms; extracellular polymeric substance; lectins; *Pseudomonas aeruginosa*

1. Introduction

Due to strong selection pressure driven by increasing antibiotic use and misuse over the last few years, a large number of bacterial species have evolved to acquire antimicrobial resistance (AMR) to a wide repertoire of antibiotics [Salam et al., 2023]. The World Health Organization (WHO) has identified AMR as one of the top ten global public health concerns, with recent estimates projecting that it can account for nearly 1.91 million deaths annually by 2050 [WHO 2020; Naghavi et al., 2024]. Bacterial biofilms substantially contribute to AMR by impeding antibiotic penetration, facilitating horizontal gene transfer among resident bacteria, and fostering metabolic heterogeneity. As a result, bacterial biofilms show increased tolerance to antibiotic agents, making them much harder to eradicate as compared to free-living planktonic cells [Hall-Stoodley et al., 2004]. Several studies report that the innate ability of ESKAPE pathogens to form biofilms results in broad-spectrum tolerance against conventional antibiotics, rendering current antimicrobial strategies almost ineffective [Sarkar et al., 2024].

Most antibacterial agents, including antibiotics, struggle to penetrate the biofilm's EPS matrix, making biofilm cells 10–1000 times more tolerant to antimicrobials than their planktonic counterparts [Shree et al., 2023]. EPS matrix serves as a reservoir for several antibiotic-chelating enzymes, rendering the antibiotics non-functional. Moreover, the EPS

matrix limits the penetration of antibiotics into the deeper layers of the biofilm [Singh and Amod et al., 2022]. Further, it promotes adhesion to the surface, cell-to-cell adhesion, and aggregation during biofilm formation [Flemming and Wingender, 2010].

The biofilm EPS matrix constitutes nearly 90% of total biofilm mass and is mainly comprised of extracellular DNA (eDNA), proteins, and polysaccharides [Fulaz et al., 2019]. One of the novel strategies to disrupt bacterial biofilms could be targeting structural proteins. Structural proteins, along with polysaccharides, play a major role in the stabilization of biofilm architecture by connecting bacterial cells to the EPS matrix [Fong and Yildiz, 2015]. Among the structural proteins, lectins represent an important subclass that mediate cell–matrix and cell–cell interactions through specific carbohydrate recognition and therefore play an important role in establishing biofilms [Napoleão et al., 2025].

P. aeruginosa is a major threat to clinical settings and plays a detrimental role in a large number of hospital-acquired infections (HAIs) [Moradali et al., 2017; Reynolds and Kollef, 2021]. It tends to cause multi-site infections, of which bacteremia is fatal, with a mortality rate ranging from 18% - 61% [Zhang et al., 2020]. LecA and LecB are the carbohydrate-binding proteins (lectins) present in the EPS matrix of *P. aeruginosa* biofilms that play a major role in maintaining its architecture. LecA shows high binding affinity with galactose and its derivatives [Garber et al. 1992]. In acute infections, LecA mediates the initial interaction with the host cells by acting as an adhesion factor, whereas in the case of chronic infections, it contributes to biofilm formation by *P. aeruginosa* [Chemani et al. 2009]. On the other hand, LecB contributes to biofilm formation by binding to Psl, a key polysaccharide present in the EPS matrix of *P. aeruginosa* biofilms. The above interaction between the LecB protein and Psl polysaccharide leads to increased retention of both bacterial cells and EPS in a developing biofilm**.** Therefore, it is evident that biofilm formation and associated virulence in *P. aeruginosa* can be prevented by inhibiting the function of the above lectins [Wagner et al., 2017].

Specific targeting by small molecules could be a potent strategy in order to disintegrate bacterial biofilms. In this context, researchers have often exploited nanoscale molecules such as nanoparticles (NPs), nanoclusters, and carbon dots (Cdots) [Amod et al., 2025; Verma et al., 2020]. Apart from nanomaterials, antimicrobial peptides (AMPs) are another class of small molecules that can be leveraged to target bacterial biofilms. These are small peptide molecules, less than 50 amino acids in length, that are endogenously produced and released in

almost all living organisms as a part of their innate immune response [Joo et al., 2016]. AMPs have been reported to show their antibacterial activity through several mechanisms, which include disruption of the cell membrane, binding to nucleic acid, among others [Singh et al., 2023]. Therefore, unlike antibiotics, the AMPs have a different set of mechanisms to target the same bacteria, which reduces the chances of bacteria to evade their bactericidal action [Yasir et al., 2018]. Importantly, AMPs show high collateral sensitivity and low cross-resistance, which makes them more promising as antimicrobial agents [Lázár et al. 2019]. They can be used in combination with other antibacterial and antibiofilm agents to achieve enhanced efficacy against bacteria and their biofilms [Anand et al., 2024a]. Additionally, AMPs are reported to possess enormous potential to exhibit prominent antibacterial and antibiofilm activity against different drug-resistant bacteria.

The human gut microbiome is a rich reservoir of AMPs and is involved in several activities such as immunomodulation, disruption of bacterial cell membrane and halting nucleic acid synthesis [Cotter et al., 2013]. As compared to other sources, AMPs tapped from human gut microbiota are expected to show low toxicity and immunogenicity since the commensal gut bacteria producing the AMPs are adapted to coexist within the human gastrointestinal environment over the course of millions of years of evolution [Dominguez-Bello et al., 2019].

Keeping the above fact in mind, in our current study, we first developed a comprehensive *in-silico* pipeline to identify putative antibacterial and antibiofilm AMPs present in the human gut microbiota. Subsequently, we performed molecular docking and molecular dynamics (MD) simulation studies to determine the interaction of these AMPs with the lectin proteins of *P. aeruginosa* (i.e., LecA and LecB). Among the selected peptides, amp6, amp21, and amp24 were found to be the most potent inhibitors of the above lectin proteins and were therefore selected for *in-vitro* validation. The amp21 showed excellent biocompatibility and the most prominent antibacterial activity. Biofilm inhibition and disruption assays showed that amp6 and amp21 brought a greater reduction in the biofilm mass of *P. aeruginosa* when compared to amp24. All three selected AMPs showed significant antibiofilm activity by stimulating the transition of bacterial cells from the biofilm to the free-living planktonic state. It is conceivable that the above-observed phenomenon was due to the interaction of the AMPs with the lectin proteins, which subsequently disturbed the integrity of the EPS matrix. In addition, the AMPs exhibited antibacterial activity by causing perturbations in the bacterial cell membrane.

2. Materials and methods

2.1. Materials

Luria Bertani (LB) broth, Muller Hinton agar (MHA), Dimethyl sulfoxide (DMSO), 99% glacial acetic acid, sodium chloride, magnesium sulphate, L-arginine, potassium phosphate, streptomycin sulphate, and tetracycline hydrochloride were procured from SRL, India. Propidium iodide (PI), trypsin, and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and RPMI 1640 media were purchased from Sigma-Aldrich. Phosphate buffer saline (PBS) and Triton X-100 were bought from Himedia, India. Antibiotic-antimycotic mix and Foetal bovine serum (FBS) were purchased from GIBCO. Glucose was purchased from Merck. *Pseudomonas aeruginosa* PAO1 strain was a kind gift from Prof. SomLata from Jamia Milia Hamdard, New Delhi, India. The HCT116 cancer line was procured from the National Centre for Cell Science (NCCS), Pune. The lead AMPs filtered from the *in-silico* studies were sent for synthesis to the Peptide Purchase Committee, Elabscience, USA. The AMPs were supplied as lyophilized powder and were stored in -20 ℃ post receipt.

2.2. Methods

2.2.1. Evolutionary study of lectin proteins: Before targeting the LecA and LecB proteins of *P. aeruginosa*, their evolutionary lineage was studied through phylogenetic analysis with the objective of (i) estimating the degree of structural conservation of these lectins among other ESKAPE pathogens (ii) determining the presence of their structural homologs in humans, to gain insights into possible cross-reactivity or toxic effects while targeting these lectins.

For this purpose, BlastP (https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE=Proteins) analysis was performed on NCBI using the FASTA sequences of LecA (PDB ID: 6YOH) and LecB (PDB ID: 1OUX) proteins [Camacho et al., 2009]. The number of hits found for LecA was 742, whereas for LecB was 1520. The FASTA sequences of the BLASTP hits were downloaded and imported into the MEGA v11 software [Tamura et al., 2021]. Sequences belonging to the ESKAPE group were selected for further analysis using the following keywords:

'Escherichia coli,' 'Staphylococcus aureus,' 'Klebsiella pneumonia,' 'Acinetobacter baumannii,' 'Pseudomonas,' *'Enterococcus faecium,'* 'Enterobacter,' and 'Enterobacteriaceae.'

Multiple sequence alignment (MSA) was performed using MUSCLE (https://www.ebi.ac.uk/jdispatcher/msa/muscle?stype=protein) [Edgar, 2004]. The phylogenetic tree was constructed using the Maximum-Likelihood method, and the tree was exported in Newick format for visualization. Finally, the phylogenetic tree was visualized and annotated using iTOL software (https://itol.embl.de/) [Letunic and Bork, 2021].

2.2.2. Selection of AMPs for screening: After the evolutionary conservation and host-specificity insights were obtained for LecA and LecB proteins through phylogenetic analysis, the next step involved the rational selection of AMPs for screening to target the above lectin proteins**.** The AMPs used in this study were taken from the study by Anand et al. (2025), where a comprehensive *in-silico* pipeline was developed to identify and characterize putative antibacterial and antibiofilm peptides produced by human gut microbiota [Anand et al., 2025].

2.2.3. Retrieval of target-protein structures: Once the AMPs were selected, the 3-D X-ray crystallographic structures of the target proteins, i.e., LecA and LecB (from *P. aeruginosa*), were downloaded from the PDB database (https://www.rcsb.org/) [Berman, 2000].

Structures of LecA protein with the PDB IDs: 7Z62/7Z63, 5MIH, and 4LK6/4LK7/4LKE either represented different chemotypes or lacked the specific inhibitor context required for docking optimization [Bruneau et al., 2023; Wagner et al., 2017; Kadam et al., 2013]. LecA structure with the PDB ID (6YOH) was selected for the docking studies owing to its high-resolution structure (1.84 Å) and ligand-bound conformation with a relevant glycomimetic inhibitor [Kuhaudomlarp et al., 2020].

On the other hand, structures of LecB protein with the PDB IDs: 5A6X, 5A6Y, 5A6Z, 5A70, and 5MB1reflected different ligand chemotypes and subtle sequence divergences and were therefore not suitable to use in a uniform docking framework [Lepsik et al., 2019; Sommer et al., 2016; Sommer et al., 2018]. LecB protein with the PDB ID 1OUX was selected for the docking studies since it provided a standard reference scaffold with well-characterized fucose/mannose binding fold [Loris et al., 2003].

2.2.4. Three-dimensional modelling of AMPs: The 3-D modelling of AMPs was performed using the PEP-FOLD3.5 software as per the protocol mentioned by Anand et al. (2025) [Anand et al., 2025].

2.2.5. Molecular docking and MM/GBSA analysis: Molecular docking and MM/GBSA analyses were performed between the selected AMPs and the 3D structures of the lectin proteins on HawkDock server (http://cadd.zju.edu.cn/hawkdock/) to evaluate their binding interactions [Weng et al., 2019]. The MM-GBSA binding free energy gives a more accurate assessment of the thermodynamic stability of each complex by considering solvation and entropic effects, whilst the docking score gives an indication of the overall binding affinity and interaction strength between the target protein and the peptides. Lower (more negative values) for both parameters indicates more stable and stronger binding between the peptides and the target protein. Further, in order to determine the contribution of each residue to the total binding free energy, MM/GBSA free energy decomposition analysis and residue analysis were performed [Dalal et al., 2021; Kumari and Dalal, 2022].

2.2.6. MD simulation analysis: MD Simulation was carried out for a duration of 100 ns using the protocol mentioned by Anand et al., 2024 [Anand et al., 2024b].

2.2.7. Analysis of MD Simulation results: GROMACS utilities were employed to obtain the trajectories for performing the MD simulation studies. For this objective, Root mean square deviation (RMSD), radius of gyration (Rg), hydrogen bond interactions, coul-SR and LJ-SR interactions were measured. Free energy landscape (FEL) plots were generated by utilising gmx-sham [Prakash et al., 2018].

2.2.8. Biocompatibility of the lead AMPs: Before studying the antibacterial and antibiofilm activity of the lead AMPs (screened through MD simulation studies), they were tested for their biocompatibility through the hemocompatibility and cytotoxicity assay:

2.2.8.1. Hemocompatibility assay: The hemocompatibility of the AMPs was investigated in accordance with our previously standardized protocol [Singh et al., 2022]. The human red blood cells (RBCs) were mixed with AMPs at varying concentrations between 6.25-200 µg mL^{-1}. Hemolytic activity of AMPs was expressed in percentage relative to the hemolysis

induced by Triton X-100, which was considered 100% that was used as a positive control. PBS served as the negative control. The experiment was performed twice independently (in triplicate), and the results are represented as the mean of both times.

2.2.8.2. Cytotoxicity assay: Post hemocompatibility assessment, the cytotoxic effects of the lead AMPs were further evaluated in mammalian cell lines through MTT dye reduction assay to evaluate their cellular safety profile. In brief, a confluent monolayer of human colorectal cancer (HCT116) cells was trypsinized using (0.25% Trypsin - 0.1% EDTA), following which the cells were harvested by centrifugation. The harvested cells were then seeded in a 96-well plate at a density of 3×10^3 cells per well in 100 μL of complete RPMI1640 media (RPMI1640 supplemented with FBS and antibiotic-antimycotic mix). The cells were allowed to grow at 37 ℃ for 24 h to permit cell attachment and further growth. The medium was then removed and replaced with 100 µL of fresh medium with or without AMPs (at varying concentrations ranging between 6.25 µg mL^{-1} to 100 µg mL^{-1}). After treatment, 10 µL of 5 mg ml^{-1} MTT solution was added to each well and was further incubated for an additional 4 h at 37 ℃. The supernatant was carefully removed, and the formazan crystals formed were dissolved in 100 µL of DMSO. The OD was estimated through a microplate reader at 570 nm, and the viability of HCT116 cells against the peptides was determined in terms of percentage. The viability of the HCT116cells treated with AMPs was expressed in percentage relative to the viability of the untreated cells, which was considered 100%. DMSO was used as the blank. The experiment was performed twice independently (in triplicate), and the results are represented as the mean of both times.

2.2.9. Antibacterial efficacy of AMPs against *P. aeruginosa:* Once the biocompatibility of the lead AMPs was determined, we subsequently investigated their antibacterial potency. Firstly, in accordance with the CLSI guidelines, we first evaluated the MIC of the lead AMPs against *P. aeruginosa* using the standard broth microdilution method, ranging from 0.25 to 64 µg mL^{-1} for 12 h. However, we did not obtain MIC up to 64 µg mL^{-1}.

Therefore, subsequently, we performed a CFU-based viability assay to evaluate concentration-dependent reductions in bacterial load upon treatment with the lead AMPs. Stationary phase cells of *P. aeruginosa* were treated with the AMPs at concentrations of 50 µg mL^{-1} and 100 µg mL^{-1}. These were then incubated at 37 ℃ for a duration of 24 h under static conditions. Before treatment, *P.aeruginosa* culture was standardized to an initial

OD$_{600}$ of 0.002. Subsequently, the untreated and *P. aeruginosa* cells treated with AMPs were diluted 10^{-6} times. 100 µL of dilution of all the samples was then spread uniformly on MHA plates and incubated at 37 °C for 24 h. The CFU mL^{-1} was calculated by the following equation:

$$CFU\ mL^{-1} = (number\ of\ colonies * dilution\ factor)\ /\ volume\ plated\ (mL)$$

In this experiment, the untreated *P. aeruginosa* cells were considered to be the control, and the bacterial load in the treated sample was plotted as % CFU mL^{-1} with respect to the untreated control. The CFU mL^{-1} in the untreated *P. aeruginosa* cells were considered to be 100%. The experiment was performed twice, and the results are represented as the mean of both times.

2.2.10. Measurement of PI fluorescence to assess permeabilization of bacterial cell membrane by AMPs: The fluorescence of propidium iodide (PI) was measured in *P. aeruginosa* treated with lead AMPs to determine whether membrane permeabilization contributed to the antibacterial activity of the AMPs. An increase in PI fluorescence corresponds to increased cell membrane permeability. For this purpose, an overnight culture of *P. aeruginosa* was incubated with lead AMPs at concentrations of 12.5 µg mL^{-1} and 100 µg mL^{-1} for 3 h at 37 °C under shaking conditions. Following incubation, the samples were centrifuged at 8000 rpm for 5 min, washed with 1X PBS, and centrifuged again under identical conditions. The supernatants were discarded, and the cell pellets were resuspended in 1X PBS containing 10 µg mL^{-1} PI. Fluorescence measurements for all the samples were recorded at an excitation wavelength of 488 nm. Herein, the untreated (control) *P. aeruginosa* cells served as the negative control.

2.2.11. Antibiofilm efficacy of AMPs against *P. aeruginosa* biofilms: Following the evaluation of antibacterial activity of the lead AMPs, their antibiofilm potential was examined against *P. aeruginosa* through both biofilm inhibition and biofilm disruption studies:

2.2.11.1 Inhibition of *P. aeruginosa* biofilm by AMPs: For studying the biofilm inhibition efficacy of the lead AMPs, stationary phase cells of *P. aeruginosa* (grown in LB broth) were treated with 50 µg mL^{-1} of all the lead AMPs. These were then incubated in polystyrene-

coated 96-well plates under static conditions at 37 ℃ for 24 h. Herein, the *P. aeruginosa* inoculum was standardized to an initial OD□□□ of 0.02 while setting up the biofilm. The untreated wells were taken as the control. Post treatment, the media was carefully aspirated without disturbing the biofilm and allowed to air-dry for 30 min. To fix the biofilm, 200 μL of methanol was added to each well and incubated at room temperature for 30 min. Following this, the media was removed, and the plate was kept for drying for 2 h. Subsequently, 0.1% (w/v) crystal violet solution was used to stain the biofilms. The staining solution was removed after 10 min of incubation at room temperature. In order to solubilize the bound dye, 30% glacial acetic acid was added, and the 96-well plate was again incubated for 10 min at room temperature. Relative quantification of biofilm mass was performed through a microplate reader by measuring the optical density at 540 nm. The mass of the AMP-treated biofilms was expressed in percentage relative to the biomass of untreated biofilms, which was considered to be 100%. The experiments were performed thrice independently (in triplicate), and the data were plotted as mean ± SD.

2.2.11.2 Disruption of *P. aeruginosa* biofilm by AMPs: For studying the biofilm disruption efficacy of the lead AMPs, mature biofilms were first obtained by incubating stationary phase cells of *P. aeruginosa* (grown in LB broth) in polystyrene-coated 96-well plates under static conditions at 37 ℃ for 48 h. Herein, the *P. aeruginosa* inoculum was standardized to an initial OD□□□ of 0.02 while setting up the biofilm. The above biofilm was then treated with 50 μg mL^{-1} of each of the lead AMPs and was further incubated for 24 h under the same conditions. Post treatment, the biofilm mass was quantitatively estimated in accordance with the standard protocol mentioned in section 2.2.11.1. The experiments were performed thrice independently (in triplicate), and the data were plotted as mean ± SD.

2.2.12. Morphological Analysis of *P. aeruginosa* biofilm treated with AMPs through SEM: Scanning electron microscopy (SEM) was used to observe the effects of lead AMPs on biofilm integrity and other morphological changes in the bacterial cells. For biofilm formation, sterile circular coverslips (18 mm diameter) were placed in the wells of a flat-bottom 6-well polystyrene tissue culture plate, and *P. aeruginosa* biofilms were allowed to develop over them.

Herein, the *P. aeruginosa* inoculum was standardized to an initial OD□□□ of 0.02 while setting up the biofilm. Herein, the 48 h biofilm of *P. aeruginosa* was treated with 100 μg mL^{-}^{1} of amp6 (as a representative) and was further incubated at 37 ℃ for 12 h. Post treatment,

biofilms were fixed with 4% PFA for 5-10 min at room temperature and were then kept overnight at 4 ℃. On the following day, biofilm samples were dehydrated with a graded series (30%, 50%, 70%, 90%, and 100%) of ethanol for 5 min each on a gel rocker. Eventually, the biofilms were coated by a gold sputter coater and examined using a FEI Quanta 250 Scanning Electron Microscope.

2.2.13. Deciphering the mode of mechanism of antibiofilm activity: In order to decipher the mode of mechanism for the antibiofilm activity of the AMPs, mature biofilms were first obtained by growing stationary phase cells of *P. aeruginosa* at 37 ℃ for 48 h in test-tubes under static conditions. Herein, the *P. aeruginosa* inoculum was standardized to an initial OD☐☐☐ of 0.02 while setting up the biofilm. The above biofilm was then treated with 200 µg mL^{-1} of all the lead AMPs and was further incubated for 24 h under the same conditions. Subsequently, the integrity of the biofilm pellicle was visually observed to study the biofilm disruption by AMPs. Post treatment, a CFU assay was performed for all the samples using the planktonic cells present in the media beneath the biofilm pellicle. The planktonic suspensions aliquoted from the media (beneath the biofilm pellicle) were diluted 10^{-6} times. These were then spread uniformly on MHA plates and incubated at 37 ℃ for 24 h. Herein, the untreated *P. aeruginosa* biofilm was considered to be the control. The experiments were performed thrice independently in triplicate.

3.   Results and discussion

3.1.   Homology search and evolutionary study of lectin proteins: The complete BLASTP output for LecA and LecB can be studied from the supplementary information (Supplementary file 1). A comparative sequence and phylogenetic analysis of LecA and LecB was performed to evaluate the evolutionary conservation and target specificity of lectin proteins across clinically relevant pathogens. The initial BlastP analysis confirmed that both LecA and LecB proteins of *P. aeruginosa* have no structural homologs in humans. This implies that targeting these lectin proteins with any antimicrobial molecule will only affect *P. aeruginosa*, thereby minimizing the risk of off-target toxicity in humans and improving their safety for therapeutic uses.

Representative or dummy trees were built such that the positioning of *P. aeruginosa* with respect to other bacteria remains least affected. The same can be visualized in Figure 1a-b.

The phylogenetic tree of LecA revealed that *P. aeruginosa* clusters closely with *K. pneumoniae*, suggesting that these two ESKAPE pathogens share a high level of sequence conservation for the LecA protein, as visible in the representative tree in Figure 1a. Given their close evolutionary relationship, it is possible that the lectin domains involved in adhesion and carbohydrate recognition may remain structurally intact and continue to play comparable functions in host-pathogen interactions. On the other hand, different clades were formed by Enterobacter species (such as *E. cloacae*, *E. hormachei*, and *E. ludwigii*), which suggested that the LecA orthologs of these strains differ from one another. Certain Enterobacter subgroups exhibited modest branch length values (7.1e–7), which reflected minor sequence changes rather than total functional loss. According to this conservation pattern, AMPs made to target LecA may be effective against several pathogens, including *P. aeruginosa* and *K. pneumoniae*, but they may also have varying binding efficiencies in Enterobacter species because of structural or sequence differences.

The phylogenetic tree of LecB showed that it exhibits a high degree of sequence conservation in *P. aeruginosa* and *A. baumannii* (branch length 0.14), which reflected their evolutionary proximity (Figure 1b). Therefore, the AMPs targeting the LecB protein in *P. aeruginosa* are expected to exhibit cross-reactivity against *A. baumannii*, another clinically significant ESKAPE pathogen. On the other hand, *Pseudomonas indica* and *Enterococcus faecium* formed distinct, distantly related branches, which suggested significant sequence divergence and potentially modified carbohydrate-binding motifs. *Pseudomonas fluorescens* and *Enterobacter asburiae* were part of the most diverse cluster; their comparatively long branches (0.74) indicated less evolutionary conservation. The effectiveness of AMP in targeting LecB was found to be pathogen-dependent. It may be strong against *Pseudomonas* and *Acinetobacter*.

Figure 1. (a) Representative image for phylogenetic tree of LecA protein among ESKAPE pathogens **(b)** Representative image for phylogenetic tree of LecA protein among ESKAPE pathogens

3.2. Selection of gut-derived AMPs

Prospective antibacterial and antibiofilm gut-derived peptides identified by Anand et al. (2025) were selected for the study [Anand et al., 2025]. The sequences of the 24 gut-derived AMPs were obtained from the same research group. Consequently, the sequences of the AMPs have been displayed in Table 1.

Table 1. Sequence of gut-derived AMPs

AMP ID	Amino Acid Sequence
amp1	NKIFKACKINKLIETITKKTKIKLCECKNVNN
amp2 (named Amp21 by Anand et al. (2025)) (Anand et al., 2025)	RVIVGHDKVNQLRHVRFFKISTAW
amp3 (named Amp32 by Anand et al. (2025)) (Anand et al., 2025)	KKIIGYIHEAGKGRILFCQ
amp4 (named Amp43 by Anand et al. (2025)) (Anand et al., 2025)	LAWQEVPLFARIIPLADVIDAIAN
amp5	PFTIWGLAGTSLTARIRNHYQKLIKYRKKRGKTIWK

amp6	RSKKLIHFHNVLHLCEVKGMAIKMKNWKTALSRRK
amp7	VLGIKIGDVFGAKFKAKAELAGGIILILMGTKILLEHLGILAF
amp8	SVKSIAWHLYWIQASNSGGPSRPLRRALEEAGIKIVEHFRR
amp9	IKSIPFSCSFHEQGMTGFWKKIPYIKKKEIKKP
amp10	GGFFFNKKNKKLYKKKKKIFFSGRVVADLPGESSSA
amp11	VEDIIKILNNIIKILILHGAVDNCAVLFYRRRLRPARK
amp12	AFKRKNGKALNLDIKTGSNIIYVLNQYKKSRKM
amp13	KKKKKKKKKPIFFFFFLGFFFFFFFFFFYNYKKIIFRARV
amp14	KKKKKKKKKKKKKKKKKKKKKKRKKRRFFFFFFFFFFFLFYIKY
amp15	GVKKGKLFGFLGGNGAGEKFNTFLFLGIFEKRKGNC
amp16 (named Amp164 by Anand et al. (2025)) (Anand et al., 2025)	GEKFGAKDLGVKIVGDLAMPGGGPLRAGKL
amp17 (named Amp175 by Anand et al. (2025)) (Anand et al., 2025)	FNTPSYFAARFKKYIGCLPTEYGKK
amp18 (named Amp186 by Anand et al. (2025)) (Anand et al., 2025)	KEKKKKKKKKNTNIFKNKKKGFFFFFFF
amp19	VCGVAVLCEGGGDIRIAARITELVSALLDLPSNRICVEQRKG
amp20 (named Amp207 by Anand et al. (2025)) (Anand et al., 2025)	KDRRGIKKRKYFFGNEKFEAVA
amp21 (named Amp218 by Anand et al. (2025)) (Anand et al., 2025)	DEVRNKIKENQKEKENETLTRRRLSRQPKL
amp22	LNDKLQPWALALGSHKKRKKIYFFKKKKKKKKKK
amp23 (named	KEVIGERKISYRYRWSIQL

Amp239 by Anand et al. (2025)) (Anand et al., 2025)	
amp24 (named Amp2410 by Anand et al. (2025)) (Anand et al., 2025)	KKKKRKKKKKKRVFFFFFFFFFFFFFFFFF

3.3. 3D modelling of the peptides on PEP-FOLD3: Following the phylogenetic analysis of the lectin proteins, the 3D structures of the AMPs identified through the *in silico* pipeline were modelled using PEP-FOLD3 and visualized in PyMOL. All the peptides showed a predominant α-helical conformation.

3.4. Molecular docking and MM/GBSA of AMPs in complex with LecA and LecB: Among all the AMPs, amp14 showed the best binding profile with LecA protein with a docking score of -4374.23 and an MM-GBSA of -85.03 kcal mol^{-1}. This indicated the formation of a very stable complex with LecA. Following this, energetically favourable interactions were obtained for amp13–LecA (MMGBSA:–60.78 kcal mol^{-1}; docking score: –3743.71), amp22–LecA (MMGBSA: –60.18 kcal mol^{-1}; docking score: –3556.73), and amp24–LecA (MMGBSA: –52.20 kcal mol^{-1}; docking score: –2974.60) complexes. Interestingly, amp8–LecA complex showed a highly negative docking score (–4063.08), indicating significant form complementarity and robust binding at the LecA interface, despite having a relatively higher MM-GBSA (–43.11 kcal mol^{-1}). The binding energy and docking score were also found to be favourably balanced in amp6–LecA (MMGBSA: –49.32 kcal mol^{-1}; docking score:–3982.64) and amp21–LecA (MMGBSA: –48.97 kcal mol^{-1}; docking score: –3820.95) complexes. To further evaluate their structural stability and dynamic behavior under physiological conditions, the seven complexes: amp14–LecA, amp13–LecA, amp22–LecA, amp24–LecA, amp6–LecA, amp21–LecA, and amp8–LecA were selected for molecular dynamics (MD) simulation studies based on thermodynamic stability and interfacial interaction strength. The MM/GBSA and Hawdock docking scores of LecA-AMP complexes have been listed in Table 2.

Table 2. MM/GBSA and Hawkdock docking score of LecA-AMP complexes

Complex	MMGBSA- Binding Free Energy (kcal mol-1)	Hawkdock docking score
amp14-LecA	-85.03	-4374.23
amp13-LecA	-60.78	-3743.71
amp22-LecA	-60.18	-3556.73
amp24-LecA	-52.2	-2974.6
amp20-LecA	-49.66	-2796.48
amp18-LecA	-49.61	-3402.78
amp6-LecA	-49.32	-3982.64
amp21-LecA	-48.97	-3820.95
amp15-LecA	-48.27	-3094.87
amp11-LecA	-47.87	-3230.46
amp5-LecA	-47.74	-2872.6
amp17-LecA	-46.06	-3336.94
amp7-LecA	-44.99	-3436.61
amp19-LecA	-44.68	-3027.59
amp2-LecA	-43.73	-3638.95
amp8-LecA	-43.11	-4063.08
amp9-LecA	-43.11	-3149.36
amp10-LecA	-40.98	-3327.84
amp12-LecA	-40.87	-3531.43
amp23-LecA	-37.65	-2601.07
amp16-LecA	-37.28	-2491.12
amp3-LecA	-35.37	-2842.58
amp4-LecA	-33.82	-3373.51
amp1-LecA	-33.02	-3389.97

With a docking score of -4554.76 and a MM-GBSA binding free energy of -115.42 kcal mol⁻¹, amp14 demonstrated excellent thermodynamic stability on interaction with the LecB protein. The docking scores of amp22–LecB (MMGBSA: –98.64 kcal mol⁻¹; docking score: –5192.46), amp13–LecB (MMGBSA: –92.03 kcal mol⁻¹; –5251.25), and amp11–LecB (–86.27 kcal mol⁻¹; –4863.99) complexes were similarly strong or even better, indicating improved interfacial complementarity and the possibility of successful binding. Furthermore, although having somewhat higher MM-GBSA energies, amp6–LecB (MMGBSA: –73.29 kcal mol⁻¹; docking score: –5881.09) and amp21–LecB (MMGBSA: –59.92 kcal mol⁻¹; docking score: –5691.76) complexes showed unusually negative docking scores, which

indicated strong peptide-lectin interactions. Finally, six complexes, i.e., amp14–LecB, amp22–LecB, amp13–LecB, amp11–LecB, amp6–LecB, and amp21–LecB, were chosen for MD simulations in order to assess their conformational stability, flexibility, and interaction persistence under physiological conditions. This was done after taking into account both energetic stability and binding affinity. The MM/GBSA and Hawdock docking scores of LecB-AMP complexes have been listed in Table 3.

Table 3. MM/GBSA and Hawkdock docking score of LecB-AMP complexes

Complex	MMGBSA- Binding Free Energy (kcal mol-1)	Hawkdock docking score
amp14-LecB	-115.42	-4554.76
amp22-LecB	-98.64	-5192.46
amp13-LecB	-92.03	-5251.25
amp11-LecB	-86.27	-4863.99
amp24-LecB	-84.72	-4896.87
amp18-LecB	-84.3	-4547.95
amp10-LecB	-76.25	-4305.42
amp6-LecB	-73.29	-5881.09
amp2-LecB	-70.6	-4297.64
amp12-LecB	-64.4	-4305.42
amp3-LecB	-63.16	-3666.27
amp9-LecB	-62.71	-3885.85
amp8-LecB	-60.74	-4301.94
amp21-LecB	-59.92	-5691.76
amp20-LecB	-57.28	-3832.56
amp1-LecB	-56.19	-3875.4
amp7-LecB	-56.17	-4301.94
amp16-LecB	-55.85	-3644.46
amp5-LecB	-55.14	-3592.51
amp15-LecB	-50.41	-3667.41
amp23-LecB	-50.37	-3393.23
amp4-LecB	-47.55	-3315.72
amp17-LecB	- 46.09	-3500.11
amp19-LecB	-36.89	-3992.06

3.5.   Protein-peptide MD simulation analysis: To assess their conformational stability and binding behavior of top-ranked AMP–lectin complexes obtained from molecular

docking, MD simulations were carried out. The simulations were evaluated using RMSD, Rg, Coul-SR and LJ-SR interaction energies, H-bonding patterns, and FEL analysis, which provide key insights into the stability and energetically favorable conformations of the complexes.

3.5.1. LecA-peptide complexes: RMSD measures the average deviation in atomic position of a structure (often the backbone or all heavy atoms) from a reference structure over time during MD simulation. Out of all the LecA-peptide complexes, amp21-LecA showed the lowest RMSD. This led to an inference that, with respect to the other AMPs, amp21 remains bounded in a constant orientation at the binding pocket of the LecA protein (Figure 2a). Following this, Rg was measured in order to assess the compactness and folding behaviour of LecA-peptide complexes over the duration of simulation. Herein, amp24-LecA showed a low and constant Rg among all the LecA-peptide complexes. This suggested that (i) binding of the amp24 with LecA led to structural tightening and compaction, (ii) the amp24-LecA complex maintained a comparatively more consistent and compact structure during MD simulation with respect to other complexes (Figure 2b). The short-range electrostatic interaction energy between the protein and peptide is represented by Coul-SR, which measures how strongly their charged or polar residues interact at a typical cutoff distance of ≤1.0–1.2 nm. Among all the LecA-peptide complexes, amp14-LecA showed the most-negative Coul-SR, which implied that the electrostatic interactions between amp14 and the LecA protein are the strongest when compared to other AMPs (Figure 2c). The short-range van der Waals or non-polar interaction energy between the protein and peptide atoms is indicated by LJ-SR, which shows how closely packed residues interact within a typical cutoff distance of ≤1.0–1.2 nm. In our study, it was observed that the amp13-LecA complex showed the lowest LJ-SR values among other LecA-peptide complexes. This observation led to an inference that among all the AMPs, amp13 showed more pronounced hydrophobic /Van der Waals interactions with the LecA protein (Figure 2d). A higher number of H-bonds in protein-peptide simulation indicates greater stability of the protein-peptide complex. The stability, strength, and specificity of the protein–peptide interaction can be assessed by counting the hydrogen bonds during MD simulation, which also provides information on the complex's structural integrity and binding affinity. It was found that amp14 and amp22 formed a greater number of H-bonds with the LecA protein. However, the other peptides formed a more or less similar number of H-bonds with the LecA protein (Figure 2e).

Figure 2. (a) RMSD profile of LecA-peptide complexes **(b)** Rg profile of LecA-peptide complexes, **(c)** Coul-SR profile for LecA-peptide complexes, **(d)** LJ-SR profile for LecA-peptide complexes, **(e)** H-Bonds for LecA-peptide complexes

Further, amp24-LecA and amp21- complexes showed a significant number of conformational states in the blue region or the lower energy state, indicating the stability of these complexes (Figure 3). However, based on overall MD simulation analysis of LecA-peptide complexes, amp24 and amp21 seem to be binding most stably with LecA.

Figure 3. FEL profile of (a) amp6-LecA **(b)** amp8-LecA **(c)** amp13-LecA **(d)** amp14-LecA **(e)** amp21-LecA **(f)** amp22-LecA **(g)** amp24-LecA

3.5.2. LecB-peptide complexes: As compared to other LecB-peptide complexes, amp21-LecB complex and amp6-LecB showed lower RMSD values. This indicated that both amp21 and amp6 maintained a stable and comparatively consistent binding at the interaction site in LecB protein (Figure 4a). The complex amp6-LecB showed the least fluctuations in Rg; significant results were not obtained in Rg plots of other LecB-peptide complexes(Figure 4b). amp14-LecB and amp22-LecB showed the lowest Coul-SR values among all the LecB-peptide complexes (Figure 4c). amp14-LecB and amp13-LecB showed the lowest LJ-SR values (Figure 4d). Among all the AMPs, amp14 formed the maximum number of H-bonds with LecB (Figure 4e).

Figure 4. (a) RMSD profile of LecB-peptide complexes, **(b)** Rg profile of LecB-peptide complexes, **(c)** Coul-SR profile for LecB-peptide complexes, **(d)** LJ-SR profile for LecB-peptide complexes, **(e)** H-Bonds for LecB-peptide complexes

The amp13-LecB, amp22-LecB, and amp21-LecB complexes showed a significant number of conformational states in the blue region or the lower energy state, indicating the stability of these complexes (Figure 5). However, amp13 and amp22 showed very large principal component analysis (PCA) dimensionality, indicating unstable conformations or scattered states of complexes (Figure 5). Only amp21 and amp6 bound LecB complexes show low PCA dimensions (Figure 5). Thus, based on overall MD simulation analysis of LecB-peptide complexes, amp6 and amp21 seem to be binding most stably with LecB.

Figure 5 (a) FEL profile of amp6-LecB **(b)** amp11-LecB **(c)** amp13-LecB **(d)** amp14-LecB **(e)** amp21-LecB **(f)** amp22-LecB

Based on MD simulation analysis results, it was inferred that amp21 and amp24 showed the best binding affinity with LecA protein, as (i) amp21-LecA complex showed minimum RMSD with a good FEL plot, (ii) amp24-LecA complex showed low and constant Rg with a good FEL plot. amp6 and amp21 showed the best binding affinity with LecB protein, as (i) amp21-LecB complex showed minimum RMSD with a good FEL plot, (ii) amp6-LecB complex showed the least fluctuations in Rg with low RMSD and LJ-SR.

3.6. Selection of topmost antibiofilm peptides: As is evident from the MD simulation analysis results, LecA protein formed the most stable complexes with amp21 and amp24, while LecB formed the most stable complexes with amp6 and amp21. These *in silico* findings guided the selection of amp6, amp21, and amp24 for subsequent experimental validation. The above peptides were sent for synthesis to Peptide Purchase Committee, Elabscience, USA, having more than 95% purity. From the chosen peptides, amp21 has been mentioned as AMP218 whereas amp24 has been mentioned as AMP2410 by Anand et al. [Anand et al., 2025].

3.7. MMGBSA residue analysis: amp21 and amp24 were identified as the strongest binders to LecA protein based on MD simulation results. Therefore, their binding residues were analyzed to assess whether they overlap with the known raffinose-binding sites of LecA. Raffinose has been previously documented to inhibit biofilm formation in *P. aeruginosa* by inhibiting the functioning of LecA protein [Kim et al., 2016]

In order to understand the binding of raffinose (antibiofilm agent) to LecA, Kim and coworkers performed *in silico* docking studies between raffinose and X-ray structure of LecA in complex with melibiose. Most of the docked raffinose poses showed common binding patterns in which the galactose moiety of raffinose was projected toward the calcium ion of the LecA active site, while the glucose and fructose moieties of raffinose were oriented toward the surface and the loop region of LecA. The galactose moiety of the raffinose interacted strongly with Tyr 36, His 50, Pro 51, Asp 100, Val 101, and Thr 104, the glucose interacted by hydrogen-bonding with Gln 53, the fructose interacted by hydrogen-bonding with Gly 37 and Asn 107 [Kim et al., 2016].

Since, only amp21 and amp24 were top candidates showing good binding to lecA, we performed a comparative analysis with raffinose binding sites of LecA, and found interesting results. Both amp21 and amp24 were found to interact with all three raffinose binding sites, that is, binding site of raffinose, fructose as well as glucose moiety. The results for same can be visualized in Figure 6, suggesting the possible role of AMPs in biofilm inhibition in a manner similar to raffinose.

Figure 6. Binding of amp21 and amp24 with raffinose-binding residues of LecA

3.8. Study of the hemocompatibility of the AMPs: Before studying the antibacterial activity of amp6, amp21, and amp24, their hemocompatibility was assessed using human RBCs. Upto a concentration of 100 µg mL^{-1}, all three AMPs induced minimal hemolysis and affected a nominal fraction of the total RBC population (Figure 7). At 200 µg mL^{-1}, amp6 and amp21 caused 2.85% and 0.99% hemolysis, respectively (Figure 7). However, amp24 exhibited comparatively higher hemolytic activity at 200 µg mL^{-1} and resulted in 7.5% hemolysis. Overall, these findings demonstrated that amp6, amp21, and amp24 were hemocompatible over a broad range of concentration.

Figure 7. Hemolysis assay of **(a)** amp6 **(b)** amp21 **(c)** amp24

3.9. Cytotoxicity assay of AMPs: Since no hemolytic activity was observed up to 200 μg mL⁻¹ of AMPs, cytotoxicity assay was performed within a concentration range of 6.25- 100 μg mL⁻¹ to avoid unnecessary exposure of the cells to high concentrations of AMPs. Out of the 3 AMPs, amp21 was found to be non-toxic to HCT116 cells across the entire tested concentration range (Figure 8a). Even at 100 μg mL^{-1}, amp21-treated cells exhibited a viability of 105.6% relative to the untreated control (100%). amp6 did not affected the viability of HCT116 cells up to 50 μg mL^{-1}. However, a marginal decrease in the viability was observed in HCT116 cells at 100 μg mL^{-1} of amp6 (Figure 8b). amp24 showed no significant effect on the cell viability up to 25 μg mL^{-1}. However, the viability of HCT116 cells decreased by nearly 20% at 50 μg mL-1 and by 28% at 100 μg mL^{-1} of amp24 (Figure 8c).

Figure 8. Cytotoxicity assay of **(a)** amp6 **(b)** amp21 **(c)** amp24

3.10. Antibacterial efficacy of AMPs against planktonic cells of *P. aeruginosa* through CFU counting assay: Following confirmation of hemocompatibility and acceptable cytotoxicity profiles, the antibacterial efficacy of the selected AMPs was evaluated using the colony-forming unit (CFU) counting assay. amp21 exhibited the strongest antibacterial activity against *P. aeruginosa*, reducing CFU counts by approximately 60% and 84% CFU mL□¹ at concentrations of 50 µg mL□¹ and 100 µg mL□¹, respectively (Figure 9). amp6 showed a comparatively moderate antibacterial activity with CFU reductions of nearly 38% and 64%, respectively, at 50 µg mL□¹ and 100 µg mL□¹, respectively (Figure 9). Interestingly, amp24 showed comparable antibacterial efficacy at both concentrations, resulting in an approximately 54% reduction in CFU counts (Figure 9).

Figure 9. CFU estimation of planktonic cells of *P. aeruginosa* on treatment with amp6,amp21, and amp24 at **(a)** 50 µg mL^{-1} and **(b)** 100 µg mL^{-1} concentration.

3.11. Membrane permeabilization assay through propidium iodide uptake: To elucidate the underlying mechanism for the observed antibacterial activity of the lead AMPs, membrane permeabilization of *P. aeruginosa* cells was evaluated through the PI uptake experiment. On comparison with the untreated cells, *P. aeruginosa* cells treated with 12.5 µg mL^{-1} and 100 µg mL^{-1} of the lead AMPs resulted in a marked increase in PI fluorescence (Figure 10). The above increase in PI fluorescence in AMP-treated cells indicated that all of the above AMPs compromised the integrity of the bacterial cell membrane. The above phenomenon led to an increase in PI uptake by the bacterial cells and subsequent increased binding to intracellular DNA.

Figure 10. Relative fluorescence of PI uptake to assess membrane permeability of *P. aeruginosa* on treatment with amp6, amp21, and amp24 at **(a)** 12.5 µg mL^{-1}, **(b)** and 100 µg mL^{-1} concentration**.** Herein, the untreated *P. aeruginosa* cells were used as the control

3.12. Assessment of antibiofilm assessment of AMPs: The antibiofilm potential of the lead AMPs against *P. aeruginosa* was assessed through biofilm inhibition and biofilm disruption assays after evaluating their antibacterial efficacy:

3.12.1. Inhibition of *P. aeruginosa* biofilm by AMPs: It was found that treatment of *P. aeruginosa* cells with 50 μg mL^{-1} of amp6 and amp21 before biofilm formation significantly inhibited biofilm development, reduced the biofilm biomass to 63.4% and 67.9% of the untreated control, respectively (Figure 11). However, amp24 showed comparatively lower biofilm inhibition, where the biofilm mass was reduced to 80.9% relative to the untreated control under identical conditions (Figure 11).

Figure 11. Quantitative estimation of *P. aeruginosa* biofilm inhibition by 50 μg mL^{-1} of amp6, amp21, and amp24. Herein, the untreated biofilm was used as the control.

3.12.2 Disruption of *P. aeruginosa* biofilm by AMPs: Treatment of preformed *P. aeruginosa* biofilms with 50 μg mL^{-1} of amp6 and amp21 resulted in substantial biofilm disruption, reducing the biofilm biomass to 68.18% and 63.6% of the untreated control, respectively (Figure 12). amp24 demonstrated relatively lower biofilm-disruption efficacy, where the biofilm mass was reduced to 75.75% relative to the control under the same conditions (Figure 12).

Figure 12. Quantitative estimation of *P. aeruginosa* biofilm disruption by 50 µg mL^{-1} of amp6, amp21, and amp24.

3.13. Morphological analysis of *P. aeruginosa* biofilm through SEM: To visually corroborate the biofilm disruption observed in section 3.12.2, we subsequently performed scanning electron microscopy (SEM) analysis. Due to methodological constraints, economic considerations, and observation of similar functional outcomes for all three AMPs in the PI fluorescence assay, SEM analysis was performed only for amp6, which served as a representative AMP. SEM analyses revealed that when compared to the untreated (control) biofilm, *P. aeruginosa* biofilms treated with amp6 showed pronounced structural disruption (Figure 13b). The untreated biofilms appeared largely intact, dense, and well-organised (Figure 13a). On the other hand, a marked reduction was observed in the bacterial cell density of amp6-treated biofilms (Figure 13b). Furthermore, treatment with amp6 not only disrupted the EPS matrix but cause prominent disruption to the bacterial cell membrane (Figure 13b). Similar effects are anticipated as both peptides, like amp6, compromise the integrity of the bacterial cell membrane and show comparable biofilm inhibition and disruptive activity.

Figure 13. SEM image of **(a)** untreated *P. aeruginosa* biofilm, **(b)** *P. aeruginosa* biofilm treated with amp6**.** The damaged bacterial cells are shown by the red arrow

3.14. Deciphering the mode of mechanism for antibiofilm activity of AMPs: Following the quantitative assessment of biofilm disruption and its morphological validation by SEM imaging, we elucidated the underlying mechanism of antibiofilm activity of the selected AMPs. Upon treatment of mature *P. aeruginosa* biofilms with amp6, amp21, and amp24, it was observed that there was a marked increase in the number of bacterial cells migrating from the biofilm pellicle into the underlying media (Figure 14a). The cells released from the biofilm matrix were subsequently quantified through the CFU assay. A substantially higher number of bacterial colonies were detected on Mueller–Hinton agar plates inoculated with AMP-treated cultures (Figure 14b). The above observations suggested that the above three AMPs exerted their antibiofilm activity by disrupting the integrity of the biofilm EPS matrix, which eventually led to the migration of underlying bacteria to the planktonic state. The above phenomenon may be attributed to the interaction of AMPs with lectin proteins that subsequently disturbed the integrity of the biofilm EPS matrix.

Figure 14. (a) Movement of bacterial cell from biofilm pellicle to the surrounding media showing transition from biofilm to the free planktonic cells **(b)** CFU assay to visualize the planktonic cell load in the media surrounding the biofilm on treatment with AMPs**.** Herein, the untreated *P. aeruginosa* cells were used as the control

4. Conclusion and future perspective:

AMPs from the human gut microbiome were computationally identified as potent inhibitors of lectins that are key targets in biofilm formation in *Pseudomonas aeruginosa*. *In vitro* studies showed that the lead AMPs filtered through *in-silico* studies showed strong antibiofilm activity by stimulating the migration of bacterial cells from the sessile biofilm stage to the mobile planktonic stage. Apart from disrupting the biofilm EPS matrix, the above AMPs exhibited prominent antibacterial activity by compromising the integrity of the bacterial cell membrane. Future studies can be done to design shorter derivatives of the above lead AMPs for enhanced antimicrobial efficacy. For practical applications, the efficacy of the above lead AMPs can also be investigated in an animal model.

Supplementary information

All relevant supplementary information has been incorporated in Supplementary file 1

Conflicts of interest

The authors have no relevant financial or non-financial interests to disclose.

Acknowledgment

AA would like to thank ICMR, Govt. of India, for his fellowship. AAA and SC would like to thank the MoE, Govt. of India, for their fellowship. SA would like to thank CSIR, Govt. of India, for his fellowship. We would like to thank IIIT Allahabad for providing the research and the Central Computation Facility (CCF). We would like to thank the Electron Microscopy Unit of CDRI Lucknow for providing the SEM imaging facility.

Funding

The authors would like to thank UP-CST for funding.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the author(s) used ChatGPT to assist with language refinement and improve clarity. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

5.   References

Amod A, Singh S, Lawrence R, Anand A A, Ranjan S, Singh S, Sahoo A K, and Samanta S K 2025. NCCD: a unique combination of MOF-scaffolded stable luminescent copper nanoclusters (Cu NCs) and carbon dots (Cdots) as a potent antimicrobial agent against *Pseudomonas aeruginosa. Nanotechnology* **36** 235102

Anurag Anand A, Amod A, Anwar S, Sahoo A K, Sethi G, and Samanta S K 2024a A comprehensive guide on screening and selection of a suitable AMP against biofilm-forming bacteria *Critical Reviews in Microbiology* **50** 859–78

Anurag Anand A, Anwar S, Yadav V and Samanta S K 2025. A comparative study of computational modeling approaches for evaluating structural dynamics and algorithmic suitability of short-length peptides *Sci Rep* **15** 34351

Anurag Anand A, Sahoo A K and Samanta S K 2024b Exploring the Potential of Designed Peptides Containing Lysine and Arginine Repeats against VIM-2 Metallo-Beta-Lactamases *Int J Pept Res Ther* **30** 41

Berman H M 2000 The Protein Data Bank *Nucleic Acids Research* **28** 235–42

Bruneau A, Gillon E, Furiga A, Brachet E, Alami M, Roques C, Varrot A, Imberty A and Messaoudi S 2023 Discovery of potent 1,1-diarylthiogalactoside glycomimetic inhibitors of Pseudomonas aeruginosa LecA with antibiofilm properties *European Journal of Medicinal Chemistry* **247** 115025

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K and Madden T L 2009 BLAST+: architecture and applications *BMC Bioinformatics* **10** 421

Chemani C, Imberty A, De Bentzmann S, Pierre M, Wimmerová M, Guery B P and Faure K 2009 Role of LecA and LecB Lectins in *Pseudomonas aeruginosa* -Induced Lung Injury and Effect of Carbohydrate Ligands *Infect Immun* **77** 2065–75

Cotter P D, Ross R P and Hill C 2013 Bacteriocins — a viable alternative to antibiotics? *Nat Rev Microbiol* **11** 95–105

Dalal V, Dhankhar P, Singh V, Singh V, Rakhaminov G, Golemi-Kotra D and Kumar P 2021 Structure-Based Identification of Potential Drugs Against FmtA of Staphylococcus aureus: Virtual Screening, Molecular Dynamics, MM-GBSA, and QM/MM *Protein J* **40** 148–65

Dominguez-Bello M G, Godoy-Vitorino F, Knight R and Blaser M J 2019 Role of the microbiome in human development *Gut* **68** 1108–14

Edgar R C 2004 MUSCLE: multiple sequence alignment with high accuracy and high throughput *Nucleic Acids Res* **32** 1792–7

Essmann U, Perera L, Berkowitz M L, Darden T, Lee H and Pedersen L G 1995 A smooth particle mesh Ewald method *The Journal of Chemical Physics* **103** 8577–93

Evans D J and Holian B L 1985 The Nose–Hoover thermostat *The Journal of Chemical Physics* **83** 4069–74

Flemming H-C and Wingender J 2010 The biofilm matrix *Nat Rev Microbiol* **8** 623–33

Fong J N C and Yildiz F H 2015 Biofilm Matrix Proteins ed M Ghannoum, M Parsek, M Whiteley and P Mukherjee *Microbiol Spectr* **3** 3.2.28

Fulaz S, Vitale S, Quinn L and Casey E 2019 Nanoparticle–Biofilm Interactions: The Role of the EPS Matrix *Trends in Microbiology* **27** 915–26

Garber N, Guempel U, Belz A, Gilboa-Garber N and Doyle R J 1992 On the specificity of the d-galactose-binding lectin (PA-I) of Pseudomonas aeruginosa and its strong binding to hydrophobic derivatives of d-galactose and thiogalactose *Biochimica et Biophysica Acta (BBA) - General Subjects* **1116** 331–3

Hall-Stoodley L, Costerton J W and Stoodley P 2004 Bacterial biofilms: from the Natural environment to infectious diseases *Nat Rev Microbiol* **2** 95–108

Joo H-S, Fu C-I and Otto M 2016 Bacterial strategies of resistance to antimicrobial peptides *Phil. Trans. R. Soc. B* **371** 20150292

Kadam R U, Garg D, Schwartz J, Visini R, Sattler M, Stocker A, Darbre T and Reymond J-L 2013 CH–π "T-Shape" Interaction with Histidine Explains Binding of Aromatic Galactosides to *Pseudomonas aeruginosa* Lectin LecA *ACS Chem. Biol.* **8** 1925–30

Kim H-S, Cha E, Kim Y, Jeon Y H, Olson B H, Byun Y and Park H-D 2016 Raffinose, a plant galactoside, inhibits Pseudomonas aeruginosa biofilm formation via binding to LecA and decreasing cellular cyclic diguanylate levels *Sci Rep* **6** 25318

Kuhaudomlarp S, Siebs E, Shanina E, Topin J, Joachim I, Da Silva Figueiredo Celestino Gomes P, Varrot A, Rognan D, Rademacher C, Imberty A and Titz A 2021 Non‑Carbohydrate Glycomimetics as Inhibitors of Calcium(II)‑Binding Lectins *Angew Chem Int Ed* **60** 8104–14

Kumari R and Dalal V 2022 Identification of potential inhibitors for LLM of *Staphylococcus aureus*: structure-based pharmacophore modeling, molecular dynamics, and binding free energy studies *Journal of Biomolecular Structure and Dynamics* **40** 9833–47

Lázár V, Martins A, Spohn R, Daruka L, Grézal G, Fekete G, Számel M, Jangir P K, Kintses B, Csörgő B, Nyerges Á, Györkei Á, Kincses A, Dér A, Walter F R, Deli M A, Urbán E, Hegedűs Z, Olajos G, Méhi O, Bálint B, Nagy I, Martinek T A, Papp B and Pál C 2018 Antibiotic-resistant bacteria show widespread collateral sensitivity to antimicrobial peptides *Nat Microbiol* **3** 718–31

Lepsik M, Sommer R, Kuhaudomlarp S, Lelimousin M, Paci E, Varrot A, Titz A and Imberty A 2019 Induction of rare conformation of oligosaccharide by binding to calcium-dependent bacterial lectin: X-ray crystallography and modelling study *European Journal of Medicinal Chemistry* **177** 212–20

Letunic I and Bork P 2021 Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation *Nucleic Acids Research* **49** W293–6

Loris R, Tielker D, Jaeger K-E and Wyns L 2003 Structural Basis of Carbohydrate Recognition by the Lectin LecB from Pseudomonas aeruginosa *Journal of Molecular Biology* **331** 861–70

Moradali M F, Ghods S and Rehm B H A 2017 Pseudomonas aeruginosa Lifestyle: A Paradigm for Adaptation, Survival, and Persistence Front. Cell. Infect. Microbiol. 7

Naghavi M, Vollset S E, Ikuta K S, Swetschinski L R, Gray A P, Wool E E, Robles Aguilar G, Mestrovic T, Smith G, Han C, Hsu R L, Chalek J, Araki D T, Chung E, Raggi C, Gershberg Hayoon A, Davis Weaver N, Lindstedt P A, Smith A E, Altay U, Bhattacharjee N V, Giannakis K, Fell F, McManigal B, Ekapirat N, Mendes J A, Runghien T, Srimokla O, Abdelkader A, Abd-Elsalam S, Aboagye R G, Abolhassani H, Abualruz H, Abubakar U, Abukhadijah H J, Aburuz S, Abu-Zaid A, Achalapong S, Addo I Y, Adekanmbi V, Adeyeoluwa T E, Adnani Q E S, Adzigbli L A, Afzal M S, Afzal S, Agodi A, Ahlstrom A J, Ahmad A, Ahmad S, Ahmad T, Ahmadi A, Ahmed A, Ahmed H, Ahmed I, Ahmed M, Ahmed S, Ahmed S A, Akkaif M A, Al

Awaidy S, Al Thaher Y, Alalalmeh S O, AlBataineh M T, Aldhaleei W A, Al-Gheethi A A S, Alhaji N B, Ali A, Ali L, Ali S S, Ali W, Allel K, Al-Marwani S, Alrawashdeh A, Altaf A, Al-Tammemi A B, Al-Tawfiq J A, Alzoubi K H, Al-Zyoud W A, Amos B, Amuasi J H, Ancuceanu R, Andrews J R, Anil A, Anuoluwa I A, Anvari S, Anyasodor A E, Apostol G L C, Arabloo J, Arafat M, Aravkin A Y, Areda D, Aremu A, Artamonov A A, Ashley E A, Asika M O, Athari S S, Atout M M W, Awoke T, Azadnajafabad S, et al 2024 Global burden of bacterial antimicrobial resistance 1990–2021: a systematic analysis with forecasts to 2050 *The Lancet* **404** 1199–226

Napoleão T H, Lira T L D S, Pontual E V, Ferreira G R S and Da Silva P M 2025 Lectins as Natural Antibiofilm Agents in the Fight Against Antibiotic Resistance: A Review *Molecules* **30** 3395

Parrinello M and Rahman A 1981 Polymorphic transitions in single crystals: A new molecular dynamics method *Journal of Applied Physics* **52** 7182–90

Prakash A, Kumar V, Meena N K and Lynn A M 2018 Elucidation of the structural stability and dynamics of heterogeneous intermediate ensembles in unfolding pathway of the N-terminal domain of TDP-43 *RSC Adv.* **8** 19835–45

Reynolds D and Kollef M 2021 The Epidemiology and Pathogenesis and Treatment of Pseudomonas aeruginosa Infections: An Update *Drugs* **81** 2117–31

Salam Md A, Al-Amin Md Y, Salam M T, Pawar J S, Akhter N, Rabaan A A and Alqumber M A A 2023 Antimicrobial Resistance: A Growing Serious Threat for Global Public Health *Healthcare* **11** 1946

Sarkar S, Roy A, Mitra R, Kundu S, Banerjee P, Acharya Chowdhury A and Ghosh S 2024 Escaping the ESKAPE pathogens: A review on antibiofilm potential of nanoparticles *Microbial Pathogenesis* **194** 106842

Shree P, Singh C K, Sodhi K K, Surya J N and Singh D K 2023 Biofilms: Understanding the structure and contribution towards bacterial resistance in antibiotics *Medicine in Microecology* **16** 100084

Singh A, Amod A, Mulpuru V, Mishra N, Sahoo A K and Samanta S K 2023 Finding Novel AMPs Secreted from the Human Microbiome as Potent Antibacterial and Antibiofilm Agents and Studying Their Synergistic Activity with Ag NCs *ACS Appl. Bio Mater.* **6** 3674–82

Singh A, Amod A, Pandey P, Bose P, Pingali M S, Shivalkar S, Varadwaj P K, Sahoo A K and Samanta S K 2022 Bacterial biofilm infections, their resistance to antibiotics therapy and current treatment strategies *Biomed. Mater.* **17** 022003

Singh A, Rani K, Tandon V, Sahoo A K and Samanta S K 2022 Ag NCs as a potent antibiofilm agent against pathogenic *Pseudomonas aeruginosa* and *Acinetobacter baumannii* and drug-resistant *Bacillus subtilis* by affecting chemotaxis and flagellar assembly pathway genes *Biomater. Sci.* **10** 6778–90

Sommer R, Wagner S, Rox K, Varrot A, Hauck D, Wamhoff E-C, Schreiber J, Ryckmans T, Brunner T, Rademacher C, Hartmann R W, Brönstrup M, Imberty A and Titz A 2018 Glycomimetic, Orally Bioavailable LecB Inhibitors Block Biofilm Formation of *Pseudomonas aeruginosa J. Am. Chem. Soc.* **140** 2537–45

Sommer R, Wagner S, Varrot A, Nycholat C M, Khaledi A, Häussler S, Paulson J C, Imberty A and Titz A 2016 The virulence factor LecB varies in clinical isolates: consequences for ligand binding and drug discovery *Chem. Sci.* **7** 4990–5001

Tamura K, Stecher G and Kumar S 2021 MEGA11: Molecular Evolutionary Genetics Analysis Version 11 ed F U Battistuzzi *Molecular Biology and Evolution* **38** 3022–7

Verma A, Shivalkar S, Sk M P, Samanta S K and Sahoo A K 2020 Nanocomposite of Ag nanoparticles and catalytic fluorescent carbon dots for synergistic bactericidal activity through enhanced reactive oxygen species generation *Nanotechnology* **31** 405704

Wagner S, Hauck D, Hoffmann M, Sommer R, Joachim I, Müller R, Imberty A, Varrot A and Titz A 2017 Covalent Lectin Inhibition and Application in Bacterial Biofilm Imaging *Angew Chem Int Ed* **56** 16559–64

Weng G, Wang E, Wang Z, Liu H, Zhu F, Li D and Hou T 2019 HawkDock: a web server to predict and analyze the protein-protein complex based on computational docking and MM/GBSA *Nucleic Acids Res* **47** W322–30

World Health Organization. 10 global health issues to track in 2021. 2020. Available from: https://www.who.int/news-room/spotlight/10-global-health-issues-to-track-in-2021 [Accessed June 11, 2023].

Yasir M, Willcox M D P and Dutta D 2018 Action of Antimicrobial Peptides against Bacterial Biofilms *Materials* **11** 2468

Zhang Y, Li Y, Zeng J, Chang Y, Han S, Zhao J, Fan Y, Xiong Z, Zou X, Wang C, Li B, Li H, Han J, Liu X, Xia Y, Lu B and Cao B 2020 Risk Factors for Mortality of In-patients with Pseudomonas aeruginosa Bacteremia in China: Impact of Resistance Profile in the Mortality *Infect Drug Resist* **13** 4115–23