Tracking interlinked microbial and geochemical succession over decades in landfilled
municipal solid waste

Kimber E. Munford[a]#, Daniel S. Grégoire[b], Laura A. Hug[a]

[a]Department of Biology, University of Waterloo, Waterloo, Ontario, Canada

[b]Department of Chemistry, Carleton University, Ottawa, Ontario, Canada

#Address correspondence to Kimber E. Munford, kmunford@uwaterloo.ca.

Abstract

Landfills are heterogeneous built environments embedded in natural freshwater systems. They
pose increasing risks of groundwater contamination from metal-bearing leachates over time. The
interlinked succession of waste decomposition processes, microbial community membership and
metal cycling across a landfill's lifespan have not been explored, reducing our ability to predict
the long-term environmental impacts of landfills. Working with 1,647 metagenome-assembled
genomes from a single landfill, from samples spanning over 39 years of waste decomposition,
we identified changes in landfill biogeochemistry and connected these changes to shifts in
microbial community composition and predicted functions over time. Comparing between Older
(aged 31 – 39 years) and Newer (aged 3 – 20 years) waste cells identified significant shifts in the
availability of labile carbon, redox-associated processes, and concentrations of mobile metals –
all higher in Newer cells. Newer cells were dominated by chemoorganoheterotrophs, while Older
cells contained higher proportions of chemolithoautotrophs and organisms with higher metabolic
versatility. Metal resistance and metal cycling genes were significantly more abundant in Older
cells. Using geochemistry data from time of filling to present and microbial membership across
six landfill cells of different ages, we developed a conceptual model of landfill characteristics
across time. This model connects redox conditions and metal fate, highlighting leachate
recirculation as a key process impacting many geochemical parameters and defining site
chemistry. Our work highlights the substantial changes occurring over the stabilization phase and
provides a conceptual framework for understanding this critical, final stage in a landfill's life
cycle.

Importance

Aging landfills pose significant risks to environmental stability and are currently poorly modeled
beyond ~20 years. Our examination of a single landfill across 39 years of waste degradation was
a unique opportunity to examine the impact of time within a connected system. Our work
connects geochemical data, microbial membership and predicted function, as well as physical
processes (*e.g.*, leachate recirculation). Our conceptual model interlinks these facets across the
lifespan of a landfill, providing an empirical data-based model of landfill aging. Previous models
were extrapolated from younger waste and did not include the microbial dimension – a critical
facet of the landfill ecosystem. Our model clarifies processes taking place in older wastes (30+
years), including oxygen infiltration, that have important implications for methane emission and
metal mobility and fate over the longer-term.

Keywords

49   Landfill, municipal solid waste, landfill leachate, metagenomics, metals, biogeochemistry,
50   leachate recirculation, microbial ecology

1.0 Introduction

Landfills are utilized across the globe to store both residential and non-residential waste, including metals from construction waste, household refuse, and electronic waste. Landfills contribute to environmental and human health concerns over the course of their expected life spans, including long-term methane emissions, punctuated and long-term groundwater contamination, and variable potential for leaching of heavy metals or other pollutants, under fluctuating redox conditions (1, 2).

Landfill refuse is generally buried in a sequential series of cells, which are each filled over a span of several years (2, 3) and expected to stably house the refuse for decades. Once closed, landfill cells go through a predictable set of biogeochemical phases as organic matter (OM) is decomposed by microbes. In phase 1 (aerobic), OM is rapidly metabolized through aerobic processes (3, 4). This phase is brief (1-2 years) as remaining oxygen is depleted via microbial respiration, leading to anoxic conditions. In phase 2 (anaerobic acid), hydrolytic, fermentative, and acetogenic bacteria continue metabolizing OM and produce organic acids, leading to decreases in pH and an overall decrease in OM (5). Phase 2 is followed by a rapid methanogenic phase (phase 3), where methane production peaks, and a slow methanogenic phase (phase 4). Methane production slows in phase 4 as labile OM sources are depleted and methanogenesis shifts to reliance on the downstream products from other organisms' hydrolysis of polymeric carbon sources (1, 3, 6). The duration of phases 2-4 are highly variable, and dependent on substrate conditions and climate, which impact the speed of OM breakdown (4, 6). Phase 4 can last for decades. It has been proposed that, beyond phase 4, landfills will enter a stabilization phase (phase 5). Some have defined this phase by the transition to microbial activity on more recalcitrant carbon sources (*e.g.*, humic and fulvic acids) (4, 6–8). At this point, due to

reduced microbial demand, oxygen entering landfill cells via passive diffusion can lead to

increased oxygen availability and more oxidizing conditions.

Shifts in redox conditions over the landfill life cycle are expected to impact microbe-

metal interactions and metal mobility as landfills age (3, 4, 9). Despite this, few have

investigated the differences in biogeochemical conditions between older and newer landfill cells,

particularly with respect to metal resistance and metal cycling (10, 11). Overwhelmingly,

existing studies compare different landfills rather than cells within one landfill, which may

impact comparisons of underlying chemistry. Additionally, they typically focus on carbon

cycling and greenhouse gas production, omitting metals. Studies have found that the early stages

of the landfill life cycle are characterized by rapid (over 5-10 years) decreases in labile organic

substrates (5, 6, 12). Others have shown that landfills experience a shift from microbial

communities associated with the breakdown of OM to those dominated by lithoautotrophs over

time (10, 13). What is less understood is how this microbial community succession impacts

metal cycling and leaching. Some studies report higher metal concentrations in newer landfills,

while others predict that metals may be mobilized in the later phases of the landfill life cycle (3,

10, 11). Increasingly oxidizing conditions in phase 5 can lead to oxidation of reduced sulfides,

releasing metals and reducing pH (3, 14). Additionally, organic substrates (*e.g.,* humic or fulvic

substances) remaining in waste during the stabilization phase (phase 5) can form soluble

complexes or colloids with metals, increasing metal mobility (3, 4).

Here we investigate microbial community structure and associated functional genes

between Older (aged 31 – 39 years) and Newer (aged 3 – 20 years) landfill cells from a landfill

in the Northeastern U.S. We connect these examinations of metabolic guilds and microbe-metal

interactions with changes in leachate chemistry collected over more than three decades. A

previous study by Grégoire *et al.* (2023) looked at microbiological and geochemical evidence for

methane cycling in the same system, where their highlighted oxygen infiltration into older

landfill cells. As a result, our study focused on differences in microbial communities and leachate

geochemistry over time, focusing on major microbial metabolic processes involved in and

controlled by redox cycling. We hypothesized that older landfill cells would host microbial

communities whose functional genes had a higher potential for utilization of inorganic electron

acceptors and cycling of metals while newer landfill cells would have microbial communities

with functional genes more strongly associated with anaerobic fermentative processes.

2.0 Results

2.1 Site description and sampling

The sampled site is a municipal landfill located in the northeastern United States (NEUS), site anonymity requested by site engineers. The site is composed of six landfill cells (A, B, C, D, E, and F), two of which are divided into sub-cells (D into D1 and D2, and F into F1 and F2), each equipped with leachate collection systems. Landfill cells were broadly separated by age into "Older" (cells A, B, and C; aged 31 – 39 years), "Intermediate" (cells D1 and D2; aged 24 – 26 years), and "Newer" (cells E, F1, and F2, 3 – 20 years) based on their predicted landfill phase (Older = phase 4-5, Newer = phase 2-3). Landfill phases were determined by Grégoire *et al.* 2023 (1) in their work investigating methane cycling at the site, utilizing measured biological oxygen demand (BOD) and chemical oxygen demand (COD) to assign phases. Leachate samples were collected in 2019 for sequencing and metagenomic analysis (see Methods, (1)). The site owners enthusiastically provided historic and current leachate geochemistry records under a Freedom Of Information Act (FOIA) request. Notably, microbial samples come from a single point in time (sampling in 2019) whereas leachate chemistry data spans the lifetime of the cells. As a result, microbial analyses focus on comparisons between the oldest ("Older") and youngest ("Newer") cells. Integrations of microbial and leachate chemistry data draw on the entire dataset.

2.2 Taxonomic composition of microbial communities

Starting from 1,647 total bins, dereplication was performed at 99% similarity to compare shared populations across cells, yielding 1,178 metagenome-assembled genomes (MAGs) across all eight metagenomes (all cells and sub-cells). Fewer total MAGs (>70% completion, <10% contamination) were detected in Older landfill cells (138 ± 48) than Intermediate (245 ± 34.7) or Newer cells (248 ± 42.7), with 93 MAGs in cell A, 188 in cell B, 134 in cell C, 220 in D1, 269 in D2, 210 in E, 294 in F1, and 239 in F2. Only 32 dereplicated MAGs were shared among all three

age groups, and only 5 dereplicated MAGs were shared between Older and Newer cells to the

exclusion of Intermediate cells (Figure 1c). Beyond limited shared MAGs, there were large

taxonomic differences between communities from Older and Newer cells, based on taxonomic

assignment by GTDB-tk (release 226) (15) and from a phylogenomic tree encompassing MAGs

from all samples (Figure 1a). Taxonomic assignment, trees, and all subsequent analyses were

performed using the set of 1,647 non-dereplicated MAGs to capture cell-by-cell differences (see

Methods). Of the 1,647 MAGs in the dataset, only 77 had too few hits to the required marker

genes to be included in the tree. Removed MAGs, on average, had a relative abundance of

<0.5%. Most of these 77 MAGs (63.6%) were within the Halobacteriota (22 MAGs) or placed

within groups with small genomes that may be missing key markers (Patescibacteriota [18

MAGs], Nanobdellota [7 MAGs], Micrarchaeota [2 MAGs]). A full table of taxonomic

classifications, relative abundance, tree membership, completeness and contamination, and

dereplication cluster membership is available on the Open Science Framework (OSF)

(https://doi.org/10.17605/OSF.IO/MSDB8)(16).

Overall, Older cells had lower Shannon Diversity (3.70 ± 0.640) than Intermediate (4.94

± 0.0322) or Newer (4.63 ± 0.0445) cells. Older cells had a strong enrichment in members of the

families Gallionellaceae and Sulfurimonadaceae, with enrichment in Arcobacteraceae in cell C.

Collectively, members of the Gallionellaceae, Sulfurimonadaceae, and Arcobacteraceae

represented 50.3% relative abundance in cell A, 30.0% in cell B, and 57.8% in cell C (Figure

1b), highlighting the overall lower microbial diversity in Older cells. The Gallionellaceae family

consisted of 15 dereplicated MAGs across one unidentified (JAQTOM01 [1 MAG]) and two

identified genera (*Gallionella* [10 MAGs] and *Sideroxyarcus* [4 MAGs]), and comprised 24.4%

(3 MAGs), 26.7% (7 MAGs), and 11.5% (6 MAGs) in cells A, B, and C, respectively. Members

of the Gallionellaceae were also present in D1 (0.327% [2 MAGs]) and F1 (0.371% [1 MAG]),

but at lower abundance than in Older cells. The family Sulfurimonadaceae was represented by 24

dereplicated MAGs across 2 genera (*Sulfuricurvum* [12 MAGs] and *Sulfurimonas* [12 MAGs]) in

all cells (Figure 1b, Table S1), and comprised 25.9% (6 MAGs), 3.36% (6 MAGs), and 1.91% (7

MAGs) in cells A, B, and C, respectively. Relative abundance of members of the

Sulfurimonadaceae was <1% in all Intermediate and Newer cells (1 MAG in D1, 1 in D2, 8 in E,

and 1 in F1). Arcobacteraceae was the most abundant family in any one sample, comprising

44.4% of the microbial community in cell C. Within cell C, three dereplicated Arcobacteraceae

MAGs were present, two from the genus *Aliarcobacter* (44.0%) and one unidentified

(CAIJNA01 [0.495%]). Arcobacteraceae were not detected in cells A or B but were present at

lower abundances in cells D2 (6.22% [4 MAGs]), E (5.19% [1 MAG]), and F1 (0.548% [1

MAG]).

Contrastingly, Newer cells showed enrichments in members of the families

Dysgonomonadaceae and Cloacimonadaceae. Dysgonomonadaceae consisted of 17 dereplicated

MAGs across 2 identified genera (*Proteiniphilum* [10 MAGs] and *Petrimonas* [2 MAGs]) and 5

MAGs unidentified at the genus level. Relative abundance of MAGs in the genus *Proteiniphilum*

was 3.04% (5 MAGs), 7.45% (7 MAGs), and 17.2% (10 MAGs) in E, F1, and F2, respectively,

while it was below 1% in Older cells (3 MAGs in C only) and D1 (1 MAG), but slightly more

abundant in D2 (2.32% [3 MAGs]). Relative abundance of MAGs in the genus *Petrimonas* was

2.12% (5 MAGs), 0.392% (7 MAGs), and 2.00% (10 MAGs) in E, F1, and F2, respectively, and

less than 1% in Older (1 MAG in A and 3 in C) and Intermediate (1 MAG in D1 and 3 in D2)

cells. In total, relative abundance of all unidentified members of the Dysgonomonadaceae family

were 3.66% (3 MAGs), 0.319% (2 MAGs), and 1.61% (4 MAGs) in E, F1, and F2, respectively,

and they were only detected in one other cell (D2 at 0.393% [2 MAGs]). The family

Cloacimonadaceae consisted of 18 dereplicated MAGs, most of which were unidentified at the

genus level (UBA5456 [10 MAGs]), with the identified genera being *Syntrophosphaera* (4

MAGs) and *Cloacimonas* (4 MAGs). Relative abundance of members of the Cloacimonadaceae

was 5.38% (10 MAGs), 12.8% (9 MAGs), and 12.7% (6 MAGs) in E, F1, and F2, respectively.

Cloacimonadaceae abundances were lower in Intermediate cells, with 2.21% in D1 (3 MAGs)

and 2.80% in D2 (6 MAGs), and at less than 1% in all Older cells (1 MAG in A and 2 in C).

As shown above, Intermediate samples tended to be variable, with microbial composition

in D1 more strongly resembling Older cells, while composition in D2 was closer to Newer cells.

Interestingly, both subcells of cell D were enriched in families whose members are typically very

small cells (≤0.8 μm) with reduced genomes (17, 18). This included the family

Profunditerraquicolaceae (D1: 10.9%, D2: 0.749%) with 43 MAGs from 2 identified genera

(*Undivivens* [D1: 0.449%, D2: 0.242%] and *Sherwoodlollariibacterium* [D1:0.353%, D2:

0.241%]), each represented by 1 dereplicated MAG, and 41 MAGs from unidentified genera

(D1: 10.14% [24 MAGs], D2: 0.266% [2 MAGs]). Members of the Profunditerraquicolaceae

were also found in Older (30 MAGs in A, B, and C) and Newer (2 MAGs in F1 only) cells at less

than 1% relative abundance in all cells except B, where relative abundance was 9.47% (22

MAGs).

Similarly, two families within the Patescibacteriota phylum were also enriched in the

Intermediate (D) cells. Members of the Minisyncoccaceae, represented by 17 MAGs across 2

identified genera (*Microsyncoccus* [3 MAGs] and *Minisyncoccus* [2 MAGs]) and 4 unidentified

genera (12 MAGs), had 2.94% (5 MAGs) and 10.6% (8 MAGs) relative abundance in D1 and

D2, respectively. Members of Minisyncoccaceae were also present in Newer cells, with 0.762%

(1 MAG), 3.44% (6 MAGs), and 2.97% (5 MAGs) relative abundance in E, F1, and F2,

respectively. Relative abundance of Minisyncoccaceae was less than 2% in all Older cells,

detected only in B and C with 3 MAGs in each cell. Finally, an uncultured order (UBA6257) in

the Patescibacteriota phylum (18 MAGs) had high relative abundance in cell D, with 7 MAGs in

D1 and 3 MAGs in D2, accounting for 9.61% and 1.04% relative abundance, respectively.

MAGs from UBA6257 were also present in Older (B: 8.75% [4 MAGs], C: 0.248% [1 MAG])

and Newer (E: 5.45% [7 MAGs], F1: 0.723% [1 MAG], F2: 0.178% [2 MAGs]) cells.

Although there were some similarities at the Family level, few dereplicated MAGs (5

total) were shared between Older and Newer cells (Figure 1c). In contrast, 42% of dereplicated

MAGs in Intermediate cells were shared with Newer, Older, or both. Thus, Intermediate cells

reflect a transitionary state between Older and Newer, highlighting how few taxa can persist

following transition between landfill phases. Due to the clear distinction between microbial

community compositions in Older and Newer cells, we opted to focus on comparisons between

these two age groups for our downstream analyses.

2.3 Microbial diversity and functional genes

There were substantial differences in predicted functional capacity for microbial

communities in landfill cells of different ages. In total, we examined 200 different functional

gene groups and pathways. We utilized Distilled and Refined Annotation of Metabolism

(DRAM) v1.0 (19) to annotate MAGs, focusing on ninety-three functional gene groups and

pathways, covering central metabolism, carbohydrate-active enzymes (CAZymes), conversion of

short-chain fatty acids (SCFAs) and alcohols, methanogenesis and methanotrophy, nitrogen

cycling, and sulfur cycling. Annotations from four functional groups were identified using

FeGenie (20), focusing on iron cycling. Ten more were annotated with an As metabolism and

resistance pipeline from Dunivin *et al.* (21), and 93 were derived from BacMetScan (22),

covering microbial metal interactions (respiration and resistance). Genes which contributed to

functional gene categories across the different annotation tools are listed in Table S2. Only genes

which were significantly different at $p < 0.05$ are reported here.

To investigate differences in functional potential between cells of differing ages, we

integrated all annotations into a presence-absence table for all functional genes, groups, or

pathways (available on the Open Science Framework at

https://doi.org/10.17605/OSF.IO/MSDB8) (16). DRAM estimates coverage of metabolic

pathways and completion of electron transport chain (ETC) complexes based on the percentage

of genes recovered. For our presence-absence assessment, we considered pathways and ETC

complexes to be present when DRAM's estimate was >70% for a given MAG. Then, abundance

of functional genes was calculated by summing the relative coverage of all MAGs that possessed

a given functional gene, group, or pathway within a cell, representing the proportion of the

microbial community with the encoded potential for a given function.

Across the full dataset of 1,647 MAGs, only 6 MAGs (from 0.107 to 0.730% relative

abundance) had no functional gene hits with the 70% completion cutoff, including for central

metabolic pathways. Four of these MAGs were members of the Patescibacteriota phylum, and

two were members of the archaeal phylum Nanobdellota (formerly "Nanoarchaeota"). Members

of both of these groups live as symbionts on prokaryotic hosts, lacking portions of some

pathways essential for independent life (18, 23, 24). Genomes for these organisms ranged from

70.8 – 78.5% completion, so it is also possible that core metabolic genes were absent from our

annotations due to incomplete genomes.

We investigated differences in functional gene and pathway abundance and prevalence

between Older and Newer cells using Wilcoxon rank sum tests ($p < 0.05$) (Table S2). To explore

the fermentative capacity of the microbial communities, we investigated the presence or absence

of Complexes I - V of the electron transport chain and DRAM annotations associated with the

breakdown of SCFAs and carbohydrates (CAZymes).

Complex IV of the ETC is associated with transfer of electrons to oxygen, the final

electron acceptor in aerobic respiration, so a lack of this complex indicates likely anaerobic

respiration. The absence of Complexes I - IV of the ETC indicates that an organism is likely an

obligate fermenter. Overall, we found that 7 out of 8 Complexes I - IV of the electron transport

chain were significantly more abundant in Older cells, including both low and high affinity

oxidases, indicating that variable oxygen availability in the system supports aerobic metabolisms

(Table S2, Figure S1). The relative lack of Complexes I - IV in Newer cells points to

fermentative metabolisms and anaerobic respiration as dominant in Newer cells. ATPases also

point to differences in metabolic processes between cell ages. F-type ATPases, which were

significantly more abundant in Older cells, often couple to an ETC for oxidative

phosphorylation, pointing to aerobic respiration when Complexes I - IV are also present. In

Newer cells, V/A-type ATPases were significantly more abundant, possibly reflecting

maintenance of proton gradients in microbes with low respiratory activity.

Functional gene groups associated with central carbon metabolism and organic matter

decomposition also indicated that fermentation was the key metabolic process in Newer cells

compared to aerobic processes in Older cells (Figure 2). Older cells had significantly higher

abundance of genes or gene groups associated with the reductive pentose phosphate (Calvin)

cycle (Older: $59.5 \pm 17.7\%$; Newer: $19.3 \pm 8.47\%$), reductive citrate (Arnon-Buchanon) cycle

(Older: 43.1 ± 10.9%; Newer: 22.5 ± 7.99%), glyoxylate cycle (Older: 14.2 ± 19.4%; Newer:

2.12 ± 3.46%), and citrate (Krebs) cycle (Older: 60.3 ± 22.4%; Newer: 15.4 ± 14.7%),

suggesting importance of both aerobic and anaerobic respiration, as well as carbon fixation in

Older cells (Figure 2). Notably, across all samples, genes for the reductive acetyl-CoA (Wood-

Ljungdahl) pathway, associated with anaerobic carbon fixation, were very uncommon, at 1.09 ±

1.36% in Older cells and 2.75 ± 1.02% in Newer cells.

273   Contrastingly, Newer cells had significantly higher abundance of gene groups associated

with breakdown of carbohydrates. These included CAZymes for breaking down complex

carbohydrates like pectin, starch, mannans, and xyloglucan, with 10/16 identified CAZymes at

significantly higher proportions in Newer cells. The only exceptions were CAZymes for

polyphenolics and mixed-linkage glucans, which were at significantly higher abundance in Older

cells. The relatively higher abundance of CAZy genes for polyphenolic and mixed-linkage

glucans in Older cells may point to more complex carbon sources remaining in Older cells.

Additionally, Newer cells had higher abundance of some genes and gene groups associated with

the breakdown of SCFAs and alcohols (5 out of 11 genes were significantly different), such as

acetate, butyrate, and pyruvate (Figure 2). Enrichment in CAZy genes and genes associated with

SCFA breakdown further point to fermentative activity in Newer cells.

284   Finally, there were also significant differences in genes and gene groups associated with

methanogenesis and methanotrophy between Older and Newer cells (Table S2; Figure S2).

Briefly, genes associated with methane oxidation were significantly more abundant in Older

cells, further indicating possible oxygen infiltration alongside remaining methane in Older cells,

though overall relative coverage of MAGs with those genes was low (<2%) (Figure S2). Trends

in methanogenesis and methanotrophy were explored in further detail in Grégoire *et al.*, 2023

(1).

Older cells had higher abundance of MAGs predicted to utilize a variety of alternative

electron donors and acceptors. Based on DRAM annotations (for nitrogen [N] and sulfur [S]) and

FeGenie results (for iron [Fe]), we found that functional genes and gene groups for reduction and

oxidation of all three elements were significantly more abundant in Older cells, pointing to

metabolic flexibility in Older cells (Figure 3). Notably, genes associated with all steps of

denitrification (nitrate -> nitrite [*napA* or *narG* gene] -> nitric oxide [respiratory nitrite reductase

gene] -> nitrous oxide [nitric oxide reductase gene] -> nitrogen [nitrous-oxide reductase gene])

were significantly more abundant in Older cells, with abundance of these genes ranging from

11.5-55.9% in Older cells and from 2.63-15.5% in Newer cells (Figure 3). Similarly, abundance

of the functional gene groups associated with oxidation of nitrite to nitrate (*nxrA* and *nxrB* genes)

(Older: $25.2 \pm 26.9\%$; Newer: $15.5 \pm 5.1\%$) and bacterial/archaeal ammonia oxidation (*amoA*

gene) (Older: $1.96 \pm 1.71\%$; Newer: $0.340 \pm 0.588\%$) were also significantly higher in Older

cells. One exception was dissimilatory nitrate reduction to ammonia (*nrfA* gene), which was

significantly more abundant in Newer cells (Older: $1.96 \pm 1.76\%$; Newer: $19.3 \pm 7.47\%$). Finally,

abundance of the nitrogen fixation gene *nifH* was significantly higher in Older cells (Older: 38.7

$\pm 21.4\%$; Newer: $7.61 \pm 4.13\%$). Taken together, the primary processes for generating

biologically available nitrogen differ between cell age categories. There was evidence for both

aerobic and anaerobic processes, including N fixation, in Older cells, whereas genes for

dissimilatory nitrate reduction to ammonia, a process common in strongly reducing

environments, were more abundant in Newer cells. These results highlight the differing

mechanisms of N acquisition for landfill microbiota as landfills age.

Functional gene groups for reduction and oxidation of S and Fe followed similar trends.

Two functional gene groups for the oxidation or reduction of sulfate were significantly different

in abundance between Older and Newer cells. The functional gene group for partial or complete

oxidation of thiosulfate to sulfate (including *soxA, soxB, soxX, soxY* and *soxZ*) was highly

abundant in Older ($22.7 \pm 25.5\%$) cells, in contrast to Newer ($4.70 \pm 8.14\%$) cells. Interestingly,

the functional gene group for dissimilatory reduction or oxidation of sulfate (*dsrA* or *dsrB*) was

also much higher in Older ($16.5 \pm 9.33\%$) than Newer ($1.84 \pm 1.35\%$) cells (Figure 3).

Functional gene groups identified by FeGenie as associated with iron oxidation (Older: $38.5 \pm$

15.5%; Newer: $0.875 \pm 1.00\%$) and possible iron oxidation or reduction (Older: $7.09 \pm 2.92\%$;

Newer: $0.890 \pm 0.154\%$) were also significantly more abundant in Older cells (Figure 3; Table

S2). The increase in sulfate and iron reduction genes in Older cells relative to nitrate reduction

genes in Newer cells may reflect shifts in terminal electron acceptors as substrates are depleted

over time. Finally, abundance of arsenic metabolism genes (*e.g.*, *arrA*, *aioA*) were not

significantly different between Older and Newer cells across either the Dunivin *et al*. HMM set

(21) or via the BacMet database. The heightened abundance of genes or gene groups for both

oxidation and reduction of Fe, S, and N compounds in Older cells highlights a greater reliance on

alternative electron donors and acceptors in Older cells.

Although there were no significant differences in abundance of respiratory arsenic

metabolism genes, genes associated with As detoxification and tolerance were significantly

different. Abundance of the arsenic methyltransferase *arsM* (Older: $17.3 \pm 4.30\%$; Newer: $4.30 \pm$

1.23%), and two cytoplasmic arsenate reductase genes (thioredoxin-dependent and glutaredoxin-

dependent) were also significantly higher abundance in Older cells. Thioredoxin-dependent

arsenate reductase was more abundant in general (Older: $17.8 \pm 15.3\%$; Newer: $7.80 \pm 5.46\%$)

than glutaredoxin-dependent arsenate reductase (Older: 8.06% ± 5.27%; Newer: 2.41 ± 3.88%).

Both are associated with cytoplasmic reduction of As(V) to As(III) so it can be pumped out of

the cell via an arsenite efflux pump (*acr3* or *arsB*). Both arsenite efflux pump genes were

significantly more abundant in Older cells, with arsenical resistance-3 protein (*acr3*) (Older: 23.6

± 8.96%; Newer: 9.78 ± 2.95%) being more abundant than *arsB*, which was detected through

both the Dunivin *et al*. HMM set and BacMetScan. The two annotation tools use different

methods to annotate genes (21, 22), and there was little overlap in annotation of *arsB* between

the two methods, with 70 MAGs identified as containing *arsB* from the Dunivin *et al*. set and 99

from BacMet, but only 4 overlapping between the two. Across both annotation tools, abundance

of the gene encoding the arsenite efflux pump (*arsB*) ranged from 1.31 to 8.43% in Older cells

and from 0.266 to 5.99% in Newer cells. The arsenic metallochaperone (*arsD*), also significantly

more abundant in Older cells (Older: 5.76 ± 1.39%; Newer: 1.84 ± 1.55%), chaperones

intracellular As(III) to *arsB*-encoded efflux pumps. Higher relative abundance of genes that

encode various parts of the As detoxification system point to more prevalent As-resistance in

Older cells relative to Newer cells. Additionally, the phosphate-transporting ATPase (*pstB*),

which can bring As(V) into cells, was also significantly more abundant in Older cells (Older:

26.1 ± 9.30%; Newer: 4.36 ± 6.21%), indicating the possibility for increased uptake of As in

Older cells as a consequence of P scavenging.

We also looked for other metal-resistance genes using BacMetScan and the BacMet

database, finding that Older cells displayed significantly higher abundance of genes associated

with mitigating chromate toxicity (*chrA*, *chrF*), copper resistance and metabolism (*copR*),

cobalt-zinc-cadmium resistance (*czcA*), cobalt, cadmium, and nickel transport (*dmeF*), tellurium

resistance (*terD*), and repairing DNA damage following chromate, tellurite, or selenite exposure

(*ruvB*) (Figure 3; Figure S3). The DNA repair gene encoding ATP-dependent DNA helicase

(*ruvB*) was the most abundant of these metal resistance genes (Older: 21.1 ± 12.8%; Newer:

0.660 ± 0.843%). All the other metal resistance genes listed above were below 3% relative

coverage, and, interestingly, five of six of these (*chrA*, *chrF*, *czcA*, *dmeF*, and *terD*) were not

found in any MAGs from Newer cells. The gene encoding the copper resistance regulator (*copR*)

was also below 3% abundance on average but present in both Older (2.66 ± 3.10%) and Newer

(0.404 ± 0.495%) cells.

2.4 Leachate chemistry

Leachate chemical parameters were investigated for the 5 years prior to sampling and the

period of filling for individual cells. Key parameters were selected that associate with microbial

functions of interest (Table 1, from full set in Table S3). Tables of all leachate chemistry

parameters over time have been uploaded to the Open Science Framework

(https://doi.org/10.17605/OSF.IO/MSDB8) (16). Further supporting the differential abundance of

redox-cycling genes discussed above, conditions were broadly less reducing in Older cells

relative to Newer cells (Table 1). BOD and COD were both significantly lower in Older cells in

the 5 years prior to sampling, and there was a massive decline in both BOD (from 9629 ± 9940

to 71.6 ± 194.2 mg/L) and COD (from 14285 ± 13298 to 446 ± 387 mg/L) since the time of

filling, indicating a clear decrease in labile carbon sources (Table 1; Table S3). Total organic

carbon (TOC) in leachate was also significantly lower in Older cells in this time window. Other

carbon sources were also at significantly lower concentrations in Older cells, including total

phenolics, and acetic acid and propionic acid, both SCFAs. The other investigated SCFAs were

not statistically significantly different but had overall lower concentrations in Older cells (Table

S3). Shifts in BOD, COD, and labile carbon sources over time and between Older and Newer

cells reflect differences in abundance of functional genes related to fermentation and carbon

metabolism over time. Interestingly, leachate concentrations of metals (As, Cr, Fe, Ni) and

redox-associated parameters (reduced sulfide and ammonia) were significantly higher in Newer

cells than Older cells in the 5 years prior to sampling, despite lower abundance of functional

genes associated with sulfate reduction and metal resistance in Newer cells (Table S3). This may

be linked to sulfate reduction decreasing soluble concentrations of metals in Older cells, with

sulfides incorporated into solid waste.

The 5-year window was selected to represent the geochemical conditions that had most

recently shaped the microbial communities examined. It remains possible that earlier conditions

have exerted a long-term impact on community membership and functional capacity. Leachate

concentrations of most metals (Cd, Cr, Co, Cu, Fe, Mn, Ni, Zn) were significantly higher ($p <$

0.05) in Older cells at the time of filling relative to the time of sampling (Table S3). For example,

concentrations of Cr in leachate decreased from 160 ± 410 to 6.17 ± 4.12 μg/L from filling to the

5 years prior to sampling, over 25 times lower. Similarly, Fe also decreased to over 25x less,

from 849 ± 2960 to 29.4 ± 101 mg/L (Table S3). Microbial communities in Older cells also had

significantly higher abundance of functional genes and gene groups associated with resistance to

some of these metals (Cr, Cd, Cu, Co, and Zn) and Fe oxidation and reduction, possibly pointing

to a link between historic metal concentrations and present-day capacity for microbe-metal

interactions.

To develop a picture of changes in overall landfill chemistry over decades, we aggregated

leachate chemistry data into 5-year rolling averages. We computed z-scores for each parameter to

assess differences in trends over time and clustered parameters into 8 k-means clusters based on

an elbow plot of within-cluster sum of squares (Figure S4, Table 2) to combine leachate

chemistry variables into clusters with similar variance over time. We separated these into those

associated with key explanatory parameters (conductivity, redox, pH, OM breakdown) and

remaining chemical groupings, highlighting changes over time by plotting averaged z-scores

(Figure 4, Figures S5-S9). This allowed us to identify key chemical parameters driving changes

in landfill conditions. A table of all z-score adjusted and averaged cluster data is available on the

Open Science Framework (https://doi.org/10.17605/OSF.IO/MSDB8) (16).

Interestingly, we found that a group of parameters (SO_4^{2-}, Cu, Mg, Ni, Zn) clustered with

the historic practice of leachate recirculation at the site, co-varying over time, with all parameters

peaking in concentration at around 10 years since cells began filling (Cluster 1, Figure 4).

Leachate recirculation occurred off and on within the landfill from 1985 to the end of 2009. To

investigate the impact of leachate recirculation as a driver of leachate chemistry, we performed

simple linear regressions on z-score-scaled 5-year rolling averages for all cell ages where there

was leachate recirculation (Table S4). Parameters significantly influenced ($p < 0.05$) by the

amount of recirculated leachate included carbon-associated parameters (BOD_5, COD), some

metals (Cu, Cr, Fe, Ni, Zn), Mg, oxidation-reduction potential, and two redox-associated

nutrients (nitrite-N, sulfate). Given that leachate recirculation is generally performed to maintain

reducing conditions and speed up waste decomposition, increases in BOD_5/COD and decreases

in oxidation-reduction potential are to be expected. The increases in metal concentrations may be

linked to faster breakdown of waste, mobilizing more metals into solution, in addition to metals

being recirculated back into the system.

The largest cluster contained parameters associated with the breakdown of carbon sources

(BOD-5, COD, TOC, SCFAs, total phenolics) and some metals (Ca, Cd, Cr, Fe) (Cluster 2,

Figure S5). Calcium concentrations and carbon parameters all followed a similar trend, broadly

peaking just prior to the 10-year mark then declining (Figure S5). For Cd, Cr, and Fe, the rise

and fall were more gradual, but still present. The peak for this cluster was earlier than the peak

for the recirculation-associated group (including Cu, Mg, Ni, and Zn) (Table 2; Figure 4, Figure

S5). Contrastingly, oxidation-reduction potential (sole parameter in Cluster 3) displayed a broad

dip at 10 years and a slow rise afterward, reflecting potential oxygen infiltration into the aging

landfill (Figure S6).

434   Conductivity was clustered with sulfide, ammonia-N, bicarbonate, As, K, and Na (Table

435   2). Conductivity and concentrations of ammonia-N, bicarbonate, K, and Na all followed a similar

436   trend with a broad peak between 10 and 20 years (Figure S7). Sulfide and As were less similar,

437   with a bump in concentration just before the 20-year mark (Figure S7). pH only clustered with

438   Mn, both of which peaked at the beginning of the landfill life cycle before dropping off (Figure

439   S8).

440   The remaining clusters, all depicted in Figure S9, were cobalt alone, nitrite-N alone, and

441   a cluster containing nitrate-N and selenium. Trends in these were more sporadic over time, and

442   total concentrations tended to be low (below 5 mg/L for nitrate-N and nitrite-N and below 1

443   mg/L for Co and Se).

444   Results for redox-associated parameters can be explained in the context of redox

metabolism broadly. Manganese peaked early in the landfill life cycle, followed by Fe, both of

which would likely be released into landfill leachate upon reduction from oxide forms. Then,

following the large peak in Fe, there was a small increase in sulfide. This redox cascade is

expected, though measured sulfide concentrations were quite low (<15 mg/L).

3.0 Discussion

This study investigated differences in geochemistry and microbial community structure,

composition and functional genes in leachate from cells in a municipal solid waste landfill across

39 years of waste deposition. Few studies have explored the chemical and biological conditions

in landfill cells in the later phases of decomposition under field conditions (phases 4 and 5) (3,

13). We compared "Older" (aged 31 – 39, phase 4/5) cells and "Newer" (aged 3 – 20, phase 2/3)

cells, finding significant differences in microbial community structure and composition as well

as microbial functional genes, and connected these differences to the underlying leachate

chemistry. To anchor this discussion, we developed a conceptual model synthesizing results from

microbial community functional predictions and landfill chemistry modeled over time (Figure 5).

3.1 Microbial contributions across the 5-phase model of the landfill life cycle

This study fills important gaps in our understanding of the role microbial assemblages

play in driving leachate chemical changes over the landfill life cycle, particularly in the later

phases. In the early phases of the landfill life cycle, one of the key changes is an accumulation of

labile organic matter (*e.g.*, organic acids) as microbes break down complex organic material (3,

12). We observed a gradual increase in leachate organic acids (acetic, propionic, butyric,

isobutyric, isovaleric, and valeric acid), biological oxygen demand, and chemical oxygen

demand from 0-7 years for landfill filling, before a decrease of these parameters to near zero by

year 12 (Figure 5b – cluster 2). Similar trends in organic acids and BOD/COD have been found

by other authors, who have observed major decreases within 10 years (12) or as little as 4 years

(5). Within this first seven years, pH declined substantially due to this accumulation of organic

acids, produced from hydrolysis and fermentation of OM during the anaerobic acid phase (phase

2) of the landfill life cycle (3–5). It is expected that organic acids produced in phase 2 then feed

the rapid methanogenesis that characterizes phase 3, explaining the overall decline in organic

acids and biological and chemical oxygen demand (4).

Our microbial samples, collected in 2019, captured two cells transitioning from phase 2

to 3 (F1 and F2) and one in phase 3 (E), making up our group of "Newer" cells. These cells had

higher Shannon Diversity and higher numbers of MAGs compared to Older cells. Very few

MAGs were shared between Older and Newer cells, reflecting differences in geochemical

context and shifts in microbial metabolism and survival strategies as landfill waste decomposes

over time (3). Due to the overall higher diversity of the microbial communities in Newer cells,

only the families Dysgonomonadaceae and Cloacimonadaceae were more relatively abundant

than other families across Newer cells (Figure 1b). MAGs from both families encoded a variety

of genes associated with carbohydrate-active enzyme (CAZyme) families and with breakdown of

SCFA and alcohol conversions. Members of the Dysgonomonadaceae have been found in a

variety of environments, including landfills (25–27) and anaerobic digesters (28, 29), and are

associated with the breakdown of a variety of organic compounds that are commonly present

early in a landfill life cycle. In our data, the genomes of members of the Dysgonomonadaceae's

encoded genes for most of the investigated CAZymes as well as for conversions of butyrate,

acetate, and pyruvate. Similarly, members of the Cloacimonadaceae have been found in landfills

(30) and anaerobic digesters (31, 32), and are associated with syntrophic propionate oxidation

(32). Genomes from the Cloacimonadaceae contained fewer genes for CAZymes, targeting

primarily chitin and starch degradation. The Cloacimonadaceae were also predicted to be active

in acetogenic metabolisms, which likely support downstream methanogenesis (1, 30).

The abundance of genes associated with fermentative processes was significantly higher

overall in Newer cells. This included CAZymes for the breakdown of many polysaccharides and

genes associated with SCFA and alcohol conversion (Figure 2). Further supporting the

predominance of fermentative metabolism, Newer cells had a near absence of genes associated

with complexes I - IV of the electron transport chain, which are necessary for respiration (Figure

5).

During phase 3 (represented by Newer cells), rapid methanogenesis is the dominant

process, utilizing breakdown products from the anaerobic acid phase (*e.g.*, SCFAs). Methane was

being generated in all Newer cells, with large decreases in total gas flared in Intermediate and

Older cells relative to Newer cells (explored in greater detail in Grégoire *et al.* [1]). In keeping

with this, Grégoire *et al.* (1) found a higher proportion of putative methanogens in Newer and

Intermediate cells than in Older cells. Supporting this shift in microbial metabolism, we also

observed large decreases in BOD, COD, TOC, and SCFAs, consistent with microbial

consumption of labile carbon sources and a likely transition from fermentation-based processes

to predominantly methanogenesis early in the landfill life cycle, corresponding to the shift from

phase 2 to 3 (Figure 5).

During phase 4, total gas production tends to decrease, but methanogenesis proceeds (4,

7, 33). Reduced concentrations of SCFAs and other more labile carbon substrates by this phase

lead microbes to rely on degradation of more complex OM, a slower process that reduces overall

methane production. In our data, Intermediate-aged cells (D1 and D2) had more recently entered

phase 4, while Older cell C was also classified to this phase. In all phase 4 cells (C and D), total

gas production was substantially lower than in phases 2-3 (Newer cells) but still dominated by

methanogenesis. We also observed a decrease in the abundance of genes or functional gene

groups associated with fermentative functions and a corresponding increase in MAGs carrying

genes that confer metabolic flexibility across a variety of redox processes.

Cell C exhibited higher abundance of functional genes and gene groups associated with

redox cycling of Fe, N and S than D1/ D2, reflecting the shift toward metabolic flexibility by the

end of the phase 4, as methanogenesis continues to slow (Figure 3; Figure 5). Interestingly, we

also observed overall lower leachate concentrations of N, S, and Fe by phase 4, indicating a shift

in microbial metabolism to lithoautotrophy as methanogenesis slowed and conditions became

less reducing (Figure 5b). These changes in functional potential were in large part due to shifts in

microbial community composition between cells D1/D2 and cell C. Cell C, representing a later

portion of phase 4, was broadly dominated by 2 MAGs from the genus *Aliarcobacter* (44.0%

relative abundance). In particular, the genomes of these MAGs encoded genes related to

denitrification (nitrate > nitrite > nitric oxide), N fixation, degradation of carbon sources

(polyphenolics, chitin, mixed-linkage glucans), oxidation of thiosulfate, and several genes related

to acetate metabolism (acetate kinase, phosphate acetyltransferase, acetate--CoA ligase [ADP-

forming] subunit alpha). The potential for metabolic flexibility, something documented across

the Arcobacteraceae family (34, 35) may explain the predominance of this group in cell C. This

metabolic flexibility may give them a competitive advantage as biological activity declines

throughout phase 4.

Phase 5 of the landfill life cycle, called the stabilization or maturation phase by some and

the humic phase by others, is characterized by an increased potential for oxygen buildup and

oxidative processes, as well as organic matter that is largely recalcitrant (*e.g.*, humic and fulvic

acids) (6–8). Because this phase may take decades to develop, few studies have investigated

microbial communities and leachate geochemical conditions in detail. By phase 5, we observed

an overall decline in carbon-associated parameters in leachate, including BOD, COD, TOC, and

concentrations of SCFAs. This decrease in BOD, COD, and the BOD to COD ratio is often seen

as indicative of the transition to "mature" landfill status (3, 5, 33). These declines also signify

reduced overall biological activity, indicating reductions in rates of decomposition as landfill

waste ages – a critical indicator of the stabilization phase. This reduction in decomposition may

be linked to the observed increase in oxygen flared from Older cells at this site, indicating that a

lower proportion of infiltrating oxygen is being consumed by microbes (1). This observed

increase in oxygen, though still at low levels, has been proposed by several authors as an

additional indicator of the stabilization phase (3, 4). We also observed an increase in oxidation-

reduction potential across in this window, a further indicator of a shift toward more oxidizing

conditions and processes in the later phases of the landfill life cycle.

Reflecting the decrease in biological oxygen demand, we observed fewer overall MAGs

and lower diversity in Older cells (A, B, Cs) relative to Newer cells. Microbial communities in

Older cells were relatively enriched in chemolithoautotrophs (*e.g.,* Gallionellaceae,

Sulfurimonadaceae), contrasting with the variety of diverse putative heterotrophs observed in

Newer cells. This shift from chemoorganoheterotrophs to chemolithoautotrophs was also

observed by Sekhohola-Dlamini *et al.* (2020), who found that a 14-year old MSW landfill with

higher total carbon and N was dominated by copiotrophs, while a nearby 36-year old landfill was

dominated by lithotrophs (13). Taxonomic shifts were also reflected in differences in functional

gene abundances between Older and Newer cells. As noted above, relative abundance of MAGs

with genes associated with the fermentation of carbon sources was significantly lower in Older

cells. Possibly associated with increases in oxygen availability, significantly more MAGs in

Older cells had genes associated with electron transport chain complexes I-IV, indicating the

potential for aerobic respiration under changing redox conditions. Furthermore, Older cells had

more metabolically flexible microbial communities, with significantly higher relative abundance

of MAGs with genes for reduction and/or oxidation of Fe, N, and S. For example, members of

the genera *Gallionella* and *Sideroxyarcus* (formerly *Sideroxydans*), which were enriched in Older

cells, are widely known to oxidize ferrous iron (36–38). Similarly, members of two other

enriched genera, *Sulfuricurvum* and *Sulfurimonas*, contain members known to be metabolically

flexible, capable of utilizing a variety of reduced S compounds as electron donors and nitrate or

oxygen as electron acceptors (39–41). Concentrations of nitrate, nitrite, and sulfide in leachate

were all relatively low (generally < 5 mg/L across all time points), indicating either low levels

available in the landfilled waste, steady conversion/uptake rates by the microbial community, or

both. In contrast, concentrations of sulfate were relatively higher (> 10 mg/L) throughout the

landfill life cycle, with no significant differences between Newer and Older cells at the time of

sampling.

3.2 Redox response across decades of landfill waste decomposition

Ammonia-N, sulfate, and Fe concentrations in leachate spiked at different points in the

landfill life cycle. Each of these redox-associated metabolites was linked to different

biogeochemical cycles in our dataset. First, other studies have reported a spike in ammonia-N

midway through the landfill life cycle as organics are depleted, mobilizing N via hydrolysis of

proteins in OM (3, 5). We observed a similar trend, with ammonia-N broadly peaking with

conductivity, bicarbonate, potassium, and sodium at around 20 years of landfill waste age (Figure

5b - cluster 4). This peak in conductivity, bicarbonate, and base cations (K and Na) has been

observed elsewhere. Conductivity has been observed to increase in young and intermediate-aged

landfills as ions are released from breakdown of organic material (3, 6, 42). Contributing to

conductivity, bicarbonate tends to increase in leachate as degradation of organic material

produces CO_2, forming carbonic acid and ultimately, bicarbonate ions (42, 43).

A notable trend during the mobilization and then reduction in labile carbon substrates

from 0 – 10 years was an increase in leachate concentrations of Fe, Ca, Cd, and Cr, with peaks

occurring around 6-8 years. (Figure 5b – Cluster 2). Although mobilization of these elements

may be linked to their pH and redox-dependent natures (3, 8), it is important to note that pH

generally remained above 6 in all samples at all time points, a phenomenon observed in other

landfills' anaerobic acid phase (14). It is therefore unlikely that mobilization of Fe, Ca, Cd, and

Cr is entirely pH-dependent, as the measured pH was not acidic enough to release these metals

into solution. Instead, it is possible that metals formed soluble complexes or colloids with OM,

increasing leachate concentrations relative to the later parts of the landfill life cycle (3).

Furthermore, reducing conditions in the anaerobic phases may have influenced metal mobility,

reducing, for example, Fe(III) to soluble Fe(II) (14). Further study is needed to fully understand

the links between organic chemistry and metal cycling in landfill leachate over time.

Finally, concentrations of sulfate, Cu, Mg, Ni, and Zn, were all linked to the historic

practice of leachate recirculation at the site (Figure 4; Figure 5b). Leachate recirculation is

common in MSW landfills, where it is used to raise moisture content and hasten anaerobic

decomposition (3, 44). At our site, leachate recirculation was practiced until the end of 2009,

with variable volumes recirculated in each cell monthly. Linear regression analysis revealed that

the volume of leachate recirculated was negatively associated with redox and nitrite-N and

positively with BOD, COD, sulfate, Cu Cr, Fe, Mg, Ni, and Zn over time. Outside of studies

specifically investigating leachate recirculation in landfills (44, 45), few mention if the practice

was performed at their sites or consider the broader impacts recirculation has on microbial guilds

that control waste decomposition, making it difficult to put our results in context. From our data,

we suggest that more reducing conditions (decreases in redox potential) are an expected response

to leachate recirculation, which is broadly associated with increased rates of fermentation of

organic matter, possibly leading to higher BOD and COD (44, 45). Increased moisture content,

brought about by recirculation of leachate, can also promote dissolution of metal-bearing

minerals (*e.g.*, Fe/Mn-oxides) and mobilize metals previously bound to solid phases – this is

particularly true for metals like Cu, Ni, and Zn that more readily solubilize or bind to organic

ligands, thus keeping them in the leachate solution (14). Increased moisture content in landfill

cells may also increase contact time, which can release a variety of ions from solid phases,

including metals and sulfate.

3.2 Metal fate within a landfill

Given concerns about the impact of increasingly oxidizing conditions on metal mobility

and metal leaching in landfills (4, 14), one of the goals of our study was to investigate changes in

metal availability and putative metal cycling over the landfill life cycle. As mentioned above, the

practice of leachate recirculation appeared to accelerate metal release into landfill leachate. Our

study identified a previously-unreported association between leachate recirculation and metal

concentrations, allowing an initial examination of this phenomenon. Leachate recirculation

ceased at this site in 2009, so we had more data for leachate recirculation from Older and

Intermediate cells (A, B, C, D1, D2), with cell E only filling from 1999 onwards, and cell F not

extant in 2009 (2014 start to filling). Overwhelmingly, metal concentrations were higher in A, B,

and C (all filled before 1993) at the time of filling than in D1 and D2 (filled between 1993 and

1998), possibly due to changes in waste streams and waste management practices over time.

More research is needed to investigate the impacts of leachate recirculation on metal mobility

and fate, as existing research focuses on recirculation's role in accelerating breakdown of OM.

We hypothesized that Older landfill cells would host microbial communities with a

higher abundance of functional genes associated with utilization of inorganic electron acceptors

and cycling of metals. We found that the abundance of metal resistance genes was significantly

higher in Older cells relative to Newer. Many different genes for metal resistance (for As, Cd,

Cu, Co, Cr, Fe, Ni, Se, Te) were significantly more abundant in Older cells than Newer cells,

despite lower metal concentrations. Genes for Fe oxidation and reduction were the only metal

metabolism genes with significantly higher proportional abundances in Older cells. Interestingly,

outside of Fe-cycling, the majority of metal resistance and metal cycling genes or gene groups

were encoded by lower abundance MAGs that were not part of the most dominant families in

Older cells (Gallionellaceae and Sulfurimonadaceae). Genes for As resistance were some of the

most ubiquitous in Older cells, present across a variety of taxa including many, but not all,

Gallionellaceae and Sulfurimonadaceae MAGs, lineages which harbour As resistance genes in

other environments (40, 46). Genes for As resistance have been found across a variety of

environments (21), including deep groundwater systems (46) and permafrost (47). While metal

exposure may partially explain the high incidence of metal resistance genes in Older cells, which

had highest metal levels historically (48), some studies have seen little overlap between metal

exposure and relevant metal-resistance genes (46, 49). It is possible that fluctuations in redox

conditions within the solid material in Older landfill cells results in spatial differences in metal

exposure that influence microbial community composition and functional gene profiles, but that

is difficult to determine without sampling solid landfill waste. Our study provided a

comprehensive look at a wide range of metals, and further study is needed to elucidate direct and

indirect biogeochemical cycling pathways for landfill metals.

3.3 Conclusions

In our study, we examined microbial communities, functional genes, and leachate

chemistry across a landfill's cells of varying ages (2-39 years). We also tracked changes in

geochemical conditions across 36 years of landfill leachate chemistry data. We found that the

early phases of the landfill life cycle were dominated by fermentative processes, with labile

carbon declining over time. Conditions became more oxidizing over time, with evidence of

oxygen buildup in Older cells. Microbial diversity declined with time, and there was a clear shift

from fermentative processes to utilization of alternative electron donors and acceptors. Our

results were largely consistent with studies that modelled potential biogeochemical cycles in

landfills in the stabilization phase, having never recorded them (3, 4).

Leachate metal concentrations were influenced by OM content and historic leachate

recirculation practices. Leachate recirculation was associated with increases in metal mobility

and the potential for metal leaching out of the system, possibly linked to organo-metal

complexes. Future research investigating organo-metal complexes and the ability of natural

landfill decomposition to pull metals into solution would be an exciting new avenue to better

understand metal fate in these environments, especially in conjunction with better modeling for

how leachate recirculation-modified moisture regimes impact metal leaching and OM cycling.

Our conceptual model of the key processes and metabolisms over a landfill's life cycle
provides a new framework, based on empirical data, and synthesizing physical processes with
microbial impacts. We note that this landfill's cells were only at the beginning of the stabilization
phase. The landfill should continue to be monitored for changes in geochemical conditions and
microbial community succession over time to better understand the unexplored dynamics of this
crucial, final stage in a landfill life cycle.

4.0 Materials and Methods

4.1 Site Description

The sampled site is a municipal landfill located in the northeastern United States (NEUS).

The site is composed of six landfill cells (A, B, C, D, E, and F), two of which are divided into

sub-cells (D into D1 and D2, and F into F1 and F2), each equipped with leachate collection

systems. Previous work at this site classified the landfill cells using a five-phase conceptual

model based on leachate chemistry and biogas data (1). Cell A, which received waste from 1980

to 1982, and Cell B, which received waste from 1982 to 1988, were both classified as in phase 5.

Cell C, which received waste from 1988 – 1993 was classified into phase 4. The authors of this

prior study identified evidence of oxygen intrusion into cells A and B, informing their

classification into phase 5 of the landfill life cycle. The two subcells of cell D were filled at

different times. D1 received waste from 1993 – 1998 and cell D2 from 1995 – 1998. Both were

classified into phase 4, due primarily to low gas production relative to cells E and F. Cell E was

filled from 1999 – 2014 and was classified into phase 3 (rapid methanogenesis) due to high gas

output, mostly composed of methane. Cell F1 filled from 2014 up to the time of sampling

(March 2019) and F2 from 2016 up to the time of sampling (March 2019). Both were identified

as transitioning from phase 2 (anaerobic acid phase) to phase 3 due to variable gas production

but still high proportions of methane (1). For this study, decreases in oxygen demand (BOD) and

chemical oxygen demand (COD) at A, B, and to a lesser extent, C, relative to the other cells

prompted us to classify A, B, and C as "Older". Cells E, F1, and F2 were categorized as

"Newer", while D1 and D2 were labelled "Intermediate".

4.2 Leachate sampling

In 2019, we collected leachate samples from each cell/sub-cell for shotgun metagenomic

sequencing. One leachate monitoring well at each cell was purged of 2-3 well volumes, and one

litre of leachate was collected using a peristaltic pump. Leachate samples were stored on ice and

transported to the lab before being filtered through a 0.22 μm Sterivex filter (Millipore Sigma,

Burlington, MA) on the day of sampling. Filters were stored at -80°C until DNA extractions

were performed. Detailed descriptions of DNA extraction, sequencing, filtering, assembly,

binning, and taxonomic assignment can be found in Grégoire *et al.* 2023 (1), which exclusively

explored the predicted methane-cycling populations across the landfill. The pipeline for

microbial community analyses is described in brief below.

4.3 DNA sequencing and prior analyses

DNA was extracted from biomass on Sterivex filters using the PowerSoil DNA Isolation

Kit (Qiagen) according to the manufacturer's protocols but with the filter added to the initial bead

tube in place of soil, and with elution into 50 μL of final buffer. DNA was quality-checked using

a NanoDrop 1000 (Thermo Scientific, Waltham, MA) and quantified using Qubit fluorimetry

(Thermo Scientific, Waltham, MA). Extracted DNA was sent to The Centre for Applied

Genomics in Toronto, Canada for shotgun metagenomic sequencing on an Illumina HiSeq

machine with paired 2x150 bp reads. Acquired metagenomic reads were quality-trimmed with

bbduk in the BBTools suite (https://sourceforge.net/projects/bbmap/) and Sickle v1.33 (50) prior

to assembly using SPAdes3 v3.15.5 (51). Scaffolds were trimmed by length ($\geq$ 2.5 kbp) and

reads were mapped onto scaffolds using Bowtie2 v2.3.4.1 (52). Scaffolds were binned using

three different binning algorithms (CONCOCT v0.4.0, MaxBin2 v2.2.6, MetaBAT2 v2.12.1) and

bins were dereplicated using DAS Tool v 1.1.1 before quality-checking in CheckM v 1.0.13 (53–

57). A total of 1,881 metagenome-assembled genomes (MAGs) were retained with >70%

completion and <10% contamination. Relative abundance of MAGs was calculated using the

mean coverage for unique scaffold identifiers within each genome bin. To evaluate the functional

potential of microbial communities, MAGs were annotated using Distilled and Refined

Annotation of Metabolism (DRAM) v1.0 (19) to look at a wide array of functional genes using

both sequence similarity and hidden Markov model (HMM) based detection.

4.4 Data analyses

All previous work by Grégoire *et al.* is outlined in brief in section 4.3 (1). For this study,

taxonomy was assigned to metagenome-assembled genomes (MAGs) using the Genome

Taxonomy Database Toolkit (GTDB-tk) r266 (15). Phylogenomic analysis for all MAGs was

performed using GToTree v1.6, using a suite of 25 hidden Markov models (HMMs) for bacteria

and archaea to identify single copy genes and generate a tree (58). 1,570 out of our 1,647 MAGs

were placed on the tree. The resultant tree was visualized using the Interactive Tree of Life tool

(59). To investigate similarity of MAGs across landfill cells, we utilized the dereplication tool

dRep, which uses Mash and genome-wide ANI (gANI) to create clusters of MAGs based on

pairwise comparisons, choosing a similarity cutoff of 99% (60).

The product file from DRAM was used to investigate the presence of genes associated

with central metabolism, carbon-cycling, nitrogen-cycling, and sulfur-cycling. DRAM combines

genes into functional gene groups based on the presence of key genes. For example, for the

function "acetate, pt 1" to be considered present, MAGs need to have genes for both phosphate

acetyltransferase and acetate kinase. We utilized these DRAM categories in all cases except the

SOX complex for oxidation of thiosulfate, which DRAM considers to be present if any *sox* gene

(*A, B, C, D, X, Y,* or *Z)* is present. Researchers have demonstrated that the presence of *soxXA,*

soxB, and *soxYZ* are necessary for partial oxidation of thiosulfate, with *soxCD* completing the

reaction (61). Accordingly, we decided to apply a stricter threshold than DRAM, considering the

combined presence of *soxA, soxB, soxX, soxY,* and *soxZ* as required to indicate the ability for

partial or complete oxidation of thiosulfate.

DRAM also annotates metabolic pathways and multi-subunit complexes (*e.g.*, electron

transport chain [ETC]) based on the proportion of genes present. We considered a pathway or

complex to be present when over 70% of genes were present. We utilized HMM-based pipelines

for Fe and As metabolism and resistance, using FeGenie (20) for iron and Dunivin *et al.*'s arsenic

resistance and metabolism pipeline (21). FeGenie uses a library of curated HMMs to group iron-

related functional genes, while Dunivin *et al.*'s pipeline looks for individual marker genes using

their HMM set. We also utilized BacMet version 2.0 for targeted annotation of metal-resistance

genes by looking for homologs to experimentally-confirmed genes from the BacMet database

(22). Genes from the BacMet database were filtered to only those which were involved in metal

resistance, excluding genes related to genes for antibiotic resistance. Because BacMet and the

Dunivin *et al*. pipeline employ different methods, we retained hits from both, even if the genes

overlapped (*e.g.*, *arsB* from Dunivin vs. *arsB* from BacMet). Further information on genes

contributing to functional groups can be found in the documentation for the tools listed above

(Table S2).

All subsequent data analyses, including statistics and generation of figures, were

performed in R version 4.3.1 (2023-06-16 ucrt) (62). We separated data into three categories for

downstream analysis: Older cells aged 31 – 39 years (cells A, B, and C) and Newer cells aged 3

– 20 years (cells E, F1, and F2). D1 and D2 (aged 24 – 26) were considered Intermediate and

were not included in statistical analyses. Base R was used for statistical tests. The packages

"phyloseq" and "microeco" were used for generating Venn diagrams of MAG overlap among

categories and for producing relative abundance bar graphs (63, 64). Relative abundance was

calculated by summing the mean coverage for all MAGs at a given taxonomic level (*e.g.*, at the

Family level). Similarly, functional gene abundance was calculated using the relative coverage of

MAGs that possessed a given functional gene, group, or pathway. Due to the variety of tools

used for annotating functional genes, we collapsed data into binary presence-absence data for

each gene, pathway, or gene group. Therefore, if 3 MAGs in a cell had coverages of 1%, 2%, and

3%, and each possessed at least one *arsA* gene, the "abundance" of the functional gene would be

6%. Wilcoxon rank sum tests were used to determine differences in functional gene abundance

and chemical parameters between Older and Newer cells. Finally, heatmaps of functional gene

abundance were generated using the package "ggplot2" (65).

4.5 Leachate chemistry

Thirty-six years of leachate chemistry records were provided by the site owners, with all

analyses performed by Brickhouse Environmental consultants. While many parameters were

provided by the site owners, we selected the following as relevant to microbial function and

showing variance over time; pH, oxidation-reduction potential, electrical conductivity,

bicarbonate, BOD, COD, TOC, sulfate, sulfide, ammonia-N, nitrate-N, nitrite-N, organic acids

(acetic acid, propionic acid, isobutyric acid, butyric acid, isovaleric acid, valeric acid), total

phenolics, base cations (Ca, K, Mg, Na) and metals (As, Cd, Co, Cr, Cu, Fe, Mn, Ni, Se). As

leachate chemistry has changed greatly over time for some cells, we reported chemistry both

during the time of filling and in the 5 years prior to our sampling to capture shifts over time.

To investigate trends in geochemical parameters over time, we computed 5-year rolling

averages across all cells by time since filling across all 36 years (year 0 – 36) of provided

leachate data. Then, we computed z-scores for each parameter, transforming data to a mean of 0

and standard deviation of 1 so trends across parameters could be compared agnostic of units.

Using an elbow plot of within-cluster sum of squares to determine the optimum numbers of

clusters (Figure S4), we clustered the z-score adjusted data into 8 k-means clusters (Table 2).

Plots of 5-year rolling averages were developed using the "ggplot2" package in R (65). Finally,

to investigate the impact of the historic practice of leachate recirculation on leachate

geochemistry, we ran linear regressions on z-score scaled 5-year rolling averages for all cells

across the time window for which there was active leachate recirculation (> 5 gallons of

leachate/month).

Acknowledgements

We are sincerely grateful to the landfill site management (anonymity by request) and

Brickhouse Environmental consulting for site access, aid with sampling, and provision of

detailed monitoring records. We thank Dr. Jennifer Biddle and her lab group for hosting our team

for sample processing prior to shipment. Thanks also to Dr. Nikhil George, Ms. Rebecca Co, and

Dr. Alexandra Sauk for help with sampling, and to Dr. Nikhil George for his work with binning

pipelines. This work was supported by an NSERC Discovery Grant (2016-03686) and an Ontario

Early Research Award (ER19-15-228) to LAH. LAH was supported by a Tier II Canada

Research Chair.

Data Availability

All data involved in this research are publicly available. Sequencing data have been

deposited to NCBI under the BioProject PRJNA900590, with raw reads available on the SRA

database. Geochemical data, microbial functional annotations, and microbial taxonomy data are

all available in the Open Science Framework (https://doi.org/10.17605/OSF.IO/MSDB8). All R

code used for statistical analyses and to generate figures are also on the Open Science

Framework.

CRediT authorship contribution statement

Kimber E. Munford: Data curation, formal analysis, investigation, methodology, visualization,
writing – original draft, writing – review & editing. **Daniel S. Grégoire:** Data curation, formal
analysis, investigation, methodology, software, writing – review & editing. **Laura A. Hug:**
Conceptualization, data curation, funding acquisition, project administration, resources, software,
supervision, writing – review & editing.

Conflict of interest statement

The authors declare that there are no conflicts of interest to disclose.

References

1. Grégoire DS, George NA, Hug LA. 2023. Microbial methane cycling in a landfill on a decadal time scale. Nat Commun 14:7402.

2. Meyer-Dombard DR, Bogner JE, Malas J. 2020. A Review of Landfill Microbiology and Ecology: A Call for Modernization With 'Next Generation' Technology. Front Microbiol 11:1127.

3. Kjeldsen P, Barlaz MA, Rooker AP, Baun A, Ledin A, Christensen TH. 2002. Present and Long-Term Composition of MSW Landfill Leachate: A Review. Crit Rev Environ Sci Technol 32:297–336.

4. Pohland FG, Harper SR. 1985. Critical review and summary of leachate and gas production from landfills. United States Environmental Protection Agency (EPA).

5. Kulikowska D, Klimiuk E. 2008. The effect of landfill age on municipal leachate composition. Bioresour Technol 99:5981–5985.

6. Wijekoon P, Koliyabandara PA, Cooray AT, Lam SS, Athapattu BCL, Vithanage M. 2022. Progress and prospects in mitigation of landfill leachate pollution: Risk, pollution potential, treatment and challenges. J Hazard Mater 421:126627.

7. Liu J, Gu Z, Wang X, Li Q. 2022. The molecular differences of young and mature landfill leachates: Molecular composition, chemical property, and structural characteristic. Chemosphere 287:132215.

8. Bozkurt S, Moreno L, Neretnieks I. 1999. Long-term fate of organics in waste deposits and its effect on metal release. Sci Total Environ 228:135–152.

9.   Östman M, Wahlberg O, Mårtensson A. 2008. Leachability and metal-binding capacity in

ageing landfill material. Waste Manag 28:142–150.

10. Zainun MY, Simarani K. 2018. Metagenomics profiling for assessing microbial diversity in

both active and closed landfills. Sci Total Environ 616–617:269–278.

11. Hosseini Beinabaj SM, Heydariyan H, Mohammad Aleii H, Hosseinzadeh A. 2023.

Concentration of heavy metals in leachate, soil, and plants in Tehran's landfill:

Investigation of the effect of landfill age on the intensity of pollution. Heliyon 9:e13017.

12. Kang K-H, Shin HS, Park H. 2002. Characterization of humic substances present in landfill

leachates with different landfill ages and its implications. Water Res 36:4023–4032.

13. Sekhohola-Dlamini L, Selvarajan R, Ogola HJO, Tekere M. 2021. Community diversity

metrics, interactions, and metabolic functions of bacteria associated with municipal solid

waste landfills at different maturation stages. MicrobiologyOpen 10:e1118.

14. Kim H, Jang Y-C, Townsend T. 2011. The behavior and long-term fate of metals in

simulated landfill bioreactors under aerobic and anaerobic conditions. J Hazard Mater

194:369–377.

15. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. 2019. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. Bioinformatics 36:1925–1927.

16. Hug LA. 2026. Munford_NEUS_Landfill_metals.

https://doi.org/10.17605/OSF.IO/MSDB8. Open Science Framework.

17. Seymour CO, Palmer M, Becraft ED, Stepanauskas R, Friel AD, Schulz F, Woyke T, Eloe-

Fadrosh E, Lai D, Jiao J-Y, Hua Z-S, Liu L, Lian Z-H, Li W-J, Chuvochina M, Finley BK,

Koch BJ, Schwartz E, Dijkstra P, Moser DP, Hungate BA, Hedlund BP. 2023. Hyperactive

nanobacteria with host-dependent traits pervade Omnitrophota. Nat Microbiol 8:727–744.

18. Srinivas P, Peterson SB, Gallagher LA, Wang Y, Mougous JD. Beyond genomics in

Patescibacteria: A trove of unexplored biology packed into ultrasmall bacteria. Proc Natl

Acad Sci U S A 121:e2419369121.

19. Shaffer M, Borton MA, McGivern BB, Zayed AA, La Rosa SL, Solden LM, Liu P, Narrowe

AB, Rodríguez-Ramos J, Bolduc B, Gazitúa MC, Daly RA, Smith GJ, Vik DR, Pope PB,

Sullivan MB, Roux S, Wrighton KC. 2020. DRAM for distilling microbial metabolism to

automate the curation of microbiome function. Nucleic Acids Res 48:8883–8900.

20. Garber AI, Nealson KH, Okamoto A, McAllister SM, Chan CS, Barco RA, Merino N.

2020. FeGenie: A Comprehensive Tool for the Identification of Iron Genes and Iron Gene

Neighborhoods in Genome and Metagenome Assemblies. Front Microbiol 11.

21. Dunivin TK, Yeh SY, Shade A. 2019. A global survey of arsenic-related genes in soil

microbiomes. BMC Biology 17:45.

22. Pal C, Bengtsson-Palme J, Rensing C, Kristiansson E, Larsson DGJ. 2014. BacMet:

antibacterial biocide and metal resistance genes database. Nucleic Acids Res 42:D737–

D743.

23. Kato S, Ogasawara A, Itoh T, Sakai HD, Shimizu M, Yuki M, Kaneko M, Takashina T,

Ohkuma M. 2022. *Nanobdella aerobiophila* gen. nov., sp. nov., a thermoacidophilic,

obligate ectosymbiotic archaeon, and proposal of *Nanobdellaceae* fam. nov., *Nanobdellales*

ord. nov. and *Nanobdellia* class. nov. Int J Syst Evol Microbiol 72:005489.

24. Kuroda K, Maeda R, Shinshima F, Urasaki K, Kubota K, Nobu MK, Noguchi TQP, Satoh

H, Yamauchi M, Narihiro T, Yamada M. 2024. Microbiological insights into anaerobic

phenol degradation mechanisms and bulking phenomenon in a mesophilic upflow anaerobic

sludge blanket reactor in long-term operation. Water Res 253:121271.

25. Co R, Hug LA. 2021. Prediction, enrichment and isolation identify a responsive,

competitive community of cellulolytic microorganisms from a municipal landfill. FEMS

Microbiol Ecol 97:fiab065.

26. Xu S, Lu W, Liu Y, Ming Z, Liu Y, Meng R, Wang H. 2017. Structure and diversity of

bacterial communities in two large sanitary landfills in China as revealed by high-

throughput sequencing (MiSeq). Waste Manag 63:41–48.

27. Hu L, Huang F, Qian Y, Ding T, Yang Y, Shen D, Long Y. 2024. Pathways and contributions

of sulfate reducing-bacteria to arsenic cycling in landfills. J Hazard Mat 473:134582.

28. Pecorini I, Rossi E, Becarelli S, Baldi F, Di Gregorio S, Iannelli R. 2023. Wet Anaerobic

Codigestion of Sewage Sludge and OFMSW in Pilot-Scale Continuously Stirred Tank

Reactors: Focus on the Reactor Microbial Communities. 4. Sustainability 15:3168.

29. Santinello D, Zampieri G, Agostini S, Müller B, Favaro L, Treu L, Campanaro S. 2024.

Process stability in anaerobic Digestion: Unveiling microbial signatures of full-scale reactor

performance. Chem Eng J 497:154962.

30.   Johnson LA, Hug LA. 2022. Cloacimonadota metabolisms include adaptations in

engineered environments that are reflected in the evolutionary history of the phylum.

Environ Microbiol Rep 14:520–529.

31.   Schwan B, Abendroth C, Latorre-Pérez A, Porcar M, Vilanova C, Dornack C. 2020.

Chemically Stressed Bacterial Communities in Anaerobic Digesters Exhibit Resilience and

Ecological Flexibility. Front Microbiol 11.

32.   Dyksma S, Gallert C. 2019. Candidatus Syntrophosphaera thermopropionivorans: a novel

player in syntrophic propionate oxidation during anaerobic digestion. Environ Microbiol

Rep 11:558–570.

33.   Luo H, Zeng Y, Cheng Y, He D, Pan X. 2020. Recent advances in municipal landfill

leachate: A review focusing on its characteristics, treatment, and toxicity assessment. Sci

Total Environ 703:135468.

34.   Li J, Xiang S, Li Y, Cheng R, Lai Q, Wang L, Li G, Dong C, Shao Z. 2024.

Arcobacteraceae are ubiquitous mixotrophic bacteria playing important roles in carbon,

nitrogen, and sulfur cycling in global oceans. mSystems 9:e00513-24.

35.   Buzzanca D, Kerkhof P-J, Alessandria V, Rantsiou K, Houf K. 2023. Arcobacteraceae

comparative genome analysis demonstrates genome heterogeneity and reduction in species

isolated from animals and associated with human illness. Heliyon 9:e17652.

36.   Zhou N, Keffer JL, Polson SW, Chan CS. 2022. Unraveling Fe(II)-Oxidizing Mechanisms

in a Facultative Fe(II) Oxidizer, Sideroxydans lithotrophicus Strain ES-1, via Culturing,

Transcriptomics, and Reverse Transcription-Quantitative PCR. Appl Environ Microbiol

88:e01595-21.

37. Kato S, Itoh T, Iino T, Ohkuma M. 2022. Sideroxyarcus emersonii gen. nov. sp. nov., a

neutrophilic, microaerobic iron- and thiosulfate-oxidizing bacterium isolated from iron-rich

wetland sediment. Int J Syst Evol Microbiol 72:005347.

38. Hallbeck L, Pedersen K. 2014. The Family Gallionellaceae, p. 853–858. *In* Rosenberg, E,

DeLong, EF, Lory, S, Stackebrandt, E, Thompson, F (eds.), The Prokaryotes. Springer

Berlin Heidelberg, Berlin, Heidelberg.

39. Han Y, Perner M. 2015. The globally widespread genus Sulfurimonas: versatile energy

metabolisms and adaptations to redox clines. Front Microbiol 6.

40. Handley KM, Bartels D, O'Loughlin EJ, Williams KH, Trimble WL, Skinner K, Gilbert JA,

Desai N, Glass EM, Paczian T, Wilke A, Antonopoulos D, Kemner KM, Meyer F. 2014.

The complete genome sequence for putative H2- and S-oxidizer Candidatus Sulfuricurvum

sp., assembled de novo from an aquifer-derived metagenome. Environ Microbiol 16:3443–

3462.

41. Kodama Y, Watanabe K. 2004. Sulfuricurvum kujiense gen. nov., sp. nov., a facultatively

anaerobic, chemolithoautotrophic, sulfur-oxidizing bacterium isolated from an underground

crude-oil storage cavity. Int J Syst Evol Microbiol 54:2297–2300.

42. Naveen BP, Mahapatra DM, Sitharam TG, Sivapullaiah PV, Ramachandra TV. 2017.

Physico-chemical and biological characterization of urban municipal landfill leachate.

Environ Pollut 220:1–12.

43. Christensen TH, Kjeldsen P, Bjerg PL, Jensen DL, Christensen JB, Baun A, Albrechtsen H-

955   J, Heron G. 2001. Biogeochemistry of landfill leachate plumes. Appl Geochem 16:659–718.

956   44. Chan GYS, Chu LM, Wong MH. 2002. Effects of leachate recirculation on biogas

957   production from landfill co-disposal of municipal solid waste, sewage sludge and marine

958   sediment. Environ Pollut 118:393–399.

959   45. Zheng J, Wang X-G, Sun Y, Wang Y-X, Sha H-Q, He X-S, Sun X-J. 2024. Natural and

960   anthropogenic dissolved organic matter in landfill leachate: Composition, transformation,

961   and their coexistence characteristics. J Hazard Mat 465:133081.

962   46. Chen M, Grégoire DS, St-Germain P, Berdugo-Clavijo C, Hug LA. 2025. Microbial

963   diversity and capacity for arsenic biogeochemical cycling in aquifers associated with

964   thermal mobilization. Sci Total Environ 977:179357.

965   47. Mindlin S, Petrenko A, Kurakov A, Beletsky A, Mardanov A, Petrova M. 2016. Resistance

966   of Permafrost and Modern Acinetobacter lwoffii Strains to Heavy Metals and Arsenic

967   Revealed by Genome Analysis. Biomed Res Int 2016:3970831.

968   48. Roosa S, Wattiez R, Prygiel E, Lesven L, Billon G, Gillan DC. 2014. Bacterial metal

969   resistance genes and metal bioavailability in contaminated sediments. Environ Pollut

970   189:143–151.

971   49. Ayala-Muñoz D, Burgos WD, Sánchez-España J, Couradeau E, Falagán C, Macalady JL.

972   2020. Metagenomic and Metatranscriptomic Study of Microbial Metal Resistance in an

973   Acidic Pit Lake. Microorganisms 8:1350.

974   50.   Joshi N, Fass J. 2025. Sickle: a sliding-window, adaptive, quality-based trimming tool for

975   FastQ files (Version 1.33).

976   51.   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020. Using SPAdes De

977   Novo Assembler. Curr Protoc Bioinformatics 70:e102.

978   52.   Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods

979   9:357–359.

980   53.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing

981   the quality of microbial genomes recovered from isolates, single cells, and metagenomes.

982   Genome Res 25:1043–1055.

983   54.   Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, Banfield JF. 2018.

984   Recovery of genomes from metagenomes via a dereplication, aggregation and scoring

985   strategy. Nat Microbiol 3:836–843.

986   55.   Alneberg J, Bjarnason BS, de Bruijn I, Schirmer M, Quick J, Ijaz UZ, Lahti L, Loman NJ,

987   Andersson AF, Quince C. 2014. Binning metagenomic contigs by coverage and

988   composition. Nat Methods 11:1144–1146.

989   56.   Wu Y-W, Simmons BA, Singer SW. 2016. MaxBin 2.0: an automated binning algorithm to

990   recover genomes from multiple metagenomic datasets. Bioinformatics 32:605–607.

991   57.   Kang DD, Froula J, Egan R, Wang Z. 2015. MetaBAT, an efficient tool for accurately

992   reconstructing single genomes from complex microbial communities. PeerJ 3:e1165.

58.   Lee MD. 2019. GToTree: a user-friendly workflow for phylogenomics. Bioinformatics

35:4162–4164.

59.   Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic

tree display and annotation. Nucleic Acids Res 49:W293–W296.

60.   Olm MR, Brown CT, Brooks B, Banfield JF. 2017. dRep: a tool for fast and accurate

genomic comparisons that enables improved genome recovery from metagenomes through

de-replication. ISME J 11:2864–2868.

61.   Friedrich CG, Rother D, Bardischewsky F, Quentmeier A, Fischer J. 2001. Oxidation of

Reduced Inorganic Sulfur Compounds by Bacteria: Emergence of a Common Mechanism?

Appl Environ Microbiol 67:2873–2882.

62.   R Core Team. 2023. R: A language and environment for statistical computing. manual. R

Foundation for Statistical Computing, Vienna, Austria.

63.   McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. PLoS One 8:e61217.

64.   Liu C, Cui Y, Li X, Yao M. 2021. microeco: An R package for data mining in microbial

community ecology. FEMS Microbiol Ecol 97:fiaa255.

65.   Wickham H. 2016. ggplot2: Elegant graphics for data analysis. Springer-Verlag New York.

1010
1011

1012 **Figures**
1013

1014

Figure 1: a) Phylogenomic tree based on 25 target gene HMMs for bacteria and archaea,
placing 1,570 out of 1,647 MAGs in relation to each other, generated in GToTree v1.6 (58)**.**
Rings depict, outer to inner: 1) family-level taxonomy, 2) cell age category. MAGs in the tree
were not dereplicated so that sample-specific relative abundances could be depicted. Colours for
families correspond to those in the b) barplot of relative abundance of all bacterial and archaeal
families with over 5% relative abundance in at least one landfill cell, with lower abundance
lineages grouped under "Other". Taxonomy was taken from GTDB-tk r266, where lineages
starting with UBA are officially recognized but not yet formally named. c) Venn diagram
indicating counts and percentages of shared MAGs (clustered at 99% similarity using DRep)
among age categories across landfill cells. Venn sections are colored by proportion of MAGs
within a category, with higher values corresponding to darker colors.

1026

1027

Figure 2: Carbon cycling differential abundance across landfill cells. Heatmap depicting the
summed relative coverage of all MAGs in each landfill cell that contain genes associated with

specific functions within carbon cycling. Only genes/pathways/processes which had significantly

different abundances between Older and Newer cells (Wilcoxon rank sum test, Benjamini-

Hochberg-corrected p-value of $p < 0.05$) are displayed. Boxes indicate whether abundance was

significantly higher in Newer or Older cells. Central C metabolism pathways from DRAM were

required to be >70% complete within a MAG for inclusion. Carbohydrate-active enzymes from

DRAM either indicate the presence of genes for both backbone and oligomer cleavage (alpha-

mannan, arabinan, beta-galactan, mixed-linkage glucans, pectin, starch, sulf-polysaccharides,

xyloglucan) or only oligomer cleavage (arabinose cleavage, fucose cleavage, rhamnose

cleavage). For polyphenolics, multiple oxidative genes are accounted for. For DRAM SCFA and

alcohol conversions, the gene(s) of interest are listed in brackets. Further information can be

found in the DRAM documentation.

1041

1042

Figure 3: Redox cycling and metal resistance differential abundance across landfill cells.
Heatmap depicting the summed relative coverage of all MAGs in each cell that contain

genes/processes associated with specific functions within redox cycling and metal resistance.

Only genes/processes which had significantly different abundances between Older and Newer

cells (Wilcoxon rank sum test, Benjamini-Hochberg-corrected p-value of $p < 0.05$) are displayed.

Genes/processes with a maximum of < 0.05% relative coverage were pruned for clarity and are

presented in Figure S3. Genes and functional groups come from four tools: FeGenie and an

HMM set from Dunivin et al. for Fe and As, respectively, BacMetScan for metal-resistance

genes and DRAM for functional groups associated with S and N cycling. For As-related genes

detected using the Dunivin et al. HMM set, the gene of interest is listed. For BacMetScan

annotations, the BacMet number is listed in brackets following the gene. For FeGenie

annotations, the process is listed, followed by the relevant genes in brackets. Finally, for DRAM

annotations, the process is listed, followed by the gene(s) of interest in brackets.

1057

1058

Figure 4: Line graphs depicting the 5-year rolling average (line) and standard deviation (shaded band around line) for landfill leachate parameters across 36 years of leachate chemistry data. All depicted parameters were within one k-means cluster (cluster 1).

1062

1063

Figure 5: Conceptual diagram of landfill processes over time and landfill phases. a) Trends in functional gene groups over landfill cells varying in age from 3 to 39 years across 4 landfill phases, overlaid on predicted methane/O_2 dynamics. Solid lines represent the beginning and end of our observations, and the dashed lines represent the ages at which all other cells were sampled. Landfill phases are depicted along the top, but it is notable that cells F1 and F2 (the two newest cells) were transitioning from phase 2 to 3. Trend lines represent three functional gene groups: Fermentation (SCFA/alcohol degradation, carbohydrate-active enzymes), Respiration (defined by abundance of complete (>70% of constituent genes present) electron transport chain complexes for ETC I-IV) and Redox Processes (comprising genes for redox transformations of Fe, N, S, and As). Gas dynamics are depicted as greyscale background, with methane dominating an anoxic environment from years 3-26, followed by oxygen infiltration in later years. b) Line graph depicting trends in key geochemical variables across 36 years of leachate chemistry data. Lines depict the means of all z-score-adjusted parameters within a given k-means cluster, with standard deviation of all z-score adjusted parameters within a cluster indicated by lighter bands. Clusters are named in the legend based on their key physical variable(s). Note that redox has no standard deviation because it was the sole member of its cluster (Cluster 3).

1080   **Tables**
1081
1082   **Table 1: All geochemical measurements (mean±sd) for key variables, collated by landfill**
1083   **cell age class during filling of the cells and during the 5 years prior to sampling in winter**
1084   **2019.** Benjamini-Hochberg corrected p-values are presented for Wilcoxon rank sum comparisons
1085   between age classes in the 5 years prior to sampling. Only parameters that were significantly
1086   different at $p < 0.05$ are presented here. Remaining variables can be seen in Table S3.

Analyte	Older (A, B, C)	Newer (E, F1, F2)	Benjamini - Hochberg corrected p - value
Conductivity (μs/cm)	4665 ± 2083	7789 ± 3629	< 0.001
Redox (mV)	-36.1 ± 71.8	-88.8 ± 69.5	< 0.001
Bicarbonate (mg/L)	1557 ± 774	1842 ± 1242	0.00257
BOD_5 (mg/L)	71.6 ± 194.2	251 ± 308	< 0.001
COD (mg/L)	446 ± 387	1052 ± 875	< 0.001
TOC (mg/L)	115 ± 56.2	272 ± 227	< 0.001
Sulfide (mg/L)	0.845 ± 0.471	5.17 ± 11.4	< 0.001
Ammonia-N (mg/L)	219 ± 123	458 ± 306	< 0.001
Acetic acid (mg/L)	12.2 ± 25.1	61.3 ± 71.4	< 0.001
Propionic acid (mg/L)	9.99 ± 21.5	34.2 ± 72.4	0.0151
Total phenolics (mg/L)	0.173 ± 0.183	0.835 ± 1.59	< 0.001
Arsenic (μg/L)	16.9 ± 14.5	61.4 ± 61.4	< 0.001
Chromium (μg/L)	6.17 ± 4.12	56.8 ± 53.8	< 0.001
Iron (mg/L)	29.4 ± 101	51.8 ± 52.1	< 0.001
Manganese (mg/L)	0.418 ± 0.494	1.29 ± 1.51	< 0.001
Nickel (μg/L)	38.5 ± 26.5	110 ± 79.8	< 0.001
Calcium (mg/L)	67.6 ± 17.3	141 ± 89.3	< 0.001
Magnesium (mg/L)	48.0 ± 23.4	91.8 ± 42.8	< 0.001
Potassium (mg/L)	195 ± 94.4	242 ± 137	0.00194
Sodium (mg/L)	469 ± 226	736 ± 422	< 0.001

1087

1088 **Table 2: Cluster membership for 8 k-means clusters of z-score adjusted 5-year rolling**
1089 **averages of 36 years of landfill leachate data.**

Cluster number	Key parameter	All parameters
1	Leachate recirculation	Leachate recirculation, sulfate, Cu, Mg, Ni, Zn
2	Organics and metals	BOD-5, COD, TOC, organic acids (acetic acid, propionic acid, isobutyric acid, butyric acid, isovaleric acid, valeric acid), total phenolics, Ca, Cd, Cr, Fe
3	Redox	Redox (oxidation-reduction potential)
4	Conductivity	Conductivity, sulfide, ammonia-N, bicarbonate, As, K, Na
5	pH	pH, Mn
6	Nitrite-N	Nitrite-N
7	Nitrate-N	Nitrate-N, Se
8	Cobalt	Co

1090

1091