Soluble LAG-3 Identifies a Dynamic Early T Cell Activation Window in self-reactivity, Type 1 Diabetes, and Broader Immune Responses.

Saptarshi Roy[1,2,*,a], Megan L. Proffer[2,10,a], Farooq Syed [3,4], Megan E. Smithmyer[5], Avinil Das Sharma[2,6,9], Abigail Twoy[7,8], Emily Soo-Hoo[7,8], Jyoti Rana[3], Jason M. Spaerth[2,6,9], Everett Meyer[7,8], Kent P Jensen[7,8], Cate Speake[5], William Hagopian[2,6], Jon D. Piganelli[1,2,10]*

1. Department of Endocrinology, Indiana University School of Medicine, Indianapolis, IN, USA.
2. Center for Diabetes and Metabolic Diseases, Indiana University School of Medicine, Indianapolis, IN, USA.
3. Department of Diabetes-Immunology, Arthur-Riggs Diabetes and Metabolism Research Institute, Beckman Research Institute at the City of Hope, Duarte, CA, USA.
4. Center for RNA Biology and Therapeutics, Beckman Research Institute at the City of Hope, Duarte, CA, USA
5. Center for Interventional Immunology, Benaroya Research Institute, Seattle, WA, USA.
6. Department of Pediatrics, Indiana University School of Medicine, Indianapolis, IN, USA.
7. Division of Blood & Marrow Transplant & Cell Therapy (BMT-CT), Stanford University School of Medicine, Stanford, CA.
8. Diabetes Immune Monitoring Core (DIMC), Stanford Diabetes Research Center (SDRC), Stanford University School of Medicine, Stanford, CA.
9. Department of Biochemistry and Molecular Biology, Indiana University School of Medicine, Indianapolis, IN, USA.
10. Department of Microbiology and Immunology, Indiana University School of Medicine, Indianapolis, IN, USA.

*Corresponding Authors:

*Jon D. Piganelli, PhD, Department of Endocrinology, Indiana University School of Medicine 1210 Waterway, 4th floor, 4100F, Indianapolis, IN 46202. Email: jdpigane@iu.edu, Phone: 317-278-6013

*Saptarshi Roy, PhD, Department of Endocrinology, Indiana University School of Medicine 1210 Waterway, 4th floor, 41007, Indianapolis, IN 46202. Email: roysapt@iu.edu , Phone: 317-278-6013

[a] Both authors equally contribute.

Abstract

Aims/hypothesis: Type 1 diabetes is a complex autoimmune disorder in which autoreactive CD4$^+$ and CD8$^+$ T cells destroy pancreatic beta-cells, resulting in insulin deficiency and hyperglycemia. Although genetic susceptibility, particularly certain HLA alleles, contributes to disease risk, not all genetically predisposed individuals develop Type 1 diabetes. Screening first degree relatives (FDRs) for islet autoantibodies (GAD65, IAA, IA-2, ZnT8) helps detect autoimmune activity. However, these serum markers arise only after T-helper cell activation, limiting early intervention opportunities. Since protein antigen recognition by B cells requires T-helper cell assistance through linked recognition, T cell activation precedes B cell activation and autoantibody production. Activation of these T cells leads to shedding of the immune-regulatory (activation) surface protein LAG-3 (Lymphocyte Activation Gene-3 or CD223), generating its soluble form, sLAG-3, that is detectable in circulation. We hypothesized that sLAG-3 may serve as an early biomarker of autoimmune activity preceding islet autoantibody development in type 1 diabetes.

Methods: Plasma sLAG-3 levels were measured longitudinally in female diabetes-prone NOD mice and analyzed in relation to islet antigen-specific CD4$^+$ T cell expansion and diabetes onset. To mechanistically link autoreactive T cell activation to sLAG-3 release. Naive autoreactive C6.6.9 TCR-transgenic (TCR-Tg) CD4$^+$ T cells were adoptively transferred into NOD.SCID mice and longitudinal assessment for plasma sLAG-3, beta-cell antigen specific CD4$^+$ T cell tetramer profiles, and circulating insulin (*Ins2*) mRNA to determine ongoing beta-cell stress. In parallel, sLAG-3 levels were analyzed from different human cohorts, including FDRs of individuals with type 1 diabetes, using cross-sectional and longitudinal approaches.

Results: In murine models, elevated sLAG-3 correlated with expansion of islet-specific CD4$^+$ T cells that preceded hyperglycemia and diabetes onset. In the adoptive transfer model, early increases in sLAG-3 and circulating *Ins2* mRNA marked immune activation and emerging beta-cell stress prior to overt diabetes. In our human cohorts, sLAG-3 was detectable in autoantibody-negative and single-autoantibody-positive FDRs, with higher

levels observed in progressors compared to non-progressors, and associated with high-
risk HLA genotypes.

Conclusions/interpretation: These findings identify sLAG-3 as a candidate biomarker
of early T cell activation in type 1 diabetes that may precede islet autoantibody
development. Integration of sLAG-3 with antigen-specific T cell and beta-cell stress
markers could improve early risk stratification and inform preventive strategies before
substantial loss of beta-cell. Prospective longitudinal studies aligned to seroconversion
are required to validate sLAG-3 as a surrogate marker of early disease activity.

Keywords: Type 1 diabetes, sLAG-3, T cell activation, autoimmunity, beta-cell
destruction, biomarker, islet autoantibodies, *Insulin* mRNA, immune infiltration.

Abbreviations

ChgA   Chromogranin A
FACS   Fluorescence-Activated Cell Sorting buffer
FDR   first degree relative
GAD65   Glutamic Acid Decarboxylase 65 (isoform of GAD)
HLA   Human Leukocyte Antigen
IA-2   Islet Antigen-2 (a pancreatic β-cell antigen)
IAA   Insulin Autoantibody
IAPP   Islet Amyloid Polypeptide (also called amylin)
IFN-γ   Interferon-gamma
mRNA   messenger Ribonucleic Acid
NOD   Non-Obese Diabetic (mouse strain)
NOD.SCID   NOD. Severe combined immunodeficient
RPM   Revolutions Per Minute (common lab term, e.g., centrifugation speed)
TCR   T Cell Receptor
ZnT8   Zinc Transporter 8
90

Research in context

What is already known about this subject?

- Before the clinical onset of hyperglycemia, type 1 diabetes is characterized by a prolonged preclinical phase in which autoreactive B and T cells mediate progressive beta-cell destruction.
- Current risk stratification strategies rely mainly on genetic susceptibility (genomic DNA) and the detection of islet autoantibodies in plasma/serum.
- Islet autoantibodies arise only after CD4$^+$ T cell activation and therefore do not capture the earliest stages of immune dysregulation.
- Consequently, biomarkers that directly reflect early pathogenic T cell activity prior to, or independent of, seroconversion remain limited and insufficiently validated.

What is the key question?

Can plasma sLAG-3 levels, beta-cell antigen-specific CD4$^+$ T cell tetramer expression, and circulating *Ins2* mRNA serve as very early biomarkers of autoimmune activity in type 1 diabetes and serve to better inform risk stratification, thereby informing preventive intervention strategies for the clinician?

What are the new findings?

- sLAG-3 increases transiently during early antigen-specific CD4$^+$ T cell activation stage, precedes hyperglycemia in mouse models, and is elevated in autoantibody-negative and single-autoantibody-positive first-degree relatives who later progress to type 1 diabetes.
- sLAG-3 was associated with beta-cell antigen-specific CD4$^+$ T cell expansion, assessment of stress induced beta cell *Ins2* mRNA release and high-risk HLA genotypes, indicating early autoimmune activation rather than established disease.

How might this impact clinical practice in the foreseeable future?

These findings support sLAG-3 as a candidate early biomarker of T cell activation, before or at the earliest stages of islet autoantibody development in some at-risk individuals. Integration of plasma sLAG-3 with beta-cell antigen specific CD4$^+$ T cell profiling and

119   insulin mRNA measurements could complement current autoantibody-based screening,

120   improve risk stratification, and enable earlier preventive interventions to preserve beta-

121   cell function in patients at-risk for type 1 diabetes.

Introduction

Type 1 diabetes is a complex, heterogeneous, and polygenic autoimmune disease characterized by CD4$^+$, CD8$^+$-T cell and B cell mediated destruction of pancreatic beta-cells, resulting in insulin deficiency and hyperglycemia [1-3]. Despite extensive research, the mechanisms driving this aberrant immune positive beta-cell interaction remain incompletely understood [4]. Type 1 diabetes affects 9.5 million people globally (2 million in USA). The global incidence is rising globally by 2-3% annually, outpacing the prevalence of high-risk HLA and other genetic alleles, indicating that environmental factors, including viral infections and oxidative stress, likely contribute to disease development [5-7]. First-degree relatives of type 1 diabetes patients, particularly those carrying HLA-DR3/-DQ2 and/or -DR4/-DQ8 HLA haplotypes, are at elevated genetic risk for type 1 diabetes. Patients and at-risk individual can be screened for islet autoantibodies in plasma/serum and HLA typing genomic DNA. Current biomarkers such as GAD65, insulin, IA-2 and ZnT8 autoantibodies are typically become detectable only after substantial beta-cell loss, indicating a clinically silent window of early immune dysregulation that remains undetected [8, 9]. Thus, there is a crucial and yet unmet need to validate this silent window for biomarkers capable of identifying type 1 diabetes at the earliest pre-seroconversion stages of autoimmunity.

The FDA approval of teplizumab, an anti-CD3 lymphodepleting immunomodulatory therapy, represents a major advance in our field that, demonstraties that disease onset can be delayed when treatment is initiated post-autoantibody seroconversion [10]. Nevertheless, full disease prevention remains challenging, as intervention at this stage occurs after significant beta-cell injury [11]. Since seroconversion occurs only after immune tolerance is breached and self-reactive CD4$^+$ T cells are activated, earlier biomarkers that detect pathogenic immune activity and distinguish progressors from non-progressors are urgently needed to inform prevention strategies [11-13]. Autoantibodies primarily reflect loss of immune tolerance but arise only after CD4$^+$ T cells provide help for B cell activation and antibody production [10, 11]. Early antigen-specific CD4$^+$ T cells deliver essential co-stimulatory signals and cytokines that drive B cell proliferation, isotype class switching, and differentiation into plasma cells that produce high-affinity

autoantibodies against beta-cell antigens [14]. While both genetic and environmental
factors likely contribute to CD4[+] T cell activation, the initiating triggers remain unclear [15,
16].

During T cell activation, a cellular metalloprotease (ADAMs-17) enzymatically cleaves the
lymphocyte activation gene-3 protein from the cell surface, releasing sLAG-3 into
circulation as a T cell activation marker [12, 14, 15]. LAG-3 is a co-inhibitory receptor
expressed on regulatory T cells that is structurally similar to CD4 [17]. Increase sLAG-3
levels, previously associated with strong immune responses in cancer, and Graves'
Disease, appears to predict emerging autoimmunity in type 1 diabetes prior to
autoantibody detection [16, 18, 19]. Concurrently, in beta cells proinflammatory stress
mediate release of *insulin* mRNA into circulation, representing another promising early
marker of beta-cell dysfunction [20-22]. However, the lack of early biomarkers that can
be used to identify at-risk individuals before seroconversion, highlights the need for
enhanced assays capable of detecting both sLAG-3 and *insulin* mRNA to monitor early
immune activity and beta-cell health.

To evaluate the feasibility of these proposed biomarkers panel, we conducted a
preliminary study integrating plasma sLAG-3 quantification via monoclonal antibody-
based ELISA assay with beta-cell antigen-specific CD4[+] T cell tetramer profiling. This
combinatorial assay was applied to longitudinal plasma samples from female diabetic-
prone NOD mice, a human cohort from the Diabetes Evaluation in Washington (DEW)
study, and cross-sectional autoantibody-positive samples from the Stanford Diabetes
Biobank. In these preclinical and clinical samples, we observed early increases in plasma
sLAG-3 coincided with rising frequencies of beta-cell antigen-specific CD4[+] T cells. Our
results accurately predicting impending diabetes onset and supporting the potential utility
of this biomarker combination. We further examined early immune activation using a
longitudinal adoptive-transfer model of autoimmune diabetes in which naïve islet-reactive
CD4[+] T cells from NOD.C6.6.9 TCR-transgenic mice were transferred into
immunodeficient NOD.SCID recipients. This reductionist system enables real-time
monitoring of autoreactive T cell antigen engagement, activation, expansion, antigen
engagement, and effector differentiation. We assessed plasma sLAG-3 concentration,

CD4+ T cell tetramer positivity, and circulating *insulin* mRNA levels as biomarkers of beta-
cell health and disease progression [20]. Notably, the early rise in circulating *Ins2* mRNA
in whole blood sapless preceded hyperglycemia and coincide with increases in immune
cell infiltration into the pancreatic islets, suggesting that immune-driven β-cell dysfunction
disrupts insulin protein synthesis and secretion far earlier than previously recognized [23-
25]. These findings support the use of metabolic markers to detect early beta cell
vulnerability, that could complement immune-activation biomarkers of immune activation
and uncover targets to preserve insulin secretion. Additionally, human studies further
confirm an increase in sLAG-3 in plasma of single autoantibody-positive first-degree
relatives of individuals with type 1 diabetes compared to individuals who are
autoantibody-negativity, reinforcing its potential as an early biomarker. Combined
monitoring of sLAG-3 and CD4+ T cell tetramer expression, along with circulating *Ins2*
mRNA, may enable earlier type 1 diabetes detection and inform type 1 diabetes
intervention trial before substantial beta-cell loss and clinical onset [20, 24].

Methods

Animal Models and Diabetes Assessment

NOD/ShiLtJ (Strain 001976), NOD.Cg-Prkdc$^{scid/J}$ (Strain 001303), and C57BL/6 (Strain
000664) male and female mice were obtained from Jackson Laboratory. Beginning at 4
weeks of age, NOD/ShiLtJ mice were routinely monitored for glucosuria in 3-day intervals
during adoptive transfer studies. Blood glucose was measured in mice positive for
glucosuria, and overt diabetes was defined as two to three consecutive readings >300
mg/dL. Blood glucose was measured by puncturing the last 1 mm of the tail and
dispensing one drop of blood on a Contour next® disposable test strip and glucometer.
All mice were bred and maintained under specific pathogen-free conditions at Indiana
University's BRTC, managed by LARC. Animal experiments were conducted in
accordance with Animal Research: Reporting of In Vivo Experiments (ARRIVE) guidelines
under protocols approved by the Indiana University (IU) School of Medicine Animal Use
Committee.

Research participant's samples

Plasma from healthy controls, type 1 diabetes participants, and FDRs were provided by Children's Hospital of Pittsburgh, Stanford University School of Medicine (IRB Protocol #35453), University of Florida, and University of Queensland. Samples were stored at −80°C until analyzed. Participants with type 1 diabetes ranged in age at onset (1 to 16 yrs old) and duration of disease (1 to 6 years post-onset). FDRs were tested for auto-antibodies to GAD65, IAA, and IA2 by standard radioimmunoassay. Plasma samples were collected from two cohorts of healthy controls, younger participants aged 25-35 and older participants aged 55-65 years old who were recruited into the Sound Life Project for measuring the sLAG-3 level from research participant before and after covid vaccination. Samples were collected with approval from Benaroya Research Institute, Seattle, WA review board (IRB19-045) and the Stanford University School of Medicine institutional review board (IRB 35453). All participants provided written informed consent. Prospective cohort longitudinal studies for measuring the sLAG-3 from Diabetes Evaluation in Washington Study (DEWIT) enrolled infants from the general population (n=39) carrying increased-risk HLA-DR-DQ genotypes identified by newborn screening, as well as first-degree relatives of individuals (n=13) with or without type 1 diabetes regerdless of HLA genotype. All participants underwent HLA screening, though eligibility criteria varied across sites. Follow-up began between 2 months and 21.6 years of age and continued at 3 to 36 month intervals for up to 26 years, with type 1 diabetes diagnosed according to American Diabetes Association criteria. All protocols were approved by local institutional review boards, and de-identified data were submitted to IBM Research. The harmonized cohort comprised median 10 visits, 8.7 years follow-up; 17% reported a family history of type 1 diabetes. For key analyses, it was examined the infant-toddler subcohort, restricted to subjects tested for insulin, IA-2, and GAD autoantibodies by age 2.5 years. The infant-toddler cohort included median 12 visits, 10.4 years follow-up [26].

Estimating soluble LAG-3 levels from NOD and human plasma

NOD/ShiLtJ and NOD.Cg-Prkdc$^{scid/J}$ mice from different experimental groups were retro-orbitally bled for plasma every other week from 4 to 16 weeks of age. Plasma was isolated

by centrifugation at 3000 × RPM for 10 minutes at 4°C. Mouse sLAG-3 was measured by
sandwich ELISA using 96-well high-affinity plates (Corning #3361) coated overnight at
4°C with 5 µg/mL monoclonal anti-LAG-3 (C9B7W, InVivoMab #BE0174). LAG-3
standard was purchased from R&D (Catalog: AF3328) antibody pairs for mouse
compatible were purchased from Jackson Immuno Research, performed according to
manufacturer's protocol. Human plasma from three institutions was analyzed using
Human LAG-3 DuoSet ELISA (R&D #DY2319B). All samples were run in triplicate, read
at OD450 nm on a SpectraMax-iD5, and analyzed with SoftMax-Pro 7.0.2.

Plasma sLAG-3 evaluation from DEWIT cohort study and Stanford Biobank

Human plasma samples were analyzed for sLAG-3 levels by the MSD U-PLEX platform
using sandwich immunoassays for protein targets and a competitive assay for neopterin,
with electrochemiluminescence detection. MULTI-ARRAY® 96-well plates contained
sLAG-3 capture antibodies, Sulfo-Tag detection antibodies, calibration standards, and
diluents. The Indiana University Translational Core processed 50 µL of each sample in
triplicate. Mean concentrations were reported, with values below the limit of detection
(LOD) assigned the LOD. Plate-specific control coefficients of variation (CVs) were
calculated for sLAG-3.

Estimating sLAG-3 from covid vacationed research participants

sLAG-3 level from covid vacationed research participants plasma was measured using
the Olink proximity extension assay on the Olink Explore 1536 platform's oncology panel
[27]. Samples were randomized across plates to balance age and sex, with longitudinal
samples from the same participant run on the same plate. Bridging controls were included
in each batch for cross-batch normalization. Protein expression values were first
normalized within wells using internal extension controls (IgG antibodies with matched
oligos), then standardized across plates using triplicate inter-plate pooled serum controls.
Data were subsequently intensified across all samples, and final normalized relative
protein quantities were reported as NPX values.

In vivo viral infection incidence study

Eight-week-old C57BL/6 mice were intraperitoneally infected with 100 PFU
Coxsackievirus B (CVB3 #ATCC® VR-30™) in HBSS under specific pathogen-free
conditions. Blood was collected via retro-orbital bleeding at baseline and on days 3, 6, 9,
12, 16, and 21 post-infections. Plasma was isolated by centrifugation and stored at -80°C.
sLAG-3 levels were quantified using standard ELISA, while intracellular IFN-γ in CD4$^+$
and CD8$^+$ T cells was measured by flow cytometry to assess immune activation during
infection.

Adoptive transfer

Spleens from 8-week-old female NOD congenic C6.6.9 TCR-Tg mice were harvested
sterilely, mechanically homogenized, passed through a 40 µm strainer, washed, and
resuspended in cold DMEM with 10% FBS. Lymphocytes were isolated via Ficoll-Paque
density gradient, washed twice with PBS, and 5×10^6 cells were retro-orbitally transferred
into 8-week-old female NOD.Cg-Prkdc$^{scid/J}$ recipients. Post-transfer, mice were monitored
for diabetes progression by blood glucose and bled every three days to measure plasma
sLAG-3, activation markers and co-inhibitory receptor (CD69, CD44, LAG-3), and CD4$^+$
T cell reactivity to beta-cell antigens (ChgA, InsB9-23, IAPP) using tetramers. Mice with
>300 mg/dL glucose on two readings were euthanized for spleen and pancreas for further
analyses.

Staining and flow cytometry analysis

To assess T cell mediated immune activation in autoimmune and viral contexts,
lymphocytes were analyzed by flow cytometry. NOD/ShiLtJ mice were longitudinally bled
during diabetes progression, and NOD.SCID mice were sampled after adoptive transfer.
For viral studies, CD4$^+$IFN-γ$^+$ and CD8$^+$IFN-γ$^+$ T cells were analyzed from C57BL/6 blood.
Lymphocytes were isolated, RBCs lysed, and cells resuspended in FACS buffer. Cells
were stained with eFluor-780 viability dye and surface markers including CD45-FITC,
CD4-PerCP-Cy5.5, CD8-APC, CD69-PE/Cy7, CD25-PE, LAG-3-BV650, CD71-APC,
CD98-PE/Cy7, and Glut-1-PE. For beta-cell antigen specific, APC-ChgA, BV421- InsB9-

23, and PE-IAPP tetramers were used. Intracellular IFN-γ-PE staining was performed

using the Fixation & Permeabilization Buffer Kit protocol (eBiosciences FOXP3 Fix/Perm

kit, Cat #00-5523-00). Data were analyzed using FlowJo-v10.

Cytokine measurement by ELISA

Plasma from CVB-infected C57BL/6 mice was collected at multiple time points and

analyzed for IFN-γ by ELISA. High-affinity 96-well plates were coated overnight at 4°C

with 25 ng/mL monoclonal rat anti-murine IFN-γ in carbonate buffer (pH 9.6). Mouse IFN-

γ ELISAs were performed using BD antibody pairs following the manufacturer's protocol.

Mean OD450 values were recorded on a SpectraMax-iD5 microplatereader, analyzed

using SoftMax Pro-7.0.2.

Histology and Immunofluorescence

Pancreata from control, diabetic, and post-adoptive transfer NOD.SCID mice (days 12

and 15) adoptively transferred NOD.SCID mice were harvested at diabetes onset, fixed

in 4% paraformaldehyde for 4 hours at 4°C, washed with distilled water, and processed

for paraffin embedding, and stained with hematoxylin and eosin to assess tissue

architecture and immune infiltration following standard protocol. For

immunofluorescence, pancreata cryoprotect overnight at 20% sucrose in PBS. Tissues

were embedded in OCT, frozen, and stored at -80°C. Cryosections (6 μm) were prepared

using a cryostat. Sections were blocked with 5% normal donkey serum and 1% BSA,

incubated overnight at 4°C with monoclonal CD3, insulin, F4/80, and B220 primary

antibody, followed by fluorophore-conjugated secondary antibodies. Similarly, APC-

ChgA, BV421- InsB9-23, and PE-IAPP tetramers antibody are also used. Nuclei were

counterstained with DAPI. Images were acquired on a Zeiss LSM-800 and analyzed using

ImageJ.

RNA isolation and ddPCR

Total RNA was isolated from the blood of adoptively transferred NOD.ScidSCID mice

using RNAprotect Animal Blood Tubes (cat. #76544) and the RNeasy Protect Animal

Blood Kit (cat. #73224) per manufacturer instructions. RNA concentration was measured
with a NanoDrop-2000 (Thermo Fisher Scientific). cDNA was synthesized using M-MLV
Reverse Transcriptase (Invitrogen) according to the manufacturer's protocol. Droplet
digital PCR (ddPCR) was performed to quantify *Ins2* mRNA. Briefly, 2 µL of cDNA was
combined with 10 µL ddPCR Supermix for Probes (1863024) and 1 µL TaqMan primer
(Cat #4427975) that target *Ins2* transcript and the volume was adjusted to 20 µL with
RNase/DNase-free water. Synthetic *Ins2* cDNA was used as a positive control and RNAse
free water was used negative controls. Droplets were generated with QX200 AutoDG,
PCR on a C1000 Thermal Cycler, and read with QX200 droplet reader. Data (copies/µL)
were analyzed using QuantaSoft software.

Statistical Analysis

Data analysis was performed using GraphPad Prism10. Pairwise comparisons were
performed using unpaired or paired two-tailed Student's t-tests. Interactions between two
groups with two variables (e.g., time or treatment, pre-post-vaccination) were determined
using a one-way ANOVA followed by an appropriate post hoc test. Sample sizes (n),
means, error bars, and statistical methods are described in figure legends. A p-value
<0.05 indicates statistical significance.

Results

Plasma sLAG-3- levels track spontaneous diabetes onset in NOD mice

LAG-3 (CD223) is a co-inhibitory receptor expressed by CD4+CD25+CD127lo regulatory
T (Treg) cells and other regulatory immune cell subsets [28, 29]. We previously proposed
that protein enzyme catalytic liberation of LAG-3 from the surface of CD4 Tregs
(solubilized or sLAG-3) is a key biomarker, indicative of loss of tolerance to pancreatic
islet beta cell antigens in autoimmune-prone NOD mice [30]. To determine whether
soluble LAG-3 In plasma/serum reflects early autoimmune activation, we followed female
NOD mice longitudinally from 4-16 weeks of age, measuring plasma sLAG-3 and blood
glucose biweekly (Fig.1a). Plasma sLAG-3 was low at 4 weeks of age, increased
significantly during 6 to12 weeks, reflecting early T cell activation and then declined to

near-baseline levels by 16 weeks of age (Fig. 1b). This window of elevated sLAG-3
coincided with the prediabetic period when insulitis is known to expand in NOD mice [31].
In contrast, blood glucose remained normal through 12 weeks and began to rise around
16 weeks of age as overt, clinical diabetes appeared (Fig. 1c). Importantly, a significant
positive correlation was observed between peak sLAG-3 levels and the timing of diabetes
onset in the female NOD cohort (Fig. 1d), indicating that higher early sLAG-3 levels are
associated with greater likelihood of progression to hyperglycemia in this model.

Beta cell antigen-specific CD4$^+$ T cells expand in parallel with sLAG-3 during
spontaneous disease

We next examined whether beta-cell antigen specific CD4$^+$ T cells show similar temporal
dynamics during spontaneous disease in female NOD mice from the same longitudinal
cohort (Fig. 1a). Using MHC class II tetramers for the pancreatic islet beta cell antigens
InsB923, ChgA, and IAPP, we observed- a significant increase in both the frequency and
absolute number of tetramer positive CD4$^+$ T cells between 6 and 12 weeks of age (Fig.
2a, b). Mean fluorescence intensity (MFI) of tetramer staining also increased over this
interval (Fig. 2c), consistent with either higher TCR occupancy or enrichment of
higher-affinity clones. Among the three specificities, InsB9-23$^+$ CD4$^+$ T cells were the
most abundant at 6 weeks and increased further by 12 weeks of age. In contrast, ChgA
and IAPP-specific cells started at a lower frequency and expanded later time points,
suggesting a progressive broadening of beta cell antigen recognition within the CD4$^+$
compartment. However, consistent with Fife et al. findings, reveal that in spontaneous
diabetes, multiple autoreactive T cell populations infiltrate the islets, including ChgA-
specific BDC2.5, and IAPP-reactive BCD6.9 T cells. Among these, BDC2.5 T cells are
the most pathogenic, likely owing to the robust production of IFN-g and TNF-a [32] ,
whereas insulin-reactive T cells, despite their abundance, exhibit limited immuno-
pathogenicity [33]. The rise in tetramer-positive cells coincided with the period of elevated
plasma sLAG-3 and preceded the clinical onset of hyperglycemia (Fig. 1b, c, and 2d).
Linear regression analysis confirmed a significant positive association between plasma
sLAG-3 levels and the frequency of beta-cell antigen–specific CD4$^+$ T cells in our NOD
murine studies (Fig. 2e), linking systemic sLAG-3 shedding to antigen-specific T cell

activation and loss of tolerance to beta cells and their protein antigen during early
autoimmune progression.

Adoptive transfer of C6.6.9 CD4$^+$ T cells reveals early sLAG-3 induction and
beta-cell stress

To further study the relationship between beta-cell injury, beta-cell antigen–specific CD4$^+$
T cells activation/expansion and release of sLAG-3 release and beta-cell injury, we
adoptively transferred splenocytes from NOD.C6.6.9 TCR transgenic mice into
NOD.SCID recipients (Fig. 3a). Donor mice express the BDC-6.9 TCR but lack the
cognate IAPP-derived hybrid peptide, so their CD4$^+$ T cells remain naïve prior to transfer.
After adoptive transfer, recipient NOD.SCID mice undergo a prediabetic phase followed
by a rapid metabolic -shift with in, blood glucose began to rise by approximately 7 weeks
after transfer and exceeded 400 mg/dL by week 10 (Fig. 3b, c and ESM Fig. 2b). Plasma
sLAG-3 levels increased early following adoptive transfer, peaking between days 12-15
after adoptive transfer and then declining as diabetes developed (Fig. 3d; ESM Fig. 2a).
In line with the data from sLAG-3 secretion, circulating levels of *Ins2* mRNA showed a
similar early increase (Fig. 3e), indicating ongoing beta-cell stress during the same
interval. *Ins2* mRNA displayed a modest positive association with sLAG-3 (Fig. 3f), but
neither marker correlated positively with blood glucose, consistent with the idea that
substantial immune and beta-cell stress precede overt hyperglycemia (Fig. 3g). However,
sLAG-3 showed an inverse correlation with blood glucose when later time points were
included (Fig. 3h), indicating high sLAG-3 in circulation early during autoreactive T cell
activation followed by a decline as beta-cell mass is lost. Flow cytometry revealed
dynamic redistribution of donor C6.6.9 CD4$^+$ T cells- over time, with expansion in the
periphery followed by accumulation in the pancreas and secondary lymphoid organs (Fig.
3i). Histopathological analysis of pancreata from diabetic recipients showed marked signs
of immune cell infiltration and insulitis, with extensive islet destruction (Fig. 3j). Together,
these data confirm that beta cells antigen specific CD4$^+$ T cells are sufficient to drive
diabetes in this model and that early activation is accompanied by a transient rise in
plasma sLAG-3 and beta-cell *Ins2* mRNA.

Longitudinal assessment of antigen specific CD4$^+$ T cell activation and beta-cell antigen tetramer profiles in adoptively transferred mice.

We next characterized the activation state and antigen specificity of CD4$^+$ T cells in peripheral blood of adoptively transferred NOD.SCID mice every three days. Shortly after transfer, CD4$^+$ T cells exhibited a pronounced activation surge defined by increased CD25$^+$ and CD69$^+$ frequencies, peaking between days 3-9 following adoptive transfer (Fig. 4a; ESM Fig. 3a). Concomitantly, the frequency of ChgA$^+$, InsB9-23$^+$, and IAPP$^+$ tetramer-positive CD4$^+$ T cells transiently rose between 9 and 15-days post-transfer and then declined as cells redistributed into tissues (Fig. 4b; ESM Fig. 3b). LAG-3 expression on CD4$^+$ T cells also increased early and then decreased among adoptively transferred CD4 T cells, consistent with the pattern of soluble LAG-3 detected in plasma. Therefore, to address the mechanism underlying the emergence of these newly detected antigen-specific T cell populations, we performed immunofluorescence staining of pancreata at days 12 and 15 post adoptive transfer. We found that tetramer-positive cells, particularly -ChgA$^+$, were more abundant at day 12 than day 15 (Fig. 4c, d), mirroring the transient early peak observed in blood. At diabetes onset, analysis pancreatic tissue revealed that CD69 expression was significantly increased on pancreatic CD4$^+$ T cells compared with blood. In contrast, CD25 expression was relatively higher in circulating CD4+ T cells (Fig. 4e; ESM Fig. 3c). Tetramer positive CD4$^+$ T cells specific for ChgA, InsB9-23, and IAPP were most enriched in the pancreas compared with spleen and blood (Fig. 4f; ESM Fig. 3d), indicating preferential homing and/or retention of autoreactive CD4$^+$ T cells at the site of beta-cell destruction. Consistent with findings by Bettini and Vignali et al., T cells that are not specific for islet beta-cell antigens do not persist within the pancreatic islets and exit once cognate antigen recognition fails to occur [34].

Beta-cell antigen driven metabolic reprogramming and immune infiltration in the pancreas

To assess metabolic changes accompanying antigen driven T cell activation, we examine the immune infiltration in pancreas and quantified nutrient transporter expression on CD4$^+$ T cells isolated from blood, spleen, and pancreas at diabetes onset in adoptively

transferred NOD.SCID mice (Fig. 5a). Immunofluorescence confirmed dense infiltrates of

CD3$^+$ T cells along with B220$^+$ B cells and F4/80$^+$ macrophages in diabetic pancreata

compared with controls, together with loss of insulin positive beta-cells (Fig. 5b), suggests

that autoreactive T cells maybe directly and indirectly by activating macrophage the local

milieu to support B cell help and drive metabolic programs sustaining effector function

[35-37]. Flowcytometry profiling demonstrated that CD4$^+$ T cells from pancreatic tissue

expressed higher frequencies of Glut-1$^+$ and CD98$^+$ cells than those from spleen or blood

(Fig. 5c; ESM Fig. 5a, b), consistent with increased glucose and amino acid uptake in

tissue-resident effector cells indicating cell activation, proliferation, and anabolic

metabolism, while decreased expression suggests a resting or differentiated state. CD98

expression was high across compartments at diabetes onset, whereas CD71$^+$ CD4$^+$ T

cells were more frequent and showed higher MFI in the spleen than in blood or pancreas

(Fig. 5d; ESM Fig. 5a, b), suggesting ongoing proliferative expansion in secondary

lymphoid organs. These data indicate that antigen encounter in this model is associated

with both immune infiltration of the pancreas and tissue-specific metabolic activation of

CD4$^+$ T cells. Targeting these metabolic checkpoints may offer a strategy to selectively

dampen pathogenic T-cell responses in early T1D [38-40].

Broadening and epitope diversification of IAPP-specific CD4$^+$ T cells Co-

recognizing ChgA and InsB9-23 antigen.

To explore how antigen recognition evolves during disease, we re-analyzed adoptive

transfer flow-cytometry data by gating on CD4$^+$IAPP$^+$ T cells and then assessing

additional binding to ChgA and InsB9-23 tetramers over time (Fig. 6a; ESM Fig. 6a, b),

This approach highlights the underutilized potential of BCR and TCR repertoire dynamics

in diagnostic medicine. At baseline, very few CD4$^+$IAPP$^+$ cells co-stained with ChgA or

InsB9-23 tetramers (Fig. 6b). After transfer into NOD.SCID recipients, the proportion and

absolute number of CD4$^+$IAPP$^+$ChgA$^+$ and CD4$^+$IAPP$^+$InsB9-23$^+$ T cells in blood

increased progressively and reached maximal levels around day 12 post-adoptive

transfer (Fig. 6c, d). At diabetes onset, both dual positive subsets were enriched in

pancreatic tissue compared with spleen (Fig. 6e, f). These findings indicate that within the

IAPP-defined autoreactive pool, there is a broadening of tetramer-defined antigen

recognition over time, with preferential accumulation of multi-positive cells in the pancreas

during overt disease.

sLAG-3 patterns in human T1D, FDRs, and longitudinal DEWIT cohorts

We then examined whether the sLAG-3 patterns observed in mice were recapitulated in

human cohorts. In a cross-sectional analysis of individuals with type 1 diabetes up to six

years after diagnosis and age-matched healthy controls, serum sLAG-3 levels did not

differ significantly between groups (Fig. 7b). This suggests, as seen in the mouse

experiments, that by the time of established disease, the early activation signal captured

by sLAG-3 is not readily detectable. In contrast, when first-degree relatives were stratified

by autoantibody status, autoantibody negative and single autoantibody FDRs displayed

significantly higher sLAG-3 levels than multi-autoantibody FDRs, T1D subjects, and non-

FDR- controls, indicating early T-cell activation (Fig. 7c). Consistent with this, sLAG-3

levels showed no significant differences among T1D patients regardless of autoantibody

status (ESM Figure 7a), reinforcing that sLAG-3 is not a marker of late-stage disease

burden.

Longitudinal analysis of the DEWIT cohort using MSD-based high sensitive assays,

chosen for their suitability in repeated-measure analyses, showed that sLAG-3 levels

were generally higher in females than males across both FDR and general population

participants. (Fig. 7e, f). When participants were stratified by outcome, progressors

exhibited a gradual rise in sLAG3 over time, culminating in a distinct peak just before

diabetes diagnosis and a subsequent decline after onset. In contrast non-progressors

showed no consistent change (Fig. 7g, j; ESM Fig. 7b). The sLAG-3 levels also tend- to

decrease with increasing disease duration among individuals with type 1 dibetes (ESM

Fig. 7c). In FDRs, sLAG-3 concentrations were highest in autoantibody negative and

single-autoantibody positive individuals and progressively lower in those with multiple

autoantibodies, with the single 4-autoantibody-positive individual showing the lowest

value (Fig. 7h). This inverse relationship between sLAG-3 and autoantibody burden was

not observed in general-population subjects (Fig. 7i), suggesting that early sLAG-3

elevation is most apparent in genetic or familial at-risk individuals. Additionally, non-

diabetic individuals showed higher sLAG-3 concentrations compared with diabetic

subjects among GADA positive, IA2A positive and ZnT8A positive participants in first-

degree relatives, but no significant difference observed in general population. In contrast,

IAA-positive groups showed no significant differences in sLAG-3 between diabetic and

non-diabetic individuals, indicating that alterations in circulating sLAG-3 are associated

with diabetes status in specific autoantibody-defined subgroups (ESM Fig. 7d-g). Finally,

sLAG-3 levels were higher in subjects carrying high risk HLA alleles such as DRB1*_04_

(DR4) _and DQA1*_03:01-DQB1*03:02 (DQ8) compared with those lacking these risk

genotypes (Fig. 7k). This result support a linking between genetic susceptibility and the

magnitude of early immune activation.

sLAG-3 dynamics during COVID-19 vaccination and CVB infection

To test whether sLAG-3 reliably marks early T cell activation in nonautoimmune contexts,

we measured sLAG-3 following COVID-19 vaccination in healthy adults and during

Coxsackievirus B3 (CVB3) infection in mice. In vaccinated adults, plasma sLAG-3

(measured by Olink as NPX values) increased significantly by 10 days after vaccination

compared with pre-vaccination levels, with most donors showing a clear upward shift (Fig.

8a, b). This transient increase occurred in the expected window for vaccine induced T cell

responses and before the mature humoral response.

In CVB infected C57BL/6 mice, plasma sLAG-3 was minimal at baseline, start to increase

by day 3, and peaked between days 6-9 before declining toward baseline by day 30 (Fig.

8c, d). Plasma IFN-γ levels followed a similar pattern with a marked increase around day

6 (Fig. 8e). Flow cytometry revealed a concurrent expansion of IFNγ-producing CD4$^+$ and

CD8$^+$ T cells and increased expression of activation markers CD25 and CD69 in

peripheral blood (Fig. 8f-i; ESM Fig. 8). These data confirm that sLAG-3 is a sensitive and

dynamic marker of early T-cell activation in both vaccine and viral infection settings.

Discussion

Our goal was to determine whether soluble LAG-3 is useful serve as a biomarker of early

T cell activation relevant to type 1 diabetes pathogenesis. Across spontaneous and

adoptive transfer NOD mouse models, human FDR and longitudinal type 1 diabetes cohorts, and independent vaccine and viral infection systems. We consistently found that sLAG-3 concentrations in plasma/serum rises significantly early during T cell activation and declines as responses contract or as overt autoimmune diabetes develops. These observations support sLAG-3 as a candidate biomarker of early T cell activation and warrant further evaluation of its utility for staging autoimmune risk [16].

In NOD mice, plasma sLAG-3 increased between 6-12 weeks of age and returned to near baseline by 16 weeks, while blood glucose remained normal until around 16 weeks of age, when overt diabetes onset occurred. Consistent with our findings, Mathews and Ludlow, demonstrated that subtle blood glucose dysregulation precedes overt diabetes and extend extends this work by providing an immunological basis for these early changes [41]. Their results suggest that early blood glucose elevations reflect increased antigen-specific, immune-mediated beta-cell stress driven in part by BDC-2.5 reactive CD4$^+$ T cells rather than global beta-cell loss, positioning early blood glucose perturbations as a functional marker of immune activation preceding clinical diabetes [41]. Interestingly, we found that, beta-cell antigen specific CD4$^+$ T cells defined by InsB923, ChgA, and IAPP tetramers expanded over the same 6-12-week window in the same animal cohort. These findings correlated positively with sLAG-3 plasma concentrations. Furthermore, in the C6.6.9 adoptive transfer model, sLAG-3 and *Ins2* mRNA rose rapidly after adoptive transfer that peaked well before hyperglycemia. Histological and flow cytometric evidence of insulitis and donor CD4$^+$ T cell accumulation in the pancreas. Together, these results indicate that sLAG-3 is induced and persists during early antigen-driven activation of autoreactive CD4$^+$ T cells and that its peak precedes frank metabolic decompensation [42].

The adoptive transfer experiments also delineate how activation, tissue localization, and metabolism among autoreactive CD4$^+$ T cells evolve during disease progression. CD25 and CD69 expression on peripheral CD4$^+$ T cells peaked early after transfer, paralleling the sLAG-3 surge, and our detection of tetramer-positive cells. Subsequently, T cells accumulated in the pancreas, where CD69 expression increases. CD4$^+$ T cells, that had infiltrated the pancreas exhibit increased expression of Glut-1, CD98, and CD71

expression relative to blood and spleen, consistent with metabolic reprogramming of
effector cells in the inflamed target organ [43-49]. Within the IAPP specific pool, the
emergence and pancreatic localization of CD4$^+$IAPP$^+$ChgA$^+$ and CD4$^+$ IAPP$^+$ InsB2-93$^+$
subsets during diabetes indicated a progressive broadening of tetramer-defined antigen
recognition over time. Our results provide a coherent picture in which early antigen
encounter drives a transient systemic sLAG-3 signal, expansion and tissue homing of
antigen-specific CD4$^+$ T-cells and increased metabolic activity in the pancreas, long
before elevated blood glucose and failure of first phase insulin loss [33].

In humans, the cross sectional T1D vs non-diabetic control comparison showed no
difference in sLAG-3 levels up to six years post diagnosis, implying that the early
activation window identified in mice is not captured when sampling individuals with
established disease. In contrast, FDRs at earlier disease stages displayed a distinct
pattern: autoantibody negative and single autoantibody FDRs had higher sLAG-3 than
multi-autoantibody positive FDRs, T1D subjects, and non-FDR controls. Thus sLAG-3
declined with increasing autoantibody burden. Longitudinal data from the DEWIT cohort
further showed that progressors, but not non-progressors, exhibited a gradual rise in
sLAG-3 culminating in a peak immediately prior to diagnosis that declined thereafter.
sLAG-3 tended to be higher in individuals carrying higher risk type 1 diabetes HLA
genotypes. Taken together, these human data are consistent with a model in which sLAG-
3 in blood serving as a biomarker of autoimmunity during a relatively narrow early stage
of immune activation that precedes or is co-incident with a broadening of autoantibody
and auto-reactive T cell expansion prior to clinical onset in genetically susceptible
individuals [50, 51].

The current human dataset has important limitations that temper how strongly sLAG-3
can be positioned as a "surrogate marker before antibodies." First, most of the human
analyses are cross-sectional. Even in the DEWIT cohort, the number of progressors with
dense longitudinal sampling is limited in. numbers. We therefore cannot yet systematically
align the timing of sLAG-3 concentration with individual seroconversion events to prove
that sLAG-3 consistently rises before the first islet autoantibody appears. Second, sLAG-
3 levels were measured using different assay platforms and in relatively small subgroups,

which constrains our ability to define robust thresholds or evaluate test characteristics
such as sensitivity, specificity, and predictive value. Third, the COVID-19 vaccine and
CVB infection experiments highlight that increases in sLAG-3 in blood occurs general
following T cell activation and may not be specific to autoimmunity. Eelevations in at-risk
individuals will likely need to be interpreted in the context of recent infections,
vaccinations, or other immune stimuli.

The COVID19 vaccination and CVB infection data nonetheless provide important context.
In both settings, sLAG-3 rose rapidly during the effector phase of the immune response,
coincided with increases in IFN-γ and T-cell activation but markers and returned to
baseline as the response resolved. These observations support the idea that sLAG-3 is
a sensitive and reversible readout of early T-cell activation. In type 1 diabetes, this
suggests that transient sLAG-3 elevations in FDRs are biologically plausible and that
serial measurements may be more informative than single time points, especially when
integrated with autoantibody profiles, islet or beta-cell antigen–specific T cell along with
detection of biorepositories, and plasma circulating insulin mRNA assessments in the
clinical context provide additional important data that may help distinguish autoimmunity
from other T cell-mediated immune responses.

Overall, our findings support as a cautious interpretation that sLAG-3 is a promising
biomarker of early T cell activation, detectable in some autoantibody-negative and single
autoantibody positive first-degree relatives and declining with increasing autoantibody
burden and after diabetes onset. When integrated with beta-cell antigen specific tetramer
profiling and markers of beta-cell stress such as circulating *Ins2* mRNA, sLAG-3 may help
define an early immunological phase characterized by autoreactive CD4[+] T cell activation
can be detected before substantial beta-cell loss is clinically evident. Larger, deeply
phenotype longitudinal cohorts will need to be tested to determine if assessment of sLAG-
3, autoantibody reactivity, markers of beta cell stress (circulating insulin mRNA in blood)
and T cell phenotypes comprise surrogate markers of type 1 diabetes risk or progression
and whether these markers are predictive of patients that go on to develop type 1 diabetes
clinically to determine how frequently it precedes seroconversion at the individual level.

Conclusion

In conclusion, our findings support sLAG-3 as a candidate early biomarker of T cell activation in type 1 diabetes, with evidence that sLAG-3 may be elevated before or at the earliest stages of islet autoimmunity in individual development in some at-risk individuals. We propose that tracking of beta-cell antigen specific CD4$^+$ T cell, circulating *Ins2* mRNA measurements, metabolic profiling, and analyses of B and T cell epitope spreading in conjunction with sLAG-3 reflects an early immunological state associated with autoreactive T-cell diversification rather than late-stage beta-cell loss. Nevertheless, larger longitudinal studies aligned to seroconversion are required to validate sLAG-3 as a surrogate marker of pre-autoantibody autoimmune disease and to define its utility for early risk stratification and therapeutic intervention.

Data and Resource Availability

The data sets generated and/or analyzed during the current study are available from the corresponding author on reasonable request.

Acknowledgements The authors thank Dr. Dario Vignali lab at University of Pittsburgh for providing the secondary antibody for the mouse sLAG-3 ELISA. We also thank to Dale Long - Facility Manager at NIH Tetramer Core Facility at Emory for providing tetramers and the Center for Diabetes and Metabolic Diseases for access to its instrumental facilities.

Funding This study was funded by the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) under project 7R01DK132583, and Breakthrough Type 1 diabetes research Strategic Research Agreement (SRA) funds, under Project 1-PNF-2025-1642-A-N to Dr. Jon D. Piganelli, and R01DK129287 to Jason M. Spaeth. Farooq Syed was supported by a Breakthrough T1D (formerly JDRF) CDA (5-CDA-2022-1176-A-N and (CDA)/3-CDA-2025-1692-A-N) and Thomas Beatson Foundation (2024-010). This study is also supported by a NIH-R21 (1R21AI193399) award to Jon D. Piganelli and Farooq Syed.

Contribution statement J.D.P., S.R., F.S., and C.S. conceptualized the study. S.R. and
J.D.P. designed all experiments; S.R. and M.L.P. performed experiments. S.R. analyzes
all data and prepares all figures. F.S and J.R performed experiments on insulin mRNA,
and M.E.S. conducted sLAG-3 assays in Covid-vaccinated participants. S. R and A.D.S.
analyzed all confocal images. W.H help to collect the DEWIT cohort samples, A.T.,
E.S.H., E.M., and K.P.J. collect human plasma sample from Stanford Biobank, J.S.
contributed to discussions and study design. S.R. and J.D.P. drafted the manuscript. S.F.,
M.L.P., J.S., A.D.S., K.P.J., J.S., and C.S. editing and finalizing it. All authors have no
conflicts of interest.

 Figure Legends

Fig. 1 Plasma sLAG-3 correlates with disease incidence in NOD mice. (**a**) Experimental
design for longitudinally monitoring blood glucose and plasma sLAG-3 level from female
NOD mice. (**b**) Plasma sLAG-3 levels in NOD mice at different ages. (**c**) Longitudinal
monitoring of blood glucose levels. (**d**) Correlation between plasma sLAG-3 levels and
diabetes incidence in NOD mice. Data are presented as mean ± SEM (n=10-12 per
group). Statistical significance was determined using one-way ANOVA followed by
Bonferroni post-hoc testing; *p<0.05*.

Fig. 2 Beta-cell antigen-specific tetramer expression in CD4$^+$ T cells corelate with disease
incidence in NOD mice. (**a**) Percentage of CD4$^+$ tetramer positivity T cells for three islet
antigens (InsB9-23, ChgA, and IAPP) (**b**) Absolute numbers of tetramer-positive CD4$^+$ T
cells. (**c**) Overlay of the MFI histograms by flow cytometry analysis of tetramer expression
in CD4$^+$ cells (**d**) Overlay representation of plasma sLAG-3 and CD4$^+$ tetramer
expression. (**e**) Linear regression analysis of plasma sLAG-3 levels and islet antigen-
specific tetramer positive CD4$^+$ T cell frequencies in CD4$^+$ T cells in NOD mice during
diabetic progression. Data are presented as mean ± SEM (n=10-12 in the study cohort)
Statistical significance was determined using one-way ANOVA with Bonferroni post-hoc
testing; *p<0.05*.

Fig. 3 Antigen-specific CD4$^+$ T cell activation and immune modulation during diabetes
progression following adoptive transfer and immune modulation. (**a**) Schematic overview
of the experimental design for adoptive transfer and tracking activation of naive C6.6.9
TCR-Tg CD4$^+$ T cells in NOD.Scid recipient mice. (**b**) Longitudinal monitoring of blood
glucose levels post-transfer. (**c**) Kaplan-Meier survival analysis of diabetes onset
following T cell transfer. (**d**) Plasma sLAG-3 levels. (**e**) Plasma *Insulin2 (Ins2)* mRNA
levels measured at defined time points post-transfer. (**f-h**) Linear regression analyses
relating plasma sLAG-3 levels, plasma Ins2-mRNA, and blood glucose levels during
diabetes progression. (**i**) Frequency of circulating C6.6.9 TCR-Tg CD4$^+$ T cells in
peripheral blood of recipient mice over time following adoptive transfer. (**j**) Representative

histological images showing progressive insulitis and islet destruction in pancreatic tissue
from NOD.Scid recipient mice. Data are presented as mean ± SEM (n=10).

Fig. 4 Longitudinal phenotypic characterization of T cell activation markers in recipients
following adoptive transfer. (a) Frequency of T-cell activation markers (CD25, CD69, and
surface LAG-3). (b) Frequency of β-cell antigen-specific tetramer-positive CD4$^+$ T cells in
peripheral blood over time. (c) Immunofluorescence staining of pancreatic sections on
days 12 and 15 post-transfer showing infiltration of ChgA, InsB9-23, and IAPP-reactive T
cells. (d) Quantification of immunofluorescence MFI for ChgA$^+$, InsB9-23$^+$, IAPP$^+$ cells
from pancreatic sections. (e, f) Percentage of different activation markers (CD25, CD69,
LAG-3) and β-cell antigen-specific tetramers (ChgA, InsB9-23, IAPP) expression on T
cells that had migrated or infiltrated into blood, pancreases and spleen at the time of
diabetes. All those values are represented as mean ± SEM (n=4-10). P <0.05 is significant
and P>0.05 not significant. Statistics were done by one-way ANOVA with multiple
comparisons, followed by the Bonferroni post hoc test.

Fig. 5 Metabolic profiling and immune infiltration in the pancreas following adoptive
transfer of C6.6.9 T cells into NOD.Scid recipients. (**a**) Schematic overview of the
experimental design used to assess metabolic status and immune infiltration in the
pancreas after adoptive transfer. (**b**) Immunofluorescence staining of pancreatic sections
from control and diabetic NOD.Scid mice, showing CD3$^+$, F4/80$^+$, and B220$^+$ immune cell
infiltration following BDC6.9 CD4$^+$ T cell transfer. (**c, d**) Absolute number and MFI of
metabolic markers (CD71, CD98, and Glut-1) expressed on CD4$^+$ T cells isolated from
adoptively transfer NOD.Scid mice peripheral blood, pancreas, and spleen at the time of
diabetes onset. All data are presented as mean ± SEM (n = 4-10). Statistical significance
was determined using one-way ANOVA with multiple comparisons, followed by Bonferroni
post hoc test. P < 0.05 represents significant.

Fig. 6. Phenotypic characterization of TCR rearrangement in IAPP-specific CD4$^+$ T cells
during diabetes progression in NOD.Scid recipient mice. (a) Schematic overview of
experimental design for monitoring the epitope spreading following adoptive transfer of
C6.6.9 TCR-Tg mouse CD4$^+$ T cells into NOD.Scid recipient mice. (b) Densitometry flow

cytometry plot representing baseline expression status of ChgA and InsB23-9 positive on C6.6.9 TCR-Tg mouse CD4$^+$IAPP$^+$ cells. (c) Frequency and absolute numbers of CD4$^+$IAPP$^+$ChgA$^+$ and (d) CD4$^+$IAPP$^+$InsB9-23$^+$ T cells in peripheral blood over time. (e, f) Frequency and absolute number of CD4$^+$IAPP$^+$ChgA$^+$ and CD4$^+$IAPP$^+$InsB$^+$ cells at the time of diabetes onset were assessed in pancreas, and spleen. All values are presented as mean ± SEM (n =5–6). Statistical significance was determined P < 0.05 represents significant using one-way ANOVA with multiple comparisons, followed by the Bonferroni post hoc test.

Fig. 7 Longitudinal and cross-sectional analyses reveal early sLAG-3 elevation prior to autoantibody expansion and clinical onset of type 1 diabetes. (a) Experimental design for sample collection. Plasma was collected from healthy controls (n=28), type 1 diabetes patients (n=39), and FDRs with varying numbers of autoantibodies (Ab-, n=37 Ab1+, n=27; Ab2+, n=29; Ab3+, n=5). (b) Serum sLAG-3 levels in individuals with type 1 diabetes up to six years post-diagnosis was comparable to healthy controls, suggesting that heterologous immune responses may mask earlier elevations. (c) Autoantibody-negative and single-autoantibody FDRs show higher sLAG-3 levels than multiple-autoantibody individuals, type 1 diabetes subjects, and controls, indicating early T cell activation. (d) Experimental design for measurement of plasma sLAG-3 levels using MSD U-PLEX Assays in DEWIT and Stanford Cohorts. (e, f) DEWIT longitudinal cohort show higher sLAG-3 expression in females than males across both cohorts. (g) Progressor vs. non-progressor trajectories in the DEWIT cohort. Progressors display a gradual increase in sLAG-3 preceding diagnosis, whereas non-progressors show no change. (h, i) sLAG-3 levels stratified by autoantibody category in FDRs and the general population. (j) Longitudinal sLAG-3 profiling in DEWIT progressors shows a distinct peak immediately before diabetes onset followed by a sharp decline after diagnosis. (k) HLA genotype analysis demonstrating higher sLAG-3 expression in individuals carrying high-risk alleles. Statistical significance was determined using one-way ANOVA with multiple comparisons, followed by the Bonferroni post hoc test, *P < 0.05 significant.

Fig. 8 sLAG-3 as an early and sensitive marker of T cell activation during vaccination and viral infection C57BL/6 Mice. (a) Schematic illustration of before and 10 days after

COVID-19 vaccination, indicating rapid T-cell activation preceding antibody development.

(b) Normalized protein expression (NPX) of sLAG-3 over time in before and after 10 days

of Covid vaccination from plasma. Each donors' samples are connected by a line. (c)

Schematic representation of time-course analysis of plasma analytes in-vivo CVB-

infected C57BL/6 mice. (d) Plasma sLAG-3 levels form C57BL/6 mice infected with CVB.

(e) Plasma IFN-γ concentrations increase after infection, with a marked rise at day 6,

reflecting concurrent immune activation. (f, g) Flow cytometry reveals time-dependent

increases in IFN-γ-producing $CD4^+$ and $CD8^+$ T cells in peripheral blood following CVB

infection. (h, I) Expression of T cell activation markers CD25 and CD69. All values are

presented as mean ± SEM (n=8). Statistical significance was determined using one-way

ANOVA, followed by the Bonferroni post hoc test. $*P < 0.05$ represent significant.

References

[1] Atkinson MA, Mirmira RG (2023) The pathogenic "symphony" in type 1 diabetes: A disorder of the immune system, beta cells, and exocrine pancreas. Cell Metab 35(9): 1500-1518. 10.1016/j.cmet.2023.06.018

[2] Burrack AL, Martinov T, Fife BT (2017) T Cell-Mediated Beta Cell Destruction: Autoimmunity and Alloimmunity in the Context of Type 1 Diabetes. Front Endocrinol (Lausanne) 8: 343. 10.3389/fendo.2017.00343

[3] Houeiss P, Luce S, Boitard C (2022) Environmental Triggering of Type 1 Diabetes Autoimmunity. Front Endocrinol (Lausanne) 13: 933965. 10.3389/fendo.2022.933965

[4] Roy S, Pokharel P, Piganelli JD (2024) Decoding the immune dance: Unraveling the interplay between beta cells and type 1 diabetes. Mol Metab 88: 101998. 10.1016/j.molmet.2024.101998

[5] Maahs DM, West NA, Lawrence JM, Mayer-Davis EJ (2010) Epidemiology of type 1 diabetes. Endocrinol Metab Clin North Am 39(3): 481-497. 10.1016/j.ecl.2010.05.011

[6] Group DP (2006) Incidence and trends of childhood Type 1 diabetes worldwide 1990-1999. Diabet Med 23(8): 857-866. 10.1111/j.1464-5491.2006.01925.x

[7] Popoviciu MS, Kaka N, Sethi Y, Patel N, Chopra H, Cavalu S (2023) Type 1 Diabetes Mellitus and Autoimmune Diseases: A Critical Review of the Association and the Application of Personalized Medicine. J Pers Med 13(3). 10.3390/jpm13030422

[8] Mallone R, Bismuth E, Thivolet C, et al. (2025) Screening and care for preclinical stage 1-2 type 1 diabetes in first-degree relatives: French expert position statement. Diabetes Metab 51(1): 101603. 10.1016/j.diabet.2024.101603

[9] Steck AK, Rewers MJ (2011) Genetics of type 1 diabetes. Clin Chem 57(2): 176-185. 10.1373/clinchem.2010.148221

[10] Thakkar S, Chopra A, Nagendra L, Kalra S, Bhattacharya S (2023) Teplizumab in Type 1 Diabetes Mellitus: An Updated Review. touchREV Endocrinol 19(2): 22-30. 10.17925/EE.2023.19.2.7

[11] Kokori E, Olatunji G, Ogieuhi IJ, et al. (2024) Teplizumab's immunomodulatory effects on pancreatic beta-cell function in type 1 diabetes mellitus. Clin Diabetes Endocrinol 10(1): 23. 10.1186/s40842-024-00181-w

[12] Li N, Wang Y, Forbes K, et al. (2007) Metalloproteases regulate T-cell proliferation and effector function via LAG-3. EMBO J 26(2): 494-504. 10.1038/sj.emboj.7601520

[13] Scherm MG, Wyatt RC, Serr I, Anz D, Richardson SJ, Daniel C (2022) Beta cell and immune cell interactions in autoimmune type 1 diabetes: How they meet and talk to each other. Mol Metab 64: 101565. 10.1016/j.molmet.2022.101565

[14] Andrews LP, Marciscano AE, Drake CG, Vignali DA (2017) LAG3 (CD223) as a cancer immunotherapy target. Immunological reviews 276(1): 80-96. 10.1111/imr.12519

[15] Anderson MS, Bluestone JA (2005) The NOD mouse: a model of immune dysregulation. Annu Rev Immunol 23: 447-485. 10.1146/annurev.immunol.23.021704.115643

[16] Cieplińska K, Niedziela E, Jagielska E, Pałyga I, Słuszniak A, Kowalska A (2025) Elevated Levels of sLAG-3 as a Possible Biomarker in Graves' Disease With and Without

Thyroid Eye Disease: A Prospective Observational Case-Control Study. Medicina
(Kaunas) 61(9). 10.3390/medicina61091664

[17] Anderson Ana C, Joller N, Kuchroo Vijay K (2016) Lag-3, Tim-3, and TIGIT: Co-
inhibitory Receptors with Specialized Functions in Immune Regulation. Immunity 44(5):
989-1004. 10.1016/j.immuni.2016.05.001

[18] Ruffo E, Wu RC, Bruno TC, Workman CJ, Vignali DAA (2019) Lymphocyte-activation
gene 3 (LAG3): The next immune checkpoint receptor. Semin Immunol 42: 101305.
10.1016/j.smim.2019.101305

[19] Botticelli A, Zizzari IG, Scagnoli S, et al. (2021) The Role of Soluble LAG3 and Soluble
Immune Checkpoints Profile in Advanced Head and Neck Cancer: A Pilot Study. J Pers
Med 11(7). 10.3390/jpm11070651

[20] SYED F, MAIER BF, ANDERSON-BAUCUM E, MASTRACCI TL, EVANS-MOLINA C,
MIRMIRA R (2020) 82-OR: Exosomes from Human ß-Cells Incorporate Stress Granule
Components and May Serve as Biomarkers of ß-Cell Stress in Type 1 Diabetes. Diabetes
69(Supplement_1). 10.2337/db20-82-OR

[21] Dekkers MC, Lambooij JM, Pu X, et al. (2024) Extracellular vesicles derived from
stressed beta cells mediate monocyte activation and contribute to islet inflammation.
Front Immunol 15: 1393248. 10.3389/fimmu.2024.1393248

[22] Dekkers MC, Pu X, Enciso-Martinez A, Zaldumbide A (2024) Beta-Cell-Derived
Extracellular Vesicles: Mediators of Intercellular Communication in the Islet
Microenvironment in Type 1 Diabetes. Cells 13(23). 10.3390/cells13231996

[23] Kracht MJ, van Lummel M, Nikolic T, et al. (2017) Autoimmunity against a defective
ribosomal insulin gene product in type 1 diabetes. Nat Med 23(4): 501-507.
10.1038/nm.4289

[24] van Tienhoven R, Jansen D, Park M, et al. (2024) Induction of islet autoimmunity to
defective ribosomal product of the insulin gene as neoantigen after anti-cancer
immunotherapy leading to autoimmune diabetes. Front Immunol 15: 1384406.
10.3389/fimmu.2024.1384406

[25] van Tienhoven R, O'Meally D, Scott TA, et al. (2025) Genetic protection from type 1
diabetes resulting from accelerated insulin mRNA decay. Cell 188(9): 2407-2416 e2409.
10.1016/j.cell.2025.02.018

[26] Anand V, Li Y, Liu B, et al. (2021) Islet Autoimmunity and HLA Markers of
Presymptomatic and Clinical Type 1 Diabetes: Joint Analyses of Prospective Cohort
Studies in Finland, Germany, Sweden, and the U.S. Diabetes Care 44(10): 2269-2276.
10.2337/dc20-1836

[27] Gong Q, Sharma M, Kuan EL, et al. (2024) Longitudinal Multi-omic Immune Profiling
Reveals Age-Related Immune Cell Dynamics in Healthy Adults. bioRxiv.
10.1101/2024.09.10.612119

[28] Aggarwal V, Workman CJ, Vignali DAA (2023) LAG-3 as the third checkpoint inhibitor.
Nature Immunology 24(9): 1415-1422. 10.1038/s41590-023-01569-z

[29] Camisaschi C, Casati C, Rini F, et al. (2010) LAG-3 expression defines a subset of
CD4(+)CD25(high)Foxp3(+) regulatory T cells that are expanded at tumor sites. J Immunol
184(11): 6545-6551. 10.4049/jimmunol.0903879

[30]   Marré ML, Profozich JL, Coneybeer JT, et al. (2016) Inherent ER stress in pancreatic
islet β cells causes self-recognition by autoreactive T cells in type 1 diabetes. J
Autoimmun 72: 33-46. 10.1016/j.jaut.2016.04.009

[31]   Catrina AM, Popa MA, Văcaru AM, Fenyo IM (2021) Inflammatory status of the
pancreas in NOD mice that do not develop overt diabetes. Rom J Morphol Embryol 62(1):
109-115. 10.47162/rjme.62.1.10

[32]   Piganelli JD, Flores SC, Cruz C, et al. (2002) A Metalloporphyrin-Based Superoxide
Dismutase Mimic Inhibits Adoptive Transfer of Autoimmune Diabetes by a Diabetogenic
T-Cell Clone. Diabetes 51(2): 347-355. 10.2337/diabetes.51.2.347

[33]   Mitchell JS, Spanier JA, Dwyer AJ, et al. (2024) CD4+ T cells reactive to a hybrid
peptide from insulin-chromogranin A adopt a distinct effector fate and are pathogenic in
autoimmune diabetes. Immunity 57(10): 2399-2415.e2398.
https://doi.org/10.1016/j.immuni.2024.07.024

[34]   Bettini M, Vignali DA (2011) T cell-driven initiation and propagation of autoimmune
diabetes. Curr Opin Immunol 23(6): 754-760. 10.1016/j.coi.2011.10.002

[35]   Clark M, Kroger CJ, Ke Q, Tisch RM (2020) The Role of T Cell Receptor Signaling in the
Development of Type 1 Diabetes. Front Immunol 11: 615371.
10.3389/fimmu.2020.615371

[36]   Hussain S, Delovitch TL (2005) Dysregulated B7-1 and B7-2 expression on nonobese
diabetic mouse B cells is associated with increased T cell costimulation and the
development of insulitis. J Immunol 174(2): 680-687. 10.4049/jimmunol.174.2.680

[37]   Tian J, Zekzer D, Hanssen L, Lu Y, Olcott A, Kaufman DL (2001) Lipopolysaccharide-
activated B cells down-regulate Th1 immunity and prevent autoimmune diabetes in
nonobese diabetic mice. J Immunol 167(2): 1081-1089. 10.4049/jimmunol.167.2.1081

[38]   Qi Q, Cavanagh MM, Le Saux S, et al. (2016) Diversification of the antigen-specific T
cell receptor repertoire after varicella zoster vaccination. Sci Transl Med 8(332):
332ra346. 10.1126/scitranslmed.aaf1725

[39]   Straub A, Grassmann S, Jarosch S, et al. (2023) Recruitment of epitope-specific T cell
clones with a low-avidity threshold supports efficacy against mutational escape upon re-
infection. Immunity 56(6): 1269-1284.e1266.
https://doi.org/10.1016/j.immuni.2023.04.010

[40]   Yi J, Miller AT, Archambault AS, et al. (2022) Antigen-specific depletion of CD4(+) T
cells by CAR T cells reveals distinct roles of higher- and lower-affinity TCRs during
autoimmunity. Sci Immunol 7(76): eabo0777. 10.1126/sciimmunol.abo0777

[41]   Ize-Ludlow D, Lightfoot YL, Parker M, et al. (2011) Progressive erosion of β-cell
function precedes the onset of hyperglycemia in the NOD mouse model of type 1
diabetes. Diabetes 60(8): 2086-2091. 10.2337/db11-0373

[42]   Syed F, Krishnan P, Chang G, et al. (2023) beta Cell microRNAs Function as Molecular
Hubs of Type 1 Diabetes Pathogenesis and as Biomarkers of Diabetes Risk. bioRxiv.
10.1101/2023.06.15.545170

[43]   Guerrero JA, Klysz DD, Chen Y, et al. (2024) GLUT1 overexpression in CAR-T cells
induces metabolic reprogramming and enhances potency. Nat Commun 15(1): 8658.
10.1038/s41467-024-52666-y

[44] Poznanski SM, Singh K, Ritchie TM, et al. (2021) Metabolic flexibility determines
human NK cell functional fate in the tumor microenvironment. Cell Metab 33(6): 1205-
1220 e1205. 10.1016/j.cmet.2021.03.023
[45] Salzberger W, Martrus G, Bachmann K, et al. (2018) Tissue-resident NK cells differ in
their expression profile of the nutrient transporters Glut1, CD98 and CD71. PLoS One
13(7): e0201170. 10.1371/journal.pone.0201170
[46] Jessop MK (1990) Homoeopathy. Vet Rec 126(20): 514
[47] Delmastro-Greenwood M, Hughan KS, Vitturi DA, et al. (2015) Nitrite and nitrate-
dependent generation of anti-inflammatory fatty acid nitroalkenes. Free Radical Biology
and Medicine 89: 333-341. https://doi.org/10.1016/j.freeradbiomed.2015.07.149
[48] Previte DM, Martins CP, O'Connor EC, et al. (2019) Lymphocyte Activation Gene-3
Maintains Mitochondrial and Metabolic Quiescence in Naive CD4(+) T Cells. Cell Rep
27(1): 129-141.e124. 10.1016/j.celrep.2019.03.004
[49] Previte DM, O'Connor EC, Novak EA, Martins CP, Mollen KP, Piganelli JD (2017)
Reactive oxygen species are required for driving efficient and sustained aerobic
glycolysis during CD4+ T cell activation. PLoS One 12(4): e0175549.
10.1371/journal.pone.0175549
[50] Sartoris S, Del Pozzo G (2024) Exploring the HLA complex in autoimmunity: From the
risk haplotypes to the modulation of expression. Clinical Immunology 265: 110266.
https://doi.org/10.1016/j.clim.2024.110266
[51] Sartoris S, Del Pozzo G (2024) Exploring the HLA complex in autoimmunity: From the
risk haplotypes to the modulation of expression. Clin Immunol 265: 110266.
10.1016/j.clim.2024.110266

Fig. 1

a

NOD Mice

longitudinal

4   6   8   12   16

Monitoring Blood Glucose Level

Measuring sLAG-3 by ELISA

Tetramer Expression

b

0.1374

<0.0001

<0.0001

0.0154

sLAG3 (µg/ml)

Age (weeks)
4   6   8   12   16

c

Blood Glucose (mg/dL)

Age (weeks)
6   8   14   16   20   24   32

d

****
****
****
****
*
****

sLAG3 (µg/ml)

% T1D Free

Age (weeks)
0   5   10   15   20   25

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7