The genome of an aneuploid cultivar facilitates elucidating the genetic bases of trait polymorphisms in *Phalaenopsis* orchids

Ruidong Jia[1‡]; Xinyi Zuo[2,3‡]; Liting Zou[2‡]; Li Wang[1]; Letian Lin[2]; Zhiqing Wang[2,9]; Yuexia Zhang[2]; Xingyu Chen[1]; Fanrui Meng[8]; Hantang Huang[10]; Lan Lan[11]; Zhimei Li[2]; Fei Wang[2,3]; Yuxuan Jin[2,12]; Hongyan Shan[4,5]; Rui Zhang[3]*; Hongzhi Kong[4,5,6]*; Peipei Wang[2,7]*

[1] State Key Laboratory of Vegetable Biobreeding, Key Laboratory of Biology and Genetic Improvement of Flower Crops (North China), Ministry of Agriculture and Rural Affairs, Institute of Vegetables and Flowers, Chinese Academy of Agricultural Sciences, Beijing 100081, China
[2] Kunpeng Institute of Modern Agriculture at Foshan, Foshan, Guangdong 528200, China
[3] College of Horticulture, Northwest A&F University, Yangling, Shaanxi 712100, China
[4] State Key Laboratory of Plant Diversity and Specialty Crops, Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences, Beijing 100093, China
[5] China National Botanical Garden, Beijing 100093, China.
[6] University of Chinese Academy of Sciences, Beijing 100049, China.
[7] Guangdong Huabo Ecological Industry Co., Ltd., Foshan, Guangdong 528241, China
[8] Bioinfo-Reach Bioinformatics Service, Guangzhou, Guangdong 510663, China
[9] College of Plant Protection, South China Agricultural University, Guangzhou, Guangdong 510642, China
[10] College of Horticulture, China Agricultural University, Beijing 100193, China
[11] State Agricultural Biotechnology Centre (SABC), College of Science, Health, Engineering and Education, Murdoch University, Perth, Western Australia, 6150, Australia
[12] Shanghai Jiaotong University, Shanghai, 200240, China

‡ These authors contributed equally to this work

Authors for correspondence:
Peipei Wang
wangpeipei02@caas.cn

Hongzhi Kong
Email: hzkong@ibcas.ac.cn

Rui Zhang
ruizhang@nwafu.edu.cn

Abstract

As an important ornamental crop, *Phalaenopsis* orchids exhibit extraordinary diversity in floral traits with unclear genetic bases. To address this problem, we assemble a haplotype-resolved, chromosome-level genome for an aneuploid cultivar 'Santiago'. We reveal pervasive variation in the number and sequence among homoeologous genes (HGs) and demonstrate proportional expression dosage effects of both chromosome and HG counts. This genome facilitates deciphering the genetic variation that contribute to trait polymorphisms in a hybrid population, some of which are also observed among HGs in *P*. 'Santiago'. Specifically, nucleotide polymorphisms in a region that includes two tandemly arranged *PsAGL6-2* genes account for lip morphology variation. Presence/absence of *PsMYB12* genes determines presence/absence of venation-associated stripes. The *b*-type HG of *PsMYB2* is associated with the pink color of sepals/petals. *PsMYBx1* represses anthocyanin biosynthesis in multiple regions, in addition to inter-spot regions. Our study advances the genetic understanding of trait diversity in *Phalaenopsis*.

Main

Phalaenopsis orchids are a major ornamental crop prized for their beauty and long flowering. Thanks to the rapid development of tissue culture technology, the production and market scale of *Phalaenopsis* have expanded significantly. Cross-breeding has yielded numerous cultivars to meet the growing market demand, which display extraordinary diversity in horticultural traits, particularly in floral size, scent, organ morphology, and color patterning. However, the growing homogenization caused by cross-breeding necessitates the urgent integration of advanced biotechnologies—such as genome editing, molecular- or genomics-assisted breeding—to develop novel, high-value cultivars. This requires comprehensive understanding of the genetic mechanisms underlying key horticultural traits and their diversity.

Key genes regulating specific traits have been identified, such as floral scent[1,2], color[3], venation-associated stripes[3], spot[3], petal size[4], and cuticular wax[5]. The genetic bases of the presence/absence (P/A) variation of floral scent[1,6] and the variation in spot/patch patterns[3,7,8] have also been studied. However, except for these two traits, the genetic mechanisms underlying variation in many other important traits remain largely uncharacterized. For example, while the background pink color on sepals/petals and venation-associated stripes are known to be regulated by *PeMYB2* and *PeMYB12*, respectively[3], the specific genetic variation responsible for the P/A of these two traits across different cultivars are unclear. Similarly, the extensive polymorphisms in lip morphology raises the question of whether and what specific changes of its key identity regulator, *PeAGL6-2* (*AGAMOUS-LIKE 6-2*)[9], lead to the transformation into petal-like organs in 'Bigfoot' cultivars. To decipher the genetic bases of trait diversity, it is essential to associate allelic polymorphisms—including single nucleotide polymorphism (SNP), insertion and deletion (indel), copy number variation (CNV), and structural variant (SV)—with trait variation. A major bottleneck in *Phalaenopsis* is the lack of a chromosome-level reference genome; the existing genomes for *P. equestris* and *P. aphrodite* are only at the scaffold levels[10,11]. Consequently, the full scope of these allelic polymorphisms and their contributions to trait diversity is still poorly understood.

In this study, we assemble a haplotype-resolved, chromosome-level genome for a highly heterozygous aneuploid cultivar, *P.* 'Santiago' (ANTHURA BV, with US patent US20150047084P1, hereafter *P.* sa, **Fig. 1A**). Analysis of this genome reveals extensive variation in the number and sequence among homoeologous genes (HGs), particularly those

of genes known to regulate key floral traits. To understand the genetic mechanisms

underlying trait diversity in *Phalaenopsis*, we generate a hybrid population (*P*. '1071-13' × *P*.

'CH1804-01', named as H80) which exhibits extensive polymorphisms in lip morphology

and floral color patterning (**Fig. S1**). By analyzing the whole genome resequencing data of

this population using *P*. sa genome as the reference, we decipher genetic variation that

account for these trait polymorphisms. Some of these genetic variation are also observed

among HGs in the *P*. sa genome. In addition, we demonstrate multiple functions of *PsMYBx1*

in repressing anthocyanin biosynthesis in diverse regions, which exhibits variation in HG

count across cultivars as well. Our study highlights the usefulness of a single genome with

high heterogeneity in elucidating the genetic bases of key trait diversity. It also facilitates

marker-assisted breeding in *Phalaenopsis* by offering critical insights for molecular marker

development.

Results

89 **Results**

The genome assembly and annotation of *P*. sa

The genome assembly and annotation of *P*. sa

The whole genome size of *P*. sa was estimated to be approximately 4.1 Gb by flow cytometry,

and 3.7 and 3.3 Gb by *k*-mer-based analysis using Genomic Character Estimator and

GenomeScope2, respectively (**Fig. S2A,B**). *K*-mer frequency analysis suggested an allo-

triploid genome (**Fig. S2C**), while karyotype analysis revealed 70 chromosomes per cell,

indicating likely aneuploidy (**Fig. S2D**) since a haplotype (1x) is supposed to contain 19

chromosomes[11]. The final assembly (**Methods**) comprised 70 pseudo-chromosomes, with 6 and

13 homoeologous chromosome groups (HChrGs) representing triploid (3x, set3) and

tetraploid (4x, set4) sets, respectively (**Fig. 1B,C**). The total assembly size was approximately

3.55 Gb, with much higher quality exceeding those of two other available *Phalaenopsis*

genome assemblies (**Table S1**). Predicted telomere sequences were identified at most

chromosome ends (**Fig. 1D**), further suggesting the relatively high assembly quality. Due to

the complex hybrid history, the *P*. sa genome did not display obvious phases of subgenomes

(**Fig. 1C, Fig. S3**). Synteny analysis revealed extensive structural variation between

homoeologous chromosomes (HChrs) (**Fig. 1D**), including approximately 140 inversions,

1079 translocations, and 513 segmental duplications (**Table S2**).

Fig. 1 Genome assembly of *Phalaenopsis* 'Santiago' (*P.* sa). (**A**) A *P.* sa flower has one middle sepal, two lateral sepals, two petals, one lip, and a gynostemium that connects the stamen with two pollinia and the ovary. (**B**) Hi-C heatmap showing chromosomal interactions, with bin size of 250 Kb. Two insets illustrate weak interaction signals between homoeologous chromosomes (HChrs) of HChr group 10 (HChrG10, pink) and HChrG19 (blue) as examples. (**C**) Circos plot displaying genomic features within 1.5-Mb bins in *P.* sa. a, chromosome sizes, with different colors indicating predicted subgenomes, which is inferred from **Fig. S3**. b, gene density, ranging from 0 to 99. c, GC content, 15.38%–53.56%. d, transposable element coverage, 37.86%–99.79%. e, Copia LTR retrotransposon coverage, 0%–17.73%. f, Gypsy LTR retrotransposon coverage, 0%–98.51%. g, tandem repeat density, 0–251. The center region shows syntenic blocks between (blue lines) or within (orange lines) chromosomes. (**D**) Collinearity among homoeologous chromosomes, with predicted telomeres indicated.

We predicted 93,708 protein-coding genes using *de novo*, homology-based, and

transcriptome-supported approaches (**Methods**, **Table S3**). Repetitive sequences constituted

67.97% (2.41 Gb) of *P. sa* genome (**Table S4**). Among these, long terminal repeats (LTRs)

represented 41.91% of the genome, consisting of Copia (4.18%), Gypsy (37.22%), and

unknown types (0.51%), while tandem repeats accounted for 2.48%. The peak LTR insertion

time was estimated at approximately 0.31 million years ago (**Fig. S2E**). Additionally, the

genome annotation included 308 microRNAs, 1,094 transfer RNAs, 8,135 ribosomal RNAs,

and 34,366 small nuclear RNAs (snRNAs) (**Table S5**).

Homoeologous gene variation in *P.* sa

During the genome annotation, we noticed substantial variation in HGs. According to

syntenic analysis (**Methods**), all the 93,708 genes were classified into 33,697 HG groups

(HGGs), which were further categorized into six types according to variation in the number

and sequence among HGs (**Fig. 2A**). Regarding variation in sequence, to facilitate the

analysis, we examined the Pfam domain constitutions of the encoded protein sequences

among HGs, with the assumption that proteins with different domain constitutions are likely

to perform divergent functions[12,13]. Type 1 contained 12,673 HGGs (37.6% of groups, 13.5%

of genes) with only single HG per gene; type 2 consisted of 7,839 HGGs (23.3%/21.8%) with

HChr-specific gene losses while ≥2 HGs were retained; type 3 included 1,964 HGGs

(5.8%/18.4%) with HChr-specific tandem duplicates; type 4 comprised 373 HGGs

(1.1%/3.3%) with same number of tandem duplicates per HG; type 5 contained 3,642 HGGs

(10.8%/12.8%) with conserved single HG per homoeologous locus (HL) but these HGs have

different protein domain constitutions; type 6 included 7,206 HGGs (21.4%/30.2%) with

single HG per HL which had conserved domain constitutions (**Fig. 2A**).

Genes in types 1-6 differed in terms of expression patterns, functional annotation (**Methods**)

and encoded protein length (**Fig. 2B,C**), and tended to belong to different transcription factor

(TF) families (**Fig. 2D**) and perform different biological functions (**Fig. 2E**). Specifically,

type 1 genes tended to be not expressed (Transcripts Per Kilobase [TPM]=0, 45.1%) or lowly

expressed (TPM≤2, 30.6%) in all 28 examined samples (**Table S6**), have no functional

annotations and have relatively short proteins (**Fig. 2B**). Among 50 type 1 genes that were

annotated as potential TFs, 13 were *FAR-RED IMPAIRED RESPONSE 1 (FAR1)* genes (**Fig.**

2D). These 13 *FAR1* genes had TPM≤2 in all examined samples (**Fig. S4**), indicating they

were undergoing pseudogenization and might be lost from the genome in the long term.

FAR1 genes are essential for phytochrome signaling and balancing plant growth and defense

responses under shade conditions[15]. The deletion of certain *FAR1* genes from *Phalaenopsis*

genomes may be results of adaptation to shady, damp forest understory. In contrast, another

Fig. 2 Homoeologous gene variation in *P*. sa genome. (**A**) Diagram illustrating the six HG types. D1 and D2: protein domain 1 and domain 2. (**B**) Frequency of six HG types (upper panel) with the number of HGGs listed within the parentheses, and their protein length distribution (lower) with the average length per type indicated as well. Box colors indicate genes with or without functional annotations. TPM: Transcripts Per Kilobase; N.E.: not expressed; L.E.: lowly expressed; E.: expressed; H.E.: highly expressed. Inset: frequency of highly expressed (TPM>10) type 1 genes with (w.) or without (w.o.) functional annotation. Different gray scales indicate whether a gene has the best non-self blastp hit on different, homoeologous or same chromosomes, or has no blastp hit. (**C**) Mixed distribution of gene expression levels, illustrating how TPM=10 was selected as the cutoff threshold for high expression (see **Methods**). Fitted normal distribution indicates true expression, while fitted exponentially decaying distribution indicates noise expression. (**D,E**) Enriched transcription factors (**D**) and biological_process GO terms (**E**) in six HG types based on Fisher's exact test. Dot size: odds ratio; dot color: -$\log_{10}$ *q*-values which are the adjusted *p*-values using the "BH" method[14]; cases with *q*-values≥0.05 or odds ratio<1 were colored gray.

519 type 1 genes were highly expressed (TPM>10, 10.3%) in certain tissues, but had no

functional annotation and no non-self blastp hit within *P*. sa genome. These genes might be

potential *de novo* new genes, as 46.8% (243) of them were preferentially expressed in stamen

(**Fig. S5A-C**), consistent with previous findings that new genes tended to arise from male

reproductive organs[16,17]. The remaining 276 genes were preferentially expressed in other

tissues (**Fig. S5D-G**), such as middle root (*Psa_17aG011940*), leaf (*Psa_16dG007910*), shoot

apical meristem and inflorescence stalk (*Psa_04bG014720*), suggestive of their potential

functions in the corresponding tissues.

Type 2 genes also tended to have short proteins (**Fig. 2B**), and be enriched for TFs that play

important roles in floral development (MIKC-type MADS-Box), cell proliferation,

determination and differentiation (Basic Helix-Loop-Helix, bHLH), lateral organ formation

and responses to environmental stimuli (Lateral Organ Boundaries Domain, LBD), and

nitrate responses (Nin-like) (**Fig. 2D**). Type 3 genes tended to have longer proteins, be

enriched for TFs involved in flowering, embryogenesis (such as M-type MADS), stress

responses and signaling (Related to ABI3 and VP1 [RAV], Calmodulin-binding transcription

activator [CAMTA]) (**Fig. 2D**). GO terms enriched in type 3 included those participating in

interaction with environments, such as secondary metabolic processes, response to biotic and

endogenous stimuli, and signal transduction (**Fig. 2E**). Type 4 genes tended to be enriched in

TFs and functions related to plant growth and stress responses, including NAM, ATAF1/2

and CUC2 (NAC), ETS2 repressor factor (ERF), MYB and WUSCHEL-related homeobox

(WOX) families (**Fig. 2D**), and biological functions in fruit ripening, flower development,

responses to biotic and endogenous stimuli (**Fig. 2E**). Genes in types 5 and 6 had even longer

proteins, were enriched for TFs responsible for and GO terms related to essential and

conserved functions (**Fig. 2B,D,E**), such as LEAFY (LFY) in type 5 and GROWTH-

REGULATING FACTOR (GRF) in type 6, and biological processes including pollination

(type 5) and cell differentiation and growth (type 6). These results indicate that the HChr-

specific gene duplicates/losses (types 2,3), together with those tandem duplicates retained in

all HLs (type 4), may have driven the high diversity of floral traits and the adaptation of

Phalaenopsis to diverse environments.

Dosage effects of homoeologous gene count

Considering the aneuploidy nature of and the substantial HG count variation in *P*. sa genome,

we asked whether the chromosome ploidy (3x vs 4x) and HG count variation affect the total

expression amount (summed expression levels of all HGs) per gene. To avoid the influences

of tandem duplicates and gene structural variation on the total expression amount per gene,

we focused on HGGs which had a single gene copy per HL if the HG is present [such as

0:0:1:1 (2/4), 1:1:1 (3/3)], with no differences in protein domain constitutions among HGs

(C1-C8, **Fig. 3A**). In addition, to ensure a valid comparison, our analysis was restricted to

reciprocal best-matched (RBM) gene pairs that differed in chromosome ploidy or HG count.

This strategy was adopted because the gene expression in the hybrid parents of *P. sa* were not

available.

Fig. 3 Dosage effects of chromosome ploidy and homoeologous gene count on gene expression.
(**A**) Diagrams showing eight comparisons, where C1 compares reciprocal best-matched (RBM) gene pairs located on HChrGs with different ploidies (3x vs 4x), while C2-C8 compare RBM genes with different HG counts. 2/3 and 2/4: only two HGs were retained for HChrGs with 3x and 4x, respectively; 3/4: three HGs were retained for HChrGs with 4x. (**B,C**) Summed (**B**) and averaged (**C**) HG expression levels in petals at S11 of RBM gene pairs between HChrGs with 3x or 4x. (**D-J**) Summed HG expression levels for RBM gene pairs in C2-C8. (**K**) Averaged HG expression levels for RBM gene pairs in C8. Regarding (**B-K**), see **Dataset 1** for expression results in other tissues.

We found that gene expression scaled with chromosome ploidy: summed expression levels of

type 6 set4 genes (4/4) were generally higher than those of their 3/3 RBM counterparts

(*p*=4.67e-04, two-sided Wilcoxon rank sum test, **Fig. 3B**), but became comparable when

averaged per chromosome count (*p*=0.43, **Fig. 3C**). This dosage effect was similar to those

found in other aneuploid organisms, where gene expression in aneuploids were compared with

their corresponding diploids, such as maize[18], budding yeast[19] and *Candida albicans*[20]. HG

counts also had dosage effects on gene expression for all but two comparisons (C3-C7, **Fig.**

3E-I): summed expressions of HGs with HChr-specific gene losses were generally lower than

those of their RBM counterparts with more HGs. However, the expression tended to be

compensated when HG from only one HL was lost (C2, C8): no significant differences in

summed expression levels between 2/3 (HG from one out of the three HLs was lost) and 3/3

genes (p=0.99, **Fig. 3D**), and between 3/4 and 4/4 genes (p=0.22, **Fig. 3J**). This was further

supported by the significantly higher averaged levels of 3/4 genes than 4/4 genes (p=0.071,

corresponding to p=0.035 of one-sided Wilcoxon rank sum test, **Fig. 3K**). These results

suggest that buffering mechanisms counteract the disruption of homeostasis caused by

immediate HChr-specific gene losses (i.e., 2/3 and 3/4 genes), but not that caused by

chromosome inequivalence.

Genetic bases of key trait polymorphisms

The extensive HG variation in *P*. sa were mainly derived from allelic polymorphisms of its

hybrid parents, which represented part of the entire polymorphism set of *Phalaenopsis* that

contribute to the trait diversity. To evaluate the utility of *P*. sa genome in revealing the

genetic mechanisms underlying those diversity, we investigated a hybrid population (H80)

that was generated in our lab and displayed substantial variation in key floral traits,

particularly in lip morphology and color patterning (**Fig. 4A, Fig. S1**). Different trait

categories can combine randomly, resulting in a unique phenotype for each individual in this

population, making it a highly representative set of *Phalaenopsis* orchids. We resequenced

the genomes of 46 H80 representative individuals and performed genome-wide association

study (GWAS) using *P*. sa genome as the reference.

Key regulators of these traits have been previously identified, including *PeAGL6-2* in lip

organ identity determination[9], *PeMYB2*, *PeMYB12* and *PeMYB11* in the formation of floral

background color, stripes and spots, respectively[3], and *MYBx1* in repressing anthocyanin

accumulation in non-patched regions[7,8]. Expression profiles of these genes (see **Table S7** for

their gene IDs in the *P*. sa genome) reflected their relatively conserved functions in *P*. sa and

H80 (**Fig. 4B, Fig. 5A,B**). Nevertheless, these genes displayed different types of HG

variation in *P*. sa (**Fig. 4C, Fig. 5C, Fig. 6A**). Therefore, we focused on identifying genetic

variants within or near these genes that were significantly associated with corresponding trait

Fig. 4 Floral trait variation in H80 and the genetic basis of lip morphology variation. (**A**) Phenotypes of 24 H80 individuals (see **Fig. S1** for all 46 individuals). The category of each individual in lip morphology (0: lip, 1: petal-like organ), stripe P/A (0: A, 1: P), background color (0: white, 1: pink), and spot/patch types (0: spotted or non-spotted, 1: patched) were indicated. Bold font: individuals with transcriptome data of the lip. Red arrowhead indicates stripe, while blue arrowhead indicates stripe-like shadow. (**B**) Heatmap showing expression profile of A-G classes of MIKC-type MADS-box genes in different floral organs at three stages (S7, S9 and S11), and in the S9 lips of 11 representative H80 individuals. Color scale indicates log-transformed expression levels. mS: middle sepals; lS: lateral sepals; P: petals; L: lip; St: stamen; O: ovary; G: gynostemium. Red and black boxes: expression of genes positively or negatively correlated with petal-to-lip transitions, respectively. (**C**) HG variation of *PsAGL6-2*. Red boxes: *PsAGL6-2* genes with directions indicated; light or dark gray boxes: neighboring genes with forward or reverse directions, respectively; boxes linked using gray lines between HChrs: HGs. The blue arrowhead indicates locations of two significant SNPs. (**D**) Manhattan plot for GWAS of lip morphology in H80, using SNPs referred to HChr set c with the additive marker-effect model. Horizontal solid and dashed lines indicate -log$_{10}$ (0.01/variant counts) and -log$_{10}$ (0.05/variant counts) (**Methods**), respectively; red and orange dots denote SNPs significantly associated with lip morphology variation; blue dots represent SNPs located within or adjacent (±2 kb) to the *PsAGL6-2* gene clusters. (**E,F**) Allele read ratio of two SNPs (count of reads supporting "T" to count of reads supporting "A" in **E**, and count of reads supporting "T" to that of "C" in **F**) within H80 individuals with different lip morphologies. *P*-values are from two-sided Wilcoxon rank-sum tests.

polymorphisms, and decoding how they contributed to trait differentiation in H80. Since H80

individuals were shown to be aneuploid as well by karyotype analysis (**Fig. S2F**, 2n=64, >3x),

they were treated as tetraploids in the GWAS to facilitate the analysis. Moreover, given that

70 chromosomes in *P.* sa can not be readily phased into subgenomes (**Fig. 1C**), the original

HChr designations (such as that HChrG01a~HChrG19a were designated as HChr set a) were

retained when serving as the reference genome during the GWAS.

Lip morphology variation and nucleotide polymorphisms nearby *PsAGL6-2* genes

PsAGL6-2 had two to four tandemly arranged duplicates on three HChrs (corresponding to

HG type 3, **Fig. 4C**), with only one copy encoding a full-length protein per HChr (**Fig. S6**).

GWAS analysis for the lip morphology (lips vs petal-like organs) identified a number of

significant variants (including SNPs, indels and SVs) on the right arms of HChrG09s,

regardless of which HChr set was used as references (**Fig. S7-9**). However, only when the

HChr set c was used as the reference (**Fig. 4D, S9A**), two significant SNPs were identified in

the intergenic region between two *PsAGL6-2* copies (**Fig. 4C**, ~65 Kb upstream of

Psa_09cG004690 and ~86 Kb downstream of *Psa_09cG004710*), which were located on the

left arms. The allele read ratios (the number of reads supporting one HG divided by the

number of reads supporting the other HG) of these two SNPs were significantly associated

with lip morphology in H80 (**Fig. 4E,F**, **Table S8**). Furthermore, expression levels of

PsAGL6-2 showed positive correlation with lip morphology. These results suggest that the

transformation of lips into petal-like organs in H80 did not result from loss-of-function

mutants in *PsAGL6-2*, as it remained expressed at substantial levels in these organs. Instead,

the morphological change is associated with reduced expression of *PsAGL6-2*, potentially

due to the introduction of an allele containing mutations in a long-distance enhancer (likely in

the intergenic region between *Psa_09cG004690* and *Psa_09cG004710*). Additionally,

expression of other proposed lip [*PsAPETALA3-4* (*PsAP3-4*)] and petal identity genes

(*PsAP3-1*, *PsAP3-3*, and *PsAGL6-1*; see **Fig. S10** for the gene tree of MADS-box genes)[21]

exhibited positive and negative correlations with petal-to-lip transitions, respectively (**Fig.**

4B). These results suggest that expression of these genes might be regulated by *PsAGL6-2*,

thereby coordinating lip morphogenesis.

Presence/absence of stripes and presence/absence of *PsMYB12* genes

As the regulator of stripe, expression of *PsMYB12* genes was highly associated with stripe

formation in *P.* sa and the presence of stripes in H80 population (**Fig. 5A,B**). It lost its entire

tandemly duplicated gene cluster from HChrG04b (HG type 2, **Fig. 5C**), indicating the P/A

of stripes in *Phalaenopsis* is potentially determined by *PeMYB12* P/A. CNV-based GWAS (**Fig. 5D, Methods**) using HChr set a as the reference identified significant variants located in or nearby *PsMYB12* genes while with the fewest false positives compared with others (**Fig. S11**). Consistent with this, stripes P/A in H80 was perfectly associated with P/A of resequencing reads mapped to *PsMYB12* genes (**Fig. 5F**, **Fig. S12, Table S9**), further confirming that stripe P/A are determined by *PeMYB12* P/A.

Fig. 5 Genetic bases of color patterning variation in H80. (**A**) Floral organs at three developmental stages showing important events of color patterning. VLS and DLS: ventral and dorsal half of the lateral sepals, respectively. (**B**) Heatmap showing expression profiles of genes participating in color patterning. The classifications of each H80 individual in the presence (0) or absence (1) of stripes, background pink colors and spots (2 indicates patches) are listed. MBW TFs: transcription factors that form the MYB-bHLH-WD40 complexes. The biosynthesis pathway for the prominent anthocyanin in *Phalaenopsis* orchids—cyanidin—is shown on the right. Str: stripe; Bac: inter-stripe background region. (**C**) HG variation of *PsMYB12*, *PsMYB2* and *PsMYB11*. (**D**) Manhattan plot for GWAS of stripe P/A in H80, using CNVs referred to HChr set a with the additive marker-effect model. (**E**) Manhattan plot for GWAS of background color in H80, using SNPs referred to HChr set b with the 1-dom-ref marker-effect model. (**F,G**) Association of P/A of *PsMYB12* genes (**F**) and P/A of *PsMYB2b*-type sequences (**G**) with stripe P/A and background color, respectively. *P*-values are from Fisher's exact test.

Since we didn't identify any genes orthologous to *PsMYB12* in the other eight orchid species
(including the non-striped *P. aphrodite*) except for *P. equestris* (**Fig. S13A**), we asked when
and how the *PsMYB12* genes and stripes originated in *Phalaenopsis*. Interestingly, stripes
were only observed for the branch leading to *P. equestris* and *P. lindenii* in wild
Phalaenopsis species (**Fig. S13B**). To test if *PsMYB12* genes originated before or after the
most recent common ancestor (MRCA) of *Phalaenopsis*, we resequenced the *Vanda*
'Pakchong Blue', an orchid cultivar outside *Phalaenopsis* which displays venation-associated
stripes on the perianth as well. We detected reads mapped to *PsMYB12* genes, albeit with a
high degree of divergence within and around these genes (**Fig. S13C**). Given that other
orchids like *Paphiopedilum* also have stripes, these results suggest that *PsMYB12* genes
predate the MRCA of *Phalaenopsis* and have undergone repeated, independent losses in
Phalaenopsis species. Alternatively, these genes may be absent in the MRCA of
Phalaenopsis and later reintroduced into the MRCA of *P. equestris* and *P. lindenii*,
potentially via gene introgression.

Background pink color and the *b*-type HG of *PsMYB2*
PsMYB2 had a single HG per HL (**Fig. 5C**), but the one on HChrG01c (*Psa_01cG009830*)
had lost its R2 domain (HG type 5, **Fig. S14A**) and was expressed at much lower levels than
other HGs (**Fig. S14B**). Instead, *PsMYB2b* (*Psa_01bG009870*) was expressed at the highest
levels among its HGs (**Fig. S14B**). GWAS using 1-dom-ref model (**Methods**) with HChr set
b as the reference identified significant SNPs and indels located in or nearby *PsMYB2* genes
while with the fewest false positives among others (**Fig. 5E**, **Fig. S15-17**). In this 1-dom-ref
model, the *PsMYB2b*-type HG was considered as dominant in determining the pink color.
Consistently, the P/A of *PsMYB2b*-type sequences can nearly perfectly predict the P/A of
background pink color for all but three individuals—H80-066, H80-072 and H80-114 (**Fig.
5G**, **Table S10**). The background color in these exceptions may be governed by other non-
PsMYB2 genes, since *PsMYB2* expression was almost undetectable in the lip of H80-072 (the
summed TPM of four HGs was 1.32, **Fig. 5B**). Given that the protein sequence of PsMYB2b
differed from PsMYB2a by only two amino acids (**Fig. S14A**), these results suggest that the
pink color in *P.* sa and most H80 individuals with pink flowers was likely determined by the
HChr set b-specific expression of *PsMYB2*, which was driven by its distinct promoter (**Table
S10**). It requires further investigation of which region of the *PsMYB2b* promoter drives its
HG-specific expression.

Spot/patch pattern variation

PsMYB11 had an extra tandem copy (*Psa_09cG010030*) on HChrG09c (HG type 3, **Fig. 5C**), which harbored a few mutations in protein sequences and the upstream sequences (**Fig. S14C,D**), compared with other HGs. *Psa_09cG010030* was highly expressed in the lip of *P. sa* at S11 where the spots formed, and in lips of H80-077 and H80-070 which had patches instead of spots (**Fig. 4A**, **Fig. S14E**). GWAS for the P/A of patches identified numerous variants distributed across the entire genome, but no variants were near the *PsMYB11* genes, regardless of variant types used and HChr sets referred to (**Fig. S18-20**). Given that the patch formation was reported to be determined by the *HORT1* insertion in the promoter sequences of *PeMYB11* genes[7], and that *P. sa* flowers harbored only small spots, the failure to identify *PsMYB11*-related variants is potentially due to the lack of *HORT1* insertion in *PsMYB11* promoters.

Functions of *PsMYBx1* in repressing anthocyanin accumulation

PsMYBx1 had only one HG (HG type 1, **Fig. 6A**). Its expression was associated with both stripe and spot/patch formation in *P. sa* and H80 (**Fig. 5A,B**), indicating potentially dual function of *PsMYBx1* in repressing anthocyanin accumulation in both inter-stripe and inter-spot/patch regions. This dual function was supported by the disruption of stripes in regions surrounding patches, such as in H80-057, H80-114, and H80-245 (**Fig. 4A**). Given this complexity of color patterning imposed by *PsMYBx1* genes, no single color patterning trait variation can be readily determined to be related to *PsMYBx1*. Thus, instead of performing GWAS, we silenced *MYBx1* genes via virus-induced gene silencing (**Methods**) in another *Phalaenopsis* cultivar, *P.* 'Jinbianlinglong' (*P.* ji) (**Fig. 6B,C**), for which substantial seedlings are available and the flowering regulation is relatively readily. *MYBx1* had four HGs in *P.* ji (**Fig. S21B**), compared with a single HG in *P. sa*. *PjPDS* (*Phytoene desaturase*) was co-silenced to indicate regions with gene silencing phenotypes where the yellow-green color was faded (**Fig. 6B**). In *PjMYBx1-PjPDS*-silenced plants, ectopic anthocyanin pigmentation was observed on the inflorescence stalk and abaxial surface of perianth, and in the inter-stripe regions (**Fig. 6C**). However, the inter-spot regions remained unchanged (blue arrowheads in **Fig. 6C**), contrasting with the proposed role of *MYBx1* in repressing anthocyanin accumulation in inter-spot regions[8]. Although expression of color patterning-related genes examined were generally decreased in *PjMYBxl-PjPDS*-silenced sepals (**Fig. 6D**) and petals (**Fig. 6E**) compared with those in mock-treated plants, a slight increase in *PjMYB12* expression was detected in petals. Taken together, these results demonstrate the multiple

functions of *MYBx1* in repressing anthocyanin accumulation in diverse regions, and suggest

variation in its functions across different *Phalaenopsis* orchids.

Fig. 6 Functions of *PjMYBx1* in repressing anthocyanin accumulation in diverse regions. (**A**) HG variation of *PsMYBx1*. (**B**) Flower and inflorescence images showing phenotypes of mock-treated (upper panel) and *PjPDS*-silenced (lower panel) plants. (**C**) Flower and inflorescence images of *PjMYBx1-PjPDS*-silenced plants. Red arrowheads indicate regions with phenotypic changes when compared with mock-treated plants, while the blue ones indicate the unchanged inter-spot regions. (**D,E**) Expression of color patterning-related genes in mock-treated and *PjMYBx1-PjPDS*-silenced sepals (**D**) and petals (**E**). *P*-values are from t-test.

Discussion

The extraordinary phenotypic diversity and the high economic importance of *Phalaenopsis*

orchids necessitate a reference genome assembly with high quality to facilitate the

fundamental research and the breeding. The haplotype-resolved, chromosome-level genome

assembly of *P.* sa provided in this study served as a suitable reference in revealing the genetic

bases of trait diversity, particularly those of lip morphology, venation-associated stripes and

pink color on sepals/petals. This is exemplified by analysis in the H80 population. Given the

high heterogeneity of *P.* sa genome and the diverse genetic mechanisms underlying different

trait variation, we demonstrated that the choice of HChr sets as the reference genome, variant

types, and marker-effect models is critical for revealing the genetic bases of target trait

variation. The ratio of true to false positives of the identified significant variants can be the

primary metric for optimization. Specifically, for lip morphology variation, stripe P/A

variation, and background color variation, the optimal models were HChr set c + SNPs +

additive model, HChr set a + CNVs + additive model, HChr set b + SNPs/indels + 1-dom-ref

model, respectively. However, for spot/patch variation, genomes of patch-bearing

Phalaenopsis cultivars are supposed to serve as superior references compared with *P*. sa

genome.

Phalaenopsis flowers display extreme complexity in color patterning, which is much more

complicated than simply superposition of background color, spots and stripes. This

complexity is well demonstrated in the H80 population, such as that spots can be distributed

along the venation (H80-057 and H80-246), and spots/patches and stripes can be mutually

exclusive (H80-245) where the stripes are disrupted by spots/patches (**Fig. 4A**). This

complexity results from the substantial allelic polymorphisms of the regulators—particularly

MYB genes, including both R2R3-type (positive regulators) and R3-type (negative regulators).

Even in the single *P*. sa genome, these *MYB* genes exhibit HG variation in copy number

(*PsMYB11*), presence and absence (*PsMBY12* and *PsMYBx1*), gene structure, sequence and

expression (*PsMYB2*). Furthermore, *PsMYBx1* exhibits functional differentiation across

cultivars by repressing anthocyanin accumulation in distinct regions. Further studies are

needed to fully decipher the genetic bases of the extensive color patterning diversity and the

interplays among these regulators in *Phalaenopsis*.

Although genes regulating some of the key traits in *Phalaenopsis* have been well

established[1–3,5,9,22,23], what and how allelic polymorphisms of these genes have contributed to

the extraordinary trait diversity remain underexplored, restricting their utilization in breeding

practices. The genetic bases of three key trait variation revealed in this study—P/A of

PsMYB12 genes, P/A of *b*-type HG of *PsMYB2*, and nucleotide polymorphisms nearby

PsAGL6-2 genes—enable design of a multiplex polymerase chain reaction (MPCR) kit for

early-stage prediction of these three traits simultaneously. More traits can be predicted using

such a single MPCR experiment when the genetic bases of their diversity are deciphered, to

further improve the cost effectiveness. Given the unusually long juvenile phases of

Phalaenopsis orchids (three to five years), such a MPCR kit has great potential in

substantially shortening breeding cycles and reducing breeding costs.

Methods

Flow cytometry analysis

Young leaves of *P*. sa seedlings were used for the flow cytometry analysis, with young leaves

of cultivated tomato one month after seed germination being used as the internal reference.

Suspensions of *P.* sa and tomato leaves were strained using propidium iodide and mixed, and
a BD FACScalibur flow cytometer was used to detect the stained cell nuclei.

Karyotype analysis

Root tips of *P.* sa were harvested between 9:00 and 10:00 a.m., and approximately 0.5 cm
segments were excised using a clean razor blade. These segments were incubated in 0.1%
colchicine solution at room temperature in the dark for 5 h to induce chromosome
condensation, and were then fixed in Carnoy's solution (absolute ethanol:glacial acetic acid =
3:1, v/v) overnight. Fixed samples were rinsed three times with distilled water, dehydrated in
75%, 85%, 95%, and 100% ethanol for 10 minutes each, and then were stored in 75% ethanol
at 4 °C. To facilitate cell wall separation, pretreated samples were incubated in 75 mM KCl
solution at 37 °C for 30–60 minutes, and treated with 1 M HCl at 60 °C in a water bath for 10
minutes. Samples were stained with carbol fuchsin solution (Coolaber, Beijing, China) for 20
minutes, squashed on microscope slides, observed and photographed using a 60× objective
lens on an OLYMPUS BX53 optical microscope.

Genome sequencing and Hi-C sequencing

For HiFi sequences, the total DNA was extracted from *P.* sa young leaves, and used for the
library construction and genome sequencing with the PacBio Sequel II/Sequel IIe platform,
following the manufacturer's instructions at Haorui Genomics (Xi'an, China). Approximately
72.1 Gb of PacBio HiFi reads were generated using the CCS mode.

For Hi-C sequences, the frozen young leaves were ground with liquid nitrogen and then fixed
with 1% formaldehyde in vacuum at room temperature for 0.5-1h to preserve the three-
dimensional chromatin architecture, and 0.15 M glycine was added in the solution for 5 min
to quench the reaction. The mixed solution was then filtered with a 40 nm cell filter.
Chromatin was digested with 200 U DPNII restriction enzyme (Qiagen, 59971) at 37 °C for
16 h. DNA ends were labeled with biotin and incubated at 37 °C for 45 min. DNA ligation
was performed by the addition of T4 DNA ligase and incubation for 1-2 h. Then proteinase K
was added to reverse cross-linking during incubation at 65 °C for 1-2 h. DNA fragments were
purified and dissolved in 86 μL of water. Purified DNA was fragmented to a size of 350-500
bp, and DNA ends were then repaired. DNA fragments labeled with biotin were finally
separated on Dynabeads® M-280 Streptavidin (Life Technologies, 60210). The sequencing
was performed with the Illumina NovaSeq 6000 for paired-end sequences with 150 bp read
length. A total of ~130.6 Gb raw data was produced. After filtering adapter sequences and

read pairs with > 10% N or with low quality (Q ≤ 5) for >50% bases in at least one read,

~123.3 Gb clean data was retained.

Genome survey

409   The genome size was estimated via *k*-mer frequency analysis using the PacBio HiFi reads.

410   The Jellyfish software (v2.2.10)[24] was used to generate the counts of 31-mers, and then the

411   GenomeScope2 software (v2.0)[25] and Genomic Character Estimator (GCE) software

412   (v1.0.2)[26] were applied to estimate the genome size and heterozygosity. SmudgePlot software

413   (v0.2.2)[25] was used to estimate the ploidy level of *P*. sa genome.

De novo genome assembly

Since it lacks a chromosome-level reference assembly for *Phalaenopsis* species, at the first

place, we attempted to assemble a set of draft representative pseudo-chromosomes to guide

the subsequent refined assembling. Three strategies were conducted to assemble the draft

genome: (**i**) all the PacBio HiFi reads were assembled using HiFiasm program (v0.19.6-

r597)[27] with the parameter 'l=0' (to disable the purge of duplication); (**ii**) the PacBio HiFi

reads were assembled using HiFiasm with the parameters 'l=3, hom-cov=57, hg-size=1.3g, n-

hap=3, s=0.3, D=8, n-weight=6, n-perturb=80,000, f-perturb=0.3', with the assumption that

the *P.* sa was a triploid, and the Hi-C reads were integrated to achieve haplotype-resolved

assemblies; (**iii**) the PacBio HiFi reads were assembled with parameters 'l=3, hom-cov=39,

hg-size=1.5g, n-hap=4, s=0.3, D=8, n-weight=6, n-perturb=80,000, f-perturb=0.3', with the

assumption that the *P.* sa was a tetraploid, and the Hi-C reads were also integrated.

The Hi-C reads were mapped to the resulting primary unitigs from assembling **i**, and

haplotigs of each haplotype from assembling **ii** and **iii**, separately, using bwa-mem2 (v2.2.1)[28].

The resulting bam files were further filtered using the 'filter_bam.py' utility in the HapHic

(v1.0.2)[29] with the parameters 'mapq=1, NM=3'. The filtered bam files were then fed into the

HapHic pipeline with parameters 'correct_nrounds=2, quick_view=T', and the output

files .hic and .assembly were imported into the Juicebox Assembly Tools (JBAT, v1.08)[30].

Primary unitigs from strategy (i) and haplotigs from (ii) and (iii) were then ordered, joined,

and manually adjusted based on Hi-C interaction patterns to obtain scaffolds, with the prior

knowledge that nearby regions within the same chromosome having strong Hi-C signals

whereas allelic regions having subdominant signals. The resulting scaffolds from all three

strategies were compared, and 19 were selected as representative pseudo-chromosomes based

on Hi-C signal continuity and Benchmarking Universal Single-Copy Orthologs (BUSCO)[31]

completeness. These 19 pseudo-chromosomes—seven from assembly **i**, ten from **ii**, and two

from **iii**—constitute the draft reference assembly (assembly v1.0).

Next, using the v1 draft assembly as a reference, we mapped all raw unitigs from assembly **ii**

and grouped them into 19 chromosomal groups. Unitigs within each group were ordered,

joined, and manually adjusted based on Hi-C interaction patterns, with the parameters

'mapq=0, NM=3' for filter_bam.py. This produced 70 preliminary pseudo-chromosomes.

Remaining unassigned unitigs were then incorporated and the entire set was reviewed and

corrected using Hi-C signals to yield the final assembly (assembly v2.0). SubPhase (v1.2.6)[32]

was used to phase subgenomes based on potential subgenome-specific k-mers, with

parameters "-k 15 -q 50 -f 12".

Genome assembly quality assessment

The *P.* sa genome assembly contiguity was accessed using Quality Assessment Tool for

Genome Assemblies (QUAST, v5.0.2)[33]. The BUSCO completeness was conducted using

BUSCO (v5.6.1)[31], against the eukaryota_odb12, viridiplantae_odb12, embrophyta_odb12,

and liliopsida_odb12 databases. The K-mer (21-mers) completeness and assembly consensus

quality value (QV) were estimated using Merqury (v1.4.1)[34]. The QV is a log-scaled

probability of errors for consensus base calls; the higher the QV values, the more accurate the

consensus base calls, with a QV value at 30 indicating 99.9% accuracy and 40 representing

99.99% accuracy[34]. The regional and structural assembly quality indicator (R-AQI and S-AQI)

were assessed using the Clipping information for Revealing Assembly Quality (CRAQ,

v1.0.9) algorithm[35]; the higher the AQI scores, the more accurate the genome assembly[35]. The

long terminal repeat retrotransposons (LTR-RTs) were first annotated using LTRharvest

(v1.6.1)[36] with the parameters '-maxlenltr 6000 -mindistltr 1500 -maxdistltr 25000 -mintsd 5

-maxtsd 5 -vic 10 -similar 85' and LTR_FINDER_parallel (v1.2)[37] with the parameters '-size

1000000 -time 300'. Then the LTR Assembly Index (LAI)[38] was calculated using

LTR_retriever (v2.9.5)[39] with the parameter '-u 0.5e-8'. The telomere and centromeric repeat

identification were conducted using quartet (v1.2.5) using default parameter settings[40].

Noncoding RNA and repetitive sequence identification

466   Noncoding RNAs were predicted using tRNAscan-SE (v2.0.9)[41] and Infernal software

467   (v1.1.4)[42] by searching the database Rfam (v14.10). The extensive *de-novo* TE annotator

468   (EDTA) (v2.1.0)[43]—that encompasses LTR_FINDER, LTRharvest, LTR_retriver, Generic

469   Repeat Finder, TIR-Learner, HelitronScanner, RepeatModeler and RepeatMasker— was used

to identify, annotate and classify repetitive elements of the genome, with parameters '--anno 1 --sensitive 1 --evaluate 1 --u 0.5e-8 --curatedlib ${SINE_LINE_lib}'. The library of short interspersed elements (SINEs) and long interspersed nuclear elements (LINEs) was downloaded from SINEBase (https://sines.eimb.ru/) as a compensation for the reference sequences. The repetitive elements identified by EDTA included long terminal repeat (LTR) retrotransposons (copia, gypsy and unknown ones) and non-LTR retrotransposons (LINE_element and unknown ones), terminal inverted repeat (TIR) transposons (CACTA, Mutator, PIF_Harbinger, Tc1_mariner and hAT), *Helitron* and other interspersed repeats. Subsequently, the DeepTE (version updated on Oct 22th, 2022) script[44] was used to reclassify the TEs that were initially annotated as 'LTR/unknown' by EDTA into copia, gypsy and unknown classes, then the output file was fed back into EDTA to enhance the comprehensiveness of TE annotation. To calculate LTR insertion time, we set the mutation rate per generation (μ) to 0.5×10^{-8}, based on the assumption that the generation times of orchids are four years[45]. The resulting hard-masked genome was converted into a soft-masked genome, which was used in the subsequent analysis. The tandem repeat sequences in the *P. sa* genome were identified using Tandem Repeats Finder (TRF, v4.09)[46] with the parameters '2 7 7 80 10 50 2000 -d -h'. Then the proportion of tandem repeats in the genome was calculated using the script buildRepeatSummary.pl (v1.36) with default parameters[47].

Transcriptome sequencing

Total RNA was extracted for roots, leaves, stems and flower buds with mixed stages (one duplicate) and other tissues at different development stages (three replicates, **Table S6**). The library preparation and PCR amplification were performed according to the manufacturer's instructions, and the RNA libraries were sequenced on the Illumina Novaseq 6000 platform and 150 bp paired-end reads were generated. Clean data were further obtained by removing reads containing adapters, reads containing poly-N and low quality reads from raw data, using fastp (v0.24.1)[48] with parameters: '-g -q 5 -u 50 -n 15 -l 150 --overlap_diff_limit 1 -- overlap_diff_percent_limit 10'.

Protein-coding gene prediction

Protein-coding genes were predicted by integrating predictions based on sequence homology and transcriptome, and the *ab initio* predictions using the soft-masked genome.

1. Homology-based prediction

Genome assembly and General Feature Format (GFF) files of six plant species were used for

homology-based prediction with Gene Model Mapper (GeMoMa, v1.9)[49]. The following

species were included: *Arabidopsis thaliana* (TAIR10.1, downloaded from NCBI), *Oryza*

sativa Japonica Group (IRGSP_1.0, from Ensembl Plant), *Asparagus officinalis* (Aspof.V1,

from Ensembl Plant), *Apostasia shenzhenica* (GCA_002786265.1, from NCBI)[50],

Dendrobium catenatum (GCF_001605985.2, from NCBI)[51], and *Phalaenopsis equestris*

(GCF_001263595.1, from NCBI)[10]. The GeMoMa parameters used were

'AnnotationFinalizer.r=NO p=false o=true tblastn=false'.

2. Transcriptome-based prediction

Clean RNA-seq reads from roots, leaves, stems, and mixed-stage flower buds were aligned to

P. sa genome using HISAT2 (hierarchical indexing for spliced alignment of transcripts,

v2.2.1)[52]. The alignments were processed by StringTie (v2.2.1)[53] to reconstruct genome-

guided transcripts. Potential open reading frames (ORFs, ≥100 amino acids) were identified

from these transcripts using TransDecoder (v5.7.1) (https://github.com/TransDecoder). A

genome-guided transcriptome was assembled using Trinity (v2.15.1)[54]. Both the StringTie

output and the Trinity assembly were then analyzed by the Program to Assemble Spliced

Alignments (PASA, v2.5.3)[55] with the parameters '--ALIGNERS blat,minimap2 --

TRANSDECODER' to generate transcriptome-supported gene models.

3. *Ab initio* prediction

The BRAKER3 pipeline (v3.0.3, incorporating Augustus and Genemark)[56], GeneID (2021

version)[57] and GlimmerHMM (2014 version)[58] were used for the *ab initio* gene prediction,

leveraging the transcriptome evidence as a guide.

4. Gene model integration and filtering

The EvidenceModeler (v2.1.0)[59,60] was used to integrate predictions from the above methods

into consensus gene structures. The following weights were assigned: GeneID (1),

GlimmerHMM (1), GeneMark (2), Augustus (4), GeMoMa (7), StringTie (5), and

PASA_transdecoder (10).

To refine the gene models, predicted protein sequences were searched against the Transposon

Protein Database (https://github.com/NBISweden/TransposonPSI/blob/master/

transposon_ORF_lib/transposon_db.pep) using TransposonPSI (v1.0.0), which employs PSI-

BLAST to identify TE-related sequences[61] with default parameters. Proteins with > 30%

similarity and > 60% coverage to known TE sequences were removed. In addition, gene

models with incomplete ORFs or ORFs ≤ 100 amino acids were excluded, if they met all the

three following conditions: lacking RNA-seq expression support, having < 50% coverage

overlap with GeMoMa models, and lacking functional annotation by InterProScan (v5.67-

99.0)[62].

5. Model optimization with extended RNA-seq data

The resulting gene models (v1.0) from above steps were further supplemented using

additional RNA-seq data from other 28 samples (**Table S6**). First, to avoid potential mapping

errors between tandem duplicates with high sequence similarities, RNA-seq reads were first

mapped to the v1.0 transcriptome, and unmapped reads were further mapped to the genome,

using TopHat2 (v2.1.1)[63] with the parameter '--max-multihits 1'. Next, individual bam files

were used to assemble transcripts (including alternative splicing isoforms) using StringTie.

The resulting GTF files from all samples were merged using StringTie to create a

comprehensive transcript annotation. TransDecoder was used to infer the ORFs, producing a

GFF file (v2.0) that included alternative splicing and five prime untranslated regions (5'UTR)

and 3'UTR information. Finally, the v2.0 annotations were compared with and used to

correct the v1.0 annotations, prioritizing v2.0 evidence in overlapping regions. This process

resulted in a final set of 93,708 protein-coding genes.

Protein-coding gene functional annotation

P. sa protein sequences were first searched against the eggNOG database (v5.0.2)[64] using

eggNOG-mapper (v2.1.12)[65], with parameters '-m diamond --seed_ortholog_evalue 1e-5 --

tax_scope Viridiplantae', for annotations including Gene Ontology (GO), Kyoto

Encyclopedia of Genes and Genomes (KEGG) Orthology (KO), and Clusters of Orthologous

Groups (COG). In order to obtain the functional description of related protein-coding genes

in model plant species, protein sequences of *P.* sa genome were searched against the UniProt

database (uniprot_sprot_plants.fasta that contains manually annotated protein sequences from

sequenced plant genomes, updated on March 27th, 2024, https://www.uniprot.org/) and

protein sequences from Arabidopsis (Araport11_pep_20240409, downloaded from The

Arabidopsis Information Resource) using BLASTP (v2.12.0+)[66] with parameters '-max_hsps

1 -max_target_seqs 1 -evalue 1e-5'. Furthermore, *P.* sa protein sequences were searched

against databases such as Pfam, SMART, CDD, Superfamily and others, using InterProScan (v5.67-99.0) with parameters '-goterms -iprlookup'. Gene Functional Annotation for Plants (GFAP, v3.1)[67] was used to annotate GO, KEGG and Pfam items by searching *P*. sa protein sequences against a plant-specific database with the parameter '-awd psd -go -kegg -pfam', and to obtain homology-based annotations by searching against the plant-specific NCBI non-redundant (nr) database with the parameter '-awd nr'. Lastly, protein sequences were searched against the Reference Protein Sequence Dataset (RPSD, release_2024_07_31) using Ensemble Enzyme Prediction Pipeline (E2P2, v5)[68] with default parameters for the enzyme function annotation. Transcription factors in *P*. sa genome were annotated in the Plant Transcription Factor Database (PlantTFDB)[69].

Homoeologous gene identification

The Synteny and Rearrangement Identifier (SyRI, v1.7.0)[70] and MCScanX v1.0.0[71] were used to conduct synteny analysis among different HChrs with default parameter settings, based on genomic and protein sequences, respectively. Later on, the homoeologous genes were identified with in-house scripts (see **Code availability**) using the outputs of SyRI preferentially and MCScanX supplementarily, to avoid improper assignment of multiple MCScanX short syntenic blocks between HChrs.

Transcriptome analysis

RNA-seq reads were mapped to the transcriptome and genome using Tophat2 (v2.1.1)[63], with parameter '--max-multihits 1', using indexes from both the genome and the final transcriptome. Read counts for protein-coding genes were obtained using htseq-count (v2.0.5)[72] and TPM was calculated using the "calculateTPM" function in R package scater (v1.0.4)[73]. The TPM values were normalized across samples using quantile normalization, followed by averaging across the three biological replicates to obtain a mean TPM value for each tissue.

To categorize genes according to their expression levels, we defined four thresholds based on TPM values: 1) TPM = 0 indicating no expression; 2) $0 < \text{TPM} \leq 2$ indicating low expression; 3) $2 < \text{TPM} \leq 10$ indicating moderate expression; 4) TPM > 10 indicating high expression. The threshold of TPM = 10 distinguishing high expression was determined by modeling the observed gene expression data as a mixed distribution. This model comprises a normal distribution, representing true biological expression, and an exponentially decaying distribution, representing technical noise or background. These two components were fitted

using the nls function from the R "stats" package[74]. The value of TPM = 10 corresponds approximately to the 50th percentile of the fitted normal distribution, suggesting that roughly half of the genuine gene expression occurs below this level.

Weighted gene co-expression network analysis (WGCNA)

Co-expression analysis for HG type 1 genes that were highly expressed (TPM > 10) in ≥ 1 tissue and had no non-self blast hist in *P. sa* genome was performed using WGCNA (v1.72.1)[75], under a signed network model, with a minimum module size of 10.

Gene tree building

To build a robust phylogeny tree for *MYB2*, *MYB11* and *MYB12* genes, coding sequences (CDS) were obtained from multiple orchid species—*P. aphrodite* (*P. ap*), *P. equestris* (*P. eq*), *Apostasia shenzhenica* (*A. sh*), *Cremastra appendiculata* (*C. ap*), *Cymbidium sinense* (*C. si*), *Dendrobium chrysotoxum* (*D. ch*), *Gastrodia elata* (*G. el*), *Platanthera guangdongensis* (*P. gu*), *Vanilla planifolia* (*V. pl*)—via the Orchid Multiomics Database (http://orchidcomics.com/, access on June 11th, 2025)[76]. For each query gene, the top 20 BLAST hits per species (top 80 hits for *P. sa*) were extracted, translated into amino acid sequences, and aligned using MEGA v7.0[77]. The resulting amino acid alignment was manually curated to retain only reliably alignable regions, which were then converted back to a corresponding CDS alignment. The best-fit nucleotide substitution model (GTR+I+G4) was selected using ModelTest-NG (v0.1.7)[78]. Finally, a maximum likelihood phylogenetic tree was constructed with RAxML-NG (v1.2.2) [79], employing 1000 bootstrap replicates.

The subclass of floral organ identity genes was determined by phylogenetic analysis. CDS of MIKC-type MADS-box genes from *P. sa* and other orchids (**Fig. S10**) were translated into amino acid sequences. These sequences were aligned using MEGA with manual adjustment. A phylogeny was inferred using the 'JTT+I+G4' model with 1000 bootstrap iterations.

Genetic variant calling

Total DNA was extracted for flower stalks of 46 H80 individuals. Sequencing services of partial H80 individuals were provided by BGI Genomics (Shenzhen, China), and the others were provided by Novogene Co., Ltd (Beijing, China). All whole-genome resequencing in this study was performed on the DNBSEQ-T7 platform using a 150-bp paired-end strategy, generating ~60 Gb of data per sample. Raw reads were processed using fastp (v0.24.1)[48] with parameters '-g -q 5 -u 50 -n 15 -l 150 --overlap_diff_limit 1 --overlap_diff_percent_limit 10'.

Clean reads were aligned to the *P*. sa genome using BWA-MEM algorithm in Burrow-Wheeler Aligner (BWA, 0.7.18-r1243-dirty)[80] with default settings, followed by sorting with SAMtools (v1.21)[81]. Duplicate reads were marked using the MarkDuplicates module in the Genome Analysis Toolkit (GATK, v4.6.1.0)[82].

Population-level SNP and indel calling was conducted using GATK. First, variants were called per sample with HaplotypeCaller using '-ERC GVCF -ploidy 4'. Individual gVCF files were merged with CombineGVCFs and jointly genotyped with GenotypeGVCFs. SNPs and indels were separated using SelectVariants and filtered with VariantFiltration using the following criteria: for SNPs, 'QD < 2.0 || MQ < 40.0 || FS > 60.0 || SOR > 3.0 || MQRankSum < -12.5 || ReadPosRankSum < -8.0'; for indels, 'QD < 2.0 || FS > 200.0 || MQ < 40.0 || SOR > 10.0 || ReadPosRankSum < -20.0'. Multiallelic sites were split into biallelic records using BCFtools norm (v1.21)[81] with '--multiallelics -both'.

SVs were called and genotyped using smoove (v0.2.8)[83] with default parameters. CNVs were detected with CNVpytor (v1.3.2)[84] using a bin size of 100. Finally, all variants (SNPs, indels, SVs and CNVs) were annotated using ANNOVAR (v2020-06-07)[85].

Genome-wide association study (GWAS) analysis

GWAS was performed using the R package GWASpoly (v2.13)[86] based on the Q + K mixed model. First, the VCF files were transformed into dosage files using VCF2dosage function with parameters '-ploidy 4 -min.DP 8 -max.missing 0.5 -min.minor 3'. To control for population structure, the covariance matrix for polygenic effect was calculated using the set.K function with the parameter 'LOCO=TRUE', where a different covariance matrix was calculated for each chromosome according to markers from all other chromosomes[87]. During this step, missing data were imputed using the k-Nearest Neighbors approach implemented by the KNNImputer function from the sklearn.impute module[88]. The imputation was performed five times with different k values (2, 5, 10, 15, and 20), and the average of the five imputed values was used. The set.params function was used to set the threshold of minor allele frequency with the parameter 'geno.freq = 1 - 5/N, MAF=0.05'. The GWASpoly function was then used to assess the significance of each marker associated with target trait variation under the additive model and two simplex dominance models (1-dom-ref and 1-dom-alt) hypothesis. In the additive model, the marker effect scales linearly with the alternate allele dosage. For the simplex dominance models, a single copy of the dominant allele (reference or alternate allele) is sufficient for the complete dominance phenotype, thereby all

657 heterozygotes possessing this allele exhibit equivalent marker effects. Two significance

658 thresholds were adopted: p-value<0.05/variant counts and p-value<0.01/variant counts,

659 according to the Bonferroni adjustment approach. Lastly, the CMplot (v4.5.1)[89] package was

660 used to generate the Manhattan plots. For better visualization, $-\log_{10}(p)$ values greater than 30

661 were truncated to 30 in the Manhattan plots.

Virus-induced gene silencing

Seedlings of *P.* 'Jinbianlinglong' (*P.* ji) were cultivated in a growth room under a 14-h

light/10-h dark photoperiod, with a light intensity of 220 μmol·m^{-2}·s^{-1} and a temperature of

25–27 °C. Partial sequences of *PjPDS* and *PjMYBx1* were introduced into the TRV2-based

pYL156 vector, as illustrated in **Fig. S21**. The resulting TRV2 plasmids were transformed

into *Agrobacterium tumefaciens* strain GV3101. Bacterial cultures containing either TRV2-

PDS or TRV2-*PDS-MYBx1*, and cultures containing the TRV1 plasmids were incubated

separately. When the OD$_{600}$ reached approximately 2.0, the cells were harvested and

resuspended in an induction buffer (10 mM morpholineethanesulfonic acid, 10 mM MgCl$_2$,

and 200 uM acetosyringone) to a final OD$_{600}$ of 2.0. Each TRV2 suspension was mixed with

an equal volume of the TRV1 suspension, and Silwet L-77 was added to a final concentration

of 0.005%. The mixture was incubated statically at room temperature for 3 h.

The resulting *Agrobacterium* mixture was infiltrated into the newly developed inflorescence

stalk and its nearest leaf. After infiltration, plants were kept in darkness for 48 h before being

returned to the growth room. Plants were monitored until phenotypic changes in the flowers

became evident.

To confirm the silencing of target genes in plants showing phenotypic changes, total RNA

was extracted from the corresponding floral tissues using the FastPure Universal Plant Total

RNA Isolation Kit (Vazyme, Nanjing, China). First-strand cDNA was synthesized using

MightyScript Plus First Strand cDNA Synthesis Master Mix (gDNA digester) (Sangon

Biotech, Shanghai, China). Quantitative reverse-transcription PCR (qRT-PCR) was

performed using SupRealQ Ultra Hunter SYBR qPCR Master Mix (U+) (Vazyme). The

$2^{-\Delta\Delta Ct}$ method was employed to calculate the relative expression of target and related genes,

with *PeActin* serving as the internal control. Primer sequences for VIGS and qRT-PCR were

listed in **Table S11.**

Data availability

The raw sequencing data used in this study have been submitted to the NCBI database under BioProject PRJNA1143085 (will be released upon publication), including PacBio HiFi, Hi-C and transcriptome sequences for *P*. sa, and whole genome sequences for 46 H80 individuals. The *P*. sa genome assembly, annotation, homoeologous gene table, and expression matrix for all protein-coding genes will be released in our database (http://www.phaldb.cn/) upon publication.

Code availability

Custom scripts and codes used in this study are provided at GitHub (https://github.com/peipeiwang6/Manuscript/tree/main/2025_Psa_genome).

Acknowledgement

We thank Dr. Huiqi Zhao for sharing the protocol for karyotype analysis, Prof. Zhiqiang Wu, Prof. Xingtan Zhang and Dr. Caiyao Zhao for their helpful suggestions on genome sequencing and assembling, Dr. Xukun Li for advice on anatomical photography and other experiments, Fu Chen for providing seedlings of *P*. 'Jinbianlinglong'. The English language of this manuscript was partially polished using DeepSeek, while all original content was written by the authors and the revised text was adopted following author review.

This work was supported by the National Natural Science Foundation of China (32370241), the Scientific Research Foundation of Kunpeng Institute of Modern Agriculture at Foshan (KIMAQD2022003), and the collaborative project (HXQS2024009 from Prof. Jue Ruan) to P.W.; National Key R&D Program of China (2023YFD1600505) to R.J.; Project supported by the Foundation for Innovative Research Groups of the National Natural Science Foundation of China (32221001) to H.K.; Chinese Universities Scientific Fund (2452024391) to R.Z.

Author contributions

PW, RJ, HK and RZ conceived and designed the study. PW, XZ, LZ, ZL, ZW, FW, YJ, YZ, and LW collected the plant materials of *P*. sa for genome and transcriptome sequencing; RJ conducted the hybridization and provided the plant materials; RJ, LW, YZ and XC conducted the karyotype analysis; PW, LZ, HH and LLan (Lan Lan) assembled the genome; LZ and PW conducted the genome annotation; LZ conducted the functional annotation for protein-coding

genes; LZ, FM and PW conducted the genome assembly quality assessment and syntenic

analysis; PW and FW conducted the transcriptome analysis; PW, XZ and FW conducted the

gene tree building; XZ conducted the genetic variant calling and GWAS analyses; ZW, LLin

(Letian Lin) and ZL conducted the VIGS experiments; LLin performed the qRT-PCR; PW

conducted all the other analyses; PW, RJ, XZ, LZ, LW, ZW and LLin wrote the draft with

contributions from all the other authors. All authors read and approved the final manuscript.

Supplementary figure legends

Fig. S1 Phenotypes of H80 individuals. Each panel shows an individual plant, with its

classifications for the P/A of stripe, background pink color and spot/patch, and lip

morphology provided on the top-left, top-right, bottom-left and bottom-right corners per

panel, respectively. Red arrowheads indicate stripes, while blue ones indicate venation-

associated shadow.

Fig. S2 Genome survey of *P. sa* and karyotype analysis for H80. (A,B) *k*-mer coverage

distribution of *P.* sa sequencing reads, under the assumptions that *P.* sa is a triploid (**A**) or

tetraploid (**B**). (**C**) *k*-mer coverage analysis indicates *P.* sa is an allo-triploid. (**D**) Karyotype

analysis reveals 70 chromosomes per cell in *P.* sa. (**E**) Inferred long terminal repeats (LTRs)

insertion time. (**F**) Karyotype analysis reveals 64 chromosomes per cell in H80 individuals.

Fig. S3 Subgenome assignments of 70 *P.* sa chromosomes according to subgenome-

specific *k*-mers. From outer to inner circles (1-9): (1) subgenome assignments; (2) significant

enrichment of subgenome-specific *k*-mers; (3) normalized proportion of subgenome-specific

k-mers; (4-7) count of subgenome-specific *k*-mers for four sets, respectively; (8) density of

long terminal repeat retrotransposons (LTR-RTs), with gray indicating nonspecific LTR-RTs;

(9) homoeologous blocks.

Fig. S4 Expression profiles of *FAR-RED IMPAIRED RESPONSE 1 (FAR1)* genes in *P.*

sa. *FAR1* genes are grouped into different HG types, as illustrated in **Fig. 2A**. Color scale in

the heatmap indicates log-transformed Transcripts Per Kilobase (TPM) values, where blue

and light blue indicate original TPM=0 and TPM<2, respectively.

Fig. S5 Expression profiles of potential *de novo* new genes. (A-G) Expression profiles of

seven co-expression modules inferred from WGCNA analysis (**Methods**) for type 1 genes

that were highly expressed (TPM>10) in ≥ 1 tissue and had no non-self blastp hists in *P.* sa

genome. *Z*-scores of TPM values among all the tissues are shown. Example genes that were

preferentially expressed in different tissues are indicated.

Fig. S6 Protein sequence alignment (A) and expression (B) of *PsAGL6-2* HGs. HGs with

truncated sequences were indicated with green shadow.

Fig. S7 Manhattan plot for GWAS of lip morphology in the H80 population using HChr

set a as the reference. (**A-D**) GWAS results when SNPs, indels, SVs and CNVs were used,

respectively, with the additive marker-effect model. Horizontal solid line and dashed line

indicate -$\log_{10}$ (0.01/variant counts) and -$\log_{10}$ (0.05/variant counts) (**Methods**), respectively;

red and orange dots denote variants significantly associated with the lip morphology variation;

blue dots represent variants located within or adjacent (±2 kb) to the tandem duplicate cluster

of *PsAGL6-2a* genes.

Fig. S8 Manhattan plot for GWAS of lip morphology in H80 using HChr set b as the

reference. (**A-D**) GWAS results when SNPs, indels, SVs and CNVs were used, respectively,

with the additive marker-effect model. Red and orange dots denote variants significantly

associated with the lip morphology variation; blue dots represent variants located within or

adjacent (±2 kb) to the tandem duplicate cluster of *PsAGL6-2b* genes.

Fig. S9 Manhattan plot for GWAS of lip morphology in H80 using HChr set c as the

reference. (**A-D**) GWAS results when SNPs, indels, SVs and CNVs were used, respectively,

with the additive marker-effect model. Red and orange dots denote variants significantly

associated with the lip morphology variation; blue dots represent variants located within or

adjacent (±2 kb) to the tandem duplicate cluster of *PsAGL6-2c* genes.

Fig. S10 Gene tree for MIKC-type MADS-box genes in orchids. NCBI accession ID (such

as JL017169) was listed between gene name (such as *CeMADS32*) and the reference (such as

2021 Ai).

Fig. S11 Manhattan plot for GWAS of stripe P/A in H80 using the additive marker-

effect model. (**A-D**) GWAS results when SNPs, indels, SVs and CNVs that were called by

referring to HChr set a were used, respectively. (**E-H**) GWAS results when SNPs, indels, SVs

and CNVs that were called by referring to HChr set b were used. Red and orange dots denote

variants significantly associated with the P/A variation of stripe; blue dots represent variants

located within or adjacent (±2 kb) to the tandem duplicate cluster of *PsMYB12a* genes.

Fig. S12 IGV images showing reads of seven H80 individuals with or without stripes

that were mapped onto the *PsMYB12* gene-containing regions. Red arrowheads indicate
stripes, while blue ones indicate venation-associated shadow in individuals which had no
intact reads mapped onto *PsMYB12* genes.

Fig. S13 Evolution of color patterning-related *MYB* genes. (**A**) Gene tree of *MYB* genes
from orchids with genome assemblies. (**B**) Evolutionary history of stripes on perianths in the
genus *Phalaenopsis*. The phylogeny of *Phalaenopsis* species were modified from our
previous review paper (Wang et al., 2025). The red star indicates the origination of stripes in
the most recent common ancestor of *P. equestris* and *P. lindenii*. Three flower images show
the venation-associated distribution of spots in other *Phalaenopsis* species. (**C**) Integrative
Genomics Viewer images showing reads of *Vanda* 'Pakchong Blue' that were mapped onto
genic regions of two *PsMYB12* genes as examples.

Fig. S14 Homoeologous gene variation in sequence and expression of *PsMYB2* and
***PsMYB11*.** (**A**) Protein sequence alignment of *PsMYB2* HGs. (**B**) HG-specific expression of
PsMYB2 in *P*. sa floral organs (upper) and the lips of selected H80 individuals (lower), with
values in the table indicating TPM values. (**C,D**) Protein sequence and promoter DNA
sequence (1 Kb upstream) alignments of *PsMYB11* HGs, respectively. (**E**) HG-specific
expression of *PsMYB11* in *P*. sa floral organs (upper) and the lips of selected H80 individuals
(lower). The HG showing high sequence divergence with others was indicated using green
shadow.

Fig. S15 Manhattan plot for GWAS of background color in H80 using the additive
marker-effect model. (**A-D**) GWAS results when SNPs, indels, SVs and CNVs that were
called by referring to HChr set a were used, respectively. Red and orange dots denote variants
significantly associated with the P/A variation of pink color; blue dots represent variants
located within or adjacent (±2 kb) to *PsMYB2a*.

Fig. S16 Manhattan plot for GWAS of background color in H80 using the 1-dom-ref
marker-effect model. (**A-D**) GWAS results when SNPs, indels, SVs and CNVs that were
called by referring to HChr set b were used, respectively. Red dots denote variants
significantly associated with the P/A variation of pink color; blue dots represent variants
located within or adjacent (±2 kb) to *PsMYB2b*.

Fig. S17 Manhattan plot for GWAS of background color in H80 using the 1-dom-alt
marker-effect model. (**A-D**) GWAS results when SNPs, indels, SVs and CNVs that were

called by referring to HChr set b were used, respectively. Red and orange dots denote

variants significantly associated with the P/A variation of pink color; blue dots represent

variants located within or adjacent (±2 kb) to *PsMYB2b*.

Fig. S18 Manhattan plot for GWAS of patch P/A in H80 using HChr set a as the

reference. (**A-D**) GWAS results when SNPs, indels, SVs and CNVs were used, respectively,

with the additive marker-effect model. Red and orange dots denote variants significantly

associated with the P/A variation of patch; blue dots represent variants located within or

adjacent (±2 kb) to *PsMYB11a*.

Fig. S19 Manhattan plot for GWAS of patch P/A in H80 using HChr set b as the

reference. (**A-D**) GWAS results when SNPs, indels, SVs and CNVs were used, respectively,

with the additive marker-effect model. Red and orange dots denote variants significantly

associated with the P/A variation of patch; blue dots represent variants located within or

adjacent (±2 kb) to *PsMYB11b*.

Fig. S20 Manhattan plot for GWAS of patch P/A in H80 using HChr set c as the

reference. (**A-D**) GWAS results when SNPs, indels, SVs and CNVs were used, respectively,

with the additive marker-effect model. Red and orange dots denote variants significantly

associated with the P/A variation of patch; blue dots represent variants located within or

adjacent (±2 kb) to the tandem duplicate cluster of *PsMYB11c* genes.

Fig. S21 *PjPDS* and *PjMYBx1* sequences and the VIGS constructions. (**A**) CDS and

3'UTR sequence alignment for four *PjPDS* HGs. (**B**) CDS sequence alignment for four

PjMYBx1 HGs. (**C**) Diagrams showing the structure of TRV2 plasmid PYL156, and the

insertion locations of *PjPDS* and *PjMYBx1* sequences. *PjPDS_1* and *PjPDS_2* are sequences

from two *PjPDS* HGs marked with green shadow.

Supplementary tables

Table S1 Comparison of basic statistics for genome assembly and annotation of three

Phalaenopsis species

Table S2 Structural variation summary

Table S3 Statistics of repetitive sequences in *P.* sa genome

Table S4 Statistics of *P.* sa genes and the functional annotations

Table S5 Statistics of noncoding RNAs in *P*. sa genome

Table S6 Information for RNA-seq samples

Table S7 IDs for genes mentioned in the manuscript

Table S8 Allele read ratios of two SNPs in H80 individuals with different lip morphology

Table S9 Association between the presence/absence of stripes and *PsMYB12* genes in 46
H80 individuals

Table S10 Association between *PsMYB2b*-type sequences with background colors

Table S11 Primer sequences used for VIGS and qRT-PCR

Reference

1. Hsiao, Y. *et al.* A novel homodimeric geranyl diphosphate synthase from the orchid

Phalaenopsis bellina lacking a DD(X)$_{2-4}$D motif. *Plant J* **55**, 719–733 (2008).

2. Huang, H. *et al. Terpene synthase-b* and *terpene synthase-e/f* genes produce

monoterpenes for *Phalaenopsis bellina* floral scent. *Front Plant Sci* **12**, 700958 (2021).

3. Hsu, C.-C., Chen, Y.-Y., Tsai, W.-C., Chen, W.-H. & Chen, H.-H. Three R2R3-MYB

transcription factors regulate distinct floral pigmentation patterning in *Phalaenopsis* spp.

Plant Physiol **168**, 175–191 (2015).

4. Chen, W., Hsu, W., Hsu, H. & Yang, C. A tetraspanin gene regulating auxin response

and affecting orchid perianth size and various plant developmental processes. *Plant

Direct* **3**, e00157 (2019).

5. Lu, H.-C. *et al. R2R3-MYB* genes coordinate conical cell development and cuticular wax

biosynthesis in *Phalaenopsis aphrodite. Plant Physiol* **188**, 318–331 (2022).

6. Chuang, Y.-C. *et al.* A dual repeat cis-element determines expression of *GERANYL

DIPHOSPHATE SYNTHASE* for monoterpene production in *Phalaenopsis* orchids. *Front

Plant Sci* **9**, 765 (2018).

7. Hsu, C.-C., Su, C.-J., Jeng, M.-F., Chen, W.-H. & Chen, H.-H. A *HORT1* retrotransposon

insertion in the *PeMYB11* promoter causes harlequin/black flowers in *Phalaenopsis*

orchids. *Plant Physiol* **180**, 1535–1548 (2019).

8. Lu, T.-W., Chen, W.-H., Chen, P.-Y., Shu, Y.-C. & Chen, H.-H. Correction: Perturbation

of periodic spot-generation balance leads to diversified pigmentation patterning of

harlequin *Phalaenopsis* orchids: in silico prediction. *BMC Plant Biol* **24**, 782 (2024).

9. Hsu, H.-F. *et al.* Multifunctional evolution of B and *AGL6* MADS box genes in orchids.

Nat Commun **12**, 902 (2021).

10. Cai, J. *et al.* The genome sequence of the orchid *Phalaenopsis equestris*. *Nat Genet* **47**,

65–72 (2015).

11. Chao, Y. *et al.* Chromosome-level assembly, genetic and physical mapping of

Phalaenopsis aphrodite genome provides new insights into species adaptation and

resources for orchid breeding. *Plant Biotechnol J* **16**, 2027–2041 (2018).

12. Bashton, M. & Chothia, C. The generation of new protein functions by the combination

of domains. *Structure* **15**, 85–99 (2007).

13. Vishwanath, S., de Brevern, A. G. & Srinivasan, N. Same but not alike: Structure,

flexibility and energetics of domains in multi-domain proteins are influenced by the

presence of other domains. *PLoS Comput Biol* **14**, e1006008 (2018).

14. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. *J R Stat Soc Series B* **57**, 289–300 (1995).

15. Liu, Y. *et al. Arabidopsis FHY3* and *FAR1* regulate the balance between growth and

defense responses under shade conditions. *Plant Cell* **31**, 2089–2106 (2019).

16. Zhao, L., Saelao, P., Jones, C. D. & Begun, D. J. Origin and spread of *de novo* genes in

Drosophila melanogaster populations. *Science* **343**, 769–772 (2014).

17. Dion-Côté, A.-M. Spermatogenesis: A hotspot for new genes. *eLife* **8**, e50136 (2019).

18. Yang, H. *et al.* Global modulation of gene expression and transcriptome size in aneuploid

combinations of maize. *Proc Natl Acad Sci U S A* **122**, e2426749122 (2025).

19. Pavelka, N. *et al.* Aneuploidy confers quantitative proteome changes and phenotypic variation in budding yeast. *Nature* **468**, 321–325 (2010).

20. Selmecki, A., Forche, A. & Berman, J. Aneuploidy and isochromosome formation in drug-resistant *Candida albicans*. *Science* **313**, 367–370 (2006).

21. Wang, F. *et al.* Molecular mechanisms underlying floral trait formation in *Phalaenopsis* orchids. *Hortic Res* uhaf340 (2025) doi:10.1093/hr/uhaf340.

22. Chen, W.-H., Jiang, Z.-Y., Hsu, H.-F. & Yang, C.-H. Silencing of *FOREVER YOUNG FLOWER* -like genes from *Phalaenopsis* orchids promotes flower senescence and abscission. *Plant Cell Physiol* **62**, 111–124 (2021).

23. Wang, R., Mao, C. & Ming, F. PeMYB4L interacts with PeMYC4 to regulate anthocyanin biosynthesis in *Phalaenopsis* orchid. *Plant Sci* **324**, 111423 (2022).

24. Marçais, G. & Kingsford, C. A fast, lock-free approach for efficient parallel counting of occurrences of k-mers. *Bioinformatics* **27**, 764–770 (2011).

25. Ranallo-Benavidez, T. R., Jaron, K. S. & Schatz, M. C. GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes. *Nat Commun* **11**, 1432 (2020).

26. Liu, B. *et al.* Estimation of genomic characteristics by analyzing k-mer frequency in de novo genome projects. Preprint at https://doi.org/10.48550/ARXIV.1308.2012 (2013).

27. Cheng, H., Concepcion, G. T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo assembly using phased assembly graphs with hifiasm. *Nat Methods* **18**, 170–175 (2021).

28. Vasimuddin, Md., Misra, S., Li, H. & Aluru, S. Efficient architecture-aware acceleration of BWA-MEM for multicore systems. in *2019 IEEE International Parallel and Distributed Processing Symposium (IPDPS)* 314–324 (IEEE, Rio de Janeiro, Brazil, 2019). doi:10.1109/IPDPS.2019.00041.

29. Zeng, X. *et al.* Chromosome-level scaffolding of haplotype-resolved assemblies using

Hi-C data without reference genomes. *Nat Plants* **10**, 1184–1200 (2024).

30. Durand, N. C. *et al.* Juicebox provides a visualization system for Hi-C contact maps with

unlimited zoom. *Cell Syst* **3**, 99–101 (2016).

31. Seppey, M., Manni, M. & Zdobnov, E. M. BUSCO: assessing genome assembly and

annotation completeness. *Methods Mol Biol* **1962**, 227–245 (2019).

32. Jia, K. *et al.* SubPhase: a robust allopolyploid subgenome phasing method based on

subgenome-specific *k*-mers. *New Phytol* **235**, 801–809 (2022).

33. Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for

genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

34. Rhie, A., Walenz, B. P., Koren, S. & Phillippy, A. M. Merqury: reference-free quality,

completeness, and phasing assessment for genome assemblies. *Genome Biol* **21**, 245

(2020).

35. Li, K., Xu, P., Wang, J., Yi, X. & Jiao, Y. Identification of errors in draft genome

assemblies at single-nucleotide resolution for quality assessment and improvement. *Nat*

Commun **14**, 1–12 (2023).

36. Ellinghaus, D., Kurtz, S. & Willhoeft, U. LTRharvest, an efficient and flexible software

for de novo detection of LTR retrotransposons. *BMC Bioinformatics* **9**, 18 (2008).

37. Ou, S. & Jiang, N. LTR_FINDER_parallel: parallelization of LTR_FINDER enabling

rapid identification of long terminal repeat retrotransposons. *Mobile DNA* **10**, 48 (2019).

38. Ou, S., Chen, J. & Jiang, N. Assessing genome assembly quality using the LTR

Assembly Index (LAI). *Nucleic Acids Res* **46**, e126 (2018).

39. Ou, S. & Jiang, N. LTR_retriever: a highly accurate and sensitive program for

identification of long terminal repeat retrotransposons. *Plant Physiol* **176**, 1410–1422

(2018).

40. Lin, Y. *et al.* quarTeT: a telomere-to-telomere toolkit for gap-free genome assembly and

centromeric repeat identification. *Hortic Res* **10**, uhad127 (2023).

41. Chan, P. P., Lin, B. Y., Mak, A. J. & Lowe, T. M. tRNAscan-SE 2.0: improved detection

and functional classification of transfer RNA genes. *Mol Biol Evol* **49**, 9077–9096 (2021).

42. Nawrocki, E. P. Annotating functional RNAs in genomes using Infernal. *Methods Mol*

Biol **1097**, 163–197 (2014).

43. Ou, S. *et al.* Benchmarking transposable element annotation methods for creation of a

streamlined, comprehensive pipeline. *Genome Biol* **20**, 275 (2019).

44. Yan, H., Bombarely, A. & Li, S. DeepTE: a computational method for de novo

classification of transposons with convolutional neural network. *Bioinformatics* **36**,

4269–4275 (2020).

45. Zhang, W. *et al.* Genome sequence of *Apostasia ramifera* provides insights into the

adaptive evolution in orchids. *BMC Genomics* **22**, 536 (2021).

46. Benson, G. Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids*

Res **27**, 573–580 (1999).

47. Smit, A. & Hubley, R. RepeatModeler Open-1.0. 2008-2015

<http://www.repeatmasker.org>.

48. Chen, S. Ultrafast one-pass FASTQ data preprocessing, quality control, and

deduplication using fastp. *iMeta* **2**, e107 (2023).

49. Keilwagen, J., Hartung, F. & Grau, J. GeMoMa: homology-based gene prediction

utilizing intron position conservation and RNA-seq data. *Methods Mol Biol* **1962**, 161–

177 (2019).

50. Zhang, G.-Q. *et al.* The *Apostasia* genome and the evolution of orchids. *Nature* **549**, 379–

383 (2017).

51. Zhang, G.-Q. *et al.* The *Dendrobium catenatum* Lindl. genome sequence provides

insights into polysaccharide synthase, floral development and adaptive evolution. *Sci Rep*
6, 19029 (2016).

52. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–
915 (2019).

53. Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from
RNA-seq reads. *Nat Biotechnol* **33**, 290–295 (2015).

54. Grabherr, M. G. *et al.* Trinity: reconstructing a full-length transcriptome without a
genome from RNA-Seq data. *Nat Biotechnol* **29**, 644–652 (2011).

55. Haas, B. J. *et al.* Improving the *Arabidopsis* genome annotation using maximal transcript
alignment assemblies. *Nucleic Acids Res* **31**, 5654–5666 (2003).

56. Gabriel, L. *et al.* BRAKER3: Fully automated genome annotation using RNA-seq and
protein evidence with GeneMark-ETP, AUGUSTUS, and TSEBRA. *Genome Res* **34**,
769–777 (2024).

57. Blanco, E., Parra, G. & Guigó, R. Using geneid to identify genes. *Curr Protoc*
Bioinformatics **18**, (2007).

58. Majoros, W. H., Pertea, M. & Salzberg, S. L. TigrScan and GlimmerHMM: two open
source ab initio eukaryotic gene-finders. *Bioinformatics* **20**, 2878–2879 (2004).

59. Haas, B. J. *et al.* Automated eukaryotic gene structure annotation using
EVidenceModeler and the program to assemble spliced alignments. *Genome Biol* **9**, R7
(2008).

60. Haas, B. J., Zeng, Q., Pearson, M. D., Cuomo, C. A. & Wortman, J. R. Approaches to
fungal genome annotation. *Mycology* **2**, 118–141 (2011).

61. Altschul, S. F. *et al.* Gapped BLAST and PSI-BLAST: a new generation of protein
database search programs. *Nucleic Acids Res* **25**, 3389–3402 (1997).

62. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification.

Bioinformatics **30**, 1236–1240 (2014).

63. Trapnell, C., Pachter, L. & Salzberg, S. L. TopHat: discovering splice junctions with

RNA-Seq. *Bioinformatics* **25**, 1105–1111 (2009).

64. Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and phylogenetically

annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids*

Res **47**, D309–D314 (2019).

65. Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J.

eggNOG-mapper v2: functional annotation, orthology assignments, and domain

prediction at the metagenomic scale. *Mol Biol Evol* **38**, 5825–5829 (2021).

66. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421

(2009).

67. Xu, D. *et al.* GFAP: ultrafast and accurate gene functional annotation software for plants.

Plant Physiol **193**, 1745–1748 (2023).

68. Schläpfer, P. *et al.* Genome-wide prediction of metabolic enzymes, pathways, and gene

clusters in plants. *Plant Physiol.* **173**, 2041–2059 (2017).

69. Jin, J. *et al.* PlantTFDB 4.0: toward a central hub for transcription factors and regulatory

interactions in plants. *Nucleic Acids Res* **45**, D1040–D1045 (2017).

70. Goel, M., Sun, H., Jiao, W.-B. & Schneeberger, K. SyRI: finding genomic

rearrangements and local sequence differences from whole-genome assemblies. *Genome*

Biol **20**, 277 (2019).

71. Wang, Y. *et al.* MCScanX: a toolkit for detection and evolutionary analysis of gene

synteny and collinearity. *Nucleic Acids Res* **40**, e49–e49 (2012).

72. Putri, G. H., Anders, S., Pyl, P. T., Pimanda, J. E. & Zanini, F. Analysing high-

throughput sequencing data in Python with HTSeq 2.0. *Bioinformatics* **38**, 2943–2945

(2022).

73. McCarthy, D. J., Campbell, K. R., Lun, A. T. L. & Wills, Q. F. Scater: pre-processing,

quality control, normalization and visualization of single-cell RNA-seq data in R.

Bioinformatics **33**, 1179–1186 (2017).

74. R Core Team. R: a language and environment for statistical computing. R foundation for

statistical computing, Vienna. (2023).

75. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 559 (2008).

76. Wei, Y. *et al.* OrchidMD: an integrated and user-interactive orchid multi-omics database

for mining genes and biological research. *Plant Biotechnol J* pbi.70445 (2025)

doi:10.1111/pbi.70445.

77. Kumar, S., Stecher, G. & Tamura, K. MEGA7: molecular evolutionary genetics analysis

version 7.0 for bigger datasets. *Mol Biol Evol* **33**, 1870–1874 (2016).

78. Darriba, D. *et al.* ModelTest-NG: a new and scalable tool for the selection of DNA and

protein evolutionary models. *Mol Biol Evol* **37**, 291–294 (2020).

79. Kozlov, A. M., Darriba, D., Flouri, T., Morel, B. & Stamatakis, A. RAxML-NG: a fast,

scalable and user-friendly tool for maximum likelihood phylogenetic inference.

Bioinformatics **35**, 4453–4455 (2019).

80. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

Preprint at https://doi.org/10.48550/ARXIV.1303.3997 (2013).

81. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008

(2021).

82. Poplin, R. *et al.* Scaling accurate genetic variant discovery to tens of thousands of

samples. Preprint at https://doi.org/10.1101/201178 (2017).

83. Pedersen, B. S. smoove: structural-variant calling and genotyping with existing tools.

(2020).

84. Suvakov, M., Panda, A., Diesh, C., Holmes, I. & Abyzov, A. CNVpytor: A tool for copy number variation detection and analysis from read depth and allele imbalance in whole-genome sequencing. *GigaScience* **10**, giab074 (2021).

85. Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic Acids Res* **38**, e164 (2010).

86. Rosyara, U. R., Jong, W. S. D., Douches, D. S. & Endelman, J. B. Software for genome-wide association studies in autopolyploids and its application to potato. *Plant Genome* **9**, plantgenome2015.8.73 (2016).

87. Yang, J., Zaitlen, N. A., Goddard, M. E., Visscher, P. M. & Price, A. L. Advantages and pitfalls in the application of mixed-model association methods. *Nat Genet* **46**, 100–106 (2014).

88. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *J Mach Learn Res* **12**, 2825–2830 (2011).

89. Yin, L. CMplot: Circle Manhattan Plot. (2024).