CAN1-mediated autophagosome size regulation fine-tunes phosphate starvation tolerance in *Arabidopsis thaliana*

David Görg[1], Fiona Smith[1], Klara Machalett[1,2], Falko Koblenz[1], Ymy Thi Ngo[1], Sascha Höhne[1], Julian Arndt[1], Sylvestre Marillonnet[3], Nenad Grujic[4], Richard Imre[4,5], Karl Mechtler[4,5], Yasin Dagdas[6], and Christin Naumann[1*]

[1]Department of Molecular Signal Processing, Leibniz Institute of Plant Biochemistry, 06120 Halle (Saale), Germany

[2]Institute of Biochemistry and Biotechnology, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

[3]Department of Cell and Metabolic Biology, Leibniz Institute of Plant Biochemistry, 06120 Halle (Saale), Germany

[4]Gregor Mendel Institute of Molecular Plant Biology, Vienna, Austria

[5]Research Institute of Molecular Pathology (IMP), Vienna BioCenter (VBC), Vienna, Austria

[6]Heidelberg University, Centre for Organismal Studies, Heidelberg, Germany

*Corresponding author e-mail: cnaumann@ipb-halle.de

ORCID IDs: 0000-0002-9712-5728 (D.G.); 0009-0004-4740-3197 (K.M.); 0000-0002-4263-797X (S.M.); 0000-0002-9502-355X (Y.D.); 0000-0001-6972-155X (C.N.)

ABSTRACT

Inorganic phosphate (Pi) availability determines root development and plant performance. In *Arabidopsis thaliana*, external Pi is sensed by root tips and Pi limitation triggers ER stress–induced autophagy, yet the physiological impact and mechanisms controlling autophagosome biogenesis remain unclear. Here, we identify CAN1 (COMPONENT OF AUTOPHAGIC NETWORK) as a novel plant-specific regulator of autophagosome size in local Pi sensing. CAN1 associates with the plasma membrane-ER tethering protein VAP27-1 and binds to Pi-responsive ATG8 isoforms through conserved interaction motifs. Loss of CAN1 augments ER stress resistance by reducing autophagosome size in Pi-deprived root tips. Our findings establish CAN1 as a determinant of autophagosome size at the ER membrane. Thus, CAN1 functions as a molecular link between Pi-dependent ER stress signaling and fine-tuning of autophagic capacity.

INTRODUCTION

Plant growth and development are tightly constrained by nutrient availability in soil. Phosphorus, primarily absorbed as inorganic phosphate (Pi), is the second most critical macronutrient, and Pi availability is often limited by complex chemical interactions with associated metals (e.g. Fe, Al) and their (Abel 2017; Abel and Naumann 2024; Lambers et al. 2015; Shen et al. 2011). To enhance Pi acquisition for maintaining cellular Pi homeostasis, plants undergo extensive developmental and metabolic reprogramming, which includes remodeling of root system architecture and chemical modification of the rhizosphere, respectively (Crombez et al. 2019; Giehl and von Wiren 2014; Kellermeier et al. 2014; Yang et al. 2024). In growing root tips of *Arabidopsis thaliana*, low external Pi rapidly reduces cell elongation and progressively inhibits cell division (Balzergue et al. 2017; Clua et al. 2024; Maniero et al. 2024; Muller et al. 2015; Naumann et al. 2022). The single ER-resident P5A-type ATPase PHOSPHATE DEFICIENCY RESPONSE 2 (PDR2) and the apoplastic ferroxidase LOW PHOSPHATE ROOT 1 (LPR1) form the core module of local Pi sensing in root tips (Svistoonoff et al. 2007; Ticconi et al. 2001; Ticconi et al. 2009). Decreasing Pi-availability activates LPR1 by facilitating access to its substrate (Fe^{2+}), triggering cell-type-specific Fe^{3+} accumulation, formation of reactive oxygen species (ROS), and cell wall modifications, which interrupts cellular communication and consequently root growth (Balzergue et al. 2017; Gutierrez-Alanis et al. 2017; Muller et al. 2015; Naumann et al. 2022). PDR2 counteracts LPR1 function by maintaining root tip Fe homeostasis through an unknown mechanism (Jakobsen et al. 2005; Muller et al. 2015; Naumann et al. 2022; Naumann et al. 2019; Ticconi et al. 2009). Biochemical and structural studies reveal that P5A-type ATPases (P5As) act as membrane-helix dislocases that assist transmembrane protein insertion and remove mistargeted proteins, thereby safeguarding ER function and organelle integrity (Ji et al. 2024; Li et al. 2024; McKenna et al. 2022; McKenna et al. 2020). Disruption of P5A activity is detrimental across eukaryotes, affecting neuronal development in animals and

causing pleiotropic ER-associated defects in yeast (Feng et al. 2020; Li et al. 2021; Qin et al.

2020; Sim and Park 2023). In plants, PDR2 function in root tips is linked to Pi-dependent ER

stress-induced autophagy (Naumann et al. 2019; Stephani et al. 2020; Ticconi et al. 2009).

Autophagy is a conserved degradation pathway essential for protein and organelle quality

control, cellular homeostasis, and stress adaptation (Gross et al. 2025; Liu and Bassham 2012;

Petersen et al. 2024; Su et al. 2020). Damaged organelles or protein aggregates are

sequestered into double-membrane autophagosomes, typically 0.5–1.5 μm in diameter, which

subsequently fuse with the vacuole for cargo degradation (Nakatogawa 2020; Zhao et al.

2022; Zheng et al. 2018). Autophagosome biogenesis originates at ER-associated phagophore

initiation sites through PI3K complex (ATG6–VPS15–VPS34–ATG14) recruitment and

progresses via ATG8 lipidation, mediated by the ATG12–ATG5–ATG16 complex.

Additional regulators, such as SH3P2, interact with ATG8 and the PI3K complex to support

phagophore expansion, cargo loading, vacuolar targeting, and possibly autophagosome size

control (Nascimbeni et al. 2017; Walczak and Martens 2013; Zhuang et al. 2013). Selective

autophagy receptors, including NBR1 (Jung et al. 2020; Svenning et al. 2011) and C53

(Stephani et al. 2020), deliver defined cargos to the phagophore through ATG8 interaction

motifs (AIMs), and cargo characteristics can directly influence autophagosome dimension

(Johansen and Lamark 2020). ATG8 isoforms enable organ- and stress-specific autophagy

responses as illustrated by the differential complementation of carbon and nitrogen starvation

phenotypes (Del Chiaro et al. 2025; Dong et al. 2025; Liu et al. 2021; Thompson et al. 2005).

Beyond nutrient starvation, local Pi sensing in root tips and Pi deficiency in green tissues

triggers iron-mediated ER stress-induced autophagy, likely originating from cargo overload

within the secretory pathway (Naumann et al. 2019; Yoshitake et al. 2022). Notably,

autophagosome size is highly variable and strongly shaped by the nature of the target, yet how

autophagosome dimensions are controlled during biogenesis remains unknown. Emerging

evidence suggests that ER-associated proteins at phagophore initiation sites are likely central

to autophagosome size determination; however, the molecular mechanism and components

involved remain elusive.

Here, using quantitative proteomics and interaction assays in Arabidopsis, we report a novel

COMPONENT OF THE AUTOPHAGIC NETWORK (CAN1) as a plant-specific regulator

of autophagosome size during Pi limitation. CAN1 associates with the ER–plasma membrane

tethering protein VAP27-1 and the ER-membrane embedded glycosylation complex

component DGL1, interacts with root tip-localized ATG8B and ATG8C isoforms via

conserved AIM motifs, and its loss of function results in smaller autophagosomes and

enhanced ER stress resistance. Through its role in Pi-mediated autophagy, CAN1 shapes

autophagosome biogenesis, establishing an ER-associated regulatory network that converges

with ATG8-mediated, organ-specific autophagy.

RESULTS

Phosphate limitation induces ATG8 isoform specific ER stress mediated autophagy

Tissue specific expression of ATG8 isoforms (ATG8A-ATG8I) in *Arabidopsis thaliana*

mediate the autophagic response under distinct stress conditions (Del Chiaro et al. 2025;

Dong et al. 2025). To investigate their roles in local root Pi sensing, we monitored native

promoter-driven GFP-ATG8 protein fusions (*pATG8x::GFP-ATG8x*) (Del Chiaro et al. 2025)

for autophagosome formation in wildtype (WT) seedlings transferred from Pi-replete (+Pi) to

Pi-deprived (-Pi) conditions as well as after exposure to carbon starvation (-C) and

tunicamycin-induced (+TM) ER stress (Fig. S1). Consistent with previous reports, GFP

fluorescence was largely absent in basal (+Pi) condition for most ATG8 isoforms with the

exception of ATG8A and ATG8F (Del Chiaro et al. 2025; Dong et al. 2025; Thompson et al.

2005). Upon carbon starvation, all tested lines exhibited GFP-positive puncta formation in the

root transition zone and, to a lower extent, in the root tip, indicative of bulk autophagy

activation independent of isoform specificity (Fig. S1). Likewise, TM treatment induced

autophagosome formation in the transition zone of all transgenic GFP-ATG8 lines. Notably,

Pi deprivation selectively activated autophagosome formation in root tips expressing GFP-

ATG8 fusion isoforms B, C, and E suggesting an ATG8 isoform-specific involvement in Pi

sensing (Fig. 1a, Fig. S1). Given that local root Pi sensing triggers ER stress-mediated

autophagy in Pi limitation via the PDR2–LPR1 module (Naumann et al. 2019; Stephani et al.

2020), we introgressed *pATG8B/C::GFP-ATG8B/C* transgenes into the Pi-sensitive

pdr2 mutant and monitored autophagosome formation following Pi deprivation. When

compared to +Pi conditions, the number of GFP-ATG8B labelled autophagosomes increased

2-fold and 2.5-fold in wildtype and *pdr2* roots, respectively, with no significant changes in

ATG8 transcript levels (Fig. 1, Fig. S2a). Interestingly, the number of GFP-ATG8C labelled

puncta increased significantly only in *pdr2,* suggesting a specific role in ER quality control,

which is supported by enhanced autophagosome formation upon TM treatment (Fig. 1).

To assess ATG8 isoform-specific functionality, we examined local Pi sensing in the ATG8-

deficient nonuple mutant *Δatg8* (Del Chiaro et al. 2025), alongside complementation lines

expressing GFP-ATG8A or GFP-ATG8H, representing the two major ATG8 clades in

Arabidopsis (Rogov et al. 2023). *Δatg8* mutants exhibited modest root growth defects in Pi

deprivation, indicative of enhanced ER stress sensitivity (Fig. S2b). Notably, ectopic

expression of either ATG8A or ATG8H did not rescue this phenotype, implying that

autophagosome induction in local Pi sensing may involve isoform-specific functions.

Local Pi sensing remodels autophagy-dependent root tip proteome

To elucidate the biological role of ER stress-induced autophagy in local Pi sensing and

uncover novel components, we employed two complementary mass spectrometry-based

screening approaches. Our working hypothesis posits that proteins regulated in an autophagy-

dependent mode during Pi sensing should (i) exhibit altered abundances in autophagy-

deficient mutants such as *atg2*, and (ii) physically interact with ATG8 isoforms. First, we

performed proteomic profiling of wildtype, *pdr2*, and *atg2* root tips using tandem mass tag-based quantitative mass spectrometry (TMT-MS). Seedlings were germinated in Pi-sufficiency and subsequently transferred to control (+Pi) or ER stress-inducing (−Pi or +TM) conditions. Comparative analysis revealed that either ER stress condition caused pronounced proteomic shifts in root tips, with *atg2* displaying a significantly higher number of differentially abundant proteins relative to wildtype and *pdr2* (Fig. 2a, Fig. S3a, Data S1). We next analyzed the protein abundances in ER stress conditions in *atg2* in comparison to WT or *pdr2*. For Pi-deprivation 89 proteins were increased in *atg2* but not in wildtype or *pdr2* root tips, suggesting autophagy-dependent regulation (Fig.2b, Fig. S3b). This set includes core autophagy components such as ATG4, multiple ATG8 isoforms, and selective autophagy receptors, NBR1 and C53, previously implicated in ER stress responses and Pi signaling (Picchianti et al. 2023; Stephani et al. 2021; Stephani et al. 2020; Wild et al. 2014). Enrichment of these proteins reinforced the validity of our experimental design and highlights a functional role of autophagy in maintaining proteostasis during local Pi perception. Second, to identify Pi-responsive ATG8 interactors, we performed immunoprecipitation coupled with mass spectrometry (IP-MS) using wildtype and *pdr2* lines expressing *p35S::GFP-ATG8A* (Fig. S3d, e; Data S2). This screen revealed over 300 potential interactors, including ATG8B and ATG8C as well as NBR1 and C53, which are established ATG8 interactors (Stephani et al. 2020; Wilfling et al. 2020; Zhao et al. 2022). To discover functionally relevant ATG8-associated proteins in Pi-deprivation, we cross-referenced the ATG8A interactome (Fig. 2c,d) with the set of 89 proteins stabilized in *atg2* (Fig. 2b), which revealed ATG8B, ATG8C, NBR1 and an uncharacterized protein (AT5G40690), among the most enriched candidates across both datasets (Fig. 2e). Consistent detection of AT5G40690, suggests a novel function in autophagy-dependent Pi sensing and root proteostasis. We therefore refer to AT5G40690 as CAN1 (COMPONENT OF AUTOPHAGIC NETWORK).

CAN1 is evolutionally conserved among land plant species

While the core autophagy machinery is conserved across plants, the identification of

modulators fine-tuning the autophagic pathway remains an active area of investigation

(Cadena-Ramos and De-la-Pena 2024; Gross et al. 2025; Stephani and Dagdas 2020).

AT5G40690 encodes a small, 210-amino-acid protein, expressed from a single exon.

Phylogenetic analysis revealed that CAN1 is exclusive to and conserved across mostly all

major land plant lineages apart from ferns and gymnosperms (Fig. 3a, Fig. S4).

Notably, in a subset of angiosperms, including both monocot and dicot species but not

bryophytes and *Amborella trichopoda*, we identified additional gene homologs encoding

truncated proteins (~110 amino acid residues), that lack the C-terminal domain present in

canonical CAN1 orthologs (Fig. 3a, Fig. S4). In Arabidopsis, two such paralogs, HRG1

(AT2G411730) and HRG2 (AT5G24640) have been implicated in reactive oxygen species

(ROS)-mediated root meristem maintenance (Gong et al. 2021). These findings suggest

functional divergence or neofunctionalization of HRG proteins within specific angiosperm

lineages such as the Brassicaceae.

Expression analysis in whole seedlings revealed consistent upregulation of *CAN1* transcript

levels in autophagy-deficient mutants (*atg2*, *atg5*, *atg7*, and *atg9*) compared to the wildtype

control (Fig. 3b). To investigate organ- and tissue-specific *CAN1* expression, we generated

transgenic *pCAN1::GUS* wildtype and *atg5* mutant lines (Fig. 3b-d). In wildtype, *CAN1*

promoter activity was predominantly detected in root tissues, including lateral roots and root

hairs, with increased activity in the root-shoot junction, while only weak GUS reporter signals

were observed in the tip region of cotyledons and true leaves as well as in flowers and

developing siliques (Fig. 3b,c). Interestingly, in the *atg5* background, the spatial distribution

of *CAN1* promoter activity was comparable to wildtype, but markedly enhanced as evidenced

by intense GUS staining, which correlates to elevated *CAN1* transcript levels measured in

autophagy-deficient lines (Fig. 3b, d). Collectively, the evolutionary conservation of *CAN1*,

its dominant expression in roots, and its transcriptional activation in autophagy mutants

support a role for *CAN1* in nutrient-dependent ER stress responses in land plants.

CAN1 interacts with ATG8 in an AIM-dependent manner

Co-immunoprecipitation of endogenous ATG8 with CAN1-mCherry expressed in

Arabidopsis roots experiencing ER stress (Fig. 4a) validated our initial IP-MS identification

of CAN1 as an ATG8 interactor (Fig. 2). We combined structural modeling with *in vitro* and

in vivo studies to determine the ATG8-binding site of CAN1. AlphaFold2 modelling indicated

CAN1 to be intrinsically disordered and predicted the presence of five putative ATG8-

interacting motif (AIM) consensus sequences (W/F/V-X-X-L/I/V), with AIM5 (aa 202-205)

being the best candidate followed by AIM4 (aa 183-189) (Fig. 4b; Fig. S5). Yeast two-hybrid

interaction assays indicated AIM5-dependent CAN1 binding to all ATG8 isoforms, and

strong interaction with ATG8G, H, and I (Fig. 4c). Transient expression of GFP fusions of

CAN1 in tobacco (*Nicotiana benthamiana*) and live cell imaging showed predominantly

cytosolic localization with partial net-like association reminiscent of ER structures (Fig. 4d).

Importantly, the subcellular localization observed for GFP-CAN1 was not altered by

introducing AIM4 and/or AIM5 deletions (Fig. 4e). Subsequent co-immunoprecipitation

analysis demonstrated the dominant role of AIM5 in CAN1 for ATG8 binding. While GFP-

CAN1 interacted with all RFP-ATG8 isoforms in tobacco leave extracts, which was also true

for GFP-CAN1$^{\Delta AIM4}$, deletion of AIM5 (GFP-CAN1$^{\Delta AIM5}$) abrogated CAN1 interaction with

most ATG8 isoforms except for ATG8E and ATG8G. As expected, deletion of AIM4 and

AIM5 abolished CAN1$^{\Delta AIM4/5}$ binding to any ATG8 isoform (Fig. 4f; Fig. S6). Collectively,

our results demonstrate differential interaction of CAN1 with ATG8 isoforms, which is

primarily mediated by AIM5.

CAN1 functions in ER stress-dependent autophagy during local Pi sensing

To investigate the relevance of CAN1–ATG8 interaction *in vivo* and its role for root

responses to ER stress, we monitored autophagosome formation in transgenic lines expressing

GFP-tagged CAN1 (*pCAN1::CAN1-GFP; pUBQ10::GFP-CAN1*) (Fig. 5a-d, Fig. S7).

Consistent with transient expression in tobacco (Fig. 4d, e), CAN1-GFP exhibited a diffuse cytoplasmic distribution under control conditions, occasionally forming low-intensity puncta, which remained unchanged when expressed in the low Pi-sensitive *pdr2* mutant (Fig. 5a, Fig. S7a ,c).

Induction of ER stress by low Pi or TM treatment caused a rapid and robust increase in CAN1-GFP puncta and partial net-like localization with further enhancement in *pdr2* root cells (Fig. 5a,b). Importantly, CAN1 transcript levels and spatial expression in root tips remained unaffected in all lines tested (Fig. S7e,f). CAN1-GFP puncta co-localized with mCherry-ATG8E, demonstrating CAN1 association with autophagosomes (Fig. 5c,d, Fig. S7; (Feng et al. 2022)). Consistently, no CAN1-GFP puncta were detected in the autophagy-deficient *atg5* mutant although basal CAN1-GFP protein abundance was increased in *atg2* and *atg5* (Fig. S8d). These findings demonstrate that CAN1 associates with the ER and autophagosomes during ER stress in root tips, with a more pronounced response under Pi limitation compared to pharmacological ER stress induction by TM (Fig. 5a–d).

To investigate CAN1 function in *Arabidopsis*, we first analyzed T-DNA insertion (3'UTR) line (*SALK_092190*), which does not affect transcript abundance (Fig. S8a, b). We then generated two independent loss-of-function alleles using CRISPR–Cas9-mediated genome editing (Stuttmann et al. 2021): *can1-c1*, carrying an insertion resulting in a frameshift and premature stop codon, and *can1-c3*, harboring a deletion predicted to have a similar effect (Fig. S8b-c). Both mutant lines were phenotypically indistinguishable from wildtype during germination and early root development (Fig. 5e, f). Under ER stress conditions, however, both *can1* loss-of-function alleles exhibited increased survival on TM-containing medium, indicating enhanced resistance (Fig. 5e). By contrast, no phenotypic differences were observed under carbon or nitrogen starvation, a condition commonly used to activate bulk autophagy (Fig. S8f-h).

We next examined the role of CAN1 in local Pi sensing. While both *can1* mutants were

significantly less sensitive to Pi deprivation, exhibiting enhanced primary root elongation,

CAN1 overexpression (*pUBQ10::GFP-CAN1*) displayed the opposite root growth phenotype

on low Pi medium (Fig. 5f). However, the *CAN1* loss- and gain-of-function lines tested

were indistinguishable with respect to iron and callose accumulation in root tips, two

histochemical markers of local Pi sensing (Naumann et al. 2022) (Fig. S9).

Thus, our observations suggest that CAN1 is dispensable for bulk autophagy but functions in

ER stress-induced autophagy modulating root growth response under local Pi limitation.

CAN1 facilitates autophagosome biogenesis through ER tethering with VAP27-1

To elucidate a role of CAN1 in autophagy, we analysed the CAN1 interactome under basal

and ER stress conditions. Under ER stress, CAN1 associates with ATG8C and factors

involved in autophagosome maturation (ATG3, FREE1, SH3P2, VPS34, NBR1) and the ER

membrane (VAP27-1, DGL1), implicating CAN1 function in phagophore expansion (Fig. 6a;

Fig. S10a). Co-immunoprecipitation confirmed CAN1 interaction with NBR1 in wildtype and

atg5 roots, restricted to ER stress in wildtype and stabilized by ConA treatment (Fig. 6b).

Considering the CAN1 localization and interactome under stress (Fig. 5c, Fig. 6a, Fig. S10a),

we tested interactions of CAN1 with DGL1, an ER quality control component of the OST

complex (Lerouxel et al. 2005), and with VAP27-1, an ER–PM tether involved in endocytosis

and mitophagy (Li et al. 2022; Man et al. 2024; Stefano et al. 2018; Wang et al. 2019). CAN1

co-immunoprecipitated with DGL1 (Fig. S10b), supporting ER recruitment of CAN1.

VAP27-1 also interacted with CAN1 in tobacco leaves and Arabidopsis roots under Pi

limitation or TM-induced ER stress (Fig. 6c,d), suggesting that CAN1 may anchor

phagophores to the ER via VAP27-1 interaction and, enable autophagosome membrane

expansion. Notably, GFP-ATG8A puncta were smaller in *can1-c1* root tips under Pi

limitation, while NBR1 turnover and autophagosome amount remained unaffected (Fig. 6e,f,

Fig. S10d, e), consistent with impaired phagophore expansion. CAN1 protein abundance

remained unchanged in ER stress (Fig. S8e, S10c).

DISCUSSION

Autophagy is essential for plant stress adaptation by maintaining cellular homeostasis under

fluctuating environmental conditions, yet the mechanisms determining autophagosome

maturation and size are largely unknown (Gross et al. 2025; Nakatogawa 2020; Zhao et al.

2022). In Arabidopsis, sensing of Pi limitation provokes iron-dependent ER stress response

followed by selective autophagy induction in growing root tips and photosynthetic tissues;

however the molecular determinants involved remain to be elucidated (Naumann et al. 2019;

Yoshitake et al. 2021; Yoshitake et al. 2022). Here we identify CAN1 as a novel

plant-specific regulator of autophagosome size during local root tip Pi-sensing. CAN1

associates with the ER through interaction with VAP27-1 and the OST complex subunit

DGL1, which is essential for ER protein quality control (Lerouxel et al. 2005) (Fig. 6).

Following ER stress perception, CAN1 associates with Pi-responsive ATG8 isoforms (B and

C) preferentially via its C-terminal AIM motif (Fig. 1, 4). Loss of CAN1 leads to significant

smaller autophagosomes and enhanced ER stress resistance (Fig. 5, 6) positioning CAN1

function in Pi-dependent ER stress signaling to the regulation of autophagosome biogenesis

and maturation.

Phagophore expansion occurs at ER-associated initiation sites where lipid flux, membrane

moulding, and cargo capture are coordinated (Gross et al. 2025; Liu et al. 2020; Liu and

Bassham 2012; Nascimbeni et al. 2017; Su et al. 2020; Zhuang et al. 2013). The ER–PM

tethering protein VAP27-1 is a key organizer of ER contact sites and supports mitophagy by

promoting ER-mitochondria connections necessary for phagophore expansion around

dysfunctional mitochondria (Li et al. 2022; Wang et al. 2019). VAP27-1 and its yeast

homologues further support ER-stress adaptation through interactions with soluble ER-phagy

receptors (Nthiga et al. 2020; Zhao et al. 2020). Although loss of VAP27-1 or DGL1 severely

compromises plant development due to their essential roles in membrane tethering and

N-glycosylation (Lerouxel et al. 2005; Wang et al. 2016), loss of CAN1 results instead in

enhanced ER stress resistance, indicating a more specialized function. The interaction of

CAN1 with VAP27-1 and DGL1 (Fig. 6) suggests that CAN1 contributes to stabilizing

phagophore–ER contacts, a process central to the controlled expansion of the autophagosome

membrane. We propose CAN1 modulates phagophore elongation by reinforcing ER–

membrane interfaces via VAP27-1 and acts together with Pi-responsive ATG8B/C isoforms

to establish a root cell-specific expansion regime for controlling autophagosome size. Such a

CAN1–VAP27-1–ATG8B/C complex may further provide the spatial framework necessary

for PI3K complex recruitment, thereby coordinating phagophore growth with cargo

encapsulation.

In the physiological context of growing root tips, the degradation of stress-induced cargo must

be balanced with the preservation of resources, especially under limited Pi availability. The

smaller autophagosomes observed in *can1* likely favour rapid turnover of small or diffusible

cargos during early ER stress perception via the LPR1-PDR2 module, conferring enhanced

resistance to low-Pi stress. Conversely, CAN1-dependent enlargement of autophagosomes in

wildtype may facilitate more efficient sequestration of larger, bulky cargo, albeit at reduced

throughput. This size-throughput trade-off aligns with the gradual response of Pi–Fe-linked

ER stress induction, where rapid turnover may facilitate cell adaptation and sustain plant

survival (Naumann et al. 2022; Naumann et al. 2019). CAN1 probably functions separate of

established regulators of autophagosome maturation and fusion, such as SH3P2 and FREE1,

whose loss is lethal, as well as autophagy inhibitors including COST1 and Rubicon-related

factors (Bao et al., 2020; Gao et al., 2015; Matsunaga et al., 2009; Walczak & Martens, 2013;

Zhuang et al., 2013). This reveals CAN1 function at a previously unrecognized regulatory

layer dedicated specifically to autophagosome size determination.

We propose that in roots CAN1 relocates to ER contact sites upon low Pi stress, engaging

ATG8B/C and VAP27-1 to stabilize PI3P complexes and regulate phagophore elongation.

Following autophagosome closure, NBR1 may recruit CAN1 for turnover, positioning CAN1

as a new local regulator of autophagosome expansion (Fig. 6g). Our findings unravelled a

complex regulatory network opening new avenues to dissect cargo selectivity and stress-

responsive remodelling of endomembrane systems for plant environmental adaptation.

METHODS

Plant Materials

All *Arabidopsis thaliana* lines used in this study originate from accession Columbia (Col-0). Mutant lines *pdr2-1 (Ticconi et al. 2009)*, *lpr1 lpr2* (Svistoonoff et al. 2007), *ire1a ire 1b* (Naumann et al. 2019), *atg5-4 (Suzuki et al. 2005)*, *atg2-1* (Wang et al. 2011), *atg7-3* (Lai et al. 2011), *atg9-4* (Shin et al. 2014), *Δatg8 (Del Chiaro et al. 2025)* and transgenic lines *p35S::GFP-ATG8A* (Thompson et al. 2005), *pdr2-1 x p35S::GFP-ATG8A* (Naumann et al. 2019), *35Spro::SpER-GFP-HDEL, pdr2-1 x 35Spro::SpER-GFP-HDEL (Ticconi et al. 2009), pUBQ10::mCherry-ATG8E, atg5-1 x pUBQ10::mCherry-ATG8E* (Stephani et al. 2020), *Δatg8* x *pUBQ10::ATG8A, Δatg8* x *pUBQ10::ATG8H* (Del Chiaro et al. 2025) were previously described. T-DNA insertion line SALK_092190 was obtained from the European Arabidopsis Stock Center (NASC). The *can1-c1* and *can1-c3* knockout lines were produced by CRISPR-Cas9 technology (Stuttmann et al. 2021). Guide RNAs for target sequences (AAGCGTGAAGGCAAACAACAC and AACCCGGAAAGCTTCAAACC) were designed with CHOPCHOP (Labun et al. 2019) and cloned into pDGE332_M1 and pDGE334_M2E constructs (Ordon et al. 2017; Stuttmann et al. 2021) and assembled in pDGE347 for transformation. Double mutants were generated by genetic crossing and genotyped by PCR. Primers used for cloning and genotyping are listed in Supplementary table 1 and all vectors are listed in Supplementary table 2.

Molecular Cloning

The *AtDGL1* coding sequence (CDS) was synthesized by Thermo Fisher Scientific with three point mutations introduced to eliminate internal BpiI recognition sites. The CDS was flanked by BpiI sites to enable Golden Gate cloning and subcloned into the MoClo vector pAGM1287, generating the level 0 *AtDGL1* module pNAU7. The *Arabidopsis thaliana* Ubiquitin 10 (Ubi10) promoter was amplified by PCR from plasmid pRedRoot (Limpens et al. 2004) and cloned as MoClo level 0 promoter modules pAGM11323, flanked by BpiI sites

GGAG and AATG for use as a standard promoter module, and pAGM11335, flanked by BpiI

sites GGAG and CCAT for assembly of transcriptional units containing N-terminal fusions.

An internal BpiI site within the promoter sequence was removed during cloning by

introducing a nucleotide substitution via the PCR primers. For N-terminal GFP fusion

constructs, modules pAGM11335 (pUbi10), pICSL30006 (N-terminal GFP tag, (Engler et al.

2014) pNAU7 (*AtDGL1*), and pAGM16841 (tNOS, (Gantner et al. 2018) were assembled in a

single BsaI Golden Gate reaction into pICH47742 (Weber et al. 2011), yielding

pUbi10::GFP-AtDGL1::tNOS.

For C-terminal GFP fusion constructs, modules pAGM11323 (pUbi10), pICSL50004 (C-

terminal GFP tag, (Engler et al. 2014) pNAU7 (*AtDGL1*), and pICH41421 (tNOS,) (Engler et

al. 2014) were assembled in a single BsaI Golden Gate reaction into pICH47742, yielding

pUbi10::AtDGL1-GFP::tNOS.

All other constructs were cloned via Gateway Technology.

Plant Growth and Chemical Treatments

Seeds were surface-sterilized and germinated on 1% (w/v) Phyto-Agar (Duchefa) containing

2.5 mM KH_2PO_4, pH 5.6 (high Pi or +Pi medium) or no Pi supplement (low Pi or –Pi

medium) as reported (Naumann et al. 2022). Agar media deficient in other nutrients (C, N, or

Fe) or supplemented with ER stress-agents were prepared as described (Naumann et al. 2022;

Naumann et al. 2019). The agar was routinely purified by repeated washing in deionized

water and subsequent dialysis using DOWEX G-55 anion exchanger (Ticconi et al. 2009).

For carbon starvation experiments on soil, 6 weeks-old plants grown under short-day

photoperiod were transferred to darkness for 6 days followed by 7 days of recovery in short-

day condition (Naumann et al. 2019; Qi et al. 2023; Thompson et al. 2005). Nutrient

limitation in aseptic conditions was performed as follows: For carbon starvation, seedlings

were transferred to carbon (sucrose) deficient media 5 days after germination and kept in the

dark for 12 days followed by 7 days of recovery on +Pi media. For nitrogen starvation, 5

days-old seedlings (after germination on +Pi) were transferred to nitrogen deficient media for

7 days.

Chemical Treatments

Tunicamycin (TM) and torin treatment were performed by transferring 5-day-old seedlings

(germinated on +Pi) to liquid media supplemented with 5 µg/mL TM, 1 µM torin or DMSO

(solvent) for 5-7 h. Concanamycin A (ConA) treatment was added to a final concentration of

1µM for the indicated time points (3-5 h).

Histochemical staining

GUS (β-glucuronidase) staining was conducted as described (Naumann et al. 2022).

Seedlings were incubated in GUS-staining solution (50 mM Na-phosphate, pH7.2; 0.5 mM

$K_3Fe(CN)_6$; 0.5 mM $K_4Fe(CN)_6$; 2mM X-Gluc; 10mM EDTA; 0,1% TritonX) at 37°C and

subsequently cleared using chloral hydrate solution (8:2:1 chloral hydrate:glycerol: ddH_2O).

Iron detection, based on Perls staining coupled to diaminobenzidine (DAB) intensification

(Perls/DAB staining) was performed as previously described (Muller et al. 2015) with minor

changes to the protocol. Plants were incubated for 10 min in 2% (v/v) HCl, 4% (w/v) K-

ferrocyanide (Perls staining), or K-ferricyanide (Turnbull staining). For DAB intensification,

plants were washed twice (ddH_2O) and incubated (15 min) in methanol containing 10 mM

Na-azide and 0.3% (v/v) H_2O_2. After washing with 100 mM Na-phosphate buffer (pH 7.4),

plants were incubated for 3 min in the same buffer containing 0.025% (w/v) DAB (Sigma-

Aldrich) and 0.005% (v/v) H_2O_2. The reaction was stopped by washing with 100 mM Na-

phosphate buffer (pH 7.4) and optically clearing with chloral hydrate, 1 g/ml 15% (v/v)

glycerol.

Confocal Microscopy

Samples were analyzed using a multizoom stereomicroscope (Nikon AZ100) for overview

images and a Zeiss AxioImager bright field microscope for detail images. Confocal

microscopy was done on a Zeiss LSM900. Imaging was performed with a ×20 air, ×40 water

or x63 water immersion objective using sequential acquisition. For each experiment, 7-10

individual samples were observed in three independent approaches. Images were processed

using ImageJ (Schindelin et al. 2012) software. Total puncta numbers were counted using the

MTB Cell Counter (Franke et al. 2015). Colocalization and Pearsons coefficient were

calculated using the BIOP FiJi Plugin.

Callose was stained for 1 h with 0.1% (w/v) aniline blue (AppliChem) in 100 mM Na-

phosphate buffer (pH 7.2) and carefully washed twice. Fluorescence was visualized using a

Zeiss LSM 900 confocal laser-scanning microscope (excitation 405 nm, emission 498 nm) in

100 mM Na-phosphate buffer (pH 7.2) (Muller et al. 2015).

Autophagosome size determination

Five-day old seedlings expressing *35S::GFP-ATG8A* in WT or *can1-c1* were transferred to

+Pi or -Pi for one day followed by liquid treatments to induce ER-stress by TM (5 μg/ml) or

torin (1 μM) treatments for 6 h. Primary root transition zones were imaged and Arivis Vision

4D 3.6.2 (arivis AG) was used for analysis. A custom pipeline to measure autophagosome

volume was set up using the Blob Finder with the following parameters: average size 2 μm,

probability threshold 55%, split sensitivity 50%. Sphericity was set to > 0.65 and a segment

feature filter was applied to ≥ 0.6 $μm^2$.

Transient expression in *N. benthamiana*

Leaves of 6-week-old tobacco (*Nicotiana benthamiana*) plants were co-infiltrated with *A.*

tumefaciens harboring the respective plasmid. After 3 d, leaf discs were analyzed by confocal

laser scanning microscopy (GFP: excitation 488 nm, emission 493-537 nm; mCherry:

excitation 555 nm, emission 582-631 nm; chlorophyll autofluorescence: excitation 633 nm,

emission, 669-722 nm).

Yeast-two-hybrid assay

To perform direct Y2H test, the full length CAN1 as well as CAN1[F183A, V186A,] CAN1[W202A,]

^{V205A} and CAN1$^{F183A, V186A, W202A, V205A}$ (CAN1$^{\Delta AIM4}$, CAN1$^{\Delta AIM5}$ and CAN1$^{\Delta AIM4,5}$

respectively), were cloned into the pDEST32 (Invitrogen) vector using gateway technology,

to create protein fusions with the Gal4 DNA binding domain. All nine Arabidopsis ATG8

isoforms were cloned into the prey pDEST22 vector to obtain Gal4 activation fusions.

The bait and prey vectors to be tested for interaction of the fusion proteins were co-

transformed into *S. cerevisiae* strain PJ69-4a using the standard LiOAc method (Gietz and

Schiestl 2007). The yeast was spotted on plates with vector selective media DDO (double

drop out media: SD/-Leu/-Trp) and were grown for 4-5 days at 30°C. Two independent

colonies of each transformation were inoculated in 1x TE buffer and incubated at 30°C at 180

rpm overnight. On the next day, 10 µL of the colonies were spotted on DDO, interaction

selective media (triple drop out media: SD/-Leu/-Trp/-His) and triple drop out media

containing 2 mM 3-aminotriazole (triple drop out media: SD/-Leu/-Trp/-His/+3-

aminotriazole). Pictures were taken 6 days after spotting. SD plates were prepared with 1.6%

(w/v) Bactoagar (AppliChem) and the SD media was obtained from TakaraBio.

Affinity-based Immunoprecipitation

Proteins were extracted from plant material with RIPA-buffer (50 mM Tris-Cl, pH 7.6, 150

mM NaCl, 20 mM NaF, 10 mM Na$_4$P$_2$O$_7$, 1 mM EDTA, 0.5 mM EGTA, 1% Nonidet-P-40,

0.5% Na-deoxycholate, 1 mM PMSF, 1 mM DTT) for 30 min at 4°C. Post centrifugation the

cleared lysate was diluted 1:1 with detergent-free RIPA and incubated with equilibrated GFP-

trap or RFP-trap Agarose beads (ChromoTek) for 2h at 8°C on a multi-rotator. After

centrifugation (2500xg, 4°C) and washing 3-times (detergent-free RIPA), proteins were eluted

at 68°C for 15 min in 2x SDS sample buffer. The supernatant was transferred to a new tube

and boiled at 95°C for 5min.

Immunoblotting

Seedlings were germinated on +Pi medium for 5 d prior to transfer to Pi-sufficient (+Pi) or

deficient (-Pi) media. Whole roots were harvested 1dpt after treatments. For basal protein

quantification whole seedlings were directly harvested 5dpg. Proteins were extracted using

RIPA-buffer (50 mM Tris-Cl, pH 7.6, 150 mM NaCl, 20 mM NaF, 10 mM $Na_4P_2O_7$, 1 mM

EDTA, 0.5 mM EGTA, 1% Nonidet-P-40, 0.5% Na-deoxycholate, 1 mM PMSF, 1 mM DTT)

for 30 min at 4°C. Protein concentrations were measured using the Pierce™ 660 nm Protein-

Assay-Kit (Thermo Fisher, 22662). Equal amounts of total proteins were loaded on 10%

SDS-gels containing TCE (2,2,2-Trichlorethanol, Sigma-Aldrich, T54801) as loading control

(Ladner et al. 2004). All gels were simultaneously immunoblotted. All antibodies used in this

study are listed in Supplementary table 3.

Quantitative proteomics

Primary root tips were excised 24 h after transfer to the indicated media and collected into

liquid nitrogen. Tissue lysis, sample preparation, protein labeling, Tandem-Mass-Tag (TMT)

spectrometry, MS/MS data analysis, and TMT-quantifications were performed as recently

described (Naumann et al. 2022; Rodriguez et al. 2020; Stephani et al. 2020).

NanoLC-MS/MS Analysis: The nano HPLC system (Vanquish *Neo* UHPLC-System, Thermo

Scientific) was coupled to an Orbitrap Astral mass spectrometer, equipped with a Nanospray

Flex ion source (all parts Thermo Scientific). Peptides were loaded onto a trap column

(PepMap Acclaim C18, 5 mm × 300 μm ID, 5 μm particle size, 100 Å pore size, Thermo

Scientific) at a flow rate of 25 μl/min using 0.1% TFA as mobile phase.

After 5 minutes, the trap column was switched in line with the analytical column (Aurora

Ultimate C18 25cm × 75 μm ID, 1.7 μm particles, 120 Å, with integrated emitter, Ionopticks

operated at 50°C). Peptides were eluted using a flow rate of 300 nl/min, starting with the

mobile phases 98% A (0.1% formic acid in water) and 2% B (80% acetonitrile, 0.1% formic

acid) and linearly increasing to 35% B over the next 60 minutes, followed by an increase to

95% B in 1.7 min, a 4-min hold at 95% B, and re-equilibration with 2% B for three column

volumes (equilibration factor of 3.0).

The Orbitrap Astral was operated in data-dependent mode, performing a full scan in the

Orbitrap every 0.7 s (m/z range 375–1500 m/z; resolution 240,000; standard AGC target,

maximum injection time 10 ms, minimum intensity 5,000). MS/MS spectra were acquired in

the Astral analyser by isolating 1.6 Da windows and fragmenting precursor ions with

normalized HCD collision energy of 30% with a maximum injection time of 5 or 10 ms

correspondingly or until a standard AGC target was reached. Fragment ions ranging from

150–2000 m/z were acquired. Precursor ions selected for fragmentation (include charge state

2-6) were put on a dynamic exclusion list for 30 s.

qRT PCR

Abundance of mRNAs was quantified using fluorescence-based real-time RT-PCR and gene-

specific amplimers listed in Supplementary table 1. Relative mRNA abundances were

normalized relative to *UBC9* transcript abundance. Samples from three independent

experiments were measured.

Phylogenetic Analysis

CAN1 sequence alignment and phylogenetic analysis were done using BLAST search of

curated sequences in the NCBI database and phytozome 13. Obtained sequences were filtered

for fragments. All phylogenetic trees were calculated by sequence alignment using MAFFT

7108 with default settings and created at the CIPRES web-portal with RAxML 8.2.10109 for

maximum likelihood analyses using the JTT PAM matrix for amino acid substitutions in

RAxML. All related sequences used for phylogenetic analyses are provided (Supplementary

Data 2).

Quantification and Statistical Analysis

All claims of statistical significance can be found in the corresponding figure legends.

Statistical tests were carried out using GraphPad Prism 10 for Windows.

Acknowledgements We thank P.J. Hussey and P. Wang for seeds; M. Ried and S. Abel for scientific discussions and critical reading of the manuscript. This work was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) projects 492493685 and 400681449/GRK 2498.; an EMBO Short-Term Fellowship to C.N. and institutional core funding (Leibniz Association) from the Federal Republic of Germany and the State of Saxony-Anhalt. Work in the K.M. group was financially supported by the EPIC-XS (project number 823839), the Horizon 2020 Program of the European Union, and the ERA-CAPS I 3686 project of the Austrian Science Fund.

Author contributions D.G. and C.N. designed research; D.G., F.S., K.M., J.A., F.K., S.M., Y.TN., N.G. and S.H. conducted experiments and analyzed data; K.M. and R.I. performed proteomic analysis; Y.D. and C.N. supervised the project; D.G. drafted the manuscript; C.N. wrote the article.

Competing interests The authors declare no competing interests.

LITERATURE CITED

Abel S (2017) Phosphate scouting by root tips. Curr Opin Plant Biol 39:168-177.
        doi:10.1016/j.pbi.2017.04.016

Abel S, Naumann C (2024) Evolution of phosphate scouting in the terrestrial biosphere.
        Philos Trans R Soc Lond B Biol Sci 379 (1914):20230355.
        doi:10.1098/rstb.2023.0355

Balzergue C, Dartevelle T, Godon C, Laugier E, Meisrimler C, Teulon JM, Creff A, Bissler
        M, Brouchoud C, Hagege A, Muller J, Chiarenza S, Javot H, Becuwe-Linka N, David
        P, Peret B, Delannoy E, Thibaud MC, Armengaud J, Abel S, Pellequer JL, Nussaume
        L, Desnos T (2017) Low phosphate activates STOP1-ALMT1 to rapidly inhibit root
        cell elongation. Nat Commun 8:15300. doi:10.1038/ncomms15300

Cadena-Ramos AI, De-la-Pena C (2024) Picky eaters: selective autophagy in plant cells. Plant
        J 117 (2):364-384. doi:10.1111/tpj.16508

Clua J, Montpetit J, Jimenez-Sandoval P, Naumann C, Santiago J, Poirier Y (2024) A
        CYBDOM protein impacts iron homeostasis and primary root growth under phosphate
        deficiency in Arabidopsis. Nat Commun 15 (1):423. doi:10.1038/s41467-023-43911-x

Crombez H, Motte H, Beeckman T (2019) Tackling Plant Phosphate Starvation by the Roots.
        Dev Cell 48 (5):599-615. doi:10.1016/j.devcel.2019.01.002

Del Chiaro A, Grujic N, Zhao J, Papareddy RK, Gao P, Ma J, Lofke C, Bhattacharya A,
        Gruetzner R, Bourguet P, Berger F, Kang BH, Marillonnet S, Dagdas Y (2025)
        Nonuple atg8 mutant provides genetic evidence for functional specialization of ATG8
        isoforms in Arabidopsis thaliana. J Cell Sci. doi:10.1242/jcs.263803

Dong K, Deng G, Liu Y, Chen KE, Wei H, Huang X, Huang W, Zheng P, Ueda T, Vierstra
        RD, Huang X, Li F (2025) Comprehensive genetic analyses of Arabidopsis
        autophagy-related 8 family reveal redundant regulatory roles during autophagy. New
        Phytol. doi:10.1111/nph.70418

Engler C, Youles M, Gruetzner R, Ehnert TM, Werner S, Jones JD, Patron NJ, Marillonnet S
        (2014) A golden gate modular cloning toolbox for plants. ACS Synth Biol 3 (11):839-
        843. doi:10.1021/sb4001504

Feng Q, De Rycke R, Dagdas Y, Nowack MK (2022) Autophagy promotes programmed cell
        death and corpse clearance in specific cell types of the Arabidopsis root cap. Curr Biol
        32 (20):4548. doi:10.1016/j.cub.2022.10.006

Feng Z, Zhao Y, Li T, Nie W, Yang X, Wang X, Wu J, Liao J, Zou Y (2020) CATP-8/P5A
        ATPase Regulates ER Processing of the DMA-1 Receptor for Dendritic Branching.
        Cell Rep 32 (10):108101. doi:10.1016/j.celrep.2020.108101

Gantner J, Ordon J, Ilse T, Kretschmer C, Gruetzner R, Lofke C, Dagdas Y, Burstenbinder K,
        Marillonnet S, Stuttmann J (2018) Peripheral infrastructure vectors and an extended
        set of plant parts for the Modular Cloning system. PLoS One 13 (5):e0197185.
        doi:10.1371/journal.pone.0197185

Giehl RF, von Wiren N (2014) Root nutrient foraging. Plant Physiol 166 (2):509-517.
        doi:10.1104/pp.114.245225

Gietz RD, Schiestl RH (2007) High-efficiency yeast transformation using the LiAc/SS carrier
        DNA/PEG method. Nat Protoc 2 (1):31-34. doi:10.1038/nprot.2007.13

Gong F, Yao Z, Liu Y, Sun M, Peng X (2021) H(2)O(2) response gene 1/2 are novel sensors
        or responders of H(2)O(2) and involve in maintaining embryonic root meristem
        activity in Arabidopsis thaliana. Plant Sci 310:110981.
        doi:10.1016/j.plantsci.2021.110981

Gross AS, Raffeiner M, Zeng Y, Ustun S, Dagdas Y (2025) Autophagy in Plant Health and
        Disease. Annu Rev Plant Biol 76 (1):197-227. doi:10.1146/annurev-arplant-060324-
        094912

Gutierrez-Alanis D, Yong-Villalobos L, Jimenez-Sandoval P, Alatorre-Cobos F, Oropeza-
Aburto A, Mora-Macias J, Sanchez-Rodriguez F, Cruz-Ramirez A, Herrera-Estrella L
(2017) Phosphate Starvation-Dependent Iron Mobilization Induces CLE14 Expression
to Trigger Root Meristem Differentiation through CLV2/PEPR2 Signaling. Dev Cell
41 (5):555-570 e553. doi:10.1016/j.devcel.2017.05.009
Jakobsen MK, Poulsen LR, Schulz A, Fleurat-Lessard P, Moller A, Husted S, Schiott M,
Amtmann A, Palmgren MG (2005) Pollen development and fertilization in
Arabidopsis is dependent on the MALE GAMETOGENESIS IMPAIRED ANTHERS
gene encoding a type V P-type ATPase. Genes Dev 19 (22):2757-2769.
doi:10.1101/gad.357305
Ji J, Cui MK, Zou R, Wu MZ, Ge MX, Li J, Zhang ZR (2024) An ATP13A1-assisted
topogenesis pathway for folding multi-spanning membrane proteins. Mol Cell 84
(10):1917-1931 e1915. doi:10.1016/j.molcel.2024.04.010
Johansen T, Lamark T (2020) Selective Autophagy: ATG8 Family Proteins, LIR Motifs and
Cargo Receptors. J Mol Biol 432 (1):80-103. doi:10.1016/j.jmb.2019.07.016
Jung H, Lee HN, Marshall RS, Lomax AW, Yoon MJ, Kim J, Kim JH, Vierstra RD, Chung T
(2020) Arabidopsis cargo receptor NBR1 mediates selective autophagy of defective
proteins. J Exp Bot 71 (1):73-89. doi:10.1093/jxb/erz404
Kellermeier F, Armengaud P, Seditas TJ, Danku J, Salt DE, Amtmann A (2014) Analysis of
the Root System Architecture of Arabidopsis Provides a Quantitative Readout of
Crosstalk between Nutritional Signals. Plant Cell 26 (4):1480-1496.
doi:10.1105/tpc.113.122101
L. Franke BS, J. L. Erickson, D. Rödel, D. Schröter, B. Möller, and M. H. Schattat (2015) The
'MTB Cell Counter' a versatile tool for the semi-automated quantification of sub-
cellular phenotypes in fluorescence microscopy images. A case study on plastids,
nuclei and peroxisomes. Journal of Endocytobiosis and Cell Research 26:31-42
Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E (2019)
CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing.
Nucleic Acids Res 47 (W1):W171-W174. doi:10.1093/nar/gkz365
Ladner CL, Yang J, Turner RJ, Edwards RA (2004) Visible fluorescent detection of proteins
in polyacrylamide gels without staining. Anal Biochem 326 (1):13-20.
doi:10.1016/j.ab.2003.10.047
Lai Z, Wang F, Zheng Z, Fan B, Chen Z (2011) A critical role of autophagy in plant
resistance to necrotrophic fungal pathogens. Plant J 66 (6):953-968.
doi:10.1111/j.1365-313X.2011.04553.x
Lambers H, Martinoia E, Renton M (2015) Plant adaptations to severely phosphorus-
impoverished soils. Curr Opin Plant Biol 25:23-31. doi:10.1016/j.pbi.2015.04.002
Lerouxel O, Mouille G, Andeme-Onzighi C, Bruyant MP, Seveno M, Loutelier-Bourhis C,
Driouich A, Hofte H, Lerouge P (2005) Mutants in DEFECTIVE
GLYCOSYLATION, an Arabidopsis homolog of an oligosaccharyltransferase
complex subunit, show protein underglycosylation and defects in cell differentiation
and growth. Plant J 42 (4):455-468. doi:10.1111/j.1365-313X.2005.02392.x
Li C, Duckney P, Zhang T, Fu Y, Li X, Kroon J, De Jaeger G, Cheng Y, Hussey PJ, Wang P
(2022) TraB family proteins are components of ER-mitochondrial contact sites and
regulate ER-mitochondrial interactions and mitophagy. Nat Commun 13 (1):5658.
doi:10.1038/s41467-022-33402-w
Li P, Bagenholm V, Hagglund P, Lindkvist-Petersson K, Wang K, Gourdon P (2024) The
structure and function of P5A-ATPases. Nat Commun 15 (1):9605.
doi:10.1038/s41467-024-53757-6

Li T, Yang X, Feng Z, Nie W, Fang Z, Zou Y (2021) P5A ATPase controls ER translocation
of Wnt in neuronal migration. Cell Rep 37 (4):109901.
doi:10.1016/j.celrep.2021.109901
Limpens E, Ramos J, Franken C, Raz V, Compaan B, Franssen H, Bisseling T, Geurts R
(2004) RNA interference in Agrobacterium rhizogenes-transformed roots of
Arabidopsis and Medicago truncatula. J Exp Bot 55 (399):983-992.
doi:10.1093/jxb/erh122
Liu F, Hu W, Li F, Marshall RS, Zarza X, Munnik T, Vierstra RD (2020) AUTOPHAGY-
RELATED14 and Its Associated Phosphatidylinositol 3-Kinase Complex Promote
Autophagy in Arabidopsis. Plant Cell 32 (12):3939-3960. doi:10.1105/tpc.20.00285
Liu W, Liu Z, Mo Z, Guo S, Liu Y, Xie Q (2021) ATG8-Interacting Motif: Evolution and
Function in Selective Autophagy of Targeting Biological Processes. Front Plant Sci
12:783881. doi:10.3389/fpls.2021.783881
Liu Y, Bassham DC (2012) Autophagy: pathways for self-eating in plant cells. Annu Rev
Plant Biol 63:215-237. doi:10.1146/annurev-arplant-042811-105441
Man Y, Zhang Y, Chen L, Zhou J, Bu Y, Zhang X, Li X, Li Y, Jing Y, Lin J (2024) The
VAMP-associated protein VAP27-1 plays a crucial role in plant resistance to ER
stress by modulating ER-PM contact architecture in Arabidopsis. Plant Commun 5
(7):100929. doi:10.1016/j.xplc.2024.100929
Maniero RA, Picco C, Hartmann A, Engelberger F, Gradogna A, Scholz-Starke J, Melzer M,
Kunze G, Carpaneto A, von Wiren N, Giehl RFH (2024) Ferric reduction by a
CYBDOM protein counteracts increased iron availability in root meristems induced
by phosphorus deficiency. Nat Commun 15 (1):422. doi:10.1038/s41467-023-43912-
w
McKenna MJ, Adams BM, Chu V, Paulo JA, Shao S (2022) ATP13A1 prevents ERAD of
folding-competent mislocalized and misoriented proteins. Mol Cell 82 (22):4277-4289
e4210. doi:10.1016/j.molcel.2022.09.035
McKenna MJ, Sim SI, Ordureau A, Wei L, Harper JW, Shao S, Park E (2020) The
endoplasmic reticulum P5A-ATPase is a transmembrane helix dislocase. Science 369
(6511). doi:10.1126/science.abc5809
Muller J, Toev T, Heisters M, Teller J, Moore KL, Hause G, Dinesh DC, Burstenbinder K,
Abel S (2015) Iron-dependent callose deposition adjusts root meristem maintenance to
phosphate availability. Dev Cell 33 (2):216-230. doi:10.1016/j.devcel.2015.02.007
Nakatogawa H (2020) Mechanisms governing autophagosome biogenesis. Nat Rev Mol Cell
Biol 21 (8):439-458. doi:10.1038/s41580-020-0241-0
Nascimbeni AC, Codogno P, Morel E (2017) Phosphatidylinositol-3-phosphate in the
regulation of autophagy membrane dynamics. FEBS J 284 (9):1267-1278.
doi:10.1111/febs.13987
Naumann C, Heisters M, Brandt W, Janitza P, Alfs C, Tang N, Toto Nienguesso A, Ziegler J,
Imre R, Mechtler K, Dagdas Y, Hoehenwarter W, Sawers G, Quint M, Abel S (2022)
Bacterial-type ferroxidase tunes iron-dependent phosphate sensing during Arabidopsis
root development. Curr Biol 32 (10):2189-2205 e2186. doi:10.1016/j.cub.2022.04.005
Naumann C, Muller J, Sakhonwasee S, Wieghaus A, Hause G, Heisters M, Burstenbinder K,
Abel S (2019) The Local Phosphate Deficiency Response Activates Endoplasmic
Reticulum Stress-Dependent Autophagy. Plant Physiol 179 (2):460-476.
doi:10.1104/pp.18.01379
Nthiga TM, Kumar Shrestha B, Sjottem E, Bruun JA, Bowitz Larsen K, Bhujabal Z, Lamark
T, Johansen T (2020) CALCOCO1 acts with VAMP-associated proteins to mediate
ER-phagy. EMBO J 39 (15):e103649. doi:10.15252/embj.2019103649

Ordon J, Gantner J, Kemna J, Schwalgun L, Reschke M, Streubel J, Boch J, Stuttmann J (2017) Generation of chromosomal deletions in dicotyledonous plants employing a user-friendly genome editing toolkit. Plant J 89 (1):155-168. doi:10.1111/tpj.13319

Petersen M, Avin-Wittenberg T, Bassham DC, Dagdas Y, Fan C, Fernie AR, Jiang L, Mishra D, Otegui MS, Rodriguez E, Hofius D (2024) Autophagy in plants. Autophagy Rep 3 (1):2395731. doi:10.1080/27694127.2024.2395731

Picchianti L, Sanchez de Medina Hernandez V, Zhan N, Irwin NA, Groh R, Stephani M, Hornegger H, Beveridge R, Sawa-Makarska J, Lendl T, Grujic N, Naumann C, Martens S, Richards TA, Clausen T, Ramundo S, Karagoz GE, Dagdas Y (2023) Shuffled ATG8 interacting motifs form an ancestral bridge between UFMylation and autophagy. EMBO J 42 (10):e112053. doi:10.15252/embj.2022112053

Qi H, Wang Y, Bao Y, Bassham DC, Chen L, Chen QF, Hou S, Hwang I, Huang L, Lai Z, Li F, Liu Y, Qiu R, Wang H, Wang P, Xie Q, Zeng Y, Zhuang X, Gao C, Jiang L, Xiao S (2023) Studying plant autophagy: challenges and recommended methodologies. Adv Biotechnol (Singap) 1 (4):2. doi:10.1007/s44307-023-00002-8

Qin Q, Zhao T, Zou W, Shen K, Wang X (2020) An Endoplasmic Reticulum ATPase Safeguards Endoplasmic Reticulum Identity by Removing Ectopically Localized Mitochondrial Proteins. Cell Rep 33 (6):108363. doi:10.1016/j.celrep.2020.108363

Rodriguez E, Chevalier J, Olsen J, Ansbol J, Kapousidou V, Zuo Z, Svenning S, Loefke C, Koemeda S, Drozdowskyj PS, Jez J, Durnberger G, Kuenzl F, Schutzbier M, Mechtler K, Ebstrup EN, Lolle S, Dagdas Y, Petersen M (2020) Autophagy mediates temporary reprogramming and dedifferentiation in plant somatic cells. EMBO J 39 (4):e103315. doi:10.15252/embj.2019103315

Rogov VV, Nezis IP, Tsapras P, Zhang H, Dagdas Y, Noda NN, Nakatogawa H, Wirth M, Mouilleron S, McEwan DG, Behrends C, Deretic V, Elazar Z, Tooze SA, Dikic I, Lamark T, Johansen T (2023) Atg8 family proteins, LIR/AIM motifs and other interaction modes. Autophagy Rep 2 (1). doi:10.1080/27694127.2023.2188523

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A (2012) Fiji: an open-source platform for biological-image analysis. Nat Methods 9 (7):676-682. doi:10.1038/nmeth.2019

Shen J, Yuan L, Zhang J, Li H, Bai Z, Chen X, Zhang W, Zhang F (2011) Phosphorus dynamics: from soil to plant. Plant Physiol 156 (3):997-1005. doi:10.1104/pp.111.175232

Shin KD, Lee HN, Chung T (2014) A revised assay for monitoring autophagic flux in Arabidopsis thaliana reveals involvement of AUTOPHAGY-RELATED9 in autophagy. Mol Cells 37 (5):399-405. doi:10.14348/molcells.2014.0042

Sim SI, Park E (2023) P5-ATPases: Structure, substrate specificities, and transport mechanisms. Curr Opin Struct Biol 79:102531. doi:10.1016/j.sbi.2023.102531

Stefano G, Renna L, Wormsbaecher C, Gamble J, Zienkiewicz K, Brandizzi F (2018) Plant Endocytosis Requires the ER Membrane-Anchored Proteins VAP27-1 and VAP27-3. Cell Rep 23 (8):2299-2307. doi:10.1016/j.celrep.2018.04.091

Stephani M, Dagdas Y (2020) Plant Selective Autophagy-Still an Uncharted Territory With a Lot of Hidden Gems. J Mol Biol 432 (1):63-79. doi:10.1016/j.jmb.2019.06.028

Stephani M, Picchianti L, Dagdas Y (2021) C53 is a cross-kingdom conserved reticulophagy receptor that bridges the gap betweenselective autophagy and ribosome stalling at the endoplasmic reticulum. Autophagy 17 (2):586-587. doi:10.1080/15548627.2020.1846304

Stephani M, Picchianti L, Gajic A, Beveridge R, Skarwan E, Sanchez de Medina Hernandez V, Mohseni A, Clavel M, Zeng Y, Naumann C, Matuszkiewicz M, Turco E, Loefke C, Li B, Durnberger G, Schutzbier M, Chen HT, Abdrakhmanov A, Savova A, Chia KS,

Djamei A, Schaffner I, Abel S, Jiang L, Mechtler K, Ikeda F, Martens S, Clausen T,
Dagdas Y (2020) A cross-kingdom conserved ER-phagy receptor maintains
endoplasmic reticulum homeostasis during stress. Elife 9. doi:10.7554/eLife.58396
Stuttmann J, Barthel K, Martin P, Ordon J, Erickson JL, Herr R, Ferik F, Kretschmer C,
Berner T, Keilwagen J, Marillonnet S, Bonas U (2021) Highly efficient multiplex
editing: one-shot generation of 8x Nicotiana benthamiana and 12x Arabidopsis
mutants. Plant J 106 (1):8-22. doi:10.1111/tpj.15197
Su T, Li X, Yang M, Shao Q, Zhao Y, Ma C, Wang P (2020) Autophagy: An Intracellular
Degradation Pathway Regulating Plant Survival and Stress Response. Front Plant Sci
11:164. doi:10.3389/fpls.2020.00164
Suzuki NN, Yoshimoto K, Fujioka Y, Ohsumi Y, Inagaki F (2005) The crystal structure of
plant ATG12 and its biological implication in autophagy. Autophagy 1 (2):119-126.
doi:10.4161/auto.1.2.1859
Svenning S, Lamark T, Krause K, Johansen T (2011) Plant NBR1 is a selective autophagy
substrate and a functional hybrid of the mammalian autophagic adapters NBR1 and
p62/SQSTM1. Autophagy 7 (9):993-1010. doi:10.4161/auto.7.9.16389
Svistoonoff S, Creff A, Reymond M, Sigoillot-Claude C, Ricaud L, Blanchet A, Nussaume L,
Desnos T (2007) Root tip contact with low-phosphate media reprograms plant root
architecture. Nat Genet 39 (6):792-796. doi:10.1038/ng2041
Thompson AR, Doelling JH, Suttangkakul A, Vierstra RD (2005) Autophagic nutrient
recycling in Arabidopsis directed by the ATG8 and ATG12 conjugation pathways.
Plant Physiol 138 (4):2097-2110. doi:10.1104/pp.105.060673
Ticconi CA, Delatorre CA, Abel S (2001) Attenuation of phosphate starvation responses by
phosphite in Arabidopsis. Plant Physiol 127 (3):963-972
Ticconi CA, Lucero RD, Sakhonwasee S, Adamson AW, Creff A, Nussaume L, Desnos T,
Abel S (2009) ER-resident proteins PDR2 and LPR1 mediate the developmental
response of root meristems to phosphate availability. Proc Natl Acad Sci U S A 106
(33):14174-14179. doi:10.1073/pnas.0901778106
Walczak M, Martens S (2013) Dissecting the role of the Atg12-Atg5-Atg16 complex during
autophagosome formation. Autophagy 9 (3):424-425. doi:10.4161/auto.22931
Wang P, Pleskot R, Zang J, Winkler J, Wang J, Yperman K, Zhang T, Wang K, Gong J, Guan
Y, Richardson C, Duckney P, Vandorpe M, Mylle E, Fiserova J, Van Damme D,
Hussey PJ (2019) Plant AtEH/Pan1 proteins drive autophagosome formation at ER-
PM contact sites with actin and endocytic machinery. Nat Commun 10 (1):5132.
doi:10.1038/s41467-019-12782-6
Wang P, Richardson C, Hawkins TJ, Sparkes I, Hawes C, Hussey PJ (2016) Plant VAP27
proteins: domain characterization, intracellular localization and role in plant
development. New Phytol 210 (4):1311-1326. doi:10.1111/nph.13857
Wang Y, Nishimura MT, Zhao T, Tang D (2011) ATG2, an autophagy-related protein,
negatively affects powdery mildew resistance and mildew-induced cell death in
Arabidopsis. Plant J 68 (1):74-87. doi:10.1111/j.1365-313X.2011.04669.x
Weber E, Engler C, Gruetzner R, Werner S, Marillonnet S (2011) A modular cloning system
for standardized assembly of multigene constructs. PLoS One 6 (2):e16765.
doi:10.1371/journal.pone.0016765
Wild P, McEwan DG, Dikic I (2014) The LC3 interactome at a glance. J Cell Sci 127 (Pt
1):3-9. doi:10.1242/jcs.140426
Wilfling F, Lee CW, Erdmann PS, Zheng Y, Sherpa D, Jentsch S, Pfander B, Schulman BA,
Baumeister W (2020) A Selective Autophagy Pathway for Phase-Separated Endocytic
Protein Deposits. Mol Cell 80 (5):764-778 e767. doi:10.1016/j.molcel.2020.10.030

Yang SY, Lin WY, Hsiao YM, Chiou TJ (2024) Milestones in understanding transport,
sensing, and signaling of the plant nutrient phosphorus. Plant Cell 36 (5):1504-1523.
doi:10.1093/plcell/koad326
Yoshitake Y, Nakamura S, Shinozaki D, Izumi M, Yoshimoto K, Ohta H, Shimojima M
(2021) RCB-mediated chlorophagy caused by oversupply of nitrogen suppresses
phosphate-starvation stress in plants. Plant Physiol 185 (2):318-330.
doi:10.1093/plphys/kiaa030
Yoshitake Y, Shinozaki D, Yoshimoto K (2022) Autophagy triggered by iron-mediated ER
stress is an important stress response to the early phase of Pi starvation in plants. Plant
J 110 (5):1370-1381. doi:10.1111/tpj.15743
Zhao D, Zou CX, Liu XM, Jiang ZD, Yu ZQ, Suo F, Du TY, Dong MQ, He W, Du LL (2020)
A UPR-Induced Soluble ER-Phagy Receptor Acts with VAPs to Confer ER Stress
Resistance. Mol Cell 79 (6):963-977 e963. doi:10.1016/j.molcel.2020.07.019
Zhao J, Bui MT, Ma J, Kunzl F, Picchianti L, De La Concepcion JC, Chen Y, Petsangouraki
S, Mohseni A, Garcia-Leon M, Gomez MS, Giannini C, Gwennogan D, Kobylinska
R, Clavel M, Schellmann S, Jaillais Y, Friml J, Kang BH, Dagdas Y (2022) Plant
autophagosomes mature into amphisomes prior to their delivery to the central vacuole.
J Cell Biol 221 (12). doi:10.1083/jcb.202203139
Zheng X, Zhao C, Liu Y (2018) Examining Autophagy in Plant by Transmission Electron
Microscopy (TEM). Bio Protoc 8 (20):e3047. doi:10.21769/BioProtoc.3047
Zhuang X, Wang H, Lam SK, Gao C, Wang X, Cai Y, Jiang L (2013) A BAR-domain protein
SH3P2, which binds to phosphatidylinositol 3-phosphate and ATG8, regulates
autophagosome formation in Arabidopsis. Plant Cell 25 (11):4596-4615.
doi:10.1105/tpc.113.118307

Figure 1

Figure 1: ER stress-induced autophagy by local Pi-sensing specifically activates ATG8B and ATG8C decorated autophagosomes. a, GFP-ATG8B and C labeled autophagosomes in Pi-deprived root tips. GFP-ATG8B/C derived fluorescence in primary root tips of transgenic (*pATG8B/C::GFP-ATG8B/C*) wildtype and *pdr2* seedlings after germination on +Pi agar (5 d) and subsequent transfer to –Pi medium for 0-24 h or treatment with 5 µg/mL tunicamycin (TM) for 6 h. Shown are representative images from three independent experiments fused Z-stacks of the transition zone. Rectangle indicate inset selection. Scale bars, 50 µm. **b** and **c**, Quantification of dataset shown in (**a**). Quantification of GFP-ATG8 puncta in root cells per normalized area. Box plots show medians and interquartile ranges of detected autophagosomes; outliers (>1.53 interquartile range) are shown as black dots. Letters denote statistical differences in condition at $p < 0.05$ (Two-way ANOVA and Tukey's HSD post hoc test) and $n \geq 27$.

Figure 2

Fold Change		ATG8 Interaction		ID	name
WT	***pdr2***	**WT**	***pdr2***		
3.52	**3.67**	yes	yes	AT5G40690.1	CAN1
2.40	**2.33**	yes	yes	AT4G04620.2	ATG8B
1.91	**1.88**	yes	yes	AT1G62040.2	ATG8C
1.81	1.78	yes	yes	AT1G44770.1	unknown protein
1.72	1.76	yes	yes	AT4G24690.1	NBR1
1.33	1.35	yes	yes	AT3G49490.1	unknown protein

Figure 2: Autophagy modulates root tip proteome composition. a, Seedlings were germinated 5 days prior to transfer to the indicated treatments. 24 h post transfer primary root tips were excised and either directly subjected to extraction and quantification by mass-spectrometry (TMT-analysis) or used for co-immunoprecipitation followed by MS analysis. Venn diagram of three overlapping pairwise comparisons for root tip TMT-MS conducted in wildtype (yellow), *pdr2* (magenta) and *atg2* (cyan) plants: wildtype (-Pi) vs wildtype (+Pi) yellow cycle, *pdr2* (-Pi) vs. *pdr2* (+Pi) magenta cycle, *atg2* (-Pi) vs. *atg2* (+Pi) cyan cycle. Protein abundances with $0.6 > FC > 1.3$. Total numbers are depicted in brackets. **b**, Overlap between proteins detected in –Pi conditions enriched in *atg2* compared to wildtype (yellow) or *pdr2-1* (magenta) identified by quantitative MS, $FC > 1.3$. Intersections show proteins with increased abundance in *atg2* but not in wildtype or *pdr2-1*. **c,d** *In vitro* interactome of GFP-ATG8A. Root extracts of GFP-ATG8A in transgenic wildtype (WT, **c**) and *pdr2* (**d**) primary roots were incubated with GFP-Trap beads and interacting proteins were analyzed by MS.

Venn diagram of proteins co-eluting with GFP-ATG8A in +Pi (light green) and -Pi (dark green) according to the genotype. **e**, Summarized results of data obtained in c and d. Fold-Change (log2) of normalized protein abundances in *atg2* compared to wildtype (first column) and *pdr2* (second column) in -Pi conditions.

Figure 3

Figure 3: CAN1 is a plant specific protein. a, Maximum-likelihood midpoint-rooted tree (400 bootstrap replicates) of full-length CAN1 orthologues (see Supplementary Data 2). **b-d**, *CAN1* gene is predominantly expressed in root tissue and elevated in autophagy mutant plants. **b**, Left panel: RT-qPCR analysis of *CAN1* transcript abundance relative to *UBC9* in seedlings of wildtype, *atg2*, *atg5*, *atg7* and *atg9,* n = 4; right panel: *pCAN1::GUS* activity in 3 week old seedlings of wildtype and *atg5,* scale bar 1 cm. **c,d**, *pCAN1::GUS* activity in 7 days old wildtype (c) and *atg5* (d) seedlings (I), root elongation zone (II), lateral root emergence (III), root-shoot junction (IV), primary root tip (V), true leaf tip (VI) and flower (VII), I-VI scale bar 100 µm, VII scale bar 1 cm, n ≥ 8.

Figure 4: CAN1 interacts with ATG8 isoforms via select AIMs. a, endogenous ATG8 interacts with CAN1. *In vivo* immunoprecipitation of extracts of Arabidopsis seedlings expressing CAN1-mCherry after germination on + Pi media for 5 days. ER stress was applied by incubation in +Pi with 5 µg/mL TM and 1 µM ConA containing liquid media for 7h. Input and bound proteins were visualized by immunoblotting with anti-RFP and anti-ATG8 antibodies, loading control Coomassie blue staining (CBB). **b**, Colabfold prediction of CAN1-ATG8 interaction via AIM4 and AIM5. *In silico* prediction of CAN1 interaction with ATG8E

revealed interaction via AIM5 (stick representation; see also Figure S6) which locates in Arabidopsis ATG8E (surface representation–grey) AIM pockets, known as W-site (site 1 – yellow) and L-site (site 2 –slate blue) (top panel) interacting with W202 and V205 (top right panel, close view – red). *In silico* mutation of W202A and V205A (CAN1$^{\Delta\text{AIM5}}$) predicts ATG8E CAN1 interaction via AIM4 (middle panel, right close view –orange) with F183 and V186 interacting with site 1 and site 2 respectively. CAN1 depleted of AIM4 and AIM5 is not predicted to interact with ATG8E via the AIM pockets (lower row panels, CAN1 $^{\Delta\text{AIM4/5}}$ – green). **c**, CAN1 interacts AIM specific *in vivo*. Y2H assay between a GAL4 DNA binding domain (DBD, bait) fused variant of CAN1 with GAL4 activation domain (AD, prey) fused variants of all nine ATG8 isoforms A-I. GAL4-DBD and AD alone were included as negative controls (EV). The growth of yeast cultures on vector-selective (−Trp/−Leu) (top), interaction-selective (−Trp/−Leu/−His) (middle) and strong interaction-selective (−Trp/−Leu/−His + 3-aminotriazol) media (bottom) six days after spotting is shown. **d**, Subcellular localization of CAN1-RFP. Co-expression of CAN1-RFP with GFP-CAN1 (Upper row) or GFP-HDEL (lower row) in transient expression assays in *N. benthamiana*. The images are single optical sections of RFP (left column), GFP (center column) and merged signals (right column). Scale bars, 10 µm. **e,f** CAN1 interacts with ATG8 in an isoform and AIM specific manner. **e**, Localization of GFP-CAN1 variants (CAN1, CAN1$^{\Delta\text{AIM4}}$, CAN1$^{\Delta\text{AIM5}}$ CAN1$^{\Delta\text{AIM4/5}}$) transiently co-expressed with RFP-ATG8B in *N. benthamiana*. The images are single optical sections of GFP (first column), RFP (second column), chlorophyll A (third column, blue) and merged (including bright field) signals (last column). Scale bars, 30 µm, n = 3. **f**, *In vitro* co-immunoprecipitation assays of all nine RFP-ATG8x isoforms co-expressed with CAN1-GFP variants in *N. benthamiana* via RFP-Trap. Presence of RFP-ATG8A-I and GFP-CAN1 variants was visualized by immunoblotting with anti-RFP and anti-GFP antibodies, loading control TCE visualization, n = 5.

Figure 5

a

pCAN1::CAN1-GFP

b

c

mCherry-ATG8E x CAN1-GFP

d

e

f

Figure 5: CAN1 functions in Pi-dependent ER stress-induced autophagy. a, CAN1-GFP
derived fluorescence in the transition zone of transgenic (*pCAN1::CAN1-GFP*) wildtype and
pdr2 seedlings after germination on +Pi agar (5 d) and subsequent transfer to + or –Pi medium

for 24 h or treatment with TM in liquid + Pi media for 6 h. Shown are representative images from three independent experiments fused Z-stacks of the root tip. Scale bars, 20 μm. **b,** Quantification of CAN1-GFP puncta in root cells per area. Box plots show medians and interquartile ranges of detected autophagosomes; outliers (>1.53 interquartile range) are shown as black dots, n ≥ 34. Significant differences compared to wildtype are indicated with * when p value ≤ 0.05, ** when p value ≤ 0.01, *** when p value ≤ 0.001, and **** when p value ≤ 0.001. Unpaired t-test with Welch's correction. **c,** CAN1-GFP puncta colocalize with mCherry-ATG8E *in vivo*. Co-localization analyses of single plane confocal images obtained from transgenic wildtype (left panel) and *atg5* (right panel) Arabidopsis roots co-expressing mCherry-ATG8E (magenta) with CAN1-GFP (green). Seedlings were germinated 5 days prior transfer to the indicated treatment for 1 day (+/-Pi) or 5 h liquid media (+ Pi + TM; - Pi + ConA; + Pi + TM + ConA). Rectangles indicate Inset. Scale bars, 20 μm. **d,** Quantification of mCherry-ATG8E and CAN1-GFP colocalized puncta in the indicated treatments as described in a. n ≥ 5. **e-f,** loss of *CAN1* increases Arabidopsis ER stress tolerance. **e,** Top: 11 days old seedlings of wildtype, *pdr2*, *can1-c1* and *can1-c3* germinated on control or ER stress (100 ng/mL, TM) media. Bottom: Fresh weight of 5 pooled seedlings per sample, n = 17-18. **f,** Top: Germination of the indicated genotypes for eight days on Pi-sufficient (+Pi) or Pi-depleted (-Pi) conditions, scale bar = 1cm. Bottom: Primary root growth inhibition in percentage of the indicated genotypes, shown are summarized data of five independent biological experiments with each n ≥ 8. Significant differences compared to wildtype are indicated with * when p value ≤ 0.05, ** when p value ≤ 0.01, *** when p value ≤ 0.001, and **** when p value ≤ 0.001. Unpaired t-test with Welch's correction.

Figure 6

Figure 6: CAN1 interacts with NBR1 and VAP27-1 to modulate autophagosomal size. a, Enrichment of proteins interacting with ATG8A, *pUBQ10::GFP-CAN1* in wildtype and *pCAN1::CAN1-GFP* in *atg5* background in ER stress (+ TM + ConA) compared with a control expressing free GFP and represented by a volcano plot. Seedlings were germinated 10 days on liquid +Pi media prior to transfer to the indicated media. The horizontal grey line

indicates the threshold above which proteins are enriched (P value <0.03, quasi-likelihood negative binomial generalized log-linear model), and the vertical dashed line indicates the threshold where the proteins' $\log_2$FC is greater than 1. **b**, Western blot analysis from immunoprecipitation (IP) experiments using wildtype and *atg5* expressing *pCAN1::CAN1–GFP* or *pCAN1::GFP-GUS* to immunoprecipitate NBR1. Seedlings were germinated 10 days on liquid +Pi media prior to transfer to + or – Pi for 24 h or ER stress treatment with 5 µg/ml TM and 1 µM ConA for 6 h. Whole roots were harvested. The blots were probed with anti-NBR1 and anti-GFP antibodies to detect the presence of NBR1. **c-d**, CAN1 interacts with VAP27;1 in *N.benthamiana* and *A. thaliana* roots in ER stress. **c**, IP of *p35S::VAP27-1-RFP* or *p35S::mCherry-HDEL* co-expressed with *pUBQ10::GFP-CAN1* in *N.benthamiana* leaves. Input and bound proteins were visualized with anti-GFP and anti-RFP antibodies. **d**, IP of extracts of F1 Arabidopsis roots co-expressing *p35S::CAN1-mCherry* and *pVAP27-1::VAP27-1-GFP* after transfer for 24 h to Pi sufficient or deficient liquid media. ER stress was applied by incubation in +Pi with 5 µg/ml TM and 1 µM ConA containing liquid media for 6 h. Input and bound proteins were visualized by immunoblotting with anti-RFP and anti-GFP antibodies. **e**, NBR1 protein amount is not influenced by CAN1 in Pi-deprived roots. Endogenous NBR1 detection by anti-NBR1 antibody in extracts from wildtype and *can1-c1* roots germinated 5 days on +Pi media prior to transfer to +Pi, –Pi or –Pi containing 1 µM ConA liquid media for 24h. **f**, Loss of CAN1 affects autophagosome size in Pi-stressed roots. Seedlings of wildtype or *can1-c1* expressing *p35S::GFP-ATG8A* were germinated for 5 days on +Pi media prior to transfer for 24 h to +Pi or –Pi media or incubated with 5 µg/ml TM or 1µM torin for 6 h for autophagic induction. Size of GFP-ATG8A labelled puncta were measured via Arivis software and calculated volume was plotted accordingly. Scatter plot, median is shown as bold line, letters denote statistical differences at p < 0.05 (Two-way ANOVA and Tukey's HSD post hoc test) and n = 20. **g**, Model of CAN1 function in autophagy. Induction of ER stress by local root tip Pi-sensing initiates CAN1 association to the ER and interaction with VAP27-1 to enable autophagosome maturation and membrane expansion for bulky cargo engulfment (e.g. ER-fractions, organelles). Potential NBR1 recruitment of CAN1 to the mature autophagosome by an unknown signal initiates ER-detachment and vacuolar degradation.

Supplementary Figure S1: Root tip ATG8 isoform specific autophagosome decoration.
Seedlings were germinated 5 days prior transfer to the indicated treatment. Representative
confocal microscopic images showing autophagosome formation of all 9 ATG8 isoforms
(*pATG8x::GFP-ATG8x*) in the transition zone (row 1,3,5,7) or primary root tip (row 2,4,6,8)
challenged with local Pi deprivation (row 3 and 4), carbon starvation (row 5 and 6) for 24 h or

5 µg/mL tunicamycin (TM) treatment (row 7 and 8) for 6-8 h, rectangle indicate Inset selection, n ≥ 10.

Supplementary Figure S2

Supplementary Figure S2: Select ATG8 isoforms are involved in root tip Pi-dependent ER stress-mediated autophagy. a, Analysis of *ATG8A-I* relative transcript levels (normalized to UBC9 expression) in wildtype and *pdr2* root tips after transfer to +Pi or –Pi medium for 24 h (germinated for 5 days on +Pi) ±SD; n = 3; Two-way ANOVA, Post-hoc Tuckey, *p value ≤ 0.05. **b**, ATG8A and ATG8H fail to complement *Δatg8* sensitivity to Pi limitation. Seedlings were germinated for 5 days on +Pi media prior to transfer to medium supplemented with or without Pi (-Pi, +Pi) and imaged after 4 days. Gain of primary root extension was measured. Box plots show medians and interquartile ranges of primary root gain; outliers (>1.53 interquartile range) are shown as black dots. Letters denote statistical differences in condition at p < 0.05 (Two-way ANOVA and Tukey's HSD post hoc test) and n ≥ 13-50.

Supplementary Figure S3

Supplementary Figure S3: ER-stress derived adaptation of root tip proteomes. a, Chemically induced ER stress impacts root proteome. Venn diagram of three overlapping pairwise comparisons for root tip TMT-MS conducted in wildtype (yellow), *pdr2* (magenta) and *atg2* (cyan) plants: wildtype (+Pi) vs wildtype (+Pi+TM) yellow cycle, *pdr2* (+Pi) vs. *pdr2* (+Pi+TM) magenta cycle, *atg2* (+Pi) vs. *atg2* (+Pi+TM) cyan cycle. Protein abundances with $0.6 > FC > 1.3$. Total numbers are depicted in brackets. **b**, Overlap between proteins identified in + Pi + TM conditions enriched in *atg2* compared to wildtype (yellow) or *pdr2-1* (magenta) identified by quantitative MS, $FC > 1.3$. **c**, Proteins of the autophagic pathway are affected in local Pi-mediated root tip proteomic adaptation. Plotted protein abundances of ATG8B/C/E/H, ATG4B, NBR1, C53 and CAN1 in +Pi (light grey), -Pi (dark grey) and +Pi +TM (black) conditions measured via TMT-MS in wildtype, *pdr2* and *atg2*. **d,e** Autophagosome induction in Pi deprived and ER stressed root tips. **d**, GFP-ATG8A derived fluorescence in primary root tips of transgenic (*p35S::GFP-ATG8A*) wildtype and *pdr2* seedlings after germination on +Pi agar (5 d) and subsequent transfer to –Pi medium for 24 h or treatment with 5 µg/mL TM for 6 h. Shown are representative images from three independent experiments fused Z-stacks of the root tip. Scale bars, 50 µm. **e**, Quantification of GFP-ATG8 puncta in root cells per normalized area and number of slices. Box plots show medians and interquartile ranges of detected autophagosomes; outliers (>1.53 interquartile range) are shown as black dots. Letters denote statistical differences in condition at $p < 0.05$ (Two-way ANOVA and Tukey's HSD post hoc test) and $n \geq 6$.

Supplementary Figure S4

Supplementary Figure S4: CAN1 is present in most land plant species. Maximum-likelihood midpoint-rooted phylogenetic tree (250 bootstraps) of 139 predicted CAN1-like proteins (Phytozome). Identified CAN1 like proteins of bryophytes (light green), amborella (turquoise), monocots (leaf green) and dicots (dark green). Full-length CAN1-like sequences are annotated by organism in a numeric order, C-terminally truncated proteins are annotated according to *A. thaliana* HRG1 and 2 respectively.

Supplementary Figure S5: CAN1 is an intrinsically disordered protein. Structure and interaction prediction of CAN1 by AlphaFold2. a, CAN1 is intrinsically disordered. Top five structure prediction by AlphaFold2 colored according to the pLDDT score of the predicted structure. **b**, CAN1 structure in green cartoon, with all five identified AIMs in stick representation (AIM1-3 green, AIM4 orange, AIM5 magenta). **c**, Specific conserved AIMs, with AIM4 and AIM5 exclusively present in full-length CAN1 proteins. **d-g** Predicted AIM-dependent interaction of full length CAN1 with ATG8E. **d**, CAN1 (cartoon, AA 192-210) interacts with ATG8E (surface representation) via AIM5 (stick, magenta), **e**, ATG8E AIM pockets, known as W-site (site 1 –yellow) and L-site (site 2 –slate blue) interacting with W202 and V205. **f**, Predicted interaction of CAN1$^{\Delta AIM5}$ (cartoon, AA 181-210) via AIM4 (stick, orange) and **g**, loss of predicted CAN1 ATG8E interaction in CAN1$^{\Delta AIM4,5}$ (cartoon, AA 181-210).

Supplementary Figure S6

Supplementary Figure S6: CAN1 co-localizes with all ATG8 isoforms in *N. benthamiana*. Localization of GFP-CAN1 variants (CAN1, CAN1$^{\Delta\text{AIM4}}$, CAN1$^{\Delta\text{AIM5}}$ CAN1$^{\Delta\text{AIM4/5}}$) transiently co-expressed with RFP-ATG8A, C-I in *N. benthamiana*. The images are single optical sections of GFP (first column), RFP (second column), chlorophyll A (third column, blue) and merged (including bright field) signals (last column). Scale bars, 10 μm; n ≥ 3.

Supplementary Figure S7: CAN1 protein functions in ER stress-induced autophagy by local Pi sensing. a, GFP-CAN1 derived fluorescence in primary root tips of transgenic (*pUBQ10::GFP-CAN1*) wildtype seedlings after germination on +Pi agar (5 d) and subsequent transfer to –Pi medium for 0-24 h or treatment with 5 μg/mL TM for 6 h. Shown are representative images from three independent experiments fused Z-stacks of the transition zone. Scale bar 20 μm. **b**, Quantification of GFP-CAN1 puncta per stack in the indicated

treatments. Box plots show medians and interquartile ranges of detected puncta; outliers (>1.53 interquartile range) are shown as black dots. Letters denote statistical differences in condition at $p < 0.05$ (Two-way ANOVA and Tukey's HSD post hoc test) and $n \geq 30$. **c**, CAN1-GFP puncta colocalize with mCherry-ATG8E *in vivo*. Co-localization analyses of single plane confocal images obtained from transgenic Arabidopsis roots co-expressing mCherry-ATG8E (magenta) with CAN1-GFP (green) in wildtype. Seedlings were germinated 5 days prior transfer to the indicated treatment for 1 day (+/-Pi) or 5 h (TM). Scale bars, 20 µm, $n \geq 10$. **d**, Pearson's Coefficient (r) analysis of the colocalization of CAN1-GFP and mCherry-ATG8E. Box plots show medians and interquartile ranges with five independent measures per biological replicate, $n \geq 10$. **e**, *CAN1* expression relative to UBC9 in gain of primary root tip 1 day after transfer of wildtype, *pdr2*, *ire1a/b* and *atg2*. Letters denote statistical differences in condition at $p < 0.05$ (Two-way ANOVA and Tukey's HSD post hoc test) and $n = 3$. **f**, *pCAN1::GUS* activity in 10 days old wildtype seedlings after transfer to the indicated media for two days or incubation with TM for 6 h, $n = 3$. Overview of the seedlings (I-III) scale bar 1 cm, primary root tip (IV-IX) with IV, VI, VIII scale bar 100 µm and V, VII, IX scale bar 1mm; $n \geq 6$.

Supplementary Figure S8

Supplementary Figure S8: CAN1 functions in ER stress-mediated autophagy. a, Transcriptional regulation of *CAN1* is not altered in SALK-Insertion line. Analysis of *CAN1* relative transcript level (normalized to UBC9 expression) of wildtype and *SALK_092190* insertion line seedlings (germinated for 5 days on +Pi media) ±SD; n ≥ 3; Students t-test. **b**, sequence analysis of *CAN1* (AT5G40690). Scheme of two *can1* CRISPR mutants, *can1-c1* and *can1-c3*. Both mutants contain deletions/insertions where *can1-c1* harbors a 118-bp insertion, while the *can1-c3* mutant possesses a 74-bp deletion. The deletions/insertions lead to amino acids changes and premature stop codons in both mutants. **c**, Expression analysis of *CAN1* by semiquantitative RT-PCR by using cDNA isolated from wildtype, *SALK_092190*, *can1-c1*, and *can1-c3* seedlings. **d**, Amount of *pCAN1::CAN1-GFP* in wildtype, *atg2* and *atg5* seedlings (germinated for 5 days on +Pi media) were analyzed by Western Blot. TCE was used as loading control, n = 4. **e**, 5-day old seedlings expressing *pCAN1::CAN1-GFP* in wildtype and *pdr2* mutant were transferred to the indicated media for 24 h, TM (300 ng/µl) ConA (1µM). Whole roots were harvested and analyzed by immunoblot. TCE was used as loading control, n = 3. **f, g** 5 days-old seedlings of the indicated genotypes were grown on full (Ctrl) and N-depleted media for 7 days. **g**, Analysis of seedling weight after treatment. Box plots show medians and interquartile ranges of primary root gain. Letters denote statistical differences in treatment condition at p < 0.05 (One-way ANOVA and Tukey's HSD post hoc test) and n = 8-16. **h**, Loss-of *CAN1* does not rescue carbon starvation in *atg5* mutant plants. 5 days-old seedlings of the indicated genotypes were germinated on full media, prior to carbon starvation for 12 days, followed by recovery on full media for one week, ±SD; n = 3; One-way ANOVA, Post-hoc Tuckey, *p value ≤ 0.05.

a

Supplementary Figure S9: LPR1-mediated local Pi response is not influenced by CAN1.
a,b After germination (5 d, +Pi), seedlings were transferred to +Pi (**a**) or –Pi (**b**) medium (25 μM Fe). Top row: Three days after transfer, root tips of wildtype, *pdr2*, *can1-c1*, *can1-c3 and pUBQ10::GFP-CAN1* were monitored for Fe^{3+} accumulation by Perls staining coupled to

diaminobenzidine (DAB) intensification. Bottom row: One day after transfer, root tips were monitored for callose deposition by aniline blue staining. Shown are representative images (n ≥ 20). Scale bars 50 μm.

Supplementary Figure S10

Supplementary Figure S10: CAN1 intersects with the autophagic pathway in ER stress.
a, Enrichment of proteins interacting with *p35S::GFP-ATG8A*, *pUBQ10::GFP-CAN1* in wildtype and *pCAN1::CAN1-GFP* in *atg5* background in + Pi compared with a control expressing free GFP and represented by a volcano plot. The horizontal grey line indicates the threshold above which proteins are enriched (P value <0.03, quasi-likelihood negative binomial generalized log-linear model), and the vertical dashed line indicates the threshold where the proteins' $\log_2$FC is > 1. **b**, Western blot analysis from immunoprecipitation (IP) experiments of tobacco leaves co-infiltrated with *p35S::CAN1–mCherry* together with

p35S::DGL1-GFP, *pUBQ10::GFP-DGL1* or *p35S::GFP-HDEL*. Input and bound proteins were visualized with anti-GFP and anti-RFP antibodies, n = 3. **c**, Amount of *pCAN1::CAN1-GFP* in wildtype, *pdr2* and *atg5* roots were analyzed by immunoblot. Seedlings were germinated 5 days on +Pi prior to transfer to the indicated media for 24 h or ER stress treatment with 5 µg/ml TM and 1 µM ConA for 6 h. TCE was used as loading control, n = 4. **d**, Autophagosome quantification in roots of wildtype and *can1-c1* expressing *p35S::GFP-ATG8A*. Seedlings were germinated for 5 days on +Pi media prior to transfer for 24 h to +Pi or –Pi media or incubated with 5 µg/ml TM or 1µM torin for 6 h for autophagic induction. Scatter plot, median is shown as bold line, letters denote statistical differences at $p < 0.05$ (Two-way ANOVA and Tukey's HSD post hoc test), n ≥ 18. **e**, Representative image of autophagosome volume quantification in Arivis. *p35S::GFP-ATG8A in* wildtype 24 h post transfer to –Pi is shown. Scale = 5 µm. Color-coded autophagosome volume according to the scale.