How Ant Genomes Repeatedly Reinvent Venom

Frank Andreas Weitz[1], Francisco Hita Garcia[2], Bjoern Marcus von Reumont[3],
Burkhard Rost[1,4,5,$], Ivan Koludarov[1,$,@]

1 School of Computation, Information, and Technology (CIT), Department of Informatics, Bioinformatics &
   Computational Biology, TUM (Technical University of Munich), 85748 Garching/Munich, Germany
2 Museum für Naturkunde Berlin, 10115 Berlin, Germany
3 Staatliches Museum für Naturkunde Karlsruhe, 76133 Karlsruhe, Germany
4 TUM School of Life Sciences Weihenstephan (WZW), Alte Akademie 8, Freising, Germany
5 Institute for Advanced Study (TUM-IAS), Lichtenbergstr. 2a, 85748 Garching/Munich, Germany
@ corresponding author
$ senior authors contributed equally to this work
Running Title: Ant Venoms
Journal: PNAS (Nature? FHG thinks we have a chance)

ABSTRACT

Background: Venoms from ant (*Formicidae*) are chemically extraordinarily diverse, yet their
genomic architecture and evolutionary dynamics remain opaque. The "*aculeatoxin*
hypothesis" assumes that most venom peptides in stinging insects (*aculeate*) evolutionarily
derive from a single gene family, the aculeatoxins. The recent refutation of this hypothesis for
bees raises fundamental questions about venom evolution in ants, which contribute most
aculeatoxins.
Methods: In order to trace venom peptide evolution, we performed synteny-aware
comparative genomics across 25 ant species spanning major subfamilies. We analyzed
phylogeny through sequences, structures, and embeddings from protein Language Model
(pLMs).
Results: We identified three conserved genomic regions (GR1-GR3) as evolutionary hotspots
for ant venom genes, each exhibiting distinct evolutionary dynamics. Most remarkably, we
discovered genuine melittin orthologs in ants at the conserved bee syntenic position (GR2),
pushing the origin of this scaffold back to early aculeates, with a possible origin deeper in
Hymenoptera. Gene copy numbers vary dramatically (0-17 genes per region), with predatory
species showing expansions and *formicine* ants (subfamily Formicinae, which rely on formic
acid spraying) showing reductions. Twenty-two distinct toxin clades emerge, with region-
specific distributions suggesting repeated recruitment to conserved platforms.
Significance: Ants evolutionarily succeed by combining single-copy conservation (bee-like
at GR2), massive gene duplication (snake-like at GR1), and repeated lineage-specific
recruitment to conserved genomic platforms (GR3). This multi-modal evolution on stable
genomic scaffolds, evidenced here, reconciles previously conflicting models of venom
evolution and reveals how genomic architecture constrains, yet repeatedly enables, molecular
innovation. Our findings highlight a general principle of genome evolution: complex adaptive
traits can arise not from a single origin or mechanism, but through recurrent reuse of
permissive genomic loci shaped by ecology. Such principles are likely relevant not only to
other venomous animals, but more broadly to the evolution of complex multi-genic traits.
29
30 **Keywords:** Gene evolution, Formicidae, Toxins, Venom Evolution

31 **Abbreviations: CDS** (coding DNA sequence); **GR** (genomic region); **K⁺** (K^+, potassium ion);
32 **MYRTX** (Myrmecitoxin, proposed ant venom toxin family); **NaV** (voltage-gated sodium
33 channel); **NCBI** (National Center for Biotechnology Information); **ORF** (open reading frame);
34 **PONTX** (Poneratoxin, proposed ant venom toxin family); **ProtT5** (Protein Transformer T5,

protein language model); **UMAP** (Uniform Manifold Approximation and Projection for dimensionality reduction); **UniProt** (Universal Protein Resource); **CCDC-p170** (Coiled-coil domain–containing protein 170).

INTRODUCTION

Venom evolution at the crossroads. Venom systems represent one of nature's most successful innovations, evolving independently over 100 times across the animal kingdom (Schendel et al., 2019). Two contrasting paradigms have emerged as explanations. The "snake model" explains venom evolution as a **duplication-driven process**, in which repeated gene duplication, positive selection, and birth-and-death dynamics generate expansive toxin gene families (Casewell et al., 2012). In contrast, the "wasps and bees model," emerging from recent genomic studies, reveals venoms largely encoded by **single-copy genes retained at conserved syntenic loci**, where **regulatory co-option rather than gene duplication** underlies venom diversification (Martinson et al., 2017; Koludarov et al., 2023). Where do ants fit in this dichotomy? With over 13,800 species exhibiting extraordinary venom diversity, from protein-rich cocktails to alkaloid-dominated secretions to formic acid sprays, ants potentially hold the key to understanding how different evolutionary strategies can operate within a single lineage (Chen, 2023; Kass et al., 2022; Touchard et al., 2016).

Aculeatoxin controversy and ant centrality. The influential "aculeatoxin hypothesis" proposed that all short aculeate venom peptides derive from a single ancestral superfamily, based primarily on similarities between signal and propeptide sequences (Robinson et al., 2018). Since *Formicidae* (ant) sequences constitute about 70% of proposed members of aculeatoxins family, testing this hypothesis requires comprehensive genomic analysis across *Formicidae*.
Our recent work on 32 bee genomes found no support for aculeatoxins: bee toxins occupy lineage-specific loci and cluster by taxonomy rather than function in protein embedding space (Koludarov et al., 2023). However, that study included limited ant genomes, one showing a melittin-like gene from Asian clonal ant *Vollenhovia emeryi* at the conserved position but with a highly divergent mature sequence, possibly even a pseudogene. The single published *venomics*-oriented ant genome (*Tetramorium bicarinatum*) revealed tandem toxin arrays but lacked comparative context (Touchard et al., 2024).

Here, we present the first comprehensive genomic analysis for the evolution of ant toxic peptides, examining 25 species across all major subfamilies. We integrate: (1) microsynteny analysis to establish orthology and trace genomic architecture; (2) phylogenetics to resolve evolutionary relationships; and (3) protein embeddings to detect remote homology and convergence patterns. This multi-faceted approach reveals how ants employ multiple evolutionary strategies simultaneously on conserved genomic platforms.

RESULTS AND DISCUSSION

Three conserved regions suggest three different evolutionary strategies

Our synteny-aware analysis revealed three main genomic regions (dubbed GR1-GR3, Figures 1, 2 and 3) repeatedly harboring ant venom genes, each exhibiting fundamentally different evolutionary dynamics that illuminate the flexibility of venom evolution within a single lineage.

GR1: Snake-like duplication factory

GR1 exemplifies classical *birth-and-death* evolution in which repeated gene duplication is coupled with frequent gene loss and pseudogenization, resulting in extreme copy-number variation (0-17 loci, Fig. 1). The region shows striking ecological correlation: predatory species with functional stings (*Tetramorium bicarinatum*: 17 copies; *Odontomachus*, *Dinoponera*, *Harpegnathos*: 4-8 copies) maintain expanded arrays, while formicine ants that spray formic acid retain single or no copies.

Within expanded genomic gene arrays generated by local duplication, we observe the full spectrum of birth-and-death dynamics: intact multi-exon genes preserving signal peptide and propeptide architectures, partially degraded single-exon variants, and fragmentary pseudogenes. The retention of introns in duplicated copies argues against retroposition (RNA-mediated duplication) and instead supports local DNA-level gene duplication as the primary mechanism of array expansion.

Phylogenetically, GR1 toxins segregate by ant lineage: *poneroid* ants share clades S and R (including the antimicrobial M-PONTX-Dq3a), whereas *myrmicine* ants (subfamily *Myrmicinae*) possess distinct clades T, U, and W. Within the genus *Tetramorium*, both ancient clades shared across the genus, including those retained in *Tetramorium alpestre*, and recent lineage-specific expansions (Clade U) are observed, illustrating ongoing diversification. This pattern identifies *Tetramorium* as a hotspot of recurrent local gene duplication, exhibiting unusually frequent GR1 copy turnover relative to other ant lineages. Such a high copy-turnover might be related to extreme survival pressure (Schlick-Steiner et al., 2010).

GR2: Melittin locus deeply conserved

The most striking discovery is the presence of two-exon venom genes at the exact syntenic position (MESR3–CCDC-p170 interval) at which bees encode melittin (Fig. 2). These ant sequences, U11-MYRTX-Tb1a and U13-MYRTX-Tb1a in *Tetramorium*, plus orthologs in *Pogonomyrmex*, preserve the standard architecture of aculeate venom peptides: the signal- and pro-peptide in exon 1, and the mature peptide in exon 2.

This conservation of position and gene model, combined with functional evidence (U11-MYRTX-Tb1a causes paralysis in insects, presumably via K^+ channel modulation or by permeabilizing membranes in a pore-forming manner (Barassé et al., 2023; Boy et al., 2025; Peigneur et al., 2026)), strongly suggests orthology with bee melittin. The implications are profound: the melittin scaffold likely existed in the *aculeate* ancestor over 200 million years ago, with lineage-specific sequence divergence obscuring the relation at the sequence level.

Some ants retain two-exon genes, others have single-exon variants or completely lost the gene. This patchy diversity across ants somewhat mirrors the bee pattern where melittin is preserved in most lineages but lost in a few others. This "use it or lose it" pattern at a conserved locus contrasts sharply with the dynamic expansions at GR1.

GR3: Recruitment platform

GR3 represents a novel evolutionary mode in which a stable syntenic genomic scaffold is repeatedly colonized by distinct toxin families in different ant lineages (Fig. 3). *Poneroid* ants recruit poneratoxins at this locus, including the NaV-modulating PC-PONTX-Pc1a responsible for extremely painful stings. In contrast, *myrmicine* ants (subfamily *Myrmicinae*) independently deploy MYRTX peptides of unknown function (U*-MYRTX) alongside lineage-specific radiations. Notably, the myrmicine species *Cardiocondyla obscurior* exhibits a unique five-gene tandem array of Clade F toxins which, although absent from other taxa, forms a basal clade relative to all other GR3-associated sequences.

This pattern, "same genomic address with different molecular tenants", suggests that certain chromosomal contexts are inherently permissive for venom gene expression. This might be due to regulatory landscapes or to the chromatin architecture. The mosaic of multi-exon genes, single-exon variants, and fragments within GR3 again supports *birth-and-death dynamics* which, in contrast to the situation for GR1, operates on taxonomically diverse substrates.

Aculeatoxins: 22 clades tell 22 stories

Our phylogenetic analysis recovered 22 distinct toxin clades with region-biased distributions. Crucially, these clades segregate by ant taxonomy (subfamily or tribe) rather than forming a single pan-aculeate superfamily. Together with syntenic evidence showing that different toxin clades occupy distinct genomic loci, this pattern is inconsistent with the aculeatoxin hypothesis, which posits a monophyletic origin of all aculeate toxins from a single ancestral gene at the stem *Aculeata*. Instead, our results support a polyphyletic origin of aculeate venom toxins, involving independent recruitment or diversification events within different aculeate lineages. This does not preclude deeper homology among venom genes at the level of stem *Hymenoptera*, but it rejects a single shared aculeate-specific toxin ancestor.

Embeddings from protein Language Models (pLMs) provide orthogonal support: when visualized in UMAP space, ant toxins cluster by genomic region and taxonomic origin, not by functional similarity (Fig. 4). Even when including signal peptides (the primary evidence for classification as *aculeatoxins*), sequences separate by lineage. This pattern, also seen in our bee analysis (Koludarov et al., 2023), indicates convergent evolution of similar functions rather than divergence from a common ancestral toxin.

Ecological drivers and functional correlations

The correlation between venom complexity and ecology is striking. Predatory ants with functional stings consistently show expanded repertoires at GR1 and GR3. *Formicine* ants, which secondarily lost the ability to sting and evolved formic acid spraying, show minimal peptide toxins, often just single genes or complete absence at these loci.

This pattern extends to specific clades: poneratoxin-related sequences (targeting vertebrate pain pathways) occur in large predatory species that use stings defensively against vertebrates. *Myrmicine*-specific expansions may target insect prey or microbes. The *Cardiocondyla*-specific radiation at GR3 could reflect unique social or ecological pressures in this genus.

Synthesis: ants reveal the full complexity of venom evolution

Our findings position ants as uniquely revealing the full complexity of venom evolution across *Hymenoptera*. Five themes emerge.

(1) Multiple evolutionary mechanisms coexist. Unlike bees, which primarily rely on regulatory co-option of single-copy genes, or snakes, which largely diversify venoms through extensive gene duplication, ants employ both duplication-driven expansion and regulatory recruitment, alongside a lineage-specific locus reuse mechanism in which different toxin families independently colonize conserved syntenic regions. Possibly the number of different evolutionary mechanisms provides a better proxy to the complexity of ants than the mere number of lineages or species (roughly similar between ants and bees).

(2) Mosaic patterns reveal deep but uneven homology. The discovery of the melittin locus in ants demonstrates that some venom positions are genuinely orthologous across *aculeates*, preserved for over 200 million years despite divergence beyond recognition by sequence similarity alone. In contrast, other venom loci show clear lineage-specific origins.

(3) Genomic architecture channels innovation. The recurrent involvement of GR1–GR3 across ant evolution suggests that these conserved syntenic loci possess intrinsic properties that facilitate venom gene expression and turnover. These could include regulatory context, chromatin accessibility, or nuclear positioning.

(4) Ancient loci are differentially retained. Functional toxin genes in one lineage correspond to pseudogenes in others (e.g. melittin, mastoparan, and poneratoxin loci (Koludarov et al., 2023)). This observation indicates ancient genomic presence followed by lineage-specific retention or loss.

(5) Ecological specialization drives genomic complexity. The strong interplay between predatory venom use, such as prey capture, immobilization, and pain induction, and toxin gene expansion suggests that ecological reliance on venom actively promotes the retention, diversification, and amplification of duplicated genes, rather than duplication being selectively neutral.

Overall, our results are incompatible with both models of entirely independent *de novo* origins of venom systems and with a single monophyletic *aculeate* toxin ancestry. Instead, advanced sequence analysis enabled by protein language models and comparative genomics reveals a dynamic evolutionary landscape characterized by deep homology, repeated recruitment, and ongoing innovation in aculeate venom systems.

Figure 1: Genomic Region 1 (GR1) shows duplication-driven birth-and-death dynamics.

GR1 is a recurrent venom locus displaying extensive copy-number variation across ant genomes (0–17 toxin loci per species). Gene models at this region span intact multi-exon genes, single-exon derivatives, and truncated fragments/pseudogenes, consistent with duplication and loss.

Arrowheads show orientation and internal notches mark exon–exon junctions (two notches → three exons; one notch → two exons; no notches → single-exon). The drawn horizontal lengths are purely schematic for clarity (not to scale) and should not be interpreted as quantitative measures of coding-sequence length; the narrow triangular shape denotes a truncated fragment, while the long arrows containing gene names denotes a representative full-length non-toxin construct.

204
205
206

Figure 2: Genomic Region 2. Deeply conserved melittin locus.

GR2 occupies the MESR3–CCDC-p170 syntenic interval and in multiple ant lineages carries two-exon venom
genes with the canonical organization (exon 1 = signal/propeptide; exon 2 = mature peptide), exemplified by U11-
MYRTX-Tb1a and U13-MYRTX-Tb1a in Tetramorium and orthologs in Pogonomyrmex. Positional and structural
conservation, together with functional evidence for at least one ant paralog (U11-MYRTX-Tb1a causes paralysis in
insects via K$^+$-channel modulation (Boy et al., 2025)), supports orthology with bee melittin; this implies a melittin-
like scaffold present in the aculeate ancestor and differentially retained, simplified (single-exon variants) or lost
across lineages.

Arrowheads show orientation and internal notches mark exon–exon junctions (two notches → three exons; one
notch → two exons; no notches → single-exon). The drawn horizontal lengths are purely schematic for clarity (not
to scale) and should not be interpreted as quantitative measures of coding-sequence length; the narrow triangular
shape denotes a truncated fragment, while the long arrows containing gene names denotes a representative full-
length non-toxin construct.

GR2 is conserved in these ants, but we could not locate any toxin genes in: Odontomachus brunneus,
Trachymyrmex zeteki, Solenopsis invicta, Pseudomyrmex gracilis, Pogonomyrmex barbatus, Ooceraea biroi,
Nylanderia fulva, Monomorium pharaonic, Linepithema humile, Harpegnathos saltator, Dinoponera quadriceps,
Camponotus vicinus, Paraponera clavata, Leptanilla sp.

223
224
225

227
228
Figure 3: Genomic Region 3. Repeated recruitment loci.

GR3 is a stable syntenic scaffold repeatedly colonized by different toxin families in different lineages. Poneroids
deploy poneratoxins here (e.g., the NaV-modulating PC-PONTX-Pc1a), myrmicines recruit MYRTX toxins and
lineage-specific radiations, and Cardiocondyla contains a unique five-gene tandem array of Clade F toxins. The
repeated occupation of the same genomic address by phylogenetically distinct toxin families, together with mixed
gene architectures (multi-exon, single-exon, fragments), indicates recurrent recruitment coupled with birth-and-
death turnover acting on a taxonomically diverse substrate.

Arrowheads show orientation and internal notches mark exon–exon junctions (two notches → three exons; one
notch → two exons; no notches → single-exon). The drawn horizontal lengths are purely schematic for clarity (not
to scale) and should not be interpreted as quantitative measures of coding-sequence length; the narrow triangular
shape denotes a truncated fragment, while the long arrows containing gene names denotes a representative full-
length non-toxin construct.

GR3 is conserved in these ants, but we could not locate any toxin genes in: Atta colombica, Ooceraea biroi,
Pseudomyrmex gracilis, Nylanderia fulva, Lasius niger, Cataglyphis hispanica, Camponotus vicinus, Formica selysi,
Mystrium camillae.

244
245
246

248
249
Figure 4: ProtT5 Embeddings show disperse clustering, with sequences from the same genomic region favoring
proximity.

Protein-sequence embeddings (including signal peptides; see Methods) projected into two-dimensional UMAP
space. Points are colored by Genomic Region. Embeddings group by lineage and genomic region rather than by
inferred function, supporting convergent recruitment of toxin-like sequences rather than a single ancestral
"aculeatoxin" family.

CONCLUSION

Ant venom evolution operates through multiple mechanisms on three conserved genomic
platforms. Our discovery of melittin orthologs at the bee syntenic position established a deep
conservation of genomic architecture despite extreme sequence divergence. The three
regions show three modes: massive duplication (GR1), single-copy conservation (GR2), and
platform recruitment (GR3). These findings reconcile conflicting models of venom evolution,
showing how different strategies can operate simultaneously within a single lineage
depending on ecological pressures and genomic context. Combined with evidence for
convergent evolution on conserved scaffolds this establishes a new paradigm: venom
evolution is channeled by genomic architecture but diversified by ecological selection. Such
a multi-modal evolutionary framework likely extends beyond ants. The three principles
revealed here - conserved platforms, lineage-specific recruitment, ecology-driven expansion
- may explain venom diversity across the tree of life.

METHODS

Data acquisition
Genome assemblies. We retrieved publicly available genome assemblies for the species
included in this study from NCBI (accession numbers and assembly versions are given in
Supplementary Table 1). Assemblies were used as provided (no additional scaffolding or
reassembly) and were chosen to prioritize high contiguity and available annotation where
possible.
Toxin sequences. We assembled a reference set of known peptide toxin sequences by
downloading reviewed and non-redundant (less than 100% sequence identity) records from
UniProt (The UniProt Consortium et al., 2025) and by manually extracting additional
sequences reported in the literature (mass-spectrometry, transcriptomics, and genomic toxin
descriptions). The combined data set (UniProt + literature) served as the query database for
all subsequent genome searches.
282
Genome searches with MMseqs2
We searched each genome for toxin query and homologs of the toxin query set using
MMseqs2 (Steinegger & Söding, 2017). Protein searches were performed using translated
search modes when necessary to accommodate exon/intron structure in draft assemblies.
MMseqs2 was run with high sensitivity to maximize recovery of divergent toxin homologs
(sensitivity = 9.5) and an e-value cutoff of 1e-3 for initial hit detection.
289
Manual curation and exon boundaries (UGENE + BDGP splice site prediction)
MMseqs2 hits were imported into UGENE (Okonechnikov et al., 2012) for manual inspection.
For each candidate locus we applied UGENE's built-in Smith–Waterman algorithm to confirm
alignment quality and to detect whether multiple adjacent peptide matches corresponded to
separate exons. Short, isolated matches (very small peptide fragments recurring at multiple
genomic locations) were considered likely repeats and discarded. Where adjacent hits from
the same query occurred on the same scaffold, these were treated as putative exon pairs and
inspected for canonical splice signals.
To define exon boundaries, we combined automated splice site prediction available at
fruitfly.org (Reese et al., 1997) with manual sequence matching. Predicted donor/acceptor
sites were used as a guide and were accepted if (1) they mostly aligned with UGENE-identified
exon boundaries via Smith–Waterman alignment, and (2) the resulting exon concatenation

produced a coherent open reading frame consistent with the query toxin. In ambiguous cases we manually adjusted exon start/stop positions to preserve reading frame and expected signal/propeptide features. Percent-identity thresholds used during manual verification were relaxed relative to standard protein searches (typically 25–35% depending on query length) to accommodate rapid toxin evolution while limiting spurious matches.

Extraction of flanking genes and orthology searches

For each confirmed toxin locus we recorded the immediate upstream and downstream protein-coding genes (flanking genes) based on the genome annotation. Coding sequences of these flanking genes were extracted separately and used as queries in a second round of MMseqs2 searches against the full set of genomes. The aim was to (1) locate homologous genomic neighborhoods in other assemblies and (2) distinguish orthologous toxin insertions from lineage-specific insertions or assembly artifacts.

Each flanking CDS was searched independently using MMseqs2 with parameters matching the toxin searches (sensitivity = 9.5; e-value ≤ 1e-3). Hits were compiled and inspected visually to determine which scaffolds carried the conserved gene order expected for orthology. In regions where both flanking genes were recovered but no annotated toxin was present, we re-examined the scaffold with exon-level Smith–Waterman searches (UGENE) to discover highly divergent toxin orthologs.

Synteny analysis and plotting

From the manual curation of flanking gene hits we inferred conserved genomic neighborhoods. We recorded the presence/absence and relative order of flanking genes and noted whether an identifiable toxin CDS co-occurred at the locus. Using these data we produced synteny figures representing the four principal toxin loci studied.

Multiple sequence alignment and outlier filtering

Toxin amino-acid sequences were aligned with MAFFT (Katoh et al., 2002) using the L-INS-i style parameters appropriate for short, divergent peptides: --localpair --maxiterate 1000. Alignments were inspected visually and a small number of clear outliers (typically ~2–3 sequences per alignment) that degraded alignment quality were removed. The alignment was then recomputed with the same MAFFT parameters to obtain the final alignment used for phylogenetic inference.

Phylogenetic inference

We inferred maximum-likelihood trees from the MAFFT alignment using IQ-TREE 2 (Nguyen et al., 2015). The analysis was run with model selection (MFP) augmented by a fixed set of mixture matrices and branch-rate options, and with intensive support estimation as follows:
iqtree2 -s toxins.fasta -st AA -m MFP -madd LG+C60+F+G -mrate R -bb 10000 -alrt 10000 -bnni -allnni -pers 0.2
Resulting trees were filtered to collapse any node with bootstrap support below 50% to avoid over-interpretation of weakly supported relationships. We then manually inspected the collapsed topology, defined robust ant-specific clades, and annotated these clades for downstream visualization.

Integration of phylogeny and synteny

Clade assignments from the IQ-TREE analysis were mapped back onto the synteny plots. Each tree-defined ant toxin clade was given a distinct color and these colors were used to

350 mark toxin occurrences in the synteny diagrams to facilitate interpretation of the relationship
351 between sequence evolution and genomic context.
352
Sequence embedding and low-dimensional visualization
To provide an orthogonal, alignment-free view of sequence relationships, we generated
per-protein embeddings with the ProtT5 (Elnaggar et al., 2021) protein language model.
ProtT5 embeddings were computed for all sequences, averaged across residues to produce
one fixed-length vector per protein, and subsequently projected to two dimensions using
UMAP via the ProtSpace (Senoner et al., 2025) workflow. Embedding plots were colored by
(1) genomic locus, (2) scaffold/species origin, and (3) phylogenetic clade to aid visual
interpretation.

REFERENCES

Barassé, V., Jouvensal, L., Boy, G., Billet, A., Ascoët, S., Lefranc, B., Leprince, J., Dejean, A., Lacotte, V., Rahioui, I., Sivignon, C., Gaget, K., Ribeiro Lopes, M., Calevro, F., Da Silva, P., Loth, K., Paquet, F., Treilhou, M., Bonnafé, E., & Touchard, A. (2023). Discovery of an Insect Neuroactive Helix Ring Peptide from Ant Venom. *Toxins*, *15*(10), 600. https://doi.org/10.3390/toxins15100600

Boy, G., Jouvensal, L., Téné, N., Carayon, J.-L., Bonnafé, E., Paquet, F., Treilhou, M., Loth, K., & Billet, A. (2025). A New Kv1.3 Channel Blocker from the Venom of the Ant Tetramorium bicarinatum. *Toxins*, *17*(8), 379. https://doi.org/10.3390/toxins17080379

Casewell, N. R., Huttley, G. A., & Wüster, W. (2012). Dynamic evolution of venom proteins in squamate reptiles. *Nature Communications*, *3*, 1066. https://doi.org/10.1038/ncomms2065

Chen, J. (2023). Chemistry and Functions of Imported Fire Ant Venom. *Toxins*, *15*(8), 489. https://doi.org/10.3390/toxins15080489

Elnaggar, A., Heinzinger, M., Dallago, C., Rehawi, G., Wang, Y., Jones, L., Gibbs, T., Feher, T., Angerer, C., Steinegger, M., & others. (2021). Prottrans: Toward understanding the language of life through self-supervised learning. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, *44*(10), 7112–7127.

Kass, J. M., Guénard, B., Dudley, K. L., Jenkins, C. N., Azuma, F., Fisher, B. L., Parr, C. L., Gibb, H., Longino, J. T., Ward, P. S., Chao, A., Lubertazzi, D., Weiser, M., Jetz, W., Guralnick, R., Blatrix, R., Lauriers, J. D., Donoso, D. A., Georgiadis, C., … Economo, E. P. (2022). The global distribution of known and undiscovered ant biodiversity. *Science Advances*, *8*(31), eabp9908. https://doi.org/10.1126/sciadv.abp9908

Katoh, K., Misawa, K., Kuma, K., & Miyata, T. (2002). MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research*, *30*(14), 3059–3066. https://doi.org/10.1093/nar/gkf436

Koludarov, I., Velasque, M., Senoner, T., Timm, T., Greve, C., Hamadou, A. B., Gupta, D. K., Lochnit, G., Heinzinger, M., Vilcinskas, A., Gloag, R., Harpur, B. A., Podsiadlowski, L., Rost, B., Jackson, T. N. W., Dutertre, S., Stolle, E., & von Reumont, B. M. (2023). Prevalent bee venom genes evolved before the aculeate stinger and eusociality. *BMC Biology*, *21*(1), 229. https://doi.org/10.1186/s12915-023-01656-5

Martinson, E. O., Mrinalini, Kelkar, Y. D., Chang, C.-H., & Werren, J. H. (2017). The Evolution of Venom by Co-option of Single-Copy Genes. *Current Biology*, *27*(13), 2007-2013.e8. https://doi.org/10.1016/j.cub.2017.05.032

Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and Evolution*, *32*(1), 268–274. https://doi.org/10.1093/molbev/msu300

Okonechnikov, K., Golosova, O., Fursov, M., & the UGENE team. (2012). Unipro UGENE: A unified bioinformatics toolkit. *Bioinformatics*, *28*(8), 1166–1167. https://doi.org/10.1093/bioinformatics/bts091

Peigneur, S., Tibery, D., & Tytgat, J. (2026). The Helix Ring Peptide U11 from the Venom of the Ant, Tetramorium bicarinatum, Acts as a Putative Pore-Forming Toxin, Not a New Kv1.3 Channel Blocker. Comment on Boy et al. A New Kv1.3 Channel Blocker from the Venom of the Ant Tetramorium bicarinatum. Toxins 2025, 17, 379. *Toxins*, *18*(1), 44. https://doi.org/10.3390/toxins18010044

Reese, M. G., Eeckman, F. H., Kulp, D., & Haussler, D. (1997). Improved splice site detection in Genie. *Proceedings of the First Annual International Conference on Computational Molecular Biology*, 232–240.

Robinson, S. D., Mueller, A., Clayton, D., Starobova, H., Hamilton, B. R., Payne, R. J., Vetter, I., King, G. F., & Undheim, E. A. B. (2018). A comprehensive portrait of the venom of the giant red bull ant, Myrmecia gulosa, reveals a hyperdiverse hymenopteran toxin gene family. *Science Advances*, *4*(9), eaau4640. https://doi.org/10.1126/sciadv.aau4640

Schendel, V., Rash, L. D., Jenner, R. A., & Undheim, E. A. B. (2019). The Diversity of Venom: The Importance of Behavior and Venom System Morphology in Understanding Its Ecology and Evolution. *Toxins*, *11*(11), 666. https://doi.org/10.3390/toxins11110666

Schlick-Steiner, B. C., Steiner, F. M., Seifert, B., Stauffer, C., Christian, E., & Crozier, R. H. (2010). Integrative Taxonomy: A Multisource Approach to Exploring Biodiversity. *Annual Review of Entomology*, *55*(1), 421–438. https://doi.org/10.1146/annurev-ento-112408-085432

Senoner, T., Olenyi, T., Heinzinger, M., Spannagl, A., Bouras, G., Rost, B., & Koludarov, I. (2025). ProtSpace: A Tool for Visualizing Protein Space. *Journal of Molecular Biology*, 168940. https://doi.org/10.1016/j.jmb.2025.168940

Steinegger, M., & Söding, J. (2017). MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nature Biotechnology*, *35*(11), 1026–1028. https://doi.org/10.1038/nbt.3988

The UniProt Consortium, Bateman, A., Martin, M.-J., Orchard, S., Magrane, M., Adesina, A., Ahmad, S., Bowler-Barnett, E. H., Bye-A-Jee, H., Carpentier, D., Denny, P., Fan, J., Garmiri, P., Gonzales, L. J. D. C., Hussein, A., Ignatchenko, A., Insana, G., Ishtiaq, R., Joshi, V., … Zhang, J. (2025). UniProt: The Universal Protein Knowledgebase in 2025. *Nucleic Acids Research*, *53*(D1), D609–D617. https://doi.org/10.1093/nar/gkae1010

Touchard, A., Aili, S., Fox, E., Escoubas, P., Orivel, J., Nicholson, G., & Dejean, A. (2016). The Biochemical Toxin Arsenal from Ant Venoms. *Toxins*, *8*(1), 30. https://doi.org/10.3390/toxins8010030

Touchard, A., Barassé, V., Malgouyre, J.-M., Treilhou, M., Klopp, C., & Bonnafé, E. (2024). The genome of the ant Tetramorium bicarinatum reveals a tandem organization of venom peptides genes allowing the prediction of their regulatory and evolutionary profiles. *BMC Genomics*, *25*(1), 84. https://doi.org/10.1186/s12864-024-10012-y

SUPPLEMENTARY MATERIAL

Supplementary Figure 1:

Maximum-likelihood phylogenetic tree of ant toxins together with Hymenopteran toxins such as melittins. Dot sizes next to nodes in the tree indicate bootstrap values as indicated in the legend. The inner annotation ring displays the phylogenetic clades of ant toxins. The middle ring displays the genomic region a toxin is located in. The outer ring contains only the toxins that did not come from the ant toxin analysis directly, but from an unpublished manuscript about Hymenopteran venoms.

Synteny of Flanking Genes Surrounding the Toxin Genes – Genomic Region 1

451
452 *Supplementary Figure 2: Synteny Figure of GR1 like Figure 1, but with more flanking genes and less detail on*
453 *toxins. Information on Scaffold / Chromosome are given.*

454
455

457
458 *Supplementary Figure 3: Synteny Figure of GR2 like Figure 2, but with more flanking genes and less detail on*
459 *toxins. Information on Scaffold / Chromosome are given.*

460
461
462
463

Synteny of Flanking Genes Surrounding the Toxin Genes – Genomic Region 3

464
465 *Supplementary Figure 4: Synteny Figure of GR3 like Figure 3, but with more flanking genes and less detail on*
466 *toxins. Information on Scaffold / Chromosome are given.*

Supplementary Table 1. Species codes, full species names, and NCBI assembly accession numbers (including version) for the genomes used in this study.

Abbrev	Species	NCBI Assembly Accession
Ains	Acromyrmex insinuator	GCA_017607455.1
Acol	Atta colombica	GCA_001594045.1
Cvic	Camponotus vicinus	GCA_025532165.1
Cobs	Cardiocondyla obscurior	GCA_019399895.2
Chis	Cataglyphis hispanica	GCA_021464435.1
Dqua	Dinoponera quadriceps	GCF_001313825.1
Fsel	Formica selysi	GCA_009859135.1
Hsal	Harpegnathos saltator	GCF_003227715.1
Amel	Apis mellifera	GCF_003254395.2
Vvel	Vespa velutina	GCF_912470025.1
Lnig	Lasius niger	GCA_041902855.1
Lhum	Linepithema humile	GCA_040581485.1
Mpha	Monomorium pharaonis	GCA_013373865.2
Nful	Nylanderia fulva	GCA_024268025.1
Obru	Odontomachus brunneus	GCF_010583005.1
Obir	Ooceraea biroi	GCA_003672135.1
Pcal	Pogonomyrmex californicus	GCA_024349325.1
Pbar	Pogonomyrmex barbatus	GCF_000187915.1
Pgra	Pseudomyrmex gracilis	GCA_002006095.1
Sinv	Solenopsis invicta	GCA_016802725.1
Talp	Tetramorium alpestre	GCA_021460005.1
Tbic	Tetramorium bicarinatum	GCA_928718305.1
Tzet	Trachymyrmex zeteki	GCA_001594055.1
Pcla	Paraponera clavata	GCA_048542505.1
Mcam	Mystrium camillae	GCA_048543715.1
Sbic	Stictoponera bicolor	GCA_048543955.1
Lsp	Leptanilla sp.	GCA_048541985.1
Psp	Probolomyrmex sp.	GCA_048541965.1