Context-dependent selection and genetic facilitation and constraint on rosette diameter and

herbivore resistance across European outdoor common gardens under ambient and

reduced precipitation in *Fragaria vesca*

Ivan M. De-la-Cruz[1,8]*, Carolina Diller[1], Femke Batsleer[2], Dries Bonte[2], Timo Hytönen[3], José

Luis Izquierdo[4], Sonia Osorio[5], David Posé[5], Aurora de la Rosa[4], Martijn L. Vandegehuchte[6]§,

Anne Muola[7]§ and Johan A. Stenberg[1]§*

[1]Department of Plant Protection Biology, Swedish University of Agricultural Sciences, Box 190,

SE-23422 Lomma, Sweden

[2]Department of Biology, Terrestrial Ecology Unit, Ghent University, Karel Lodewijk

Ledeganckstraat, 9000 Gent, Belgium

[3]Department of Agricultural Sciences, Viikki Plant Science Centre, University of Helsinki, 0014

Helsinki, Finland

[4]Centro de Investigación, Seguimiento y Evaluación, Parque Nacional Sierra de Guadarrama,

28740 Rascafría, Spain

[5]Instituto de Hortofruticultura Subtropical y Mediterránea La Mayora, Departamento de

Biología Molecular y Bioquímica, Universidad de Málaga-Consejo Superior de Investigaciones

Científicas, Campus de Teatinos, 29750 Algarrobo, Spain

[6]Department of Biology, Norwegian University of Science and Technology, Høgskoleringen 5,

7491 Trondheim, Norway

[7]Division of Biotechnology and Plant Health, Invertebrate Pests and Weeds in Forestry,

Agriculture and Horticulture, Norwegian Institute of Bioeconomy Research, 9016 Tromsø,

Norway

[8]Evolutionary Biology Centre, Department of Ecology and Genetics, Plant Ecology and

Evolution Division, Uppsala University, 752 36 Uppsala, Sweden

***Corresponding authors: Ivan M. De-la-Cruz, ivan.de.la.cruz.arguello@slu.se; Johan A.

Stenberg, johan.stenberg@slu.se.

§Shared senior authorship.

Abstract

The expression of plant defensive traits against herbivores often incurs costs to other essential

functions, such as growth and reproduction. Understanding how selection acts on putatively

functional traits that are expected to trade off across space and time is therefore critical for

predicting evolutionary responses to ongoing and future environmental change. Here, we used

multiple replicated genotypes of woodland strawberry (*Fragaria vesca*; Rosaceae) grown over

two years in three outdoor common gardens in Spain, Belgium, and Sweden. In each garden,

genotypes were exposed to both a reduced-precipitation treatment simulating drought and an

ambient precipitation treatment. We estimated directional and correlational selection on rosette

diameter (a proxy for growth) and herbivore resistance (measured as the inverse of chewing

damage) using fruit and stolon production as proxies for sexual and asexual fitness across all

environments (i.e., every site × year × treatment combination). We then combined selection

gradients with environment-specific genetic (co)variance among genotypes to quantify the

expected response to selection ($\Delta z = G\beta$) and to identify covariance-driven constraints or

facilitation therein. Selection consistently favored larger rosette diameter for both fitness proxies

across nearly all environments, which, in combination with genetic covariances among

genotypes, resulted in a general evolutionary response toward increased rosette diameter, with

the strongest response at the wettest site (Belgium). In contrast, selection on resistance and the

corresponding among-genotype evolutionary responses were strongly context-dependent.

Correlational selection on rosette diameter × resistance occurred in only a few environments,

primarily under reduced precipitation. Environment-dependent genetic covariances constrained

or facilitated selection on both traits only at the site with the highest herbivory (Sweden) under

drought conditions. Overall, our results reveal a context-dependent interplay between selection

and genetic architecture, underlining the difficulty of predicting evolutionary trajectories under

environmental change, and highlighting how spatially and temporally variable conditions may

maintain standing genetic variation in plant traits.

Keywords: growth-defense trade-offs, G-matrix, herbivore resistance, *Fragaria vesca*,

phenotypic selection, genetic constraint and facilitation

Introduction

Plants must often balance their allocation of limited resources between growth and defense against herbivores and pathogens (Coley et al. 1985; Züst and Agrawal 2017; Monson et al. 2022). Growth enhances resource acquisition and competitive ability, whereas defense protects existing tissues through mechanisms such as the production of toxic secondary metabolites that deter, harm or kill herbivores (Züst and Agrawal 2017; Monson et al. 2022). Because nutrients are often limited in nature, investment in defense typically reduces resources available for growth or reproduction, leading to common—though not universal—growth–defense trade-offs due to allocation costs (Coley et al. 1985; Züst and Agrawal 2017; Giolai and Laine 2021; Monson et al. 2022). The trade-offs between growth and defense may also have a genetic basis through pleiotropy or linkage disequilibrium (genetic costs; Züst and Agrawal 2017). Alternatively, simultaneous expression of growth and defense traits may be constrained by environmental conditions (ecological costs; Macel 2011; Züst and Agrawal 2017). For instance, drought or heat stress can suppress plant defenses and alter the cost–benefit balance of plant resistance (Zhao et al. 2025). Thus, trade-offs between defense and growth are often shaped by resource-allocation strategies, genetic correlations, and environmental heterogeneity, which together can generate a mosaic of evolutionary trajectories for these traits across populations (Thompson 2005; He et al. 2022; Monson et al. 2022). However, although growth–defense trade-offs are well documented, comparatively little is known about whether selection gradients align with the underlying genetic (co)variation in these traits, and whether such alignment persists across space, time, and environmental stressors such as drought (but see Züst and Agrawal 2017 and references therein).

Predicting the evolutionary trajectories (response to selection; $G\beta$) of growth and defense requires quantifying selection on each trait and on their combinations, together with the genetic architecture linking them; i.e., evaluating the genetic correlations among defense and growth (Eroukhmanoff and Svensson 2011; Milocco and Salazar-Ciudad 2022; Mallard et al. 2023). These genetic correlations can be detected via the variance–covariance among genotype mean traits (the G matrix) (Agrawal and Stinchcombe 2009; Wise and Rausher 2013; Mallard et al. 2023). Thus, assessing how selection aligns with the G-matrix provides crucial insights into the predicted evolutionary trajectories of growth and defense. For example, even when selection favors both growth and increased defense, a negative genetic covariance (e.g., larger rosettes genetically linked with lower resistance) can counteract these responses and channel evolution along the line of least genetic resistance, thereby slowing down adaptation toward increased growth (i.e., genetic constraint; Coley et al. 1985; Züst and Agrawal 2017). In contrast, covariance aligned with selection can facilitate or increase the multivariate change (Wise and Rausher 2013). There is no universal rule: genetic correlations sometimes constrain adaptation, and other times facilitate it (Wise and Rausher 2013). Moreover, if the G matrix shifts across environments (genotype-by-environment interaction) (Eroukhmanoff and Svensson 2011; Mallard et al. 2023), a tight trade-off in one environment may relax or reverse in another. A genotype's capacity to respond to selection may thus be itself environment-dependent (Eroukhmanoff and Svensson 2011; Agrawal et al. 2012; Mallard et al. 2023). Nevertheless, few studies have tested whether environment-dependent genetic constraints shape selection gradients on growth and resistance when fitness is partitioned into sexual and asexual components, as in many clonal plants.

In many clonal plants, fitness accrues through both sexual reproduction and asexual

propagation, and selection on growth and defense may therefore differ between these

components of fitness (Monson et a. 2022). Yet phenotypic selection analyses that explicitly

quantify selection via both sexual and asexual fitness proxies remain comparatively scarce,

limiting our ability to predict how growth–defense interactions evolve when reproduction occurs

through multiple modes (Huot et al. 2014; He et al. 2022; Monson et a. 2022). Here, we used the

woodland strawberry, *Fragaria vesca* (Rosaceae) to (1) estimate whether directional and/ or

correlational selection acted on plant growth and/or resistance using fruit and stolon production

as proxies for sexual and asexual fitness, respectively, (2) test whether selection aligns with the

among-genotype genetic variance–covariance for both traits; i.e., the predicted evolutionary

response, and (3) evaluate instances in which selection gradients on growth and resistance are

constrained or facilitated across different environments, including a reduced-precipitation

treatment. To this end, we grew 16 wild *F. vesca* genotypes for two years in three outdoor

common gardens located in Spain, Belgium, and Sweden, which vary in environmental

conditions such as temperature, precipitation, and herbivory (Figure 1A,B,C,D,E). The woodland

strawberry genotypes originate from natural populations spanning the species' European

latitudinal range, providing broad standing genetic variation with which to evaluate the

evolutionary response potential in growth and resistance, as well as the underlying genetic

synergies or constraints.

Materials and methods

The woodland strawberry, *Fragaria vesca*, is a perennial plant species that occurs throughout the

Northern Hemisphere in semi-open habitats such as forest clearings and along forest edges

(Hancock 1999). It reproduces both sexually via hermaphroditic flowers and asexually (clonally)

by forming aboveground stolons (Hancock 1999; Schulze et al. 2012; Muola and Stenberg

2018). The main growing season is between March and August, depending on the geographic

location (see Supplementary Table S1).

Three study sites (Fig. 1A) were located along a south–north gradient, spanning the

southern and mid-northern portions of the continental European distribution of *F. vesca*

(Supplementary Table S1). The study sites were central Spain (Rascafría 40°54′17.941″N,

3°52′46.31″W), Belgium (Gontrode 50°59′0.581″N, 3°47′50.248″E), and southern Sweden

(Alnarp 55°38′59.99″N, 13°03′60.00″E). This selection ensured a diverse range of precipitation

regimes, temperatures, and herbivore communities across sites (Figure 1C, D, E and

Supplementary Table S1). The 16 plant genotypes used in this study originated from distinct wild

populations along the continental European south-north gradient (Supplementary Table S2).

Complete details on plant growth conditions (including sowing and propagation of plant

material), maintenance, and experimental setup are provided in De-la-Cruz et al. (2025a,b).

Twenty clonally propagated plantlets per genotype were planted at each site, each in a separate

pot, except for one genotype (GER3), for which only 10 replicate plants per site were available.

Experimental set-up

The plants ($n = 310$ per study site; $n = 930$ in total) were divided into two treatments, reduced

precipitation and control, and exposed to the sites' natural conditions (Figure 1B). At each site,

10 plants from each of 15 of the genotypes, and 5 from genotype GER3, received a reduced

precipitation treatment, while the other 10 plants per genotype (5 for genotype GER3) served as

controls. A split-plot design was used, with the genotype serving as a split-plot factor (one plant

per genotype within a plot; hereafter referred to as a block) and the reduced precipitation

treatment applied to whole plots (blocks; Figure 1B). Pots were spaced 50 cm apart in all

directions in a regular grid formation (Figure 1B). The reduced precipitation treatment consisted

of rainout shelters that reduced the incoming precipitation by 50% (Figure 1B; Yahdjian and Sala

2002). Plants receiving the control treatment were placed in blocks without shelters (Figure 1B).

The entire experimental area was covered with MyPex® weed membrane (Don and Low Ltd) to

prevent weed growth and fenced with fine-mesh chicken wire (also dug into the soil) to prevent

mammalian herbivory. During the summer (June − August), the humidity was measured after 6 −

8 days without rain (or after 3 days without rain if the maximum daily temperatures were +30°C

or above) with a soil moisture meter (Fieldscout TDR 150, Spectrum Technologies, Inc.). If the

average soil moisture of 20 randomly chosen pots (10 per treatment) was lower than 10% of the

volumetric water content (VWC), additional water was supplied: 1 L and 0.5 L to plants under

control and reduced precipitation conditions, respectively. Thus, plants in the reduced

precipitation treatment always received ~50% of the amount of water compared to the control

plants. Soil moisture was also measured during the growing season (June, July and August) in

two different sections of all plant pots during both years. The average soil moisture (%) for the

control and the reduced precipitation treatment across all sites was as follows: control$^{(2021)}$ =

25.44 ± 0.52, reduced precipitation$^{(2021)}$ = 11.56 ± 0.40, control$^{(2022)}$ = 28.40 ±0.48 and reduced

precipitation$^{(2022)}$ = 12.08 ± 0.37 (see also Supplementary Table S1). These data confirmed that

the shelters effectively served as a reduced precipitation treatment (Supplementary Table S1).

Data collection

Leaf chewing damage

The leaf area damaged by chewing herbivores was scored visually for 10 randomly chosen

leaves per plant once in June, July and August of 2021 and 2022. We then calculated the mean

plant damage across the three months (n = 30). Chewing herbivore damage was scored as

proportional leaf area removed and recorded as fractions of a leaflet or, for smaller areas, as

decimal values following these criteria: 1/3 of a leaf (i.e., one leaflet), 1/6 of a leaf (i.e., half a

leaflet), 1/12 or 1/24 of a leaf (i.e., one-quarter or one-eighth of a leaflet), and specific decimal

proportions for smaller areas (0.01, 0.02, or 0.03). To minimize observer bias, participants were

trained during bespoke workshops to assess chewing damage on woodland strawberry leaves

until estimates converged across participants. In addition, each observer was trained by using the

web application https://zaxherbivorytrainer.com to estimate the amount of leaf damage on

randomized images of leaves. Estimates within $0 - 5\%$ of the actual answer were considered

accurate. See Figure 1E for the mean leaf damage values per site across treatments and years.

<u>Rosette diameter and reproductive traits</u>

For each plant, rosette diameter was used as a proxy for vegetative growth and measured

horizontally with a ruler (cm). Proxies of sexual and asexual fitness were estimated as the

number of ripe fruits and stolons (Muola and Stenberg 2017; De-la-Cruz et al. 2025a),

respectively. Rosette diameter and counts of fruits (including those fallen into the soil pots) and

stolons were recorded monthly in June, July, and August of each study year (2021 and 2022).

Only fruits without visible herbivore damage or pathogen infection were included, as such

damage can reduce seed number and quality and thereby bias fitness estimates. Fruits and stolons

were systematically removed after counting to prevent double counting and, in the case of

stolons, to avoid growth into neighboring pots. Fruit removal is unlikely to induce compensatory

responses (Hilty et al. 2021), as ripe fruits naturally detach from the plant over time. Rosette

diameter was averaged across the three sampling times within each year for downstream

analyses (see below). Fruit and stolon counts were summed across the three sampling times

within each year and used as annual proxies of plant fitness for subsequent analyses (see below).

<u>Statistical analysis</u>

<u>Phenotypic selection on resistance and rosette diameter across environments</u>

All analyses were conducted in R v4.4.0 (R Core Team 2021). We used the inverse of plant

damage as a proxy for plant resistance (calculated using the operational definition 1 – mean

proportion of total leaf area damaged by chewing herbivores; Simms and Rausher 1987; Núñez-

Farfán and Dirzo 1994). This operational resistance measurement reflects only realized chewing

damage (Simms and Rausher 1987; Núñez-Farfán and Dirzo 1994). However, because

herbivores often preferentially attack larger, healthier plants that offer greater biomass and

nutrient availability (i.e., increased apparency; Price 1991; Hella Schlinkert et al. 2015), and

because plant resistance may align with a latitudinal cline (Woods et al. 2012), we adjusted our

measure of plant resistance by using a generalized linear mixed model (GLMM) to obtain size-

and latitude-independent estimates of herbivore resistance within each environment (i.e., each

site × treatment × year combination). To this end, for each environment, we fitted a GLMM (beta

error distribution with a logit link), using proportional plant damage as the response variable and

rosette diameter and latitude as fixed effects or predictors. Genotype and block were included as

random effects to account for genetic and spatial replication. From each model, we extracted

environment-specific slopes for rosette diameter ($b_{diameter}$) and latitude ($b_{latitude}$), quantifying the

effects of plant size and geographic origin on herbivore damage. We then calculated individual-

level adjusted damage by subtracting the predicted contributions of rosette diameter and latitude

from the observed logit-transformed damage: Adjusted damage = logit($Damage_{observed}$) − ($b_{diameter}$

× rosette diameter) − ($b_{latitude}$ × latitude). This value represents the expected damage a plant

would experience if it had the mean rosette diameter and latitude of its origin (Stinchcombe et al.

2008). Adjusted damage was back-transformed using the inverse logit function, and adjusted

resistance was defined as $1 -$ adjusted damage. This metric enabled selection analyses on

herbivore resistance that are unbiased by predictable variation in plant apparency and geographic

origin (Price 1991; Hella Schlinkert et al. 2015). Likewise, to remove potential geographic clines

in growth (as observed by De-la-Cruz et al. 2025a in the same dataset analyzed here), rosette

diameter was residualized against latitude within each environment using a mixed-effects model

with genotype and block included as random effects, yielding a latitude-independent estimate of

plant size (Woods et al. 2012).

We then performed proxies of directional and correlational phenotypic selection analyses

on resistance and rosette diameter (adjusted; see above) using two different GLMMs, one for

each fitness measures (i.e. for fruit and stolon production). In each model, the response variable

was fruit or stolon production. To estimate selection on *relative* fitness proxies within each

environment (i.e., every site × year × treatment combination), we computed the environment-

specific mean for fruit and stolon production and used its natural logarithm as an offset in the

two models. The GLMMs were fit using a negative binomial distribution with a log link

(variance function "nbinom2") using glmmTMB v1.1.13 (Brooks et al. 2017; McGillycuddy et

al. 2025). The structure of the models was specified as follows:

Fruits/stolons = Offset (Log (Mean fruits/stolons)) + (resistance + rosette diameter + resistance ×

rosette diameter) * site * year * treatment.

Resistance and rosette diameter values were standardized to mean 0 and standard

deviation of 1 within each environment (i.e., site × treatment × year combination). In these

models the independent effect (or additive effect) of resistance or rosette corresponds to

directional selection gradient proxies, while the interaction (synergistic/antagonistic effects)

between resistance and rosette corresponds to the correlational selection proxies. We included the

block, genotype × site, genotype × year, genotype × treatment, genotype × site × year, genotype

× year × treatment, genotype × site × treatment, and genotype × site × year × treatment as

random effects. Incorporating the random effects into these GLMMs allowed each genotype to

have its own baseline shift in fitness in each environment.

Because the model uses a log link and an offset, coefficients on the latent (log) scale

correspond to multiplicative effects on expected fruit or stolon production relative to the

environment mean (McCullagh and Nelder 1989; de Villemereuil et al. 2016). Thus, we used

emtrends v1.3.5.1 to obtain, for each environment, the marginal slopes ("trends") for resistance,

rosette diameter, and their interaction on the latent scale, along with standard errors and Wald

tests. We back-transformed the latent-scale gradients to the raw scale by using: $\beta raw = e^{\beta} - 1$

(McCullagh and Nelder 1989). Standard errors and 95% confidence intervals were also

calculated for the raw scale selection gradients. The generalized linear coefficients (i.e., the

selection gradients; βi, Lande and Arnold 1983) obtained from the models represent the

magnitude and direction of selection acting on resistance, rosette, resistance × rosette in

comparable units (standard deviations) across years, sites, and treatments (Lande and Arnold

1983; Wise and Rauser 2013).

<u>Among-genotype variance–covariance matrix per environment</u>

For each environment (every site × year × treatment combination), we performed a two-response

bivariate bayesian GLMM in the R package brms v2.23.0 (Bürkner 2017) with the cmdstanr

v0.9.0 interface (Gabry et al. 2025). Resistance was modeled with a Beta distribution and logit

link as follows: Resistance ~ Block + (1 | Genotype). Rosette diameter was modeled assuming a

Gaussian distribution and identity link as follows: Rosette diameter ~ Block + (1 | Genotype).

Block was included as a fixed effect for both traits to control for micro-environmental

heterogeneity/structure variance within sites. Genotype as random effect for the multivariate

GLMM allowed the model to estimate the among-genotype covariance between traits. Residual

cross-trait correlation was disabled, ensuring that cross-trait association is attributed to the

genotype effects rather than to residual error (de Villemereuil et al. 2016). For each environment-

specific model, we ran four chains, with 8,000 iterations per chain, warm-up to 25% of the total,

adapt_delta = 0.95 and max_treedepth = 15 and fixed random seed of 1001. For each fitted

environment-specific model, posterior draws (each draw is one possible 2×2 variance-

covariance matrix from the MCMC bayesian GLMM; ~288,000 draws in total across all

environments) were converted to a latent-scale G-matrix (G_{lat}); a 2×2 genotype-level variance–

covariance matrix G on the latent scale (logit scale for resistance; identity for rosette diameter).

We converted the latent covariance and variances matrices (G_{lat}) to observed scale (i.e., the

natural measurement scale of each trait; G_{obs}) using *Qgmvparams* function of QGglmm v0.8.0 R

package (de Villemereuil et al. 2016). Thus, we obtained an observed-scale genetic (co)variance

matrices (G_{obs}) per environment, consistent with the link functions and distributional assumptions

of the two traits (de Villemereuil et al. 2016).

The observed G matrices (G_{obs}) were further standardized to enable comparisons across

environments. For each environment, we computed the phenotypic standard deviation vector and

formed a diagonal scaling matrix (Khachiyan and Kalantari 1990). The observed-scale matrices

(G_{obs}) were then standardized as $G_{st} = D^{-1}G_{obs}D^{-1}$ (Khachiyan and Kalantari 1990), where D^{-1} is

the diagonal scaling matrix for resistance and rosette diameter for each environment and G_{obs} is

the observed G matrix for that environment (see above). The G_{st} matrices were summarized by

calculating the mean among-genotype variance-covariance matrix ($G_{st\text{-}mean}$) of resistance and

rosette diameter together with their 95% credible intervals for each environment. Diagonal

elements quantify genetic variance for each trait; the off-diagonal element reflects the sign and

magnitude of the genetic covariance between resistance and rosette.

<u>Evolutionary response potential</u>

We estimated the among-genotype evolutionary response potential for each environment using

Lande's equation, $\Delta z = G\beta$ (Lande 1979; Milocco and Salazar-Ciudad 1999), where G is the 2 ×

2 environment-specific among-genotype variance–covariance matrix (G_{st}; see above) and β is the

vector of directional selection gradients on resistance and rosette size for that environment. From

the Lande's equation, we retained both components: Δresistance and Δrosette diameter (the

among genotype evolutionary response for resistance and rosette diameter for each

environment). Δresistance and Δrosette diameter are expressed in phenotypic standard deviation

units. We stress that we interpret $\Delta z = G\beta$ as the among-genotype evolutionary response potential

in every environment rather than population-specific evolutionary predictions (Milocco

and Salazar-Ciudad 1999). In other words, Δz describes the expected change in the trait mean if

selection acted on differences among the genotypes studied within a given environment (every

site × treatment × year combination). These values are not forecasts of population-level

evolutionary trajectories, as they do not account for genotype/allele frequencies or other

population-specific demographic processes (Milocco and Salazar-Ciudad 1999). Positive values

indicate a comparative (among-genotype) predicted increase in resistance or rosette diameter

mean, while negative values indicate a decrease.

<u>Facilitation and genetic constraints</u>

We estimated facilitation and genetic constraints following the approach of Agrawal and

Stinchcombe (2009) and Wise and Rausher (2013). From the bayesian GLMMs, we extracted the

posterior draws (posterior G_{st} matrices) for each environment (see above). We obtained the

directional selection gradients (β) for resistance and rosette diameter per environment from the

negative binomial GLMM analyses that we described above. For each posterior draw, we again

calculated the $\Delta z = G \times \beta$ under two scenarios: 1) Full G matrix or Response Vector$_1$: RV$_1$ = $G \times$

β, and 2) Diagonalised G matrix (G_0, covariances set to zero) or Response Vector$_0$: RV$_0$ =

diag(G) $\times \beta$ (Wise and Rausher 2013). Thus, RV$_1$ estimates the among-genotype evolutionary

response for resistance and rosette diameter considering the genetic covariances among them for

each posterior draw, whereas RV$_0$ estimates what the evolutionary response would have been if

there had been no genetic covariances among them (Smith and Rausher 2008; Agrawal and

Stinchcombe 2009). We then quantified the extent to which the predicted response of resistance

or rosette size changed when genetic covariances were included (Wise and Rausher 2013). To do

so, we compared the predicted change under the full G scenario with that obtained under the no-

covariance scenario, as follows: Trait difference (TDiff) = RV$_1$ trait − RV$_0$ trait (Wise and

Rausher 2013). We then interpreted the results as "facilitation" or "constraint" based on whether

covariances increased or decreased the evolutionary response in the direction favored by

selection, rather than relying solely on the raw sign of TDiff. We therefore defined a direction-

aware covariance effect for each trait by projecting the TDiff onto the corresponding selection

gradient, as follows: Facilitation/Constraint (FC) = TDiff $\times \beta$. Under this definition, genetic

covariances were classified as facilitating evolutionary change for resistance or rosette diameter

when FC > 0, indicating that covariances increase the magnitude of the predicted response in the

direction of selection (Agrawal and Stinchcombe 2009; Wise and Rausher 2013). Conversely,

covariances were classified as constraining when FC < 0, indicating that covariances

reduce/constraint the predicted response in the direction of selection (Agrawal and Stinchcombe

2009; Wise and Rausher 2013). In other words, we asked if the predicted phenotypic selection

was aligned or misaligned with the structure (e.g., antagonistic or synergistic covariance between

traits) of the genetic matrix (Wise and Rausher 2013). For each environment, we computed the

proportion of draws where each trait experienced facilitation or constraint. An effect was

considered significant when ≥ 90% of the posterior draws supported either facilitation or

constraint (Lambert 2018).

Results

Phenotypic selection on resistance

Significant phenotypic selection on resistance was detected only in Sweden and Spain. In

Sweden, under control conditions in 2021, plants with higher resistance produced more fruits

(Fig. 2A, Supplementary Table S3). In contrast, in Sweden in 2022 under reduced precipitation,

plants with lower resistance had higher fruit production (Figure 2A, Supplementary Table S3).

We also found that plants with higher resistance consistently produced fewer stolons in Sweden

across both years and treatments and in Spain (only in 2022 under reduced precipitation) (Figure

2B, Supplementary Table S3).

Phenotypic selection on rosette diameter

For fruit production, gradients of selection on rosette diameter were positive in 11 of the 12

environments (i.e., in almost every site × treatment × year combination; Figure 2AB,

Supplementary Table S3). This pattern suggests that plants with larger rosette diameter had more

fruits across sites, treatments, and years (Figure 2A, Supplementary Table S3). The exception

occurred in Spain in 2022 under reduced-precipitation conditions, where plants with higher fruit

production had smaller rosette diameter (Figure 2A, Supplementary Table S3). In terms of stolon

production, selection on rosette diameter showed positive gradients in nearly all environments

(almost all significant too), indicating that larger plants had more stolons across sites, treatments,

and years (Figure 2B, Supplementary Table S3).

<u>Among-genotype evolutionary response potential for resistance</u>

Among-genotype evolutionary response for resistance, using fruit production as the fitness

proxy, showed a predicted response for higher resistance in Sweden in 2021 under reduced

precipitation (Figure 3A). However, in 2022, a predicted response for lower resistance was

detected in Sweden under reduced precipitation (Figure 3A), and the same pattern was observed

when stolon production was used as a fitness proxy at the same site, year, and treatment (Figure

3B).

<u>Among-genotype evolutionary response potential for rosette diameter</u>

We detected a consistent among-genotype evolutionary response favoring larger rosette diameter

across all sites, treatments, and years, using both fitness proxies (Figure 3C,D). The strongest

evolutionary response for rosette diameter (both fitness measures) was observed in Belgium

across both treatments and years (Figure 3C,D). The exception was found in Spain in 2022 under

reduced precipitation using fruit production as fitness measure, where a decrease in rosette

diameter was predicted (Figure 3C,D).

<u>Selection on resistance × rosette diameter interaction</u>

The GLMM for fruit production revealed that in Sweden in 2022 under reduced-precipitation,

plants with higher resistance and larger rosettes produced more fruits (Supplementary Figure

S1A, Supplementary Table S3). In contrast, in Spain in 2022 under reduced precipitation, plants

with lower resistance and smaller rosettes had higher fruit production (Supplementary Figure

S1A, Supplementary Table S3). In Belgium, under reduced precipitation, plants with larger

rosette diameter and lower resistance produced more fruits in 2021, and the same pattern was

observed in 2022, although only marginally significant (Supplementary Figure S1A,

Supplementary Table S3). Significant correlational selection on stolon production was only

detected in Spain in 2022 under reduced precipitation, indicating that plants exhibiting both

higher resistance and larger rosette diameter produced more stolons (Supplementary Figure S1B,

Supplementary Table S3).

<u>Genetic facilitations and constraints</u>

Using fruit production as fitness proxy, we found that in Sweden under reduced precipitation in

2021, the genetic covariances facilitated the selection for an increased resistance and rosette

diameter (Figure 4A, Supplementary Table S4). However, in 2022, in Sweden under reduced

precipitation, the covariances facilitated selection for reduced resistance (i.e., intensified

selection against resistance) and also facilitated selection for larger rosettes (Figure 4A,

Supplementary Table S4).

Using stolon production as a fitness proxy, genetic covariances in 2021 in Sweden under

reduced precipitation constrained selection favoring resistance and also constrained selection for

reduced rosette diameter (i.e., slowed selection for smaller rosettes; Figure 4B, Supplementary

Table S4). In contrast, in 2022 in Sweden under reduced precipitation, covariances facilitated

selection for reduced resistance and facilitated selection for larger rosette diameter (Figure 4B).

Discussion

This study reveals how environmental heterogeneity interacts with genetic architecture to shape among-genotype evolutionary trajectories of resistance and growth across different environments spanning much of the European range of *F. vesca*.

Across nearly all conditions examined (year × site × treatment combinations), plants with larger rosette diameters consistently showed higher fruit and stolon production. Because genotypes differed in rosette diameter, this pattern implies that, if heritable, mean rosette size would be expected to increase. The strongest predicted among-genotype evolutionary response for larger rosettes occurred in Belgium, likely due to consistently high precipitation across both years, which may have increased resource availability for growth and strengthened selection for larger plants (Hilty et al. 2021). Larger rosette diameters may confer higher fitness through greater photosynthetic capacity, improved resource acquisition, and enhanced competitive ability (Poorter et al. 2012; Ainsworth and Bush 2011; Hilty et al. 2021). An exception was found in Spain under the reduced-precipitation treatment in 2022, where smaller rosettes had higher fruit production. Spain was the driest site in 2022, hence plants may have experienced a more severe drought stress at that site. Under intense water limitation, large rosettes may incur high metabolic costs that reduce fruit production (Eziz et al. 2017). Under such conditions, reduced growth can function as a strategy to limit transpiration and metabolic demand (Eziz et al. 2017; Takahashi et al. 2020).

In contrast to the consistent selection for larger rosette diameter, selection on herbivore resistance was absent or non-significant when either stolon or fruit production was used as a fitness proxy. Likewise, the predicted among-genotype evolutionary response for resistance generally resulted in a non-significant change, except in Sweden under reduced precipitation in

2021, where resistance was predicted to increase. Plants in Sweden experienced higher chewing-

herbivore damage than at the other sites in both years and treatments. Under control conditions in

2021 in Sweden, higher resistance was associated with increased fruit production, and we

detected a marginally significant positive selection gradient under reduced precipitation,

suggesting that a higher resistance conferred higher fitness when herbivore pressure was elevated

in 2021. However, in 2022, selection on resistance was not significant under control conditions,

whereas under reduced precipitation, higher resistance was associated with lower fruit

production. We also detected a predicted evolutionary response against resistance (a predicted

shift toward lower resistance) in Sweden in 2022 under reduced precipitation using both fitness

proxies. It is possible that the metabolic costs of maintaining defensive traits could outweigh

their protective benefits when water/resources are limited (Coley et al. 1985; Züst and Agrawal

2017; Monson et al. 2021). In other words, producing defenses under reduced precipitation

when herbivory was high appeared to impose fitness costs. Moreover, we hypothesize that

increasing plant size, under both treatments, provided greater benefits for fruit and stolon

production than allocating in defense at the herbivory levels observed in Sweden. At these

damage levels (~10% leaf loss), resistance may be more costly than beneficial, particularly under

reduced precipitation, and plants may instead cope with herbivory through compensatory growth

or tolerance (Nuñez-Farfán et al. 2007; Muola et al. 2010; Züst and Agrawal 2017; Blumenthal et

al. 2020), a possibility that needs further study.

Phenotypic selection analyses using stolon production as the fitness proxy further showed

that plants with higher resistance produced fewer stolons especially in Sweden across all

treatments and years and in Spain in 2022 under reduced precipitation. Using stolon production

as a fitness measure adds insights into how resistance was selected against and stresses the

importance of allocating resources to growth and clonal propagation rather than to resistance

traits. Our findings appear to align with the resource-allocation theory, which suggests that

selection for rapid reproduction and larger size can favor fast-growing but weakly defended

genotypes (Coley et al. 1985; Monson et al. 2021).

Correlational selection on resistance × rosette diameter (for both fitness proxies) was

largely absent in all environments, with four exceptions. In Sweden in 2022 under reduced

precipitation, plants with both higher resistance and larger rosettes produced more fruits,

suggesting a possible synergistic benefit of investing in growth and defense when water is

limited, and herbivory is high (Li et al. 2024). There is empirical evidence from other study

systems that under intense herbivory, plants with both higher growth and higher resistance show

higher fitness (Carmona and Fornoni 2013; De-la-Cruz et al. 2023). However, it is important to

recall that we also found directional selection for reduced resistance in this environment (see

above). Thus, we hypothesize that resistance appears beneficial only when accompanied by an

allocation for a larger rosette diameter. In other words, we hypothesize that increasing resistance

alone is costly—especially under water limitation—unless plants can also allocate resources to

growth (Carmona and Fornoni 2013).

In Spain in 2022 under reduced precipitation, correlational selection pointed in opposite

directions depending on the fitness proxy. When fruit production was used, the favored trait

combination consisted of smaller rosettes and lower resistance; when stolon production was

used, the favored combination consisted of larger rosettes and higher resistance. In other words,

plants investing more in growth and defense appeared less successful sexually but more

successful clonally. These results suggest a trade-off between reproductive modes at the site

where the most severe drought and high temperatures were observed. It is possible that larger,

better-defended plants allocated all their resources to stolon production, a strategy that may

enhance persistence under extreme conditions (Vallejo-Marín et al. 2010; Orive et al. 2017;

Blumenthal et al. 2020). In Belgium under reduced precipitation, plants with larger rosettes and

lower resistance produced more fruits in 2021, with a similar but marginally significant pattern in

2022. These results from Belgium reinforce that reduced precipitation can promote selective

trade-offs (see above). In summary, correlational selection on growth and resistance is not fixed

but shaped by reproductive modes and/or ecological context, including spatial variation in

herbivory and precipitation.

Genetic correlations can either enhance or constrain the evolutionary adaptive potential,

with positive correlations among traits under positive selection accelerating adaptation and

negative correlations inhibiting it (Agrawal and Stinchcombe 2009). We detected both genetic

constraints and facilitation only in Sweden, and the direction and magnitude of these effects were

strongly dependent on environmental context. For fruit production, in Sweden under reduced

precipitation in 2021, genetic covariances facilitated the selection for increased resistance and

larger rosette diameter, indicating that the G-matrix (e.g., via pleiotropy, linkage disequilibrium;

Eroukhmanoff and Svensson 2011; Svensson 2023) may boost their joint adaptive evolution.

When stolon production was used as the fitness proxy in Sweden in 2021 under reduced

precipitation, genetic covariances constrained evolution for resistance and toward smaller

rosettes. Larger rosettes may enable greater allocation to stolon production than smaller rosettes

(Orive et al. 2017), and clonality can distribute risk among ramets, promote division of labor, and

enhance persistence in stressful environments (Vallejo-Marín et al. 2010; Orive et al. 2017).

Thus, plants may have allocated more resources to vegetative growth (thereby reducing

evolutionary change toward smaller rosettes), which in the first year could have increased stolon

production to persist under drier conditions and herbivory. At the same site in 2022 under

reduced precipitation, we found that selection would tend to decrease resistance (facilitation for a

reduced resistance), while simultaneously facilitating an increase in rosette diameter. This

temporal reversal in the evolutionary responses of resistance highlights how the evolutionary

consequences of genetic architecture can shift rapidly with changing selection regimes

(genotype-by-environment interactions; Eroukhmanoff and Svensson 2011). These results also

show that the reproductive mode plays an important role in shaping the evolution of these traits

and highlights the complexity plants face when coping with multiple stressors while allocating

resources to both reproductive modes, growth, and resistance (Züst and Agrawal 2017).

In summary, the genetic constraints and facilitations identified only at the site with the

highest herbivory (Sweden), together with the context-dependent nature of selection, highlight

the difficulty of predicting among-genotype potential responses to climate change without

integrating both environmental variation (biotic and abiotic factors) and the genetic architecture.

Acknowledgements

We thank Juan Antonio Vielva Juez, Director of the Centro de Investigación, Seguimiento y

Evaluación del Parque Nacional de la Sierra de Guadarrama, Madrid, Spain for hosting the

southernmost (Rascafría/Spain) study site and for logistic support during the experiment. We

thank Hans Matheve for helping with fieldwork and data digitalization and Sanne de Jong,

Matilda Jützeler and several other assistant researchers who helped us to collect the data during

fieldwork. We thanks to UPPMAX (Uppsala Multidisciplinary Center for Advanced

Computational Science) of Uppsala University, for providing us access to their computational

resources for the Bayesian analyses.

Author contributions

The study was conceived by IMDC, MLV, AM, JAS, DB, TH, SO, DP. The experiments were

performed by IMDC, CD, FB, MLV, ADL, JLI, AM and JAS. The data were analyzed by IMDC.

The draft manuscript was written by IMDC with considerable input and revisions from all the

authors.

Conflict of interests

The authors declare that they have no conflicts of interest.

Open Research

Data supporting this manuscript, and all the scripts/workflow will be deposited in Figshare or

Zenodo upon manuscript acceptance.

Figure legends

Figure 1. (A) The location of the three experimental sites [orange triangles, from south to mid-

and northern Europe: Spain (Rascafría), Belgium (Gontrode), and southern Sweden (Alnarp)]

and the origin of each of the 16 genotypes used in the study (black circles; for genotype labels

and their latitude and longitude of origin, see Supplementary Table S2). (B) The experimental

design for each study site. Each colour represents one of the 16 genotypes. Rainout shelters (blue

shaded areas) excluded 50% of the incoming precipitation. The blue arrows indicate the flow

direction of the rainwater through the drainage pipes. (C, D) Mean temperature (C) and

precipitation (D) and their standard errors in the period April–August at each site (see also

Supplementary Table S1). (E) Mean plant chewing damage and its error bars are shown for each

site under the control and reduced-precipitation treatments for 2021 and 2022. Credit for picture

1B: Martijn L. Vandegehuchte.

Figure 2. (A) Selection gradients (β) for fruit production across the study sites, Spain, Belgium,

and Sweden, under control and reduced-precipitation treatments in 2021 and 2022 (site ×

treatment × year combinations). Purple bars denote selection on chewing herbivore resistance,

and teal bars denote selection on rosette diameter. 95% confidence intervals indicate uncertainty

around the selection gradients. Asterisks indicate statistical significance ($p \leq 0.05$). (B) Selection

gradients (β) for stolon production, with colors and annotations as in panel A.

Figure 3. Panels (A–D) show means and 95% confidence intervals (CIs) of the among-genotype

predicted evolutionary response to selection ($\Delta z = G\beta$) under control and reduced-precipitation

treatments in 2021 and 2022 across the three study sites: Spain, Belgium, and Sweden. (A)

Predicted evolutionary responses (Δz) of herbivore resistance based on selection via fruit

production. (B) Predicted evolutionary responses (Δz) of herbivore resistance based on selection

via stolon production. (C) Predicted evolutionary responses (Δz) of rosette diameter based on

selection via fruit production. (D) Predicted evolutionary responses (Δz) of rosette diameter

based on selection via stolon production. Colors indicate whether each mean Δz ($\pm$95% CI) is

positive (purple), negative (orange), or non-significant (grey). The vertical dashed line denotes

$\Delta z = 0$.

Figure 4. Panels (A) and (B) illustrate per-trait genetic constraints and facilitation, quantified as

the difference between predicted evolutionary responses calculated using the full genetic

variance–covariance matrix (G) and those calculated with covariances removed (G0). Thus,

Difference = Response with G – Response with G0 (see also methods). (A) Median differences

and 95% confidence intervals (CIs) based on selection via fruit production across precipitation

treatments (control *vs*. reduced precipitation) for each focal trait (herbivore resistance and rosette

diameter) in 2021 and 2022 across Spain, Belgium, and Sweden. (B) Median differences and

95% CIs based on selection via stolon production across precipitation treatments and focal traits

for 2021 and 2022 across the same sites. The vertical dashed line indicates no effect of genetic

covariances (Difference = 0). Colors indicate whether covariances constrain the selection

response (blue), facilitate the selection response (green), or have no detectable effect (grey). See

also methods and Supplementary Table S4.

References

Agrawal, A. A., Hastings, A. P., Johnson, M. T. J., Maron, J. L., and Salminen, J.-P. (2012). Insect herbivores drive real-time

ecological and evolutionary change in plant populations. Science, 338(6103), 113–116.

https://doi.org/10.1126/science.1225977

Agrawal, A. F., and Stinchcombe, J. R. (2009). How much do genetic covariances alter the rate of adaptation? Proceedings.

Biological Sciences, 276(1659), 1183–1191. https://doi.org/10.1098/rspb.2008.1671

Ainsworth, E. A., and Bush, D. R. (2011). Carbohydrate export from the leaf: A highly regulated process and target to enhance

photosynthesis and productivity. Plant Physiology, 155(1), 64–69. https://doi.org/10.1104/pp.110.167684

Blumenthal, D. M., Mueller, K. E., Kray, J. A., Ocheltree, T. W., Augustine, D. J., and Wilcox, K. R. (2020). Traits link drought

resistance with herbivore defence and plant economics in semi-arid grasslands: The central roles of phenology and leaf

dry matter content. Journal of Ecology, 108(6), 2336–2351. https://doi.org/10.1111/1365-2745.13454

Brooks, M. E., Kristensen, K., Benthem, K. J. van, Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Mächler, M., and

Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear

mixed modeling. The R Journal, 9(2), 378–400. https://doi.org/10.32614/RJ-2017-066

Bürkner, P.-C. (2017). brms: An R package for bayesian multilevel models using Stan. Journal of Statistical Software, 80, 1–28.

https://doi.org/10.18637/jss.v080.i01

Carmona, D., and Fornoni, J. (2013). Herbivores can select for mixed defensive strategies in plants. New Phytologist, 197(2),

576–585. https://doi.org/10.1111/nph.12023

Coley, P. D., Bryant, J. P., and Chapin, F. S. (1985). Resource availability and plant antiherbivore defense. Science, 230(4728), 895–899. https://doi.org/10.1126/science.230.4728.895

de Villemereuil, P., Schielzeth, H., Nakagawa, S., and Morrissey, M. (2016a). General methods for evolutionary quantitative genetic inference from generalized mixed models. Genetics, 204(3), 1281–1294. https://doi.org/10.1534/genetics.115.186536

De-la-Cruz, I. M., Batsleer, F., Bonte, D., Diller, C., Hytönen, T., Luis Izquierdo, J., Osorio, S., Posé, D., de la Rosa, A., Vandegehuchte, M. L., Muola, A., and Stenberg, J. A. (2025a). Genotypic responses to different environments and reduced precipitation reveal signals of local adaptation and phenotypic plasticity in woodland strawberry. Annals of Botany, 136(3), 611–621. https://doi.org/10.1093/aob/mcaf025

De-la-Cruz, I. M., Batsleer, F., Bonte, D., Diller, C., Izquierdo, J. L., Still, S., Osorio, S., Posé, D., de la Rosa, A., Vandegehuchte, M. L., Muola, A., Hytönen, T., and Stenberg, J. A. (2025b). Flowering responses of the woodland strawberry to local climate and reduced precipitation along a European latitudinal gradient. Journal of Plant Ecology, 18(5), rtaf105. https://doi.org/10.1093/jpe/rtaf105

De-la-Cruz, I. M., and Núñez-Farfán, J. (2023). Inter-annual variation in the abundance of specialist herbivores determines plant resistance in *Datura stramonium*. Ecology and Evolution, 13(12), e10794. https://doi.org/10.1002/ece3.10794

Eroukhmanoff, F., and Svensson, E. I. (2011). Evolution and stability of the G-matrix during the colonization of a novel environment. Journal of Evolutionary Biology, 24(6), 1363–1373. https://doi.org/10.1111/j.1420-9101.2011.02270.x

Eziz, A., Yan, Z., Tian, D., Han, W., Tang, Z., and Fang, J. (2017). Drought effect on plant biomass allocation: A meta-analysis. Ecology and Evolution, 7(24), 11002–11010. https://doi.org/10.1002/ece3.3630

Gabry, J., Češnovar, R., Johnson, A., and Bronder, S. (2025). *cmdstanr: R Interface to 'CmdStan'*. R package version 0.9.0, https://discourse.mc-stan.org, https://mc-stan.org/cmdstanr/

Giolai, M., and Laine, A.-L. (2024). A trade-off between investment in molecular defense repertoires and growth in plants. Science, 386(6722), 677–680. https://doi.org/10.1126/science.adn2779

Hancock, J. F. (1999). Strawberries. CABI Pub.

He, Z., Webster, S., and He, S. Y. (2022). Growth-defense trade-offs in plants. Current Biology: CB, 32(12), R634–R639. https://doi.org/10.1016/j.cub.2022.04.070

Hilty, J., Muller, B., Pantin, F., and Leuzinger, S. (2021). Plant growth: the what, the how, and the why. New Phytologist, 232(1), 25–41. https://doi.org/10.1111/nph.17610

Huot, B., Yao, J., Montgomery, B. L., and He, S. Y. (2014). Growth-defense tradeoffs in plants: A balancing act to optimize fitness. Molecular Plant, 7(8), 1267–1287. https://doi.org/10.1093/mp/ssu049

627   Khachiyan, L., and Kalantari, B. 1992. Diagonal matrix scaling and linear programming. SIAM Journal on Optimization, 2(4).

628   https://epubs.siam.org/doi/10.1137/0802034

629   Lambert, B. (2018). A Student's guide to bayesian statistics. SAGE Publications Ltd. https://doi.org/10.4135/9781036234546

630   Lande, R. (1979). Quantitative genetic analysis of multivariate evolution, applied to brain:body size allometry. Evolution,

631   33(1Part2), 402–416. https://doi.org/10.1111/j.1558-5646.1979.tb04694.x

632   Lande, R., and Arnold, S. J. (1983). The measurement of selection on correlated characters. Evolution, 37(6), 1210–1226.

633   https://doi.org/10.2307/2408842

634   Li, Z.-X., Tan, J.-F., Yao, N., and Xie, R.-H. (2024). From trade-off to synergy: How nutrient status modulates plant resistance to

635   herbivorous insects? Advanced Biotechnology, 2(4), 37. https://doi.org/10.1007/s44307-024-00045-5

636   Macel, M. (2011). Attract and deter: A dual role for pyrrolizidine alkaloids in plant–insect interactions. Phytochemistry Reviews,

637   10(1), 75–82. https://doi.org/10.1007/s11101-010-9181-1

638   Mallard, F., Afonso, B., and Teotónio, H. (2023). Selection and the direction of phenotypic evolution. eLife, 12, e80993.

639   https://doi.org/10.7554/eLife.80993

640   McCullagh, P. (2019). Generalized linear models (2nd ed.). Routledge. https://doi.org/10.1201/9780203753736

641   McGillycuddy, M., Popovic, G., Bolker, B. M., and Warton, D. I. (2025). Parsimoniously fitting large multivariate random effects

642   in glmmTMB. Journal of Statistical Software, 112, 1–19. https://doi.org/10.18637/jss.v112.i01

643   Milocco, L., and Salazar-Ciudad, I. (2022). Evolution of the g matrix under nonlinear genotype-phenotype maps. The American

644   Naturalist, 199(3), 420–435. https://doi.org/10.1086/717814

645   Monson, R. K., Trowbridge, A. M., Lindroth, R. L., and Lerdau, M. T. (2022). Coordinated resource allocation to plant growth–

646   defense tradeoffs. New Phytologist, 233(3), 1051–1066. https://doi.org/10.1111/nph.17773

647   Muola, A., Mutikainen, P., Laukkanen, L., Lilley, M., and Leimu, R. (2010). Genetic variation in herbivore resistance and

648   tolerance: The role of plant life-history stage and type of damage. Journal of Evolutionary Biology, 23(10), 2185–2196.

649   https://doi.org/10.1111/j.1420-9101.2010.02077.x

650   Muola, A., and Stenberg, J. A. (2018). Folivory has long-term effects on sexual but not on asexual reproduction in woodland

651   strawberry. Ecology and Evolution, 8(23), 12250–12259. https://doi.org/10.1002/ece3.4687

652   Núñez-Farfán, J., and Dirzo, R. (1994). Evolutionary ecology of *Datura stramonium* l. in Central Mexico: natural selection for

653   resistance to herbivorous insects. Evolution, 48(2), 423–436. https://doi.org/10.1111/j.1558-5646.1994.tb01321.x

654   Núñez-Farfán, J., Fornoni, J., and Valverde, P. L. (2007). The evolution of resistance and tolerance to herbivores. Annual Review

655   of Ecology, Evolution, and Systematics, 38(Volume 38, 2007), 541–566.

656   https://doi.org/10.1146/annurev.ecolsys.38.091206.095822

Orive, M. E., Barfield, M., Fernandez, C., and Holt, R. D. (2017). Effects of clonal reproduction on evolutionary lag and

evolutionary rescue. The American Naturalist, 190(4), 469–490. https://doi.org/10.1086/693006

Poorter, H., Niklas, K. J., Reich, P. B., Oleksyn, J., Poot, P., and Mommer, L. (2012). Biomass allocation to leaves, stems and

roots: Meta-analyses of interspecific variation and environmental control. New Phytologist, 193(1), 30–50.

https://doi.org/10.1111/j.1469-8137.2011.03952.x

Price, P. W. (1991). The plant vigor hypothesis and herbivore attack. Oikos, 62(2), 244–251. https://doi.org/10.2307/3545270

Schlinkert, H., Westphal, C., Clough, Y., Grass, I., Helmerichs, J., and Tscharntke, T. (2016). Plant size affects mutualistic and

antagonistic interactions and reproductive success across 21 Brassicaceae species. Ecosphere, 7(12), e01529.

https://doi.org/10.1002/ecs2.1529

Schulze, J., Oeschger, L., Gross, A., Mueller, A., Stoll, P., and Erhardt, A. (2012). Solitary bees – Potential vectors for gene flow

from cultivated to wild strawberries. Flora - Morphology, Distribution, Functional Ecology of Plants, 207(10), 762–

767. https://doi.org/10.1016/j.flora.2012.07.007

Simms, E. L., and Rausher, M. D. 1987. Costs and Benefits of Plant Resistance to Herbivory. The American Naturalist, 130(4).

https://www.journals.uchicago.edu/doi/10.1086/284731

Smith, R. A., and Rausher, M. D. (2008). Experimental evidence that selection favors character displacement in the ivyleaf

morning glory. The American Naturalist, 171(1), 1–9. https://doi.org/10.1086/523948

Stinchcombe, J. R., Agrawal, A. F., Hohenlohe, P. A., Arnold, S. J., and Blows, M. W. (2008). Estimating nonlinear selection

gradients using quadratic regression coefficients: double or nothing? Evolution, 62(9), 2435–2440.

https://doi.org/10.1111/j.1558-5646.2008.00449.x

Svensson, E. I. (2023). Phenotypic selection in natural populations: What have we learned in 40 years? Evolution, 77(7), 1493–

1504. https://doi.org/10.1093/evolut/qpad077

Takahashi, F., Kuromori, T., Urano, K., Yamaguchi-Shinozaki, K., and Shinozaki, K. (2020). Drought stress responses and

resistance in plants: from cellular responses to long-distance intercellular communication. Frontiers in Plant Science,

11, 556972. https://doi.org/10.3389/fpls.2020.556972

Thompson, J. N. (2005). The Geographic Mosaic of Coevolution. University of Chicago Press.

https://press.uchicago.edu/ucp/books/book/chicago/G/bo3533766.html

Vallejo-Marín, M., Dorken, M. E., and Barrett, S. C. H. (2010). The ecological and evolutionary consequences of clonality for

plant mating. Annual Review of Ecology, Evolution, and Systematics, 41(41), 193–213.

https://doi.org/10.1146/annurev.ecolsys.110308.120258

Wise, M. J., and Rausher, M. D. (2013). Evolution of resistance to a multiple-herbivore community: Genetic correlations, diffuse

coevolution, and constraints on the plant's response to selection. Evolution, 67(6), 1767–1779.

https://doi.org/10.1111/evo.12061

Woods, E. C., Hastings, A. P., Turley, N. E., Heard, S. B., and Agrawal, A. A. (2012). Adaptive geographical clines in the growth

and defense of a native plant. Ecological Monographs, 82(2), 149–168. https://doi.org/10.1890/11-1446.1

Yahdjian, L., and Sala, O. E. (2002). A rainout shelter design for intercepting different amounts of rainfall. Oecologia, 133(2),

95–101. https://doi.org/10.1007/s00442-002-1024-3

Zhao, C., Han, W.-H., Xiong, Y.-D., Ji, S.-X., Du, H., Chi, Y.-J., Chen, N., Wu, H., Liu, S.-S., and Wang, X.-W. (2025). Drought

suppresses plant salicylic acid defence against herbivorous insects by down-regulating the expression of ICS1 via NAC

transcription factor. Plant Stress, 16, 100887. https://doi.org/10.1016/j.stress.2025.100887

Züst, T., and Agrawal, A. A. (2017). Trade-offs between plant growth and defense against insect herbivory: an emerging

mechanistic synthesis. Annual Review of Plant Biology, 68, 513–534. https://doi.org/10.1146/annurev-arplant-042916-

040856

Figure 1

A

B

C

D

E

Figure 2

Figure 3

Figure 4

Context-dependent selection and genetic facilitation and constraint on rosette diameter and herbivore resistance across European outdoor common gardens under ambient and reduced precipitation in *Fragaria vesca*

Ivan M. De-la-Cruz[1,8]*, Carolina Diller[1], Femke Batsleer[2], Dries Bonte[2], Timo Hytönen[3], José Luis Izquierdo[4], Sonia Osorio[5], David Posé[5], Aurora de la Rosa[4], Martijn L. Vandegehuchte[6]§, Anne Muola[7]§ and Johan A. Stenberg[1]§*

[1]Department of Plant Protection Biology, Swedish University of Agricultural Sciences, Box 190, SE-23422 Lomma, Sweden

[2]Department of Biology, Terrestrial Ecology Unit, Ghent University, Karel Lodewijk Ledeganckstraat, 9000 Gent, Belgium

[3]Department of Agricultural Sciences, Viikki Plant Science Centre, University of Helsinki, 0014 Helsinki, Finland

[4]Centro de Investigación, Seguimiento y Evaluación, Parque Nacional Sierra de Guadarrama, 28740 Rascafría, Spain

[5]Instituto de Hortofruticultura Subtropical y Mediterránea La Mayora, Departamento de Biología Molecular y Bioquímica, Universidad de Málaga-Consejo Superior de Investigaciones Científicas, Campus de Teatinos, 29750 Algarrobo, Spain

[6]Department of Biology, Norwegian University of Science and Technology, Høgskoleringen 5, 7491 Trondheim, Norway

[7]Division of Biotechnology and Plant Health, Invertebrate Pests and Weeds in Forestry, Agriculture and Horticulture, Norwegian Institute of Bioeconomy Research, 9016 Tromsø, Norway

[8]Evolutionary Biology Centre, Department of Ecology and Genetics, Plant Ecology and Evolution Division, Uppsala University, 752 36 Uppsala, Sweden

*Corresponding authors: Ivan M. De-la-Cruz, ivan.de.la.cruz.arguello@slu.se; Johan A. Stenberg, johan.stenberg@slu.se.

§Shared senior authorship.

Supplementary Table S1. Main abiotic environmental conditions occurring at the three
experimental sites. Average soil moisture (% of humidity) measured with a FieldScout TDR 150,
Spectrum Technologies, Inc. Soil moisture measurements were taken from two opposite sides on
the surface of each plant's pot at two different time points during the growing season of *F. vesca*
at each site in 20211 and 2022. Mean annual temperature and precipitation values were retrieved
from the closest weather stations: Spain: Agencia Estatal de Meteorología; Belgium: The Royal
Meteorological Institute of Belgium; Sweden: Swedish Meteorological and Hydrological
Institute. MASL = Meters above sea level. See also Figure 1A-D.

Site	Year	Mean soil moisture (% of humidity) during growing season		Mean temperature (ºC) during growing season	Mean rainfall (mm) during growing season	Length of growing season	Flowering period	Altitude MASL
		Control	Reduced precipitation					
Spain	2021	14.23	4.97	17.53	44.26	**April–August**	Late **April–July**	1200
	2022	27.57	18.81	19.49	17.06			
Belgium	2021	42.86	24.52	23.18	110	Late **March–August**	Late **April–July**	23
	2022	38.46	12.89	22.63	304.6			
Sweden	2021	16.85	5.79	18.63	62.60	Late **March–August**	Late **April–July**	9
	2022	12.83	3.03	18.56	52.03			

Supplementary Table S2. Genotype labels and latitude and longitude of origin. Order is from lowest to highest latitudes.

Genotype	Latitude	Longitude
ES2	37.7796	-3.7849
ES20	40.2938	-5.0091
ES13	43.1344	-4.888
IT3	45.94	10.81
GER100	47.96666667	7.83333333
FR2	48.801407	2.130122
FR3	50.015654	2.6973567
GER3	51.4192	11.128
LIT3	54.5729	24.6722
DK1	55.5703	9.7466
SE6	60.1018	18.3433333
FIN13	60.2292	25.0233
FIN39	60.4051	25.1562
NOR21	70.1671	24.7561
NOR19	70.0301	22.0653
NOR8	70.0324	23.4012

Supplementary Table S3. Directional and correlational selection gradients on herbivore resistance and rosette diameter across environments (every site × year × treatment combination). The linear selection gradients (β) for resistance and rosette diameter and the correlational selection coefficient (Resistance × Rosette) were estimated separately for fruit production (sexual fitness proxy) and stolon production (asexual fitness proxy). Standard errors (SE) and p-values testing the null hypothesis of no selection (coefficient = 0) are also shown. Positive β indicates selection favouring increases in the focal trait, whereas negative β indicates selection favouring decreases. A positive interaction term indicates selection favouring matched trait combinations (high–high and low–low), whereas a negative interaction indicates selection favouring mismatched combinations (high–low and low–high or a trade-off). Environments are defined by year (2021 and 2022) × site (Spain, Belgium, Sweden) × treatment (control *vs.* reduced precipitation). See Figure 2 and Supplementary Figure S1.

Year	Site	Treatment	β Resistance	SE Resistance	p	β Rosette	SE Rosette	p	Resistance × Rosette	SE Resistance × Rosette	p
						Fruits					
2021	Sweden	Control	0.273	0.123	0.013	0.856	0.185	0.000	0.027	0.106	0.796
2022	Sweden	Control	0.060	0.135	0.650	0.096	0.139	0.472	-0.011	0.125	0.928
2021	Belgium	Control	-0.004	0.079	0.964	0.401	0.143	0.001	-0.048	0.082	0.572
2022	Belgium	Control	0.034	0.079	0.660	0.859	0.192	0.000	0.004	0.081	0.961
2021	Spain	Control	-0.060	0.076	0.444	0.792	0.170	0.000	0.088	0.089	0.303
2022	Spain	Control	0.123	0.093	0.159	0.341	0.109	0.000	0.081	0.084	0.319
2021	Sweden	Reduced precipitation	0.103	0.094	0.252	0.375	0.130	0.001	0.155	0.102	0.100
2022	Sweden	Reduced precipitation	-0.278	0.072	0.001	0.875	0.199	0.000	0.151	0.097	0.093
2021	Belgium	Reduced precipitation	0.019	0.082	0.810	0.788	0.168	0.000	-0.094	0.074	0.227
2022	Belgium	Reduced precipitation	0.068	0.076	0.356	0.551	0.130	0.000	-0.086	0.064	0.198
2021	Spain	Reduced precipitation	0.042	0.083	0.602	0.625	0.148	0.000	0.040	0.086	0.636
2022	Spain	Reduced precipitation	-0.076	0.100	0.466	-0.455	0.056	0.000	0.293	0.168	0.048
						Stolons					
2021	Sweden	Control	-0.167	0.034	0.000	0.413	0.062	0.000	-0.032	0.042	0.444
2022	Sweden	Control	-0.186	0.057	0.003	0.393	0.106	0.000	-0.005	0.073	0.951
2021	Belgium	Control	-0.039	0.033	0.241	0.504	0.064	0.000	0.014	0.035	0.688
2022	Belgium	Control	0.114	0.092	0.191	0.716	0.110	0.000	-0.051	0.071	0.486
2021	Spain	Control	0.042	0.042	0.312	0.350	0.059	0.000	0.014	0.039	0.717
2022	Spain	Control	-0.059	0.067	0.398	0.511	0.104	0.000	0.033	0.071	0.631
2021	Sweden	Reduced precipitation	-0.170	0.033	0.000	0.180	0.051	0.000	-0.016	0.035	0.649
2022	Sweden	Reduced precipitation	-0.263	0.052	0.000	0.621	0.110	0.000	0.011	0.059	0.851
2021	Belgium	Reduced precipitation	0.030	0.031	0.322	0.207	0.043	0.000	-0.048	0.030	0.122
2022	Belgium	Reduced precipitation	0.049	0.055	0.361	0.462	0.083	0.000	-0.039	0.044	0.390
2021	Spain	Reduced precipitation	0.026	0.040	0.504	0.254	0.055	0.000	0.016	0.038	0.673
2022	Spain	Reduced precipitation	-0.214	0.053	0.000	0.129	0.079	0.084	0.134	0.069	0.039

Supplementary Figure S1. Correlational selection surfaces between rosette diameter and
herbivore resistance across environments; sites (Spain, Belgium and Sweden), treatments
(control and reduced precipitation) and years (2021 and 2022). (A) Predicted fruit production and
(B) predicted runner production are shown as functions of herbivore resistance (x-axis) and
rosette diameter (y-axis). For each environment (site × treatment × year), the heatmap shows
predicted fitness (lighter colours indicate higher values; see colour bars), and contour lines
denote iso-fitness contours. Dashed and dotted diagonals indicate reference axes representing
contrasting trait combinations (high–high *vs*. high–low). White asterisks depict significance of
the correlational selection. See Supplementary Table S3.

Supplementary Table S4. Genetic facilitations and constraints across environments (i.e., every site × treatment × year combination). Genetic covariances were classified as facilitating evolutionary change for resistance or rosette diameter when FC > 0, indicating that covariances increase the magnitude of the predicted response in the direction of selection. This is indicated by the *p_facil* column, which is showing the posterior probability of *direction-aware facilitation* (FC > 0). Conversely, covariances were classified as constraining when FC < 0 or p_const showing the posterior probability of *direction-aware constraint* (FC < 0). Posterior median of Tdiff is showing the *effect size* or how much covariances change the predicted response of that trait. 2.5% and 97.5% quantiles of TDiff (a 95% credible interval for TDiff). Significant facilitation: p_facil ≥ 0.90, significant constraint: p_const ≥ 0.90. See also methods and Figure 4.

Trait	Year	Site	Treatment	p_facil	p_const	Median TDiff	Low 95% IC	High 95% IC	Trait	Year	Site	Treatment	p_facil	p_const	Median TDiff	lo	hi
Resistance	2021	Sweden	Control	0.51354167	0.48645833	0.00584574	-0.4997448	0.50612821	Resistance	2021	Sweden	Control	0.48645833	0.51354167	0.00316947	-0.270954	0.27441494
Rosette	2021	Sweden	Control	0.51354167	0.48645833	0.00178889	-0.1529304	0.15488381	Rosette	2021	Sweden	Control	0.48645833	0.51354167	-9.31E-04	-0.0806114	0.07959475
Resistance	2021	Sweden	Reduced precipitation	0.93420833	0.06579167	0.17144299	-0.0555186	0.57449432	Resistance	2021	Sweden	ReduControoled preControlipitation	0.06579167	0.93420833	0.04523518	-0.0146486	0.15158016
Rosette	2021	Sweden	Reduced precipitation	0.93420833	0.06579167	0.02649645	-0.0085804	0.0887879	Rosette	2021	Sweden	ReduControled preControlipitation	0.06579167	0.93420833	-0.0389095	-0.1303831	0.01260009
Resistance	2021	Belgium	Control	0.580125	0.419875	0.02637259	-0.5176852	0.83881046	Resistance	2021	Belgium	Control	0.419875	0.580125	0.01699335	-0.3335739	0.54049307
Rosette	2021	Belgium	Control	0.580125	0.419875	4.90E-04	-0.0096201	0.01558747	Rosette	2021	Belgium	Control	0.419875	0.580125	-0.0011249	-0.0357801	0.02208228
Resistance	2021	Belgium	Reduced precipitation	0.4735	0.5265	-0.0061208	-0.7158679	0.67609953	Resistance	2021	Belgium	ReduControled preControlipitation	0.4735	0.5265	-0.0012757	-0.1492056	0.14091682
Rosette	2021	Belgium	Reduced precipitation	0.4735	0.5265	-4.83E-04	-0.0565062	0.05336714	Rosette	2021	Belgium	ReduControled preControlipitation	0.4735	0.5265	-1.32E-04	-0.0154433	0.01458536
Resistance	2021	Spain	Control	0.79191667	0.20808333	-0.1731552	-1.0179534	0.24232964	Resistance	2021	Spain	Control	0.20808333	0.79191667	-0.0618047	-0.3633408	0.08649536
Rosette	2021	Spain	Control	0.79191667	0.20808333	1.22E-04	-1.70E-04	7.15E-04	Rosette	2021	Spain	Control	0.20808333	0.79191667	-0.0081007	-0.0476227	0.01133686
Resistance	2021	Spain	Reduced precipitation	0.728875	0.271125	0.04367797	-0.1293544	0.3524122	Resistance	2021	Spain	ReduControled preControlipitation	0.728875	0.271125	0.02349194	-0.0695725	0.18954279
Rosette	2021	Spain	Reduced precipitation	0.728875	0.271125	0.00132716	-0.0039304	0.01070806	Rosette	2021	Spain	ReduControled preControlipitation	0.728875	0.271125	0.00149365	-0.0044235	0.01205139
Resistance	2022	Sweden	Control	0.21225	0.78775	-0.0330393	-0.2121247	0.0503349	Resistance	2022	Sweden	Control	0.78775	0.21225	-0.0880865	-0.5655491	0.13419866
Rosette	2022	Sweden	Control	0.21225	0.78775	-0.0411516	-0.2642087	0.06269385	Rosette	2022	Sweden	Control	0.78775	0.21225	0.03826451	-0.0582955	0.2456727
Resistance	2022	Sweden	Reduced precipitation	0.99104167	0.00895833	-0.4399823	-1.2414158	-0.0710315	Resistance	2022	Sweden	ReduControled preControlipitation	0.99104167	0.00895833	-0.5215912	-1.4716764	-0.0842066
Rosette	2022	Sweden	Reduced precipitation	0.99104167	0.00895833	0.20113948	0.03247229	0.56751762	Rosette	2022	Sweden	ReduControled preControlipitation	0.99104167	0.00895833	0.22275862	0.03596252	0.62851632
Resistance	2022	Belgium	Control	0.37825	0.62175	-0.0273806	-0.5190528	0.30278481	Resistance	2022	Belgium	Control	0.37825	0.62175	-0.0268388	-0.5087825	0.29679371
Rosette	2022	Belgium	Control	0.37825	0.62175	-0.0021018	-0.0398436	0.0232424	Rosette	2022	Belgium	Control	0.37825	0.62175	-0.0024805	-0.0470233	0.02743063
Resistance	2022	Belgium	Reduced precipitation	0.24183333	0.75816667	-0.1362531	-0.9752412	0.28211789	Resistance	2022	Belgium	ReduControled preControlipitation	0.24183333	0.75816667	-0.1094875	-0.7836646	0.22669857
Rosette	2022	Belgium	Reduced precipitation	0.24183333	0.75816667	-0.0037429	-0.0267904	0.00774993	Rosette	2022	Belgium	ReduControled preControlipitation	0.24183333	0.75816667	-0.0067074	-0.0480089	0.01388801
Resistance	2022	Spain	Control	0.25379167	0.74620833	-0.067654	-0.472357	0.1764258	Resistance	2022	Spain	Control	0.74620833	0.25379167	-0.0812544	-0.5673146	0.21189257
Rosette	2022	Spain	Control	0.25379167	0.74620833	-0.0143733	-0.1003537	0.03748222	Rosette	2022	Spain	Control	0.74620833	0.25379167	0.00789774	-0.0205955	0.05514163
Resistance	2022	Spain	Reduced precipitation	0.3395	0.6605	0.03242438	-0.1859818	0.38096255	Resistance	2022	Spain	ReduControled preControlipitation	0.6605	0.3395	-0.0030388	-0.0357038	0.01743021
Rosette	2022	Spain	Reduced precipitation	0.3395	0.6605	0.01292856	-0.0741564	0.15190106	Rosette	2022	Spain	ReduControled preControlipitation	0.6605	0.3395	0.02307946	-0.1323806	0.27116661