A conserved photosynthetic cytochrome enhances growth of
Chlamydomonas reinhardtii in fluctuating light

Darius Kosmützky[1,†,*], Laura T Wey[2], Lauri Nikkanen[2], Aron Ferenczi[3], Joshua M Lawrence[1,4],
Alberto Scarampi[1,‡], Attila Molnar[3], Yagut Allahverdiyeva[2], Christopher J Howe[1,*]

1: Department of Biochemistry, University of Cambridge, Cambridge CB2 1QW, UK

2: Molecular Plant Biology, Department of Life Technologies, University of Turku, Turku 20014,
Finland

3: Institute of Molecular Plant Sciences, University of Edinburgh, Edinburgh EH9 3BF, UK

4: Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, UK

†: Present address: Sainsbury Laboratory, University of Cambridge, Cambridge CB2 1LR, UK

‡: Present address: School of Life Sciences, University of Warwick, Coventry CV4 7AL, UK

*: Corresponding authors

Abstract

Having at least one of the small soluble electron carriers cytochrome c_6 (c_6) and plastocyanin is indispensable for photosynthesis. It was believed that c_6 had been lost from plants until a ubiquitous and highly conserved homologue, now named cytochrome c_{6A} (c_{6A}), was found in *Arabidopsis thaliana*. In spite of an early suggestion that c_{6A} functioned like c_6 in the photosynthetic electron transport chain it was soon shown to be unable to do so despite the strong structural homology between the two proteins. The function of c_{6A}, and why it is apparently universal in plants and green algae, therefore remained unknown. Here, we show that, in the green alga *Chlamydomonas reinhardtii*, c_{6A} confers a growth advantage under fluctuating light and is crucial for maintaining the light harvesting balance between photosystems I and II in photomixotrophic conditions. We show that in the absence of c_{6A}, the light harvesting balance shifts towards PSII, leading to a more reduced plastoquinone pool and increased photooxidative stress. This study provides insights into how photosynthetic organisms acclimatize to stressful light conditions, indicating that c_{6A} is important for this adaptation. These findings provide a basis for further mechanistic studies on a hypothesized role for c_{6A} in thiol-based redox regulation in the thylakoid lumen, with implications for photoprotection mechanisms such as state transitions.

Main Text

Introduction

The photosynthetic electron transport chain (PETC) is essential for nearly all life on Earth (Fig. 1a). At least one of the soluble electron carriers plastocyanin and cytochrome c_6 is indispensable for the PETC to function (1, 2). A close homologue of c_6, cytochrome c_{6A}, was first discovered in *Arabidopsis thaliana* and later in other plants and eukaryotic algae (3, 4). Whereas plants lost the gene for c_6 and contain only c_{6A}, some eukaryotic algae such as *Chlamydomonas reinhardtii* (*Chlamydomonas*) contain genes for both (5), although c_{6A} is expressed at a low level compared to c_6 or plastocyanin (6, 7). Since its discovery, c_{6A} has been hypothesized to have various functions, including replacement of plastocyanin (3), acting as a safety valve for excess electrons of photosynthesis (7), regulation of generation of reactive oxygen species (ROS) in PSI (8), and being

involved in high light signalling (9) or thiol-based redox regulation (10, 11). However, the function

of cytochrome c_{6A} remains elusive (7).

While c_6 and c_{6A} are homologues, there are structural differences between the two (Fig. 1b). c_{6A}

contains a 'loop insertion peptide' (LIP), an insertion with a conserved disulfide bridge in a surface-

located and well-defined loop. The LIP is absent from c_6 (10). Assessing the stability of various

mutant proteins showed that the LIP is structurally important (12, 13). While a catalytic role for the

conserved disulfide bridge within the LIP is uncertain, it is predicted to contribute to the protein's

structural stability (13, 14). In addition to the LIP, there are differences in the heme environment

of the two proteins, and these contribute to c_{6A} (+71 mV (15)) having a 254 mV lower midpoint

potential than plastocyanin or c_6 (+325 mV (15)).

Phylogenetic analyses revealed that c_{6A} represents a monophyletic group distinct from c_6 and likely

originates from cyanobacterial cytochrome c_{6B}, another homologue of c_6 that has a similar midpoint

potential to c_{6A} but lacks the LIP (16). The sequence of c_{6A} is highly conserved across the green

lineage, as with plastocyanin, suggesting an important role for c_{6A} within photosynthetic

eukaryotes (16). However, the large difference in midpoint potentials renders it highly unlikely that

c_{6A} acts as a functional replacement for plastocyanin in plants and algae, as c_{6A} would not be

expected to accept electrons from the cytochrome b_6f complex (Fig. 1a). Further, plastocyanin was

found to be indispensable even when c_{6A} was overexpressed in *A. thaliana* (1). Additionally, the

surfaces of c_6 and plastocyanin are both strongly negatively charged, whereas c_{6A} has a strongly

positive surface charge (17). Together with the difference in isoelectric points (plastocyanin: 5.1,

c_{6A}: 4.1 in *A. thaliana* (18)) this indicates different interactions.

Interactions involving c_{6A} further highlight its distinct role (Fig. 1a). It cannot effectively reduce

PSI *in vitro*, making direct interaction between c_{6A} and PSI unlikely (17). Additionally, c_{6A} has

been shown to reduce plastocyanin *in vitro* (19) and weak interaction (K_d = 2.01 mM) was detected

with plastocyanin using NMR titrations (6), consistent with a transient interaction. Yeast Two-

Hybrid assays also revealed interactions of c_{6A} with itself (20), the lumen thiol oxidase LTO1 (21),

and the redox-regulated immunophilin FKBP13 (3). LTO1 has been reported to be involved in

regulating state transitions via the kinase STT7 (22). Interaction with LTO1 and FKBP13 therefore

suggests a luminal localization for c_{6A}, consistent with its detection in the soluble fraction of the

thylakoid lumen in *A. thaliana* (23). In *Chlamydomonas*, however, only a chloroplastic localization

has been confirmed (6). In *A. thaliana*, c_{6A} was also found to be uniquely localized to sensory

plastids in the epidermis and vascular parenchyma rather than mesophyll chloroplasts (24).

Photosynthetic organisms require flexibility in their photosynthetic apparatus because they grow

under a diverse set of environmental conditions. These include varying light conditions from

diurnal cycles to more rapid fluctuations caused by the sun, clouds, canopy, and in the case of

Chlamydomonas, also soil and water movements. For *Chlamydomonas*, exposure to a 12 s on/12 s

off cycle with light intensities of 600-700 µmol photons m^{-2} s^{-1} in photoautotrophic conditions

resulted in growth rates reaching only about 30% of those achieved under continuous

illumination (25). Additionally, a growth penalty was revealed under fluctuating light with

frequencies in the range of 10-100 s on/off (26). In mutant assays using *Chlamydomonas*, various

fluctuating light regimes have been employed to determine the function of proteins associated with

photosynthesis. These include the important proteins Proton Gradient Regulation 5 (PGR5) and

Proton Gradient Regulation Like 1 (PGRL1), both implicated in cyclic electron flow, its regulation

or ATP synthase regulation (27), as well as flavodiiron proteins (FLVA/B), which facilitate the

Mehler-like reaction (28). Additionally, the qE quenching effector LHCSR3 (29, 30) and plastid

terminal oxidase PTOX2 (31) have been implicated in the response to fluctuating light, as mutants

of all of these proteins showed a growth deficit under fluctuating light conditions.

Here, we present the first in-depth analysis of the connection between c_{6A} and photosynthesis. We

show that this highly conserved protein is important for growth under a harsh fluctuating light

regime we call DISCO Light and helps maintain the redox balance of the plastoquinone (PQ) pool

through its effect on the light harvesting balance between photosystems I (PSI) and II (PSII).

Results

Cytochrome c_{6A} confers a growth advantage under a fluctuating light regime

To investigate the role of c_{6A} in *Chlamydomonas* we used CRISPR/Cpf1 (32) to create a knock-out

of the c_{6A} gene (*CYC4*, Cre16.g670950) in the background strain CC-1883 (WT). The knock-out

strain ("c_{6A}-KO") was confirmed through PCR and sequencing (Supplementary Methods &

Supplementary Fig. 1a-d). We also generated a c_{6A} overexpression c_{6A}-OE) strain by transforming

the c_{6A}-KO strain with a construct that drives the expression of FLAG-tagged c_{6A} from a *PSAD*

promoter. Overexpression was confirmed by immunoblot and RNAseq analysis (Supplementary

Methods & Supplementary Fig. 1d-e).

To investigate the role of cytochrome c_{6A} in growth, we tested how *Chlamydomonas* strains grow

under various light and trophic regimes. Under continuous light (30 μmol photons m^{-2} s^{-1}), both

c_{6A}-KO and WT strains reached similar maximum optical densities (OD$_{750nm}$) of around 2 in

photomixotrophic and ~0.6 in photoautotrophic conditions, with no statistically significant

differences (Fig. 1c). We defined continuous light (30 μmol photons m^{-2} s^{-1}) and photomixotrophic

conditions as standard conditions for this study.

118
Figure 1: Cytochrome c_{6A} confers a growth advantage under a fluctuating light regime. a, Schematic of
the photosynthetic electron transport chain with known (black) and putative interactions (purple) of c_{6A} marked.
Proton movement (grey) and the Q-cycle (blue) are shown as arrows. **b,** Overlay of the crystal structures of
cytochrome c_6 from *Chlamydomonas* (orange, PDB: 1CYJ (57)) and c_{6A} from *A. thaliana* (purple, PDB: 2CE0
(19)). The position of the LIP is noted with the disulfide bridge formed between the two conserved cysteine
residues (yellow sticks). The amino acid residues at position 52 were found to affect the heme environment
contributing to the low midpoint potential of c_{6A} compared to c_6 or plastocyanin (stick model and labelled with
a green diamond) (15). **c,** Optical density (OD$_{750nm}$) growth curve of c_{6A} mutant strains and WT under
continuous light (30 μmol$_{photons}$ m^{-2} s^{-1}) in either photoautotrophy or -mixotrophy and DISCO Light and
photomixotrophy (n = 3 biological replicates ±SD). Statistical testing was carried out with repeated measures
ANOVA. Post hoc pairwise two-sided t-tests at each time point were performed if the interaction effect of time
and strain on OD$_{750nm}$ in the ANOVA was significant (p$_{time:strain}$ < 0.05) comparing WT (pink) or c_{6A}-OE (green)
against c_{6A}-KO (p < 0.05: *, p ≤ 0.1: #). **d,** DISCO Light (Darkness Interrupted by Short COnstant Light) consists

of alternating 2 min darkness followed by 2 min of high light pulses (12 s high light (700 $\mu mol_{photons}$ m^{-2} s^{-1}) &
12 s darkness). **e**, Visual differences of the c_{6A} mutant and WT cultures of at day 3 of growth in DISCO Light.
134

We also tested multiple stressful light conditions for their effect on c_{6A}-KO and WT. Only a light

regime we named DISCO (Darkness Interrupted by Short COnstant) Light and photomixotrophy

led to an observable phenotypic growth difference (Fig. 1c). DISCO Light was devised to expose

Chlamydomonas to a complex fluctuating light regime covering multiple frequencies, as it consists

of two alternating phases: 2 min of darkness followed by 2 min of alternating 12 s high light pulses

of 700 $\mu mol_{photons}$ m^{-2} s^{-1} and 12 s darkness (Fig. 1d). c_{6A}-KO grew more slowly and reached lower

cell densities than WT under photomixotrophic DISCO conditions (Fig. 1c & e): after one day c_{6A}-

KO cultures showed significantly reduced growth ($P = 0.029$), and by day 3 they were

approximately 1.5 times less dense than WT cultures ($P = 0.10$). The growth penalty in c_{6A}-KO

was recovered by overexpressing c_{6A}, with c_{6A}-OE showing significantly higher cell densities than

c_{6A}-KO from day 1 ($P = 0.024$, Fig. 1c & e). In the DISCO light regime without acetate in the

medium (photoautotrophy), no strain could grow (Supplementary Fig. 2e).

In photomixotrophy and darkness or continuous high light at the same light intensity as in the

DISCO regime (700 $\mu mol_{photons}$ m^{-2} s^{-1}) WT and c_{6A}-KO grew at similar rates (Supplementary

Fig. 2a-b). Furthermore, fluctuating light regimes corresponding to elements of the DISCO

protocol ("2 min high light & 2 min darkness", and "12 s high light & 12 s darkness"), showed only

small differences between WT and c_{6A}-KO that were not part of a consistent trend under either

photomixotrophic or photoautotrophic conditions (Supplementary Fig. 2c-d).

Measurements of total chlorophyll concentration and acetate consumption confirmed the

phenotypic difference between the strains observed from OD_{750nm} measurements in both standard

and DISCO Light conditions (Supplementary Fig. 2f-h). The correlation between total chlorophyll

concentration and optical density was similar between WT and c_{6A}-KO. This indicated no

difference in chlorophyll concentration per cell between the strains under standard conditions or

DISCO Light (Supplementary Fig. 2g). Additionally, measurements of acetate concentration in the

growth medium indicated that c_{6A}-KO consumed less acetate, likely because of lower cell densities

when exposed to DISCO Light, with differences apparent from day 3 (Supplementary Fig. 2h).

These results indicate that c_{6A} is important for growth of *Chlamydomonas* under the high intensity

and rapidly fluctuating DISCO Light regime and acetate availability in photomixotrophic

conditions.

<u>c_{6A} is likely localized in the thylakoid lumen and does not have strongly bound interaction partners</u>

c_{6A} was detected in the soluble fraction of the thylakoid lumen in *A. thaliana* (23). In

Chlamydomonas, however, only chloroplastic localization has been confirmed (6). To determine

the subcellular localization of c_{6A} in *Chlamydomonas* further, we performed immunofluorescence

microscopy on the c_{6A}-FLAG overexpressing strain (c_{6A}-OE) and its background strain (c_{6A}-KO)

grown under standard conditions. In the negative control (c_{6A}-KO), the AlexaFluor488 signal

corresponding to FLAG was weak and non-specific, i.e. appearing throughout the cell including in

the flagella (Supplementary Fig. 3a). In contrast, the c_{6A}-FLAG overexpressing sample (c_{6A}-OE)

showed a more intense (2 to 3-fold pixel intensity) and more localized fluorescence signal,

confirming chloroplast-specific localization of c_{6A} (Supplementary Fig. 3b). Two cells covering the

variation of AlexaFluor488 signal intensity were chosen as representatives to capture cell to cell

differences that could arise from technical variation from fixing and staining or c_{6A}-FLAG

expression level differences between cells. Nevertheless, the absence of AlexaFluor488 signal from

cell compartments other than the chloroplast in c_{6A}-OE compared to c_{6A}-KO supports the FLAG-

specificity of the signal. The detection of anti-FLAG signal in structures resembling pyrenoid-

traversing thylakoids supports a thylakoid luminal localization of the soluble

c_{6A} (Supplementary Fig. 3c-d). While it cannot be excluded that c_{6A} is associated with the thylakoid

membrane on the stromal side, the structure of c_{6A} (Fig. 1b) gives no indication of membrane-

association.

To detect possible covalent or strongly bound interaction partners, we performed co-

immunoprecipitation (co-IP) analyses of the c_{6A}-FLAG overexpressing strain (c_{6A}-OE) and the

untransformed background strain (c_{6A}-KO) using anti-FLAG-agarose beads. Immunodetection of

FLAG in the precipitate revealed bands at ~15 kDa in c_{6A}-OE but not in c_{6A}-KO, corresponding to

mature c_{6A}-FLAG (red arrows in Supplementary Fig. 4), with additional FLAG signals that were

not c_{6A}-FLAG specific in the 25-35 kDa range (black arrows in Supplementary Fig. 4). Under both

standard (Supplementary Fig. 4c) and DISCO Light (Supplementary Fig. 4f) conditions, the

immunoprecipitate did not reveal any bands unique to c_{6A}-OE except the c_{6A}-FLAG band,

indicating that no specific interaction partners of c_{6A} were detected.

The combined immunofluorescence microscopy and co-IP results indicate that c_{6A} is likely to be

localized in the thylakoid lumen, as for *A. thaliana* (23) and are consistent with any hypothesized

function relying primarily on weak or transient interactions.

<u>c_{6A} is important to maintain the light harvesting balance between photosystems I and II in</u>

<u>photomixotrophy</u>

Given the light-dependent growth phenotype observed in DISCO conditions, the likely localization

of c_{6A} in the thylakoid lumen, and c_{6A}'s high sequence similarity to c_6 which is involved in the

PETC, we investigated the photosystems and light harvesting complexes to explore further a

potential role of c_{6A} in photosynthesis.

We recorded low temperature (77K) chlorophyll fluorescence emission spectra. When grown in

continuous light (30 μmol$_{photons}$ m^{-2} s^{-1}) under photoautotrophic conditions WT and c_{6A}-KO

exhibited very similar fluorescence spectra (Fig. 2a, PSI/PSII fluorescence ratios 1.25 and 1.3,

respectively, $P = 0.374$), consistent with similar growth (Fig. 1c). However, when grown in

photomixotrophic conditions and continuous light (30 μmol$_{photons}$ m^{-2} s^{-1}, standard conditions), and

despite the two strains showing similar growth (Fig. 1b), WT maintained a PSI/PSII fluorescence

ratio of about 1.2, while c_{6A}-KO showed a significantly lower ratio (0.96, $P = 0.008$), which was

accompanied by a more prominent CP47-associated peak at 695 nm (33) (Fig. 2a). The PSI/PSII

fluorescence ratio was restored to nearly WT levels in the c_{6A}-OE strain (Fig. 2a). When samples

were grown in DISCO Light and photomixotrophy, the differences seen in standard conditions

were amplified. By day 4, the PSI/PSII fluorescence ratio in c_{6A}-KO dropped drastically to 0.66 in

comparison to WT and c_{6A}-OE (1.05 and 1.33, respectively & $P = 0.027$ and 0.0005, respectively).

Interestingly, WT showed a shoulder peak at ~681 nm (Fig. 2a), which may reflect partial

detachment of light harvesting complex II (LHCII) from PSII (34). The lower level of normalized

PSI fluorescence in c_{6A}-KO in all photomixotrophic samples did not originate from lower PSI

protein levels, as evidenced by PSI subunit immunoblot analyses (Supplementary Fig. 5).

To investigate if this c_{6A}-dependent light harvesting phenotype was due to alterations in state

transitions in photomixotrophy, for example due to c_{6A}-KO being locked in state 1, we induced

either state 1 (darkness and oxia) or state 2 (darkness and anoxia) in samples grown under standard

conditions (Fig. 2b). Both strains showed, as expected, a lower PSI/PSII fluorescence ratio in state

1 and a higher PSI/PSII fluorescence ratio in state 2, reflecting the increased sizes of the antenna

cross section of PSII and PSI, respectively. However, c_{6A}-KO consistently exhibited a lower

PSI/PSII fluorescence ratio than WT (Fig. 2c). Whereas WT exhibited PSI fluorescence of

1.13 (normalized to PSII fluorescence) in state 2 conditions, c_{6A}-KO showed a significantly lower

value of 1.01 ($P = 0.046$). Similarly, under state 1 inducing conditions, normalized PSI fluorescence

in WT was 0.85, compared to 0.77 in c_{6A}-KO ($P = 0.071$). While the CP47-associated peak at

695 nm (33) showed different intensity between the two states in WT, in c_{6A}-KO the same peak

intensity was recorded in both states.

As the light harvesting balance is affected by photosystem complex abundance and composition,

we next analyzed the thylakoid membrane complexes from cells grown in standard conditions using

BN-PAGE. Gels showed a distinct green band containing a PSI complex without LHCs (band III

marked red, Fig. 2d&e) that was consistently more prominent in c_{6A}-KO compared to WT and c_{6A}-

OE. The other bands showed a similar pattern and thus presumably similar photosystem complex

levels across strains (Fig. 2d). 2D-PAGE analyses and gel simulations to calculate the predicted

mobility in the gel (R_f) revealed similar subunit compositions of the complexes between

strains (Fig. 2e & Supplementary Fig. 7a-b).

Together, these results show that c_{6A} is important for maintaining the light harvesting balance

between the photosystems in photomixotrophic conditions, especially in combination with DISCO

light, and indicate that an altered distribution of LHCs could underly this observation.

249
Figure 2: *c*6A affects the light harvesting balance between photosystems I and II in photomixotrophy.
a, Low temperature (77 K) chlorophyll fluorescence spectra (excitation: 440 nm) of c_{6A} mutant strains and WT
from cells grown under continuous light (30 $\mu mol_{photons}$ m^{-2} s^{-1}) in photoautotrophy -or mixotrophy and cells
grown photomixotrophically under DISCO Light for 2 or 4 days. The prominent PSI and PSII fluorescence
peaks of note are indicated by black arrows. **b**, Low temperature (77 K) chlorophyll fluorescence spectra of
c_{6A}-KO and WT exposed to state 1 or 2 inducing conditions (n = 3 biological replicates ±SE). **c**, Quantification
of the PSI to PSII fluorescence ratio at 713 nm from **b**. Statistical testing was carried out with a two-sided t-
test per state (p < 0.05: *, p < 0.1: #). **d**, BN-PAGE analysis of c_{6A} mutant strains and WT thylakoid membrane
complexes isolated from cells grown in photomixotrophy and continuous light (30 $\mu mol_{photons}$ m^{-2} s^{-1}). The inset
shows the area of interest with band III from a separate longer BN-PAGE run with the same samples. The full
gels with three biological replicates per strain can be found in Supplementary Figure 6. Thylakoid membrane
complexes were assigned to bands according to the literature (53–55). **e,** 2D-PAGE analysis of a BN-PAGE
lane from c_{6A} -KO with the boxes indicating assignment of spots to protein complexes based on comparison
with calculated R$_f$ values of known thylakoid membrane complex components (Supplementary Fig. 7c).
264

Absence of c_{6A} leads to a more reduced PQ pool

Given the tight link between light harvesting balance and the redox state of the PQ pool, we

quantified the redox state of the PQ pool using chlorophyll fluorescence measurements.

Firstly, the chlorophyll fluorescence induction (OJIP) traces of cells grown in standard conditions

revealed a 10% increase in the F_J parameter in c_{6A}-KO compared to WT and c_{6A}-OE ($P = 0.012$ and

0.008, respectively) (Fig. 3a,b). Since F_J reflects the extent of Q_A reduction and is positively

correlated with PQ pool reduction (35), this increase indicates that the absence of c_{6A} led to a more

reduced PQ pool.

To investigate further the extent of c_{6A}-dependent PQ pool overreduction, we exposed cells grown

in standard conditions to DISCO light and recorded chlorophyll fluorescence. The value of the

steady-state fluorescence yield F' under actinic light is influenced by the redox state of the PQ pool,

as the PQ pool is in a redox equilibrium with Q_A in PSII (36). During the high light periods of

DISCO Light, F' was on average 36% higher in c_{6A}-KO than in WT ($P = 0.0006$), indicating a more

reduced PQ pool in the absence of c_{6A} in DISCO Light as well (Fig. 3c). Also during the high light

periods of the DISCO Light regime, 1-qL (a measure of Q_A reduction) was high in WT (0.9)

indicating that Q_A and by implication the PQ pool was highly reduced (37). In c_{6A}-KO 1-qL was

significantly higher (0.975) again compared to WT ($P = 6.8 \times 10^{-5}$) (Fig. 3d), indicating a more

reduced PQ pool, consistent with the F' finding.

These results suggest that the absence of c_{6A} leads to a more reduced PQ pool in *Chlamydomonas*

under photomixotrophic conditions, exacerbated by DISCO Light exposure. Consequently, the

more reduced Q_A pool could lead to increased damage to PSII, potentially contributing to the

observed growth penalty in DISCO Light.

Redox kinetics of *b* hemes are affected and PSII quantum yield lowered in absence of c_{6A}

Given the effects of the absence of c_{6A} on light harvesting and PQ pool redox balance, we also

assessed the function of PETC complexes. We first measured the redox kinetics of cytochrome b_6f

using a Joliot-type spectrometer (JTS).

In photoautotrophic samples grown under continuous light (30 $\mu mol_{photons}$ m^{-2} s^{-1}), all strains

exhibited a *b* hemes reduction peak within 100 ms of light exposure, but in c_{6A}-KO the peak was

only half the intensity of WT (Supplementary Fig. 8). In samples grown in standard

photomixotrophic conditions, the *b* hemes in WT and c_{6A}-OE were reduced within 100 ms of light

exposure followed by transient oxidation and re-reduction. In contrast, the *b* hemes in c_{6A}-KO were

oxidized within 100 ms of illumination (Fig. 3e) and showed nearly no re-reduction in the following

500 ms. Interestingly, no differences in the cytochrome *f* redox kinetics were observed in either

photoautotrophy or photomixotrophy (Supplementary Fig. 8), indicating no effect of the absence

of c_{6A} on the function of the high potential chain.

304
Figure 3: c_{6A} affects the redox state of the PQ pool, the redox kinetics of *b* hemes in cytochrome b_6f
and the quantum yield of PSII. a, Chlorophyll fluorescence induction measurements (OJIP curves) of c_{6A}
mutant strains and WT grown under photomixotrophic continuous light (30 $\mu mol_{photons}$ m^{-2} s^{-1}) conditions and
normalized to F_0 and F_P (n = 3 biological replicates ±SD). **b**, Comparison of the normalized fluorescence at
inflection point F_J (2.4 ms) from **a**. Asterisks indicate a significant difference (p < 0.05) as determined by one-
way ANOVA and post-hoc pairwise two-sided t-test (n = 3 biological replicates ±SD). **c**, Raw steady-state
fluorescence yield F' in Volts (V) of c_{6A}-KO and WT grown under standard conditions and exposed to a DISCO
Light regime (on top of box) in a DUAL-PAM-100 spectrometer (n = 3 biological replicates ±SD). **d**, 1-qL values
in the light phases of c_{6A}-KO and WT grown under standard conditions and exposed to the same light protocol
as in **c** (n = 3 biological replicates ±SD, asterisk: significant difference (p < 0.001) as determined by two-sided
t-test). **e**, Redox changes of *b* hemes in c_{6A} mutant strains and WT grown under standard conditions and
exposed to DCMU (competitive inhibitor of the Q_B site in PSII), methyl viologen (MV, electron acceptor of PSI)
or dibromothymoquinone (DBMIB, b_6f Q_P site blocker). Samples were exposed to green light (500 $\mu mol_{photons}$
m^{-2} s^{-1}, green bar) in a JTS-10 spectrometer (n = 3 biological replicates ±SD). **f**, Photosystem II quantum yield,
Y(II), of c_{6A}-KO and WT grown under standard conditions and exposed to the same light protocol as in **c** (n =
3 biological replicates ±SD).

A more reduced PQ pool leads to lower reduction and even apparent oxidation of the *b* hemes upon

light exposure (38–40). The latter is likely due to *b* hemes being pre-reduced in conditions with a

highly reduced PQ pool. Thus, the effect of the absence of c_{6A} on the redox kinetics of *b* hemes are

consistent with the observed change in the redox state of the PQ pool. Artificial modulation of the

PQ pool redox state should therefore lead to smaller differences in *b* hemes redox kinetics between

the strains. In the presence of dichlorophenyl-dimethylurea (DCMU), a PSII inhibitor that leads to

PQ pool oxidation, samples grown in standard conditions showed that the b hemes in c_{6A}-KO were

indeed reduced upon illumination but to a lower magnitude than in WT/ c_{6A}-OE (Fig. 3e). Similarly,

when we added methyl viologen (MV), an artificial electron acceptor from PSI that also leads to

PQ pool oxidation, the b hemes reduction was partially restored in in c_{6A}-KO (Fig. 3e).

Furthermore, when we added dibromothymoquinone (DBMIB), which blocks the Q_P site of

cytochrome b_6f and leads to PQ pool reduction, the b hemes only showed oxidation upon

illumination without differences between the strains (Fig. 3e).

The observed differences in b hemes redox kinetics can therefore be explained by the more reduced

PQ pool in c_{6A}-KO. Further, these results indicate that in the absence of c_{6A} the cytochrome b_6f

complex including the Q-cycle is in principle functional if the PQ pool is oxidized. Nevertheless,

small differences between c_{6A}-KO and WT/ c_{6A}-OE remained when the PQ pool was artificially

oxidized, warranting further investigation of the Q-cycle.

To investigate the possibility of a disturbed Q-cycle underlying the b hemes phenotype, we

measured the proton motive force (pmf) and its components of *Chlamydomonas* cells grown in

standard conditions using electrochromic shift (ECS) measurements. No substantial difference

between presence and absence of c_{6A} was detected in pmf, proton flux (vH^+) and thylakoid

conductivity (gH^+) when samples were exposed to either continuous red light (53 $\mu mol_{photons}$ m^{-2}

s^{-1}) or DISCO Light conditions (Supplementary Fig. 9a-f). Further, no differences between the

strains were found in the response of the ECS signal to a single turnover flash and the partitioning

of pmf into ΔpH and $\Delta\Psi$ (Supplementary Fig. 9g&h). Together, this is consistent with c_{6A} not

affecting Q-cycle function.

We also analyzed the efficiency of photosystems I and II of *Chlamydomonas* cells grown in

standard conditions when exposed to continuous or DISCO Light. In continuous light, we did not

find any significant differences between WT and c_{6A}-KO in the quantum yield of PSII

(Supplementary Fig. 10a) and PSI (Supplementary Fig. 10b) over a wide range of intensities. There

was, however, an elevated photosystem II quantum yield Y(II) for c_{6A}-OE in comparison to c_{6A}-

KO at higher light intensities (above 210 $\mu mol_{photons}$ s^{-1} m^{-2}) resulting in an increased relative

electron transport rate (rETRII). Conversely, during the high light periods of DISCO Light, Y(II)

was less than half in c_{6A}-KO compared to WT (Fig. 3f). In contrast to PSII, PSI quantum yield Y(I)

was not majorly affected by absence of c_{6A} and DISCO Light and showed a decreasing trend in the

light phases from 0.2 to about 0.125 over the course of 40 min in both WT and c_{6A}-KO

(Supplementary Fig. 11). For both strains, DISCO Light led to an increase in acceptor side

limitation Y(NA), indicating that in DISCO Light downstream electron acceptors are not readily

available (Supplementary Fig. 11). Consistent with the more reduced PQ pool in absence of c_{6A},

c_{6A}-KO exhibited a lower donor-side limitation Y(ND) than WT during DISCO Light treatment

(Supplementary Fig. 11).

The DISCO Light condition therefore seems to lead to inhibition of photosynthesis at PSII in c_{6A}-

KO which is consistent with the highly reduced PQ pool in absence of c_{6A}.

c_{6A} partly alleviates the stressful effect of DISCO Light on PSII

To determine if the more reduced PQ pool in the absence of c_{6A} causes stress for *Chlamydomonas*

and could explain the observed growth phenotype, we measured PSII energy partitioning of cells

grown in standard conditions and then exposed to DISCO Light using a DUAL-PAM.

Increased stress levels in the absence of c_{6A} were evident when examining the quantum yield of

non-regulated dissipation of excitation energy, Y(NO), which describes the fraction of energy that

is neither used for photochemistry nor dissipated as heat in a regulated manner via NPQ. At the

start of DISCO Light exposure, over 90% of light energy was dissipated in a non-regulated fashion

in c_{6A}-KO, while WT had values below 80% (Fig. 4a). Over 40 minutes of DISCO Light, Y(NO)

levels in c_{6A}-KO relaxed, nearing WT levels of about 70%. High Y(NO) levels indicate PSII

photoinhibition that is imminent or has already occurred due to over-reduction (41). When probing

for PsbA, C-terminal fragments forming two bands at around 15 kDa were visible (Fig. 4b). An

increased ratio of PsbA (D1) degradation products to full length PsbA (D1) protein, which is

indicative of PSII photoinhibition (42), was observed in c_{6A}-KO on Day 4 of DISCO Light exposure

(Fig. 4b & Supplementary Fig. 12). In c_{6A}-KO, 20% of the observed PsbA (D1) protein signal

originated from the C-terminal fragment bands, while in WT and c_{6A}-OE the C-terminal fragments

only contributed 8% and 7% to the total PsbA (D1) signal, respectively (Supplementary Fig. 12).

The yield of regulated non-photochemical quenching Y(NPQ) was negligible in dark-adapted WT

and c_{6A}-KO samples (Fig. 4b). However, over the first 40 min of DISCO Light exposure, c_{6A}-KO

exhibited approximately twice the Y(NPQ) levels as WT during all phases of the DISCO Light

protocol. This sustained Y(NPQ) response, even in the dark, indicates increased stress levels in the

absence of c_{6A}, leading to an amplified engagement of NPQ mechanisms to allow energy dissipation

in a regulated manner. Interestingly, increased Y(NPQ) in the absence of c_{6A} was also observed

under a range of continuous light intensities, which was partially restored in the c_{6A}-OE strain

(Supplementary Fig. 10a). Elevated NPQ levels in the absence of c_{6A} were further supported by an

approximately 1.5-fold protein level of the NPQ effector LHCSR3 (43) in c_{6A}-KO compared to WT

after two days of growth in DISCO Light (Fig. 4b & Supplementary Fig. 12).

Together, these data indicate that PSII experiences more photoinhibition in c_{6A}-KO than WT and

c_{6A}-OE under DISCO Light consistent with the observed reduced Y(II) in c_{6A}-KO under DISCO

Light (Fig. 3f). Increased photoinhibition is likely contributing to the observed growth penalty (Fig.

1c).

403
Figure 4: c_{6A} partly alleviates the stressful effect of DISCO Light on *Chlamydomonas*. a, Regulated non-
photochemical quenching Y(NPQ), and non-regulated energy dissipation Y(NO) of c_{6A}-KO and WT grown
under standard conditions and exposed to a DISCO Light regime in a DUAL-PAM-100 spectrometer (n = 3
biological replicates ±SD). **b**, Immunoblot analyses probing for PsbA in whole cell protein samples from c_{6A}
mutant strains and WT grown in DISCO Light for four days (n = 3 biological replicates, PonceauS stain as
total protein loading control). **c**, Immunoblot analyses probing for LHCSR3 in whole cell protein samples from
c_{6A} mutant strains and WT grown in DISCO Light for two days (n = 3 biological replicates, PonceauS stain as
total protein loading control). Relative quantification of the share of signal from C-terminal fragments for PsbA
or the signal intensities normalized to total protein for LHCSR3 and can be found in Supplementary Figure 12.
413

Discussion

Here we present an in-depth characterization of the physiological role of the highly conserved cytochrome c_{6A} protein in relation to photosynthesis and show that it is important for optimal growth in fluctuating light due to its effect on the light harvesting and PQ pool redox state balances of the model alga *Chlamydomonas* (summary see Supplementary Table 1).

Analysis of the redox state of the PQ pool using complementary chlorophyll fluorescence approaches indicated that the PQ pool was more reduced in c_{6A}-KO, and this was particularly marked in the high light periods of DISCO Light. JTS measurements of the redox state of the b hemes of the cytochrome b_6f complex in response to light exposure showed disturbance of the redox kinetics in the c_{6A}-KO that were consistent with a more reduced PQ pool in c_{6A}-KO. Addition of DCMU or MV, which would be expected to lead to PQ oxidation, or DBMIB, which leads to PQ reduction, revealed effects consistent with an increased pre-reduction of the b hemes in c_{6A}-KO through a more reduced PQ pool. c_{6A}-KO exhibited no change in the proton motive force, measured by electrochromic shift, rendering unlikely a direct effect of c_{6A} on the Q cycle. Under the high light periods of DISCO Light, c_{6A}-KO showed a dramatic decrease in PSII quantum yield compared to WT, which would also be consistent with a more reduced PQ pool. The c_{6A}-KO did not exhibit any significant difference in PSI quantum yield under DISCO light, although it showed a decrease in donor-side limitation of PSI, again consistent with a more reduced PQ pool. c_{6A}-KO also showed some indication of increased acceptor-side limitation.

c_{6A}-KO exhibited an increase in the non-regulated dissipation of excitation energy [Y(NO)] and regulated non-photochemical quenching [Y(NPQ)] with the onset of DISCO Light, as well as a relative increase in the amount of the NPQ effector LHCSR3 protein and degradation products of PsbA after two and four days of growth in DISCO Light, respectively. These results indicate an increased tendency to photoinhibition in c_{6A}-KO, which might be accounted for by the increased PQ reduction. A general impairment of PSII function or repair in the absence of any other effects from loss of c_{6A} might be expected to lead to a decreased reduction in PQ. This suggests that the PSII impairment is a consequence of the increased PQ reduction resulting from loss of c_{6A}, rather than the primary result of c_{6A} loss.

What might cause the increase in reduction of the PQ pool in the absence of c_{6A}? The 77K chlorophyll fluorescence data indicated a significantly lower PSI/PSII fluorescence ratio in c_{6A}-KO compared to WT under photomixotrophic conditions, and this difference was amplified under DISCO Light conditions. In BN-PAGE, a band corresponding to a LHC-less PSI was more

prominent in c_{6A}-KO than WT and c_{6A}-KO showed a lower PSI/PSII fluorescence value than the

wild type under both state 1 and state 2 conditions. These data are consistent with the photosystems

being preferentially in the equivalent of state 1 in c_{6A}-KO compared to WT. However, there was

no evidence for the c_{6A}-KO being permanently 'locked' in either state 1 or state 2.

Overall, it therefore seems likely that c_{6A}-KO is disturbed in the light harvesting balance between

PSI and PSII, in favor of PSII, especially under DISCO Light. This leads to a more reduced PQ

pool, and consequent photoinhibition, with impaired growth of c_{6A}-KO in DISCO conditions. c_{6A}

is likely to be located in the thylakoid lumen and has been shown to interact with the lumen thiol

oxidase LTO1 (21), which is closely involved in thiol-based regulation of the STT7 (STN7 in

Arabidopsis) state transition kinase, keeping STT7 oxidized (22). STT7 is active only when

oxidized and, in response to a reduced PQ pool, catalyzes phosphorylation of LHCII and causing

LHCII to move from PSII to PSI. c_{6A} has been proposed to catalyze the formation of disulfide

bridges in the lumen (10, 11), transferring electrons to plastocyanin, with which it has been shown

to react (19). Although we cannot exclude a more complex explanation, with the shift in PSI/PSII

balance and change in state of the PQ pool being indirect consequences of loss of c_{6A}, we propose

that c_{6A} contributes to oxidation of STT7. c_{6A} is not essential for the oxidation as state transitions

remain in c_{6A}-KO. However, it may be that c_{6A} assists in the transfer of electrons from STT7 to

LTO1 (with which it interacts (21)) or another acceptor. Thus, in the absence of c_{6A}, STT7 would

be more reduced than in WT, leading to increased light harvesting at PSII and a more reduced PQ

pool compared to WT. This model could be tested in the future by making substitutions to the

conserved cysteine residues in c_{6A}. If correct, this will be a significant contribution to our

understanding of the complex thiol-based regulatory network of the thylakoid lumen of

photosynthetic organisms (44). Furthermore, the demonstration of the importance of c_{6A} for growth

under fluctuating light conditions presents a significant contribution to understand how organisms

cope with rapidly changing light conditions (45, 46). The connection of c_{6A} to state transitions also

make it a significant target for optimizing crop productivity.

Materials and Methods

Growth conditions

Chlamydomonas strains were maintained on Tris-Acetate-Phosphate (TAP) (47) medium with

Hutner's trace element solution (48) supplemented with 1.5% agar at 30 $\mu mol_{photons}$ m^{-2} s^{-1} and

25 °C. For longer-term storage cultures were kept on TAP slants in ambient light at room

temperature. Liquid cultures were grown at 30 $\mu mol_{photons}$ m^{-2} s^{-1}, 25 °C and 120 rpm in AlgaeTron

AG130 or AG230 incubators (Photon System Instruments) unless otherwise specified.

For experimental cultures, triplicates of 30 mL medium (Tris-phosphate (TP) medium (47) for

photoautotrophic, and TAP for photomixotrophic conditions) were inoculated with

Chlamydomonas strains to OD$_{750nm}$ = 0.1. The cultures were then subjected to the experimental

conditions (Table 1). All experimental cultures were subcultured at least twice in standard

conditions before exposure to experimental conditions.

Table 1: Experimental light conditions in the AlgaeTron AG130 and AG230 growth chambers.

Name	Light regime
Standard growth light	continuous 30 $\mu mol_{photons}$ m^{-2} s^{-1}
High light	continuous 700 $\mu mol_{photons}$ m^{-2} s^{-1}
Darkness	continuous 0 $\mu mol_{photons}$ m^{-2} s^{-1}
2 min on/off	alternating 2 min 700 $\mu mol_{photons}$ m^{-2} s^{-1} & 2 min darkness
12 s on/off	alternating 12 s 700 $\mu mol_{photons}$ m^{-2} s^{-1} & 12 s darkness
DISCO Light	alternating 2 min darkness & 2 min of 12 s 700 $\mu mol_{photons}$ m^{-2} s^{-1}/ 12 s off (Fig. 1d)

491
492

Algal strains

The cytochrome c_{6A} knock-out strain "c_{6A}-KO" (CC-1883 background) was generated as previously

described (32) using the gRNAs 5'-ATTTACAGTTAGTGCATCCACTTG-3' and

5'-TCCGAGGGTAAGCAAGCACGGCAC-3' to target the 5'UTR and Exon III for Cpf1-

mediated double strand breaks, respectively. A repair template with the following sequence was

provided at the same time to introduce a multiple stop codon (lowercase letters):

5'-TCATGCTCGGCCAGTTTTCTTTGATTTCAAATCCTCGCTAATTTActaattaactaaCAGCT

TCCTTGCTTCATGGACCCAGTAACGGTGCATGGGTACGCG-3'.

The c_{6A}-overexpression strain c_{6A}-OE was created by transforming c_{6A}-KO with the constructs

pDKM-000 by electroporation as described before (49) and specified in the Supplementary

Methods.

Protein extraction and Western Blot

A known volume of *Chlamydomonas* cells at a desired growth stage was harvested by

centrifugation at 1000 xg for 5 min. The pellet was subsequently resuspended in RIPA lysis buffer

with freshly added PMSF (1mM final concentration) to reach a calculated OD_{750nm} of 50. The

chlorophyll content was measured as above, but here 10 µL of sample were mixed with 40 µL of

acetone and 1 mL of 80% acetone, resulting in a dilution factor D of 105. Protein samples were

then adjusted to the same total chlorophyll concentration with RIPA buffer and frozen at -20 °C for

at least 1.5 h or overnight, followed by thawing on ice to complete cell lysis. Protein concentration

was determined using a Bradford assay (Merck) according to manufacturer's instructions diluting

the protein samples 5- to 10-fold.

For SDS-PAGE, all samples were diluted to 1 µg/µL protein with RIPA buffer before being mixed

with 4x Lämmli buffer and incubated for 30 min (50 °C, 600 rpm). After centrifuging the samples

(12000 xg, 1 min) to remove insoluble material, samples were loaded to the desired amount into

wells of polyacrylamide gels (Mini-PROTEAN TGX Stain-Free, BioRad). PageRuler Plus (7 µL,

Thermo Scientific) was used as size marker. Electrophoresis was carried out at 4 °C and 200 V for

about 50 min. The gel was imaged in a ChemiDoc MP imaging system (BioRad) using the Stain

Free setting and 5 min activation time. Some gels were stained with Pierce™ Silver Stain for Mass

Spectrometry (Thermo Scientific).

Transfer of protein from polyacrylamide gels onto polyvinylidene fluoride (PVDF) membranes was

carried out using the iBlot dry transfer apparatus, following the manufacturer's instructions and

using program P3 for 5 min. Membranes were then probed for successful transfer with PonceauS

(Thermo Scientific) as per manufacturer's instructions and imaged with the ChemiDoc MP imaging

system (BioRad). Following the PonceauS destaining, membranes were blocked in 5% milk in Tris-

buffered saline with 0.1% Tween 20 (TBS-T) for 1 h at room temperature with gentle shaking.

After washing membranes once with TBS-T to remove remnants of blocking agent, membranes

were incubated with specific primary antibody solutions in 5% milk TBS-T overnight at 4 °C (PsaB

(AS10695, Agrisera) at 1:1000, PsbA D1 (AS05084, Agrisera) at 1:10000, LHCSR3 (AS142766,

Agrisera) at 1:2000). Subsequently, membranes were washed with TBS-T (5x 5 min) and incubated

for 1.5 h at room temperature with goat anti-rabbit horseradish peroxidase-linked secondary

antibody in 5% milk TBS-T (1:10000, 1706515, BioRad). After washing membranes thrice in TBS-

T and once in TBS (10 min each), they were incubated with chemiluminescence substrate

(SuperSignal™ West Pico PLUS Chemiluminescent Substrate, Thermo Scientific) and imaged

using a ChemiDoc MP imaging system (BioRad).

<u>Low temperature chlorophyll (77K) fluorescence</u>

77K fluorescence emission spectra were recorded with a QE Pro spectrophotometer (Ocean Optics,

Dunedin, FL, USA). Cell samples (200 µL) frozen in liquid N_2 were excited with blue light (440

nm). Raw spectra were normalized to the photosystem II (PSII) fluorescence peak at 685 nm.

Chlamydomonas cells were harvested and total chlorophyll concentration was adjusted to 5 µg/mL

with fresh medium (TP or TAP). Cells were then frozen in liquid N_2 for measurement.

For the state transition experiment, *Chlamydomonas* cells grown in TAP under standard conditions

were adjusted to 5 µg/mL total chlorophyll with fresh medium. Anoxia (state 2 inducing) samples

were adjusted with medium containing glucose (final concentration 20 mM) and Glucose Oxidase

(final concentration 50 U/mL) that was purged with N2 for 3 min. The final samples were again

purged with N_2 before being sealed. Oxia (state 1 inducing) samples were adjusted with unchanged

medium and left unsealed. All samples were then shaken for 30 min in darkness before being frozen

in liquid N_2 for fluorescence measurement.

<u>Blue native PAGE and second dimension SDS-PAGE</u>

Thylakoid membranes from *Chlamydomonas* cultures were isolated over a discontinuous Percoll

gradient as described before (50). BN-PAGE was then carried out largely as previously (51).

Thylakoid samples were diluted with resuspension buffer (25 mM Bis-Tris/HCl (pH 7.0), 20%

(w/v) glycerol, 0.25 mg/ml Pefabloc, and 10 mM sodium fluoride) to a chlorophyll concentration

of 0.53 µg/µL and to a total of 6 µg chlorophyll. An equal volume of β -dodecyl maltoside (β -DM,

2% in resuspension buffer) was added to each sample and the thylakoid membranes were

solubilized by gentle pipetting for 2 min. After centrifugation (4 °C, 18000 xg, 20 min) samples

were mixed with a one-tenth volume of sample loading buffer (100 mM Bis-Tris/HCl (pH 7.0), 0.5

M ACA, 30% (w/v) sucrose and 50 mg/ml Serva Blue G) and loaded into the wells of a

NativePAGE 3-12% Bis-Tris protein gel (Invitrogen). Electrophoresis was carried out with specific

anode buffer (50 mM Bis-Tris/HCl, pH 7.0) and cathode buffer containing Serva Blue G dye (50

mM Tricine, 15 mM Bis-Tris/HCl, pH 7.0, and 0.01% Serva Blue G) at 4 °C for 20 min at 75 V,

30 min at 100 V, 30 min at 125 V, 60 min at 150 V, 30 min at 175 V and 60 min at 200 V. After

the 30 min at 125 V, the blue cathode buffer was replaced by the same cathode buffer but without

Serva Blue G. The last step was prolonged to 5 h if additional separation of bands of interest was

needed.

Gel electrophoretic separation of complexes in BN-PAGE lanes on a second dimension SDS-PAGE

was carried out as previously described (51, 52), although 12% polyacrylamide gels were used

here. Gels were stained with Pierce™ Silver Stain (Thermo Scientific).

The predicted mobility in the gel (R_f) values of known thylakoid membrane complex components

were calculated using their known molecular weight and a linear regression of the distance of the

protein standard in relation to the running front (r^2 = 0.98). The 2D-PAGE spots were compared

with the R_f values and with published BN-PAGE analyses from *Chlamydomonas* (53–55) to assign

the bands and proteins.

<u>Chlorophyll fluorescence</u>

Chlorophyll fluorescence induction (OJIP) of *Chlamydomonas* was measured with a FL3500

fluorometer (Photon System Instruments). Prior to measurement, cells were resuspended in TAP

and total chlorophyll concentration was adjusted to 20 µg/mL. After recovery under standard

growth conditions for at least 30 min, chlorophyll concentration was readjusted to 5 µg/mL. After

15 min incubation in darkness, fluorescence induction in response to red light (984 $µmol_{photons}$ m^{-2}

s^{-1}, 45% A-voltage) was recorded for 120 s. Raw fluorescence data was double normalized to

fluorescence at time point 0 and the maximum fluorescence value to compare the kinetics between

samples (37).

Chlorophyll fluorescence of *Chlamydomonas* was measured with a pulse amplitude-modulated

spectrometer (Dual-PAM-100, Walz). Prior to measurements, cells were resuspended in TAP and

the chlorophyll concentration was adjusted to 20 µg/mL. After recovery under standard growth

conditions for at least 30 min, measurements were performed in a quartz cuvette at 25 °C with

stirring and samples were initially incubated in darkness for 5 min before the blue fluorescence

measuring light (Intensity 12, 28 $µmol_{photons}$ m^{-2} s^{-1}) was turned on. Saturating pulses (Intensity 5,

5000 $µmol_{photons}$ m^{-2} s^{-1}, 500 ms width) for PSII parameter determination were applied with stirring

turned off. DISCO Light conditions in the DUAL-PAM-100 were achieved by alternating 2 min

darkness followed by 2 min consisting of 12 s high light (690 $µmol_{photons}$ m^{-2} s^{-1}) and 12 s darkness

alternations. Saturating pulses were applied in the middle of each 12 s phase or three times during

the 2 min dark phases. F' was recorded as the fluorescence yield during illumination with the actinic

light treatment before application of a saturating pulse. Photosynthetic parameters were calculated

as described previously (41).

<u>Cytochrome b_6f redox kinetics</u>

Cytochrome *f* and *b* heme redox kinetics were determined in *Chlamydomonas* cells by

deconvoluting absorbance changes at 546, 554, 563, and 573 nm that were measured using a JTS-

10 pump probe spectrophotometer (BioLogic) and appropriate 10 nm FWHM (full width at half

maximum) interference filters (Edmund Optics). BG39 filters (Schott) were used to shield the light detectors from scattered light. Prior to measurements, cells were resuspended in fresh medium to a total chlorophyll concentration of 20 µg/mL. After recovery under standard growth conditions for at least 30 min, Ficoll 400 was added to a final concentration of 10%. For some samples, DCMU, methyl viologen, or DBMIB were added to a final concentration of 10 µM, 10 mM, or 20 µM respectively, at the same time as Ficoll. Cells were dark-adapted for 3 min before measurement with each interference filter and then illuminated with 5 s of green light (500 $\mu mol_{photons}$ m^{-2} s^{-1}). Flashes of white detection light were administered during 300 µs dark intervals in actinic illumination. Every recording was corrected for any detection light impact by subtracting a trace with actinic light turned off. Deconvolution was performed with the JTS-10 software by drawing a baseline between 573 and 546 nm. During analysis, the signal-to-noise ratio was improved with a Savitzky-Golay filter (56).

Acknowledgments

We thank Dr. K. Geissler (University of Cambridge) for the provision of the pCM0 MoClo parts and Dr. J. Walter (University of Cambridge) for assistance with blue native PAGE. We are also grateful to Dr. M. Jokel (University of Turku) for discussions on cell culture. All photosynthesis experiments were performed within the PHOTOSYN infrastructure at the University of Turku. We gratefully acknowledge the support of this work by: the Gates Cambridge Trust with support from the Benn W Levy Trust, and the Professor John Glover Memorial Fund (to DK); the Novo Nordisk Foundation (PhotoCat, project no. NNF20OC0064371, to YA; and Photo-e-microbes project no. NNF22OCOO79717 to LTW); the Research Council of Finland no. 355335 to YA, no. 354876 to LN; the Biotechnology and Biological Sciences Research Council (BB/M011194/1 to AS, BB/M011194/1 to JML, BB/R506163/1 to AF, and BB/Y001370/1 to CJH), and Trinity Hall, Cambridge (to JML).

Author Contributions: DK conceived the project, designed and performed experiments under the supervision of CJH and YA, and analyzed data. DK, LTW, LN, and YA developed the biophysical part of the project. LTW and LN supervised the biophysical measurements and assisted with data analysis and interpretation. AF created the knock-out mutant line under the supervision of AM. JML and AS assisted with data collection, analysis and interpretation. AM,

639 YA, and CJH secured funding, assisted with data interpretation, and provided resources. DK

640 wrote the manuscript with comments and input from all authors.

641 **Competing Interest Statement:** The authors declare no competing interests.

642
643
References

1. M. Weigel, *et al.*, Plastocyanin is indispensable for photosynthetic electron flow in Arabidopsis
thaliana. *J. Biol. Chem.* **278**, 31286–31289 (2003).

2. G. A. Peschek, C. Obinger, M. Paumann, The respiratory chain of blue-green algae
(cyanobacteria). *Physiol. Plant.* **120**, 358–369 (2004).

3. R. Gupta, Z. He, S. Luan, Functional relationship of cytochrome c6 and plastocyanin in
Arabidopsis. *Nature* **417**, 567–571 (2002).

4. J. Wastl, D. S. Bendall, C. J. Howe, Higher plants contain a modified cytochrome c6. *Trends*
Plant Sci. **7**, 244–245 (2002).

5. J. Wastl, S. Purton, D. S. Bendall, C. J. Howe, Two forms of cytochrome c6 in a single eukaryote.
Trends Plant Sci. **9**, 474–476 (2004).

6. I. Nimmo, Location and function of cytochrome c6A. *PhD Thesis Univ. Camb.* (2011).

7. C. J. Howe, R. H. Nimmo, A. C. Barbrook, D. S. Bendall, "Cytochrome c6A of Chloroplasts" in
Cytochrome Complexes: Evolution Structures, Energy Transduction and Signaling, Advances in
Photosynthesis and Respiration., W. A. Cramer, T. Kallas, Eds. (Springer Science+Business
Media, 2016).

8. U. Hani, *et al.*, A complex and dynamic redox network regulates oxygen reduction at photosystem
I in Arabidopsis. *Plant Physiol.* (2024). https://doi.org/10.1093/plphys/kiae501.

9. B. A. Slater, The ancestry and function of cytochrome c6A. *PhD Thesis Univ. Camb.* (2020).

10. C. J. Howe, B. G. Schlarb-Ridley, J. Wastl, S. Purton, D. S. Bendall, The novel cytochrome c6 of
chloroplasts: a case of evolutionary bricolage? *J. Exp. Bot.* **57**, 13–22 (2006).

11. B. G. Schlarb-Ridley, R. H. Nimmo, S. Purton, C. J. Howe, D. S. Bendall, Cytochrome c 6A is a
funnel for thiol oxidation in the thylakoid lumen. *FEBS Lett.* **580**, 2166–2169 (2006).

12. H. Chida, *et al.*, Crystal structure of oxidized cytochrome c 6A from Arabidopsis thaliana. *FEBS*
Lett. **580**, 3763–3768 (2006).

13. J. A. R. Worrall, B. F. Luisi, B. G. Schlarb-Ridley, D. S. Bendall, C. J. Howe, Cytochrome c 6A:
discovery, structure and properties responsible for its low haem redox potential. *Biochem. Soc.*
Trans. **36**, 1175–1179 (2008).

14. J. M. Mason, D. S. Bendall, C. J. Howe, J. A. R. Worrall, The role of a disulfide bridge in the
stability and folding kinetics of Arabidopsis thaliana cytochrome c6A. *Biochim. Biophys. Acta*
BBA - Proteins Proteomics **1824**, 311–318 (2012).

15. J. A. R. Worrall, *et al.*, Modulation of Heme Redox Potential in the Cytochrome c 6 Family. *J.*
Am. Chem. Soc. **129**, 9468–9475 (2007).

16. B. Slater, D. Kosmützky, R. E. R. Nisbet, C. J. Howe, The Evolution of the Cytochrome c6
Family of Photosynthetic Electron Transfer Proteins. *Genome Biol. Evol.* **13**, 1–12 (2021).

17. F. P. Molina-Heredia, *et al.*, A new function for an old cytochrome? *Nature* **424**, 33–34 (2003).

18. J. Wastl, *et al.*, Redox properties of Arabidopsis cytochrome c6 are independent of the loop
extension specific to higher plants. *Biochim. Biophys. Acta BBA - Bioenerg.* **1657**, 115–120
(2004).

19. M. J. Marcaida, *et al.*, Structure of Cytochrome c6A, a Novel Dithio-cytochrome of Arabidopsis
thaliana, and its Reactivity with Plastocyanin: Implications for Function. *J. Mol. Biol.* **360**, 968–
977 (2006).

20. P. Pesaresi, *et al.*, Mutants, overexpressors, and interactors of arabidopsis plastocyanin isoforms:
Revised roles of plastocyanin in photosynthetic electron flow and thylakoid redox state. *Mol.*
Plant **2**, 236–248 (2009).

21. Y. Lu, *et al.*, Identification of Potential Targets for Thylakoid Oxidoreductase AtVKOR/LTO1 in
Chloroplasts. *Protein Pept. Lett.* **22**, 219–225 (2015).

22. J. Wu, *et al.*, Functional redox links between lumen thiol oxidoreductase1 and serine/threonine-
protein kinase STN7. *Plant Physiol.* **186**, 964–976 (2021).

23. P. J. Gollan, A. Trotta, A. A. Bajwa, I. Mancini, E.-M. Aro, Characterization of the Free and
Membrane-Associated Fractions of the Thylakoid Lumen Proteome in Arabidopsis thaliana. *Int.*
J. Mol. Sci. **22**, 8126 (2021).

24. J. Beltrán, *et al.*, Specialized Plastids Trigger Tissue-Specific Signaling for Systemic Stress
Response in Plants. *Plant Physiol.* **178**, 672–683 (2018).

25. M. Janssen, *et al.*, Efficiency of light utilization of Chlamydomonas reinhardtii under medium-
duration light/dark cycles. *J. Biotechnol.* **78**, 123–137 (2000).

26. P. J. Graham, B. Nguyen, T. Burdyny, D. Sinton, A penalty on photosynthetic growth in
fluctuating light. *Sci. Rep.* **7**, 1–11 (2017).

27. L. Nikkanen, *et al.*, PGR5 is needed for redox-dependent regulation of ATP synthase both in
chloroplasts and in cyanobacteria. [Preprint] (2025). Available at:
https://www.biorxiv.org/content/10.1101/2024.11.03.621747v2 [Accessed 29 March 2026].

28. M. Jokel, X. Johnson, G. Peltier, E. Aro, Y. Allahverdiyeva, Hunting the main player enabling
Chlamydomonas reinhardtii growth under fluctuating light. *Plant J.* **94**, 822–835 (2018).

29. C. J. Steen, A. Burlacot, A. H. Short, K. K. Niyogi, G. R. Fleming, Interplay between LHCSR
proteins and state transitions governs the NPQ response in Chlamydomonas during light
fluctuations. *Plant Cell Environ.* **45**, 2428–2445 (2022).

30. T. Roach, LHCSR3-Type NPQ Prevents Photoinhibition and Slowed Growth under Fluctuating
Light in Chlamydomonas reinhardtii. *Plants* **9**, 1604 (2020).

31. W. J. Nawrocki, *et al.*, Chlororespiration Controls Growth Under Intermittent Light. *Plant
Physiol.* **179**, 630–639 (2019).

32. A. Ferenczi, D. E. Pyott, A. Xipnitou, A. Molnar, Efficient targeted DNA editing and replacement
in Chlamydomonas reinhardtii using Cpf1 ribonucleoproteins and single-stranded DNA. *Proc.
Natl. Acad. Sci.* **114**, 13567–13572 (2017).

33. E. G. Andrizhiyevskaya, *et al.*, Origin of the F685 and F695 fluorescence in Photosystem II.
Photosynth. Res. **84**, 173–180 (2005).

34. J. Kargul, J. Nield, J. Barber, Three-dimensional reconstruction of a light-harvesting complex I-
photosystem I (LHCI-PSI) supercomplex from the green alga Chlamydomonas reinhardtii:
Insights into light harvesting for PSI. *J. Biol. Chem.* **278**, 16135–16141 (2003).

35. S. Z. Tóth, G. Schansker, R. J. Strasser, A non-invasive assay of the plastoquinone pool redox
state based on the OJIP-transient. *Photosynth. Res.* **93**, 193 (2007).

36. B. A. Diner, Dependence of the deactivation reactions of Photosystem II on the redox state of
plastoquinone pool a varied under anaerobic conditions. Equilibria on the acceptor side of
Photosystem II. *Biochim. Biophys. Acta BBA - Bioenerg.* **460**, 247–258 (1977).

37. H. M. Kalaji, *et al.*, Frequently asked questions about chlorophyll fluorescence, the sequel.
Photosynth. Res. **132**, 13–66 (2017).

38. P. Joliot, A. Joliot, Proton pumping and electron transfer in the cytochrome bf complex of algae.
Biochim. Biophys. Acta BBA - Bioenerg. **849**, 211–222 (1986).

39. F. Buchert, M. Scholz, M. Hippler, Electron transfer via cytochrome b 6 f complex displays
sensitivity to antimycin A upon STT7 kinase activation. *Biochem. J.* **479**, 111–127 (2022).

40. A. Malnoë, F.-A. Wollman, C. de Vitry, F. Rappaport, Photosynthetic growth despite a broken Q-
cycle. *Nat. Commun.* **2**, 301 (2011).

41. C. Klughammer, U. Schreiber, Complementary PS II quantum yields calculated from simple
fluorescence parameters measured by PAM fluorometry and the Saturation Pulse method. *PAM
Appl. Notes* **1**, 27–35 (2008).

42. W. J. Nawrocki, *et al.*, Molecular origins of induction and loss of photoinhibition-related energy
dissipation q I. *Sci. Adv.* **7**, 1–13 (2021).

43. M. Ballottari, *et al.*, Identification of pH-sensing Sites in the Light Harvesting Complex Stress-
related 3 Protein Essential for Triggering Non-photochemical Quenching in Chlamydomonas
reinhardtii. *J. Biol. Chem.* **291**, 7334–7346 (2016).

44. D. Hoh, J. E. Froehlich, D. M. Kramer, Redox regulation in chloroplast thylakoid lumen: The pmf
changes everything, again. *Plant Cell Environ.* **47**, 2749–2765 (2024).

45. N. A. Eckardt, *et al.*, Lighting the way: Compelling open questions in photosynthesis research.
Plant Cell **14**, 1–16 (2024).

46. S. P. Long, *et al.*, Into the Shadows and Back into Sunlight: Photosynthesis in Fluctuating Light.
Annu. Rev. Plant Biol. **73**, 617–648 (2022).

47. D. S. Gorman, R. P. Levine, Cytochrome f and plastocyanin: their sequence in the photosynthetic
electron transport chain of Chlamydomonas reinhardi. *Proc. Natl. Acad. Sci.* **54**, 1665–1669
(1965).

48. S. H. Hutner, L. Provasoli, A. Schnatz, C. P. Haskins, Some Approaches to the Study of the Role
of Metals in the Metabolism of Microorganisms. *Proc. Am. Philos. Soc.* **94**, 152–170 (1950).

49. P. Crozet, *et al.*, Birth of a Photosynthetic Chassis: A MoClo Toolkit Enabling Synthetic Biology
in the Microalga *Chlamydomonas reinhardtii*. *ACS Synth. Biol.* **7**, 2074–2086 (2018).

50. C. B. Mason, T. M. Bricker, J. V. Moroney, A rapid method for chloroplast isolation from the
green alga Chlamydomonas reinhardtii. *Nat. Protoc.* **1**, 2227–2230 (2006).

51. S. Järvi, M. Suorsa, V. Paakkarinen, E. M. Aro, Optimized native gel systems for separation of
thylakoid protein complexes: Novel super- and mega-complexes. *Biochem. J.* **439**, 207–214
(2011).

52. M. Rantala, V. Paakkarinen, E. M. Aro, Analysis of thylakoid membrane protein complexes by
blue native gel electrophoresis. *J. Vis. Exp.* **2018**, 1–6 (2018).

53. E. Devadasu, *et al.*, Long- and short-term acclimation of the photosynthetic apparatus to salinity
in Chlamydomonas reinhardtii. The role of Stt7 protein kinase. *Front. Plant Sci.* **14**, 1–17
(2023).

54. J. G. García-Cerdán, *et al.*, A thylakoid membrane-bound and redox-active rubredoxin (RBD1)
functions in de novo assembly and repair of photosystem II. *Proc. Natl. Acad. Sci. U. S. A.* **116**,
16631–16640 (2019).

55. B. Spaniol, *et al.*, Complexome profiling on the Chlamydomonas lpa2 mutant reveals insights
into PSII biogenesis and new PSII associated proteins. *J. Exp. Bot.* **73**, 245–262 (2022).

56. M. J. E. Savitzky, A.; Golay, Smoothing and Differentiation of Data. *Anal Chem* **36**, 1627–1639
(1964).

57. C. A. Kerfeld, H. P. Anwar, R. Interrante, S. Merchant, T. O. Yeates, The Structure of Chloroplast
Cytochrome c6 at 1.9 Å Resolution: Evidence for Functional Oligomerization. *J. Mol. Biol.* **250**,
627–647 (1995).