A conserved photosynthetic cytochrome enhances growth in fluctuating light

Darius Kosmützky[1,‡,*], Laura T Wey[2], Lauri Nikkanen[2], Aron Ferenczi[3], Joshua M Lawrence[1,4], Alberto Scarampi[1,‡‡], Attila Molnar[3], Yagut Allahverdiyeva[2], Christopher J Howe[1,*]

1: Department of Biochemistry, University of Cambridge, Cambridge CB2 1QW, UK

2: Molecular Plant Biology, Department of Life Technologies, University of Turku, Turku 20014, Finland

3: Institute of Molecular Plant Sciences, University of Edinburgh, Edinburgh EH9 3BF, UK

4: Yusuf Hamied Department of Chemistry, University of Cambridge, Cambridge CB2 1EW, UK

‡: Present address: Sainsbury Laboratory, University of Cambridge, Cambridge CB2 1LR, UK

‡‡: Present address: School of Life Sciences, University of Warwick, Coventry CV4 7AL, UK

*: Corresponding authors

Abstract

Having at least one of the small soluble electron carriers cytochrome c_6 (c_6) and plastocyanin is indispensable for photosynthesis. It was believed that c_6 had been lost from plants until the identification of a ubiquitous and highly conserved homologue, now named cytochrome c_{6A} (c_{6A}), led to a paradigm shift. However, c_{6A} was soon shown to be unable to replace c_6 (or plastocyanin) functionally in the photosynthetic electron transport chain, despite their significant structural homology. The function of c_{6A}, and why it is apparently universal in plants and green algae, remain unknown. Here, we show that, in the green alga *Chlamydomonas reinhardtii*, c_{6A} confers a growth advantage under fluctuating light and is crucial for maintaining the light harvesting balance between photosystems I and II in photomixotrophic conditions. We show that in the absence of c_{6A}, the light harvesting balance shifts towards PSII, leading to a more reduced plastoquinone pool and increased photooxidative stress. This study provides new insights into how photosynthetic organisms acclimatize to stressful light conditions, indicating that c_{6A} is important for this adaptation. These findings provide a basis for further mechanistic studies on a hypothesized role for c_{6A} in thiol-based redox regulation in the thylakoid lumen, with implications for photoprotection mechanisms such as state transitions.

Introduction

The photosynthetic electron transport chain (PETC) is essential for nearly all life on Earth (Fig. 1a).

At least one of the soluble electron carriers plastocyanin and cytochrome c_6 is indispensable for the

PETC to function (1, 2). A close homologue of c_6, cytochrome c_{6A}, was first discovered in

Arabidopsis thaliana and later in other plants and eukaryotic algae (3, 4). Whereas plants lost the gene

for c_6 and contain only c_{6A}, some eukaryotic algae such as *Chlamydomonas reinhardtii*

(*Chlamydomonas*) contain genes for both (5), although c_{6A} is expressed at a low level compared to c_6

or plastocyanin (6, 7). Since its discovery, c_{6A} has been hypothesized to have various functions,

including replacement of plastocyanin (3), acting as a safety valve for excess electrons of

photosynthesis (7), regulation of generation of reactive oxygen species (ROS) in PSI (8), and being

involved in high light signalling (9) or thiol-based redox regulation (10, 11). However, the function of

cytochrome c_{6A} remains elusive (7).

While c_6 and c_{6A} are homologues, there are structural differences between the two (Fig. 1b). c_{6A}

contains a 'loop insertion peptide' (LIP), an insertion with a conserved disulfide bridge in a surface-

located and well-defined loop. The LIP is absent from c_6 (10). Assessing the stability of various

mutant proteins showed that the LIP is structurally important (12, 13). While a catalytic role for the

conserved disulfide bridge within the LIP is uncertain, it is predicted to contribute to the protein's

structural stability(13, 14). Furthermore, differences in heme environment contribute to c_{6A} (+71 mV

(15)) having a 254 mV more negative midpoint potential than plastocyanin or c_6 (+325 mV (15)).

Phylogenetic analyses revealed that c_{6A} represents a monophyletic group distinct from c_6 and likely

originates from cyanobacterial cytochrome c_{6B}, another homologue of c_6 that has a similar midpoint

potential to c_{6A} but lacks the LIP (16). The sequence of c_{6A} is highly conserved across the green

lineage, as with plastocyanin, suggesting an important role for c_{6A} within photosynthetic eukaryotes

(16). However, the large difference in midpoint potentials renders it highly unlikely that c_{6A} acts as a

functional replacement for plastocyanin in plants and algae, as the former cannot accept electrons

from the cytochrome b_6f complex (Fig. 1a). Further, plastocyanin was found to be indispensable even

when c_{6A} was overexpressed in *A. thaliana*[1]. Additionally, the surfaces of c_6 and plastocyanin are

strongly negatively charged, whereas c_{6A} has a strongly positive surface charge (17). Together with

the difference in isoelectric points (plastocyanin: 5.1, c_{6A}: 4.1 in *A. thaliana* (18)) this indicates

different interactions.

Interactions involving c_{6A} further highlight its distinct role (Fig. 1a). It cannot effectively reduce PSI

in vitro, making direct interaction between c_{6A} and PSI unlikely (17). Additionally, c_{6A} has been

shown to reduce plastocyanin *in vitro* (19) and weak interaction (K_d = 2.01 mM) was detected with

plastocyanin using NMR titrations (6), consistent with a transient interaction. Yeast Two-Hybrid

assays also revealed interactions of c_{6A} with itself (20), the lumen thiol oxidase LTO1 (21), and the

redox-regulated immunophilin FKBP13 (3). For LTO1 an involvement in regulating state transitions

via the kinase STT7 has been described (22). Interaction with LTO1 and FKBP13 suggests a luminal

localization for c_{6A}, consistent with its detection in the soluble fraction of the thylakoid lumen in *A.*

thaliana (23). In *Chlamydomonas*, however, only chloroplastic localization has been confirmed (6). In

A. thaliana, c_{6A} was also found to be uniquely localized to sensory plastids in the epidermis and

vascular parenchyma rather than mesophyll chloroplasts (24).

Photosynthetic organisms require flexibility in their photosynthetic apparatus because they grow

under a diverse set of environmental conditions. These include varying light conditions from diurnal

cycles to more rapid fluctuations caused by the sun, clouds, canopy, and in the case of

Chlamydomonas, also soil and water movements. For *Chlamydomonas*, exposure to a 12 s on/12 s off

cycle with light intensities of 600-700 μmol photons m^{-2} s^{-1} in photoautotrophic conditions resulted in

growth rates reaching only about 30% of those achieved under continuous illumination (25).

Additionally, a growth penalty was revealed under fluctuating light with frequencies in the range of

10-100 s on/off (26). In mutant assays using *Chlamydomonas*, various fluctuating light regimes have

been employed to determine the function of proteins associated with photosynthesis. Prominent

members include Proton Gradient Regulation 5 (PGR5) and Proton Gradient Regulation Like 1

(PGRL1) proteins, both implicated in cyclic electron flow, its regulation or ATP synthase regulation

(27), as well as flavodiiron proteins (FLVA/B), which facilitate the Mehler-like reaction (28).

Additionally, the qE quenching effector LHCSR3 (29, 30) and plastid terminal oxidase PTOX2 (31)

have been implicated in the response to fluctuating light, as mutants of all of these proteins showed a

growth deficit under fluctuating light conditions.

Here, we present the first in-depth analysis of the connection between c_{6A} and photosynthesis. We

show that this highly conserved protein is important for growth under a harsh fluctuating light regime

we called DISCO Light and helps maintain the redox balance of the plastoquinone (PQ) pool through

its effect on the light harvesting balance between photosystems I (PSI) and II (PSII).

Results

<u>Cytochrome c_{6A} confers a growth advantage under a fluctuating light regime</u>

To investigate the role of c_{6A} in *Chlamydomonas* we used CRISPR/Cpf1 (32) to create a knock-out of

the c_{6A} gene (*CYC4*, Cre16.g670950) in the background strain CC-1883 (WT). The knock-out strain

("c_{6A}-KO") was confirmed through PCR and sequencing (Methods & Supplementary Fig. 1a-d). We

also generated a c_{6A} overexpression (c_{6A}-OE) strain by transforming the c_{6A}-KO strain with a construct

that drives the expression of FLAG-tagged c_{6A} from a *PSAD* promoter. Overexpression was confirmed

by immunoblot and RNAseq analysis (Methods & Supplementary Fig. 1d-e).

To investigate the role of cytochrome c_{6A} in growth, we tested how *Chlamydomonas* strains grow

under various light and trophic regimes. Under continuous light (30 μmol photons m^{-2} s^{-1}), both c_{6A}-

KO and WT strains reached similar maximum optical densities (OD_{750nm}) of around 2 in

photomixotrophic and ~0.6 in photoautotrophic conditions, with no statistically significant differences

(Fig. 1c). We defined continuous light (30 μmol photons m^{-2} s^{-1}) and photomixotrophic conditions as

standard conditions for this study.

We also tested multiple stressful light conditions for their effect on c_{6A}-KO and WT. Only a novel

light regime we named DISCO (Darkness Interrupted by Short COnstant) Light and photomixotrophy

led to an observable phenotypic growth difference (Fig. 1c). DISCO Light was devised to expose

Chlamydomonas to a complex fluctuating light regime covering multiple frequencies, as it consists of

two alternating phases: 2 min of darkness followed by 2 min of alternating 12 s high light pulses of

700 $μmol_{photons}$ m^{-2} s^{-1} and 12 s darkness (Fig. 1d). c_{6A}-KO grew more slowly and reached lower cell

densities than WT under photomixotrophic DISCO conditions (Fig. 1c & e): after one day c_{6A}-KO

cultures showed significantly reduced growth ($P = 0.029$), and by day 3 they were approximately 1.5

times less dense than WT cultures ($P = 0.10$). The growth penalty in c_{6A}-KO was recovered by

overexpressing c_{6A}, with c_{6A}-OE showing significantly higher cell densities than c_{6A}-KO from day 1

($P = 0.024$, Fig. 1c & e). In the DISCO light regime without acetate in the medium (photoautotrophy),

no strain could grow (Supplementary Fig. 2e).

In photomixotrophy and darkness or continuous high light at the same light intensity as in the DISCO

regime (700 $μmol_{photons}$ m^{-2} s^{-1}) WT and c_{6A}-KO grew at similar rates (Supplementary Fig. 2a-b).

Furthermore, fluctuating light regimes corresponding to elements of the DISCO protocol ("2 min high

light & 2 min darkness", and "12 s high light & 12 s darkness"), showed only small differences

between WT and c_{6A}-KO that were not part of a consistent trend under either photomixotrophic or

photoautotrophic conditions (Supplementary Fig. 2c-d).

Figure 1: Cytochrome c_{6A} confers a growth advantage under a fluctuating light regime. a, Schematic of the
photosynthetic electron transport chain with known (black) and putative interactions (purple) of c_{6A} marked. Proton
movement (grey) and the Q-cycle (blue) are shown as arrows. **b,** Overlay of the crystal structures of cytochrome c_6 from
Chlamydomonas (orange, PDB: 1CYJ (33)) and c_{6A} from *A. thaliana* (purple, PDB: 2CE0 (19)). The position of the LIP is
noted with the disulfide bridge formed between the two conserved cysteine residues (yellow sticks). The amino acid residues
at position 52 were found to affect the heme environment contributing to the low midpoint potential of c_{6A} compared to c_6 or
plastocyanin (stick model and labelled with a green diamond) (15). **c,** Optical density (OD_{750nm}) growth curve of c_{6A} mutant
strains and WT under continuous light (30 $\mu mol_{photons}$ m^{-2} s^{-1}) in either photoautotrophy or -mixotrophy and DISCO Light
and photomixotrophy (n = 3 biological replicates ±SD). Statistical testing was carried out with repeated measures ANOVA.
Post hoc pairwise two-sided t-tests at each time point were performed if the interaction effect of time and strain on OD_{750nm}
in the ANOVA was significant ($p_{time:strain} < 0.05$) comparing WT (pink) or c_{6A}-OE (green) against c_{6A}-KO (p < 0.05: *, p ≤
0.1: #). **d,** DISCO Light (Darkness Interrupted by Short COnstant Light) consists of alternating 2 min darkness followed by
2 min of high light pulses (12 s high light (700 $\mu mol_{photons}$ m^{-2} s^{-1}) & 12 s darkness). **e,** Visual differences of the c_{6A} mutant
and WT cultures of at day 3 of growth in DISCO Light.

Measurements of total chlorophyll concentration and acetate consumption confirmed the phenotypic

difference between the strains observed from OD_{750nm} measurements in both standard and DISCO

Light conditions (Supplementary Fig. 2f-h). The correlation between total chlorophyll concentration

and optical density was similar between WT and c_{6A}-KO. This indicated no difference in chlorophyll

concentration per cell between the strains under standard conditions or DISCO Light (Supplementary

Fig. 2g). Additionally, measurements of acetate concentration in the growth medium indicated that

c_{6A}-KO consumed less acetate, likely because of lower cell densities when exposed to DISCO Light,

with differences apparent from day 3 (Supplementary Fig. 2h).

These results indicate that c_{6A} is important for growth of *Chlamydomonas* under the high intensity and

rapidly fluctuating DISCO Light regime and acetate availability in photomixotrophic conditions.

<u>c_{6A} is likely localized in the thylakoid lumen and does not have strongly bound interaction partners</u>

c_{6A} was detected in the soluble fraction of the thylakoid lumen in *A. thaliana* (23). In

Chlamydomonas, however, only chloroplastic localization has been confirmed (6). To determine the

subcellular localization of c_{6A} in *Chlamydomonas* further, we performed immunofluorescence

microscopy on the c_{6A}-FLAG overexpressing strain (c_{6A}-OE) and its background strain (c_{6A}-KO)

grown under standard conditions. In the negative control (c_{6A}-KO), the AlexaFluor488 signal

corresponding to FLAG was weak and non-specific, i.e. appearing throughout the cell including in the

flagella (Supplementary Fig. 3a). In contrast, the c_{6A}-FLAG overexpressing sample (c_{6A}-OE) showed

a more intense (2 to 3-fold pixel intensity) and more localized fluorescence signal, confirming

chloroplast-specific localization of c_{6A} (Supplementary Fig. 3b). Two cells covering the variation of

AlexaFluor488 signal intensity were chosen as representatives to capture cell to cell differences that

could arise from technical variation from fixing and staining or c_{6A}-FLAG expression level

differences between cells. Nevertheless, the absence of AlexaFluor488 signal from cell compartments

other than the chloroplast in c_{6A}-OE compared to c_{6A}-KO supports the FLAG-specificity of the signal.

The detection of anti-FLAG signal in structures resembling pyrenoid-traversing thylakoids supports a

thylakoid luminal localization of the soluble c_{6A} (Supplementary Fig. 3c-d). While it cannot be

excluded that c_{6A} is associated with the thylakoid membrane on the stromal side, the structure of c_{6A}

(Fig. 1b) gives no indication of membrane-association.

To detect possible covalent or strongly bound interaction partners, we performed co-

immunoprecipitation (co-IP) analyses of the c_{6A}-FLAG overexpressing strain (c_{6A}-OE) and the

untransformed background strain (c_{6A}-KO) using anti-FLAG-agarose beads. Immunodetection of

FLAG in the precipitate revealed bands at ~15 kDa in c_{6A}-OE but not in c_{6A}-KO, corresponding to

mature c_{6A}-FLAG (red arrows in Supplementary Fig. 4), with additional FLAG signals that were not

c_{6A}-FLAG specific in the 25-35 kDa range (black arrows in Supplementary Fig. 4). Under both

standard (Supplementary Fig. 4c) and DISCO Light (Supplementary Fig. 4f) conditions, the

immunoprecipitate did not reveal any bands unique to c_{6A}-OE except the c_{6A}-FLAG band, indicating

that no specific interaction partners of c_{6A} were detected.

The combined immunofluorescence microscopy and co-IP results indicate that c_{6A} is likely to be

localized in the thylakoid lumen and are consistent with any hypothesized function relying primarily

on weak or transient interactions.

<u>c_{6A} is important to maintain the light harvesting balance between photosystems I and II in</u>

<u>photomixotrophy</u>

Given the light-dependent growth phenotype observed in DISCO conditions, the likely localization of

c_{6A} in the thylakoid lumen, and c_{6A}'s high sequence similarity to c_6 which is involved in the PETC, we

investigated the photosystems and light harvesting complexes to explore further a potential role of c_{6A}

in photosynthesis.

We recorded low temperature (77K) chlorophyll fluorescence emission spectra. When grown in

continuous light (30 $\mu mol_{photons}$ m^{-2} s^{-1}) under photoautotrophic conditions WT and c_{6A}-KO exhibited

very similar fluorescence spectra (Fig. 2a, PSI/PSII fluorescence ratios 1.25 and 1.3, respectively, $P =$

0.374), consistent with similar growth (Fig. 1c). However, when grown in photomixotrophic

conditions and continuous light (30 $\mu mol_{photons}$ m^{-2} s^{-1}, standard conditions), and despite the two

strains showing similar growth (Fig. 1b), WT maintained a PSI/PSII fluorescence ratio of about 1.2,

while c_{6A}-KO showed a significantly lower ratio (0.96, $P = 0.008$), which was accompanied by a more

prominent CP47-associated peak at 695 nm (34) (Fig. 2a). The PSI/PSII fluorescence ratio was

restored to nearly WT levels in the c_{6A}-OE strain (Fig. 2a). When samples were grown in DISCO

Light and photomixotrophy, the differences seen in standard conditions were amplified. By day 4, the

PSI/PSII fluorescence ratio in c_{6A}-KO dropped drastically to 0.66 in comparison to WT and c_{6A}-OE

(1.05 and 1.33, respectively & $P = 0.027$ and 0.0005, respectively). Interestingly, WT showed a

shoulder peak at ~681 nm (Fig. 2a), which may reflect partial detachment of light harvesting complex

II (LHCII) from PSII (35). The lower level of normalized PSI fluorescence in c_{6A}-KO in all

photomixotrophic samples did not originate from lower PSI protein levels, as evidenced by PSI

subunit immunoblot analyses (Supplementary Fig. 5).

To investigate if this c_{6A}-dependent light harvesting phenotype was due to alterations in state

transitions in photomixotrophy, for example due to c_{6A}-KO being locked in state 1, we induced either

state 1 (darkness and oxia) or state 2 (darkness and anoxia) in samples grown under standard

conditions (Fig. 2b). Both strains showed, as expected, a lower PSI/PSII fluorescence ratio in state 1

and a higher PSI/PSII fluorescence ratio in state 2, reflecting the increased sizes of the antenna cross

section of PSII and PSI, respectively. However, c_{6A}-KO consistently exhibited a lower PSI/PSII

fluorescence ratio than WT (Fig. 2c). Whereas WT exhibited PSI fluorescence of 1.13 (normalized to

PSII fluorescence) in state 2 conditions, c_{6A}-KO showed a significantly lower value of 1.01 ($P =$

0.046). Similarly, under state 1 inducing conditions, normalized PSI fluorescence in WT was 0.85,

compared to 0.77 in c_{6A}-KO ($P = 0.071$). While the CP47-associated peak at 695 nm (34) showed

different intensity between the two states in WT, in c_{6A}-KO the same peak intensity was recorded in

both states.

Figure 2: c_{6A} affects the light harvesting balance between photosystems I and II in photomixotrophy. a, Low temperature (77 K) chlorophyll fluorescence spectra (excitation: 440 nm) of c_{6A} mutant strains and WT from cells grown under continuous light (30 $\mu mol_{photons}$ m^{-2} s^{-1}) in photoautotrophy -or mixotrophy and cells grown photomixotrophically under DISCO Light for 2 or 4 days. The prominent PSI and PSII fluorescence peaks of note are indicated by black arrows. **b**, Low temperature (77 K) chlorophyll fluorescence spectra of c_{6A}-KO and WT exposed to state 1 or 2 inducing conditions (n = 3 biological replicates ±SE). **c**, Quantification of the PSI to PSII fluorescence ratio at 713 nm from **b**. Statistical testing was carried out with a two-sided t-test per state (p < 0.05: *, p < 0.1: #). **d**, BN-PAGE analysis of c_{6A} mutant strains and WT thylakoid membrane complexes isolated from cells grown in photomixotrophy and continuous light (30 $\mu mol_{photons}$ m^{-2} s^{-1}). The inset shows the area of interest with band III from a separate longer BN-PAGE run with the same samples. The full gels with three biological replicates per strain can be found in Supplementary Figure 6. Thylakoid membrane complexes were assigned to bands according to the literature (36–38). **e,** 2D-PAGE analysis of a BN-PAGE lane from c_{6A}-KO with the boxes indicating assignment of spots to protein complexes based on comparison with calculated R_f values of known thylakoid membrane complex components (Supplementary Fig. 7c).

As the light harvesting balance is affected by photosystem complex abundance and composition, we

next analyzed the thylakoid membrane complexes from cells grown in standard conditions using BN-

PAGE. Gels showed a distinct green band containing a PSI complex without LHCs (band III marked

red, Fig. 2d&e) that was consistently more prominent in c_{6A}-KO compared to WT and c_{6A}-OE. The

other bands showed a similar pattern and thus presumably similar photosystem complex levels across

strains (Fig. 2d). 2D-PAGE analyses and gel simulations to calculate the predicted mobility in the gel

(R_f) revealed similar subunit compositions of the complexes between strains (Fig. 2e &

Supplementary Fig. 7a-b).

Together, these results show that c_{6A} is important for maintaining the light harvesting balance between

the photosystems in photomixotrophic conditions, especially in combination with DISCO light, and

indicate that an altered distribution of LHCs could underly this observation.

<u>Absence of c_{6A} leads to a more reduced PQ pool</u>

Given the tight link between light harvesting balance and the redox state of the PQ pool, we

quantified the redox state of the PQ pool using chlorophyll fluorescence measurements.

Firstly, the chlorophyll fluorescence induction (OJIP) traces of cells grown in standard conditions

revealed a 10% increase in the F_J parameter in c_{6A}-KO compared to WT and c_{6A}-OE ($P = 0.012$ and

0.008, respectively) (Fig. 3a,b). Since F_J reflects the extent of Q_A reduction and is positively

correlated with PQ pool reduction (39), this increase indicates that the absence of c_{6A} led to a more

reduced PQ pool.

To investigate further the extent of c_{6A}-dependent PQ pool overreduction, we exposed cells grown in

standard conditions to DISCO light and recorded chlorophyll fluorescence. The value of the steady-

state fluorescence yield F' under actinic light is influenced by the redox state of the PQ pool, as the

PQ pool is in a redox equilibrium with Q_A in PSII (40). During the high light periods of DISCO Light,

F' was on average 36% higher in c_{6A}-KO than in WT ($P = 0.0006$), indicating a more reduced PQ pool

in the absence of c_{6A} in DISCO Light as well (Fig. 3c). Also during the high light periods of the

DISCO Light regime, 1-qL (a measure of Q_A reduction) was high in WT (0.9) indicating that Q_A and

by implication the PQ pool was highly reduced (41). In c_{6A}-KO 1-qL was significantly higher (0.975)

again compared to WT ($P = 6.8 \times 10^{-5}$) (Fig. 3d), indicating a more reduced PQ pool, consistent with

the F' finding.

These results suggest that the absence of c_{6A} leads to a more reduced PQ pool in *Chlamydomonas*

under photomixotrophic conditions, exacerbated by DISCO Light exposure. Consequently, the more

reduced Q_A pool could lead to increased damage to PSII, potentially contributing to the observed

growth penalty in DISCO Light.

Figure 3: c_{6A} affects the redox state of the PQ pool, the redox kinetics of b hemes in cytochrome b_6f and the quantum yield of PSII. a, Chlorophyll fluorescence induction measurements (OJIP curves) of c_{6A} mutant strains and WT grown under photomixotrophic continuous light (30 $\mu mol_{photons}$ m^{-2} s^{-1}) conditions and normalized to F_0 and F_P (n = 3 biological replicates ±SD). **b**, Comparison of the normalized fluorescence at inflection point F_J (2.4 ms) from **a**. Asterisks indicate a significant difference (p < 0.05) as determined by one-way ANOVA and post-hoc pairwise two-sided t-test (n = 3 biological replicates ±SD). **c**, Raw steady-state fluorescence yield F' in Volts (V) of c_{6A}-KO and WT grown under standard conditions and exposed to a DISCO Light regime (on top of box) in a DUAL-PAM-100 spectrometer (n = 3 biological replicates ±SD). **d**, 1-qL values in the light phases of c_{6A}-KO and WT grown under standard conditions and exposed to the same light protocol as in **c** (n = 3 biological replicates ±SD, asterisk: significant difference (p < 0.001) as determined by two-sided t-test). **e**, Redox changes of b hemes in c_{6A} mutant strains and WT grown under standard conditions and exposed to DCMU (competitive inhibitor of the Q_B site in PSII), methyl viologen (MV, electron acceptor of PSI) or dibromothymoquinone (DBMIB, b_6f Q_P site blocker). Samples were exposed to green light (500 $\mu mol_{photons}$ m^{-2} s^{-1}, green bar) in a JTS-10 spectrometer (n = 3 biological replicates ±SD). **f**, Photosystem II quantum yield, Y(II), of c_{6A}-KO and WT grown under standard conditions and exposed to the same light protocol as in **c** (n = 3 biological replicates ±SD).

Redox kinetics of b hemes are affected and PSII quantum yield lowered in absence of c_{6A}

Given the effects of the absence of c_{6A} on light harvesting and PQ pool redox balance, we also assessed the function of PETC complexes. We first measured the redox kinetics of cytochrome b_6f using a Joliot-type spectrometer (JTS).

In photoautotrophic samples grown under continuous light (30 $\mu mol_{photons}$ m^{-2} s^{-1}), all strains exhibited a b hemes reduction peak within 100 ms of light exposure, but in c_{6A}-KO the peak was only half the intensity of WT (Supplementary Fig. 8). In samples grown in standard photomixotrophic conditions, the b hemes in WT and c_{6A}-OE were reduced within 100 ms of light exposure followed by transient oxidation and re-reduction. In contrast, the b hemes in c_{6A}-KO were *oxidized* within 100 ms of illumination (Fig. 3e) and showed nearly no re-reduction in the following 500 ms. Interestingly, no differences in the cytochrome f redox kinetics were observed in either photoautotrophy or photomixotrophy (Supplementary Fig. 8), indicating no effect of the absence of c_{6A} on the function of the high potential chain.

299 A more reduced PQ pool leads to lower reduction and even apparent oxidation of the b hemes upon

300 light exposure (42–44). The latter is likely due to b hemes being pre-reduced in conditions with a

301 highly reduced PQ pool. Thus, the effect of the absence of c_{6A} on the redox kinetics of b hemes are

302 consistent with the observed change in the redox state of the PQ pool. Artificial modulation of the PQ

303 pool redox state should therefore lead to smaller differences in b hemes redox kinetics between the

304 strains. In the presence of dichlorophenyl-dimethylurea (DCMU), a PSII inhibitor that leads to PQ

305 pool oxidation, samples grown in standard conditions showed that the b hemes in c_{6A}-KO were indeed

306 reduced upon illumination but to a lower magnitude than in WT/ c_{6A}-OE (Fig. 3e). Similarly, when we

307 added methyl viologen (MV), an artificial electron acceptor from PSI that also leads to PQ pool

308 oxidation, the b hemes reduction was partially restored in in c_{6A}-KO (Fig. 3e). Furthermore, when we

309 added dibromothymoquinone (DBMIB), which blocks the Q_P site of cytochrome b_6f and leads to PQ

310 pool reduction, the b hemes only showed oxidation upon illumination without differences between the

311 strains (Fig. 3e).

312 The observed differences in b hemes redox kinetics can therefore be explained by the more reduced

313 PQ pool in c_{6A}-KO. Further, these results indicate that in the absence of c_{6A} the cytochrome b_6f

314 complex including the Q-cycle is in principle functional if the PQ pool is oxidized. Nevertheless,

315 small differences between c_{6A}-KO and WT/ c_{6A}-OE remained when the PQ pool was artificially

316 oxidized, warranting further investigation of the Q-cycle.

318 To investigate the possibility of a disturbed Q-cycle underlying the b hemes phenotype, we measured

319 the proton motive force (pmf) and its components of *Chlamydomonas* cells grown in standard

320 conditions using electrochromic shift (ECS) measurements. No substantial difference between

321 presence and absence of c_{6A} was detected in pmf, proton flux (vH^+) and thylakoid conductivity (gH^+)

322 when samples were exposed to either continuous red light (53 $\mu mol_{photons}$ m^{-2} s^{-1}) or DISCO Light

323 conditions (Supplementary Fig. 9a-f). Further, no differences between the strains were found in the

324 response of the ECS signal to a single turnover flash and the partitioning of pmf into ΔpH and $\Delta \Psi$

325 (Supplementary Fig. 9g&h). Together, this is consistent with c_{6A} not affecting Q-cycle function.

327 We also analyzed the efficiency of photosystems I and II of *Chlamydomonas* cells grown in standard

328 conditions when exposed to continuous or DISCO Light. In continuous light, we did not find any

329 significant differences between WT and c_{6A}-KO in the quantum yield of PSII (Supplementary Fig.

330 10a) and PSI (Supplementary Fig. 10b) over a wide range of intensities. There was, however, elevated

331 photosystem II quantum yield Y(II) for c_{6A}-OE in comparison to c_{6A}-KO at higher light intensities

332 (above 210 $\mu mol_{photons}$ s^{-1} m^{-2}) resulting in an increased relative electron transport rate (rETRII).

333 Conversely, during the high light periods of DISCO Light, Y(II) was less than half in c_{6A}-KO

compared to WT (Fig. 3f). In contrast to PSII, PSI quantum yield Y(I) was not majorly affected by

absence of c_{6A} and DISCO Light and showed a decreasing trend in the light phases from 0.2 to about

0.125 over the course of 40 min in both WT and c_{6A}-KO (Supplementary Fig. 11). For both strains,

DISCO Light led to an increase in acceptor side limitation Y(NA), indicating that in DISCO Light

downstream electron acceptors are not readily available (Supplementary Fig. 11). Consistent with the

more reduced PQ pool in absence of c_{6A}, c_{6A}-KO exhibited a lower donor-side limitation Y(ND) than

WT during DISCO Light treatment (Supplementary Fig. 11).

The DISCO Light condition therefore seems to lead to inhibition of photosynthesis at PSII in c_{6A}-KO

which is consistent with the highly reduced PQ pool in absence of c_{6A}.

<u>c_{6A} partly alleviates the stressful effect of DISCO Light on PSII</u>

To determine if the more reduced PQ pool in the absence of c_{6A} causes stress for *Chlamydomonas* and

could explain the observed growth phenotype, we measured PSII energy partitioning of cells grown in

standard conditions and then exposed to DISCO Light using a DUAL-PAM.

Increased stress levels in the absence of c_{6A} were evident when examining the quantum yield of non-

regulated dissipation of excitation energy, Y(NO), which describes the fraction of energy that is

neither used for photochemistry nor dissipated as heat in a regulated manner via NPQ. At the start of

DISCO Light exposure, over 90% of light energy was dissipated in a non-regulated fashion in c_{6A}-

KO, while WT had values below 80% (Fig. 4a). Over 40 minutes of DISCO Light, Y(NO) levels in

c_{6A}-KO relaxed, nearing WT levels of about 70%. High Y(NO) levels indicate PSII photoinhibition

that is imminent or has already occurred due to over-reduction (45). When probing for PsbA, C-

terminal fragments forming two bands at around 15 kDa were visible (Fig. 4b). An increased ratio of

PsbA (D1) degradation products to full length PsbA (D1) protein, which is indicative of PSII

photoinhibition (46), was observed in c_{6A}-KO on Day 4 of DISCO Light exposure (Fig. 4b &

Supplementary Fig. 12). In c_{6A}-KO, 20% of the observed PsbA (D1) protein signal originated from the

C-terminal fragment bands, while in WT and c_{6A}-OE the C-terminal fragments only contributed 8%

and 7% to the total PsbA (D1) signal, respectively (Supplementary Fig. 12).

The yield of regulated non-photochemical quenching Y(NPQ) was negligible in dark-adapted WT and

c_{6A}-KO samples (Fig. 4b). However, over the first 40 min of DISCO Light exposure, c_{6A}-KO

exhibited approximately twice the Y(NPQ) levels as WT during all phases of the DISCO Light

protocol. This sustained Y(NPQ) response, even in the dark, indicates increased stress levels in the

absence of c_{6A}, leading to an amplified engagement of NPQ mechanisms to allow energy dissipation

in a regulated manner. Interestingly, increased Y(NPQ) in the absence of c_{6A} was also observed under

a range of continuous light intensities, which was partially restored in the c_{6A}-OE strain

(Supplementary Fig. 10a). Elevated NPQ levels in the absence of c_{6A} were further supported by an

approximately 1.5-fold protein level of the NPQ effector LHCSR3 (47) in c_{6A}-KO compared to WT

after two days of growth in DISCO Light (Fig. 4b & Supplementary Fig. 12).

Together, these data indicate that PSII experiences more photoinhibition in c_{6A}-KO than WT and c_{6A}-

OE under DISCO Light consistent with the observed reduced Y(II) in c_{6A}-KO under DISCO Light

(Fig. 3f). Increased photoinhibition is likely contributing to the observed growth penalty (Fig. 1c).

Figure 4: c_{6A} partly alleviates the stressful effect of DISCO Light on *Chlamydomonas*. a, Regulated non-photochemical quenching Y(NPQ), and non-regulated energy dissipation Y(NO) of c_{6A}-KO and WT grown under standard conditions and exposed to a DISCO Light regime in a DUAL-PAM-100 spectrometer (n = 3 biological replicates ±SD). **b**, Immunoblot analyses probing for PsbA in whole cell protein samples from c_{6A} mutant strains and WT grown in DISCO Light for four days (n = 3 biological replicates). **c**, Immunoblot analyses probing for LHCSR3 in whole cell protein samples from c_{6A} mutant strains and WT grown in DISCO Light for two days (n = 3 biological replicates). Relative quantification of the share of signal from C-terminal fragments for PsbA or the signal intensities normalized to total protein for LHCSR3 and can be found in Supplementary Figure 12.

Discussion

Here we present an in-depth characterization of the physiological role of the highly conserved

cytochrome c_{6A} protein in relation to photosynthesis and show that it is important for optimal growth

in fluctuating light due to its effect on the light harvesting and PQ pool redox state balances of the

model alga *Chlamydomonas* (summary see Supplementary Table 1).

Analysis of the redox state of the PQ pool using complementary chlorophyll fluorescence approaches

indicated that the PQ pool was more reduced in c_{6A}-KO, and this was particularly marked in the high

light periods of DISCO Light. JTS measurements of the redox state of the *b* hemes of the cytochrome

b_6f complex in response to light exposure showed disturbance of the redox kinetics in the c_{6A}-KO that

were consistent with a more reduced PQ pool in c_{6A}-KO. Addition of DCMU or MV, which would be

expected to lead to PQ oxidation, or DBMIB, which leads to PQ reduction, revealed effects consistent

with an increased pre-reduction of the b hemes in c_{6A}-KO through a more reduced PQ pool. c_{6A}-KO

exhibited no change in the proton motive force, measured by electrochromic shift, rendering unlikely

a direct effect of c_{6A} on the Q cycle. Under the high light periods of DISCO Light, c_{6A}-KO showed a

dramatic decrease in PSII quantum yield compared to WT, which would also be consistent with a

more reduced PQ pool. The c_{6A}-KO did not exhibit any significant difference in PSI quantum yield

under DISCO light, although it showed a decrease in donor-side limitation of PSI, again consistent

with a more reduced PQ pool. c_{6A}-KO also showed some indication of increased acceptor-side

limitation.

c_{6A}-KO exhibited an increase in the non-regulated dissipation of excitation energy [Y(NO)] and

regulated non-photochemical quenching [Y(NPQ)] with the onset of DISCO Light, as well as a

relative increase in the amount of the NPQ effector LHCSR3 protein and degradation products of

PsbA after two and four days of growth in DISCO Light, respectively. These results indicate an

increased tendency to photoinhibition in c_{6A}-KO, which might be accounted for by the increased PQ

reduction. A general impairment of PSII function or repair in the absence of any other effects from

loss of c_{6A} might be expected to lead to a decreased reduction in PQ. This suggests that the PSII

impairment is a consequence of the increased PQ reduction resulting from loss of c_{6A}, rather than the

primary result of c_{6A} loss.

What might cause the increase in reduction of the PQ pool in the absence of c_{6A}? The 77K chlorophyll

fluorescence data indicated a significantly lower PSI/PSII fluorescence ratio in c_{6A}-KO compared to

WT under photomixotrophic conditions, and this difference was amplified under DISCO Light

conditions. In BN-PAGE, a band corresponding to a LHC-less PSI was more prominent in c_{6A}-KO

than WT and c_{6A}-KO showed a lower PSI/PSII fluorescence value than the wild type under both state

1 and state 2 conditions. These data are consistent with the photosystems being preferentially in the

equivalent of state 1 in c_{6A}-KO compared to WT. However, there was no evidence for the c_{6A}-KO

being permanently 'locked' in either state 1 or state 2.

Overall, it therefore seems likely that c_{6A}-KO is disturbed in the light harvesting balance between PSI

and PSII, in favor of PSII, especially under DISCO Light. This leads to a more reduced PQ pool, and

consequent photoinhibition, with impaired growth of c_{6A}-KO in DISCO conditions. c_{6A} is likely to be

located in the thylakoid lumen and has been shown to interact with the lumen thiol oxidase LTO1

(21), which is closely involved in thiol-based regulation of the STT7 (STN7 in Arabidopsis) state

transition kinase, keeping STT7 oxidized (22). STT7 is active only when oxidized and, in response to

a reduced PQ pool, catalyzes phosphorylation of LHCII and causing LHCII to move from PSII to PSI.

c_{6A} has been proposed to catalyze the formation of disulfide bridges in the lumen (10, 11), transferring

electrons to plastocyanin, with which it has been shown to react (19). Although we cannot exclude a

more complex explanation, with the shift in PSI/PSII balance and change in state of the PQ pool being
indirect consequences of loss of c_{6A}, we propose that c_{6A} contributes to oxidation of STT7. c_{6A} is not
essential for the oxidation as state transitions remain in c_{6A}-KO. However, it may be that c_{6A} assists in
the transfer of electrons from STT7 to LTO1 (with which it interacts (21)) or another acceptor. Thus,
in the absence of c_{6A}, STT7 would be more reduced than in WT, leading to increased light harvesting
at PSII and a more reduced PQ pool compared to WT. This model could be tested in the future by
making substitutions to the conserved cysteine residues in c_{6A}. If correct, this will be a significant
contribution to our understanding of the complex thiol-based regulatory network of the thylakoid
lumen of photosynthetic organisms (48). Furthermore, the demonstration of the importance of c_{6A} for
growth under fluctuating light conditions presents a significant contribution to understand how
organisms cope with rapidly changing light conditions (49, 50). The connection of c_{6A} to state
transitions also make it a significant target for optimizing crop productivity.

Methods

<u>Growth conditions</u>
Chlamydomonas strains were maintained on Tris-Acetate-Phosphate (TAP) (51) medium with
Hutner's trace element solution (52) supplemented with 1.5% agar at 30 $\mu mol_{photons}$ m^{-2} s^{-1} and 25 °C.
For longer-term storage cultures were kept on TAP slants in ambient light at room temperature. Liquid
cultures were grown at 30 $\mu mol_{photons}$ m^{-2} s^{-1}, 25 °C and 120 rpm in AlgaeTron AG130 or AG230
incubators (Photon System Instruments) unless otherwise specified.

For experimental cultures, triplicates of 30 mL medium (Tris-phosphate (TP) medium (51) for
photoautotrophic, and TAP for photomixotrophic conditions) were inoculated with *Chlamydomonas*
strains to OD$_{750nm}$ = 0.1. The cultures were then subjected to the experimental conditions (Table 1).
All experimental cultures were subcultured at least twice in standard conditions before exposure to
experimental conditions.

Table 1: Experimental light conditions in the AlgaeTron AG130 and AG230 growth chambers.

Name	Light regime
Standard growth light	continuous 30 $\mu mol_{photons}$ m^{-2} s^{-1}
High light	continuous 700 $\mu mol_{photons}$ m^{-2} s^{-1}
Darkness	continuous 0 $\mu mol_{photons}$ m^{-2} s^{-1}
2 min on/off	alternating 2 min 700 $\mu mol_{photons}$ m^{-2} s^{-1} & 2 min darkness
12 s on/off	alternating 12 s 700 $\mu mol_{photons}$ m^{-2} s^{-1} & 12 s darkness
DISCO Light	alternating 2 min darkness & 2 min of 12 s 700 $\mu mol_{photons}$ m^{-2} s^{-1}/ 12 s off (Fig. 1d)

<u>Algal strains</u>

The cytochrome c_{6A} knock-out strain "c_{6A}-KO" (CC-1883 background) was generated as previously described (32) using the gRNAs 5'-ATTTACAGTTAGTGCATCCACTTG-3' and 5'-TCCGAGGGTAAGCAAGCACGGCAC-3' to target the 5'UTR and Exon III for Cpf1-mediated double strand breaks, respectively. A repair template with the following sequence was provided at the same time to introduce a multiple stop codon (lowercase letters): 5'-TCATGCTCGGCCAGTTTTCTTTGATTTCAAATCCTCGCTAATTTActaattaactaaCAGCTTCCTT GCTTCATGGACCCAGTAACGGTGCATGGGTACGCG-3'. The knock-out in the strain "c_{6A}-KO" was confirmed through whole locus PCR, showing a truncated locus in the c_{6A}-KO strain compared to WT (Supplementary Fig. 1b). Sanger sequencing validated the deletion of the *CYC4* sequence from 5'UTR to exon III and the introduction of a sequence with multiple stop codons. Whole genome sequencing with high read depth in both strains showed the c_{6A} locus was disrupted in the targeted region in c_{6A}-KO but not in WT (Supplementary Fig. 1c). Transcriptomic analysis showed no c_{6A} gene transcripts in the c_{6A}-KO strain, indicating effective disruption of transcription (Supplementary Fig. 1d).

The c_{6A}-overexpression strain c_{6A}-OE was created by transforming c_{6A}-KO with the constructs pDKM-000 by electroporation as described before (53) with the following specifications. A *Chlamydomonas* culture from standard growth conditions with $OD_{750} \sim 1$ was concentrated 100-fold in TAP with 60 mM sucrose. 250 μL were then incubated with 1 μg of linearized DNA (PmeI) for 10 min on ice in an electroporation cuvette (0.4 cm gap, BioRad) prior to electroporation (BioRad Gene Pulser). Cells were recovered in 10 mL TAP with 60 mM sucrose for 16 h in a shaking incubator (2 μmol photon m^{-2} s^{-1}, 120 rpm, 25°C) prior to plating for selection on TAP-agar plates with Paromomycin (10 μg/mL). Following selection, immunoblot analysis confirmed the expression of c_{6A}-FLAG in three PCR-confirmed transformants, showing a band just under 15 kDa, which corresponds well to the expected size of c_{6A}-FLAG without its predicted signal peptide (13.5 kDa, cut site VLA/AEA, TargetP2.0) (Supplementary Fig. 1e). One colony (number 3 in Supplementary Fig. 1e) was selected as the representative c_{6A}-OE strain. RNAseq analysis showed a ~14-fold increase in c_{6A} transcripts compared to WT, confirming overexpression (Supplementary Fig. 1d).

The construct pDKM-000 was assembled following the molecular cloning syntax with the parts pCM0-016 (PSAD promoter) (53), pDK0-002 (*CYC4* gene with introns, this study), pCM0-094 (C-terminal triple FLAG tag) (53), and pCM0-114 (PSAD terminator) (53). The *CYC4* gene coding for c_{6A} with two intronic BsaI restriction sites altered (GAGACC→GAAACC) was synthesized (Twist Bioscience).

 <u>PCR genotyping and whole genome sequencing</u>

The genotype regarding the *CYC4* allele of the strains used was determined using the Phire Plant Direct PCR kit (Thermo Fisher) and following the manufacturer's instructions. Primers were used that spanned the whole c_{6A} gene binding 245 bp upstream and 75 bp downstream of *CYC4* (fw: 5'-GAGACGCTACGCTGCATGT-3' & rev: 5'-ACTTCCTCGCGCTGCATA-3', annealing: 65 °C, extension: 45 s). For WT, this should lead to an amplicon of 2469 bp while for c_{6A}-KO a 1773 bp amplicon is expected, following the CRISPR induced deletion of 708 bp and addition of a 12 bp multiple stop codon.

For the confirmation of transformation with pDKM-000, colony PCR with the primers spanning the transgene was performed as above (fw: 5'-TCTTTCTCCATGGGCGGGCT-3', rev: 5'-CCCTCGGTCACATGTGCATCC-3', annealing: 69 °C, extension: 70 s).

For whole genome sequencing, genomic DNA from photomixotrophically grown cultures under standard conditions was obtained by phenol:chloroform extraction. The DNA was found to be of sufficient quality via spectrophotometric quantification (NanoDrop) and gel electrophoresis and was subsequently subjected to Illumina sequencing (PE-150 technology, Novogene, Cambridge, UK). After confirming the quality of the raw reads using FastQC (54), BBDuk in Geneious Prime was used to trim adapter sequences, low quality (< 30 quality value) and short reads (<10 bp). Subsequently, reads were aligned to the latest reference genome assembly for *Chlamydomonas* from JGI (CC-4532 v6.1 (55)) with the Geneious Mapper. Coverages of 97.8% (WT) and 97.4% (c_{6A}-KO) were reached with average read depths of 133.6x (WT) and 72.5x (c_{6A}-KO), and average mean MAPQ values of 202.3 (WT) and 200.3 (c_{6A}-KO). Integrative genomics viewer was used for coverage visualization (56).

<u>RNA sequencing</u>

Mid-exponential phase *Chlamydomonas* cultures (5 mL) were harvested by centrifugation at 1000 xg for 5 min. After disposal of the supernatant the cell concentrate was snap-frozen in liquid nitrogen. The frozen pellet was resuspended in 1 mL of TRIzol reagent (Thermo Fisher Scientific) and stored until further use at -80 °C. The TRIzol suspension was mixed with 1 mL of ethanol in preparation for RNA purification using the Direct-zol RNA Miniprep kit (Zymo Research) following the manufacturer's instructions. The optional DNase treatment and two additional wash steps were included: one wash with 0.4 mL of RNA Wash Buffer before the DNase treatment and another wash with 0.4 mL of Direct-zol RNA PreWash Buffer after the first pre-wash. The resulting concentrations and the 260/280, 260/230 ratios were subsequently quantified spectrophotometrically (NanoDrop1000, Thermo Fisher Scientific). Integrity of the extracted RNA was confirmed by gel electrophoresis.

The RNA samples were sent to Novogene for quality control, directional poly-A enriching library

preparation, and Illumina sequencing (PE-150 technology). After confirming the quality of raw reads

using FastQC (54), the latest genome assembly for *Chlamydomonas* from JGI (CC-4532 v6.1 (55))

was used as reference genome onto which raw reads were mapped with hisat2 (57). Alignment was

confirmed (between 82 and 95% aligned reads for all samples) and read assignment statistics were

calculated with picard tools (58). Count data for genetic features was extracted using featureCounts

(59) and the latest gff3 reference file for *Chlamydomonas* from JGI. The featureCounts table was

analyzed in R, where analysis of differential expression was performed with DEseq2 (60).

<u>Chlorophyll quantification</u>

A 200 µL sample from a *Chlamydomonas* culture was mixed with 800 µL acetone, vigorously mixed

for 20 s and centrifuged for 3 min at 13000 xg. The supernatant was isolated and its absorbance at 663

nm and 645 nm was measured against an 80% acetone blank. The concentration of chlorophyll a, b

and total chlorophyll was calculated with the dilution factor D (here D = 5) as Chl a (µg/mL) = $(OD_{663}$

$*12.7 - OD_{645} *2.69) *D$; Chl b (µg/mL) = $(OD_{645} *22.9 - OD_{663} *4.68) *D$; Total chl (µg/mL)= chl a

+ chl b (61).

<u>Protein extraction and Western Blot</u>

A known volume of *Chlamydomonas* cells at a desired growth stage was harvested by centrifugation

at 1000 xg for 5 min. The pellet was subsequently resuspended in RIPA lysis buffer with freshly

added PMSF (1mM final concentration) to reach a calculated OD_{750nm} of 50. The chlorophyll content

was measured as above, but here 10 µL of sample were mixed with 40 µL of acetone and 1 mL of

80% acetone, resulting in a dilution factor D of 105. Protein samples were then adjusted to the same

total chlorophyll concentration with RIPA buffer and frozen at -20 °C for at least 1.5 h or overnight,

followed by thawing on ice to complete cell lysis. Protein concentration was determined using a

Bradford assay (Merck) according to manufacturer's instructions diluting the protein samples 5- to

10-fold.

For SDS-PAGE, all samples were diluted to 1 µg/µL protein with RIPA buffer before being mixed

with 4x Lämmli buffer and incubated for 30 min (50 °C, 600 rpm). After centrifuging the samples

(12000 xg, 1 min) to remove insoluble material, samples were loaded to the desired amount into wells

of polyacrylamide gels (Mini-PROTEAN TGX Stain-Free, BioRad). PageRuler Plus (7 µL, Thermo

Scientific) was used as size marker. Electrophoresis was carried out at 4 °C and 200 V for about 50

min. The gel was imaged in a ChemiDoc MP imaging system (BioRad) using the Stain Free setting

and 5 min activation time. Some gels were stained with Pierce™ Silver Stain for Mass Spectrometry

(Thermo Scientific).

Transfer of protein from polyacrylamide gels onto polyvinylidene fluoride (PVDF) membranes was

carried out using the iBlot dry transfer apparatus, following the manufacturer's instructions and using

program P3 for 5 min. Membranes were then probed for successful transfer with PonceauS (Thermo

Scientific) as per manufacturer's instructions and imaged with the ChemiDoc MP imaging system

(BioRad). Following the PonceauS destaining, membranes were blocked in 5% milk in Tris-buffered

saline with 0.1% Tween 20 (TBS-T) for 1 h at room temperature with gentle shaking. After washing

membranes once with TBS-T to remove remnants of blocking agent, membranes were incubated with

specific primary antibody solutions in 5% milk TBS-T overnight at 4 °C (PsaB (AS10695, Agrisera)

at 1:1000, PsbA D1 (AS05084, Agrisera) at 1:10000, LHCSR3 (AS142766, Agrisera) at 1:2000).

Subsequently, membranes were washed with TBS-T (5x 5 min) and incubated for 1.5 h at room

temperature with goat anti-rabbit horseradish peroxidase-linked secondary antibody in 5% milk TBS-

T (1:10000, 1706515, BioRad). After washing membranes thrice in TBS-T and once in TBS (10 min

each), they were incubated with chemiluminescence substrate (SuperSignal™ West Pico PLUS

Chemiluminescent Substrate, Thermo Scientific) and imaged using a ChemiDoc MP imaging system

(BioRad).

<u>Low temperature chlorophyll (77K) fluorescence</u>

77K fluorescence emission spectra were recorded with a QE Pro spectrophotometer (Ocean Optics,

Dunedin, FL, USA). Cell samples (200 µL) frozen in liquid N_2 were excited with blue light (440 nm).

Raw spectra were normalized to the photosystem II (PSII) fluorescence peak at 685 nm.

Chlamydomonas cells were harvested and total chlorophyll concentration was adjusted to 5 µg/mL

with fresh medium (TP or TAP). Cells were then frozen in liquid N_2 for measurement.

For the state transition experiment, *Chlamydomonas* cells grown in TAP under standard conditions

were adjusted to 5 µg/mL total chlorophyll with fresh medium. Anoxia (state 2 inducing) samples

were adjusted with medium containing glucose (final concentration 20 mM) and Glucose Oxidase

(final concentration 50 U/mL) that was purged with N2 for 3 min. The final samples were again

purged with N_2 before being sealed. Oxia (state 1 inducing) samples were adjusted with unchanged

medium and left unsealed. All samples were then shaken for 30 min in darkness before being frozen

in liquid N_2 for fluorescence measurement.

<u>Blue native PAGE and second dimension SDS-PAGE</u>

Thylakoid membranes from *Chlamydomonas* cultures were isolated over a discontinuous Percoll

gradient as described before (62). BN-PAGE was then carried out largely as previously (63).

Thylakoid samples were diluted with resuspension buffer (25 mM Bis-Tris/HCl (pH 7.0), 20% (w/v)

glycerol, 0.25 mg/ml Pefabloc, and 10 mM sodium fluoride) to a chlorophyll concentration of 0.53

µg/µL and to a total of 6 µg chlorophyll. An equal volume of β -dodecyl maltoside (β -DM, 2% in

resuspension buffer) was added to each sample and the thylakoid membranes were solubilized by

gentle pipetting for 2 min. After centrifugation (4 °C, 18000 xg, 20 min) samples were mixed with a

one-tenth volume of sample loading buffer (100 mM Bis-Tris/HCl (pH 7.0), 0.5 M ACA, 30% (w/v)

sucrose and 50 mg/ml Serva Blue G) and loaded into the wells of a NativePAGE 3-12% Bis-Tris

protein gel (Invitrogen). Electrophoresis was carried out with specific anode buffer (50 mM Bis-

Tris/HCl, pH 7.0) and cathode buffer containing Serva Blue G dye (50 mM Tricine, 15 mM Bis-

Tris/HCl, pH 7.0, and 0.01% Serva Blue G) at 4 °C for 20 min at 75 V, 30 min at 100 V, 30 min at

125 V, 60 min at 150 V, 30 min at 175 V and 60 min at 200 V. After the 30 min at 125 V, the blue

cathode buffer was replaced by the same cathode buffer but without Serva Blue G. The last step was

prolonged to 5 h if additional separation of bands of interest was needed.

Gel electrophoretic separation of complexes in BN-PAGE lanes on a second dimension SDS-PAGE

was carried out as previously described (63, 64), although 12% polyacrylamide gels were used here.

Gels were stained with Pierce™ Silver Stain (Thermo Scientific).

The predicted mobility in the gel (R_f) values of known thylakoid membrane complex components

were calculated using their known molecular weight and a linear regression of the distance of the

protein standard in relation to the running front ($r^2 = 0.98$). The 2D-PAGE spots were compared with

the R_f values and with published BN-PAGE analyses from *Chlamydomonas* (36–38) to assign the

bands and proteins.

<u>Chlorophyll fluorescence</u>

Chlorophyll fluorescence induction (OJIP) of *Chlamydomonas* was measured with a FL3500

fluorometer (Photon System Instruments). Prior to measurement, cells were resuspended in TAP and

total chlorophyll concentration was adjusted to 20 µg/mL. After recovery under standard growth

conditions for at least 30 min, chlorophyll concentration was readjusted to 5 µg/mL. After 15 min

incubation in darkness, fluorescence induction in response to red light (984 $\mu mol_{photons}$ m^{-2} s^{-1}, 45% A-

voltage) was recorded for 120 s. Raw fluorescence data was double normalized to fluorescence at

time point 0 and the maximum fluorescence value to compare the kinetics between samples (41).

Chlorophyll fluorescence of *Chlamydomonas* was measured with a pulse amplitude-modulated

spectrometer (Dual-PAM-100, Walz). Prior to measurements, cells were resuspended in TAP and the

chlorophyll concentration was adjusted to 20 µg/mL. After recovery under standard growth conditions

for at least 30 min, measurements were performed in a quartz cuvette at 25 °C with stirring and

samples were initially incubated in darkness for 5 min before the blue fluorescence measuring light

(Intensity 12, 28 $\mu mol_{photons}$ m^{-2} s^{-1}) was turned on. Saturating pulses (Intensity 5, 5000 $\mu mol_{photons}$ m^{-2}

s^{-1}, 500 ms width) for PSII parameter determination were applied with stirring turned off. DISCO

Light conditions in the DUAL-PAM-100 were achieved by alternating 2 min darkness followed by 2

min consisting of 12 s high light (690 $\mu mol_{photons}$ m^{-2} s^{-1}) and 12 s darkness alternations. Saturating

pulses were applied in the middle of each 12 s phase or three times during the 2 min dark phases. F'

was recorded as the fluorescence yield during illumination with the actinic light treatment before

application of a saturating pulse. Photosynthetic parameters were calculated as described previously

(45).

<u>P700 redox measurements</u>

Simultaneous whole-cell P700 absorbance and chlorophyll fluorescence of *Chlamydomonas*

(Supplementary Figure 10) were measured with a pulse amplitude-modulated spectrometer (Dual-

PAM-100, Walz). Prior to measurements, cells were resuspended in TAP and the total chlorophyll

concentration was adjusted to 20 µg/mL. After recovery under standard growth conditions for at least

30 min, 2 mL of the cell suspension were filtered onto glass fiber filters (WHA1822025, Whatman)

and 100 µL fresh TAP were added to keep the filter moist. Measurements were performed using leaf

clamps holding the filter and samples were initially incubated in darkness for 5 min before

fluorescence (Intensity 12, 28 $\mu mol_{photons}$ m^{-2} s^{-1}) and P700 measuring light (Intensity 15) were turned

on.

Saturating pulses (Intensity 5, 5000 $\mu mol_{photons}$ m^{-2} s^{-1}, 500 ms width) were applied that allowed

calculation of photosynthetic parameters as before (45, 65) with P700 SP-evaluation settings adjusted

to 7 ms delay and 10 ms width.

<u>Cytochrome b$_6$f redox kinetics</u>

Cytochrome *f* and *b* heme redox kinetics were determined in *Chlamydomonas* cells by deconvoluting

absorbance changes at 546, 554, 563, and 573 nm that were measured using a JTS-10 pump probe

spectrophotometer (BioLogic) and appropriate 10 nm FWHM (full width at half maximum)

interference filters (Edmund Optics). BG39 filters (Schott) were used to shield the light detectors from

scattered light. Prior to measurements, cells were resuspended in fresh medium to a total chlorophyll

concentration of 20 µg/mL. After recovery under standard growth conditions for at least 30 min,

Ficoll 400 was added to a final concentration of 10%. For some samples, DCMU, methyl viologen, or

DBMIB were added to a final concentration of 10 µM, 10 mM, or 20 µM respectively, at the same

time as Ficoll. Cells were dark-adapted for 3 min before measurement with each interference filter

and then illuminated with 5 s of green light (500 $\mu mol_{photons}$ m^{-2} s^{-1}). Flashes of white detection light

were administered during 300 µs dark intervals in actinic illumination. Every recording was corrected

for any detection light impact by subtracting a trace with actinic light turned off. Deconvolution was

performed with the JTS-10 software by drawing a baseline between 573 and 546 nm. During analysis,

the signal-to-noise ratio was improved with a Savitzky-Golay filter (66).

<u>Immunofluorescence microscopy</u>

Immunofluorescence staining of *Chlamydomonas* cells was carried out as previously described (67),

however the methanol incubation step was left out to retain chlorophyll in the samples serving as a

thylakoid marker. The antibodies used were primary mouse anti-FLAG (1:2500, B3111, Sigma) and

secondary goat anti-Mouse IgG Alexa Fluor™ 488 (1:200, A11029, Invitrogen).

An inverted fluorescence microscope (Nikon Eclipse Ti) was used to investigate the stained slides.

Brightfield and epifluorescence images were taken with a 20X or 100X oil immersion lens.

AlexaFluor 488 signal was recorded in a YFP channel (500 nm excitation (73% intensity), yellow

emission filter, 500 ms exposure) while another channel was used for chlorophyll autofluorescence

(635 nm excitation (31% intensity), red emission filter, 500 ms exposure). Images were acquired as z-

stacks (acquisition every 0.3 µm) and the fluorescence signals were deconvoluted using the

Richardson-Lucy method and 20 iterations in the NIS-Elements Advanced Research Software

(Nikon).

<u>Co-immunoprecipitation</u>

Chlamydomonas cultures from mid exponential growth phase were harvested by centrifugation (1000

xg, 5 min) and the pellets were resuspended in RIPA lysis buffer with freshly added PMSF (1 mM

final concentration) and DNAse I (final concentration 100 U/mL). Cells were lysed by bead-beating

(~100 µL of 425-600 µm beads) for 5 cycles of 1 min bead-beating and 1 min on ice. The lysate was

pelleted (1 min, 2000 xg) before the supernatant was centrifuged again for 5 min at 15000 xg. The

cleared lysate was then diluted with dilution buffer (10 mM Tris/Cl pH 7.5, 150 mM NaCl, 0.5 mM

EDTA) and protein concentration was determined with a Bradford assay. For co-immunoprecipitation

of FLAG-tagged proteins and putative interaction partners in the lysates, DYKDDDDK Fab-Trap™

Agarose beads (Chromotek) were utilized as per manufacturer's instructions with at least 4 washes

and elution with 2x SDS sample buffer. For the co-immunoprecipitation of samples grown in standard

conditions, a different colony than number 3 but containing the same construct (pDKM-000) was

used.

<u>Electrochromic shift measurements of proton motive force</u>

Changes in the proton motive force and components, vH+ and gH+, were determined with a Dual-

PAM-100 (Walz) by measuring absorbance difference signal at the 550-515 nm, known as

electrochrormic shift (ECS) (68). Dark interval relaxation kinetics (DIRKs) were recorded by

interrupting actinic light exposure for 450 ms with darkness. Prior to measurements, *Chlamydomonas*

cells were resuspended in fresh medium and the total chlorophyll concentration was adjusted to 20

μg/mL. After recovery under standard growth conditions for at least 30 min, Ficoll 400 was added to a

final concentration of 10%. Cells were dark-adapted for 10 min before measurement and

measurements started with a single turnover flash (100 μs) used later for normalization of the ECS

signal. For the standard growth condition induction experiment red actinic light (53 $\mu mol_{photons}$ m^{-2} s^{-1})

was turned on and DIRKs recorded every 5 s for 30 s and every 10 s thereafter. For the DISCO Light

experiment the light regime as in described in the PSII/PSI parameter measurements section was used,

but instead of saturating pulses, DIRKs were recorded. At the end of the protocol, slow relaxation of

the ECS signal in darkness was recorded for 1 min and used later for analysis of partitioning the pmf

into ΔpH and ΔΨ (69). Analysis of kinetic data was carried out using OriginPro (Version 2024,

OriginLab Corporation). The first 20 ms of the DIRK were fitted with a linear regression, the absolute

slope of which corresponds to the proton flux (vH+) (70), while the inverse of the time constant of the

first order exponential decay fit of a DIRK corresponds to the activity of the thylakoid conductivity

(gH+) (71). ECS signals were normalized to the ECS magnitude induced by a single-turnover

saturating flash of 20 μs in dark-adapted cells.

References

1. M. Weigel, *et al.*, Plastocyanin is indispensable for photosynthetic electron flow in Arabidopsis
thaliana. *J. Biol. Chem.* **278**, 31286–31289 (2003).
2. G. A. Peschek, C. Obinger, M. Paumann, The respiratory chain of blue-green algae
(cyanobacteria). *Physiol. Plant.* **120**, 358–369 (2004).
3. R. Gupta, Z. He, S. Luan, Functional relationship of cytochrome c6 and plastocyanin in
Arabidopsis. *Nature* **417**, 567–571 (2002).
4. J. Wastl, D. S. Bendall, C. J. Howe, Higher plants contain a modified cytochrome c6. *Trends*
Plant Sci. **7**, 244–245 (2002).
5. J. Wastl, S. Purton, D. S. Bendall, C. J. Howe, Two forms of cytochrome c6 in a single eukaryote.
Trends Plant Sci. **9**, 474–476 (2004).
6. I. Nimmo, Location and function of cytochrome c6A. *PhD Thesis Univ. Camb.* (2011).
7. C. J. Howe, R. H. Nimmo, A. C. Barbrook, D. S. Bendall, "Cytochrome c6A of Chloroplasts" in
Cytochrome Complexes: Evolution Structures, Energy Transduction and Signaling, Advances in
Photosynthesis and Respiration., W. A. Cramer, T. Kallas, Eds. (Springer Science+Business
Media, 2016).
8. U. Hani, *et al.*, A complex and dynamic redox network regulates oxygen reduction at photosystem
I in Arabidopsis. *Plant Physiol.* (2024). https://doi.org/10.1093/plphys/kiae501.
9. B. A. Slater, The ancestry and function of cytochrome c6A. *PhD Thesis Univ. Camb.* (2020).
10. C. J. Howe, B. G. Schlarb-Ridley, J. Wastl, S. Purton, D. S. Bendall, The novel cytochrome c6 of
chloroplasts: a case of evolutionary bricolage? *J. Exp. Bot.* **57**, 13–22 (2006).
11. B. G. Schlarb-Ridley, R. H. Nimmo, S. Purton, C. J. Howe, D. S. Bendall, Cytochrome c 6A is a
funnel for thiol oxidation in the thylakoid lumen. *FEBS Lett.* **580**, 2166–2169 (2006).
12. H. Chida, *et al.*, Crystal structure of oxidized cytochrome c 6A from Arabidopsis thaliana. *FEBS*
Lett. **580**, 3763–3768 (2006).

13. J. A. R. Worrall, B. F. Luisi, B. G. Schlarb-Ridley, D. S. Bendall, C. J. Howe, Cytochrome c 6A:
discovery, structure and properties responsible for its low haem redox potential. *Biochem. Soc.*
Trans. **36**, 1175–1179 (2008).
14. J. M. Mason, D. S. Bendall, C. J. Howe, J. A. R. Worrall, The role of a disulfide bridge in the
stability and folding kinetics of Arabidopsis thaliana cytochrome c6A. *Biochim. Biophys. Acta*
BBA - Proteins Proteomics **1824**, 311–318 (2012).
15. J. A. R. Worrall, *et al.*, Modulation of Heme Redox Potential in the Cytochrome c 6 Family. *J.*
Am. Chem. Soc. **129**, 9468–9475 (2007).
16. B. Slater, D. Kosmützky, R. E. R. Nisbet, C. J. Howe, The Evolution of the Cytochrome c6
Family of Photosynthetic Electron Transfer Proteins. *Genome Biol. Evol.* **13**, 1–12 (2021).
17. F. P. Molina-Heredia, *et al.*, A new function for an old cytochrome? *Nature* **424**, 33–34 (2003).
18. J. Wastl, *et al.*, Redox properties of Arabidopsis cytochrome c6 are independent of the loop
extension specific to higher plants. *Biochim. Biophys. Acta BBA - Bioenerg.* **1657**, 115–120
(2004).
19. M. J. Marcaida, *et al.*, Structure of Cytochrome c6A, a Novel Dithio-cytochrome of Arabidopsis
thaliana, and its Reactivity with Plastocyanin: Implications for Function. *J. Mol. Biol.* **360**, 968–
977 (2006).
20. P. Pesaresi, *et al.*, Mutants, overexpressors, and interactors of arabidopsis plastocyanin isoforms:
Revised roles of plastocyanin in photosynthetic electron flow and thylakoid redox state. *Mol.*
Plant **2**, 236–248 (2009).
21. Y. Lu, *et al.*, Identification of Potential Targets for Thylakoid Oxidoreductase AtVKOR/LTO1 in
Chloroplasts. *Protein Pept. Lett.* **22**, 219–225 (2015).
22. J. Wu, *et al.*, Functional redox links between lumen thiol oxidoreductase1 and serine/threonine-
protein kinase STN7. *Plant Physiol.* **186**, 964–976 (2021).
23. P. J. Gollan, A. Trotta, A. A. Bajwa, I. Mancini, E.-M. Aro, Characterization of the Free and
Membrane-Associated Fractions of the Thylakoid Lumen Proteome in Arabidopsis thaliana. *Int.*
J. Mol. Sci. **22**, 8126 (2021).
24. J. Beltrán, *et al.*, Specialized Plastids Trigger Tissue-Specific Signaling for Systemic Stress
Response in Plants. *Plant Physiol.* **178**, 672–683 (2018).
25. M. Janssen, *et al.*, Efficiency of light utilization of Chlamydomonas reinhardtii under medium-
duration light/dark cycles. *J. Biotechnol.* **78**, 123–137 (2000).
26. P. J. Graham, B. Nguyen, T. Burdyny, D. Sinton, A penalty on photosynthetic growth in
fluctuating light. *Sci. Rep.* **7**, 1–11 (2017).
27. L. Nikkanen, *et al.*, PGR5 is needed for redox-dependent regulation of ATP synthase both in
chloroplasts and in cyanobacteria. [Preprint] (2024). Available at:
http://biorxiv.org/lookup/doi/10.1101/2024.11.03.621747 [Accessed 27 August 2025].
28. M. Jokel, X. Johnson, G. Peltier, E. Aro, Y. Allahverdiyeva, Hunting the main player enabling
Chlamydomonas reinhardtii growth under fluctuating light. *Plant J.* **94**, 822–835 (2018).
29. C. J. Steen, A. Burlacot, A. H. Short, K. K. Niyogi, G. R. Fleming, Interplay between LHCSR
proteins and state transitions governs the NPQ response in Chlamydomonas during light
fluctuations. *Plant Cell Environ.* **45**, 2428–2445 (2022).
30. T. Roach, LHCSR3-Type NPQ Prevents Photoinhibition and Slowed Growth under Fluctuating
Light in Chlamydomonas reinhardtii. *Plants* **9**, 1604 (2020).
31. W. J. Nawrocki, *et al.*, Chlororespiration Controls Growth Under Intermittent Light. *Plant*
Physiol. **179**, 630–639 (2019).
32. A. Ferenczi, D. E. Pyott, A. Xipnitou, A. Molnar, Efficient targeted DNA editing and replacement
in Chlamydomonas reinhardtii using Cpf1 ribonucleoproteins and single-stranded DNA. *Proc.*
Natl. Acad. Sci. **114**, 13567–13572 (2017).
33. C. A. Kerfeld, H. P. Anwar, R. Interrante, S. Merchant, T. O. Yeates, The Structure of Chloroplast
Cytochromec6at 1.9 Å Resolution: Evidence for Functional Oligomerization. *J. Mol. Biol.* **250**,
627–647 (1995).
34. E. G. Andrizhiyevskaya, *et al.*, Origin of the F685 and F695 fluorescence in Photosystem II.
Photosynth. Res. **84**, 173–180 (2005).

35. J. Kargul, J. Nield, J. Barber, Three-dimensional reconstruction of a light-harvesting complex I-photosystem I (LHCI-PSI) supercomplex from the green alga Chlamydomonas reinhardtii: Insights into light harvesting for PSI. *J. Biol. Chem.* **278**, 16135–16141 (2003).

36. E. Devadasu, *et al.*, Long- and short-term acclimation of the photosynthetic apparatus to salinity in Chlamydomonas reinhardtii. The role of Stt7 protein kinase. *Front. Plant Sci.* **14**, 1–17 (2023).

37. J. G. García-Cerdán, *et al.*, A thylakoid membrane-bound and redox-active rubredoxin (RBD1) functions in de novo assembly and repair of photosystem II. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 16631–16640 (2019).

38. B. Spaniol, *et al.*, Complexome profiling on the Chlamydomonas lpa2 mutant reveals insights into PSII biogenesis and new PSII associated proteins. *J. Exp. Bot.* **73**, 245–262 (2022).

39. S. Z. Tóth, G. Schansker, R. J. Strasser, A non-invasive assay of the plastoquinone pool redox state based on the OJIP-transient. *Photosynth. Res.* **93**, 193 (2007).

40. B. A. Diner, Dependence of the deactivation reactions of Photosystem II on the redox state of plastoquinone pool a varied under anaerobic conditions. Equilibria on the acceptor side of Photosystem II. *Biochim. Biophys. Acta BBA - Bioenerg.* **460**, 247–258 (1977).

41. H. M. Kalaji, *et al.*, Frequently asked questions about chlorophyll fluorescence, the sequel. *Photosynth. Res.* **132**, 13–66 (2017).

42. P. Joliot, A. Joliot, Proton pumping and electron transfer in the cytochrome bf complex of algae. *Biochim. Biophys. Acta BBA - Bioenerg.* **849**, 211–222 (1986).

43. F. Buchert, M. Scholz, M. Hippler, Electron transfer via cytochrome b 6 f complex displays sensitivity to antimycin A upon STT7 kinase activation. *Biochem. J.* **479**, 111–127 (2022).

44. A. Malnoë, F.-A. Wollman, C. de Vitry, F. Rappaport, Photosynthetic growth despite a broken Q-cycle. *Nat. Commun.* **2**, 301 (2011).

45. C. Klughammer, U. Schreiber, Complementary PS II quantum yields calculated from simple fluorescence parameters measured by PAM fluorometry and the Saturation Pulse method. *PAM Appl. Notes* **1**, 27–35 (2008).

46. W. J. Nawrocki, *et al.*, Molecular origins of induction and loss of photoinhibition-related energy dissipation q I. *Sci. Adv.* **7**, 1–13 (2021).

47. M. Ballottari, *et al.*, Identification of pH-sensing Sites in the Light Harvesting Complex Stress-related 3 Protein Essential for Triggering Non-photochemical Quenching in Chlamydomonas reinhardtii. *J. Biol. Chem.* **291**, 7334–7346 (2016).

48. D. Hoh, J. E. Froehlich, D. M. Kramer, Redox regulation in chloroplast thylakoid lumen: The pmf changes everything, again. *Plant Cell Environ.* **47**, 2749–2765 (2024).

49. N. A. Eckardt, *et al.*, Lighting the way: Compelling open questions in photosynthesis research. *Plant Cell* **14**, 1–16 (2024).

50. S. P. Long, *et al.*, Into the Shadows and Back into Sunlight: Photosynthesis in Fluctuating Light. *Annu. Rev. Plant Biol.* **73**, 617–648 (2022).

51. D. S. Gorman, R. P. Levine, Cytochrome f and plastocyanin: their sequence in the photosynthetic electron transport chain of Chlamydomonas reinhardi. *Proc. Natl. Acad. Sci.* **54**, 1665–1669 (1965).

52. S. H. Hutner, L. Provasoli, A. Schnatz, C. P. Haskins, Some Approaches to the Study of the Role of Metals in the Metabolism of Microorganisms. *Proc. Am. Philos. Soc.* **94**, 152–170 (1950).

53. P. Crozet, *et al.*, Birth of a Photosynthetic Chassis: A MoClo Toolkit Enabling Synthetic Biology in the Microalga *Chlamydomonas reinhardtii. ACS Synth. Biol.* **7**, 2074–2086 (2018).

54. S. Andrews, FastQC: A Quality Control Tool for High Throughput Sequence Data. (2010). Available at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

55. R. J. Craig, *et al.*, The Chlamydomonas Genome Project, version 6: Reference assemblies for mating-type plus and minus strains reveal extensive structural mutation in the laboratory. *Plant Cell* **35**, 644–672 (2023).

56. J. T. Robinson, *et al.*, Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

57. D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915 (2019). Available at: http://dx.doi.org/10.1038/s41587-019-0201-4.

58. Broad Institute, Picard Tools. Available at: http://broadinstitute.github.io/picard.

59. Y. Liao, G. K. Smyth, W. Shi, FeatureCounts: An efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

60. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

61. D. I. Arnon, COPPER ENZYMES IN ISOLATED CHLOROPLASTS. POLYPHENOLOXIDASE IN *BETA VULGARIS*. *Plant Physiol.* **24**, 1–15 (1949).

62. C. B. Mason, T. M. Bricker, J. V. Moroney, A rapid method for chloroplast isolation from the green alga Chlamydomonas reinhardtii. *Nat. Protoc.* **1**, 2227–2230 (2006).

63. S. Järvi, M. Suorsa, V. Paakkarinen, E. M. Aro, Optimized native gel systems for separation of thylakoid protein complexes: Novel super- and mega-complexes. *Biochem. J.* **439**, 207–214 (2011).

64. M. Rantala, V. Paakkarinen, E. M. Aro, Analysis of thylakoid membrane protein complexes by blue native gel electrophoresis. *J. Vis. Exp.* **2018**, 1–6 (2018).

65. C. Klughammer, U. Schreiber, Saturation Pulse method for assessment of energy conversion in PS I. *PAM Appl. Notes* 11–14 (2008).

66. M. J. E. Savitzky, A.; Golay, Smoothing and Differentiation of Data. *Anal Chem* **36**, 1627–1639 (1964).

67. T. Yamano, H. Fukuzawa, Indirect Immunofluorescence Assay in Chlamydomonas reinhardtii. *BIO-Protoc.* **6**, 1–6 (2016).

68. B. Bailleul, P. Cardol, C. Breyton, G. Finazzi, Electrochromism: A useful probe to study algal photosynthesis. *Photosynth. Res.* **106**, 179–189 (2010).

69. J. A. Cruz, C. A. Sacksteder, A. Kanazawa, D. M. Kramer, Contribution of electric field ($\Delta\psi$) to steady-state transthylakoid proton motive force (pmf) in vitro and in vivo. Control of pmf parsing into $\Delta\psi$ and ΔpH by ionic strength. *Biochemistry* **40**, 1226–1237 (2001).

70. P. Joliot, A. Joliot, Cyclic electron transfer in plant leaf. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 10209–10214 (2002).

71. A. Kanazawa, D. M. Kramer, In vivo modulation of nonphotochemical exciton quenching (NPQ) by regulation of the chloroplast ATP synthase. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 12789–12794 (2002).

72. S. Rexroth, J. M. W. Meyer zu Tittingdorf, F. Krause, N. A. Dencher, H. Seelert, Thylakoid membrane at altered metabolic state: Challenging the forgotten realms of the proteome. *Electrophoresis* **24**, 2814–2823 (2003).

Acknowledgements

We thank Dr. K. Geissler (University of Cambridge) for the provision of the pCM0 MoClo parts and Dr. J. Walter (University of Cambridge) for assistance with blue native PAGE. We are also grateful to Dr. M. Jokel (University of Turku) for discussions on cell culture. All photosynthesis experiments were performed within the PHOTOSYN infrastructure at the University of Turku. We gratefully acknowledge the support of this work by: the Gates Cambridge Trust with support from the Benn W Levy Trust, and the Professor John Glover Memorial Fund (to DK); the Novo Nordisk Foundation (PhotoCat, project no. NNF20OC0064371, to YA; and Photo-e-microbes project no. NNF22OCOO79717 to LTW); the Research Council of Finland no. 355335 to YA, (no. 354876) to LN; the Biotechnology and Biological Sciences Research Council (BB/M011194/1 to AS, BB/M011194/1 to JML, BB/R506163/1 to AF, and BB/Y001370/1 to CJH), and Trinity Hall, Cambridge (to JML).

Author contributions

DK conceived the project, designed and performed experiments under the supervision of CJH and YA, and analyzed data. DK, LTW, LN, and YA developed the biophysical part of the project. LTW and LN supervised the biophysical measurements and assisted with data analysis and interpretation. AF created the knock-out mutant line under the supervision of AM. JML and AS assisted with data

899 collection, analysis and interpretation. AM, YA, and CJH secured funding, assisted with data
900 interpretation, and provided resources. DK wrote the manuscript with comments and input from all
901 authors.

902 **Competing interests**
903 The authors declare no competing interests.