Temporally resolved glutamate and GABA responses measured in human medial frontal cortex during working memory encoding and recall

Daniel Cocking[1†], Ryan Elson[2†], Katherine Dyke[2], Mohammad Zia Ul Haq Katshu[3,4], Adam Berrington[1⊗], Claudia Danielmeier[2⊗]

1. Sir Peter Mansfield Imaging Centre, School of Physics and Astronomy, University of Nottingham, Nottingham, United Kingdom
2. School of Psychology, University of Nottingham, Nottingham, United Kingdom
3. Institute of Mental Health, School of Medicine, University of Nottingham, Nottingham, United Kingdom
4. Nottinghamshire Healthcare NHS Foundation Trust, United Kingdom

[†] Co-first authors
[⊗] Co-principal investigators

*Corresponding Author: Daniel Cocking
Address: Sir Peter Mansfield Imaging Centre, University of Nottingham, University Park, Nottingham, UK, NG7 2RD
E-mail: daniel.cocking@nottingham.ac.uk
Phone Number: +44 115 9514737

*Corresponding Author: Ryan Elson
Address: School of Psychology, University of Nottingham, University Park, Nottingham, UK, NG7 2RD
E-mail: ryan.elson@nottingham.ac.uk
Phone Number: +44 115 7486970

Keywords: functional magnetic resonance spectroscopy (fMRS), glutamate dynamics, GABA, working memory, medial frontal cortex, event-related design, 7T, cognitive neurochemistry

Abstract

Resolving rapid neurochemical changes during cognition is key to linking neurophysiology with behaviour. We developed a mixed block/event-related design of a working memory (WM) task for functional magnetic resonance spectroscopy (fMRS) at 7T to resolve glutamate and GABA fluctuations during encoding and recall phases of WM. Participants performed the WM task interleaved with rest and control conditions while semi-LASER spectra were acquired from the medial frontal cortex. Group-level phase-specific analyses revealed significant glutamate, but not GABA, increases during encoding and recall relative to control. However, larger individual increases in GABA during encoding predicted higher WM accuracy. Glutamate exhibited distinct, temporally resolved peaks for encoding (~1.1s) and recall (~1.4s). These time courses extend beyond previously reported post-stimulus windows and align closely with behavioural response latencies. Our findings provide temporally high-resolved characterisations of glutamate responses during distinct phases of WM and highlight differential roles of glutamate and GABA in WM processes.

1.Introduction

The cognitive processes underlying working memory (WM) emerge from rapid fluctuations in excitatory and inhibitory neural activity, yet these dynamics are challenging to quantify *in vivo*. Probing these dynamic processes in humans offers a critical link between neural activity and behaviour. Using proton (^{1}H) magnetic resonance spectroscopy (MRS), resting concentrations of the major excitatory and inhibitory neurotransmitters, glutamate (Glu) and gamma-aminobutyric acid (GABA), have been associated with varied measures of cognitive function[1–5]. Time-resolved measurements during functional tasks (fMRS) have revealed transient changes in Glu and GABA concentrations to cognitive demands [6–8], allowing more direct associations with cognition. Such studies typically rely on blocked-designs, averaging measurements over long durations (>15 s) of a task paradigm [9–17]. This approach limits the ability to assign neurochemical changes to specific cognitive processes within trials, such as the different phases within working memory (WM) paradigms. Although tracking changes to individual events on a sub-second scale with time-locking to stimulus and event-related designs is feasible [6,18,19], short timescale temporal dynamics of Glu and GABA, which may reflect synaptic activity [20], remain elusive.

Canonical 'time courses' of Glu and GABA concentration are currently lacking. Glu has been proposed to peak in concentration around ~0.5 s after stimulus onset [17] and return to baseline after three to four seconds, [17,19]. The lack of an established response complicates study design in determining the optimal time to acquire spectroscopy data post-stimulus. Compared to the postulated peak at 0.5 s [17], some prior work may have sampled too early, [6,21–23] or too late [24]. Small concentration changes make measuring time courses of neurochemicals difficult, thus fMRS studies often measure changes over long durations. Cognitively demanding tasks such as WM paradigms, however, offer a possibility to elicit strong responses, and hence detectable signal changes, over timescales on the order of single events.

Working memory is a crucial cognitive ability allowing us to attend to and manipulate information held in short-term memory [25]. It comprises of three core phases: encoding, maintenance and retrieval. A commonly used WM paradigm is the n-back task [7,8] which conflates encoding and recall, obscuring which neurochemical shifts occur during different phases of WM. Therefore, to characterise the temporal

dynamics of WM-related neurochemical responses with fMRS in different phases, a
task paradigm separating encoding, maintenance, and retrieval is required.

In the current work, we developed a unique event-related fMRS acquisition at
7T, sensitive to the temporal dynamics of Glu by sampling at 16 time points over a
2.5 s window post-stimulus. Our experimental design allowed separate estimation of
Glu concentration time courses associated with encoding and recall phases of an
associative WM task. Participants encoded the colour, shape and location of four
abstract shapes, then recalled the colour from a shape or location cue. Normalising
to control trials allowed us to isolate WM-specific processes over and above motor
and visual responses within an fMRS voxel situated in the pre-SMA region of medial
frontal cortex (MFC), guided by a separate fMRI localiser study (see Supplementary
Materials). The MFC is involved in many of the cognitive functions required for
associative WM, such as executive function and attention [26–30]. Different
subpopulations of neurons in the pre-SMA region have also been associated with
different stages of WM [31].

We found evidence for a transient Glu response in the MFC of healthy human
participants at 7T, displaying peak increases in concentration around 1.1s and 1.4s,
following encoding and recall, respectively. At an individual level, Glu change was
not related to task performance, whereas changes in GABA concentrations
compared to control trials were positively correlated with WM performance,
particularly during encoding, suggesting that inhibitory processes, potentially to
inhibit task-irrelevant information, are important during demanding WM tasks.

103 2. Results

104 2.1. Time courses of Glu reveal distinct peaks during
105   encoding and recall

We estimated Glu time courses from our MRS voxel during WM encoding and
recall and in control trials. Across the trials, data were acquired at 16 sampling points
between stimulus onset (0 s) and stimulus offset (2.5 s) in 0.17 s intervals. To
improve neurochemical estimation for Glu, spectra were binned into eight timepoints
prior to spectral fitting (see Fig. 1). Within each run, concentrations for each time bin
were converted to a percentage change from the run's initial rest period (the

"baseline", comprising eight spectra with Cramer-Rao Lower Bounds [CRLBS] =

9.1% [SE = 0.8%]), and then averaged across runs.

During encoding, Glu was significantly increased compared to baseline, prior

to correction, at 1.08 s post stimulus onset (4th time bin [1.00-1.17 s]; mean [SE] =

3.87% [1.45%], t(18) = 2.67, p = .016, 95% CI = [0.82, 6.93], d = 0.61, see Fig.1a).

During recall, Glu levels were significantly increased, prior to correction, at 1.42 s

post probe onset (5th time bin [1.33-1.50 s]; mean [SE] = 3.61% [1.68%], t(18) =

2.15, p = .045, 95% CI = [0.09, 7.13], d = 0.49). No other timepoint was significant for

either WM phase or the control task. Bonferroni-corrected α = .002.

During encoding, Glu was significantly increased compared to control at 1.08

s (4th time bin; mean [SE] = 5.41% [1.57%], t(18) = 3.44, p = .003, 95% CI = [2.11,

8.71], d = 0.79), measured by converting raw concentrations during encoding (and

recall) to percent changes relative to control, demonstrating a Glu increase specific

to WM-related processes, independent from visual or motor demands in control trials

(Fig. 1b). This normalisation was performed separately for each time point, with

changes averaged over runs.

During recall, Glu was significantly increased compared to control at 1.42 s

(5th time bin; mean [SE] = 5.30% [1.42%], t(18) = 3.74, p = .002, 95% CI = [2.32,

8.28], d = 0.86) and, prior to correction, was also increased at 1.08 s (4th time bin;

mean [SE] = 4.72% [1.87%], t(18) = 2.52, p = .021, 95% CI = [0.79, 8.65], d = 0.58).

Bonferroni-corrected α = .003.

Figure 1

Time courses of Glu concentration changes for Control, Encoding and Recall phases.

Note: (**a**) Glu time course measured during control trials (blue, top), encoding (orange, middle) and recall (red, bottom) phase as percentage change from the baseline. Vertical dotted lines represent the average reaction times for control (blue, 0.91 s) and working memory trials (red, 1.38 s). Linked dots at the bottom of each graph represent the pair of sampling points in each time bin. (**b**) Glu response for encoding (orange) and recall (red) normalised to control Glu (blue) for each time bin separately (% change). (**c**) average % increases in Glu compared to control trials during encoding (orange) and recall (red) across all 16 sampling points (left), the 'peak' window including the middle eight sampling points (middle) and the outside window including the first four and last four sampling points (right). All error bars and

shaded regions show the standard error across participants. * Significant uncorrected, ** significant Bonferroni-corrected.

134   The mean (SE) Glu Cramér-Rao Lower Bound (CRLB) values across each
135 time point for control, encoding and recall WM phases were 12.25% (1.24%),
136 12.05% (1.16%) and 12.01 (1.11%) respectively. CRLBs did not significantly differ by
137 time point or phase indicating no difference in data quality.

2.2. Glu, but not GABA, increases during encoding and recall

139
140

141
The novel time course reconstructions allowed us to interrogate the Glu
response over time. We employed subsequent analyses to compare the overall Glu
and GABA concentration changes averaged over longer post-stimulus time windows
(two windows ~1.25 s, one 2.5 s). To isolate WM-specific changes in neurochemical
concentrations, results below are reported as the percentage change from control
trials.
We aimed to test three questions. 1) Did the concentration changes during
encoding and recall differ from control trials? 2) Did the concentration changes differ
between encoding and recall? 3) Were the concentration changes significantly
correlated across encoding and recall?
On average, Glu increased by 1.40% (SE: 0.65%) during encoding and 1.82%
(SE: 0.84%) during recall relative to control (Fig. 1c). This was shown by initially
averaging spectra for all 16 sampling points prior to fitting (32 spectra per fit per run)
These increases were significant prior to Bonferroni correction (α = .025, encoding:
t(18) = 2.17, p = .044, 95%CI = [0.05, 2.76], d = 0.50, recall: t(18) = 2.18, p = .043,
95%CI = [0.06, 3.58], d = 0.50). Concentration changes during encoding and recall
did not significantly differ (t(18) = 0.55, p = .591, 95%CI = [-1.18, 2.02], d = 0.13) and
were significantly correlated (r = 0.50, p = .030). The mean (SE) Glu CRLB values
did not significantly differ between control, encoding and recall WM phases: 5.72%
(0.44%), 5.66% (0.41%) and 5.65 (0.42%)

2.2.1. Windowing (0.67-1.83 s) increases the Glu response

Given the inverted U-shaped rise and fall in the time course of Glu, we employed a windowed averaging approach centred around the peaks (~1.08 and ~1.42 s) of the Glu response, assessing Glu in a critical window without the influence of earlier and later time points where the concentration appears to be at baseline. We separated the 'peak' (middle) eight sampling points (0.67-1.83 s) focusing on the peak of Glu responses and the 'outside' eight sampling points (the first four and last four sampling points). For each analysis, spectra were averaged across the eight sampling points prior to fitting (16 spectra per fit per run; Fig. 1c). 'Peak' and 'outside' Glu concentrations during encoding and recall were separately normalised to the corresponding windows of the control task (Fig. 1c).

A 2x2 within-subjects ANOVA showed a main effect of time window (peak/outside: $F(1, 18) = 6.41$, $p = .021$, $\eta_p^2 = 0.26$), with significantly higher Glu changes in the peak window (mean [SE] = 2.83% [0.77%]) than in the outside time windows (mean [SE] = 0.22% [0.85%]). The effect of WM phase (encoding/recall) was not significant ($F(1, 18) = 0.92$, $p = .351$, $\eta_p^2 = 0.05$) nor was the interaction ($F(1, 18) = 0.01$, $p = .922$, $\eta_p^2 = 0.00$). This suggests WM-related Glu changes are restricted to approximately between 0.67-1.83 s post stimulus onset.

In this 'peak' time window (0.67-1.83 s), we observed significant Glu increases during encoding (mean [SE] = 2.49% [0.97%], $t(18) = 2.58$, $p = .019$, 95%CI = [0.46, 4.52], $d = 0.59$) and recall (mean [SE] = 3.17% [1.03%], $t(18) = 3.06$, $p = .007$, 95%CI = [1.00, 5.34], $d = 0.70$) compared to the control condition. Bonferroni-corrected $\alpha = .025$. Interestingly, unlike the full window analysis, Glu concentration changes were not significantly correlated across encoding and recall ($r = 0.17$, $p = .483$), suggesting more independent Glu responses related to the different WM phases when the influence of baseline Glu levels is reduced. CLRBs did not significantly differ between control, encoding and recall WM phases using the 'peak' sampling points: 6.97% (0.54%), 6.98% (0.57%) and 6.97% (0.51%) respectively.

As the results highlighted the benefits of cropping to the peak time window for Glu, all subsequent analyses, including those involving GABA are also restricted to this middle window. Results incorporating all 16 sampling points can be seen in the Supplementary Materials (e.g., see Supplementary Fig.1).

2.2.2. Group-level GABA concentrations are not modulated during WM phases

GABA concentrations did not significantly differ from control trials during encoding (mean [SE] = -1.11% [3.33%], $t(17)$ = -0.33, p = .743, 95%CI = [-8.15, 5.92], d = 0.08) or recall (mean [SE] = 1.04% [2.41%], $t(17)$ = 0.43, p = .672, 95%CI = [-4.02, 6.12], d = 0.10). Bonferroni-corrected α = .025. The concentration changes did not significantly differ between encoding and recall ($t(17)$ = 0.66, p = .517, 95%CI = [-4.70, 9.00], d = 0.16) and were highly correlated (r = 0.63, p = .004). GABA CLRBs did not significantly differ between control, encoding and recall WM phases: 21.48% (1.40%), 23.01% (1.60%) and 22.52% (1.29%), respectively. The GABA CRLB values are larger than that of Glu analysis, and therefore, to maintain reliability of measurements, spectra were not split into smaller bin sizes.

2.3. Changes in GABA, but not Glu, predict WM performance

We investigated the relationship between task performance (percent accuracy and RT) and the change in Glu and GABA levels during encoding and recall (averaged around the Glu peak as described above) relative to control trials. Correlations can be seen in Fig. 2 (for completeness, correlations using all sampling points can be seen in the Supplementary Materials and Supplementary Fig. 2).

Interestingly, increases in GABA during WM encoding predicted better task performance (Fig. 2b), with participants with more GABA increases showing higher accuracy (r = 0.60, p = .008; 1 outlier removed with Cook's distance >1 in all correlations) and faster reaction times, significant prior to correction (r = -0.47, p = .049). Trends in the same direction were present for GABA during recall (accuracy: r = 0.42, p = .083, RT: r = -0.43, p = .074). Bonferroni-corrected α = .025. Thus, while group-level GABA concentrations were not significantly increased during WM phases compared to control, individual-level GABA dynamics predicted task performance. In contrast, changes in Glu during encoding or recall were not significantly correlated with accuracy or RT on the WM trials (*all p* >= .455, see Fig. 2a).

Figure 2

Correlations between the Glu (**a**) and GABA (**b**) changes from control with working memory performance.

Note: Percent signal change from control trials during encoding (orange) and recall (red), showing changes in Glu (**a**) and GABA (**b**), showing the group-level changes (left) and the correlations with working memory accuracy (middle) and working memory reaction time (right). Plots show data using the middle eight sampling points. * Significant prior to correction, ** significant after correction.

228   ## 2.4.   Trend towards diminishing Glu increase across

229   ## runs

Figure 3

Behavioural data, Glu and GABA across runs.

Note: Data across runs. (**a**) Mean accuracy and (**b**) reaction time during the working memory trials. Mean percent Glu (**c**) and GABA (**d**) change within memory trials (averaged across encoding and recall) from control trials using the middle eight sampling points. (**e**) Raw Glu concentration at baseline. Error bars show +/- 1 SE across participants. * Significant prior to correction, ** significant after correction.

To assess the effect of run on neurochemical responses (see Fig. 3), we

averaged the percent changes in glutamate and GABA across encoding and recall.

See supplementary materials (Supplementary Fig. 3) for results using all 16

sampling points.

For Glu, mean percent signal changes (SE) per run were: Run 1: 5.73%

(1.78%), Run 2: 2.59% (1.36%), Run 3: -0.27% (1.21%), Run 4: 3.26% (1.70%). The

within-subjects ANOVA showed a trending effect of run ($F(3, 56) = 2.60$, $p = .062$, η_p^2

= 0.13, see Fig. 3c). The decrease in Glu change from Run 1 to Run 3 was

significant prior to correction ($\alpha = .008$, $t(18) = -2.57$, $p = .019$, 95% CI = [-10.88, -

1.10], $d = 0.59$). No other differences between runs were significant (all $p >= .102$).

The assumption of sphericity was met.

Compared to control trials, Glu showed a significant increase for Run 1 only

($t(18) = 3.23$, $p = .005$, 95% CI = [2.00, 9.46], $d = 0.74$). The increases in Run 2

($t(18) = 1.90$, $p = .073$, 95% CI = [-0.27, 5.45], $d = 0.44$) and Run 4 ($t(18) = 1.92$, $p =$

.071, 95% CI = [-0.31, 6.83], d = 0.44) only trended towards significance prior to
correction (α = .013).

For additional context, there was no significant effect of run for the raw Glu
concentrations obtained during the initial rest spectra at the start of each run (eight
spectra per fit, F(3, 54) = 0.07, p = .975, η_p^2 = 0.00, see Fig. 3e) showing that Glu
decreases over time were related to participants performing the task but returned to
similar concentrations in between runs.

For GABA (Fig. 3d), the effect of run was not significant (F(2.01, 36.26) =
1.58, p = .220, η_p^2 = 0.08, Greenhouse-Geisser correction applied) and
concentrations during WM blocks did not differ from control (all p >= .175). The mean
percent changes were: Run 1: 9.58% (7.96%), Run 2: -6.00% (4.25%), Run 3: 1.21%
(4.10%), Run 4: 4.29% (4.93%).

2.5.　WM performance improves across runs

The mean (SE) accuracy rate across all participants and runs was 75.74%
(2.66%) and the mean reaction time (RT) was 1.38 s (0.02 s).

For memory trials, we explored the main effect of run (1-4) to check if
performance decreases in line with the trend seen for Glu (Fig. 3c). There were main
effects for both accuracy (F(3, 54) = 4.80, p = .005, η_p^2 = 0.21) and reaction time (F(3,
54) = 7.11, p < .001, η_p^2 = 0.28).

Mean accuracies (SE) per run were: Run 1: 70.72% (3.23%), Run 2: 77.14%
(2.51%), Run 3: 75.82% (3.42%), Run 4: 79.28% (2.85%). Accuracy was significantly
higher in Run 2 (t(18) = 3.17, p = .005, 95% CI = [2.17, 10.66], d = 0.73) and Run 4
(t(18) = 3.02, p = .007, 95% CI = [2.59, 14.51], d = 0.69) compared to Run 1. The
difference between Runs 1 and 3 trended towards significance prior to correction
(t(18) = 2.05, p = .055, 95% CI = [-10.32, 0.12], d = 0.47). No other comparisons
were significant. Bonferroni-corrected α = .008. See Fig. 3a.

Mean RTs per run were: Run 1: 1.46 s (0.03 s), Run 2: 1.37 s (0.03 s), Run 3:
1.35 s (0.03 s), Run 4: 1.33 s (0.03 s). Reaction times were significantly faster in Run
2 (t(18) = -3.95, p < .001, 95% CI = [-0.14, -0.04], d = 0.91), Run 3 (t(18) = -3.05, p =
.007, 95% CI = [-0.18, -0.03], d = 0.70), and Run 4 (t(18) = -3.37, p = .003, 95% CI =
[-0.21, -0.05], d = 0.77) compared to Run 1. No other comparisons were significant.
Bonferroni-corrected α = .008. See Fig. 3b.

Behavioural data in control trials was at ceiling and therefore not included in
any analyses above. Mean (SE) accuracy was 98.48% (1.02%) and reaction time
was 0.91 s (0.04 s).
280

3. Discussion
Functional MRS studies offer new insights into the dynamics of neural
processes that occur during cognitive tasks. Here, we developed an event-related
fMRS approach to characterise dynamic neurochemical changes in the posterior
MFC during an associative WM task. Glutamate concentration measured post-
stimulus exhibited distinct peaks in concentration following the onset of the encoding
(~1.08 s) and recall periods (~1.42 s), with both WM phases showing significant
increases in concentration of Glu compared to a control condition that accounted for
visual and motor demands, indicating a direct link to cognitive processes involved in
WM. Importantly, GABA modulation during encoding predicted task accuracy, with
greater increases in GABA concentration associated with better performance.
Together, these findings provide highly temporally precise evidence of phase-specific
excitatory and inhibitory dynamics during WM.
Our reconstructed time courses of Glu concentration during a cognitive task
are the most comprehensive to date, with a temporal resolution of ~0.3 s spanning 0-
2.5 s post-stimulus. They show a rise in Glu, peaking 1.08 s after stimulus
presentation during encoding and 1.42 s after probe presentation during recall and
returning to baseline within the 2.5 s. These dynamics extend beyond previous work
which identified peaks in concentration in the 0.3–1.0 s range [21,24]. Based on this, for
cognitive processes in MFC our results would recommend setting the acquisition
delay to ~1.0-1.5 s post-stimulus. Our results suggest that prior work investigating
cognitive processes [6,22,23] may have sampled Glu too early. Indeed, we did not find a
significant increase within 0-0.67 s.
Interestingly, the Glu peak at 1.42 s during recall aligns with reaction times
during memory trials (1.38 s), however increases in Glu cannot solely be attributed to
motor responses since no response was required during encoding. The close
alignment between Glu peak and RT could reflect processes following memory
retrieval, for instance internal monitoring processes which have been associated with

activity in the posterior MFC [30]. The time of the Glu peak during encoding (1.08s)

corresponds to the later end of a previously hypothesized Glutamatergic Response

Function (GRF) peaking 0.3-1.0 s after stimulus onset [19] and later than the peak of

0.5 s suggested by Koolschijn et al [17].

The apparent return towards baseline of the Glu response within 2.5 s during

encoding potentially suggests a transient response which may not reflect

maintenance. Maintenance may be served by the dlPFC instead [32–34]. Moreover, the

return to baseline within each WM phase indicates that we could detect

distinguishable responses to each phase, rather than a single prolonged response

across the WM block.

In our analysis of longer time windows, windowing improved the visibility of

the Glu response. When incorporating all sampling points, we observed some

evidence of increased Glu during encoding (1.4%) and recall (1.8%) compared to

control, significant prior to Bonferroni correction. As expected from the observed time

course, windowing to the central eight sampling points (0.67-1.83 s), thereby

focussing on the peaks of the Glu responses, revealed larger detectable increases in

Glu during encoding (2.5%) and recall (3.2%). The changes in Glu averaged across

this window are less than previous work in the hippocampus, which similarly

compared encoding and recall to a visuo-motor control task (4-10%) [35] and in the

visual cortex when comparing remembered to forgotten items (~5%) [18]. However,

similar magnitude changes were reported in the dlPFC during a WM task (2.7%) [16].

Over our measured time courses, we find maximal percent Glu changes of 5.4% and

5.3% during encoding and retrieval, respectively, relative to control. We may have

observed larger % increases had we performed a more classic event-related design,

intermixing control and WM trials, rather than a mixed-design. While we observed a

rise and fall of Glu over the 2.5 window, the magnitude of this transient response

may have been partially hidden by slower changes across the block of trials,

speculated to be involved in homeostatic changes or plasticity [36].

The increase in Glu during WM trials compared to control trials is consistent

with the increased fMRI activation for working memory over control trials in our fMRI

localiser (see Methods and Supplementary Methods). In contrast, Jelen and

colleagues [7] did not find a main effect of working memory load (0-back versus 2-

back) on Glx (Glu and glutamine) in the MFC, however, here we employed a different

type of WM task. We also acquired fMRS at 7T, which offers a larger signal-to-noise

ratio (SNR) and improved sensitivity to separate signals such as Gln and Glu [37].

344   Interestingly, individual increases in Glu did not correlate with WM

performance. The increase in Glu might therefore reflect either general increases in

attentional demands during the WM task, not directly tied to performance and/or it

could be linked to cognitive control processes [38] that are not (or to a lesser extent)

required in the control condition. The posterior MFC has been associated with

cognitive control processes before [e.g., 30]. The WM task used in this study involves

associative binding of different colours, shapes and locations, thereby creating a

representational competition and potential response conflict. Higher Glu levels during

the WM task compared to the simpler control task indicate that the metabolic

demand is tied to encoding and recalling complex associations, not just visual

perception or motor responses. Similar observations of increased Glx (Glu +

glutamine) in medial prefrontal cortex during an associative memory task involving a

longer delay period have been reported [39]. The increases in Glu with cognitive

control demands are supported by the raw Glu time courses (Fig. 1a), with increases

during WM encoding and recall (compared to baseline) but little change in the control

task. The control task used here elicited minimal or no conflict because the response

mapping was direct and stable. If increased cognitive control demands are managed

successfully, as in adaptive proactive control processes [40] leading to a resolution of

internal response conflicts before a response is executed, a direct association with

behavioural task performance is not expected. The decreasing size of Glu responses

over blocks (Fig. 3c) would also be in line with cognitive control processes: as

participants get more practiced to execute the WM task, some processes might

become more automatised (e.g. due to the development of strategies to memorise

associations), thereby decreasing cognitive control demands. An increase with

cognitive demands is consistent with evidence showing a positive correlation

between increased Glx and BOLD (blood oxygenation level dependent) signal in the

dlPFC and lower WM performance, which could reflect the increased neurovascular

demands of challenging cognitive processes [41].

372   The diminishing increase in Glu from the first to the third run, could not be

explained by build-up of Glu over the session as there was no difference in the

baseline Glu concentration measured over subsequent runs during the initial rest

period (8 spectra). The result may also be explained by repetition suppression –

repeating stimuli can result in a decrease in neural response that is detectable within

neuroimaging [42]. This effect has more recently been detected in the glutamate signal

using ^{1}H-MRS [21,24]. While no exact combination of shape, location, colour and probe

was repeated in our study, we may have observed repetition suppression to the

general type of WM trials.

In addition to its role as an excitatory neurotransmitter, Glu plays a key role in

cerebral metabolism through its involvement in the tricarboxylic acid (TCA) cycle,

where it can be used as a buffer. Glu is distributed intracellularly between the cytosol

and synaptic vesicles, with the majority residing in the cytosol [43]. These

compartments have been suggested to exhibit distinct T_2 relaxation times, with

vesicular Glu relaxing rapidly (~10 ms) compared to cystolic Glu (180–220 ms) at 3T

[44]. Consequently, given the relatively long echo time used in our sLASER acquisition

(TE = 80 ms), the observed Glu signal changes may be dominated by cystolic Glu

pools rather than vesicular components. Other fMRS studies employing cognitive

paradigms [16,18,35] have reported sustained increases in glutamate (Glu)

concentration that, in some cases, are larger than those observed in the present

study (2.7–10%). Notably, all these studies used short-TE MRS sequences (TE =

23–36 ms). In our work, we combined long-TE MRS with a rapidly sampled event-

related paradigm in an effort to probe this faster acting synaptic component of Glu.

However, we cannot separate the potential contribution of a metabolic Glu

component, which may also explain the absence of a correlation with cognitive

measures.

We did not observe significant group-level changes in GABA during either

encoding or recall compared to control trials. Similar findings of no change in GABA

levels have been reported in medial [39] and dorsolateral [41] prefrontal cortex during

performance of an associative memory task. However, at an individual level, we

observed that larger increases in GABA during encoding were associated with

enhanced task performance, with similar trends present during recall. This suggests

that dynamic, individual-level GABA fluctuations underpin aspects of WM

performance.

This study is the first to show a relationship between GABA dynamics within

MFC during the encoding phase of a WM task and performance. Previous work with

static MRS has highlighted a positive relationship between WM performance and

GABA quantified at rest from another frontal region - the dlPFC [45,46 although see 47,48].

Interestingly, Yoon et al [46] found that higher GABA levels were predictive of improved

performance at higher WM loads. Higher GABA may therefore be particularly

beneficial for accurate performance on demanding WM tasks such as the one used

here, potentially via suppression of task irrelevant information – a role already

observed in MFC through measuring the BOLD signal with fMRI [26,27]. In MFC, GABA

increases have been observed during interference tasks requiring inhibition [49] and

higher hippocampal GABA levels have been linked to improved performance during

a think/no-think paradigm in which task irrelevant information is suppressed [50].

Since Glu and GABA seem to be related to different types of cognitive

processes (see also review by Pasanta and colleagues [36]), it could be beneficial to

investigate Glu and GABA dynamics separately, rather than combining these

measures into excitatory/inhibition ratios [e.g., see 8] in the context of WM tasks.

To conclude, based on our findings, we can now better characterise the time

course of Glu responses in relation to different events in cognitive tasks, with the

change from control trials peaking around 1.08 s during encoding and 1.42 s during

recall. Our results also demonstrated the benefits and importance of selecting the

appropriate acquisition time to be maximally sensitive to Glu changes. Comparing

WM-related metabolite changes to a control condition allowed us to isolate the

changes associated with cognitive functions involved in WM over and above visual

and motor responses. Future theoretical work on the roles of Glu and GABA in WM

should integrate the findings that Glu levels in MFC are closely tied to WM demands

but do not predict performance, whereas GABA increases during the encoding phase

do predict task performance.

433 4. Methods

434 4.1. Participants

Twenty-one participants took part in the study, two were excluded due to poor

data quality (see data analysis), leaving a final sample of 19 participants (9 female,

10 male; one left-handed), aged 20-37 years (mean: 28.95, SD: 5.68 years). One

participant was red-green colourblind, however they could distinguish the colours

used in the task sufficiently and were not a significant outlier from either their working

memory performance or neurochemical levels so were retained for analysis.

No participants reported having a current neurological or psychiatric disorder,
recent medication use that could influence neurochemicals of interest, or a history of
significant head injury. All participants provided informed consent. The study was
approved by the local ethics committee (ethical approval number F1465).
445

4.2. Task

During the fMRS scan, participants completed four runs of an associative
visuospatial WM task, adapted from Belekou et al [51], with two blocks of memory and
two blocks of control trials per run, with each block containing 16 trials (see Fig. 4).
Participants completed 24 practice trials (12 memory task, 12 control task) before
scanning.
Custom shapes without obvious semantic associations were used as stimuli in
the associative working memory task. Each stimulus spanned an equal width and
height and could be presented in one of four colours: blue (RGB: 62, 132, 254),
green (RGB: 2, 200, 2), ochre (RGB: 208, 166, 3) or purple (RGB: 187, 106, 254).
Each WM trial started with a variable delay (grey screen) between 0.57 and
3.73 s. Participants were then shown four shapes presented at four horizontal
locations spread across the width of the screen in the upper half of the display (see
Fig. 4b, the encoding phase, 2.5 s). Three shapes were coloured, with participants
needing to encode the shape, location and colour of the items. The fourth shape was
grey, with the shape and location of the grey item varying across trials. The grey item
was a placeholder for the location and did not need to be actively encoded.
Participants were then presented with a scintillating, coloured mask (0.6 s) followed
by a variable inter-stimulus interval (ISI). Each of the 16 trials within a block had a
unique ISI ranging between 0.23 and 4.40 s. This was followed by the recall phase,
in which either a white shape or location cue was presented above four response
colour options (recall phase, 2.5 s). Participants had to recall the colour of the shape
or the colour presented at the cued location from the encoding phase (response
window: 2.5 s). The recall phase was again followed by a coloured mask (0.6 s)
before the next trial began. More details about the ITIs and ISIs can be found in the
data acquisition section. Memory trials spanned an average of 10 s, with a total block
duration of 160 s.

The control task used the same stimuli as the associative WM task. Within a
control trial participants were presented with three grey items and one coloured item
above the four response options. Participants responded to the colour of the item.
The control task was designed to elicit similar visual processes and motor responses
as the WM task to control for these processes in the planned analysis. This screen
was presented for 2.5 s followed by a 0.6 s mask. Control trials spanned an average
of 5 s including a variable ITI (0.57-4.07 s), with a total block duration of 80 s.
Timings of the control trials followed the timings of the encoding phase in the
memory trials.
Within each of the four runs of the task, participants were presented with two
blocks of 16 memory trials and two blocks of 16 control trials, presented in a
counterbalanced order. Blocks were separated by a 30 s rest period, and each run
began with 37.4 s rest and ended with a 30 s rest. Each run therefore took a total of
637.4 s (10 minutes, 37.4 seconds).
The order of the trials was pseudo-randomised across participants such that
no participant encountered the same trial twice, and different participants were
presented with different subsets of trials. Each memory block contained an equal
number of shape and location cues, and an equal number of the four options as the
target colour. Timings of the trials within the block were pre-defined by two sets of
possible timings (see fMRS data acquisition for more details, see also Fig. 4c).

Figure 4

Plot of run and trial structures.

Note: (**a**) A depiction of two runs of the task, showing two of the four counterbalancing options used in the study. Purple diamonds and red stars refer to the trial timing sets depicted in **b**. (**b**) Depictions of a memory trial (left) and control trial (right). In the memory trial, participants may be presented with a location probe (top) or shape probe (bottom). Arrows showing the correct answer are for illustration purposes only. ITI = inter-trial interval, ISI = inter-stimulus interval. (**c**) Coloured boxes show timings of stimuli in relation to sample acquisition (dashed vertical lines) for the memory trials (left) and control trials (right). The upper and lower plots show the two sets of predefined timings. Each row represents a trial within a block, starting with the top row. Textured boxes represent the mask duration. S1 = acquisition of the first MRS sample within a trial, which occurs during encoding in the memory trials, and the probe in the control trials. S2 = acquisition of the second MRS sample in the memory trials. The grey lines show when then previous trial ended.

4.3. Stimulus Apparatus

The task was presented using an MR compatible projector (PROPixx MRI
from VPixx), which displayed the stimuli at 120 Hz on a screen suspended vertically
above the foot end of the scanner bed, with participants using mirrored glasses to
view the display. Screen resolution was 1980 x 1020 pixels, with the projection
subtending 62 by 32 cm at approximately 2.4 m, although the lower half of the
screen was generally not visible due to the participant's body partly obstructing the
field of view. The researchers ensured participants could see the stimuli before
commencing the experiment. Participants responded to the task using the four
fingers of their right-hand and the RESPONSEPixx 5-button response box, with each
button corresponding to one of the four response options presented on the screen.
505

4.4. MR Data acquisition

Data were acquired on a 7T MR scanner (Phillips Achieva) using a 32 channel
receive and 2 channel transmit head coil (Nova Medical).
An anatomical 3D T1-weighted gradient echo acquisition was first acquired
and used for subsequent positioning of MRS voxel, TR = 4.9 s, voxel size = 0.85 x
0.85 x 1 mm^3, TE = 2.17 ms, flip angle = 7°, field of view = 246 x 246 x 180 mm^3, T_{acq}

= 119 s. All single-voxel spectroscopy measurements were acquired with semi-

LASER localisation and TR = 5s, voxel size = 20 x 20 x 20 mm^3, spectral points =

2048, bandwidth = 6000 Hz, TE = 80 ms, NSA = 128. TE=80 ms was chosen to

optimise sensitivity to both Glu and GABA at 7 T in line with a recent publication [52].

Refocussing was performed using GOIA-WURST pulses [53]. A DOTCOPS crusher

gradient scheme [54] with interleaved outer volume suppression (OVS) and VAPOR

water suppression (BW = 140 Hz) [55] were performed.

Voxel placement was informed by an fMRI localiser (see Supplementary

Methods) captured in five separate participants using the task, with slightly modified

timings and trial numbers to better suit fMRI acquisition. The MRS voxel was placed

in the pre-SMA region of the MFC, parallel to but just superior to the cingulate cortex.

The most posterior edge of the voxel was positioned just anterior (~2mm) of a

vertical line drawn from the AC, perpendicular to the AC-PC line, although voxel

placement varied slightly per-participant to avoid larger blood vessels (see Fig. 5 for

positioning).

Figure 5

FMRI-informed positioning the fMRS voxel.

Note: Heatmap of voxel placements over participants (green), superimposed with fMRI activation from the localiser study (see Supplementary Materials). The orange regions of activation show a greater BOLD response to memory trials over control trials. Blue regions show a greater BOLD signal to control over memory trials.

In each of the four runs, a total of 128 spectra were acquired. Within each

memory trial, two spectra were acquired, one during the encoding phase and one

during recall. During the control trials, a single acquisition was made per trial.

Examples of spectral quality and metabolite fits can be found in Fig. 6.

The onsets of the stimuli were varied such that across the 16 trials in a block,

the spectral acquisitions aligned with 16 equally spaced intervals between stimulus

onset (0 s) and stimulus offset (2.5 s); every 0.17 s, acquired in a non-sequential

order. The timings were pre-defined, with two unique solutions designed to be

maximally different (see Fig. 4c and Supplementary Materials). The sampling method

of the spectral acquisitions was designed and applied to acquire neurochemical

responses at different time lags after stimulus onset and to be able to estimate the

glutamate response function associated with this task. The same order of sampling

times from the encoding phase was used for the control trials as well (see Fig. 4c).

Within a run, one memory and one control block followed each of the two solutions,

with the order of the solutions being counterbalanced across runs.

Figure 6

Sample spectra from one participant using 8 (**a**) and 128 (**b**) averages per fit, and composition of neurochemicals (**c**).

Note: Example spectra acquired from a single subject, averaging the initial 8 spectra in the first rest period (**a**), and averaging over all 128 spectra acquired in a single run (**b**). All spectra have been pre-processed (black) and then fit (red). (**c**) shows the individual fits for each metabolite for included in the analysis of (b) with GABA and Glu highlighted in blue.

4.5. fMRS data analysis

All spectral processing was performed in FSL-MRS V2.4.0 [56] in Python

V3.12.7. Each run was acquired as separate scan with corresponding raw data

saved for processing. Pre-processing of each run involved coil combination, eddy

current-correction, frequency and phase alignments, Hankel Lanczos Singular Value

Decomposition (HLSVD) and zero-order automatic phase corrections. The frequency

and phase alignments were performed around the NAA singlet (1.7-2.3 ppm). Water

reference spectra were used for pre-processing steps and also fit to obtain linewidth

(mean: 8, SD: 3 Hz). Spectral data with water reference linewidths >20 Hz were

rejected (one participant).

Fitting was performed using a truncated-Newton algorithm. Basis sets were

generated using FSL-MRS via density matrix simulations using a sLASER sequence

with real pulse information and interpulse timings. with TE=80 ms, and a 90°

spredrex and 180° adiabatic GOIA pulses. The metabolites generated were Alanine

(Ala), Ascorbate (Asc), Aspartate (Asp), Creatine (Cr), Creatine CH2 group (CrCH2),

Gamma-aminobutyric acid (GABA), Glucose (Glc), Glutamine (Gln), Glutamate

(Glu), Glycine (Gly), Glycerophosphocholine (GPC), Glutathione (GSH), Myo-Inositol

(Ins), Lactate (Lac), N-acetylaspartate (NAA), N-acetylaspartylglutamate (NAAG),

Phosphocholine (PCh), Phosphocreatine (PCr), Phosphoethanolamine (PE), Scyllo-

Inositol (Scyllo) and Tau. No baseline was used and simulated macromolecular

components at 0.9 and 2.1 ppm were implemented. Each macromolecule was

assigned to a separate group for linewidth and phase compared to the rest of the

other metabolites. To form a high SNR fit during initialisation for further analysis, all

spectra from the four runs per participant are averaged together and fit together, an

appropriate scaling is obtained from this data as well per participant

4.5.1. Time course analysis:

The time course analysis was achieved by assigning spectra for the 16

sampling points into to eight bins (or timepoints), with the mean sampling time for

each bin as follows: 0.083, 0.417, 0.75, 1.083, 1.417, 1.750, 2.083, 2.417 s. For

each bin, that resulted in a total of four spectral acquisitions per run. The time

courses were initially converted to a percentage change compared to the initial rest

spectra for each run and then averaged across runs, to observe the dynamic change

in Glu during each WM phase (encoding, recall) and during the control trials. One-

sample t-tests were used to assess whether the concentration during each of the

WM phases and control trials was significantly different from the initial rest period

(baseline) at any timepoint. Bonferroni-corrected α = .002.

For each timepoint and run separately, the Glu concentrations during

encoding and recall were additionally calculated as a percentage change from the

control condition, without any normalisation to the initial rest spectra, to isolate

processes involved in WM. Percent changes were then averaged over run

separately for each participant. One-sample t-tests were again employed to assess if
Glu levels during encoding and recall were significantly different at any timepoint
compared to control. T-tests were two-sided. Bonferroni-corrected α = .003.

4.5.2.   WM phase analysis:

In addition to the time course analysis, we performed event-based analysis
averaging over larger time windows. Spectra for each run and task phase (control,
encoding, recall) separately were averaged over either all 16 sampling points
(number of spectral averages, NSA=32/run/phase), the middle eight
(NSA=16/run/phase), or the first and last four sampling points (NSA=16/run/phase,
see Results) prior to fitting. The change in Glu and GABA concentrations were
calculated as the percentage change during encoding and recall compared to the
control task to capture memory-related changes in concentrations over the visual
and motor responses in the control trials. No internal, within-spectra normalisation,
such as to total creatine, was performed. The normalisation to control was performed
for each run separately and then averaged over runs to assess for effects of WM
phase, or averaged over encoding and recall assessing the main effect of run.
We assessed whether 1) the concentration changes during encoding and
recall differed from control trials (one-sample t-tests), 2) the concentration changes
differed between encoding and recall (paired-sample t-tests), 3) the concentration
changes were significantly correlated across encoding and recall (Pearson's
correlation). T-tests were two-sided.
GABA levels for one participant were removed due to being a significant
outlier during recall (>3 SD from group mean).
610

4.6. Behavioural data analysis

All trials were retained in the behavioural data analysis as no trials showed a
pre-emptive response (<0.3 s). Of 256 trials, including both memory and control
trials, a maximum of eight trials were missed by a single participant. Missed trials in
a WM task may reflect a lapse in attention or may reflect subjective task difficulty.
RT-based outliers in individual trials were negligible (max. 5 trials for an individual
participant). Including trial wise outliers should therefore not compromise data
quality.

To compare accuracy and reaction time over the four runs, within-subjects one-way ANOVAs and post-hoc paired samples t-tests were employed. T-tests were two-sided.

5. Acknowledgements

This work was supported by the Wellcome Trust [226715/Z/22/Z]. AB would like to acknowledge the Royal Academy of Engineering. We thank Prof. Helen Barron for their insight into study design during the early stages of the project, Dr. Josh Khoo for creating an initial version of the experimental script, and Siavash Rakhtshah for assisting with MR scanning and with eligibility queries.

6. Conflict of interest

The authors declare no competing interests.

7. Author contributions

All authors were involved in task-design. AB and DC optimised fMRS data acquisition and analysis. RE created the experimental script. DC and RE collected and analysed the data with supervision from AB, CD and KD. RE, DC, CD, KD and AB wrote the manuscript. All authors reviewed and approved the manuscript. DC and RE have contributed equally to the manuscript and therefore share first authorship.

8. References

1. Patel, T. & Talcott, J. B. Moderate relationships between NAA and cognitive ability in healthy adults: implications for cognitive spectroscopy. *Front. Hum. Neurosci.* **8**, (2014).

2. Reid, M. A. *et al.* 7T Proton Magnetic Resonance Spectroscopy of the Anterior Cingulate Cortex in First-Episode Schizophrenia. *Schizophr. Bull.* **45**, 180–189 (2019).

3. Rowland, L. M. *et al.* Frontal Glutamate and γ-Aminobutyric Acid Levels and Their Associations With Mismatch Negativity and Digit Sequencing Task Performance in Schizophrenia. *JAMA Psychiatry* **73**, 166–174 (2016).

4. White, T. L. *et al.* The neurobiology of wellness: 1H-MRS correlates of agency,

flexibility and neuroaffective reserves in healthy young adults. *NeuroImage* **225**,

117509 (2021).

5. Jelen, L. A., King, S., Mullins, P. G. & Stone, J. M. Beyond static measures: A

review of functional magnetic resonance spectroscopy and its potential to

investigate dynamic glutamatergic abnormalities in schizophrenia. *J.*

Psychopharmacol. (Oxf.) **32**, 497–508 (2018).

6. Craven, A. R. *et al.* GABA, glutamatergic dynamics and BOLD contrast assessed

concurrently using functional MRS during a cognitive task. *NMR Biomed.* **37**,

e5065 (2024).

7. Jelen, L. A. *et al.* Functional magnetic resonance spectroscopy in patients with

schizophrenia and bipolar affective disorder: Glutamate dynamics in the anterior

cingulate cortex during a working memory task. *Eur. Neuropsychopharmacol.* **29**,

222–234 (2019).

8. Saviola, F. *et al.* Disentangling metabolic and neurovascular timescales

supporting cognitive processes. *Proc. Natl. Acad. Sci.* **122**, e2506513122 (2025).

9. Bednařík, P. *et al.* Neurochemical responses to chromatic and achromatic stimuli

in the human visual cortex. *J. Cereb. Blood Flow Metab.* **38**, 347–359 (2018).

10. Chen, C. *et al.* Activation induced changes in GABA: Functional MRS at 7 T with

MEGA-sLASER. *NeuroImage* **156**, 207–213 (2017).

11. DiNuzzo, M. *et al.* Perception is associated with the brain's metabolic response to

sensory stimulation. *eLife* **11**, e71016 (2022).

12. Ip, I. B. *et al.* Combined fMRI-MRS acquires simultaneous glutamate and BOLD-

fMRI signals in the human brain. *NeuroImage* **155**, 113–119 (2017).

13. Mangia, S. *et al.* Sustained Neuronal Activation Raises Oxidative Metabolism to a

New Steady-State Level: Evidence from 1H NMR Spectroscopy in the Human

Visual Cortex. *J. Cereb. Blood Flow Metab.* **27**, 1055–1063 (2007).

14. Mekle, R. *et al.* Detection of metabolite changes in response to a varying visual

stimulation paradigm using short-TE 1H MRS at 7 T. *NMR Biomed.* **30**, e3672

(2017).

15. Volovyk, O. & Tal, A. Increased Glutamate concentrations during prolonged motor

activation as measured using functional Magnetic Resonance Spectroscopy at

3T. *NeuroImage* **223**, 117338 (2020).

16. Woodcock, E. A., Anand, C., Khatib, D., Diwadkar, V. A. & Stanley, J. A. Working

Memory Modulates Glutamate Levels in the Dorsolateral Prefrontal Cortex during

1H fMRS. *Front. Psychiatry* **9**, (2018).

17. Koolschijn, R. S., Clarke, W. T., Ip, I. B., Emir, U. E. & Barron, H. C. Event-related

functional magnetic resonance spectroscopy. *NeuroImage* **276**, 120194 (2023).

18. Koolschijn, R. S. *et al.* Memory recall involves a transient break in excitatory-

inhibitory balance. *eLife* **10**, e70071 (2021).

19. Mullins, P. G. Towards a theory of functional magnetic resonance spectroscopy

(fMRS): A meta-analysis and discussion of using MRS to measure changes in

neurotransmitters in real time. *Scand. J. Psychol.* **59**, 91–103 (2018).

20. Lea-Carnall, C., El-Deredy, W., Stagg, C. J., Williams, S. R. & Trujillo-Barreto, N.

A mean-field model of glutamate and GABA synaptic dynamics for functional

MRS. *NeuroImage* **266**, 1–18 (2023).

21. Apšvalka, D., Gadie, A., Clemence, M. & Mullins, P. G. Event-related dynamics of

glutamate and BOLD effects measured using functional magnetic resonance

694  spectroscopy (fMRS) at 3 T in a repetition suppression paradigm. *NeuroImage*

695  **118**, 292–300 (2015).

696 22. Craven, A. R. *et al.* GABA, glutamate dynamics and BOLD observed during

697  cognitive processing in psychosis patients with hallucinatory traits. *Sci. Rep.* **15**,

698  19466 (2025).

699 23. Mohammadi, H. *et al.* Unveiling Glutamate Dynamics: Cognitive Demands in

700  Human Short-Term Memory Learning Across Frontal and Parieto-Occipital

701  Cortex: A Functional MRS Study. *J. Biomed. Phys. Eng.* **14**, 519–532 (2024).

702 24. Lally, N. *et al.* Glutamatergic correlates of gamma-band oscillatory activity during

703  cognition: A concurrent ER-MRS and EEG study. *NeuroImage* **85**, 823–833

704  (2014).

705 25. Baddeley, A. D. & Hitch, G. Working Memory. in *Psychology of Learning and*

706  *Motivation* (ed. Bower, G. H.) vol. 8 47–89 (Academic Press, 1974).

707 26. Danielmeier, C., Eichele, T., Forstmann, B. U., Tittgemeyer, M. & Ullsperger, M.

708  Posterior Medial Frontal Cortex Activity Predicts Post-Error Adaptations in Task-

709  Related Visual and Motor Areas. *J. Neurosci.* **31**, 1780–1789 (2011).

710 27. Danielmeier, C. *et al.* Acetylcholine Mediates Behavioral and Neural Post-Error

711  Control. *Curr. Biol.* **25**, 1461–1468 (2015).

712 28. Klein, T. A. *et al.* Neural correlates of error awareness. *NeuroImage* **34**, 1774–

713  1781 (2007).

714 29. Rushworth, M. F., Buckley, M. J., Behrens, T. E., Walton, M. E. & Bannerman, D.

715  M. Functional organization of the medial frontal cortex. *Curr. Opin. Neurobiol.* **17**,

716  220–227 (2007).

717 30. Ullsperger, M., Danielmeier, C. & Jocham, G. Neurophysiology of Performance

718  Monitoring and Adaptive Behavior. *Physiol. Rev.* **94**, 35–79 (2014).

31. Kamiński, J. *et al.* Persistently active neurons in human medial frontal and medial

temporal lobe support working memory. *Nat. Neurosci.* **20**, 590–601 (2017).

32. Barbey, A. K., Koenigs, M. & Grafman, J. Dorsolateral prefrontal contributions to

human working memory. *Cortex* **49**, 1195–1205 (2013).

33. Goldman-Rakic, P. S. Circuitry of the frontal association cortex and its relevance

to dementia. *Arch. Gerontol. Geriatr.* **6**, 299–309 (1987).

34. Goldman-Rakic, P. S. Regional and cellular fractionation of working memory.

Proc. Natl. Acad. Sci. **93**, 13473–13480 (1996).

35. Stanley, J. A. *et al.* Functional dynamics of hippocampal glutamate during

associative learning assessed with in vivo 1H functional magnetic resonance

spectroscopy. *NeuroImage* **153**, 189–197 (2017).

36. Pasanta, D. *et al.* Functional MRS studies of GABA and glutamate/Glx – A

systematic review and meta-analysis. *Neurosci. Biobehav. Rev.* **144**, 104940

(2023).

37. Tkáč, I., Öz, G., Adriany, G., Uğurbil, K. & Gruetter, R. In vivo 1H NMR

spectroscopy of the human brain at high magnetic fields: Metabolite

quantification at 4T vs. 7T. *Magn. Reson. Med.* **62**, 868–879 (2009).

38. Botvinick, M. M., Braver, T. S., Barch, D. M., Carter, C. S. & Cohen, J. D. Conflict

monitoring and cognitive control. *Psychol. Rev.* **108**, 624–652 (2001).

39. Thielen, J.-W. *et al.* The increase in medial prefrontal glutamate/glutamine

concentration during memory encoding is associated with better memory

performance and stronger functional connectivity in the human medial prefrontal–

thalamus–hippocampus network. *Hum. Brain Mapp.* **39**, 2381–2390 (2018).

40. Braver, T. S. The variable nature of cognitive control: a dual mechanisms

framework. *Trends Cogn. Sci.* **16**, 106–113 (2012).

41. Oh, H. *et al.* A preliminary study of dynamic neurochemical changes in the

dorsolateral prefrontal cortex during working memory. *Eur. J. Neurosci.* **59**, 2075–

2086 (2024).

42. Grill-Spector, K. & Malach, R. fMR-adaptation: a tool for studying the functional

properties of human cortical neurons. *Acta Psychol. (Amst.)* **107**, 293–321

(2001).

43. Walker, M. C. & van der Donk, W. A. The many roles of glutamate in metabolism.

J. Ind. Microbiol. Biotechnol. **43**, 419–430 (2016).

44. Emeliyanova, P., Parkes, L. M., Williams, S. R. & Lea-Carnall, C. Evidence for

biexponential glutamate T2 relaxation in human visual cortex at 3T: A functional

MRS study. *NMR Biomed.* **37**, e5240 (2024).

45. Ragland, J. D. *et al.* Disrupted GABAergic facilitation of working memory

performance in people with schizophrenia. *NeuroImage Clin.* **25**, 102127 (2020).

46. Yoon, J. H., Grandelis, A. & Maddock, R. J. Dorsolateral Prefrontal Cortex GABA

Concentration in Humans Predicts Working Memory Load Processing Capacity.

J. Neurosci. **36**, 11788–11794 (2016).

47. Andersson, P., Li, X. & Persson, J. Hippocampal and prefrontal GABA and

glutamate concentration contribute to component processes of working memory

in aging. *Cereb. Cortex* **35**, bhaf105 (2025).

48. Michels, L. *et al.* Frontal GABA Levels Change during Working Memory. *PLOS*

ONE **7**, e31933 (2012).

49. Kühn, S. *et al.* Neurotransmitter changes during interference task in anterior

cingulate cortex: evidence from fMRI-guided functional MRS at 3 T. *Brain Struct.*

Funct. **221**, 2541–2551 (2016).

50. Schmitz, T. W., Correia, M. M., Ferreira, C. S., Prescot, A. P. & Anderson, M. C.

Hippocampal GABA enables inhibitory control over unwanted thoughts. *Nat.*

Commun. **8**, 1311 (2017).

51. Belekou, A., Katshu, M. Z. U. H., Dundon, N. M., d'Avossa, G. & Smyrnis, N.

Spatial and non-spatial feature binding impairments in visual working memory in

schizophrenia. *Schizophr. Res. Cogn.* **32**, 100281 (2023).

52. Finkelman, T., Furman-Haran, E., Paz, R. & Tal, A. Quantifying the excitatory-

inhibitory balance: A comparison of SemiLASER and MEGA-SemiLASER for

simultaneously measuring GABA and glutamate at 7T. *NeuroImage* **247**, 118810

(2022).

53. Tannús, A. & Garwood, M. Improved Performance of Frequency-Swept Pulses

Using Offset-Independent Adiabaticity. *J. Magn. Reson.* **120**, 133–137 (1996).

54. Landheer, K. & Juchem, C. Dephasing optimization through coherence order

pathway selection (DOTCOPS) for improved crusher schemes in MR

spectroscopy. *Magn. Reson. Med.* **81**, 2209–2222 (2019).

55. Tkáč, I., Starčuk, Z., Choi, I.-Y. & Gruetter, R. In vivo 1H NMR spectroscopy of rat

brain at 1 ms echo time. *Magn. Reson. Med.* **41**, 649–656 (1999).

56. Clarke, W. T., Ligneul, C., Cottaar, M., Ip, I. B. & Jbabdi, S. Universal dynamic

fitting of magnetic resonance spectroscopy. *Magn. Reson. Med.* **91**, 2229–2246

(2024).

788