Aberrant GPX4 processing reveals its critical roles in maintaining ROS homeostasis in

Citrus

Barry, Shannon[1,2], Wang, Xin[2], & Wang, Nian[1,2]

[1] Citrus Research and Education Center, Institute of Food and Agricultural Sciences, University

of Florida, Lake Alfred, FL 33850, U.S.A.

[2] Department of Microbiology and Cell Science, Institute of Food and Agricultural Sciences,

University of Florida, Gainesville, FL 32611. U.S.A.

Corresponding author: nianwang@ufl.edu

Abstract

Glutathione peroxidases (GPXs) are widely recognized as key antioxidants that mitigate

oxidative stress by detoxifying reactive oxygen species (ROS). However, GPXs are largely

uncharacterized in citrus. Here, we demonstrated that *Citrus sinensis* contains four GPX proteins

(CsGPX1–4). Unexpectedly, overexpression of *CsGPX4*, a homolog of *AtGPX8* in *Arabidopsis*,

in citrus resulted in typical oxidative stress phenotypes including severe growth inhibition,

chlorosis, and elevated intracellular ROS accumulation. Transmission electron microscopy

(TEM) analysis further revealed stress responses at cellular level. Whole genome shot gun

sequencing analysis showed that T-DNA insertion occurs in the UTR of *SWEET2* gene, which is

unlikely to be responsible for the oxidative stress phenotypes. Immunoblotting revealed that

CsGPX4 accumulates as a truncated protein in citrus, in contrast to the full-length version

expressed in *Nicotiana benthamiana*. MALDI-TOF assays further confirmed the truncation of

CsGPX4 in the transgenic line with the predicted cleavage site between L115-K117. This

truncation was associated with altered subcellular localization, shifting from cytoplasmic and

nuclear distribution in *N. benthamiana* to membrane association in citrus. Proteomic profiling

further indicated extensive reprogramming of pathways involved in detoxification, cytoskeletal

stability, hormone signaling, and cell wall modification. Our data suggests that *de facto*

overexpression of truncated CsGPX4 may have dominant-negative effects on proteins interacting

with CsGPX4, thus interfering with their normal functions. In conclusion, our study

demonstrates CsGPX4 as a critical regulator of redox homeostasis and ROS homeostasis in

citrus and reveals selective truncation of CsGPX4 as a unique proteolytic or regulatory strategies

in such processes.

Keywords: Citrus sinensis, Overexpression, CsGPXs, CsGPX4, AtGPX8, AtGPXs, transient

expression, proteomics, post-translational truncation, redox homeostasis, species-specific

Introduction

Pathogens, insects, and environmental stressors contribute significantly to global crop yield

losses, posing an escalating threat to food security (Bailey-Serres et al., 2019; Lesk et al., 2016).

As the world population is projected to approach 10 billion by 2050, agricultural productivity

must increase by an estimated 60% to meet future food demands (FAO, 2019; Fedoroff, 2015).

This challenge underscores the urgent need to understand plant stress responses and develop

more resilient crop varieties. A common physiological consequence of both biotic and abiotic

stress is the overaccumulation of reactive oxygen species (ROS), including superoxide anion

($O_2 \cdot^-$), hydrogen peroxide (H_2O_2), singlet oxygen ($^1O_2$), and hydroxyl radicals ($\cdot OH$). These

molecules, though generated as byproducts of core metabolic processes such as photosynthesis,

respiration, and nitrogen assimilation, accumulate to cytotoxic levels under stress conditions

(Mittler, 2017; Wang et al., 2023). While excessive ROS can damage nucleic acids, proteins,

lipids, and membranes, they also function as key signaling molecules that regulate plant

development, immunity, and acclimation responses (Suzuki et al., 2012; Mitter et al., 2022).

Thus, maintaining redox homeostasis is essential for plant survival and adaptation under adverse

conditions.

Plants have evolved a complex antioxidant defense system comprising enzymatic and non-

enzymatic components to mitigate ROS accumulation. Major enzymatic scavengers include

superoxide dismutase (SOD), catalase (CAT), ascorbate peroxidase (APX), glutathione

peroxidase (GPX), glutathione reductase (GR), peroxiredoxins (PRXs), thioredoxins (TRXs),

and dehydroascorbate reductase (DHAR), which operate across distinct subcellular

compartments to fine-tune ROS levels and maintain cellular redox balance (Wang et al., 2023).

Among these, GPXs are thiol-based peroxidases that detoxify peroxides by using either

glutathione (GSH) or thioredoxins (Trx) as electron donors (Bela et al., 2015; You and Chan,

2015). Notably, plant GPXs differ from their animal counterparts in both structure and cofactor

usage—plant GPXs use cysteine (Cys) as the active site residue and prefer Trx, whereas animal

GPXs use selenocysteine (SeCys) and primarily depend on GSH (Islam et al., 2015; Wang et al.,

2023). Functionally, GPXs play important roles in plant responses to abiotic stresses such as

drought, salinity, extreme temperatures, and oxidative conditions. For example, overexpression

of *GPX5* in *Salvia miltiorrhiza* improved drought tolerance (Zhang et al., 2019), while knockout

of *GPX3* in *Oryza sativa* resulted in excessive H_2O_2 accumulation (Passaia et al., 2013).

GPX8 in particular has been implicated in protecting nuclear DNA and other organelles from

oxidative stress in *Arabidopsis* (Gaber et al., 2012). Overexpression of *AtGPX8* enhanced

oxidative tolerance, whereas its loss impaired root development and reduced stress resilience.

However, in woody species such as *C. sinensis*, functional studies of *GPX* genes remain limited.

For instance, Faltin et al. (2010) attempted to overexpress a phospholipid hydroperoxide GPX

(PHGPx) in *C. sinensis* but were only successful after abolishing enzymatic activity, indicating

potential toxicity or instability under overexpression conditions. In *Nicotiana benthamiana*,

overexpression of GPX8 alleviated cadmium-induced oxidative stress and preserved

photosynthetic pigments (Peng et al., 2024), highlighting the context-dependent functions of

GPXs.

In this study, we initially aimed to overexpress *CsGPX4*, a homologue of *AtGPX8*, to mitigate

oxidative stress caused by *Candidatus* Liberibacter asiaticus (CLas), which causes citrus

Huanglongbing (HLB). CLas is known to cause ROS accumulation, which contributes to phloem

cell death and subsequent HLB symptoms (Ma et al., 2022). GPXs were reported to be involved

in preventing oxidant-induced programmed cell death (PCD) (Passaia et al., 2014).

Unexpectedly, CsGPX4 overexpression resulted in the accumulation of a post-translationally

truncated protein isoform specific to *C. sinensis*, but not in *N. benthamiana*. This truncation was

associated with increased H_2O_2 accumulation, altered subcellular localization, chlorosis,

dwarfism, and ultrastructural cellular abnormalities. To our knowledge, this is the first report of a

plant antioxidant enzyme undergoing species-specific post-translational truncation in response to

overexpression, leading to a distinct stress-related phenotype.

Results

Identification of *GPX* genes in *C. sinensis*

To identify *GPX* genes in *C. sinensis*, we conducted BLAST search with the eight GPX proteins

in *Arabidopsis*. Only 4 *GPX* genes were identified in *C. sinensis* (hereafter *CsGPX1-4*) (**Fig. 1A,**

Supplementary Table 1). Three of the four *CsGPX* genes (*CsGPX1*, *CsGPX2*, and *CsGPX4*)

reside on chromosome 5, whereas *CsGPX3* is located on chromosome 4 (**Fig. 1B**). The close

proximity of *CsGPX2* and *CsGPX4* on chromosome 5 suggests a possible duplication event.

CsGPX proteins range from 170 to 245 amino acids, with a corresponding molecular weight of

19.3-27 kDa. CsGPX4 is the smallest GPX protein in *C. sinensis.* All CsGPXs have negative

GRAVY values which indicate hydrophilic characteristic consistent with GPX families in other

plant species (Khan et al., 2024). Interestingly, CsGPX4 displayed an acidic pI of 4.76, in

contract to the basic pI values of >7.9 observed in other CsGPX proteins. In addition, CsGPX1,

CsGPX2, CsGPX3, and CsGPX4 were predicted to localize in the endoplasmic reticulum (ER),

mitochondria, plastids and cytoplasm, respectively (**Supplementary Table 1**).

On average, the CsGPXs have approximately 50% to 75% similarity (**Fig. 1C**). Multiple

sequence alignment revealed conserved regions of CsGPX proteins including the three conserved

motifs in plants: VNAS[R/K/Q]CG, LAFPCNQF, and WNF(S/T)KF (Churin et al., 1999). The

catalytic tetrad (cysteine (C), glutamine (Q), tryptophan (W), and asparagine (N)) of GPX was

observed (Tosatto et al., 2008). The three conserved Cys residues in the active sites of plant

GPXs were also present in citrus GPXs (**Fig. 1D**). All the four CsGPX proteins contain a

conserved thioredoxin (TRX)-like fold (**Fig. 1E**), consistent with that plant GPXs prefer TRX

rather than glutathione as a reducing agent (Gaber et al., 2012). AlphaFold 3 structural

predictions further revealed a well-resolved catalytic core with high confidence of the CsGPX

proteins, while the N-terminal regions exhibited low confidence.

Overexpression of *CsGPX4* in citrus leads to oxidative stress phenotypes

CLas triggers ROS accumulations, which contributes to phloem cell death and subsequent HLB

symptoms (Ma et al., 2022). GPXs are known to play important roles in preventing oxidant-

induced programmed cell death (Passaia 2014). A previous study reported that transgenic

overexpressing *AtGPX8*, a homologue of *CsGPX4* in *Arabidopsis* (**Fig. 1A**), increases plant

tolerance to oxidative stress (Gaber et al., 2012). Here, we investigated whether overexpression

of *CsGPX4* in *C. sinensis* can inhibit oxidative stress triggered by CLas. Specifically, CsGPX4

shares 79% identity and 90% similarity with AtGPX4 (**Fig. S1A**). The tertiary structures of

CsGPX4 and AtGPX8 predicated by Alphafold3 showed that they share a highly conserved TRX

fold (**Fig 1F; S1B**). Superimposition of these two models showed a root mean square deviation

(RMSD) of 0.364 Å, indicating a strong three-dimensional similarity (**Fig. S1B**).

We generated two stable *C. sinensis* lines overexpressing *CsGPX4* (GPX4-OX) driven by the

Cauliflower mosaic virus (CaMV) 35S promoter (**Fig. 2A, B**) via *Agrobacterium* mediated

transformation, which were identified by GFP fluorescence and further confirmed through

reverse transcription PCR (RT-PCR) (**Fig. 2D)**. Surprisingly, both GPX4-OX transgenic plants

exhibited oxidative stress phenotypes including chlorosis and reduced growth compared to

empty vector transformed controls (EV) (**Fig. 2B**). Transmission electron microscopy (TEM)

analysis revealed striking ultrastructural difference in midrib cells of CsGPX4-OX lines

compared with EV controls (**Fig. 3**). The EV control plant cells displayed typical healthy

architecture including compact cytoplasm, continuous membranes, and well-organized vacuoles.

In contrast, both CsGPX4-OX lines exhibited a progressive structural deterioration. At the

whole-cell level, CsGPX4-OX cells showed irregular cell boundaries and reduced cytoplasmic

density relative to the well-defined walls and dense cytoplasm of control plants (**Fig. 3A**).

Vacuolar morphology was also disrupted with CsGPX4-OX lines displaying clear deformities in

the tonoplast and intracellular membranes (**Fig. 3B**). Cytoplasmic organization was

compromised by the extensive vesiculation and disordered intracellular contents in CsGPX4-OX

(**Fig. 3C**). These abnormalities were accompanied by fragmentation of membrane systems, loss

of membrane continuity, and accumulation of electron dense materials, hallmarking of severe

cellular stress in CsGPX4-OX, but not in the EV control (**Fig. 3D**).

CsGPX4 undergoes citrus-specific truncation and membrane-associated localization

Previous studies demonstrate that overexpression of *AtGPX8* driven by the 35S promoter in

Arabidopsis (Gaber et al., 2012) and knockout of *AtGPX8* had no significant effects on plant

growth and development (Passaia et al., 2014). Because of the high similarity between AtGPX8

and CsGPX4, we anticipated that the abnormal phenotypes of GPX4-OX lines unlikely resulted

from overexpression of functional CsGPX4. Next, we examined whether GPX4-OX insertion

disrupts the gene function at the insertion site and whether the GPX4 protein was expressed as

expected.

Whole Genome Shot Gun Sequencing analysis identified the CsGPX4 T-DNA insertion site at

position chr2:1,646,389 in the *C. sinensis* genome, which was located within the 5′ untranslated

region (UTR) of the gene *SWEET2* (Cs_ont_2g002980.3) (**Fig. S2**). Since mutation of *SWEET2*

has no negative effects in growth or development of plants (Chen et al., 2015), it is unlikely that

the abnormal phenotypes of the CsGPX4-OX line resulted from insertion in the UTR of

SWEET2.

Next, we investigated the CsGPX4 expression in the CsGPX4-OX lines. Immunoblotting with an

anti-HA antibody confirmed expression of the tagged CsGPX4 protein in two independent

GPX4-OX lines (**Fig. 2C**). However, the observed protein band in the transgenic lines

corresponded to a molecular weight of approximately 11 kDa, significantly lower than the

predicted 22.6 kDa including 3HA tag (**Fig. 2C**). PCR amplification using cDNA as template

revealed the expected 505 bp fragment in both EV and GPX4-OX lines (**Fig. 2D**). Sequencing of

the amplified fragment eliminated the possibility of mutation of *CsGPX4* in the GPX4-OX lines

(**Fig. S3**).

To investigate the reasons underlying the truncated CsGPX4 (**Fig. 2C**), we transiently expressed

CsGPX4 in *C. sinensis* and *Nicotiana benthamiana* (**Fig. S4A**). Immunoblot analysis of

transiently expressed CsGPX4 in *C. sinensis* detected the truncated CsGPX4 (**Fig. 4A**) similar to

transgenic expression (**Fig. 2E**), whereas a full length CsGPX4 of approximately 22.6 kDa

including the 3HA tag was observed in *N. benthamiana* (**Fig. 4A; S5**). This result confirmed the

integrity of CsGPX4 overexpression construct and suggested overexpression of CsGPX4 could

undergo a citrus specific truncation, but not in *N. benthamiana*.

To further confirm the truncation of CsGPX4 in the CsGPX4-OX transgenic plants, Matrix-

Assisted Laser Desorption/Ionization - Time-Of-Flight Mass Spectrometry (MALDI-TOF MS)

was performed. MALDI-TOF analysis of CsGPX4 purified through immunoprecipitation from

the CsGPX4-overexpressing *C. sinensis* line identified two major HA-tagged protein isoforms

with the m/z of 10,148.260 and 10,654.116, respectively (**Fig. 4B**), neither corresponding to the

predicted full-length CsGPX4-3HA (22.6 kDa). Based on the MALDI-TOF results, the predicted

cleavage window for the overexpressed CsGPX4-3HA was calculated to be between L115-K117.

Additionally, we transiently expressed CsGPX4-mVenus in *C. sinensis* and *N. benthamiana*

fused with Monomeric Venus *(*mVenus*)* (**Fig. S4B**) to determine if there are any differences in

the subcellular localization (**Fig. 4C**). Confocal microscopy analysis of CsGPX4-mVenus

showed a diffuse cytoplasmic and nuclear distribution in *N. benthamiana* leaf cells (**Fig. 4C;**

S6), which is consistent with the subcellular localization of AtGPX8 (Gaber et al., 2012). In
contrast, CsGPX4 in *C. sinensis* displayed a strong membrane-association. It is probable that
truncation affects the subcellular localization of CsGPX4.

CsGPX4 overexpression disrupts intracellular redox homeostasis and reprograms the
proteome

Because the CsGPX4-OX lines showed oxidative stress phenotypes, we quantified ROS
accumulation (**Fig. 5A-B**). H2DCFDA fluorescence revealed a significant elevation of
intracellular ROS in CsGPX4-OX leaves relative to EV controls ($p < 0.05$) (**Fig. 5A**). In contrast,
the extracellular H_2O_2 measured via the potassium iodide (KI) assay showed no significant
difference between OX-GPX4 leaves and EV lines (**Fig. 5B**).

To assess the broader physiological impact of CsGPX4-OX, we performed a global proteomic
profiling of leaves of CsGPX4-OX lines comparison to EV transformed plants (**Fig. 5C**). We
identified 150 downregulated proteins and 91 upregulated proteins that were significantly
regulated in CsGPX4-OX compared with EV. There were 16 GO biological processes that were
downregulated and two GO biological processes that were upregulated. Downregulated
PANTHER pathways were enriched for developmental pathways, including root and shoot
system development, broader multicellular organism and anatomical structure developmental
processes (**Fig. S7**). Upregulated PANTHER pathways were enriched for metabolic and
biosynthetic pathways (**Fig. S7**).

The significant results were associated with growth, cytoskeletal organization, hormone
signaling and redox detoxification including N-myristoyltransferase 1 (NMT1;
Cs_ont_1g019730.1), glutathione S-transferase U7 (GSTU7; Cs_ont_9g010810.1), microtubule-
associated protein 65 (MAP65; Cs_ont_0419670.1), transport inhibitor response 1 (TIR1;
Cs_ont_2g00600.1), UDP-glycosyltransferase 74E2 (UGT74E2; Cs_ont8g025420.1), and
sucrose synthase 4 (SUS4; Cs_ont_9g001770.2). The upregulated proteins included SNF1-
Related Protein Kinase 2.4 (SnRK2.4; Cs_ont_1g021000.1), and beta-D-xylosidase 2 (BXL2;
Cs_ont_9g001770.2).

Discussion

In this study, we performed systematic characterization of *C. sinensis* GPX family, comprising

CsGPX1–CsGPX4. Previous studies demonstrated that *Arabidopsis* contains eight GPX genes

(Rodriguez Milla et al. 2003), and tobacco has 14 *GPX* genes (Peng et al. 2024). GPXs play

important roles in detoxification of ROS in different compartments. For instance, GPXs were

reported to function in chloroplasts, cytosol, endoplasmic reticulum (ER), mitochondria, nucleus

and membrane (Attacha et al., 2017; Ferro et al. 2003, Rodriguez Milla et al. 2003, Jean-Benoit

et al. 2004; Chang et al. 2009; Gaber et al., 2012). Despite the small amount of GPXs in citrus,

citrus GPXs were predicted or observed to function in most of the known GPX locations

including ER, mitochondrion, plastid, cytosol and nucleus. Nevertheless, it is intriguing why

citrus contains so few GPX genes compared to other plant species.

Multiple studies showed that overexpression of *GPX8* increases plant tolerance against stresses

such as oxidative stress (Gaber 2012; Peng et al. 2024). Unexpectedly, overexpression of

CsGPX4, a homologue of *AtGPX8*, leads to phenotypes commonly associated with oxidative

stress, including growth inhibition, chlorosis, elevated ROS levels, and widespread cellular

disruption (Savchenko and Tikhonov, 2021). It is unlikely that the abnormal phenotypes are due

to overexpression of functional CsGPX4, interfere with the detoxification enzyme activity of

CsGPX4 or disruption of the gene function of insertion site. Overexpression of *GPX8* in

Arabidopsis and tobacco as well as mutation of *GPX8* has no negative effect on plant shoot

growth and development and does not cause oxidative stress phenotypes (Gaber 2012; Peng et al.

2024). In *Salvia miltiorrhiza*, overexpression of *GPX5* enhanced drought and oxidative stress

tolerance without causing negative effects on plant growth and development (Zhang et al., 2018).

Likewise, overexpression of *TaGPX3.2A* and *TaGPX3.4A* in wheat under salt and drought stress

enhanced ROS scavenging capacity (Jiang et al., 2023). WGS analysis showed that T-DNA

insertion occurs in the UTR of *SWEET2* gene. Prior studies indicate that *SWEET2* knockout

mutants in *A. thaliana* do not exhibit significant phenotypic alterations under normal growth

conditions (Chen et al., 2015). Instead, the oxidative stress phenotypes likely result from that

overexpression of *CsGPX4* causes truncation as evidenced by Western blotting and MALDI-

TOF analyses. Overexpression of truncated proteins in plants frequently results in dominant-

negative effects, where the truncated protein disrupts the function of the full-length proteins

including itself or interacting proteins. For instance, overexpression of truncated ERECTA

protein in *Arabidopsis* disrupts signalling of the whole ERECTA gene family and leads to

reduced vegetative growth (Villagarcia et al., 2012). GPX proteins are known to interact with

other proteins and have signaling functions (Delaunay et al., 2002; Miao et al., 2006). It is

probable that the overexpressed truncated CsGPX4 can still interact with other proteins, thus

interfering with their normal functions. Such disruptive effects were not observed in the *AtGPX8*

mutant, suggesting other protein(s) can complement its function. In addition, we could not

exclude other possibilities since the truncated CsGPX4 seems to locate to the membrane instead

of cytosol and nucleus of the full size CsGPX4 and its homologue AtGPX8 (Gaber et al., 2012).

Proteomic profiling revealed downregulation of key proteins involved in detoxification and

cellular structure, including NMT1, GSTU7, MAP65, TIR1, UGT74E2, and SUS4, and

upregulation of SnRK2.4 and BXL2, linking CsGPX4 overexpression misregulation to impaired

antioxidant defense, cytoskeletal destabilization, and altered hormone and cell wall signaling.

Specifically, NMT1 is essential for post-translational lipid modification; its knockout causes

severe growth arrest post-germination (Pierre et al., 2007). GSTU7 is involved in detoxifying

ROS and enhancing stress resilience (Ugalde et al., 2021). MAP65 facilitates microtubule

bundling and axial cell expansion (Lucas et al., 2011). TIR1 is a core auxin receptor; loss of

function impairs hormone-mediated growth (Ruegger et al., 1998). UGT74E2 modulates

oxidative stress response and water stress tolerance (Tognetti et al., 2010). Lastly, SUS4 is a key

to sucrose metabolism and cell wall biosynthesis in citrus (Wang et al., 2018). On the other hand,

upregulated proteins in CsGPX4-OX include several stress-response regulators. For instance,

SnRK2.4 is known to mediate ROS signaling, antioxidant gene expression (e.g., catalase,

ascorbate), and salt stress tolerance (Szymańska et al., 2019; Fujii & Zhu, 2009). These changes

in CsGPX4-OX plants indicate a disruption in core metabolic pathways as observed in oxidative

stress response (Savchenko and Tikhonov 2021), likely contributing to the observed growth

inhibition, microtubule instability, and reduced redox buffering in CsGPX4-OX plants.

In conclusion, our study identifies CsGPX4 as a critical regulator of redox balance in citrus

based on the analysis of the overexpression of truncated CsGPX4 and the consequences (**Fig. 6**).

How overexpression of *CsGPX4* leads to its truncation in citrus but not in tobacco remains to be

explored. The selective truncation of CsGPX4 underscores a unique proteolytic or regulatory

strategies related to redox signaling and ROS homeostasis.

Materials and Methods

Computational analysis of GPX orthologs

GPX homologs were identified from the *Arabidopsis* TAIR database (https://www.arabidopsis.org/) and searched against the *Citrus sinensis* genome via BLAST using the Citrus Genome Database (https://www.citrusgenomedb.org/). Full-length GPX sequences from Arabidopsis, Citrus, and an outgroup were aligned using MAFFT (Katoh et al., 2019) and trimmed with Gblocks on NGPhylogeny.fr (Talavera & Castresana, 2007; Castresana, 2000). Maximum likelihood trees were constructed in MEGA 12 (Kumar et al., 2024) with 1,000 bootstrap replicates. Jalview (Troshin et al., 2011) was used for alignment visualization.

GPX proteins were domain searched using NCBI Conserved Domain Database (CDD) (Marchler-Bauer et al., 2015). Identified GPX proteins were chromosome-mapped using MG2C (Chao et al., 2021). Multiple sequence alignments were performed with MUSCLE (Edgar, 2004) and visualized in Jalview. Protein structure predictions for *C. sinensis* GPXs and AtGPX8 were performed using AlphaFold 3 (Jumper et al., 2021; Varadi et al., 2022), with confidence visualized via pLDDT scores.

Vector construction

The coding sequence of *CsGPX4* was amplified from *C. sinensis* cDNA using primers incorporating XbaI and KpnI restriction sites (**Supplementary Table 2**). The PCR product was gel-purified using the Monarch DNA Gel Extraction Kit and cloned into the binary vector RCsVMV-erGFP-pCAMBIA-1380N-35S-BXKES (**Fig. 2A**), previously digested with XbaI and KpnI (New England Biolabs), using the In-Fusion cloning system (Takara Bio). The resulting construct included a C-terminal 3HA tag and GFP under control of the CaMV 35S promoter.

The constructs pCAMBIA1300S-35S-L279 (**Fig. S4A**) for transient expression of *CsGPX4* and *AtGPX8* (Wang et al., 2017) and pCAMBIA1380-35S-mVenus (**Fig. S4B**) for subcellular localization of *CsGPX4* were similarly assembled, but SacI was used for digestion (**Supplementary Table 2**). Constructs were verified by Sanger sequencing.

Plant transformation

A. tumefaciens strain EHA105 carrying binary constructs was cultured to OD600 = 0.4–0.5 in MS liquid medium (pH 5.4). *C. sinensis* seeds were surface-sterilized with 50% sodium hypochlorite, plated on Murashige and Skoog (MS) medium, and kept in darkness for 6–8

weeks. Epicotyl segments were excised, immersed in *Agrobacterium*-containing medium for 20

min, dried on sterile filter paper, and transferred to co-cultivation medium containing 6-

Benzylaminopurine (BA), Indole-3-acetic acid (IAA), 2,4-Dichlorophenoxyacetic acid (2,4-D),

and acetosyringone. After three days in darkness, explants were transferred to selection medium

with kanamycin and timentin and maintained in darkness for four weeks. Emerging GFP-positive

shoots were micrografted onto Carrizo rootstock and transferred to MS medium supplemented

with IAA and IBA. Upon establishment, plants were moved to soil and acclimatized in growth

chambers.

Agroinfiltration assay

C. sinensis and *N. benthamiana* leaves were infiltrated using the strain *A. tumefaciens* EHA105

strain. *A. tumefaciens* carrying the plasmid was suspended in 10 mM $MgCl_2$, 10 mM 2-N-

morpholino ethanesulfonic acid potassium salt (MES-K), pH 5.6 with 100 µM of acetosyringone.

OD600 of 0.6 was obtained and mixed with *A. tumefaciens* carry vector pCAMBIA1300S-P19 in

a 1:1 ratio. Roughly six-week-old *N. benthamiana* was inoculated and kept in Plant Growth

Chambers from Geneva Scientific (#E-36HO Williams Bay, Wisconsin, United States). The

plants were grown at 23°C under high humidity in 16 h light/8 h dark and leaves were collected

two days later (Acanda et al., 2021). *C. sinensis* leaves were inoculated in a climate controlled

cool greenhouse with dim lighting.

Confocal localization analysis

316 Leaves were imaged for EGFP fluorescence using Confocal Laser Scanning Microscopy TCS-

317 SP5 from Leica (Mannheim, Germany, Europe). Samples were excited with 488 nm and detected

318 using photomultiplier tubes (PMT) tuned to 493-540 nm for EGFP.

Western blot analysis

C. sinensis and *N. benthamiana* leaves were collected and placed in liquid nitrogen and

TissueLyser II (Cat. #85300) from Qiagen (Hilden, Germany, Europe) was used to homogenize

the plant tissue. Total protein was extracted using CelLyticTM P Cell Lysis Reagent (#C2360)

and 1x Protease Inhibitor Cocktail (#P9599) from Sigma (Kawaski, Japan). Lysates were

centrifuged at 11,000 g for 15 minutes at 4°C. Supernatant was taken out and process was

repeated two more times. Laemmli Sample Buffer (#1610747) from Bio-Rad (Hercules, CA,

United States) with 10% of Beta-mercaptoethanol (βME) was mixed with lysates in 1:3 and

boiled at 99°C for 10 minutes. The denatured proteins of CsGPX4, AtGPX8 and EV were

separated using a 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis gel (SDS-

PAGE). Immunoblots were washed in Tris Buffered Saline with Tween 20 (TBST) with 5%

Bovine serum albumin (BSA). Antibody used in this study was Anti-HA High Affinity

(#11867423001) from Roche (Basal, Switzerland, Europe). To visualize the immunoblots,

SuperSignalTM West Pico PLUS Chemiluminescent Substrate (#34580) from Thermo Fisher

was used. Loading was confirmed by Ponceau S staining (#P7170) from Sigma-Aldrich (St.

Louis, Missouri, United States) of the Rubisco large subunit (RbcL), which served as a total

protein loading control.

TEM analysis

Small leaf sections were collected and fixed in cold 3% Gluteraldehyde in 0.1M Sorensen's

Phosphate buffer (SPBS), pH 6.8 for 24 hours. Tissues were then rinsed three times for 30

minutes in 0.1M Sorensen's Phosphate Buffer. Tissues were Osmicated in 1% osmium tetroxide

for four hours room temperature. Tissues were then rinsed 3 times for one hour each in 0.1M

SPBS. Tissues were dehydrated in acetone of concentrations 25%, 50%, 90% 100%, 100%,

100% for 20 minutes. Tissues were infiltrated in a 1:1 mixture of acetone wrapped in plastic

overnight at RT. Infiltrated tissues were made in a 1:3 mixture of acetone and infiltrated again

for six hours. Tissues were transferred to embedded molds, oriented and sat in RT until settled.

The polymerize plastic was heated at 60°C for 12 hours.

The embedded tissue was placed on Epoxy resin blocks that were provided from University of

Florida Interdisciplinary Center Electron Microscopy (ICBR-EM) Core for ultramicrotomy and

TEM examination. Tissues were made into semi-thick resin sections and collected onto super

frost plus glass slides from Fisher Scientific (#22-037-246) and stained with toluidine blue.

Ultrathin resin sections of 100 nm were collected on carbon coated Formvar 2x1 mm nickle slot

grids-TB from Electron Microscopy Sciences (EMS) (Hatfield, PA, United States). Ultrathin

resin sections were treated with 1% meta periodate for 15 minutes and washed with H_2O. Tissues

were air dried and then post-stained with 2% aqueous uranyl acetate and citrate from EMS.

Sections were then examined with a Field Electron and Ion (FEI) Teecnai G2 Spirit Twin TEM

(Hillsboro, OR, United States) and Digital Micrograph Software Gatan (Pleasanton, CA, United

States).

H_2O_2 quantification

Extracellular H_2O_2 concentrations were measured based on Kumar et al., (2011). Samples were

adjusted to fit on 96-well plate. Infiltrated *C. sinensis* tissues were weighed and grinded and 1:4

of 0.1% (w/v) of trichloroacetic acid (TCA) and 5 µL of samples or standard was mixed with

28.25 µL of 1.0 phosphate buffer at a pH of 7.0 and 16.25 µL of 1.0 potassium iodide (KI)

solution. Plates were incubated in room temperature in the dark for five minutes. Absorbance of

290 nm was measured for the oxidized product. H_2O_2 concentrations were then calculated using

the standard curve with H_2O_2 concentration and fresh weight measured in µmol/g.

For H2DCFDA Fluorescence Intensity, H2DCFDA stained was used from *C. sinensis* infiltrated

with the following protocol with minor adjustments (Kaur et al., 2016) and manufactures

instructions from Invitrogen (Eugene, OR, United States). Leaf size of 25 mm^2 were collected

and cleaned using distilled H_2O and incubated in 10 µM H2DCFDA from Thermo Fisher

(#D399) and were vacuum infiltrated for five minutes. Samples were washed using distilled

H_2O. Fresh PBS, pH 7.2 was added and incubated in the dark for 30 minutes. Fluorescence was

measured using BMG LABTECH VANTAstar (Ortenberg, Germany) at Ex/Em= 485/535 nm in

end point mode and analysis was performed and RFU was calculated (Aimonen et al., 2022).

mRNA expression analysis by RT-qPCR

Total RNA was extracted using RNeasy Plant Mini Kit (Cat. #74904) following manufacturer's

instructions. QScript cDNA Supermix (#95048-025) from Quantabio (Beverly, MA, United

States) was used to synthesize cDNA. cDNA was diluted 10 times for RT-qPCR and reactions

were carried out using 1 µL of cDNA, 1 µL of specific primers (**Supplementary Table. 2**), 7 µL

of H_2O, and 10 uL of Thermo-Fisher Scientific Fast SYBR Green Master Mix (Cat. #4385617).

RT-qPCR was conducted using 95°C for 20s followed by 40 cycles at 95°C for 1s, and 60°C for

20s. Relative gene expression was calculated for fold change using the $2^{-\Delta\Delta C_T}$ method (Livak &

Schmittgen, 2001). *CsFBOX* (encoding a F-Box family) was used as the endogenous

housekeeping control (Mafra et al, 2012)

Whole-genome sequencing and T-DNA insertion site identification

To identify the genomic insertion site of the CsGPX4 overexpression construct, whole-genome

resequencing was performed using genomic DNA extracted from a transgenic plant line.

Genomic DNA was extracted using the Promega Genomic DNA Kit (#A1120, Madison, WI,

USA) and paired-end libraries were prepared and sequenced on the Illumina platform to generate

150 bp paired-end reads. Raw sequencing reads were assessed for quality using FastQC, and

adapters were removed using Trimmomatic. High-quality reads were aligned to the *C. sinensis*

genome (Zhong et al., 2024) concatenated with the full CsGPX4 T-DNA plasmid sequence using

BWA-MEM. SAM files were converted to BAM, sorted, and indexed using SAMtools.

To determine the T-DNA insertion site, read pairs mapping partially to the T-DNA plasmid and

partially to the *C. sinensis* genome were identified. Integrative Genomics Viewer (IGV) was

used to visualize junction reads, soft-clipping events, and mate-pair connections (Thorvaldsdottir

et al., 2013). BEDTools was used to intersect candidate insertion coordinates with the *C. sinensis*

genome annotation in GFF3 format to assess gene disruption.

Shotgun proteomics

Infiltrated *C. sinensis* leaves were collected and placed in liquid nitrogen and using the

TissueLyser II to homogenize. Total protein was extracted using 50 mM Tris-HCL, pH 8.0,

0.01% n-dodecyl- β-maltoside (DDM), and 1x of Plant Proteaserrest Protease Inhibitor Cocktail

(#786) from G-BioSciences (St. Louis, MO, United States). Lysis preparation consists of

centrifuging sample at 4°C for 15 minutes and removing supernatant. This process was repeated

3 times. Laemmli sample buffer (#1610747) was mixed with lysates in 1:3 and boiled at 99°C for

10 minutes. An 12% SDS-PAGE gel was made with 50 μg of lysis protein loaded with three

biological replicates of CsGPX4 and EV and was run very briefly. Each lane was excised for in-

gel trypsin digestion. Trypsin (#V5280) from Promega (Madison, WI, United States) was used

for digestion. The peptides were desalted using Pierce C18 Spin Tips & Columns (#87784) from

Thermo Fisher following manufacturer's manual and samples were analyzed using a Thermo

Orbitrap Exploris 240 combined Vanquish Neo UHPLC with a 120-minute LC gradient (Jiang et

al, 2025). Mass spectrometry data were acquired in data-dependent acquisition (DDA) mode.

Full MS scans (MS1) were collected over an m/z range of 375-1475 at a resolution of 120,000.

MS/MS fragmentation was performed at a resolution of 15,000, targeting the most abundant

precursor ions within a 3-second duty cycle. The normalized AGC target is set to 300% for MS1,

and a standard MS2 AGC target was applied using a normalized HCD collision energy of 30%.

Dynamic exclusion was enabled with a 45-second exclusion duration. PatternLab V was used for

data analysis with extracted ion chromatograms for protein expression profiles (Santos et a.,

2022). Protein expression changes were evaluated using extracted-ion chromatograms

(XIC). Briefly, two-sample Student's t-test was used for differential analysis with p cutoff being

0.05. To control false discovery rate, Banjamini-Hochberg FDR estimator was included in

the TFold module. F-stringency, L-stringency and q-value were set to 0.1, 0, and 0.05 as

default settings. Minimal absolute fold change for proteins with significance was 1.7.

MALDI-TOF MS assay

Total protein was processed as stated in Shotgun Proteomics section. Pierce[TM] Anti-HA Agarose

(#26181) from Thermo Fisher was used for immunoprecipitation. Lysates were incubated

rotating with Anti-HA Agarose slurry at 4°C for four hours. Reaction was washed using 50 mM

Tris-HCL, pH 8.0. Proteins were eluted using 0.1M Glycine, pH 2.0 and 1:10 1M

Monoammonium Phosphate (ABC). C18 Spin Tips & Columns (#87784) from Thermo Fisher

was applied for desalting following manufacturer's instructions. Protein at 50 µL was

concentrated at 10 uM.

For Matrix-assisted laser desorption/ionization time-of flight (MALDI-TOF), 1 µL aliquot of

purified protein was mixed 1:1 with saturated sinapinic acid (SA) matrix prepared in 70% CAN

and 0.1% TFA. A 1 µL aliquot of this mixture was spotted onto a stainless steel MALDI target

plate and air dried. Spectra were acquired on a Bruker autoflex maX MALDI-TOF (Billerica,

Massachusetts, United States) mass spectrometer operating in linear positive ion mode over an

m/z range of 5,000-22,000. Spectra were acquired using FlexControl and analyzed with

PolyTools 2.0. Signal intensities were reported in arbitrary units (a.u.) scales $x10^4$.

Acknowledgements

We thank Dr. Karen Kelley at University of Florida Interdisciplinary Center for Biotechnology

Research for TEM expertise and Dr. Kari Basso for running our sample into the MALDI-TOF.

Additionally, we would like to thank Dr. Zhonglin Mou for providing us the vector for transient

expression. We would like to thank our funding sources of the Emergency Citrus Disease

Research and Extension Program grants 2022-70029-38471 and 2021-67013-34588 and the U.S. Department of Agriculture's National Institute of Food and Agriculture grant 2023-70029-41280. MALDI-TOF funding was provided from NIH grant award numbers S10 OD021758-01A1 and S10 OD030250-01A1.

Author Contributions

S.B. and N.W. conceived the work and drafted manuscript. S.B. also designed and preformed experiments. X.W. ran the CsGPX4 and EV samples in LC-MS for global proteomics.

Completing interests

All authors declare no conflict of interest and competing financial interest.

Data Availability

Whole genome resequencing data for CsGPX4 were deposited to NCBI under project ID (PRJNA1419271). Global proteomic data of CsGPX4 and EV are deposited in MassIVE repository (MSV000100757) and ProteomeXchange Consortium (PXD074185) (Vizcaino et al., 2015).

References

1. Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A. J., Bambrick, J., Bodenstein, S. W., Evans, D. A., Hung, C. C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A., Arvaniti, E., Beattie, C., … Jumper, J. M. (2024). Accurate structure prediction of biomolecular interactions with AlphaFold 3. Nature, 630(8016), 493–500. https://doi.org/10.1038/s41586-024-07487-w

2. Aimonen, K., Imani, M., Hartikainen, M., Suhonen, S., Vanhala, E., Moreno, C., Rojas, O. J., Norppa, H., & Catalán, J. (2022). Surface functionalization and size modulate the formation of reactive oxygen species and genotoxic effects of cellulose nanofibrils. Particle and fibre toxicology, 19(1), 19. https://doi.org/10.1186/s12989-022-00460-3

3. Bailey-Serres, J., Parker, J. E., Ainsworth, E. A., Oldroyd, G. E., & Schroeder, J. I. (2019). Genetic strategies for improving crop yields. Nature, 575(7781), 109–118.

4. Bela, K., Horváth, E., Gallé, Á., Szabados, L., Tari, I., & Csiszár, J. (2015). Plant glutathione peroxidases: emerging role of the antioxidant enzymes in plant development and stress responses. Journal of Plant Physiology, 176, 192–201.

5. Bela, K., Riyazuddin, R., & Csiszár, J. (2022). Plant glutathione peroxidases: non-heme peroxidases with large functional flexibility as a core component of ROS-processing mechanisms and signalling. Antioxidants (Basel, Switzerland), 11(8), 1624. https://doi.org/10.3390/antiox11081624

6. Bové, J. M. (2006). Huanglongbing: a destructive, newly-emerging, century-old disease of citrus. Journal of Plant Pathology, 7–37.

7. Cao, S., Hepowit, N., & Maupin-Furlow, J. A. (2015). Ubiquitin-Like Protein SAMP1 and JAMM/MPN+ Metalloprotease HvJAMM1 constitute a system for reversible regulation of metabolic enzyme activity in Archaea. PloS One, 10(5), e0128399. https://doi.org/10.1371/journal.pone.0128399

8. Castresana, J. (2000). Selection of Conserved Blocks from Multiple Alignments for Their
Use in Phylogenetic Analysis. Molecular Biology and Evolution, 17(4), 540–
552. https://doi.org/10.1093/oxfordjournals.molbev.a026334

9. Chae, H. B., Bae, S. B., Paeng, S. K., Wi, S. D., Phan, K. A. T., Kim, M. G., Kim, W. Y.,
Yun, D. J., & Lee, S. Y. (2023). The physiological role of thiol-based redox sensors in
plant defense signaling. The New phytologist, 239(4), 1203–1211.
https://doi.org/10.1111/nph.19018

10. Chao, J., Li, Z., Sun, Y., Aluko, O. O., Wu, X., Wang, Q., & Liu, G. (2021). MG2C: a
user-friendly online tool for drawing genetic maps. Molecular horticulture, 1(1), 16.
https://doi.org/10.1186/s43897-021-00020-x

11. Chen, H. Y., Huh, J. H., Yu, Y. C., Ho, L. H., Chen, L. Q., Tholl, D., Frommer, W. B., &
Guo, W. J. (2015). The Arabidopsis vacuolar sugar transporter SWEET2 limits carbon
sequestration from roots and restricts Pythium infection. The Plant journal : for cell and
molecular biology, 83(6), 1046–1058. https://doi.org/10.1111/tpj.12948

12. Churin, Y., Schilling, S., & Börner, T. (1999). A gene family encoding glutathione
peroxidase homologues in Hordeum vulgare (barley). *FEBS letters*, *459*(1), 33–38.
https://doi.org/10.1016/s0014-5793(99)01208-9

13. Dum, M. T., Teufel, F., Thumuluri, V., Armenteros, J. J. A., Johansen, A. R., Winther,
O., & Nielsen, H. (2024). DeepLoc 2.1: multi-label membrane protein type prediction
using protein language models. Nucleic Acids Research, 52(W1), W215–
W220. https://doi.org/10.1093/nar/gkae237

14. Ebel, R. C. (2017). Huanglongbing: Mechanism of citrus decline and horticulture
management in Florida. Integrated Plant Health Services, Fort Myers, FL, 17, 2019.
www.IPHSLLC.com

15. Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and
high throughput. Nucleic Acids Research, 32(5), 1792–1797.
https://doi.org/10.1093/nar/gkh340

16. Estrella-Maldonado, H., González-Cruz, C., Matilde-Hernández, C., Adame-García, J.,
Santamaría, J. M., Santillán-Mendoza, R., & Flores-de la Rosa, F. R. (2023). Insights into

the molecular basis of Huanglongbing tolerance in Persian lime (Citrus latifolia Tan.) through a transcriptomic approach. International Journal of Molecular Sciences, 24(8), 7497. https://doi.org/10.3390/ijms24087497

17. Faltin, Z., Holland, D., Velcheva, M., Tsapovetsky, M., Roeckel-Drevet, P., Handa, A. K., Abu-Abied, M., Friedman-Einat, M., Eshdat, Y., & Perl, A. (2010). Glutathione peroxidase regulation of reactive oxygen species level is crucial for in vitro plant differentiation. Plant & cell physiology, 51(7), 1151–1162. https://doi.org/10.1093/pcp/pcq082

18. FAO. (2019). The State of Food and Agriculture 2019. Moving forward on food loss and waste reduction. Rome. Licence: CC BY-NC-SA 3.0 IGO.

19. Fedoroff, N. V. (2015). Food in a future of 10 billion. Agriculture & Food Security, 4(1), 11.

20. Fujii, H., & Zhu, J. K. (2009). Arabidopsis mutant deficient in 3 abscisic acid-activated protein kinases reveals critical roles in growth, reproduction, and stress. Proceedings of the National Academy of Sciences of the United States of America, 106(20), 8380–8385. https://doi.org/10.1073/pnas.0903144106

21. Fujita, Y., Nakashima, K., Yoshida, T., Katagiri, T., Kidokoro, S., Kanamori, N., Umezawa, T., Fujita, M., Maruyama, K., Ishiyama, K., Kobayashi, M., Nakasone, S., Yamada, K., Ito, T., Shinozaki, K., & Yamaguchi-Shinozaki, K. (2009). Three SnRK2 protein kinases are the main positive regulators of abscisic acid signaling in response to water stress in Arabidopsis. Plant & Cell Physiology, 50(12), 2123–2132. https://doi.org/10.1093/pcp/pcp147

22. Gaber, A., Ogata, T., Maruta, T., Yoshimura, K., Tamoi, M., & Shigeoka, S. (2012). The involvement of Arabidopsis glutathione peroxidase 8 in the suppression of oxidative damage in the nucleus and cytosol. Plant & Cell Physiology, 53(9), 1596–1606. https://doi.org/10.1093/pcp/pcs100

23. Gottwald, T. R., Da Graça, J. V., & Bassanezi, R. B. (2007). Citrus Huanglongbing: the pathogen and its impact. Plant Health Progress, 8(1). https://doi.org/10.1094/php-2007-0906-01-rv

24. Gutscher, M., Sobotta, M. C., Wabnitz, G. H., Ballikaya, S., Meyer, A. J., Samstag, Y., &
Dick, T. P. (2009). Proximity-based protein thiol oxidation by H2O2-scavenging
peroxidases. The Journal of biological chemistry, 284(46), 31532–31540.
https://doi.org/10.1074/jbc.M109.059246

25. Herbette, S., Roeckel-Drevet, P., & Drevet, J. R. (2007). Seleno-independent glutathione
peroxidases. More than simple antioxidant scavengers. The FEBS journal, 274(9), 2163–
2180. https://doi.org/10.1111/j.1742-4658.2007.05774.

26. Holmgren, A., & Morgan, F. J. (1976). Enzyme reduction of disulfide bonds by
thioredoxin. The reactivity of disulfide bonds in human choriogonadotropin and its
subunits. *European journal of biochemistry*, *70*(2), 377–383.
https://doi.org/10.1111/j.1432-1033.1976.tb11027.x

27. Iqbal, A., Yabuta, Y., Takeda, T., Nakano, Y., & Shigeoka, S. (2006). Hydroperoxide
reduction by thioredoxin-specific glutathione peroxidase isoenzymes of Arabidopsis
thaliana. The FEBS journal, 273(24), 5589–5597. https://doi.org/10.1111/j.1742-
4658.2006.05548.x

28. Islam, T., Manna, M., & Reddy, M. K. (2015). Glutathione peroxidase of Pennisetum
glaucum (PgGPx) is a functional Cd^{2+} dependent peroxiredoxin that enhances tolerance
against salinity and drought stress. PLoS One, 10(11), e0143344.

29. Jiang, B., Su, C., Wang, Y., Xu, X., Li, Y., & Ma, D. (2023). Genome-wide identification
of Glutathione peroxidase (GPX) family genes and silencing TaGPX3.2A reduced
disease resistance in wheat. Plant physiology and biochemistry : PPB, 204, 108139.
https://doi.org/10.1016/j.plaphy.2023.108139

30. Jiang, Z., Islam, K. N., Wolfe, M., O'Connell, M., Williams, D., Florance, A., Vinyard,
D. J., Zhang, X., Brenner, M., Kiss, A. J., Liu, X., & Wang, X. (2025). Enhancing
photosynthesis under salt stress via directed evolution in cyanobacteria. PLANT
PHYSIOLOGY. https://doi.org/10.1093/plphys/kiaf209

31. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,
Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C.,
Kohl, S. A. A., Ballard, A. J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R.,

Adler, J., Back, T., … Hassabis, D. (2021). Highly accurate protein structure prediction

with AlphaFold. Nature, 596(7873), 583–589. https://doi.org/10.1038/s41586-021-

03819-2

32. Kaur, N., Sharma, I., Kirat, K., & Pati, P. K. (2016). Detection of reactive oxygen species

in Oryza sativa L. (Rice). Bio-protocol, 6(24), e2061.

https://doi.org/10.21769/BioProtoc.2061

33. Khan, W. U., Khan, L. U., Khan, N. M., Wenquan, W., & Chen, F. (2024). Evolutionary

insights and expression dynamics of the glutathione peroxidase (GPX) gene family in

water lily (Nymphaea colorata) in response to multiple abiotic stresses. Plant Stress, 14,

100699. https://doi.org/10.1016/j.stress.2024.100699

34. Kumar, N., Ebel, R. C., & Roberts, P. D. (2011). Superoxide dismutase activity in

kumquat leaves infected with *Xanthomonas axonopodis* pv. citri. The Journal of

Horticultural Science and Biotechnology, 86(1), 62-68.

35. Kumar, S., Stecher, G., Suleski, M., Sanderford, M., Sharma, S., & Tamura, K. (2024).

MEGA12: Molecular Evolutionary Genetic Analysis Version 12 for Adaptive and Green

Computing. Molecular biology and evolution, 41(12), msae263.

https://doi.org/10.1093/molbev/msae263

36. Lesk, C., Rowhani, P., & Ramankutty, N. (2016). Influence of extreme weather disasters

on global crop production. Nature, 529(7584), 84–87.

37. Li, J., Li, L., Pang, Z., Kolbasov, V. G., Ehsani, R., Carter, E. W., & Wang, N. (2019).

Developing Citrus Huanglongbing (HLB) management strategies based on the severity of

symptoms in HLB-endemic citrus-producing regions. Phytopathology, 109(4), 582–592.

https://doi.org/10.1094/PHYTO-08-18-0287-R

38. Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using

real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods (San Diego,

Calif.), 25(4), 402–408. https://doi.org/10.1006/meth.2001.1262

39. Lucas, J. R., Courtney, S., Hassfurder, M., Dhingra, S., Bryant, A., & Shaw, S. L. (2011).

Microtubule-associated proteins MAP65-1 and MAP65-2 positively regulate axial cell

growth in etiolated Arabidopsis hypocotyls. The Plant Cell, 23(5), 1889–1903.

https://doi.org/10.1105/tpc.111.084970

40. Ma, W., Pang, Z., Huang, X., Xu, J., Pandey, S. S., Li, J., Achor, D. S., Vasconcelos, F.

N. C., Hendrich, C., Huang, Y., Wang, W., Lee, D., Stanton, D., & Wang, N. (2022).

Citrus Huanglongbing is a pathogen-triggered immune disease that can be mitigated with

antioxidants and gibberellin. Nature communications, 13(1), 529.

https://doi.org/10.1038/s41467-022-28189-9

41. Mafra, V., Kubo, K. S., Alves-Ferreira, M., Ribeiro-Alves, M., Stuart, R. M., Boava, L.

P., Rodrigues, C. M., & Machado, M. A. (2012). Reference Genes for Accurate

Transcript Normalization in Citrus Genotypes under Different Experimental

Conditions. PLoS ONE, 7(2), e31263. https://doi.org/10.1371/journal.pone.0031263

42. Marchler-Bauer, A., Derbyshire, M. K., Gonzales, N. R., Lu, S., Chitsaz, F., Geer, L. Y.,

Geer, R. C., He, J., Gwadz, M., Hurwitz, D. I., Lanczycki, C. J., Lu, F., Marchler, G. H.,

Song, J. S., Thanki, N., Wang, Z., Yamashita, R. A., Zhang, D., Zheng, C., & Bryant, S.

H. (2015). CDD: NCBI's conserved domain database. Nucleic Acids Research,

43(Database issue), D222–D226. https://doi.org/10.1093/nar/gku1221

43. Margis, R., Dunand, C., Teixeira, F. K., & Margis-Pinheiro, M. (2008). Glutathione

peroxidase family - an evolutionary overview. The FEBS journal, 275(15), 3959–3970.

https://doi.org/10.1111/j.1742-4658.2008.06542.x

44. Mittler R. (2017). ROS Are Good. Trends in plant science, 22(1), 11–19.

https://doi.org/10.1016/j.tplants.2016.08.002

45. Mittler, R., Zandalinas, S. I., Fichman, Y., & Van Breusegem, F. (2022). Reactive

oxygen species signalling in plant stress responses. *Nature reviews. Molecular cell*

biology, *23*(10), 663–679. https://doi.org/10.1038/s41580-022-00499-2

46. Morris, K., Linkies, A., Müller, K., Oracz, K., Wang, X., Lynn, J. R., Leubner-Metzger,

G., & Finch-Savage, W. E. (2011). Regulation of seed germination in the close

Arabidopsis relative Lepidium sativum: a global tissue-specific transcript analysis. Plant

Physiology, 155(4), 1851–1870. https://doi.org/10.1104/pp.110.169706

47. Navrot, N., Collin, V., Gualberto, J., Gelhaye, E., Hirasawa, M., Rey, P., Knaff, D. B.,
Issakidis, E., Jacquot, J. P., & Rouhier, N. (2006). Plant glutathione peroxidases are
functional peroxiredoxins distributed in several subcellular compartments and regulated
during biotic and abiotic stresses. Plant physiology, 142(4), 1364–1379.
https://doi.org/10.1104/pp.106.089458

48. Pan, J. L., & Bardwell, J. C. (2006). The origami of thioredoxin-like folds. Protein
Science, 15(10), 2217–2227. https://doi.org/10.1110/ps.062268106

49. Passaia, G., Queval, G., Bai, J., Margis-Pinheiro, M., & Foyer, C. H. (2014). The effects
of redox controls mediated by glutathione peroxidases on root architecture in Arabidopsis
thaliana. Journal of Experimental Botany, 65(5), 1403–1413.
https://doi.org/10.1093/jxb/ert486

50. Peng, X., Ma, T., Song, K., Ji, X., Xiang, L., Chen, N., Zu, R., Xu, W., Zhu, S., & Liu,
W. (2024). Overexpression of NtGPX8a improved cadmium accumulation and tolerance
in tobacco (Nicotiana tabacum L.). Genes, 15(3), 366.
https://doi.org/10.3390/genes15030366

51. Pierre, M., Traverso, J. A., Boisson, B., Domenichini, S., Bouchez, D., Giglione, C., &
Meinnel, T. (2007). N-myristoylation regulates the SnRK1 pathway in Arabidopsis. The
Plant Cell, 19(9), 2804–2821. https://doi.org/10.1105/tpc.107.051870

52. Qi, Y., & Grishin, N. V. (2005). Structural classification of thioredoxin-like fold proteins.
Proteins, 58(2), 376–388. https://doi.org/10.1002/prot.20329

53. Rodriguez Milla MA, Maurer A, Rodriguez Huete A, Gustafson JP. (2003) Glutathione
peroxidase genes in *Arabidopsis* are ubiquitous and regulated by abiotic stresses through
diverse signaling pathways. Plant J. 36(5):602-15. https://doi.org/10.1046/j.1365-
313x.2003.01901.x

54. Ruegger, M., Dewey, E., Gray, W. M., Hobbie, L., Turner, J., & Estelle, M. (1998). The
TIR1 protein of Arabidopsis functions in auxin response and is related to human SKP2
and yeast grr1p. Genes & Development, 12(2), 198–207.
https://doi.org/10.1101/gad.12.2.198

55. Savchenko, T., & Tikhonov, K. (2021). Oxidative Stress-Induced Alteration of Plant

Central Metabolism. *Life (Basel, Switzerland)*, *11*(4), 304.

https://doi.org/10.3390/life11040304

56. Suzuki, N., Koussevitzky, S., Mittler, R., & Miller, G. (2012). ROS and redox signalling

in the response of plants to abiotic stress. Plant, cell & environment, 35(2), 259–270.

https://doi.org/10.1111/j.1365-3040.2011.02336.x

57. Szymańska, K. P., Polkowska-Kowalczyk, L., Lichocka, M., Maszkowska, J., &

Dobrowolska, G. (2019). SNF1-Related Protein Kinases SnRK2.4 and SnRK2.10

Modulate ROS Homeostasis in Plant Response to Salt Stress. International journal of

molecular sciences, 20(1), 143. https://doi.org/10.3390/ijms20010143

58. Talavera, G., & Castresana, J. (2007). Improvement of phylogenies after removing

divergent and ambiguously aligned blocks from protein sequence alignments. Systematic

biology, 56(4), 564–577. https://doi.org/10.1080/10635150701472164

59. Tholey, A., & Becker, A. (2017). Top-down proteomics for the analysis of proteolytic

events - Methods, applications and perspectives. Biochimica et biophysica acta.

Molecular cell research, 1864(11 Pt B), 2191–2199.

https://doi.org/10.1016/j.bbamcr.2017.07.002

60. Thorvaldsdóttir, H., Robinson, J. T., & Mesirov, J. P. (2013). Integrative Genomics

Viewer (IGV): high-performance genomics data visualization and exploration. *Briefings

in bioinformatics*, *14*(2), 178–192. https://doi.org/10.1093/bib/bbs017

61. Tognetti, V. B., Van Aken, O., Morreel, K., Vandenbroucke, K., van de Cotte, B., De

Clercq, I., Chiwocha, S., Fenske, R., Prinsen, E., Boerjan, W., Genty, B., Stubbs, K. A.,

Inzé, D., & Van Breusegem, F. (2010). Perturbation of indole-3-butyric acid homeostasis

by the UDP-glucosyltransferase UGT74E2 modulates Arabidopsis architecture and water

stress tolerance. *The Plant cell*, *22*(8), 2660–2679.

https://doi.org/10.1105/tpc.109.071316

62. Tosatto, S. C., Bosello, V., Fogolari, F., Mauri, P., Roveri, A., Toppo, S., Flohé, L.,

Ursini, F., & Maiorino, M. (2008). The catalytic site of glutathione

peroxidases. Antioxidants & redox signaling, 10(9), 1515–1526.

https://doi.org/10.1089/ars.2008.2055

63. Troshin, P. V., Procter, J. B., & Barton, G. J. (2011). Java bioinformatics analysis web

services for multiple sequence alignment--JABAWS:MSA. Bioinformatics (Oxford,

England), 27(14), 2001–2002. https://doi.org/10.1093/bioinformatics/btr304

64. Ugalde, J. M., Lamig, L., Herrera-Vásquez, A., Fuchs, P., Homagk, M., Kopriva, S.,

Müller-Schüssele, S. J., Holuigue, L., & Meyer, A. J. (2021). A dual role for glutathione

transferase U7 in plant growth and protection from methyl viologen-induced oxidative

stress. Plant Physiology, 187(4), 2451–2468. https://doi.org/10.1093/plphys/kiab444

65. Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan,

D., Stroe, O., Wood, G., Laydon, A., Žídek, A., Green, T., Tunyasuvunakool, K.,

Petersen, S., Jumper, J., Clancy, E., Green, R., Vora, A., Lutfi, M., Figurnov, M., …

Velankar, S. (2022). AlphaFold Protein Structure Database: massively expanding the

structural coverage of protein-sequence space with high-accuracy models. Nucleic Acids

Research, 50(D1), D439–D444. https://doi.org/10.1093/nar/gkab1061

66. Villagarcia, H., Morin, A. C., Shpak, E. D., & Khodakovskaya, M. V. (2012).

Modification of tomato growth by expression of truncated ERECTA protein from

Arabidopsis thaliana. *Journal of experimental botany*, *63*(18), 6493–6504.

https://doi.org/10.1093/jxb/ers305

67. Vizcaíno, J. A., Csordas, A., del-Toro, N., Dianes, J. A., Griss, J., Lavidas, I., Mayer, G.,

Perez-Riverol, Y., Reisinger, F., Ternent, T., Xu, Q. W., Wang, R., & Hermjakob, H.

(2016). 2016 update of the PRIDE database and its related tools. *Nucleic acids*

research, *44*(D1), D447–D456. https://doi.org/10.1093/nar/gkv1145

68. Wang, C., Zhou, M., Zhang, X., Yao, J., Zhang, Y., & Mou, Z. (2017). A lectin receptor

kinase as a potential sensor for extracellular nicotinamide adenine dinucleotide

in *Arabidopsis thaliana*. *eLife*, *6*, e25474. https://doi.org/10.7554/eLife.25474

69. Wang, N., & Trivedi, P. (2013). Citrus huanglongbing: A newly relevant disease presents

unprecedented challenges. Phytopathology, 103(7), 652–665.

70. Wang, W., Cheng, Y., Ruan, M., Ye, Q., Yao, Z., Wang, R., Zhou, G., Liu, D., & Wan, H.

(2023). Comprehensive identification of glutathione peroxidase (GPX) gene family in

response to abiotic stress in pepper (Capsicum annuum L.). Gene, 881, 147625.

https://doi.org/10.1016/j.gene.2023.147625

71. Wang, X., Lin, L., Tang, Y., Xia, H., Zhang, X., Yue, M., Qiu, X., Xu, K., & Wang, Z.

(2018). Transcriptomic insights into citrus segment membrane's cell-wall components

relating to fruit sensory texture. BMC Genomics, 19(1), 280.

https://doi.org/10.1186/s12864-018-4669-y

72. Wu, G. A., Terol, J., Ibanez, V., López-García, A., Pérez-Román, E., Borredá, C.,

Domingo, C., Tadeo, F. R., Carbonell-Caballero, J., Alonso, R., Curk, F.,

Du, D., Ollitrault, P., Roose, M. L., Dopazo, J., Gmitter, F. G., Rokhsar, D. S., & Talon, M.

(2018). Genomics of the origin and evolution of Citrus. Nature, 554(7692), 311–316.

https://doi.org/10.1038/nature25447

73. Xiao, C., Zhang, H., Xie, F., Pan, Z. Y., Qiu, W. M., Tong, Z., Wang, Z. Q., He, X. J.,

Xu, Y. H., & Sun, Z. H. (2022). Evolution, gene expression, and protein–protein

interaction analyses identify candidate CBL-CIPK signalling networks implicated in

stress responses to cold and bacterial infection in citrus. BMC Plant Biology, 22(1), 420.

https://doi.org/10.1186/s12870-022-03809-0

74. You, J., & Chan, Z. (2015). ROS regulation during abiotic stress responses in crop plants.

Frontiers in Plant Science, 6, 1092.

75. Zhai, C. Z., Zhao, L., Yin, L. J., Chen, M., Wang, Q. Y., Li, L. C., Xu, Z. S., & Ma, Y. Z.

(2013). Two wheat glutathione peroxidase genes whose products are located in

chloroplasts improve salt and H_2O_2 tolerances in Arabidopsis. PLoS One, 8(10), e73989.

https://doi.org/10.1371/journal.pone.0073989

76. Zhang, L., Wu, M., Teng, Y., Jia, S., Yu, D., Wei, T., Chen, C., & Song, W. (2019).

Overexpression of the glutathione peroxidase 5 (RcGPX5) gene from Rhodiola crenulata

increases drought tolerance in Salvia miltiorrhiza. Frontiers in Plant Science, 9, 1950.

https://doi.org/10.3389/fpls.2018.01950

77. Zhang, M., Zi, Y., Yang, X., Yang, X., Zhu, L., Cai, H., Yin, T., Liu, X., & Zhang, H.

(2025). Genome-wide analysis of the CsAP2/ERF gene family of sweet orange (Citrus

745   sinensis) and joint analysis of transcriptional metabolism under salt stress. Annals of

746   botany, 135(6), 1165–1180. https://doi.org/10.1093/aob/mcaf006

747  78. Zhang, X., Li, X., Zhao, R., Zhou, Y., & Jiao, Y. (2020). Evolutionary strategies drive a

748   balance of the interacting gene products for the CBL and CIPK gene families. The New

749   Phytologist, 226(5), 1506–1516. https://doi.org/10.1111/nph.16445

750  79. Zhao, X., Qiao, L., & Wu, A. M. (2017). Effective extraction of Arabidopsis adherent

751   seed mucilage by ultrasonic treatment. Scientific Reports, 7(1), 40672.

752  80. Zhong, C., Yang, X., Niu, J., Zhou, X., Zhou, J., Pan, G., Sun, Z., Chen, J., Cao, K., &

753   Luan, M. (2024). Transcriptome analysis of *Citrus Aurantium* L. to study synephrine

754   biosynthesis during developmental stages. *PeerJ*, *12*, e17965.

755   https://doi.org/10.7717/peerj.17965

757 **Figure legends**

758 **Figure 1. Characterization of the CsGPX proteins in *C. sinensis*.** (**A**) Neighbor-Joining

759 phylogenetic tree of CsGPX and AtGPX proteins, constructed using MEGA-X with 1,000

760 bootstrap replicates. The bacterial protein SeBtuE was used as the outgroup. Cs: *C. sinensis*. At:

761 *Arabidopsis thaliana*. (**B**) Chromosomal positions of *CsGPX1-CsGPX4* in *C. sinensis*. Genomic

762 positions were obtained from the *C. sinensis* reference genome and visualized using custom

763 annotation scripts. (**C**) Pairwise amino acid identity analysis of CsGPX proteins. (**D**) Multiple

764 sequence alignment of CsGPX proteins generated with Clustal omega. The catalytic residues are

765 represented by black arrows. The three conserved motifs are highlighted by red boxes. The three

766 conserved Cys are marked by triangles. (**E**) Conserved domain analysis of CsGPX proteins.

767 TRX: Thioredoxin domain. Bar diagrams illustrate domain length and positions in each protein.

768 (**F**) Predicted tertiary structures of CsGPX proteins generated using AlphaFold3. All proteins

769 exhibit a well-resolved TRX fold in dark blue while N-terminal regions show intrinsically

770 disordered or flexible conformations (low confidence in orange).

771 **Figure 2. Verification of GPX4 overexpression and evidence of post-translational**

772 **truncation in transgenic *C. sinensis* lines. (A)** Schematic representation of plasmid construct

773 used for *CsGPX4* overexpression. LB and RB, the left and right borders of the T-DNA region;

CsVMV, the cassava vein mosaic virus promoter; GFP, green fluorescent protein; 35T, the

cauliflower mosaic virus 35S terminator; NosP and NosT, the nopaline synthase gene promoter

and its terminator; NptII, the coding sequence of neomycin phosphotransferase II. HA:

Hemagglutinin tag. (**B**) Phenotypic comparison of empty vector (EV) and *CsGPX4*

overexpression (OX-CsGPX4) *C. sinensis* plants. The picture was taken 1 year after planting in

soil. (**C**) Immunoblot detection of HA-tagged CsGPX4 using an anti-HA antibody. Rubisco was

used as a loading control. (**D**) Reverse transcription PCR confirmation of *CsGPX4* transgene

expression in the OX-CsGPX4 lines. cDNA synthesized from total RNA was amplified using

CsGPX4-specific primers. Amplicons of identical size were observed in both OX and EV plant,

confirming correct transcript expression. PCR products were resolved by agarose gel

electrophoresis.

Figure 3. Transmission electron microscopy (TEM) analysis of *C. sinensis* leaves

overexpressing *CsGPX4*. (**A**) Overall cell organization. CsGPX4-OX plant cells exhibit

irregular boundaries and reduced cytoplasmic density compared to the empty vector (EV)

transformed lines. (**B**) Vacuolar structure. CsGPX4-OX cells show pronounced tonoplast (green

arrow) and membrane (blue arrow) deformation compared to EV. (**C**) Cytoplasmic organization.

CsGPX4-OX cells display extensive vesiculation (arrow) and disorganization of intracellular

contents (dashed arrow) compared to EV. (**D**) Membrane integrity. OX lines show fragmented

membranes, loss of structural continuity, and accumulation of electron-dense material (arrow),

with more severe defects observed in OX-GPX4-2. Scale bars: (**A**) = 2 µm; (**B**) = 1 µm; (**C**) = 2

µm; (**D**) = 0.5 µm.

Figure 4. Host specific processing and subcellular localization of CsGPX4 in *C. sinensis*

and *N. benthamiana*. (**A**) Immunoblot analysis of CsGPX4-3HA transiently expressed in *C.*

sinensis and *N. benthamiana.* A truncated 11 kDa protein was detected in *C. sinensis* and a full-

length protein of 22.5 kDa was detected in *N. benthamiana*. Rubisco was used as a loading

control. (**B**) MALDI-TOF analysis reveals truncated CsGPX4 proteoform in CsGPX4-OX lines.

MALDI-TOF mass spectrum of immunopurified CsGPX4 from citrus leaf tissues expressing

HA-tagged CsGPX4 ranging (m/z) 6000-22000. Two prominent peaks were observed at m/z

10,548.260 and 10,654.116. Both masses are substantially lower than the predicted full-length

protein (22.5 kDa). The 10,548.260 m/z peak dominates in intensity, suggesting this truncated

form is the most stable isoform. Intensity values are presented in arbitrary units (a.u.) scales at

x104. (**C**) Subcellular localization of CsGPX4 in *C. sinensis* and *N. benthamiana*. Confocal

microscopy of leaf epidermis expressing CsGPX4-EGFP-mVenus. Fluorescence displays a

strong membrane-associated pattern in *C. sinensis*. Fluorescence displays GPX4 localization in

cytoplasm and nucleus in *N. benthamiana*. Scale bars: *C. sinensis* = 25 µm; *N. benthamiana* = 50

µm.

Figure 5. Quantification of ROS levels and proteomic analysis of the CsGPX4-OX lines and

wild type. (**A**) Intracellular ROS levels measured using the fluorescent probe H2DCFDA. Three

biological replicates were used, $p < 0.05$, student's t test. (**B**) Quantification of extracellular H_2O_2

using the KI assays. Three biological replicates were used, $p < 0.05$, student's t test. (**C**) Volcano

plot showing differently expressed proteins in CsGPX4-OX leaves relative to EV controls.

Significantly downregulated proteins (red) labeled include NMT1, GSTU7, MAP65, TIR1,

UGT74E2, and SUS4. Significantly upregulated proteins (teal) labeled include SnRK2.4 and

BXL2. Non-significant proteins (gray). Three biological replicates were used.

Figure 6. Proposed model of CsGPX4 metabolic shift results of some involved proteins and

CsGPX4 truncation affects before and after. Top: Schematic summary illustrating the

contrasting roles of CsGPX4 in its functional (left) and truncated (right) states. Functional state

(green): Intact CsGPX4 supports reactive oxygen species (ROS) detoxification and redox

balance, sustaining cytoskeletal organization (MAP6, NMT1), hormonal signaling (UGT42E,

SnRK2.4), and balanced carbohydrate metabolism (SUSA, BXL2). These processes maintain

healthy cytoplasmic structure, intact organelles, and thickened cell walls. Truncated state (red):

Loss of full-length CsGPX4 compromises peroxidase activity, leading to ROS accumulation and

downregulation of redox-related and cytoskeletal proteins (GSTU7, MAP6, NMT1, SUSA).

Stress-associated proteins (SnRK2.4, BXL2) become upregulated, promoting vacuolization,

organelle swelling, and cell wall degradation.

Figure 1. Characterization of the CsGPX proteins in *C. sinensis*. (**A**) Neighbor-Joining phylogenetic tree of CsGPX and AtGPX proteins, constructed using MEGA-X with 1,000 bootstrap replicates. The bacterial protein SeBtuE was used as the outgroup. Cs: *C. sinensis*. At: *Arabidopsis thaliana*. (**B**) Chromosomal positions of *CsGPX1-CsGPX4* in *C. sinensis*. Genomic positions were obtained from the *C. sinensis* reference genome and visualized using custom annotation scripts. (**C**) Pairwise amino acid identity analysis of CsGPX proteins. (**D**) Multiple sequence alignment of CsGPX proteins generated with Clustal omega. The catalytic residues are represented by black arrows. The three conserved motifs are highlighted by red boxes. The three conserved Cys are marked by triangles. (**E**) Conserved domain analysis of CsGPX proteins. TRX: Thioredoxin domain. Bar diagrams illustrate domain length and positions in each protein. (**F**) Predicted tertiary structures of CsGPX proteins generated using AlphaFold3. All proteins exhibit a well-resolved TRX fold in dark blue while N-terminal regions show intrinsically disordered or flexible conformations (low confidence in orange).

Figure 2. Verification of GPX4 overexpression and evidence of post-translational truncation in transgenic *C. sinensis* lines. (A) Schematic representation of plasmid construct used for *CsGPX4* overexpression. LB and RB, the left and right borders of the T-DNA region; CsVMV, the cassava vein mosaic virus promoter; GFP, green fluorescent protein; 35T, the cauliflower mosaic virus 35S terminator; NosP and NosT, the nopaline synthase gene promoter and its terminator; NptII, the coding sequence of neomycin phosphotransferase II. HA: Hemagglutinin tag. **(B)** Phenotypic comparison of empty vector (EV) and *CsGPX4* overexpression (OX-CsGPX4) *C. sinensis* plants. The picture was taken 1 year after planting in soil. **(C)** Immunoblot detection of HA-tagged CsGPX4 using an anti-HA antibody. Rubisco was used as a loading control. **(D)** Reverse transcription PCR confirmation of *CsGPX4* transgene expression in the OX-CsGPX4 lines. cDNA synthesized from total RNA was amplified using CsGPX4-specific primers. Amplicons of identical size were observed in both OX and EV plant, confirming correct transcript expression. PCR products were resolved by agarose gel electrophoresis.

Figure 3. Transmission electron microscopy (TEM) analysis of *C. sinensis* leaves overexpressing *CsGPX4*. (**A**) Overall cell organization. CsGPX4-OX plant cells exhibit irregular boundaries and reduced cytoplasmic density compared to the empty vector (EV) transformed lines. (**B**) Vacuolar structure. CsGPX4-OX cells show pronounced tonoplast (green arrow) and membrane (blue arrow) deformation compared to EV. (**C**) Cytoplasmic organization. CsGPX4-OX cells display extensive vesiculation (arrow) and disorganization of intracellular contents (dashed arrow) compared to EV. (**D**) Membrane integrity. OX lines show fragmented membranes, loss of structural continuity, and accumulation of electron-dense material (arrow), with more severe defects observed in OX-GPX4-2. Scale bars: (**A**) = 2 µm; (**B**) = 1 µm; (**C**) = 2 µm; (**D**) = 0.5 µm.

Figure 4. **Host specific processing and subcellular localization of CsGPX4 in *C. sinensis* and *N. benthamiana*.** (**A**) Immunoblot analysis of CsGPX4-3HA transiently expressed in *C. sinensis* and *N. benthamiana*. A truncated 11 kDa protein was detected in *C. sinensis* and a full-length protein of 22.5 kDa was detected in *N. benthamiana*. Rubisco was used as a loading control. (**B**) MALDI-TOF analysis reveals truncated CsGPX4 proteoform in CsGPX4-OX lines. MALDI-TOF mass spectrum of immunopurified CsGPX4 from citrus leaf tissues expressing HA-tagged CsGPX4 ranging (m/z) 6000-22000. Two prominent peaks were observed at m/z 10,548.260 and 10,654.116. Both masses are substantially lower than the predicted full-length protein (22.5 kDa). The 10,548.260 m/z peak dominates in intensity, suggesting this truncated form is the most stable isoform. Intensity values are presented in arbitrary units (a.u.) scales at x104. (**C**) Subcellular localization of CsGPX4 in *C. sinensis* and *N. benthamiana*. Confocal microscopy of leaf epidermis expressing CsGPX4-EGFP-mVenus. Fluorescence displays a strong membrane-associated pattern in *C. sinensis*. Fluorescence displays GPX4 localization in cytoplasm and nucleus in *N. benthamiana*. Scale bars: *C. sinensis* = 25 µm; *N. benthamiana* = 50 µm.

Figure 5. Quantification of ROS levels and proteomic analysis of the CsGPX4-OX lines and wild type. (**A**) Intracellular ROS levels measured using the fluorescent probe H2DCFDA. Three biological replicates were used, $p < 0.05$, student's t test. (**B**) Quantification of extracellular H_2O_2 using the KI assays. Three biological replicates were used, $p < 0.05$, student's t test. (**C**) Volcano plot showing differently expressed proteins in CsGPX4-OX leaves relative to EV controls. Significantly downregulated proteins (red) labeled include NMT1, GSTU7, MAP65, TIR1, UGT74E2, and SUS4. Significantly upregulated proteins (teal) labeled include SnRK2.4 and BXL2. Non-significant proteins (gray). Three biological replicates were used.

Figure 6. Proposed model of CsGPX4 metabolic shift results of some involved proteins and CsGPX4 truncation affects before and after. Top: Schematic summary illustrating the contrasting roles of CsGPX4 in its functional (left) and truncated (right) states. Functional state (green): Intact CsGPX4 supports reactive oxygen species (ROS) detoxification and redox balance, sustaining cytoskeletal organization (MAP6, NMT1), hormonal signaling (UGT42E, SnRK2.4), and balanced carbohydrate metabolism (SUSA, BXL2). These processes maintain healthy cytoplasmic structure, intact organelles, and thickened cell walls. Truncated state (red): Loss of full-length CsGPX4 compromises peroxidase activity, leading to ROS accumulation and downregulation of redox-related and cytoskeletal proteins (GSTU7, MAP6, NMT1, SUSA). Stress-associated proteins (SnRK2.4, BXL2) become upregulated, promoting vacuolization, organelle swelling, and cell wall degradation.