Title: Conserved cellular architecture and developmental mechanisms of the zebrafish meninges

Authors:

Ashley L. Arancio[1], Kathryn Wilhem[1], Hung-Jhen Chen[1], Brandon M. Hernandez[1], Percy J.
Raggi[1], D'Juan T. Farmer[1,2]

[1]Department of Molecular, Cell and Developmental Biology, University of California, Los Angeles

[2]Howard Hughes Medical Institute

Abstract

The meninges are a multilayered connective tissue that supports and protects the brain and skull,
yet their developmental origins and signaling functions remain poorly understood. Here, we
establish zebrafish as a tractable model for defining meningeal development and function across
larval and adult stages. Using a restricted *foxc1b*:Gal4 reporter, we resolve the spatiotemporal
emergence of meningeal fibroblasts and demonstrate a conserved, dosage-sensitive requirement
for Foxc1 activity during meningeal formation. A targeted pharmacological screen identifies Wnt
signaling as essential for timely establishment of the primary meninx. To define adult meningeal
diversity, we integrate single-cell multiome profiling with spatial validation and identify multiple
transcriptionally distinct populations organized into layered compartments, including pial,
arachnoid, dural, and periosteal dura fibroblasts. Finally, inducible larval ablation of meningeal
cells reveals limited regenerative capacity following widespread loss and leads to persistent
defects in calvarial osteogenesis and brain architecture, including reduced osteoblast
differentiation at bone fronts and disrupted tissue organization at sites lacking meningeal recovery.
Together, these findings define key features of zebrafish meninges and provide a framework for
dissecting meningeal development, regeneration, and meninges-dependent signaling in vivo.

Keywords
zebrafish, meninges, intratissue communication, craniofacial skeleton,

Summary Statement

This study uses zebrafish to show how the tissues surrounding the brain develop early and guide
proper formation of both the brain and skull.

Introduction

The meninges are a multilayered connective tissue that encase the central nervous system and, in the head, lie between the brain and the skull. They are comprised of transcriptionally distinct fibroblast populations, with resident immune cells and an extensive vasculature that support tissue nourishment and waste removal (Dasgupta and Jeong, 2019; Decimo et al., 2012; O'Rahilly and Muller, 1986). Meningeal fibroblasts are broadly organized into three layers: the pia mater, which lies adjacent to the brain and is separated from neural tissue by a basement membrane; the arachnoid mater, which contributes to cerebrospinal fluid regulation; and the dura mater, which anchors the meninges to the overlying skull.

Recent single-cell sequencing studies have begun to resolve the cellular composition of embryonic and adult mouse and human meninges (DeSisto et al., 2020; Ebert et al., 2025; Kearns et al., 2023; Pietila et al., 2023). These analyses reveal substantial fibroblast heterogeneity by embryonic day 14 (E14) and extensive diversity within both the leptomeninges (pia and arachnoid) and the dura mater. Within the dura, transcriptionally distinct inner and outer (periosteal) layers have been identified. These studies highlight an unexpected degree of cellular complexity within vertebrate meninges.

Beyond their structural role, the meninges play essential functions during development by regulating neighboring tissues such as the brain and the skull (Como et al., 2023; Dasgupta and Jeong, 2019). Meningeal-derived retinoic acid signaling modulates cortical neurogenesis (Como et al., 2025; Haushalter et al., 2017; Matrongolo et al., 2023; Mishra et al., 2016; Siegenthaler et al., 2009). Signaling cues including TGFβ family ligands (Greenwald et al., 2000; Slater et al., 2009) from meninges have also been shown to influence cranial suture maintenance and calvarial growth (Cooper et al., 2012; Greenwald et al., 2000; Kwan et al., 2008; Opperman et al., 1993; Slater et al., 2008). Despite these roles, the mechanisms governing meningeal development remain incompletely understood. The forkhead box transcription factor FOXC1 is required for meningeal formation in mice and humans, as hypomorphic or null alleles disrupt meningeal development (Aldinger et al., 2009; Siegenthaler et al., 2009; Zarbalis et al., 2007). Similarly, ZIC family transcription factors are required for meningeal development, as Zic1/3 double mutants exhibit basement membrane defects and impaired fibroblast proliferation (Inoue et al., 2008). In both contexts, genetic perturbations broadly affect all meningeal layers, and only recently have

studies begun to resolve how distinct meningeal cell types are specified during development (Derk

et al., 2023).

Although most functional studies of the meninges have focused on rodent models, ultrastructural

analyses reveal stratified meningeal layers across vertebrate and invertebrate species,

suggesting deep evolutionary conservation (Kappers, 1925; Nakao, 1979). Recent work has

shown that zebrafish possess stratified meninges containing vasculature and immune cell

populations comparable to those of mammals (Galanternik et al., 2025). These findings highlight

zebrafish as a promising system for studying meningeal development, yet the relationships

between zebrafish and mammalian meningeal layers, the transcriptional diversity of zebrafish

meningeal fibroblasts, and the conservation of meningeal signaling functions remain unresolved.

Here, we define the developmental timing and molecular identities of meningeal fibroblasts in

zebrafish. Using *foxc1a* and *foxc1b* mutant alleles, we demonstrate a conserved, dosage-

sensitive requirement for FOXC1 during meningeal formation, with region-specific sensitivity to

gene dosage. Pharmacological perturbation identifies Wnt signaling as essential for timely

establishment of the embryonic meninx. Single-cell genomics resolves multiple transcriptionally

distinct meningeal layers with clear parallels to murine populations. Finally, larval ablation of

meninges reveals limited regenerative capacity following large-scale loss and uncovers persistent

defects in both brain and skull development. Together, these findings establish zebrafish as a

powerful model for dissecting the developmental and regulatory programs governing meningeal

biology.

Results

A *foxc1b*:Gal4 reporter marks developing meningeal populations in zebrafish

To assess whether the zebrafish paralogs for Foxc1 (*foxc1a* and *foxc1b*) are active in meninges,

we examined zebrafish carrying the *foxc1b*:Gal4;UAS:mCherryNTR transgene (Whitesell et al.,

2019). In contrast to the previously described *foxc1b*:GFP reporter, which is broadly expressed

throughout the face and craniofacial skeleton, mCherry fluorescence was restricted to regions

surrounding the brain (**Fig. 1A**). Orthogonal sections revealed that mCherry and GFP signals

were co-detected within a single cellular layer positioned adjacent to the brain surface. Co-

imaging with the epithelial marker *cdh1*:mlanYFP confirmed that mCherry[+] cells reside beneath

the epithelium (**Fig. 1B**). To assess mesenchymal identity, *foxc1b*:Gal4;UAS:mCherryNTR fish

were crossed with *fli1a*:GFP, a broad marker of craniofacial mesenchyme. mCherry$^+$ cells

encasing the brain co-expressed *fli1a*:GFP, whereas *fli1a*:GFP-labeled vasculature lacked

detectable mCherry signal (**Fig. 1C**). A developmental time course revealed no detectable

mCherry signal at 24 hpf. By 36–48 hpf, mCherry$^+$/*fli1a*:GFP$^+$ cells emerged within the rostral

meninges, while *fli1a*:GFP$^+$ cells appeared in caudal regions prior to mCherry detection. By 72

hpf, double-positive cells were broadly distributed around the brain (**Fig. 1D**). Time-lapse imaging

confirmed early mCherry activation in the rostral meninges, primarily over the midbrain, with

fli1a:GFP expression preceding mCherry in all regions (Movie 1). Together, these data define the

spatiotemporal emergence of *foxc1b*$^+$ meningeal cells and establish *foxc1b*:Gal4 as a restricted

reporter of meningeal development.

To further define the identity of mCherry$^+$ meningeal cells, we examined markers from the

Daniocell single-cell transcriptomic dataset (Sur et al., 2023). Subsetting and reclustering of

meningeal populations resolved two clusters of meningeal precursors, one cluster of meningeal

fibroblasts, and three leptomeningeal clusters (**Fig. 1E**). Meningeal precursor clusters expressed

canonical mesenchymal markers, including *fli1a* and *lum*, as well as the transcription factor *meox1*

(**Fig. 1F**). Consistent with previous reports (Galanternik et al., 2025), all leptomeningeal clusters

expressed ependymin (*epd*) and gamma-glutamylcyclotransferase b (*ggctb*), whereas the

meningeal fibroblast cluster was distinguished by expression of solute carrier family 4 member 7

(*slc4a7*) and transferrin receptor protein 1a (*tfr1a*) (**Fig. 1F**). To validate these transcriptomic

assignments in vivo, we performed RNAscope analysis, which confirmed expression of both

slc4a7 and *ggctb* transcripts within mCherry$^+$ meningeal cells by 5 dpf (**Fig. 1G**). These markers

were frequently co-expressed and did not resolve into discrete layers, consistent with incomplete

lineage commitment at larval stages.

Foxc1 paralogs exhibit conserved requirements during meningeal development

To test whether requirements for meningeal development are conserved across vertebrates, we

examined the function of the transcription factors *foxc1a* and *foxc1b*, both of which were detected

in meningeal precursor, meningeal fibroblast, and leptomeningeal populations (**Fig. 2A**).

RNAscope analysis revealed co-expression of *foxc1a* and *foxc1b* in the rostral meninges by 36

hpf, including within mCherry$^+$ meningeal cells (**Fig. 2B**). Transcripts for both paralogs extended

beyond the domain of mCherry expression, suggesting that reporter accumulation follows

foxc1a/b transcriptional activation. At this stage, mCherry$^+$ meningeal cells were detected along

both the dorsal and ventral surfaces of the brain, whereas neither reporter nor *foxc1a/b* transcripts

were observed in the dorsal hindbrain or the most caudal ventral meninges. By 5 dpf, mCherry

and both foxc1 paralogs were broadly co-expressed throughout the meningeal layer encasing the

brain (**Fig. 2B**).

To determine the functional requirements of *foxc1a* and *foxc1b*, we generated an allelic series

using established mutant alleles (Xu et al., 2018) in combination with the

foxc1b:Gal4;UAS:mCherryNTR reporter. Loss of *foxc1b* alone did not alter meningeal reporter

distribution, as *foxc1a*$^{+/+}$; *foxc1b*$^{-/-}$ and *foxc1a*$^{+/-}$; *foxc1b*$^{-/-}$ larvae appeared phenotypically normal

(**Fig. 2C**). In contrast, loss of *foxc1a* resulted in reduced mCherry signal in the hindbrain

meninges, and removal of *foxc1b* in the *foxc1a* mutant background further exacerbated this

phenotype, consistent with a dosage-sensitive requirement for Foxc1 activity during meningeal

development (**Fig. 2C**).

Quantification of dorsal mCherry signal revealed regional differences in sensitivity to Foxc1

dosage (**Fig. 2D**). Midbrain meninges were comparatively resilient, with significant reductions in

mCherry signal detected only in *foxc1a*$^{-/-}$; *foxc1b*$^{+/-}$ and *foxc1a*$^{-/-}$; *foxc1b*$^{-/-}$ mutants. In contrast,

hindbrain meningeal signal was significantly reduced in *foxc1a*$^{-/-}$; *foxc1b*$^{+/+}$ larvae and further

diminished in *foxc1a*$^{-/-}$; *foxc1b*$^{+/-}$ mutants, with little additional loss upon complete *foxc1b*

removal. Spatiotemporal analysis demonstrated that reduced mCherry signal in *foxc1a*$^{-/-}$;

foxc1b$^{-/-}$ was evident by 36 hpf and persisted through 48 hpf and 72 hpf (**Fig. 2E**). Time-lapse

imaging confirmed that mCherry expression initiates rostrally over the midbrain but fails to expand

across the brain surface in mutant embryos (**Movie 2**).

To further resolve regional defects, mCherry$^+$ meningeal cells were quantified across five dorsal–

ventral domains in sectioned larvae. While dorsal midbrain meninges were largely preserved in

foxc1a$^{-/-}$; *foxc1b*$^{-/-}$ mutants, all other regions exhibited significant depletion (**Fig. 2F,G**). To

assess meningeal integrity using additional markers, RNAscope was performed for the broad

meningeal marker *igfbp2a* and the leptomeningeal marker *ggctb*. In contrast to mCherry loss,

igfbp2a expression was maintained around the mutant brain, including in regions lacking

detectable mCherry signal (**Fig. 2H**). By comparison, *ggctb* expression was restricted to small

residual meningeal domains near the midbrain and was largely absent from the forebrain and

hindbrain, where mCherry loss was most pronounced (**Fig. 2H**). Together, these data

demonstrate a conserved, dosage-sensitive requirement for Foxc1 genes during vertebrate
meningeal development and reveal region-specific dependence on Foxc1 activity across the
developing brain.

Pharmacological inhibition identifies pathways required for meningeal formation

To identify developmental pathways required for meningeal establishment, we performed a
targeted pharmacological screen using *foxc1b*:Gal4;UAS:mCherryNTR embryos. Inhibitors were
applied beginning at 24 hpf, prior to mCherry$^+$ meningeal emergence, and embryos were analyzed
at 72 hpf, when mCherry$^+$ cells normally encase the brain (**Fig. 3A**). Inhibition of Hedgehog
(Cyclopamine), Notch (DAPT), retinoic acid (DEAB), FGF (Infigratinib), or TGFβ (LY2109761)
signaling did not disrupt formation of a continuous meningeal layer (**Fig. 3B**). In contrast, inhibition
of BMP signaling (Dorsomorphin) or Wnt signaling (XAV-939) prevented complete encasement
of mCherry$^+$ meninges, with defects most pronounced in the dorsal hindbrain, the last meningeal
population to form (**Fig. 3B**).

To determine whether early pathway inhibition was sufficient to disrupt meningeal development,
embryos were transiently treated with XAV-939 or Dorsomorphin between 24 and 48 hpf and
allowed to develop until 5–6 dpf (**Fig. 3C**). Treated larvae were comparable in size to controls but
exhibited persistent depletion of mCherry signal within medial regions of the hindbrain meninges
for both treatments (**Fig. 3D**). Analysis of *foxc1b*:Gal4;UAS:mCherryNTR; *fli1a*:GFP larvae
revealed that mesenchymal cells persisted in the hindbrain following XAV-939 treatment, whereas
mCherry reporter activation was selectively impaired. In contrast, Dorsomorphin-treated larvae
lacked both mCherry and *fli1a*:GFP signal in this region, consistent with broader loss of
mesenchymal cells (**Fig. 3D**).

XAV-939 treatment also appeared to impair forebrain meninges (**Fig. 3D**). To further assess this
phenotype, we examined larvae carrying the forebrain meningeal reporter *tbx18*:dsRed2, which
co-expresses with *foxc1b*:Gal4;UAS:nlsGFP (**Fig. 3E**). Transient XAV-939 treatment from 24 to
48 hpf resulted in a marked reduction of dsRed2 signal, particularly at the forebrain–midbrain
boundary, indicating that Wnt signaling is required for proper establishment of forebrain meninges
(**Fig. 3F**). Together, these data identify BMP and Wnt signaling as essential regulators of
meningeal development and underscore the utility of zebrafish for dissecting conserved pathways
governing formation of the vertebrate meninges.

Adult zebrafish meninges are comprised of transcriptionally distinct fibroblasts

During embryonic stages, zebrafish meninges appear as a single cellular (**Fig. 1G**), whereas prior studies suggest that adult zebrafish meninges are multilayered (Galanternik et al., 2025). To define the transcriptional identities of adult meningeal populations, we performed single-nucleus multiomic profiling.

To assess persistence of *foxc1b* reporter expression in adults, brains and skullcaps were dissected from *foxc1b*:Gal4;UAS:mCherryNTR fish. Robust mCherry signal was detected surrounding the brain and along the ventral surface of the skullcap, with enrichment in midbrain and hindbrain regions (**Fig. 4A**). RNA analysis confirmed co-expression of mCherry and *foxc1b* in meninges associated with both brain and skull (**Fig. 4B**). Notably, pan-laminin immunostaining selectively marked mCherry$^+$ cells associated with the brain but not those adjacent to the skull, suggesting molecular heterogeneity within adult meninges (**Fig. 4C**).

To define transcriptional and epigenetic profiles of adult meningeal populations, connective tissues associated with the brain or skull were isolated from *foxc1b*:Gal4;UAS:nlsGFP and *fli1a*:GFP fish, which we previously identified as labeling meningeal populations associated with the calvaria (Farmer et al., 2024), and subjected to multiome analysis (**Fig. 4D**). After quality control, 1,224 cells from the *fli1a*:GFP dataset and 1,188 cells from the *foxc1b*:Gal4;UAS:nlsGFP dataset were recovered. Unsupervised clustering of merged datasets identified 17 distinct cell populations that passed filtering (**Fig. 4E, Fig. S1A, Table S1**). Clusters included mesenchymal populations expressing connective tissue markers such as *pdgfrb* and *col1a1a* (**Fig. S1B**), as well as pericytes, smooth muscle/mural cells, endothelial cells, epithelial cells, immune populations, pigment cells, and neuronal cell types (**Fig. 4E, Fig. S1C**). Relative abundance of non-mesenchymal populations differed modestly between datasets, whereas endothelial, epithelial, and T cell populations were present at comparable levels (**Fig. S1D**).

Within mesenchymal populations, dermal fibroblasts were enriched in the *fli1a*:GFP dataset, while *krt94*$^+$ mesenchymal cells, captured in our prior datasets (Farmer et al., 2024) and validated to be a connective tissue that sits at the borders of skull (**Fig. S1E**), were preferentially enriched in the *foxc1b*:Gal4;UAS:nlsGFP dataset (**Fig. 4F, G**). With the exception of a single cluster (arachnoid fibroblasts), the remaining mesenchymal clusters exhibited comparable contributions

from both datasets (**Fig. 4F, G**). To further resolve the identities of previously uncharacterized
fibroblast populations, we analyzed cluster-enriched genes and assigned preliminary cell-type
identities, which we validate in this study. These populations include arachnoid fibroblasts, dural
fibroblasts, and two transcriptionally distinct periosteal dura fibroblast populations (**Fig. 4E-G**).

To relate zebrafish meningeal populations to mammalian counterparts, we reanalyzed published
mouse leptomeningeal and dura mater single-cell datasets (Pietila et al., 2023) and performed
cross-species comparisons. We identified genes enriched within each zebrafish cluster and
calculated module scores based on zebrafish genes and their mouse paralogs, enabling
systematic comparison across species (**Fig. S2A–D**). The *coch*$^+$ cluster corresponded most
closely to mouse pial fibroblasts and inner arachnoid cells (BFB1/BFB2), whereas the *slc6a2*$^+$
cluster matched a previously undescribed dura population enriched for *Pappa* (**Fig. S2E-F**). The
ccn2a$^+$ periosteal dura fibroblast cluster aligned with mouse outer dura populations expressing
Matn4 and *Nppc* (Farmer et al., 2021) (**Fig. S2G**). In contrast, the *rspo2*$^+$ periosteal dura fibroblast
cluster lacked clear correspondence to known mouse meningeal populations (**Fig. S2H**).

Analysis of chromatin accessibility revealed cluster-specific regulatory programs. Arachnoid
fibroblasts were enriched for ETV, EGR, and GATA motifs, dural fibroblasts for TFAP, STAT, and
nuclear receptor motifs, and both populations for ZIC family motifs, consistent with known roles
in meningeal biology (Inoue et al., 2008) (**Fig. 4H**). Periosteal dura fibroblast clusters shared
enrichment for IRF, NFAT, and NFI motifs, while all meningeal populations exhibited enrichment
for ZNF family motifs, indicating distinct but overlapping regulatory architectures (**Fig. 4H**).

To spatially localize transcriptionally defined populations, RNAscope in situ hybridization was
performed in adult zebrafish. *ggctb*$^+$*igfbp2a*$^+$ cells persisted adjacent to the brain but were largely
absent beneath the skull, despite poor representation in single-cell datasets and co-expression
of *ggctb* and *foxc1b*:Gal4;UAS:GFP (**Fig. 5A; Fig. S3A**). Cells adjacent to the brain also
expressed transcripts encoding the secreted chemokine *cxcl12a* and the retinoic acid–
synthesizing enzyme *aldh1a2* (**Fig. S3B**). Arachnoid cells co-expressing *coch* and *igfbp2a* formed
a spatially discrete population restricted to brain-associated meninges and were positioned
superficial to *ggctb*$^+$ cells (**Fig. 5B,C**). In contrast, dural populations marked by *slc6a2* and
periosteal dura fibroblasts marked by *ccn2a* and *rspo2* localized preferentially to the skull-
associated side of the meninges **(Fig. 5D,E)**. *ccn2a*$^+$ and *rspo2*$^+$ cells were intermingled within a

single bone-adjacent layer, consistent with transcriptionally distinct periosteal dura fibroblast
populations occupying the same anatomical compartment.

Together, these data demonstrate that adult zebrafish meninges comprise a multilayered and
transcriptionally heterogeneous structure consisting of at least four distinct populations: a pial
layer (*ggctb*[+]*igfbp2a*[+]), a discrete arachnoid layer (*coch*[+]), an inner dura mater layer (*slc6a2*[+]),
and a periosteal dura layer composed of intermingled *ccn2a*[+] and *rspo2*[+] fibroblasts (**Fig. 5F**).

Larval meningeal ablation reveals long-term consequences on calvarial development

To assess the functional requirement for the meninges during development, meningeal cells were
ablated using the *foxc1b*:Gal4;UAS:mCherryNTR line. Because ablation at juvenile or adult
stages resulted in lethality within 24 hours, subsequent analyses focused on larval ablation.
Larvae were treated with metronidazole (MTZ) or DMSO to induce apoptosis specifically in
mCherry[+] meningeal cells. MTZ treatment at 5 dpf resulted in widespread loss of mCherry signal,
confirming efficient meningeal ablation (**Fig. S4A,B**). Consistent with loss of meningeal integrity,
expression of the broad meningeal marker *igfbp2a* was markedly reduced around the brain at 0
and 1 days post-treatment (dpt), with discontinuous recovery by 2 dpt, indicating partial
regeneration of the meninges (**Fig. S4C**). MTZ-treated larvae exhibited reduced long-term
survival, with 54% surviving to 18 dpt compared to 87% of DMSO-treated controls, demonstrating
lasting consequences of larval meningeal ablation (**Fig. S4D**).

To examine long-term skeletal outcomes, MTZ-treated fish were raised to adulthood and their
craniofacial skeletons analyzed. Alizarin Red staining revealed widespread calvarial defects, with
the most prevalent phenotype consisting of a persistent opening in the skull (**Fig. 6A, B**). These
openings were frequently associated with misdirected bone growth, with bone folding on itself and
growing away from the opening (**Fig. 6B**). In addition, an ectopic bone occasionally formed
between the frontal and parietal bones (**Fig. 6B**). Defects were reproducibly localized to the
bregma, where cranial sutures converge (**Fig. 6C**).

To determine whether skeletal phenotypes were sensitive to the timing of meningeal ablation,
MTZ treatment was initiated at 3, 4, or 5 dpf. Ablation at 3 dpf, when the brain is already encased
by meninges, resulted in the lowest incidence of skeletal defects (~33%) (**Fig. 6D**). In contrast,
ablation at 4 or 5 dpf produced defects in ~80% of animals (**Fig. 6D**). Later ablation increased

both the frequency and size of calvarial openings and was associated with a higher incidence of
ectopic bone formation (**Fig. 6E,F**). These findings indicate that meningeal disruption during larval
stages causes persistent, stage-dependent defects in calvarial patterning.

To investigate the basis of calvarial openings, osteogenesis was examined during skull
development. Prior to calvarial bone coverage, pigmented dura mater cells normally populate the
juvenile head (Galanternik et al., 2025). In control animals at 8 mm, before calvarial bones encase
the brain (Topczewska et al., 2016), these cells uniformly covered the midbrain and hindbrain
(**Fig. 7A**). In contrast, MTZ-treated fish exhibited pronounced depletion of pigmented dura cells
at the same stage, consistent with meningeal disruption preceding skull formation (**Fig. 7A**).

Repeated live imaging of *RUNX2*:EGFP fish revealed no differences between control and MTZ-
treated animals at 9 mm, when calvarial bones have initiated growth but remain separated (**Fig.
7B**). However, by 14 mm, when bones normally approximate, MTZ-treated fish exhibited
persistent openings at the prospective bregma. *RUNX2*$^+$ bone fronts failed to converge and
instead appeared folded or misdirected, growing over adjacent bone. These defects persisted
through 18 mm, a stage when cranial sutures are normally established, indicating failure of defect
resolution. To determine whether osteoblast differentiation was impaired, the osteoblast:nlsEOS
reporter was used to track de novo osteoblast formation (Farmer et al., 2024) (**Fig. 7C**). In control
animals, newly differentiated osteoblasts accumulated as bone fronts approached and decreased
after bones met (**Fig. 7D**). In contrast, MTZ-treated fish showed a marked reduction in de novo
osteoblast differentiation at open bone fronts, with few newly differentiated osteoblasts detected
near calvarial openings (**Fig. 7D, E**).

Given the reproducible localization of defects at the bregma, we tested whether localized
meningeal ablation was sufficient to induce long-term craniofacial abnormalities. Targeted two-
photon ablation of meningeal cells at the dorsal midbrain–hindbrain boundary resulted in efficient
but transient loss of meningeal fluorescence, with robust regeneration by 30 hours post-ablation
(**Fig. S5A**). Fish subjected to focal ablation and raised to adulthood exhibited no craniofacial
abnormalities or calvarial openings (**Fig. S5B**). These findings indicate that large-scale, rather
than localized, disruption of meningeal tissues is required to compromise calvarial development.

Impaired meninges regeneration leads to impaired calvarial and brain development

To define tissue-level disruptions underlying impaired osteogenesis and calvarial opening

formation, histological and whole-mount analyses were performed on skulls from DMSO- and

MTZ-treated animals. To preserve skull–brain architecture, zebrafish heads were plastic-

embedded, sectioned, and stained with toluidine blue. In control samples, the brain remained

closely apposed to the skull with intact intervening tissues (**Fig. 8A**). In contrast, MTZ-treated

skulls failed to fully encase the brain, which instead adhered to the overlying dermis and exhibited

disorganization of the underlying neural tissue (**Fig. 8A**).

To assess meningeal regeneration at sites of calvarial opening, RNAscope was performed on

foxc1b:Gal4;UAS:mCherryNTR;UAS:GFP animals as mChery expression does not reactivate

after MTZ treatment. In controls, GFP$^+$ meningeal cells lined the ventral skull surface and

igfbp2a$^+$GFP$^+$ cells encased the brain (**Fig. 8B**). In MTZ-treated tissue, brain tissue remained

adherent to the skin at calvarial openings, and neither GFP nor *igfbp2a* signal was detected at

the defect itself, despite persistence of GFP$^+$ cells adjacent to surrounding bone and *igfbp2a*$^+$

GFP$^+$ around brain associated meninges away from the opening (**Fig. 8B**). These data indicate

impaired meningeal regeneration at sites of failed bone formation.

Because some MTZ-treated skulls exhibited ectopic bone formation, meningeal regeneration was

examined at these locations. RNAscope revealed irregular brain–skull adhesion and absent

igfbp2a expression beneath ectopic bone, indicating that bone formation can occur in the absence

of meningeal recovery but results in aberrant calvarial patterning (**Fig. S6**).

Consistent with histological and molecular analyses, imaging of the *wu:fj16a03*:nlsEOS reporter

that labels brain-associated meningeal cells (**Fig. S7**), revealed depletion of EOS$^+$ cells overlying

calvarial openings, with enrichment along the periphery of the defect on the skullcap (**Fig. 8C**).

Single-cell transcriptomic analysis of DMSO- and MTZ-treated skullcaps showed largely

comparable cellular compositions, with the exception of a cluster uniquely enriched in MTZ-

treated samples expressing markers characteristic of arachnoid fibroblasts normally associated

with brain meninges (**Fig. S8A-D**; **Tables S2**).

Transmission electron microscopy further confirmed loss of pigmentation and disorganization of

fibroblast populations at calvarial openings, despite preservation of overlying dermal and epithelial

structures (**Fig. S9**). Disruption of the underlying brain parenchyma was also observed, consistent

with impaired meningeal support of neural tissue organization.

In control animals, bilateral optic tecta were clearly demarcated based on the distribution of

wu:fj16a03:nlsEOS[+] cells, whereas this organization was lost in MTZ-treated brains (**Fig. 8C,**

arrows). Whole-mount staining for acetylated tubulin confirmed disrupted brain architecture in

regions lacking meningeal regeneration, including loss of proper separation between the optic

tecta (**Fig. 8D**). Together, these findings demonstrate that impaired meningeal regeneration leads

to persistent defects in both skull and brain development, underscoring the critical role of

meninges in coordinating the organization and growth of adjacent tissues.

Discussion

The meninges have long been recognized as a critical connective tissue supporting and protecting

the brain and skull (Dasgupta and Jeong, 2019), yet their cellular composition, developmental

origins, and signaling functions remain incompletely understood. Here, we define the

developmental emergence, molecular heterogeneity, and functional importance of zebrafish

meningeal populations across larval and adult stages. By integrating genetic reporters, mutant

analyses, single-cell multiomic profiling, spatial validation, pharmacological perturbation, and

targeted ablation, we uncover conserved roles for the meninges in coordinating calvarial growth

and brain organization.

Developmental emergence and heterogeneity of zebrafish meninges

Using a restricted *foxc1b*:Gal4 reporter, we show that zebrafish meningeal populations emerge in

a spatiotemporally ordered manner, with rostral meninges forming before caudal populations,

paralleling patterns described in mammals (Zarbalis et al., 2007). A notable distinction is the

concurrent emergence of foxc1[+] meninges along both dorsal and ventral surfaces of the zebrafish

brain, in contrast to predominantly basal initiation reported in mammals. During larval stages,

meningeal cells exhibit incomplete lineage commitment, with co-expression of fibroblast and

leptomeningeal markers within a single cellular layer, suggesting a prolonged plastic state. This

may reflect delayed specialization in zebrafish that facilitates rapid growth and remodeling during

early development.

In adults, single-cell profiling combined with spatial validation reveals a multilayered meningeal

architecture composed of at least four transcriptionally and spatially distinct populations. These

findings provide molecular resolution to prior anatomical descriptions (Galanternik et al., 2025)

and demonstrate that, as in mammals, zebrafish meninges comprise specialized layers with
distinct identities. Interestingly, although cells corresponding to each meningeal layer could be
identified based on positional and transcriptional features, certain specialized cell types, such as
arachnoid barrier and dural border cells, lacked clear zebrafish counterparts. This suggests that
zebrafish represent a simplified meningeal system, potentially accompanied by distinct structural
and functional differences compared with mammalian meninges. Nonetheless, the experimental
accessibility of zebrafish provides a powerful and tractable model for dissecting the core
mechanisms that govern meningeal stratification and layer-specific functions.

Conserved regulatory programs governing meningeal development
Functional analyses demonstrate a conserved, dosage-sensitive requirement for Foxc1 during
meningeal development, with region-specific dependencies across the brain. Loss of *foxc1a*
preferentially affects hindbrain meninges, while combined reduction of *foxc1a* and *foxc1b*
exacerbates defects, consistent with mammalian studies linking FOXC1 dosage to meningeal
integrity (Aldinger et al., 2009; Siegenthaler et al., 2009; Zarbalis et al., 2007). Interestingly,
although both paralogs show comparable expression during meningeal development, foxc1a
appears more critical, suggesting limited paralog-specific functions. Persistence of some
meningeal populations in Foxc1-deficient contexts suggests that additional transcriptional
regulators contribute to early meningeal specification.

Pharmacological perturbation identifies BMP and Wnt signaling as essential for meningeal
establishment, particularly for late-forming hindbrain populations. Disruption of Wnt signaling
selectively impaired meningeal differentiation without broadly affecting mesenchymal survival,
suggesting stage- or region-specific roles. These findings are consistent with prior work
implicating Wnt signaling in meningeal expansion (Choe et al., 2014) and highlight conserved
signaling pathways regulating meningeal development. Motif enrichment analyses further
implicate transcription factor families previously associated with meningeal biology, including ZIC
factors, suggesting that layered combinations of signaling inputs and transcriptional programs
generate meningeal diversity. The identification of population-specific enrichment of GATA,
TFAP, and NFAT motifs highlights new potential regulatory pathways for future study.

Meninges as regulators of calvarial osteogenesis
A central finding of this study is that meningeal integrity is required for proper calvarial bone
formation. Larval ablation of meninges results in persistent calvarial openings, misdirected bone

growth, ectopic bone formation, and defects localized to the bregma. These phenotypes are stage-dependent, with later ablation producing more severe defects, suggesting a progressive decline in meningeal regenerative capacity. Live imaging reveals that impaired calvarial development primarily reflects reduced de novo osteoblast differentiation at bone fronts rather than defects in initial bone specification. Notably, focal meningeal ablation is insufficient to induce long-term skeletal defects, indicating that large-scale or sustained disruption of meningeal tissue is required to compromise calvarial patterning. Together with the occurrence of ectopic bone formation in regions lacking canonical meningeal markers, these observations support a model in which meninges provide positional or organizational cues that support bone growth, rather than acting solely as inductive sources of osteogenic signals. Similar roles for meninges in maintaining suture patency have been described in mammals (Cooper et al., 2012; Greenwald et al., 2000; Kwan et al., 2008; Opperman et al., 1993; Slater et al., 2008), and hypomorphic *Foxc1* mutations produce calvarial openings reminiscent of those observed here (Zarbalis et al., 2007). It is possible that promiscuous adhesion of the meninges to adjacent tissues serves to re-establish a closed CSF system necessary for brain function, albeit at the expense of creating a suboptimal local environment for brain and bone.

Meningeal regeneration and tissue coordination

Despite partial regeneration following ablation, regenerated meninges fail to re-establish normal tissue architecture at calvarial openings. Loss of meningeal markers, abnormal brain–dermis adhesion, fibroblast disorganization, and disruption of underlying neural tissue correlate with defects in brain morphology, including impaired separation of the optic tecta. These findings underscore the role of meninges as both structural and signaling intermediates between the brain and skull.

Consistent with mouse studies, loss of meningeal tissue likely disrupts delivery of key secreted factors, such as retinoic acid (Como et al., 2025; Haushalter et al., 2017; Matrongolo et al., 2023; Mishra et al., 2016; Siegenthaler et al., 2009) and CXCL12 (Borrell and Marin, 2006), which regulate neural development and were also detected in zebrafish meninges adjacent to brain tissue. The pronounced brain phenotypes observed here highlight the conservation of meningeal–neural interactions and establish zebrafish as a powerful system for studying tissue coordination during craniofacial development.

Implications and future directions

Together, these findings position the meninges as dynamic, heterogeneous tissues essential for

coordinating brain and skull development across multiple stages in zebrafish. Given the

conservation of meningeal architecture and regulatory programs, insights from this model are

likely to inform our understanding of human disorders influenced by meningeal-derived signals,

including craniofacial disorders like craniosynostosis and calvarial dysplasia.

504 **Figures and Figure Legends**

Figure 1. A *foxc1b* reporter specifically labels larval meninges and reveals early spatial patterns of meningeal formation. (A–C) Confocal images of 5 dpf zebrafish larvae showing detection of the *foxc1b*:Gal4;UAS:mCherryNTR reporter in combination with additional transgenic markers. Lateral (top) and dorsal (bottom) views are shown for each condition, with orthogonal sections provided to illustrate relative cell distribution (n=5 per condition). **(A)** Larvae carrying *foxc1b*:GFP together and *foxc1b*:Gal4;UAS:mCherryNTR show restricted mCherry⁺ expression

surrounding the brain, with GFP and mCherry co-expression enriched in cells closest to the brain

surface (arrowheads). **(B)** Co-imaging of *foxc1b*:Gal4;UAS:mCherryNTR with the epithelial

marker *cdh1*:mlanYFP shows non-overlapping expression. **(C)** Larvae carrying *fli1a*:GFP and

foxc1b:Gal4;UAS:mCherryNTR display co-expression of mCherry and GFP in cells surrounding

the brain. Arrowheads indicate representative double-positive cells. **(D)** Developmental time

course (24–72 hpf) of *fli1a*:GFP and *foxc1b*:Gal4;UAS:mCherryNTR expression. Arrows denote

mCherry$^+$GFP$^+$ cells, and arrowheads indicate GFP-only cells (n=5 per stage). **(E)** UMAP

clustering of subsetted meningeal populations from the Daniocell single-cell RNA-sequencing

dataset (Sur et al., 2023). **(F)** Dot plot showing marker genes enriched in each meningeal cluster

identified in (E). **(G)** In situ hybridization for leptomeningeal marker *ggctb* and meningeal fibroblast

marker *slc4a7*, combined with mCherry immunostaining in *foxc1b*:Gal4;UAS:mCherryNTR larvae.

Insets show higher-magnification views highlighting co-detection (n=3). Scale bars, 100 μM.

Figure 2. *foxc1a* and *foxc1b* exhibit dosage-sensitive requirements for meningeal

formation. (A) Violin plots showing expression of *foxc1a* and *foxc1b* across meningeal clusters.

(B) In situ hybridization for *foxc1a* and *foxc1b*, combined with mCherry immunostaining in

foxc1b:Gal4;UAS:mCherryNTR fish at 36 hpf and 5 dpf (n=3 per stage). **(C)** Dorsal-view confocal

images of *foxc1b*:Gal4;UAS:mCherryNTR larvae across an allelic series of *foxc1a* and *foxc1b*

mutants (n=5 per genotype). Yellow dotted lines outline the larval brain, and white dotted lines

demarcate the forebrain-midbrain and midbrain-hindbrain boundaries. **(D)** Quantification of dorsal

mCherry$^+$ meningeal coverage of the midbrain and hindbrain across genotypes. **(E)**

Developmental time course (36 hpf–3 dpf) of *foxc1b*:Gal4;UAS:mCherryNTR expression in wild-

type and *foxc1a$^{-/-}$*; *foxc1b$^{-/-}$* embryos (n=5 per genotype). Yellow dotted lines outline the larval

brain**. (F)** Representative histological sections of control and mutant larvae stained with Hoechst,

highlighting regions used for quantification of mCherry signal. Color coded dashed lines indicate

assigned zones for mCherry quantitation (n=5 per genotype). **(G)** Quantification of mCherry signal

surrounding the brain from histological sections shown in (F). **(H)** In situ hybridization for *igfbp2a*

or *ggctb*, combined with mCherry immunostaining in *foxc1b*:Gal4;UAS:mCherryNTR wildtype and

foxc1a$^{-/-}$; *foxc1b$^{-/-}$* larvae (n=3 per genotype). Arrowhead marks *igfpb2a$^+$* region devoid of

mCherry signal in mutant. P-values were calculated using two-tailed non-parametric Student's t-

tests. Error bars represent s.e.m. Scale bars, 100 μm.

Figure 3. Drug screening of developmental pathways reveals a requirement for Wnt and

BMP signaling during meningeal formation. (A) Schematic illustrating the short-term drug

treatment strategy. **(B)** Dorsal-view confocal images of *foxc1b*:Gal4;UAS:mCherryNTR embryos

treated with inhibitors of developmental signaling pathways using the long-term drug treatment

paradigm and imaged at 72 hpf. Asterisks indicate reduced mCherry signal following

Dorsomorphin and XAV-939 treatment (n = 10 embryos per condition). **(C)** Schematic illustrating

the short-term drug treatment strategy. **(D)** Dorsal-view confocal images of

foxc1b:Gal4;UAS:mCherryNTR larvae treated with DMSO, XAV-939 or Dorsomorphin using the

short-term drug treatment paradigm and imaged at 5 or 6 dpf (n = 10 larvae per condition).

Asterisk marks rostral regions where mCherry$^+$ cells appear reduced. The dotted box marks a

representative region shown at higher magnification, including co-imaging with *fli1a*:GFP.

Arrowhead indicates region containing *fli1a*:GFP single-positive cells in XAV-939–treated

embryos. **(E)** Dorsal-view confocal images of 5 dpf zebrafish larvae showing

foxc1b:Gal4;UAS:nlsGFP expression in combination with *tbx18*:dsRed2 (n = 5 larvae). Dorsal

(top) and magnified (bottom) views are shown, with orthogonal sections illustrating co-expression

within the rostral meninges. Dotted lines indicate regions used for orthogonal sections. **(F)** Dorsal-

view confocal images of *tbx18*:dsRed2 larvae treated with DMSO or XAV-939 using the long-term

drug treatment paradigm and imaged at 6 dpf. Asterisk shows disruption of dsRed2$^+$ cells at the

forebrain-midbrain interface (n = 5 larvae per condition). Scale bars, 100 μm.

Figure 4. Single-cell genomics reveals multiple fibroblast cell types with zebrafish meninges. (A) Dorsal view of a dissected brain and ventral-view of a dissected skullcap from adult *foxc1b*:Gal4;UAS:mCherryNTR animals, showing mCherry expression associated with cranial meninges (n = 5). **(B)** In situ hybridization for *foxc1b* and mCherry in adult *foxc1b*:Gal4;UAS:mCherryNTR animals shows co-expression of *foxc1b* and mCherry in meningeal tissue located beneath the cranial bones (arrowhead) and surrounding the brain (arrow) (n = 3). **(C)** Immunostaining for laminin and mCherry in adult

foxc1b:Gal4;UAS:mCherryNTR animals shows co-detection in meninges surrounding the brain

(arrowhead) (n = 3). F= frontal bone, P= parietal bone, B= brain. **(D)** Schematic illustrating the

transgenic lines and tissues used for single-cell genomic experiments. **(E)** UMAP clustering of

sorted cells from merged datasets with cluster annotations. Red dashed lines delineate fibroblast

clusters, and predicted fibroblast identities are indicated within the dotted red box. **(F)** Dot plot

showing marker genes enriched in each fibroblast cluster identified in (E). **(G)** Graphical depiction

of the fraction of each fibroblast cluster captured from each dataset, with colors corresponding to

the schematic in (D). **(H)** Graphical depiction of the frequency of enriched transcription factor

family motifs within each representative cluster. Scale bars, 100 μm.

Figure 5. Meningeal fibroblasts are spatially organized into distinct layers between the skull and brain in the adult zebrafish. (A) In situ hybridization for *ggctb* and *igfbp2a* shows co-expression of both genes in cells surrounding the brain (arrows), with *igfbp2a*+ cells (arrowhead) interspersed among double-positive cells. **(B)** In situ hybridization for *coch* and *igfbp2a* shows scattered co-expression of both genes around the brain (arrows), along with *igfbp2a*+ cells (arrowhead) distributed along the brain surface. **(C)** In situ hybridization for *ggctb* and *coch* reveals non-overlapping expression domains, with *coch*+ cells (arrowheads) positioned superficial to *ggctb*+ cells. **(D)** In situ hybridization for *slc6a2* and *ccn2a* shows *ccn2a*high cells (arrowheads) located closest to the bone, while *slc6a2*high cells (arrows) are positioned farther from the bone. Low levels of *slc6a2* expression are also detected around the brain. **(E)** In situ hybridization for *ccn2a* and *rspo2* shows expression of both genes within the same layer adjacent to the bone, with double-positive cells (arrow) distributed along the skull. Signal for both genes is absent around the brain. **(F)** Model illustrating the spatial organization of meningeal subtypes within the adult zebrafish skull. Boxes indicate magnified regions corresponding to locations highlighted in the images. All stains were performed in biological triplicates.

Figure 6. Larval ablation of meninges leads to persistent calvarial defects. (A) Dorsal views of Alizarin-stained skulls from DMSO- and MTZ-treated adult zebrafish reveal openings in the calvaria of treated fish (n = 13 fish per condition). **(B)** Representative images of the bregma region in Alizarin-stained skulls from DMSO- and MTZ-treated fish show a range of skeletal phenotypes in treated animals. Arrows indicate irregular lines around bones, suggestive of misdirected bone growth. Arrowhead indicated ectopic bone. Asterisks mark sites of calvarial openings **(C)** Hand-drawn outlines of suture borders in control and MTZ-treated zebrafish reveal reproducible defects at the bregma. **(D)** Quantification of MTZ-treated zebrafish exhibiting skeletal defects following ablation at different developmental time points. **(E)** Quantification of skeletal phenotype classes

observed at different treatment times. **(F)** Quantification of open calvarial defects expressed as

the ratio of opening area to total calvarial area. P-values were calculated using two-tailed non-

parametric Student's t-tests. Error bars represent s.e.m. Scale bars, 100 μm.

Figure 7. Osteogenesis is disrupted by early ablation of meninges. (A) Transmitted light

images of control- and MTZ-treated zebrafish at 8 mm (n = 5 fish per condition). The dotted line

demarcates the region of pigment depletion. **(B)** Dorsal-view confocal images from serial imaging

of *foxc1b*:Gal4;UAS:mCherryNTR; *RUNX2*:GFP zebrafish treated with DMSO or MTZ between 9

mm and 18 mm (n = 6 fish per condition). Arrows indicate regions of abnormal overlapping bone,

and asterisks mark RUNX2-negative openings in the calvaria. **(C)** Schematic illustrating the

treatment and imaging scheme used to quantify de novo osteoblast rates in the juvenile calvaria.

The red box indicates the region imaged in subsequent panels. **(D)** Serial confocal imaging of

individual DMSO- and MTZ-treated fish carrying the osteoblast:nlsEOS transgene shows reduced

generation of de novo osteoblasts in treated animals. The same fish were photoconverted at

approximately 8 mm and imaged at 10 mm, photoconverted again and imaged at 13 mm,

photoconverted a final time and imaged at 16 mm (n = 5 fish per condition). F= frontal bone, P=

parietal bone **(E)** Quantification of de novo osteoblasts at bone fronts.. P-values were calculated

using two-tailed non-parametric Student's t-tests. Error bars represent s.e.m. Scale bars, 100 μm.

Figure 8. Fish fail to regenerate dorsal meninges, resulting in irregular skeletal and brain morphology. (A) Toluidine blue staining of plastic-embedded heads from DMSO- and MTZ-treated fish shows absence of calvarial bone and abnormal attachment of the brain to the overlying dermis and skin in treated animals (n = 3 fish per condition). Dotted boxes indicate regions shown at higher magnification in the lower panels. Arrowheads mark bone ends in MTZ treated skull. **(B)** In situ hybridization for *igfbp2a* combined with GFP immunostaining in *foxc1b*:Gal4;UAS:mCherryNTR;UAS:GFP heads (n = 4 fish per condition). Numbered dotted boxes indicate regions shown at higher magnification in the lower panels. **(C)** Confocal imaging of dissected skullcaps and brains from DMSO- or MTZ-treated fish carrying the *wu:fj16a03*:nlsEOS reporter (n = 3 fish per condition). Arrowheads mark EOS+ cells adhered to the skullcap. Arrows indicate EOS+ signal between optic tecta. **(D)** Confocal imaging of whole brains stained for acetylated tubulin reveals irregular brain morphology in MTZ-treated fish (n = 3

fish per condition). Dotted lines indicate regions used for orthogonal sections. F= frontal bone, P=
parietal bone. Asterisks mark sites of calvarial openings. Scale bars, 100 μm.

Movie 1. Time-lapse of embryo carrying *fli1a*:GFP and *foxc1b*:Gal4;UAS:mCherryNTR reporters,
beginning at 24hpf and imaged for approximately 72 hours.

Movie 2. Time-lapse of *foxc1a$^{-/-}$*; *foxc1b$^{-/-}$* embryo carrying *fli1a*:GFP and
foxc1b:Gal4;UAS:mCherryNTR reporters, beginning at 24hpf and imaged for approximately 72
hours.

Methods

Zebrafish
All experiments were approved by the Institutional Animal Care and Use Committee at the
University of California, Los Angeles (Protocol #ARC-2022-044). Published lines include
Tg(*fli1a*:eGFP)^{y1} (Lawson and Weinstein, 2002), Tg(Hsa.*RUNX2*-Mmu.Fos:EGFP)^{zf259} (Kague et
al., 2016), *Tg(-5.0kbfoxc1b:EGFP)^{mw44}* (Miesfeld and Link, 2014), *Tg(foxc1b:gal4FF)^{ca72}*
(Whitesell et al., 2019)*, cdh1-mlanYFP^{xt17}* (Cronan and Tobin, 2019)*, Tg(tbx18:DsRed2)^{pd22}*
(Kikuchi et al., 2011)*, Tg(UAS:GFP)(Asakawa et al., 2008), Tg(UAS-NTR:mCherry)^{C264}* (Davison
et al., 2007), *Tg(UAS:nlsGFP)^{el609}* (Barske et al., 2020)*, osteoblast*:nlsEOS (Farmer et al., 2024),
foxc1a^{el543} (Xu et al., 2018)*,* and *foxc1b^{el620}* (Xu et al., 2018). One transgenic line was created
using CRISPR/Cas9-based genomic integration of nlsEOS (Kimura et al., 2014; Thomas and
Raible, 2019): wu:fj16a03 (guide RNAs targeting the upstream of the 5' UTR: 5′-
AATGATGAAATATCTCTGCG-3). One-cell embryos were co-injected with guide RNAs
(200 ng/μL), mbait guide RNA (200 ng/μL), IDT Cas9 protein (150 ng/μL), and a mbait-nlsEOS
plasmid (20 ng/μL). Founders were identified by screening progeny for nlsEOS fluorescence.

Single nuclei isolation
Adult zebrafish skullcaps were dissected away (*foxc1b*:gal4;UAS:nlsGFP, *fli1a*:GFP and DMSO
and MTZ-treated *foxc1b*:Gal4;UAS:mCherryNTR) and fluorescent meninges attached to the brain
were scraped away (*foxc1b*:gal4; UAS:nlsGFP) in Ringer's solution. Tissue was mechanically
minced with a razor and all tissue was transferred to a 1.5 mL tube for enzymatic dissociation
(0.25% trypsin (Life Technologies, 15090-046), 1 mM EDTA, and 400 mg/mL Collagenase D
(Sigma, 11088882001) in PBS). Tissue was incubated on a nutator at 28.5 °C for ~45 min and

the enzymatic reaction was stopped by adding 6X stop solution (6 mM CaCl2 and 30% fetal

bovine serum (FBS) in PBS). Cells were pelleted by centrifugation (300 g) at 4 °C and

resuspended in a final volume of 500 uL of 0.03% BSA in PBS. DAPI was added to resuspended

cells, and cells were fluorescence-activated cell sorted to isolate live cells that were DAPI negative

and GFP or mCherry positive for enrich for fibroblast populations or just DAPI negative cells for

DMSO/MTZ skullcaps. Single nuclei suspension was collected by following 10X Genomics nuclei

isolation protocol with low cell input for Multiome. FACS cells were pelleted for 15 min (300 rcf at

4°C) and incubated with lysis buffer for 90 s on ice. Isolated nuclei were washed by Wash buffer

and Nuclei buffer and checked for nuclear integrity with a fluorescence confocal microscope with

DAPI staining before use in library construction.

Single cell-omic library construction, sequencing, and alignment

Multi-omic libraries (*foxc1b*:gal4;UAS:nlsGFP and *fli1a*:GFP) of snATACseq and snRNAseq from

the same barcoded single nuclei were constructed per manufacturer's instructions (10X

Genomics, Chromium Next GEM Single Cell Multiome ATAC + Gene Expression). Single-Cell

RNA sequencing libraries (DMSO or MTZ-treated *foxc1b*:Gal4;UAS:mCherryNTR) were

constructed by 10X Genomics Chromium Single Cell 3′ Library and Gel Bead Kit v.2. To capture

accessible chromatin and transcripts from the same nuclei in Multiome, accessible chromatin and

polyadenylated mRNA from the same nuclei were pulled down and barcoded with the same

sequences within isolated GEMs to achieve single-nuclei separation. cDNAs were synthesized

through reverse transcription from polyadenylated mRNA. Quality control of libraries was

assessed with the 4200 TapeStation system and Qubit dsDNA HS assay kit. Libraries were

sequenced on Illumina NovaSeq (Multiome) and NextSeq (scRNAseq) platforms. For Multiome

libraries, sequencing reads were aligned to customized genome built with GRCz11.fa,

GRCz11.104.gtf, and JASPAR2022.pfm with addition of GFP gene information and called peak

by Cell Ranger ARC v2.0.0 per manufacturer's instructions to generate peak-by-cell and gene-

by-cell count matrices. ScRNA-seq sequencing reads were aligned to customized genome built

from GRCz11 with addition of mCherry gene information by Cell Ranger v3.0.0 to generate gene-

by-cell count matrices.

Single cell-omics data analysis

scRNA-seq data using the 10X Genomics Chromium Single Cell 3′ Library and Gel Bead Kit v.2

was analyzed as previously described (Farmer et al., 2024). Multi-omics data were analyzed by

Seurat v5 and Signac v1.4 packages following the standard workflow with optimization. Cell

Ranger ARC-output count matrices of individual libraries were used to create Seurat objects by
"CreateSeuratObject" and "CreateChromatinAssay" functions. Quality control was performed by
setting the thresholds of accessible region counts (nCount_ATAC) < 40000 (foxc1b) or < 30000
(fli1a), transcript counts (nCount_RNA) < 6000 (foxc1b) or < 7500 (fli1a), ratio of
mononucleosome to nucleosome-free region (nucleosome_signal) < 3, and enrichment at TSS
(TSS.enrichment) > 0.5. In order to compare accessible chromatin regions (peaks) between
libraries, peaks-called were combined by "reduce" function that merges all intersecting peaks
between libraries and filtered by new peak widths (peakwidths < 10000 & > 20). The reduced-
peak list was used to recalculate peak-by-cell count matrices by "CreateFragmentObject",
"FeatureMatrix", and "CreateChromatinAssay" functions. "Merge" function was used to combine
count matrices of two libraries. Merged ATAC data were processed by "FindTopFeatures"
(min.cutoff = 5) and LSI linear dimension reduction; and merged RNA data were processed by
"SCTransform" and dimensionally reduced by PCA. 2:29 dimensions of LSI and 1:30 dimensions
of PCA were used to cluster cells with both ATAC and RNA information by
"FindMultiModalNeighbors" and "FindClusters" (resolution = 1, graph.name="wsnn", algorithm =
3). UMAP visualization was generated by "RunUMAP" (nn.name = "weighted.nn"). After filtering
for artificial clusters of dying cells , 19 clusters were identified. Differentially enriched genes were
calculated by "FindAllMarkers" (test.use = "wilcox", logfc.threshold = 0.25, return.thresh = 0.01,
min.pct = 0.25). Differentially accessible regions were calculated by "FindAllMarkers" (test.use =
"LR", logfc.threshold = 0.25, return.thresh = 0.01, min.pct = 0.1). To calculate motif enrichments
of each cell by chromVAR, "AddMotifs" was first used to calculate motif enrichments of each
accessible chromatin region (peak) followed by "RunChromVAR" to calculate motif activity in each
cell. "FindAllMarkers" test.use = "wilcox", mean.fxn = rowMeans, logfc.threshold = 0.25,
return.thresh = 0.01, min.pct = 0.1) was performed to identify differentially enriched motifs.
Enriched motifs from the same family were grouped together to calculate the frequency of
enrichment for desired fibroblast clusters.
742

Published dataset analysis
The processed Seurat object and associated annotation list were downloaded directly from the
Daniocell website(Sur et al., 2023) and meninges clusters were subsetted and reclustered to
capture transcriptionally distinct clusters by refining the resolution. For mouse datasets, individual
leptomeningeal and dura mater datasets were downloaded from GEO and reanalyzed
independently using standard Seurat methods. Module scores were calculated using the
AddModuleScore function in Seurat with 10 control features. Scores were derived from 7–8 genes

differentially expressed in zebrafish meningeal clusters with clear mouse homologs. FeaturePlots

visualizing module scores were displayed with a minimum cutoff of the 10th percentile (q10) and

a maximum cutoff of the 90th percentile (q90).

RNAscope in situ hybridization and immunofluorescence

Embryonic and adult fish were measured and fixed individually in 4% PFA overnight at 4°C. Whole

embryos processed in 10% sucrose in PBS for one day and 30% sucrose in PBS for at least one

day before embedding tissue in Optimal Cutting Temperature (OCT) medium. Frozen sections

were cut at 10 µm. Adult heads were decalcified for 2 weeks and processed by paraffin embedding

as previously described (Farmer et al., 2024). RNAscope reagents were purchased from

Advanced Cell Diagnostics, and experiments were performed using the RNAscope Fluorescent

Multiplex V2 Assay according to the manufacturer's protocol for formalin-fixed paraffin-embedded

(adult) or for fixed frozen (embryos) sections. For cryosections, heat antigen retrieval was omitted.

The following probes were used for this study: Channel 1, dr-*gttcb*, dr-*krt94*, dr-*slc6a2,* dr-*foxc1b*;

Channel 2, dr-*coch*, dr-*foxc1a*, *mcherry*, dr-*slc4a7*; Channel 3, dr-*ccn2a*, dr-*igfbp2a*; Channel 4,

dr-*rspo2*. RNAscope experiments were followed by immunofluorescence or performed

independently using the same antigen retrieval reagents as the RNAscope Fluorescent Multiplex

kit, and slides were blocked for 1 h in 10% (adult stains) or 5% (embryonic stains) goat serum,

stained with primary overnight (anti-laminin, 1:250, Millipore #L9393; anti-GFP, 1:1000, Novus

NB100-1614; anti-mCherry, 1:1000, Novus NBP2-25157) and stained with Alexa-Fluor

secondaries from Thermo Scientific (1:250, Thermo) and Hoechst 33342 at room temperature for

1 h. Wholemount staining of adult brains were performed by permeabilizing tissues with 0.25%

Triton for 3 hours at room temperature followed by blocking in anti-acetylated α-tubulin antibody

(1:500, Sigma T6793) overnight, and the secondary for two hours at room temperature with

secondary antibody and Hoechst 33342.

Transmission Electron Microscopy

Whole heads were fixed with 2.5% glutaraldehyde and 4% paraformaldehyde in 0.1M sodium

cacodylate buffer overnight at 4°C, and decalcified for 2 weeks in 0.5 M EDTA. The hole area was

dissected away, including surrounding calvaria tissue and underlying brain. Dissected tissues

were then post-fixed in 1% osmium tetroxide in 0.1M sodium cacodylate or 0.1M PB, and

dehydrated through a graded series of ethanol concentrations. After infiltration with Eponate 12

resin, the samples were embedded in fresh Eponate 12 resin and polymerized at 60°C for 48

hours. Ultrathin sections of 70 nm thickness were prepared and placed on formvar/Carbon coated

copper grids and stained with uranyl acetate and Reynolds' lead citrate. The grids were examined
using a JEOL 100CX transmission electron microscope at 60 kV and images were captured by
an AMT digital camera (Advanced Microscopy Techniques Corporation, model XR611) (Electron
Microscopy Core Facility, UCLA Brain Research Institute).
788

Plastic embedding
Zebrafish heads were dissected and fixed in 70% ethanol without decalcification. After
dehydration, samples were gradually infiltrated with increasing concentrations of methyl
methacrylate resin and embedded in 2.5% methyl methacrylate. Longitudinal sections (5 μm thick)
were obtained using a Polycut microtome fitted with a tungsten carbide knife and stained with
toluidine blue for histological analysis. Sections were subsequently stained with Toluidine Blue
staining and imaged using an ECHO Revolution microscope.
796

2-photon ablation
5 dpf larvae were mounted in low-melting-point agarose and imaged on an inverted LSM 880
microscope equipped with a two-photon laser. Using a 20× air objective, a region at the midbrain–
hindbrain boundary was manually delineated using the ROI tool to encompass approximately 20
nuclei. Targeted ablation was performed at 450 mW with a scan speed of 400 Hz for a total
duration of 9 s (three cycles of 3 s each). Ablation efficacy was assessed by imaging in the
transmitted photomultiplier tube (T-PMT) channel to confirm loss of pigmented cells. Z-stacks
were acquired immediately after ablation and again 6 h post-treatment to verify sustained signal
loss. Following imaging, larvae were individually housed and re-imaged at 1 and 2 days post-
treatment (dpt) before being raised to adulthood.
807

808 **Adult skeletal preparations**
809 Skeletal preparations were performed as previously described (Teng et al., 2018), and zebrafish
810 were imaged on a Zeiss Stemi with Zeiss Labscope software in 100% glycerol.
811

Imaging
All imaging was performed on a Zeiss LSM880 or a Zeiss LSM980 microscope using ZEN
software. For repeated live imaging experiments, fish were anesthetized in Tricaine, mounted,
and imaged using a 10X objective. For whole calvaria nlsEOS conversion experiments, live fish
were transferred to a 6- or 12-well dish and exposed to a handheld UV light (UV Flashlight Black

Light, 3-in-1 Magnetic Flashlight Rechargeable, AdamStar) for 5–30 min. Converted fish were

individually housed and re-imaged for up to 2 months following initial conversion.

Metronidazole treatment

821   Zebrafish carrying *foxc1b*:gal4;UAS:mCherryNTR were outcrossed to wildtype AB zebrafish and

822   sorted by 5 dpf. mCherry+ fish were split into two groups to receive DMSO in Embryonic Media

823   (EM) or 10 mM Metronidazole in EM. Larvae were treated overnight and washed 3 times in EM

824   before entering the nursery at 6 dpf. Tanks for both conditions were assessed daily to assess

825   survival rate and imaged to confirm loss or return of fluorescence signal.

Inhibitor drug screen

Zebrafish embryos were screen for *foxc1b:*Gal4;UAS:mCherryNTR fluorescence at

approximately 18 hpf and chorions were digested enzymatically with 2 mg/mL Pronase. Fish

carrying *tbx18:*DsRed2 could not be screened before drug treatments. DAPT (Tocris, 2634), 4-

Diethylaminobenzaldehyde (Sigma-Aldrich, D86256), Dorsomorphin (Selleckchem S7840),

Infigratinib (Selleckchem, S2183), LY2109761 (Selleckchem, S2704), and XAV-939

(Selleckchem, S1180) were dissolved in DMSO and aliquoted as stock solutions. DAPT,

Infigratinib, and LY2109761 stock solutions were added to heated E3 and cooled to room

temperature before treating embryos. DEAB and Dorsomorphin stocks were diluted in room

temperature Embryo Medium. XAV-939 stock was added to heated Embryo Medium and cooled

to room temperature. Cyclopamine (Selleckchem S1146) was dissolved in Ethanol, and stock was

diluted into heated E3 and cooled to room temperature. PTU media was added to all drug

solutions, regardless of whether prepared in EM or E3, to a final concentration of 197 µM. Fresh

drug was added daily for animals treated from 24-72 hpf. For transient inhibition experiments,

drug media was replaced with 197 µM PTU in embryo medium every 24 hours until experiment

termination. Animals were euthanized and fixed in 4% PFA at 4ºC overnight prior to imaging.

Quantification and statistical analysis

For *foxc1a/*b allelic series, regions were defined by manually drawing ROIs around brain domains.

Fluorescent signal in the brain or deeper tissues was removed from the z-stack to avoid

misidentifying as dorsal meninges signal. Max projections of edited z-stacks were then

thresholded and measured for fraction of mCherry+ cells using ImageJ. For quantifying mCherry

expression around the perimeter of 5 dpf cryosections, brain domains and mCherry+ meninges

were manually defined in ImageJ by establishing ROI perimeters in ImageJ. For meninges

ablation experiments, entire calvaria were defined by manually drawing a ROI around the frontal

and parietal bones using ImageJ. Hole regions were defined by manually drawing a ROI around

the area where bone was absent in the calvaria. The measure tool was then used to obtain the

areas of the entire calvaria and the hole regions in pixels. To quantify the size of each hole relative

to the whole calvaria size, a ratio of hole area divided by entire calvaria area was calculated for

each sample. For osteoblast:nlsEOS experiments, regions were defined by manually drawing a

ROI across bone fronts or cranial sutures using ImageJ. The channels were then split and nuclei

within each channel were counted manually. De novo osteoblasts were defined as green only

cells. All statistical analyses were completed using PRISM.

Data availability

The raw and processed sequencing data for single-cell genomics datasets in this study are

available on GEO (accession number: GSE318179).

Acknowledgements

The authors thank the Bone Histomorphometric Laboratory, the Brain Research Institute

Microscopic Techniques and Electron Microscopy, and Broad Stem Cell Research Center's Flow

Cytometry Core and Sequencing Core for their services. We thank you Susan Childs for providing

the crucial *foxc1b*: Gal4;UAS:mCherryNTR fish line.

Contributions

A.L.A, K.W., H.C., and D.T.F conceived and designed the study. A.A. and B.M.H. performed larval

experiments. K.W. and D.T.F. performed adult experiments. P.R. and K.W. generated and

described the *wu:fj16a03*:nlsEOS reporter. H.C performed single-cell and bioinformatic analysis.

D.T.F. supervised the research. D.T.F. wrote the manuscript with support from A.L.A, K.W., and

H.C.

Funding:

HHMI Hanna H. Gray Fellows Program (D.T.F), HHMI Freeman Hrabowski Program (D.T.F.)

Competing interests

The authors have no competing interests to declare.

References

1. Aldinger, K.A., Lehmann, O.J., Hudgins, L., Chizhikov, V.V., Bassuk, A.G., Ades, L.C., Krantz, I.D., Dobyns, W.B., Millen, K.J., 2009. FOXC1 is required for normal cerebellar development and is a major contributor to chromosome 6p25.3 Dandy-Walker malformation. Nat Genet 41, 1037-1042.

2. Asakawa, K., Suster, M.L., Mizusawa, K., Nagayoshi, S., Kotani, T., Urasaki, A., Kishimoto, Y., Hibi, M., Kawakami, K., 2008. Genetic dissection of neural circuits by Tol2 transposon-mediated Gal4 gene and enhancer trapping in zebrafish. Proc Natl Acad Sci U S A 105, 1255-1260.

3. Barske, L., Fabian, P., Hirschberger, C., Jandzik, D., Square, T., Xu, P., Nelson, N., Yu, H.V., Medeiros, D.M., Gillis, J.A., Crump, J.G., 2020. Evolution of vertebrate gill covers via shifts in an ancient Pou3f3 enhancer. Proc Natl Acad Sci U S A 117, 24876-24884.

4. Borrell, V., Marin, O., 2006. Meninges control tangential migration of hem-derived Cajal-Retzius cells via CXCL12/CXCR4 signaling. Nat Neurosci 9, 1284-1293.

5. Choe, Y., Zarbalis, K.S., Pleasure, S.J., 2014. Neural crest-derived mesenchymal cells require Wnt signaling for their development and drive invagination of the telencephalic midline. PLoS One 9, e86025.

6. Como, C.N., Kim, S., Siegenthaler, J., 2023. Stuck on you: Meninges cellular crosstalk in development. Curr Opin Neurobiol 79, 102676.

7. Como, C.N., O'Rourke, R., Winkler, C., Mitchell, D., Tran, L., Lorberbaum, D., Sussel, L., Franco, S., Siegenthaler, J., 2025. Meningeal-derived retinoic acid regulates neurogenesis via suppression of Notch and Sox2. Cell Rep 44, 115637.

8. Cooper, G.M., Durham, E.L., Cray, J.J., Jr., Siegel, M.I., Losee, J.E., Mooney, M.P., 2012. Tissue interactions between craniosynostotic dura mater and bone. J Craniofac Surg 23, 919-924.

9. Cronan, M.R., Tobin, D.M., 2019. Endogenous Tagging at the cdh1 Locus for Live Visualization of E-Cadherin Dynamics. Zebrafish 16, 324-325.

10. Dasgupta, K., Jeong, J., 2019. Developmental biology of the meninges. Genesis 57, e23288.

11. Davison, J.M., Akitake, C.M., Goll, M.G., Rhee, J.M., Gosse, N., Baier, H., Halpern, M.E., Leach, S.D., Parsons, M.J., 2007. Transactivation from Gal4-VP16 transgenic insertions for tissue-specific cell labeling and ablation in zebrafish. Dev Biol 304, 811-824.

12. Decimo, I., Fumagalli, G., Berton, V., Krampera, M., Bifari, F., 2012. Meninges: from protective
membrane to stem cell niche. Am J Stem Cells 1, 92-105.

13. Derk, J., Como, C.N., Jones, H.E., Joyce, L.R., Kim, S., Spencer, B.L., Bonney, S., O'Rourke, R.,
Pawlikowski, B., Doran, K.S., Siegenthaler, J.A., 2023. Formation and function of the meningeal
arachnoid barrier around the developing mouse brain. Dev Cell 58, 635-644 e634.

14. DeSisto, J., O'Rourke, R., Jones, H.E., Pawlikowski, B., Malek, A.D., Bonney, S., Guimiot, F., Jones,
K.L., Siegenthaler, J.A., 2020. Single-Cell Transcriptomic Analyses of the Developing Meninges
Reveal Meningeal Fibroblast Diversity and Function. Dev Cell 54, 43-59 e44.

15. Ebert, S.N., Eisner, C., Karamboulas, K., Bernier, L.P., Kaplan, D.R., MacVicar, B.A., Miller, F.D.,
2025. Single-Cell Approaches Define the Murine Leptomeninges: Cortical Brain Interface as a
Distinct Cellular Neighborhood Composed of Neural and Non-neural Cell Types. eNeuro 12.

16. Farmer, D.T., Dukov, J.E., Chen, H.J., Arata, C., Hernandez-Trejo, J., Xu, P., Teng, C.S., Maxson, R.E.,
Crump, J.G., 2024. Cellular transitions during cranial suture establishment in zebrafish. Nat
Commun 15, 6948.

17. Farmer, D.T., Mlcochova, H., Zhou, Y., Koelling, N., Wang, G., Ashley, N., Bugacov, H., Chen, H.J.,
Parvez, R., Tseng, K.C., Merrill, A.E., Maxson, R.E., Jr., Wilkie, A.O.M., Crump, J.G., Twigg, S.R.F.,
2021. The developing mouse coronal suture at single-cell resolution. Nat Commun 12, 4797.

18. Galanternik, M.V., Castranova, D., Gober, R.D., Nguyen, T., Kenton, M., Margolin, G., Kraus, A.,
Sur, A., Dye, L.E., Pham, V., Maese, A., Holmgren, M., Gore, A.V., Samasa, B., Goldstein, A., Davis,
A.E., Swearer, A.A., Iben, J., Li, T., Coon, S.L., Dale, R.K., Farrell, J.A., Weinstein, B.M., 2025.
Anatomical and Molecular Characterization of the Zebrafish Meninges. bioRxiv.

19. Greenwald, J.A., Mehrara, B.J., Spector, J.A., Warren, S.M., Crisera, F.E., Fagenholz, P.J.,
Bouletreau, P.J., Longaker, M.T., 2000. Regional differentiation of cranial suture-associated dura
mater in vivo and in vitro: implications for suture fusion and patency. J Bone Miner Res 15, 2413-
2430.

20. Haushalter, C., Schuhbaur, B., Dolle, P., Rhinn, M., 2017. Meningeal retinoic acid contributes to
neocortical lamination and radial migration during mouse brain development. Biol Open 6, 148-
160.

21. Inoue, T., Ogawa, M., Mikoshiba, K., Aruga, J., 2008. Zic deficiency in the cortical marginal zone
and meninges results in cortical lamination defects resembling those in type II lissencephaly. J
Neurosci 28, 4712-4725.

22. Kague, E., Roy, P., Asselin, G., Hu, G., Simonet, J., Stanley, A., Albertson, C., Fisher, S., 2016.
Osterix/Sp7 limits cranial bone initiation sites and is required for formation of sutures. Dev Biol
413, 160-172.

23. Kappers, C.U.A., 1925. The meninges in Cyclostomes Selachians and teleosts compared with
those in man. P K Akad Wet-Amsterd 28, 72-80.

24. Kearns, N.A., Iatrou, A., Flood, D.J., De Tissera, S., Mullaney, Z.M., Xu, J., Gaiteri, C., Bennett, D.A.,
Wang, Y., 2023. Dissecting the human leptomeninges at single-cell resolution. Nat Commun 14,
7036.

25. Kikuchi, K., Gupta, V., Wang, J., Holdway, J.E., Wills, A.A., Fang, Y., Poss, K.D., 2011. tcf21+
epicardial cells adopt non-myocardial fates during zebrafish heart development and
regeneration. Development 138, 2895-2902.

26. Kimura, Y., Hisano, Y., Kawahara, A., Higashijima, S., 2014. Efficient generation of knock-in
transgenic zebrafish carrying reporter/driver genes by CRISPR/Cas9-mediated genome
engineering. Sci Rep 4, 6545.

27. Kwan, M.D., Wan, D.C., Wang, Z., Gupta, D.M., Slater, B.J., Longaker, M.T., 2008. Microarray
analysis of the role of regional dura mater in cranial suture fate. Plast Reconstr Surg 122, 389-
399.

28. Lawson, N.D., Weinstein, B.M., 2002. In vivo imaging of embryonic vascular development using
transgenic zebrafish. Dev Biol 248, 307-318.

29. Matrongolo, M.J., Ho-Nguyen, K.T., Jain, M., Ang, P.S., Reddy, A., Schaper, S., Tischfield, M.A.,
2023. Loss of Twist1 and balanced retinoic acid signaling from the meninges causes cortical
folding in mice. Development 150.

30. Miesfeld, J.B., Link, B.A., 2014. Establishment of transgenic lines to monitor and manipulate
Yap/Taz-Tead activity in zebrafish reveals both evolutionarily conserved and divergent functions
of the Hippo pathway. Mech Dev 133, 177-188.

31. Mishra, S., Choe, Y., Pleasure, S.J., Siegenthaler, J.A., 2016. Cerebrovascular defects in Foxc1
mutants correlate with aberrant WNT and VEGF-A pathways downstream of retinoic acid from
the meninges. Dev Biol 420, 148-165.
32. Nakao, T., 1979. Electron microscopic studies on the lamprey meninges. J Comp Neurol 183, 429-
453.
33. O'Rahilly, R., Muller, F., 1986. The meninges in human development. J Neuropathol Exp Neurol
45, 588-608.
34. Opperman, L.A., Sweeney, T.M., Redmon, J., Persing, J.A., Ogle, R.C., 1993. Tissue interactions
with underlying dura mater inhibit osseous obliteration of developing cranial sutures. Dev Dyn
198, 312-322.
35. Pietila, R., Del Gaudio, F., He, L., Vazquez-Liebanas, E., Vanlandewijck, M., Muhl, L., Mocci, G.,
Bjornholm, K.D., Lindblad, C., Fletcher-Sandersjoo, A., Svensson, M., Thelin, E.P., Liu, J., van
Voorden, A.J., Torres, M., Antila, S., Xin, L., Karlstrom, H., Storm-Mathisen, J., Bergersen, L.H.,
Moggio, A., Hansson, E.M., Ulvmar, M.H., Nilsson, P., Makinen, T., Andaloussi Mae, M., Alitalo,
K., Proulx, S.T., Engelhardt, B., McDonald, D.M., Lendahl, U., Andrae, J., Betsholtz, C., 2023.
Molecular anatomy of adult mouse leptomeninges. Neuron 111, 3745-3764 e3747.
36. Siegenthaler, J.A., Ashique, A.M., Zarbalis, K., Patterson, K.P., Hecht, J.H., Kane, M.A., Folias, A.E.,
Choe, Y., May, S.R., Kume, T., Napoli, J.L., Peterson, A.S., Pleasure, S.J., 2009. Retinoic acid from
the meninges regulates cortical neuron generation. Cell 139, 597-609.
37. Slater, B.J., Kwan, M.D., Gupta, D.M., Amasha, R.R., Wan, D.C., Longaker, M.T., 2008. Dissecting
the influence of regional dura mater on cranial suture biology. Plast Reconstr Surg 122, 77-84.
38. Slater, B.J., Kwan, M.D., Gupta, D.M., Lee, J.K., Longaker, M.T., 2009. The role of regional posterior
frontal dura mater in the overlying suture morphology. Plast Reconstr Surg 123, 463-469.
39. Sur, A., Wang, Y., Capar, P., Margolin, G., Prochaska, M.K., Farrell, J.A., 2023. Single-cell analysis
of shared signatures and transcriptional diversity during zebrafish development. Dev Cell 58,
3028-3047 e3012.
40. Teng, C.S., Ting, M.C., Farmer, D.T., Brockop, M., Maxson, R.E., Crump, J.G., 2018. Altered bone
growth dynamics prefigure craniosynostosis in a zebrafish model of Saethre-Chotzen syndrome.
Elife 7.

41. Thomas, E.D., Raible, D.W., 2019. Distinct progenitor populations mediate regeneration in the
zebrafish lateral line. Elife 8.

42. Topczewska, J.M., Shoela, R.A., Tomaszewski, J.P., Mirmira, R.B., Gosain, A.K., 2016. The
Morphogenesis of Cranial Sutures in Zebrafish. PLoS One 11, e0165775.

43. Whitesell, T.R., Chrystal, P.W., Ryu, J.R., Munsie, N., Grosse, A., French, C.R., Workentine, M.L., Li,
R., Zhu, L.J., Waskiewicz, A., Lehmann, O.J., Lawson, N.D., Childs, S.J., 2019. foxc1 is required for
embryonic head vascular smooth muscle differentiation in zebrafish. Dev Biol 453, 34-47.

44. Xu, P., Balczerski, B., Ciozda, A., Louie, K., Oralova, V., Huysseune, A., Crump, J.G., 2018. Fox
proteins are modular competency factors for facial cartilage and tooth specification.
Development 145.

45. Zarbalis, K., Siegenthaler, J.A., Choe, Y., May, S.R., Peterson, A.S., Pleasure, S.J., 2007. Cortical
dysplasia and skull defects in mice with a Foxc1 allele reveal the role of meningeal differentiation
in regulating cortical development. Proc Natl Acad Sci U S A 104, 14002-14007.

1011