Winding-Up of Fibrin Fibers as a Novel Mechanism of Platelet-Mediated Fiber Compaction

Alexei Grichine[1], Tatiana Kovalenko[2,3], Florence Appaix[1], Anne-Sophie Ribba[4], Anita Eckly[5], Jean-Yves Rinckel[5], Mikhail Panteleev[2,3,6], Laurence Lafanechère[4] and Karin Sadoul[4]

1 University Grenoble Alpes, INSERM U1209, CNRS UMR 5309, Optical microscopy and cell imaging (MicroCell) facility, Institute for Advanced Biosciences, 38000 Grenoble, France
2 Center for Theoretical Problems of Physico-Chemical Pharmacology, Russian Academy of Sciences, 30 Srednyaya Kalitnikovskaya str., 109029, Moscow, Russia
3 Dmitry Rogachev National Medical Research Center of Pediatric Hematology, Oncology and Immunology, 1 Samory Mashela str., 117997, Moscow, Russia
4 Université Grenoble Alpes, INSERM U1209, CNRS UMR 5309, Institute for Advanced Biosciences (IAB), 38000 Grenoble, France
5 University of Strasbourg, INSERM, EFS Grand Est, BPPS UMR-S 1255, FMTS, F-67065 Strasbourg, France
6 Physics Faculty, M. V. Lomonosov Moscow State University, Moscow, Russian Federation

corresponding author:
Karin Sadoul
Institute for Advanced Biosciences
Research Center UGA / Inserm U 1209 / CNRS 5309
Team: Cytoskeletal Dynamics and Nuclear Functions
Site Albert Bonniot - Allée des Alpes
38700 La Tronche, France
Phone : +33 (0)4 76 54 95 71
e-mail: karin.sadoul@univ-grenoble-alpes.fr

Abstract

This study reveals a previously unrecognized mechanism by which platelets retract and compact fibrin fibers. Using a newly developed 2D fiber-retraction assay, we observed an initial "gearwheel" pattern of actin–myosin organization in spread platelets with associated, extracellular fibrin patches that appear to form an initiation complex for fibrin fiber attachment and rearrangement. The final outcome of this process results in platelets surrounded by tightly packed fibrin fibers, mirroring the architecture of platelets and adjacent fibers within a retracted clot. Thus, the observed compaction process might also take place during clot retraction in order to reduce clot volume, stiffen the clot and enhance wound repair. Apart from pulling on fibers like on a rope, platelets actively wind-up fibrin fibers into compact structures, similar to balls of wool. Besides DNA packaging, this represents a new example of a natural fiber compaction mechanism. Using a combination of 3D clot-retraction and 2D fiber-retraction assays, expansion and electron microscopy, live imaging and mathematical modeling, we show that platelets use an actomyosin-driven swirling motion to gather and loop fibrin fibers around the base of bulbous protrusions ("bulbs"). These bulbs form when a platelet becomes trapped between fibrin fibers during clot retraction or 2D fiber-retraction assays. These findings complement and extend earlier models of platelet-mediated fibrin fiber retractions, offering new insight into how platelets mechanically organize fibrin fibers.

Introduction

Platelets are absolutely essential in case of vessel injury. They are not only necessary to initiate hemostasis but also to retract the forming clot thereby pulling together the edges of the ruptured vessel and flattening the clot for improved blood flow. Different approaches to visualize fibrin fibers in whole blood or plasma clots have shown that fibrin fibers represent the main mechanical and structural framework of clots,[1,2] occupying a substantial portion of the clot volume. Previous studies[3-6] have shown that platelets are able to retract an unconstrained plasma clot to a tiny volume. The extent and rate of clot retraction are influenced by several factors, including platelet and red blood cell counts, fibrin and thrombin concentrations, and the geometric context of clot formation.[7] However, platelets are the primary cellular agents driving such extensive clot compaction, although erythrocytes also contribute.[8] How do platelets accomplish this tremendous task? Two key studies offer insight: While Fletcher's lab demonstrated that platelets generate contractile forces comparable to muscle cells,[9] Weisel's group provided mechanistic insight, showing that platelets extend filopodia to bind and pull on fibrin fibers, much like tugging on ropes[10] causing localized fiber accumulation.

However, pulling alone may not fully account for the dramatic reduction in clot volume. We hypothesized that efficient clot retraction requires substantial densification of the fibrin fiber network. Using expansion microscopy, we observed that platelets wind-up fibrin fibers similar to balls of wool around individual bulbs. Using a "2D fiber-retraction assay" we found that fibrin patches are initially arranged in a ring-like pattern in close proximity to the circular organization of the actin cytoskeleton in spread platelets. Over time, wound-up fibrin fibers accumulate above the plasma membrane of platelets in a myosin dependent way.

To explore the mechanistical aspects of the platelet mediated fiber compaction we used computational modeling and live-cell video microscopy. Our results suggest that actomyosin-driven swirling within spread platelets on a 2D surface or within individual bulbs of platelets in the 3D clot space may underlie the observed fibrin coiling and compaction.

Results

Platelets become labeled by fluorescent fibrin fibers during clot retraction

We first estimated the capacity of platelets to retract diluted plasma clots (12% plasma) under unconstrained conditions (Fig. 1A). Under these conditions, 4×10^7 platelets are able to retract the clot volume to approximately 1.5% of its initial volume (400 µl) within 60 min. Even platelets kept overnight at room temperature before the retraction assay were still retracting the clot to the same extent, however after a longer lag phase. We then used live imaging to follow the reorganization of fibrin fibers during unconstrained clot retraction. Platelet rich plasma (PRP) was spiked with fibrinogen-Alexa 488 before thrombin induced clot formation and image acquisition. Very rapidly one can observe the formation of fluorescent fibrin nodes during the retraction process. Since under the assay conditions the clot is composed only of platelets and fibrin fibers, we hypothesized that the nodes may correspond to platelets surrounded by fibrin fibers (Fig. 1B and video 1). To test this hypothesis, we performed a time course analysis of clot retraction in presence of fluorescent fibrinogen under constrained conditions preventing complete retraction. This approach allowed us to stain and visualize individual platelets in a clot even after prolonged retraction times. Hemostatic clots that form in vivo following vessel injury are also likely to experience mechanical constraints, as they remain anchored to the damaged vessel walls while spanning the breach. Retracted clots were fixed and stained for the αIIb-integrin subunit to detect the localization of platelets within the clot. At the position of each platelet, we consistently observe colocalization of a fibrin node at all time points examined (Fig. 1C), confirming our hypothesis that the fibrin nodes observed in figure 1B form at platelet sites. Furthermore, the results show that the build-up of fibrin fibers around individual platelets occurs very rapidly, within 10 min. The fibrin fibers forming the nodes did not decorate the entire platelet surface but surrounded the center of platelets embedded in the clot. Please note, that platelets occupy a larger space after longer retraction times, but the fiber nodes persist even after 80 min of retraction.

To gain a better insight about the organization of fibrin fibers around platelets within a clot, we used 4x isotropic expansion microscopy. We found that fibrin fibers form a cage-like structure around the platelet center, clearly visible when scrolling through the image stacks (Fig. 2 A-D and animation, video 2). This pattern was consistently observed across all platelets in the clot and after different clot retraction times (not shown). Most platelets had formed large bulbs, thin filopodia, or both with fibrin fibers typically localized around the base of the bulbs or along filopodia. It remains unclear whether all fibrin segments reside on the external platelet membrane or whether some of them extend into channels of the open canalicular system (OCS). Some of these channels persist in activated platelets.[11] Please note that straight fibers radiate from the platelet, indicating a high tension in the platelet clot environment. We want to emphasize that changes in temperature, or plasma/platelet concentrations change the kinetics of retraction but we always observe the cage-like fibrin fiber organization around individual platelets (not shown).

Platelets, spread on a 2D surface, organize fibrin fibers above them

So far, our results demonstrate that platelets possess a remarkable capacity to precisely organize fibrin fibers around themselves when embedded in a clot. To investigate the mechanism by which platelets arrange the fibers around their center we sought to use a model system with reduced complexity. We, therefore, developed a "2D fiber-retraction assay", which is a reductionist approach that does not replicate full clot retraction, but can provide insight into how platelets interact with and organize fibrin fibers. To this end, a

diluted suspension of PRP (2.5×10^6 platelets/ml, 4 μl plasma/ml PBS) was incubated with fluorescent fibrinogen and thrombin for 15 min to induce platelet activation and fibrin polymerization. Platelets were then allowed to spread on a 2D surface for 30 min before fixation. Under these conditions, we observe four categories of spread platelets and associated fibrin fiber organizations. Platelets without fibrin fiber contact (28.3±2%) and platelets with a small fibrin dot at their center, which could be a fibrin initiation complex (40±3%). Another category of platelets had an accumulation of coiled fibers above them (fiber winding 18.3±0.5%) and a fourth category showed strong fiber densification around them (fiber compaction 13.6±0.9%) (Fig. 3, left image of A, quantification B). We then investigated, whether these fiber accumulations depend on myosin actions as it is the case for clot retraction. Myosin inhibition using blebbistatin strongly diminished the build-up of fibers around the spread platelets (fiber-winding 6.6±0.9% and fiber compaction 0.2±0.2%), while a higher percentage of platelets with fiber initiations or fiber contacts was observed (63.4±2%) (Fig. 3, right image of A and quantification B).

Then, we used the 2D fiber-retraction assay followed by expansion microscopy to investigate in more detail how the fibers above spread platelets are organized. Platelets were stained either for the αIIb integrin subunit or for myosin. We observed that fibrin fibers are wound-up above platelets (category "fiber winding" in fig. 3), frequently forming ball-like structures wrapped around a bulbous protrusion in the middle of the spread platelet, often referred to as pseudo-nucleus in the literature (Fig. 4A-E and animation, Video 3).

In addition to these spread platelets with coiled fibrin fibers above them, we also observed platelets entangled by many fibers and therefore unable to spread completely. These platelets had formed filopodia and bulbs similar to platelets within a clot and a dense fiber accumulation is observed around the base of their bulbs (category "fiber compaction" in fig. 3A). These fibers appeared to get coiled into multiple loops, tightly compacted around the base of platelet bulbs similar to individual balls of wool (six small "balls" can be distinguished in Fig. 5A, focal planes at 4 and 5.25 μm and animation, video 4, see also Fig. 6 and animation, video 5). Winding-up of fibers by platelets implies that fibers around individual platelets may get twisted and thereby bundled. Local twisting of fibers near platelets is indeed observed in most of the presented images, although the resolution of optical microscopy is often insufficient to resolve regions of densely packed fibers. Particularly clear examples are shown in Figure 6. It should be noted that fiber bundles between two platelets can also be twisted (Fig. 6C-E). The enhanced fibrin labeling observed around these platelets, compared to those in a typical constrained clot (Fig. 2A-D), may be due to multiple winding-up of free, unconstrained fibrin fibers.

A ring-like organization of fibrin fiber patches can be observed at the center of spread platelets

We next increased the plasma volume (7 μl/ml instead of 4 μl/ml) in order to enhance the amount of formed fibrin fibers. This modification led to the formation of a fragile fibrin gel, which was partially lost during the fixation or staining steps. As a result, only very few fibers remained attached to the platelets or to the coverslip.

The residual fibrin fluorescence revealed a ring-like arrangement of fibrin patches at the platelet center. This fibrin "rosette" co-localized with a circular actin organization, forming a gearwheel-like pattern with a compacted fibrin mass at the inner edge of the circle. The actin cytoskeleton was either composed of radial and transverse fibers (Fig. 7 A,B and animation, video 6) or radial fibers and actin nodules (Fig. 7C and animation, video 6) depending on the degree of platelet spreading.[12,13] A faint rosette with a compact fibrin mass and an attached fibrin fiber is shown in figure 7D and animation, video 6).

We then investigated the localization of myosin II under these conditions and found that it is
distributed throughout the platelet. However, a denser ring-like organization of myosin is
present in the center of the spread platelet, closely associated with the fibrin rosette (Fig. 8
and animation, video 7). Some myosin spots are intercalated with compacted fibrin (Fig.8 A,
focal plane at z=7.8 µm), while wound-up fibrin fibers can be observed at the periphery and
above the myosin ring structure (Fig. 8A-F). The intensity of fibrin rosettes is highly variable
between platelets of different donors. Intense rosettes as shown in figure 7 were observed
for platelets from two donors (approximately 3-5% of spread platelets with fiber initiation
complexes), while faint rosettes (as in Fig. 8AB) were observed for 50% of spread platelets
with initiation complexes using blood from three other donors.
190
Modeling platelet-mediated winding-up of fibrin fibers
To address the mechanical aspects of the observed platelet-mediated fibrin fiber
organizations, we developed a model based on our experimental observations. We first
quantified fiber curvature and the number of platelet extensions. Each platelet (in clots or on
a glass surface surrounded by fibers and therefore unable to spread) showed 7-10
bulbs/filopodia (Fig. 9A-C). Fiber curvature was determined using transmission electron
microscopy (TEM) and fluorescent (FLUO) images. Curvatures were similar for both image
types (mean values for κ_{TEM}=10.09 ± 6.6 μm^{-1} and κ_{FLUO}=6.77 ± 4.7μm^{1}) with differences likely
due to the higher resolution of TEM (Fig. 9D-F).
Our model incorporates these observations, published parameters (Table 1, methods
section) and builds on the framework established by Bershadsky and colleagues.[14] They
demonstrated that in fibroblasts, cultured on circular substrates, the actin cytoskeleton is
composed of radial actin fibers extending towards the cell center and circular transverse
fibers enriched in myosin-IIA. Live-cell imaging in their study revealed that the transverse
fibers slide centripetally along the radial fibers, generating a chiral swirling motion directed
towards the cell center. Fibronectin-coated beads bound to integrins on the cell surface are
dragged along with the intracellular swirling of the cytoskeleton suggesting that
extracellularly bound fibrin fibers may experience a similar rotational motion. Chiral rotations
of the cytoskeleton are not only observed at the level of entire cells, but also in subcellular
regions, such as individual filopodia of the neuronal growth cone.[15,16]
Along this line, our simulations suggest that fibrin fibers are anchored to integrins in the
plasma membrane of bulbs and filopodia, are pulled towards the platelet center by
retraction of filopodia, as demonstrated by Kim et al.[10] Spiral cytoskeletal motions within
each bulb drive integrins, and thus the attached fibrin fibers, towards the platelet center,
wrapping the fibers around the base of the bulbs (Fig. 10A and animation, video 8). The
efficiency of loop formation depends on the initial fiber position (Fig. 10B-D). The process
effectively compacts long fibers into loops with a radius of ~420-nm near the base of bulbs
(Fig. 10D).
219
Transmission electron microscopy (TEM) provides evidence of curved and cross-sectioned
fibrin fibers at the base of platelet bulbs
To confront the results obtained by expansion microscopy and model simulations with those
using TEM, we analyzed TEM images of platelets and fibrin fibers within a clot. We observed
numerous instances of cross-sectioned fibrin fibers at the base of bulbs as illustrated in
Figure 11. This figure also shows two examples suggesting fibers that curve along bulbs from
top to bottom and fibers attached to a filopodium.
227
228

Live imaging of platelet-mediated reorganization and compaction of fibrin fibers

We performed live video microscopy using platelets from three different donors to observe how platelets wind-up fibrin fibers (Fig. 12 and video 9). In many cases, fibrin fibers were already accumulated above platelets at the start of imaging, with minimal further fiber movements or compactions; likely due to fibers binding to each other, to the dish or to the platelet surface (Fig. 12A). However, dynamic fiber movements, both clockwise and counterclockwise, were observed above two platelets present within the rectangle indicated in figure 12A (see zoom images lower panel). An arrow in Fig. 12A marks the formation of a kink in a fibrin fiber. For the second experiment (blood from a different donor), we reduced the incubation temperature to 21°C in order to slow down the platelet and fiber movements. In the rectangle indicated in figure 12B, a fibrin fiber gets progressively curved and eventually ruptures (at time point 00:48). The resulting free end of the fiber then undergoes a clockwise rotation (see zoom images lower panel). Another platelet producing a fiber densification is marked by an arrow in figure 12B. The video shows a counterclockwise rotation of the central part of the platelet during the fiber densification process. A third experiment (platelets from another donor), again performed at room temperature, shows the rotational (counterclockwise) movement of the pseudo-nucleus and alongside the winding-up of fibers around this bulbous protrusion. This apparently generates strong tension that pulls the pseudo-nucleus towards the edge of the spread platelet, where the movement ceases (Fig. 12C, see zoom images lower panel). These fibrin fiber movements occurred without visible filopodia-fiber interactions, suggesting that the bound fibers are dragged by intracellular movements of the cytoskeleton. Since still images cannot fully illustrate the fiber movements or the rotation of the platelet center, we invite the reader to consult the accompanying time-lapse videos (video 9). After the time-lapse acquisitions, samples were fixed and stained for the αIIb-integrin subunit to localize the position of platelets on the dish surface. The final fibrin fiber organization around platelets in these live videos of the 2D fiber-retraction assays closely resembles the arrangement of fibers around platelets in a constrained clot, except that some platelets are completely spread on the surface with no or very few fiber contacts. Platelets surrounded by densely packed fibers had formed bulbs and filopodia, but remained attached to the dish surface (Fig. 12D).

To minimize pre-imaging fiber buildup, we next added preformed fibrin fibers to pre-spread platelets and started imaging immediately. Under these conditions, surface coating is observed and only a few fibers are present, yet a central fibrin accumulation is consistently observed on each platelet, possibly representing a nucleation site. Among 35 platelets analyzed, 15 displayed a central fibrin mass rotating counterclockwise (viewed from below), 4 rotating clockwise, and 16 showed no consistent directional movement (Fig. 12E). In some instances, the rotating complexes stalled and stopped moving. These observations support the idea that cytoskeletal swirling drives fibrin reorganization above spread platelets.

Discussion

Using a 2D fiber-retraction assay, we identified a previously unrecognized capacity of platelets to compact fibrin fibers into a small volume by winding them into coiled structures during platelet induced fiber retractions. To our knowledge the only example of natural fiber compaction is coiling of DNA molecules.[17] In contrast to intracellular, enzyme-mediated DNA compaction, extracellular fibrin fiber compaction requires the cellular activity of platelets. Our investigation began with the observation that platelets become labeled with fluorescent fibrinogen during clot retraction as previously shown by Brzoska et al.[18] They demonstrated that fibrin binding to thrombin activated platelets depends on αIIbβ3 integrins and the actin cytoskeleton. While fibrin fiber retraction by platelets in a clot has been attributed to non-muscle myosin II,[19,20] the mechanism underlying fiber organization and compaction has remained unclear.

One key mechanism, revealed in previous studies, is the extension of platelet filopodia that bind to fibrin fibers and pull them inwards in a hand-over-hand motion.[10] This is clearly the main process by which platelets retract fibrin fibers in a clot. However, this mechanism alone cannot explain the cage-like fiber organization observed around platelets in a constrained clot (Fig. 2). Our findings introduce an additional, distinct mechanism by which platelets can organize fibrin fibers. Platelets wind-up fibrin fibers into dense, ball-like structures, which is an efficient form of compaction (Fig. 3-6). Thus, our model reveals an additional mechanistic aspect of platelet mediated fiber retraction beyond those described by Michael et al.[21] and Kliuchnikov et al.,[22] so far observed only in the 2D fiber-retraction assay. Nevertheless, the platelet morphology and the organization of fibers around them after the retraction process closely resemble those of platelets in a clot and both, 2D and 3D fiber retractions, depend on myosin actions.

The 2D fiber-retraction assay may recapitulate early post-injury events, such as activation of platelets in suspension triggered by agonists released from neighboring platelets, ultimately leading to the spreading of activated platelets onto the subendothelial matrix to anchor the developing clot at the site of vessel injury. This assay revealed two key observations. First, when many fibers are present, some platelets spread with the typical fried-egg phenotype and coil-up fibers above their pseudo-nucleus (Fig. 3, 4). Others form large bulbs in several directions, similar to platelets in a clot, and compacted fibers are observed around the base of each bulb (Fig. 5, 6). Under these unconstrained 2D conditions, fiber accumulation around the bulbs is much higher than in the case of a constrained 3D clot (Fig. 2), where internal tensions limit fiber winding.

Second, when fewer fibers are retained on the culture surface, several spread platelets display a fibrin rosette around their pseudo-nucleus (Fig. 7,8) that colocalizes with actin, myosin and integrins. A dense fibrin mass is seen at the inner periphery of this rosette, sometimes with an attached fiber. This gearwheel-like architecture may represent an early priming/nucleation step of platelet-induced winding of fibrin fibers. The actin cytoskeleton in platelets exhibits a chiral organization[23] which, combined with the action of myosin, could be responsible for the swirling motion observed in spread platelets (Fig.12 E-G and video 9). Integrin-bound extracellular fibers are pulled along by this vortex inducing their winding-up above the pseudo-nucleus (Fig. 12C). Fibrin rosettes may fade as coiled fibers become more prominent, suggesting a transition from an initiation phase to active coiling and compaction.

Ideas and Speculations

How this 2D "gearwheel" translates to the 3D clot environment remains unclear. One possibility is that the pseudo-nucleus may correspond to a bulb around which fibers are coiled and accumulate. Platelets with bulbs within a clot have been described previously.[24-27]

We wondered whether platelet bulbs might represent plasma membrane blebs similar to
membrane protrusions formed by megakaryocytes during clot retraction.[28] However, the
consistent presence of 8–9 bulbs per platelet (Fig. 9A-C) suggests that they reflect a "three-
dimensional spreading" morphology, reminiscent of the eight corners of a cube, allowing
platelets to sense the clot's 3D environment along the main directions. Unlike blebs, these
bulbs often contain organelles such as mitochondria (Fig. 9D, upper image and Fig. 11B)
implying that each bulb may have its own energy supply and actomyosin organization. This
could enable swirling within each bulb and explain the observed fibrin fiber accumulation at
their base. Thus, at the onset of clot retraction, each emerging bulb may act as a discrete
functional unit onto which the gearwheel assembles, enabling fibrin fibers to attach and
initiating their subsequent winding and compaction (see scheme, Fig. 13). Indeed, blebs of
megakaryocytes retracting a clot may behave similarly as platelet bulbs since fibrin fibers can
be seen surrounding the base of blebs during megakaryocyte-mediated clot retraction, as
can be observed in the study by Kim et al. (see the supplementary movie S3 of this
publication).[28]
333

334
335
Figure 13: Schematic illustration of the hypothesis explaining the potential mechanism by which
platelets may wind-up fibrin fibers and compact them.
Represented is a platelet in the 2D fiber-retraction assay (upper part) or in a clot (lower part). At the
beginning of platelet fiber interactions, a fibrin rosette is present around the pseudo-nucleus and in
each bulb with a dense fiber mass at the internal periphery of the rosette possibly representing an
initiation complex. Radial actin fibers extend to the center of the platelet and an actomyosin
rotational movement along the radial actin fibers may lead to the winding-up and the downward
sliding of the extracellular attached fibers to encircle the pseudo-nucleus or the base of the bulbs at
the final state of compaction (3D rendering of the scheme was produced using an open AI).
345

Our live time-lapse videos of platelet-mediated fiber movements in a 2D environment
support the concept of a cytoskeletal vortex that drives the coiling and compaction of
extracellular attached fibers. Additionally, the other time-lapse video capturing the rotation
of compacted fibrin at the platelet center further suggests the presence of cytoskeletal
swirling within spread platelets. This swirling may play a role in scanning the extracellular
space for forming fibers that can be anchored to the initiation complex. The process occurs
on a notably slow time scale, which could facilitate effective fiber capture by the rotating
complex. However, this rate appears insufficient to account for the rapid labeling of platelets
during clot formation, suggesting that regulatory mechanisms, possibly involving mechanical
feedback, may be at play. Indeed, as mechanosensory cellular particles, platelets adjust their
contractile forces and dynamics in response to environmental cues such as fiber tension and
clot stiffness.[9,10,29,30,22]

In conclusion, our study identifies a novel mechanistic aspect of platelet function: the active
winding-up and compaction of fibrin fibers. This complements the existing model of platelet-
mediated clot retraction[10] and provides a new perspective on how platelets may contribute
to clot architecture, mechanical stability, and wound healing.

Materials and methods

Reagents and antibodies

Reagents: Neutral formalin (Sigma-Aldrich, Saint-Quentin Fallavier, France), Pluronic F-127 (Sigma-Aldrich, P2443), thrombin (Sigma-Aldrich, T4648), Jasp-K4-HA (a HAK-actin probe, kind gift of Paul Guichard/Virginie Hamel),[31] apyrase (Sigma-Aldrich, A6535), heparin (Sigma-Aldrich, H3393), PGI2 (Sigma-Aldrich, P6188)

Antibodies: mouse anti-integrin αIIb (Santa-Cruz, sc-7310), monoclonal rabbit anti-HA (Cell Signaling, C29F4), rabbit anti-non muscle myosin II heavy chain (Covance, PRB-440P), rabbit anti-phospho S19 myosin light chain (Abcam, ab2480), monoclonal rabbit anti-myosin light chain 2 (Abcam, ab92721), Cy3 goat anti-mouse IgG (Jackson, 115-165-166), Cy3 goat anti-rabbit IgG (Jackson, 111-165-003), Alexa 647 goat anti-rabbit IgG (Jackson, 111-605-003)

Preparation of human platelet rich plasma (PRP) and platelet free plasma (PFP)

Whole blood was obtained from the French blood bank (age-range 18-65, mixed genders; pooled blood from multiple donors has not been used). The IRB has authorised our institute's research projects using human samples under the number DC-2022-4976. PRP was prepared from whole blood drawn into Na-citrate Vacutainers (3 tubes of 2.7 ml per donor). The blood is pooled into a 15 ml falcon tube and centrifuged for 12 min 400g, RT, no brake. The upper phase is the platelet rich plasma (PRP), in which platelets are distributed in a gradient (low concentration on top, high concentration at the bottom). In order to increase the platelet concentration in the final PRP, the upper 3 ml are removed and used to prepare platelet free plasma (PFP, by centrifugation at 12000g, 1 min, RT). The remaining upper phase is collected as the final PRP and the platelet concentration determined.

Platelet washing procedure

PRP was diluted 1/4 with Tyrode/0.35% BSA buffer and platelet activation inhibitors (0.02 U/ml apyrase, 1 μm PGI2, 10 U/ml Heparin, final concentrations) were added. Samples were then centrifuged for 15 min at 900g at RT. The supernatant was aspirated, and platelets were gently resuspended in fresh Tyrode/BSA buffer, 0.02 U/ml apyrase, 1 μm PGI2. The suspension was centrifuged as before and the platelet pellet was resuspended in fresh Tyrode/BSA buffer without inhibitors.

Clot retraction assays

We used three different clot retraction assays illustrated in supplementary figure 1.

A) Unconstrained clot retraction (Fig. 1AB): See paragraph "Live video microscopy" below for method details.

B) Constrained clot retraction between two holders (Fig. 1C): In order to perform immunofluorescence stainings of individual platelets within a clot, we used a clot retraction assay around two sterile inoculation loops as previously described.[32,33] Briefly, PRP was adjusted to 1×10^8 platelets per ml in 50% plasma/50% PBS (to reduce fibrin fiber density), fibrinogen-Alexa 488 final concentration 12.5 μg/ml and 4 μl/ml erythrocytes for color contrast. 400 μl of this suspension was used for clot induction by addition of thrombin (2.5U/ml final) and immediately transferred to 2 ml Eppendorf tubes coated with 2% Pluronic® and containing the inoculation loops. After different retraction times, clots were fixed for 1 h with isotonic formalin (9 volumes formalin / 1 volume 10xPBS). Clots were then washed with PBS and incubated ON at 4°C in 1 ml PBS/15% sucrose. Clots were further incubated in 1 ml PBS/15% sucrose/7.5 % gelatin at 37°C for 4 h and then inclusion blocks were formed in the same solution on ice. Blocks were then trimmed and snap frozen for 1

min in isopentan at -60°C and stored at -80°C until 14 μm thick transversal sections were cut
using a cryostat.

C) Constrained clot retraction within an inoculation loop: Small clots of 30 μl platelet
suspension (50% plasma/50% PBS; $1x10^7$ platelets/ml; fibrinogen-Alexa 488 final
concentration 12.5 μg/ml; 4 μl/ml erythrocytes for color contrast) were induced to form
within plastic inoculation loops (4 mm internal diameter, attached to a glass coverslip) by
addition of thrombin to a final concentration of 2.5 U/ml as described previously.[13] Five
minutes after clot induction, clots were covered with 500 μl PBS and kept in a cell culture
incubator for 10 min. Clots were then fixed with 500 μl isotonic formalin for 15 min, RT, in
the dark and then kept in PBS at 4°C until further use. Forming a clot within a small
inoculation loop, allowed us to do expansion microscopy without performing cryosections of
larger clots, which are difficult to detach from slides during the expansion process.

425
426
Supplementary Figure 1: Illustration of clot retraction assays used in this study
A) Unconstrained clot retraction
B) Constrained clot retraction between two holders
C) Constrained clot retraction within an inoculation loop

2D fiber retraction assay
PRP was diluted in PBS ($2.5x10^6$ platelets/ml, plasma concentrations were kept at 4 or 7
μl/ml depending on the amount of fibrin fibers to be formed; see explanation below) and
fibrinogen-Alexa 488 was added to a final concentration of 12.5 μg/ml. To induce fibrin fiber
polymerization and platelet activation, thrombin was added (final concentration of 2.5U/ml).
The suspension was gently mixed once (to prevent platelet aggregation) and after 15 min of
incubation at RT in the dark, 400 μl was transferred into wells of a 24 well plate containing
glass coverslips. The plate was centrifuged 3 min, 600 g, RT to allow synchronised contact of
all platelets with the glass surface. The plate was then placed in a cell culture incubator for
30 min to allow platelet spreading. Samples were then fixed with isotonic formalin for 15 min
at RT and then kept in PBS at 4°C until further use.

To promote fibrin fiber formation, while preventing the development of a clot, the plasma
concentration must be kept low (4 μl plasma per ml PBS). This can be achieved by adding a
volume of PRP containing $2.5x10^6$ platelets to 1 ml PBS, provided the platelet concentration
of PRP is at least $6.25x10^8$/ml (condition used for Fig. 3, 4, 5, 6, 8ABEF). If the platelet
concentration in the PRP is lower, platelets must be washed and the appropriate amount of
plasma added separately.

At higher plasma concentrations (7 μl plasma per ml PBS), more fibers polymerize leading to
the formation of a fragile clot which is aspirated during fixation/washing procedures. Under
these conditions very few fibers are left behind. However, a fibrin rosette can be observed in
the centre of several spread platelets (condition used for Fig. 7, 8CD).

The 2D fiber-retraction assay followed by immunofluorescence staining and expansion has
been repeated 10x with blood from different donors. A total of 18 image stacks stained for
fibrin/integrin, 20 stained for fibrin/integrin/actin and 62 stained for fibrin/myosin have been
acquired.
458
Immunofluorescence
Fixed platelets were permeabilized with PBS/0.2% Triton X-100 for 15 min at RT and then
incubated with blocking buffer (3% BSA and 10% goat serum in PBS) for 1h at RT. Coverslips
were then incubated for 2 h with a primary antibody diluted in blocking buffer, then washed
three times and incubated with a secondary antibody diluted in blocking buffer for 2h at RT.
After three washing steps, coverslips were either mounted in Mowiol or processed for
expansion.
466
Expansion microscopy
Expansion microscopy was essentially performed according to Gambarotto et al.[34] with
minor modifications as described previously[33]. Briefly, fixed, immunostained platelets or clots
on coverslips (12 mm diameter) were incubated in PBS/0.7% formaldehyde/1% acrylamide
over night at room temperature. The coverslip was then placed upside down onto an ice cold
35 µl drop of gelation solution (PBS/19% Na-acrylate/10% acrylamide/0.1% bisacrylamide)
pipetted onto parafilm on ice immediately after addition of TEMED and ammonium
persulfate both to a final concentration of 0.5%. After 5 min of incubation on ice, the samples
were transferred to 37°C for 1 h. The coverslip plus gel facing up was then incubated in
denaturation buffer (5.7% SDS/0.2M NaCl/50mM Tris, pH9) for 15 min at RT with gentle
agitation to detach the gel from the coverslip. The gel was then transferred into an
Eppendorf tube with fresh denaturation buffer and boiled for 30 min at 95°C. To expand the
gel, it was placed into a large volume of H_2O and the water was changed twice before
incubation ON at RT in fresh H_2O leading to 4x isometric expansion.
481
Image acquisition
For image acquisition of some of the expanded samples (Fig. 2 , 5CD, 6, 7, 8), a confocal
microscope ConfoBright (Nikon A1R+MP) equipped with a home-made adaptive optics
module was used with a 40x/1.15 long distance water immersion objective. The geometric
optical aberrations were corrected both in excitation and detection light paths in the open
loop mode with a deformable mirror (AlpAO DM97-15) inserted between the confocal
scanning head and the microscope inverted stand (Nikon Ti2E). Local aberrations were
sensed using the metrics of molecular brightness, derived from fluorescence fluctuations,
and were iteratively corrected as individual Zernike modes. This configuration allowed to
compensate for the refractive index mismatch, the specimen tilt and the eventual light
scattering in depth of expanded samples and to bring back the confocal resolution to its
diffraction limit in the vicinity of the imaged platelet.
Image acquisition of other expanded samples (Fig. 4, 5AB) was performed with a confocal
microscope Dynascope (LSM 710 ConfoCor 3; Zeiss), 63×/1.4 NA Plan Apochromat objective
and the acquisition software Zen 2010. The same microscope and software were used for live
imaging (Fig. 12) using the Apochromat 40×/1.2 NA water-immersion objective. Axial Z-stacks
(step 0.45 µm) were imaged in a time series using the avalanche photodiode detector in
photon counting mode at low 488 nm excitation power (0.1% transmission) to limit
photobleaching and phototoxicity, the 633 nm laser was set to 0.6% in order to obtain the
transmission light images of the platelets. The pinhole was closed to 0.25 Airy units (14 µm)

to increase the spatial resolution. The acquisition of transmitted light was concomitant to the
fluorescence.
The FIJI software was used for image processing and analysis.[35] No treatment that modifies
the raw images was applied. FIJI software was used for the Maximum Intensity Projection
(MIP) images, 3D reconstructions of image stacks and for the depth-color coded projection
image using the rainbow LUT (Fig. 4D) as well as for the depth-color coded images fire LUT
(Fig. 6) and the temporal-color coded image spectral LUT (Fig.12E).
509
Transmission electron microscopy (TEM)
Clots were formed around two holders as described previously.[32] Briefly, PRP was adjusted
with PBS to 10 % plasma, 1×10^8 platelets/ml and 4ml erythrocytes per ml were added for
color contrast. Clot formation was induced by addition of thrombin (final concentration of
2.5 U/ml). After 1h of retraction at 37°C, clots were fixed using 1,5 ml of 2,5 % glutaraldehyde
for 1h at RT. After fixation, a contrast-enhancing step was performed by incubating the
samples in a solution containing 1.5% potassium ferrocyanide and 1% osmium tetroxide in
0.1 M sodium cacodylate buffer. Following this, the samples were embedded in EPON resin
according to the protocol described by Eckly et al.[36] Thin sections (100 nm) were cut and
stained with uranyl acetate and lead citrate and examined using a Jeol 1200 Plus
transmission electron microscope operating at 120 kV.
521
Live video microscopy
- Unconstrained clot retraction (Fig. 1B):
Thrombin (final concentration of 2.5 U/ml) was added to 0.4 ml of a platelet suspension (50%
plasma/50% PBS; 1×10^8 platelets/ml; fibrinogen-Alexa 488 final concentration 12.5 µg/ml; 4
µl/ml erythrocytes for color contrast) to induce clot formation. After quick and gentle mixing,
the suspension was transferred immediately into a well (8-well Lab-Tek, coated 2% Pluronic®
to prevent clot attachment to well walls). Acquisition was started immediately at 100 µm
above the bottom of the well using the confocal microscope ConfoBright (for microscope
details see paragraph image acquisition).
- 2D fiber-retraction assay (Fig. 12A-D):
Platelets (washed Fig. 12AB or not Fig. 12C) were adjusted to 2.5×10^6 platelets/ml PBS and
2.5 µl plasma was added and fibrinogen-Alexa488 (final concentration of 12.5 µg/ml). 15
minutes after thrombin induced fibrin fiber formation and platelet activation, 1800 µl of the
suspension was transferred into a petri dish (WPI Fluoro Dish, FD35–100). The dish was
centrifuged 3 min, 600 g, RT and installed in the microscope incubator at 37°C (Fig. 12A) or
room temperature (Fig. 12BCD).
- 2D preformed fiber-retraction assay (Fig. 12E):
Preformed fluorescent fibrin fibers were prepared in the absence of platelets (2.5 µl
plasma/ml PBS; fibrinogen-Alexa 488, final concentration 12.5 µg/ml; thrombin, final
concentration 2.5U/ml; incubation 15 min RT in the dark). PRP was adjusted to 2.5×10^6
platelets/ml with PBS and 1800 µl was transferred into a petri dish (WPI Fluoro Dish, FD35–
100). The dish was centrifuged 3 min, 600 g at RT to allow spreading of the platelets. After
centrifugation the supernatant was replaced by the preformed fibrin fiber suspension and
the dish was installed in the microscope incubator at 37°C.
For all time-lapse videos, image acquisition was started immediately after adjusting the focal
position and image parameters (<1 min) using a confocal laser-scanning microscope
(ConfoBright or Dynascope, for microscope details see paragraph image acquisition).
549
550

The model of fibrin winding-up

The platelet bulb was modelled as a straight cylinder (the radius R = 400 nm, the length = 1500 nm). We assumed that the internal cytoskeleton of the bulb underwent the swirling motion with the angular velocity Rate_swirl as it was shown in[14] and simultaneously moved down to the bottom of the bulb with the velocity Rate. Integrins αIIbβ3 on the bulb surface were associated with internal cytoskeletal proteins,[37] thus, they underwent the swirling and downstream motion with the same rates.

The fibrin fiber was modelled as it was done previously[21,22] with several simplifications. The fiber composed of N equally spaced nodes connected by elastic springs (see supplementary Fig. 2A). The spring equilibrium length was L and it was equal to the initial distance between nodes. The fiber was located in the XY-plane and the angle α between fiber and x-axis was variable (Table 1). The motion of the node i in solution was described by the second Newton's law of motion. The system was assumed to be in an overdamped regime (mä = 0):

(1) $\eta \cdot \dot{r}_i = F_{i,i-1} + F_{i,i+1} + F_i^B$

where r_i was the position vector of the node i in three-dimensional coordinates (x,y,z); η was the drag coefficient for the fiber node in the blood serum; $F_{i,i-1} = K_r \cdot (\Delta r_{i,i-1} - L)$ and $F_{i,i+1} = K_r \cdot (\Delta r_{i,i+1} - L)$ were the vector forces which acted on the node i from the node i-1 or i+1 according to the Hook's law due to the spring elongation or shortening ($\Delta r_{i,i-1}$ and $\Delta r_{i,i+1}$ was the distance between nodes i and i-1 or the nodes i and i+1; Kr was the fiber spring constant); F_i^B was the Brownian force which acted on the node i due to thermal fluctuations. To solve the equations for the nodes in solution, the system of equations was discretized and the Heun's method was used. According to this method, the evolution of coordinates in time for the differential equation

(2) $\dot{x} = f(x, t)$

was calculated using a two-step algorithm:

(3) $\tilde{x}^{n+1} = x^n + \Delta t \cdot f(x^n, t^n)$

(4) $x^{n+1} = x^n + \frac{\Delta t}{2} \cdot \{f(x^n, t^n) + f(\tilde{x}^{n+1}, t^{n+1})\}$

where Δt was the time step used in calculation and discretization scheme; tn+1 = tn + Δt; xn was coordinate at the tn moment of time.

Thus, the evolution of the node i coordinates in time was calculated according to a two-step process (in a vector form):

(5) $\tilde{r}_i^{n+1} = r_i^n + \frac{\Delta t}{\eta} \cdot (F_{i,i-1}(r_i^n, r_{i-1}^n) + F_{i,i+1}(r_i^n, r_{i+1}^n) + F_i^{n,B})$

(6) $\quad r_i^{n+1} = r_i^n + \frac{\Delta t}{2\eta} \cdot (F_{i,i-1}(r_i^n, r_{i-1}^n) + F_{i,i+1}(r_i^n, r_{i+1}^n) + F_i^{n,B} + F_{i,i-1}(\tilde{r}_i^{n+1}, \tilde{r}_{i-1}^{n+1}) + F_{i,i+1}(\tilde{r}_i^{n+1}, \tilde{r}_{i+1}^{n+1}) + \tilde{F}_i^{n+1,B})$

The Brownian force in three dimensions was calculated as described previously.[21] Briefly, the mean square displacement of a particle over a single time step was $\langle x^2 \rangle = \frac{6k_bT}{\eta} dt$, and the

discretized Brownian force acting on the node i was $F_i^{n,B} = \varepsilon_i \cdot \sqrt{\frac{6k_bT}{\eta}} dt$, where εi was

sampled from the standard normal distribution.

At the initial time point, the node 0 was bound to the integrin receptor on the bulb surface (top view, supplementary Fig. 2B). The distance of the binding point from the bulb top was D. The height of the binding point from the bulb surface was equal to the integrin αIIbβ3 molecule length (H = 20 nm) in its elongated conformation.[38] The equations for the node 0 swirling and downstream motion were:

(7) $x_0^{n+1} = (R + H) \cdot \cos(Rate\_swirl \cdot \Delta t + \varphi^n)$

(8) $y_0^{n+1} = (R + H) \cdot \sin(Rate\_swirl \cdot \Delta t + \varphi^n)$

(9) $\varphi^{n+1} = Rate\_swirl \cdot \Delta t + \varphi^n$

(10) $z_0^{n+1} = z_0^n - Rate \cdot \Delta t$

When the node i entered the light-magenta zone (supplementary Fig. 2B), and its distance
from the bulb surface became smaller than H, it bound the integrin molecule and its
equations of motion became similar to the node 0 equations (7) – (10).

The integrin αIIbβ3 surface density on the bulb surface was relatively high (the distance
between the neighbouring molecules was approximately 15 nm). Given such a high density,
we assumed that when the node became close enough to the bulb, it immediately
encountered the integrin molecule and bound it. Thus, when the distance between the node
i and the bulb surface became smaller than H, node i bound to the integrin molecule and the
node's equations of motion became similar to the node 0 equations (7) – (10). The binding
was assumed to be irreversible.

The model parameters were listed in the Table 1 below. The model equations were solved
using Python 3.11.

A

B

611
612

Supplementary Figure 2: Model of fibrin fiber binding to the platelet bulb and fibrin
winding-up.

A) The model of a fibrin fiber in solution. The fiber was modeled as a sequence of N nodes
(black dots) which were connected by elastic springs (blue). $F_{i,i-1}$/ $F_{i,i+1}$ were vector forces
which acted on the node i according to the Hook's law, the black vectors show the force
directions.

B) The system configuration at time point T=0. Magenta circle: internal volume of a bulb.
Light-magenta: the binding region. If the node i entered the binding region, it bound to the
integrin molecule. The black arrow shows the rotation direction of the cytoskeleton and
integrin molecules.

Parameter	Value	Citation
R, the bulb radius	400 nm	CS
Bulb length	1500 nm	CS
H, the height of the binding sites on integrins for fiber nodes from the bulb surface	20 nm	[38]
L, initial distance between modeled nodes in fibrin fiber	2×45 nm (2 fibrin molecules)	[39]
α, angle between fiber and X-axis in 3D coordinates	π/4 or π/2.5	CS
Number of nodes	60	CS
D, distance from the top of the bulb to the point where the node 0 was bound at T = 0	500 nm	CS
Δt, time step	0.00015 μs	CS
Rate, rate of downstream movement of integrins	1 nm/μs	CS
Rate of swirling	π/500 rad/μs	CS
Temperature, T	300 K	CS
Drug coefficient for the node, η	3.77×10^{-3} nN*μs/nm	[21]
Stiffness of fibrin fiber, Kr	11.5 nN/nm	[22]

626 *Table 1. The model parameters used in the simulation. CS – current study.*

Acknowledgement

We are grateful to Jacques Mazzega of the microscopic platform and to Arnold Fertin for image analyses. We gratefully acknowledge the MicroCell facility (GIS IBiSA, ISdV, IAB) for its technical assistance and the microscopy system. The MicroCell facility is a member of the national infrastructure France-BioImaging (https://france-bioimaging.org/). We thank Rémy Sadoul and Alexander Bershadsky for critical reading of the manuscript. This work was supported by the Fondation Recherche Médicale (FRM, grant number DEI20151234416) and the University Grenoble Alpes (UGA, grant number AGIR-POLE FRAG15CS08).

Declaration of Interest statement

The authors declare that they have no competing financial interests.

Authorship contributions

A.G. developed the microscope with adaptive optics corrections and acquired images T.K. and M.P. established the model and simulations. A.-S.R. performed experiments, F.A. helped with image analyses and rendering. A.E. and J.-Y.R. acquired TEM images. L.L. participated in manuscript writing. K.S. initiated and planned the study, performed experiments, interpreted the results and wrote the manuscript. All authors read the manuscript and agreed to its content.

Data sharing statement

All raw data are stored on the server of our microscope platform and can be accessed upon request.

References

1. Litvinov RI, Weisel JW. Fibrin mechanical properties and their structural origins. *Matrix Biol*. 2017;60-61:110-123.
2. Tutwiler V, Maksudov F, Litvinov RI, Weisel JW, Barsegov V. Strength and deformability of fibrin clots: Biomechanics, thermodynamics, and mechanisms of rupture. *Acta Biomater*. 2021;131:355-369.
3. Adair BD, Alonso JL, van Agthoven J, et al. Structure-guided design of pure orthosteric inhibitors of alphaIIbbeta3 that prevent thrombosis but preserve hemostasis. *Nat Commun*. 2020;11(1):398.
4. Litvinov RI, Weisel JW. Blood clot contraction: Mechanisms, pathophysiology, and disease. *Res Pract Thromb Haemost*. 2023;7(1):100023.
5. Peshkova AD, Weisel JW, Litvinov RI. A novel technique to quantify the kinetics of blood clot contraction based on the expulsion of fluorescently labeled albumin into serum. *J Thromb Haemost*. 2024;22(6):1742-1748.
6. Tucker KL, Sage T, Gibbins JM. Clot Retraction. In: Gibbins J., Mahaut-Smith M. (eds) Platelets and Megakaryocytes. Methods in Molecular Biology (Methods and Protocols). 2012;788.
7. Sun Y, Oshinowo O, Myers DR, Lam WA, Alexeev A. Resolving the missing link between single platelet force and clot contractile force. *iScience*. 2022;25(1):103690.
8. Peshkova AD, Rednikova EK, Khismatullin RR, et al. Red blood cell aggregation within a blood clot causes platelet-independent clot shrinkage. *Blood Adv*. 2025.
9. Lam WA, Chaudhuri O, Crow A, et al. Mechanics and contraction dynamics of single platelets and implications for clot stiffening. *Nat Mater*. 2011;10(1):61-66.
10. Kim OV, Litvinov RI, Alber MS, Weisel JW. Quantitative structural mechanobiology of platelet-driven blood clot contraction. *Nat Commun*. 2017;8(1):1274.
11. Leistikow EA, Barnhart MI, Escolar G, White JG. Receptor–ligand complexes are cleared to the open canalicular system of surface-activated platelets. *British Journal of Haematology*. 1990;74(1):93-100.
12. Poulter NS, Pollitt AY, Davies A, et al. Platelet actin nodules are podosome-like structures dependent on Wiskott-Aldrich syndrome protein and ARP2/3 complex. *Nat Commun*. 2015;6:7254.
13. Joubert C, Grichine A, Dolega M, et al. Spatial and temporal characterization of cytoskeletal reorganizations in adherent platelets. *Platelets*. 2024;35(1):2422437.
14. Tee YH, Shemesh T, Thiagarajan V, et al. Cellular chirality arising from the self-organization of the actin cytoskeleton. *Nat Cell Biol*. 2015;17(4):445-457.
15. Tamada A, Igarashi M. Revealing chiral cell motility by 3D Riesz transform-differential interference contrast microscopy and computational kinematic analysis. *Nat Commun*. 2017;8(1):2194.
16. Tamada A, Kawase S, Murakami F, Kamiguchi H. Autonomous right-screw rotation of growth cone filopodia drives neurite turning. *J Cell Biol*. 2010;188(3):429-441.
17. Estévez-Torres A, Baigl D. DNA compaction: fundamentals and applications. *Soft Matter*. 2011;7:6746.
18. Brzoska T, Suzuki Y, Mogami H, Sano H, Urano T. Binding of thrombin-activated platelets to a fibrin scaffold through $\alpha(IIb)\beta_3$ evokes phosphatidylserine exposure on their cell surface. *PloS one*. 2013;8(2):e55466-e55466.
19. Tutwiler V, Litvinov RI, Lozhkin AP, et al. Kinetics and mechanics of clot contraction are governed by the molecular and cellular composition of the blood. *Blood*. 2016;127(1):149-159.

20. Leon C, Eckly A, Hechler B, et al. Megakaryocyte-restricted MYH9 inactivation dramatically affects hemostasis while preserving platelet aggregation and secretion. *Blood*. 2007;110(9):3183-3191.

21. Michael C, Pancaldi F, Britton S, et al. Combined computational modeling and experimental study of the biomechanical mechanisms of platelet-driven contraction of fibrin clots. *Commun Biol*. 2023;6(1):869.

22. Kliuchnikov E, Peshkova AD, Vo MQ, et al. Exploring effects of platelet contractility on the kinetics, thermodynamics, and mechanisms of fibrin clot contraction. *NPJ Biol Phys Mech*. 2025;2(1):6.

23. Hagmann J. Pattern formation and handedness in the cytoskeleton of human platelets. *Proc Natl Acad Sci U S A*. 1993;90(8):3280-3283.

24. Morgenstern E, Ruf A, Patscheke H. Ultrastructure of the interaction between human platelets and polymerizing fibrin within the first minutes of clot formation. *Blood Coagul Fibrinolysis*. 1990;1(4-5):543-546.

25. White JG, Krivit W, Vernier RL. The Platelet-Fibrin Relationship in Human Blood Clots: An Ultrastructural Study Utilizing Ferritin-Conjugated Anti-Human Fibrinogen Antibody. *Blood*. 1965;25:241-257.

26. Cohen I, Gerrard JM, White JG. Ultrastructure of clots during isometric contraction. *J Cell Biol*. 1982;93(3):775-787.

27. Morgenstern E, Daub M, Dierichs R. A new model for in vitro clot formation that considers the mode of the fibrin(ogen) contacts to platelets and the arrangement of the platelet cytoskeleton. *Ann N Y Acad Sci*. 2001;936:449-455.

28. Kim OV, Litvinov RI, Gagne AL, French DL, Brass LF, Weisel JW. Megakaryocyte-induced contraction of plasma clots: cellular mechanisms and structural mechanobiology. *Blood*. 2024;143(6):548-560.

29. Williams EK, Oshinowo O, Ravindran A, Lam WA, Myers DR. Feeling the Force: Measurements of Platelet Contraction and Their Diagnostic Implications. *Semin Thromb Hemost*. 2019;45(3):285-296.

30. Myers DR, Qiu Y, Fay ME, et al. Single-platelet nanomechanics measured by high-throughput cytometry. *Nat Mater*. 2017;16(2):230-235.

31. Mercey O, Reymond L, Lemaître F, et al. HAK-actin, U-ExM-compatible probe to image the actin cytoskeleton. *bioRxiv*. 2025:2025.2008.2026.672318.

32. Kovalenko TA, Giraud MN, Eckly A, et al. Asymmetrical Forces Dictate the Distribution and Morphology of Platelets in Blood Clots. *Cells*. 2021;10(3).

33. Grichine A, Jacob S, Eckly A, et al. The fate of mitochondria during platelet activation. *Blood Adv*. 2023;7(20):6290-6302.

34. Gambarotto D, Hamel V, Guichard P. Ultrastructure expansion microscopy (U-ExM). *Methods Cell Biol*. 2021;161:57-81.

35. Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for biological-image analysis. *Nat Methods*. 2012;9(7):676-682.

36. Eckly A, Strassel C, Cazenave JP, Lanza F, Leon C, Gachet C. Characterization of megakaryocyte development in the native bone marrow environment. *Methods Mol Biol*. 2012;788:175-192.

37. Bidone TC, Skeeters AV, Oakes PW, Voth GA. Multiscale model of integrin adhesion assembly. *PLoS Comput Biol*. 2019;15(6):e1007077.

38. Xu XP, Kim E, Swift M, Smith JW, Volkmann N, Hanein D. Three-Dimensional Structures of Full-Length, Membrane-Embedded Human alpha(IIb)beta(3) Integrin Complexes. *Biophys J*. 2016;110(4):798-809.

751   39.   Weisel JW, Litvinov RI. Fibrin Formation, Structure and Properties. *Sub-cellular*
752   *biochemistry*. 2017;82:405-456.
753
754
755

Figure 1

Figure 1: Platelets become labelled by fluorescent fibrin fibers during clot retraction

A) Time course of unconstrained clot retraction by platelets (4×10^7 in 400 μl of 12% plasma in PBS) for

a total of 60 minutes at room temperature. Experiments have been performed 6x using blood from

different donors and typical examples of blood from two different donors are shown (experiments III

and IV are repetitions of I and II the day after; although a longer lag-phase is observed, the final

retraction volume is similar for the four experiments).

B) Unconstrained clot retraction (1×10^8 platelets per ml in 50% plasma/50% PBS) in presence of

fibrinogen-Alexa 488. Image acquisition was started immediately after thrombin addition at a focal

plane 100 μm above the bottom of the well and image stacks were collected (61 focal planes, step size
0.5 μm) for a time period of 20:27 min (100 frames with a time interval of 12.5 sec). See also
associated video 1. The experiment has been performed twice using platelets of the same donor.
Shown are the first, intermediate and last time points of a depth color-coded time-lapse video (left
panel; scale bar 10 μm) and the maximal intensity projections (MIPs) of the same time points (right
panel; scale bar 10 μm).
C) Constrained clot retraction (1x10^8 platelets per ml in 50% plasma/50% PBS, fibrinogen-Alexa 488
final concentration 12.5 μg/ml) between two holders. Clots were induced by addition of thrombin
(2.5U/ml final), fixed at the indicated retraction times, embedded in gelatin, flash frozen and
cryosections (14 μm) were stained for the integrin subunit αIIb (magenta; scale bar 5 μm). The time
course was performed twice using PRP from two different donors (retraction assays were repeated,
although not for all time points, more than eight times using blood from different donors, with
consistent results). Image acquisition was performed using a wide-field epi fluorescence microscope
(BX41; Olympus) equipped with a Plan 100x/1.25 NA oil objective, a camera (DP70; Olympus), and the
acquisition software analySIS (Olympus).
783
784

 Figure 2

Figure 2: Fibrin fibers are organized in a "cage-like" fashion around platelets within a constrained
clot
A-D: Two typical examples (out of 39 acquisitions of four experiments using blood from different
donors) of platelets in a constrained clot with attached fibrin fibers are shown ($1x10^7$ platelets per ml
PBS/50% plasma and fibrinogen-Alexa 488). Clot retraction was allowed to take place in an
inoculation loop (see methods) for 15 min before fixation and immunofluorescence staining of the
integrin subunit αIIb (magenta). Samples were then processed for expansion (scale bars 10 μm = 2.5
μm after correction for expansion; indicated z-levels are not corrected for expansion; see also
associated animation, video 2).
A) Four focal planes from a stack of images (z-level as indicated) showing a platelet in a clot (fibrin
fibers in green, plasma membrane staining using an antibody against the integrin subunit αIIb in
magenta and the merge).
B) 3D reconstruction of the image stack used in A (46 planes, step size 0.5 μm), a bulb is indicated by
an arrow.
C) Four focal planes show another typical platelet in a clot (z-level as indicated).
D) 3D image reconstruction of the image stack used in C (28 planes, step size 1 μm), a bulb is
indicated by an arrow.
804

Figure 3

Figure 3: Spread platelets accumulate fibrin fibers above them in a myosin dependent way

A) Shown is a 2D fiber-retraction assay in the presence of DMSO (control) or 50 μM blebbistatin for
myosin inhibition. After a 30 min preincubation of platelets (4 μl plasma per ml PBS, see methods
section) with or without blebbistatin, thrombin is added and the assay is continued as described in the
methods section. Scale bar 10 μm, arrowheads indicate platelets without fiber contacts, thin arrows
with fiber initiation or contact, larger arrows indicate platelets starting to wind-up fibers and even
larger arrows point to platelets with compacted fibers around them (see legend below bar graph in
B).
B) Quantification of the percentage of platelets in the different categories using image acquisitions of
four experiments using blood from different donors (for control conditions a total of 636 platelets
were counted and for blebbistatin conditions a total of 524). Shown are means ± sem, evaluation of
significance using the Student's t-test.

Figure 4

Figure 4: Coiled-up fibrin fibers above spread platelets
Two representative examples (out of 18 acquisitions from three independent experiments using blood
from different donors) of fibrin fiber accumulations above spread platelets in the 2D fiber-retraction
assay (4 μl plasma per ml PBS, see methods section) are shown (see also animation, video 3). Samples
were stained for the αIIb integrin subunit and expanded (scale bars 10 μm = 2.5 μm after correction
for expansion, indicated z-levels are not corrected for expansion).
A) Three different focal planes (z-levels as indicated) from an image stack showing two spread
platelets and attached fibrin fibers.
B) 3D reconstruction of the image stack used in A (51 planes, step size 0.25 μm), shown are three
different view angles.
C) Three different focal planes from an image stack illustrating a spread platelet and attached, coiled-
up fibrin fibers.

D) 3D reconstruction of the image stack used in C (50 planes, step size 0.25 μm), shown are three different view angles. The fibrin fibers are additionally displayed as depth-color coded (Fire LUT) reconstructions to highlight the straight fiber parts along the z-axis.

E) Same fiber image stacks as in C and D. Left panels are MIPs of the green fibrin fibers along the z, x and y axes. Right panels are the same projections color coded in the HSB space (Hue, Saturation, Brightness) according to the orientation of individual fiber segments. The fibers are well aligned along the z-axis, while they crisscross horizontally.

Figure 5

Figure 5: Strongly compacted fibrin fibers around the center of platelets in the 2D fiber-retraction
assay

Two representative examples of platelets are shown with bulbs encircled by fibrin fibers in the 2D
fiber-retraction assay (4 μl plasma/ml, see methods section) behaving similar to platelets in a clot
(see also animation, video 4). Samples were stained for the αIIb integrin subunit (A, B) or the myosin
light chain (MLC; C,D) and processed for expansion (scale bars 10 μm = 2.5 μm after correction for
expansion, indicated z-levels are not corrected for expansion). Experiment was repeated 4x with blood
from different donors with a total of 18 image stacks acquired for fibrin/integrin and 62 for
fibrin/myosin staining.

A) Three different focal planes from an image stack showing two platelets, one spread and one with
bulbs and attached, compacted fibrin fibers.

B) 3D image reconstruction of the image stack used in A (34 planes, step size 0.25 μm), shown are
three different view angles.

C) Three different focal planes from an image stack illustrating another platelet with bulbs and
attached, rolled-up, compacted fibrin fibers.

D) 3D image reconstruction of the image stack used in C (55 planes, step size 0.25 μm), shown are
three different view angles.

     Figure 6

Figure 6: Fibrin fibers are twisted in the proximity of platelets

Two representative examples of platelets with compacted fibrin fibers are shown in the 2D fiber-

retraction assay (4 µl plasma/ml, see also animation, video 5). Samples were stained for myosin II (A,

B) or phosphorylated myosin light chain (p-MLC; C-E) and processed for expansion (scale bars 10 µm =

2.5 μm after correction for expansion, indicated z-levels are not corrected for expansion). Please note
also the strong fiber accumulation around the platelets similar to platelets shown in figure 5.
A) 3D image reconstruction of an image stack (72 planes, step size 0.25 μm) illustrating a platelet
with bulbs and wound-up compacted fibrin fibers.
B) Three different focal planes (z-level as indicated) from the image stack used in A, depth color-
coded. Rectangles indicate twisted fibrin fibers and a corresponding zoom is shown below.
C) Three different focal planes (z-level as indicated) from an image stack illustrating two platelets with
bulbs and attached, wound-up, compacted fibrin fibers. Twisted thick fiber bundles are observed
between the two platelets.
D) 3D image reconstruction of the image stack (91 planes, step size 0.25 μm) used in C, shown are
three different view angles.
E) The same focal planes as in C and the top view of the 3D image reconstruction shown in D of the
fibrin fibers, depth color-coded to better distinguish the twist of the thick fiber bundles between the
two platelets.
884

Figure 7

Figure 7: A fibrin rosette is located close to a ring-like actin organization in spread platelets

Four representative examples (out of 20 acquisitions, experiment repeated 4x using blood from

different donors) of a fibrin rosette associated with spread platelets in the 2D fiber-retraction assay (7

μl plasma per ml PBS, see methods section) are shown (see also animation, video 6). Samples were

stained for the αIIb integrin subunit as well as for actin and processed for expansion (scale bars 10 μm

= 2.5 μm after correction for expansion).

A) 3D image reconstruction of an image stack (30 planes, step size 0.33 μm) of fibrin fibers (green)

and integrin or actin staining (upper and lower panels respectively, magenta).

B) Similar example as in A (17 planes, step size 0.33 μm).

C) Another example of a fibrin rosette with intercalated actin nodules (34 planes, step size 0.33 μm).

D) An example showing a long fibrin fiber associated with the fibrin rosette (35 planes, step size 0.38

μm).

900
901
902
903

Figure 8

Figure 8: The fibrin rosette and attached fibers are in close proximity to myosin II in spread platelets

Three representative examples (out of 62 acquisitions, experiment repeated 4x using blood from different donors) of fibrin and myosin localization in spread platelets (see also animation, video 7) in the 2D fiber-retraction assay (4 μl plasma/ml for A, B, E, F or 7 μl plasma/ml C, D; see methods section). Samples were stained for myosin (A, B, C, D) or phosphorylated myosin light chain (p-MLC; E, F) and expanded (scale bars 10 μm = 2.5 μm after correction for expansion, indicated z-levels are not corrected for expansion).

A) Three different focal planes (z-levels as indicated) from an image stack showing a spread platelet and an attached fibrin fiber in the center of a faint fibrin rosette.

B) 3D image reconstruction of the image stack used in A (36 planes, step size 0.3 μm), shown are three different view angles.

C) Three different focal planes from an image stack illustrating a spread platelet with a fibrin rosette and an attached, rolled-up fibrin fiber.

D) 3D image reconstruction of the image stack used in C (31 planes, step size 0.33 μm), shown are three different view angles.

E) Three different focal planes from an image stack illustrating a spread platelet with attached, rolled-up fibrin fibers.

F) 3D image reconstruction of the image stack used in E (54 planes, step size 0.25 μm), shown are three different view angles.

Figure 9

924
925
926
Figure 9: Determination of platelet protrusions and fibrin fiber curvatures in proximity of platelets
A) Three examples of platelets within a clot stained for the αIIb integrin subunit (including platelets
shown in figure 2). MIPs are shown, the number of platelet extensions is indicated and has been
determined manually by scrolling through the image z-stack. Scale bars 10 μm = 2.5 μm after
correction for expansion.
B) Three examples of platelets attached to a glass surface (including platelets shown in figure 5) and
stained for the MLC (upper panel) or the αIIb integrin subunit (middle and lower panel). MIPs are
shown, the number of platelet extensions is indicated and has been determined manually by scrolling
through the image z-stack. Scale bars 10 μm = 2.5 μm after correction for expansion.
C) The comparison of all data points between platelets in clots or attached to a glass surface does not
show a statistically significant difference between protrusions of platelets in a clot or when spread on
a glass surface (Mann-Whitney U test).

D-F: The degree of fiber curvature in examples shown in D and E is determined as $\kappa=1/r$ using the
radius of a circle fitting the region of strongest curvature of the fibers.
D) Three examples of TEM images used to determine the curvature of fibrin fibers close to platelets
within a clot (measured examples are indicated by arrows).
E) Three examples of fluorescent images of fibers wound around platelets attached to a glass surface
(including platelets shown in figure 5). A single focal plane of each image stack is shown (scale bars 10
$\mu m = 2.5\ \mu m$ after correction for expansion).
F) Comparison of all data points obtained using TEM images and fluorescent images (curvature in κ
μm^{-1}). For quantification, 31 EM acquisitions of three clots from different donors were used as well as
12 Fluo acquisitions of platelets from 4 different donors.
949

950 Figure 10
951

Figure 10: Simulation of platelet mediated winding-up of fibrin fibers
A) Time evolution of the fibrin fiber configuration around the platelet bulb. Two different angles
between the fiber and the x-axis were used ([1]: α = π/4, lime color; [2]: α = π/2.5, dark green color).
Left: initial positions, top view and side view. Center: intermediate position, top view and side view.
The fibrin fiber began to form a loop around the bulb. Right: final position, top view and side view. A
compact fibrin loop was formed around the bottom of the bulb (see also animation, video 8).
B) Time when the first node bound to the bulb and thus the formation of the fibrin loop started. The
fibrin winding-up started earlier for a larger angle between fiber and x-axis (and consequently smaller
angle between fiber and tangent line at the point of the node 0 binding). The number of nodes that
were bound to the bulb at the end of the simulation was slightly higher for larger α. Thus, larger α led
to a more compact fibrin fiber organization.
C) The distance between nodes and the bottom of bulb along z-axis. The distance was smaller for
larger α and the fibrin organization was slightly more compact.
D) Comparison of final fiber configurations around the bulb for two different α.
966

Figure 11

Figure 11: Features predicted by the model/simulation are observed on TEM images

Interpreted, potential fiber organisations are highlighted in green on an identical image on the right.

A) Fibrin fibers appear to encircle a bulb.

B) Cross sectioned fibrin fibers at the base of a bulb, which potentially curve around the bulb.

C) Several cross sectioned fibers at the base of a bulb, which may surround the bulb.

D) Several cross sectioned fibers along a platelet extension potentially forming a spiral.

E) Cross sectioned fibrin fibers around a forming bulb.

F) Cross sectioned fibers at the base of a bulb and two fibers along a filopodial extension.

Figure 12

Figure 12: Live imaging of spread platelets winding-up fluorescent fibrin fibers

A-C: Shown are three experiments using blood from different donors.

A) Washed platelets were adjusted to 5x10⁶ platelets per 2 ml PBS and 5 µl plasma was added as well
as fibrinogen-Alexa 488. Thrombin was added to induce fibrin fiber formation and platelet activation.
After 10 min, the suspension was transferred into a petri dish (WPI Fluoro), centrifuged and installed
in the microscope incubator at 37°C to start image acquisition (see also video 9, upper panel). The first
and last time points of the time-lapse video are shown; acquired using the fluorescence and

transmission channels. Projections of four focal fluorescent image planes of fibrin fibers (green) and projections of the transmission planes as well as the merged channels are shown. An arrow indicates a forming kink in a fibrin fiber (scale bar 10 μm). Below are shown Individual time points of the time-lapse video zooming on the rectangle indicated in the panels above. This region shows rotational fiber movements.

B) Independent experiment under conditions as described in A except that the petri dish was kept at room temperature to slow down the platelet mediated fiber reorganizations (scale bar 10 μm). An arrow shows platelet mediated fiber compaction. Below are shown Individual time points of the time-lapse video zooming on the rectangle indicated in the panels above. This region shows a fibrin fiber getting curved by the platelet and finally ruptures followed by a rotational movement of the ruptured fiber.

C) Platelets of a third donor (not washed, i.e., 5 μl PRP with $4x10^6$ platelets per 2 ml PBS) and fibrinogen-Alexa 488 were added to 2 ml of PBS and the experiment was continued as in B (scale bar 10 μm). Below are shown Individual time points of the time-lapse video zooming on the rectangle indicated in the panels above. This region shows fibrin fibers getting coiled around the pseudo-nucleus of the spread platelet. The last zoomed image shows a temporal color-coded projection of the transmission channel to illustrate the rotational movement of the pseudo-nucleus (scale bar 5 μm).

D) Samples shown in B and C were fixed 3h after the start of time-lapse acquisitions and stained for the integrin subunit aIIb. Shown are fibrin fibers (green) integrin staining (magenta) and the merge (samples B and C, upper and lower panels, respectively; scale bar 10 μm).

E) Preformed fluorescent fibrin fibers were prepared in the absence of platelets. PRP was adjusted to $2.5x10^6$ platelets per ml with PBS and 1800 μl was transferred into a petri dish (WPI Fluoro). The dish was centrifuged to allow spreading of the platelets and the supernatant was replaced by the preformed fibrin fibers. The dish was installed in the microscope incubator to start image acquisition (see also video 9, last panel). The experiment was performed twice with blood of the same donor. The last fluorescent image of the time-lapse video is shown (left image, scale bar 10 μm). Fifteen of the fibrin accumulations in the middle of the platelets are rotating in a counterclockwise direction (indicated as "1"), four rotate clockwise ("2") and for 16 no clearly visible turn is observed ("3"). A projection of all fluorescent time points, temporal color-coded using the LUT spectral, is shown. White corresponds to the sum of all the colors at each time point, meaning no movement during the time-lapse video. (right image, scale bar 10 μm).

Supplemental Videos (Time-lapse and Animations)

Video 1 (associated to figure 1B)

Depth color-coded time-lapse of an unconstrained clot retraction (1x10^8 platelets per ml in 50% plasma/50% PBS) in presence of fibrinogen-Alexa 488. Fibrin fibers (green) present in z-projections of each time point are also shown upper right corner. Acquisition was started immediately after thrombin addition at 100 µm above the bottom of the well and image stacks were acquired (61 focal planes, step size 0.5 µm) for a time period of 20:27 min (100 frames with a time interval of 12.5 seconds, scale bar 10 µm).

Video 2 (associated to figure 2)

Two typical examples of platelets in a constrained clot with attached fibrin fibers (green) are shown (1x10^7 platelets per ml PBS/50% plasma and fibrinogen-Alexa 488). Clot retraction was allowed to take place for 15 min before fixation and immunofluorescence staining of the integrin subunit αIIb (magenta). Samples were then processed for expansion (scale bars 10 µm = 2.5 µm after correction for expansion). Left panels, scrolling through the image stack; right panels, 3D reconstruction.

Video 3 (associated to figure 4)

Two typical examples of fiber accumulations (green) above spread platelets in the 2D fiber-retraction assay (4 µl plasma per ml PBS, see methods section) are shown. Samples were stained for the αIIb integrin subunit (magenta) and expanded (scale bars 10 µm = 2.5 µm after correction for expansion). Upper panels, scrolling through the image stack; lower panels, 3D reconstruction.

Video 4 (associated to figure 5)

Two examples of platelets with bulbs encircled by fibrin fibers (green) in the 2D fiber-retraction assay (4 µl plasma/ml, see methods section) are shown. Samples were stained for the αIIb integrin subunit (A, B; magenta) or the myosin light chain (MLC; C,D; magenta) and processed for expansion (scale bars 10 µm = 2.5 µm after correction for expansion). Upper panels, scrolling through the image stack; lower panels, 3D reconstruction.

Video 5 (associated to figure 6)

Two examples of platelets with twisted fibrin fibers (green) are shown in the 2D fiber-retraction assay (4 µl plasma/ml, see methods section). Samples were stained for myosin II (A; magenta) or phosphorylated myosin light chain (p-MLC; C,D; magenta) and processed for expansion (scale bars 10 µm = 2.5 µm after correction for expansion). Upper panel 3D reconstruction (Fig. 6A). Middle panel, scrolling through the image stack (Fig. 6C); lower panel, 3D reconstruction (Fig. 6D).
Please note also the strong fiber accumulation around the platelets similar to platelets shown in video 4/figure 5.

Video 6 (associated to figure 7)

Four examples of a fibrin rosette associated with spread platelets in the 2D fiber-retraction assay (7 µl plasma per ml PBS, see methods section) are shown. Samples were stained for the αIIb integrin subunit as well as for actin and processed for expansion (scale bars 10 µm = 2.5 µm after correction for expansion). 3D image reconstructions of image stacks. Fibrin fibers (green) and integrin or actin staining (upper and lower panels respectively, magenta).

Video 7 (associated to figure 8)

Three examples of fibrin (green) and myosin (magenta) localization for spread platelets in the 2D fiber-retraction assay (4 μl plasma/ml for A, B, E, F or 7 μl plasma/ml C, D; see methods section). Samples were stained for myosin (A, B, C, D) or phosphorylated myosin light chain (p-MLC; E, F) and expanded (scale bars 10 μm = 2.5 μm after correction for expansion). Upper panels, scrolling through the image stack; lower panels, 3D reconstruction.

Video 8 (associated to figure 10)

Simulation of fiber winding-up around a platelet bulb. Angle between the fiber and the x-axis $\alpha = \pi/2.5$

Video 9 (associated to figure 12)

Live imaging of spread platelets (transmission gray) and fluorescent fibrin fibers (green), scale bars 10 μm.

Upper three panels : Spread platelets from three different donors are shown, which reorganize and compact fluorescent fibrin fibers.

Lower panel : Preformed fluorescent fibrin fibers were added to spread platelets. Surface coating is observed and a dense fibrin mass in the platelet center is rotating in several platelets.