1
**QuantCell: machine learning based cell annotation from qualitative and quantitative imaging profiles**

Wade Boohar[1], Bowen Wang[1], Zachary Thomas[1, 2], Anna Nogalska[1], Rong Lu[1, 2, 3, 4*]

[1]Department of Stem Cell Biology and Regenerative Medicine, Eli and Edythe Broad Center for
Regenerative Medicine and Stem Cell Research, Keck School of Medicine, [2]Department of
Biomedical Engineering, [3]Department of Medicine, [4]Leonard Davis School of Gerontology,
University of Southern California, Los Angeles, CA 90033, USA

* **Correspondence:**

Rong Lu

Keck School of Medicine, University of Southern California

1425 San Pablo Street, BCC 306, Los Angeles, CA 90033

Office: (323) 442 0169

Email: ronglu@usc.edu

**Abstract**

Recent advances in spatial omics enable high-resolution, multiplexed *in situ* imaging of gene and protein expression. A major challenge in analyzing these data is cell annotation, especially in complex tissues with limited molecular markers, overlapping marker expression, or rare cell types such as stem cells. Here, we present QuantCell, a machine learning (ML) based framework that leverages quantitative imaging data to improve cell annotation derived from qualitative data. QuantCell evaluates multiple ML models to identify the best-performing approach and applies a user-defined false discovery rate (FDR) to ensure reliable annotation. Using PhenoCycler imaging of mouse bone marrow, we show that QuantCell increases the number of annotated cells from 33.1% to 90.2% at a 5% FDR, achieving 96.5% accuracy based on the test dataset from conventional annotation. Benchmarking results demonstrate that QuantCell outperforms existing annotation methods in predictive accuracy. It also minimizes performance bias toward abundant cell types, enabling robust detection of rare cell populations. Unlike many existing methods, QuantCell does not require manual cell annotation, which can be labor-intensive for multi-marker cell types and unfeasible for rare cell populations. In summary, QuantCell provides an effective ML solution for spatial omics analysis, with a workflow that is broadly compatible with various imaging platforms and adaptable to project-specific needs.

## Introduction

Recent advances in spatial omics technologies have revolutionized our ability to investigate
cellular organization and function within native tissue environments[1–4]. These technologies
generate high-resolution, spatially resolved molecular profiles that quantify gene or protein
expression at a single-cell or subcellular level[5–9]. A critical step in analyzing such data is to
assign cell identities based on spatial molecular profiles, a process known as cell annotation or
classification. This step sets a foundation for downstream analyses that infer cellular
interactions, spatial organization, and functional states.

Several approaches have been developed to annotate cells based on spatial molecular profiles.
Semi-supervised methods are widely used, relying on prior knowledge of marker genes[10–13] or
predefined combinations of positive and negative markers for specific cell types[14,15] to guide the
annotation of cell clusters generated by unsupervised algorithms. In contrast, unsupervised
methods cluster cells without prior knowledge or labels, and infer cell identities directly from the
data[16–18]. More recently, supervised approaches have emerged for large-scale datasets,
leveraging manual annotations of a subset of cells to train classifiers that subsequently annotate
the remaining cells[19,20].

However, existing cell annotation methods face significant challenges when applied to complex
tissues containing diverse cell populations with complex molecular markers. Rare cell types,
such as stem cells and cancer stem cells, play pivotal roles in physiological and pathological
contexts[21–24], but are difficult to manually identify for training existing annotation models. The
challenges are further exacerbated in spatial proteomics, where the number of measurable
protein markers is constrained by cost and technical limits[25,26]. Yet, understanding protein
expression and localization is essential for uncovering fundamental biological mechanisms and
disease pathology[27–29]. Recent advances in artificial intelligence have shown great promise in
addressing these challenges by leveraging the large-scale data generated by spatial imaging
technologies[30–32]. However, the available models and platforms are often not generalizable
beyond their original experimental context, as their performance is highly optimized for a
specific set of tissues, markers, and technical procedures[33,34].

In this study, we present QuantCell, a machine learning (ML) based framework designed to
improve cell annotation using qualitative and quantitative features derived from multiplexed
imaging. QuantCell leverages cell segmentation and image processing by QuPath[35,36], which
extracts intensity-based summary statistics across imaging channels and cellular
compartments. Using predefined sets of positive and negative markers for expected cell types,
QuantCell identifies cells with marker expression profiles unambiguously matching one of the
marker sets. These cells serve as "ground truth" for training and evaluating multiple ML models,
from which QuantCell selects the best-performing model to annotate previously unlabeled cells.
To ensure classification confidence, QuantCell uses a false discovery rate (FDR) based
thresholding strategy, allowing low-confidence or ambiguous cells to remain unannotated.

To demonstrate the utility of QuantCell, we imaged mouse bone marrow (BM) using the
PhenoCycler (formerly CODEX) platform (Akoya Biosciences). PhenoCycler leverages

DNA-barcoded antibodies to enable high-resolution, multiplexed imaging[37]. Mouse BM
contains over 20 distinct cell types with a wide range of frequencies[38–42]. Notably,
hematopoietic stem and progenitor cells (HSPCs) in the BM are well characterized by
complex combinations of cell surface markers[43–45], making BM an ideal system for
evaluating the accuracy and robustness of QuantCell.

**Results**

**QuantCell concept and design**

We demonstrated the performance of QuantCell using data acquired from the PhenoCycler[37]
platform. Mouse BM sections were imaged after staining with barcoded antibodies targeting 25
distinct proteins that are established cell type markers (Fig. 1a, Supplementary Fig. 1 and 2).
These markers were able to annotate 21 BM cell types (Supplementary Table 1), which
collectively comprise the vast majority of cells in the BM. Following the PhenoCycler
manufacturer's recommendation, cell segmentation was performed using nuclear staining in
QuPath[35], an open-source bioimage analysis software platform (Fig. 1b, Supplementary Fig.
3a). QuantCell performs equally well using nuclei-based and membrane-based cell
segmentation (Supplementary Fig. 3). Subsequently, the presence or absence of each protein
marker was annotated in QuPath for each segmented cell based on manual qualitative
assessment of marker expression for dozens of cells (Fig. 1c).

Cell annotation in QuPath requires manually identifying representative cells for each cell type,
which is labor-intensive, especially for rare cell types or those defined by multiple markers.
QuantCell leverages qualitative marker expression profiles from QuPath to classify cells based
on expected combinations of positive and negative markers for individual cell types (Fig. 1d,
Supplementary Table 1). Using this conventional approach based on qualitative marker
expression profiles, 33.1% of BM cells were annotated.

To classify the remaining cells, QuantCell leverages quantitative imaging data to improve cell
annotation derived from qualitative data. Using conventional annotations as the ground truth,
QuantCell trains ML models for cell classification on the quantitative imaging measurements of
individual cells (Fig. 1e). This process involves selecting appropriate ML models (Fig. 1f) and
fine-tuning their hyperparameters to optimize performance (Fig. 1g). For model selection and
training, 80% of the cells annotated by the conventional approach were used, with the remaining
20% reserved as a testing dataset. The testing dataset was used to identify the best-performing
model (Fig. 1h), which was then applied to classify cells that were not annotated by the
conventional approach (Fig. 1i).

**Selection and optimization of ML models for cell classification**

QuantCell systematically compares a variety of classification models based on their macro
average precision (AP) score (Fig. 1f). The AP score is derived from the precision–recall curve
(Fig. 2a), where precision reflects the accuracy of cell type prediction by the model, and recall
represents the fraction of ground-truth cells that are correctly annotated. The AP score, which

corresponds to the area under the precision–recall curve, provides a summary measure of
model performance. Precision, recall, and AP are calculated separately for each cell type, and
the unweighted average across all cell types defines the macro AP score.

As a demonstration, we compared 10 models: random forest, multilayer perceptron, extra trees,
logistic regression, linear SVC, nearest neighbors, decision tree, ridge classifier, LightGBM, and
gaussian naive-Bayes (Fig. 2b). The QuantCell framework readily accommodates additional
models as appropriate. Based on the AP score comparison, three best-performing ML models
were identified (Fig. 2a,b) and further optimized through hyperparameter tuning (Supplementary
Table 2). This fine-tuning process is essential to optimize each model's performance, ensuring
the selection of the most effective model. During model selection and parameter tuning,
principal component analysis (PCA) plots are generated to enable visual assessment. For each
cell type, we used (i) all markers in the panel and (ii) only the markers used for its conventional
annotation, to produce PCA plots showing conventionally annotated cells, QuantCell-annotated
cells, and cells not assigned to that cell type (Supplementary Fig. 4). These PCA plots help
guide model selection and parameter tuning by allowing researchers to identify potential
domain-shift issues.

While the optimal model selected by QuantCell may vary depending on the dataset, our analysis
using PhenoCycler data showed that the random forest model outperformed all other models
tested (Fig. 2c) and required substantially less runtime than the second-best model, the
multilayer perceptron (Supplementary Fig. 5a). The random forest model operates by
aggregating the predictions of multiple decision trees trained on subsets of the data, selecting
the most common classification outcome. This ensemble approach reduces overfitting while
maintaining strong predictive performance[46]. Random forest models are widely used for
classification tasks in many applications due to its robustness, versatility, and efficacy[47,48].

**135    Impact of the data sources and statistical descriptors on prediction accuracy**

During the ML model selection and optimization, the model was trained on all imaging
measurements obtained from QuPath's cell segmentation and marker annotation. This
comprehensive quantitative dataset included data captured from multiple cellular compartments,
such as membrane, cytoplasm, nucleus, and whole-cell region. For each compartment, multiple
statistical descriptors were collected, including the minimum, maximum, standard deviation,
median, and mean of fluorescence intensity for each protein marker. Understanding the impact
of data quantity, data source, and statistical descriptor on prediction accuracy could help identify
data types with the highest and lowest predictive value, including those with detrimental effects
that should be excluded to enhance QuantCell performance (Fig. 2d-f). This analysis was
conducted using the hyperparameter-optimized random forest model, which demonstrated the
highest macro AP score in our evaluation (Fig. 2a-c). Measurements of all markers were
included when comparing cellular compartments and statistical descriptors. Cellular
compartments were defined using default StarDist[49]-based cell segmentation in QuPath.

Our analyses suggest that prediction precision depends on the amount of training data used.
However, we observed minimal difference between using the full training dataset and using only

75% of it, suggesting that our current training dataset provides sufficient predictive power (Fig.
2d). Comparing various data sources and statistical descriptors, our results suggest that the
model performed the best when incorporating all cellular compartments and statistical
measurements (Fig. 2e). Among the statistical measurements, the mean produced the best
prediction accuracy, followed by the median, while the minimum performed markedly worse
(Fig. 2e). This is expected as the minimum primarily reflects background intensity. Interestingly,
standard deviation outperformed the maximum (Fig. 2e), suggesting that variability in signal
may carry more predictive value than peak intensity alone.

Comparing data from different cellular compartments, we found that the intensity measured
across the entire cell generated the best prediction accuracy (Fig. 2e). Although most of the
protein markers analyzed are localized to the cell surface, our results show that features derived
from the cell membrane exhibited the lowest predictive performance (Fig. 2e). This is likely due
to the dense cellular architecture of the BM, where closely packed cells can lead to overlapping
membrane signals from adjacent cells. As a result, membrane measurements may not
accurately represent protein expression levels in individual cells. In addition, our imaging data
were acquired using a wide-field microscope rather than a confocal microscope. As a result, the
nuclear region likely includes signals originating from the cytoplasm and membrane in the z-
dimension.

We also performed a feature exclusion analysis to evaluate whether excluding any type of
imaging measurement would improve model performance. Overall, the impact of excluding any
single feature, for all 25 protein markers, was negligible (Fig. 2f). Notably, removing features
with high individual importance, such as the median or mean intensity of all cellular
compartments, or all statistical measures across the whole-cell region (Fig. 2e), resulted in only
minor reductions in the macro AP (Fig. 2f). This suggests that the QuantCell prediction remains
robust and can compensate using other spatial measurements, even in the absence of the most
informative features.

**Evaluation of QuantCell prediction performance**

Using the hyperparameter-optimized random forest model and imaging data from all cellular
compartments and statistical measurements, we next evaluated the predictive performance of
QuantCell. We began by examining potential misclassifications using the reserved testing
dataset annotated by the conventional method (Fig. 3a). While the conventional classification
may contain errors, these annotations were used to train QuantCell and therefore served as the
ground truth for this analysis.

Our results show that 96.5% of cells classified by QuantCell were consistent with annotations
from the conventional method, demonstrating high prediction accuracy across all cell types (Fig.
3a, Supplementary Fig. 6). Precision was also high across all cell types, with a minimum cell
type precision rate of 87.5%, underscoring the model's robustness even in unbalanced cell
populations (Fig. 3a, Supplementary Table 1, Supplementary Fig. 7). With high precision rates,
QuantCell was able to increase the number of annotated cells from 33.1% to 90.2% at a false
discovery rate of 5% (Fig. 3b). The cell type frequencies predicted by QuantCell were consistent

with those obtained from the conventional annotation (Fig. 3c, Supplementary Fig. 8a). To
independently validate QuantCell's classifications, we performed flow cytometry on
unfractionated BM cells from the same mice, focusing on select hematopoietic cell types. The
results showed consistent cell type frequencies between QuantCell's annotation and flow
cytometry data, further validating QuantCell's classification accuracy (Fig. 3d, Supplementary
Fig. 8b).

**Comparison of QuantCell with existing annotation frameworks**

To evaluate QuantCell's performance, we compared it with other cell annotation models
developed for proteomics data, which are difficult to annotate due to limited marker availability
(Table 1). Astir is a probabilistic model that uses deep recognition neural networks to annotate
cell types based on user-defined markers for each cell type[10]. AnnoSpat is a recently proposed
semi-supervised approach that first clusters cells, then identifies a subset of confidently
annotated cells using predefined positive and negative protein markers[15]. These annotated cells
are used to train a neural network, which subsequently annotates the remaining cells. MAPS is
another recently introduced model that uses a feedforward neural network trained on a ground
truth dataset to perform cell type annotation[19].

We compared each model's performance on the training dataset using balanced accuracy, the
unweighted average accuracy across all cell types. This metric was used instead of macro AP
because some models do not provide cell type probabilities required for AP calculation. Our
analysis shows that QuantCell achieves higher prediction accuracy than existing models (Fig.
3e) and operates 45 times faster than the second most accurate model, MAPS (Supplementary
Fig. 5b). QuantCell also demonstrates robust performance across all cell types (Supplementary
Fig. 9) and successfully annotates a substantial number of cells that cannot be classified by
conventional approaches (Fig. 3b). QuantCell is well suited for analyzing data from experiments
where the marker panels may not capture all expected cell types, allowing for flexible and high-
confidence annotation even in partially characterized biological systems.

Multilayer perceptron and MAPS are feed-forward neural networks, and both demonstrated
relatively strong performance in our evaluation (Fig. 2b, 3e). However, MAPS relies on manually
annotated cells for training, which is a major limitation[19] (Table 1). Manual cell annotation is
labor-intensive for cell types defined by multiple markers and unfeasible for rare cell
populations. In contrast, QuantCell automatically generates training datasets based on
molecular marker profiles, eliminating the need for manual cell type annotation.

The other two benchmarked methods, Astir and AnnoSpat, also have limitations in generating
training datasets. AnnoSpat's reliance on clustering compromises its performance in identifying
rare cell types. While clustering is useful for pattern discovery, this unsupervised approach lacks
the sensitivity of supervised learning, particularly in imbalanced datasets where rare cell types
can be absorbed into larger, heterogeneous clusters[50]. Furthermore, clustering treats all
markers as equally informative. Noise from uninformative markers can compromise the
clustering and annotation accuracy. Astir is constrained by its use of only positive markers for
annotation (Table 1), assuming that marker expression is higher in the corresponding cell types

than in all other cells. This assumption is not valid for markers shared by multiple cell types,
which is common among stem/progenitor cells and diseased cells.

**Cell type specific prediction performance**

A major strength of QuantCell is its strong performance across cell types with widely varying
frequencies. We further evaluated QuantCell's performance across individual cell types using
the F1 score, the harmonic mean of precision and recall which summarizes predictive
performance for each cell type. Our results suggested that the F1 score was significantly
influenced by cell frequency and cell circularity (Fig. 4a), likely due to its effect on the sample
size and quality for model training. In contrast, features such as cell size and the number of
protein markers used in conventional annotation did not show direct correlation with F1 score
(Fig. 4a), indicating the robustness and consistency of QuantCell's performance across cell
types.

To assess the contribution of individual protein markers to cell type classification, we
systematically excluded one marker at a time from the training dataset. This analysis revealed
the markers that are indispensable for accurately annotating specific cell types (Fig. 4b,
Supplementary Fig. 10). These essential markers often, though not always, correspond to those
used in conventional annotation (Supplementary Table 1). This discrepancy reflects how the
model weighs informative features for each cell type and the redundancy among markers. For
example, removing one lineage marker may have little effect on HSPCs because other lineage
markers provide similar information. Together, these results validate QuantCell's predictions
and highlight its reliability and effectiveness in classifying diverse cell types.

**Discussion**
In this study, we developed QuantCell, an ML based framework designed to improve cell
annotation using qualitative and quantitative imaging data. We validated the performance of
QuantCell using multiple complementary approaches. First, we assessed its accuracy on an
independently held-out test dataset generated through conventional annotation (Fig. 3a,
Supplementary Fig. 7). Second, we confirmed QuantCell predictions using flow cytometry
measurements (Fig. 3d, Supplementary Fig. 8b). Third, all marker and cell annotations were
independently reviewed by three field experts. Fourth, we evaluated QuantCell on three
externally annotated datasets from published studies, demonstrating robust, high-precision
performance (Supplementary Fig. 11). Finally, PCA analyses showed that QuantCell-annotated
cells tightly colocalize with conventionally annotated cells and are clearly separated from cells
not assigned to the corresponding cell type (Supplementary Fig. 4).

QuantCell outperforms existing methods in predictive accuracy and offers significant
advancements in the following key areas. First, QuantCell optimizes annotation performance
across all cell types (Fig. 3a, Supplementary Fig. 7, 9). By avoiding performance bias toward
abundant cell types, it provides a more balanced and accurate classification. This is particularly
important in biomedical datasets where rare cell types, such as stem cells or diseased cells, are
often the primary focus but are poorly detected by existing methods. Second, QuantCell

performs annotation based on a user-defined FDR, enabling stringent control over classification
confidence (Fig. 3b). It reduces the risk of misclassification by leaving uncertain cells
unannotated, which improves the reliability of biological interpretation. Third, QuantCell does not
require manual cell type annotation, which is labor-intensive for cell types defined by multiple
markers and unfeasible for rare cell populations. It uses both positive and negative markers to
guide training dataset generation, enabling evaluation of each marker's contribution (Fig. 4b,
Supplementary Fig. 10). Nonetheless, QuantCell's performance is constrained by the quality of
the training data. It may underperform on datasets with sparse or low-dimensional features (Fig.
2e) or for cell types with limited or low-quality training data (Fig. 4a). To improve the quality of
the training and testing datasets, one potential approach is to filter out outlier cells annotated by
the conventional method. Because the conventional method assigns cell identities based on
predefined marker combinations (Supplementary Table 1), cells of the same type are expected
to cluster in the corresponding marker space (Supplementary Fig. 4, left panels). By comparing
the density of cells assigned to a given type with the densities of other types in this space, it is
possible to estimate the probability that each cell truly belongs to its assigned identity. The cell
distributions can be visually inspected in PCA plots to determine whether low-probability cells
should be removed. Implementing such procedures in the future could further improve model
predictions.

Despite this limitation, QuantCell is able to increase the number of annotated cells from 33.1%
to 90.2% at a 5% FDR, achieving 96.5% overall accuracy based on the test dataset from
conventional annotation (Fig. 3b). Our benchmarking analysis shows that QuantCell
outperforms existing platforms in predictive accuracy (Fig. 3e). It has the potential to be
compatible with a wide range of spatial omics platforms, including two- and three-dimensional
imaging, RNA-based technologies such as seqFISH[51] and MERFISH[52], and sequencing-based
approaches like Visium[53], Slide-seq[54], and Stereo-seq[9]. The framework is also adaptable to
project-specific priorities. For example, QuantCell is configured by default to optimize
performance equally across all cell types, but users can customize weighting schemes to
prioritize specific cell populations of interest. Users can also incorporate their preferred
supervised ML models in QuantCell for model selection (Fig. 1f). Future studies could further
expand QuantCell's capabilities by incorporating transfer learning to enhance performance in
data-sparse settings. In addition, incorporating domain adaptation techniques may enable
QuantCell to generalize across tissue types, experimental platforms, or imaging modalities with
minimal retraining.

In conclusion, QuantCell represents a powerful application of ML in the rapidly advancing field
of spatial omics. It improves cell annotation by leveraging large-scale data from high-resolution
multiplexed imaging. We demonstrate that QuantCell not only accurately annotates a
substantial number of cells left unclassified by conventional methods, but also performs robustly
across diverse BM cell types, underscoring its potential to resolve complex cellular phenotypes
in heterogeneous tissues.

**Methods**

<u>Mice</u>

C57BL/6J mice (Jackson Laboratory stock #000664) were maintained in the Research Animal
Facility at the University of Southern California (USC) under a 12:12 hour light-dark cycle, with
controlled room temperature ($22 \pm 1\,°C$) and humidity ($55 \pm 10\%$). All animal procedures were
approved by the Institutional Animal Care and Use Committee of USC.

PhenoCycler antibodies

The following primary antibodies were obtained from Akoya Biosciences: RX025-AF488 anti-
CD11b (M1/70), RX030-Cy5 anti-CD11c (N418), RX020-ATTO550 anti-CD19 (6D5), RX021-Cy5
anti-CD3 (17A2), RX002-ATTO550 anti-CD31 (MEC13.3), RX026-ATTO550 anti-CD4 (RM4-5),
RX005-ATTO550 anti-CD44 (IM7), RX007-AF488 anti-CD45 (30-F11), RX010-AF488 anti-
CD45R (RA3-6B2), RX027-Cy5 anti-CD71 (RI7217), RX029-ATTO550 anti-CD8a (53-6.7),
RX024-Cy5 anti-Ly6g (1A8), RX014-ATTO550 anti-MHCII (M5/114.15), RX003-Cy5 anti-TCRαβ
(H57-597), and RX004-AF488 anti-TER119 (TER-119). The following primary antibodies were
obtained from ThermoFisher Scientific: BX035-ATTO550 anti-CD105 (209701), BX050-AF647
anti-CD115 (AFS98), BX045-Cy5 anti-CD117 (2B8), RX013-AF488 anti-CD34 (RAM34), BX037-
ATTO550 anti-CD41 (MWReg30), BX040-AF488 anti-endomucin (polyclonal), BX036-Cy5 anti-
F4/80 (CI:A3-1), and BX041-ATTO550 anti-CD127 (A7R34). The following primary antibody was
obtained from Leinco Technologies: BX028-AF488 anti-CD32/CD16 (2.4G2). The following
primary antibody was obtained from R&D Systems: BX049-ATTO550 anti-LEPr (polyclonal). All
primary antibodies obtained from vendors other than Akoya Biosciences were conjugated to
Akoya barcodes by Akoya Biosciences.

Antibody validation and quality control

All antibodies were conjugated and validated by Akoya Biosciences. Successful conjugation was
verified by sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS–PAGE), confirming
that each conjugated antibody migrated at the expected molecular weight. After verification of
conjugation, Akoya staff stained our tissue sections with the antibodies and evaluated antibody
performance using a standardized scoring system: –1 (non-specific binding), 0 (no signal/no
positive cells), 1 (weak signal), 2 (good staining), and 3 (saturated). Antibody clones and/or
dilutions were adjusted based on these evaluations. A blank imaging cycle was used as a negative
control. All antibodies included in the final panel received a rating of 2.

Bone marrow preparation for flow cytometry analysis

Bone marrow cells were isolated from crushed bones of the arms, spine, pelvis, and sternum.
Cells were separated from bone debris using a density gradient (Sigma-Aldrich, Cat. #11191).
The resulting cell suspension was then divided into two fractions for antibody staining: one for
analyzing mature blood cells and the other for identifying hematopoietic stem and progenitor cells
(HSPCs).

The following antibodies were used to identify mature blood cells: PE anti-TCRαβ (H57-597), PE-
Cy5 anti-CD8α (53-6.7), PE-Cy5.5 anti-CD19 (eBio1D3), PE-Cy7 anti-CD45.2 (104), eFluor660

anti-B220 (RA3-6B2), APC-eFluor780 anti-CD11b (M1/70), BV510 anti-Gr-1 (RB6-8C5), PE-
eFluor610 anti-CD4 (GK1.5), and PerCP-eFluor710 anti-CD115 (AF598).

The following antibodies were used to identify HSPCs: PE anti-CD150 (TC15-12F12.2), PE-Cy5
anti-CD135 (A2F10), PE-Cy7 anti-Ly-6A/E (D7), AF700 anti-CD45.2 (104), APC-eFluor780 anti-
CD117 (2B8), BV510 anti-CD127 (A7R34), eFluor450 Lineage (anti-CD3 (17A2), anti-CD19
(eBio1D3), anti-CD11b (M1/70), anti-Gr-1 (RB6-8C5), anti-TER-119 (TER-119), anti-CD4
(GK1.5), anti-CD8α (53-6.7), anti-B220 (RA3-6B2)), Biotin anti-CD32/CD16 (93), and BV605
Streptavidin (secondary).

Bone marrow cells were incubated with the mature blood cell antibody mix for 1 hour, or with the
HSPC antibody mix for 1.5 hours, followed by a 20-minute incubation with the secondary antibody.
After antibody incubation, samples stained for mature blood cells were resuspended in Dulbecco's
Phosphate Buffered Saline (DPBS) (ThermoFisher, Cat. #21600010), supplemented with 2% fetal
bovine serum (FBS; Avantor, Cat. #76419-584) and 4′,6-diamidino-2-phenylindole (DAPI).
Samples stained for HSPCs were resuspended in DPBS containing 2% FBS and propidium
iodide. All samples were analyzed using a FACSAria II (Becton Dickinson) as previously
described[55–57]. Flow cytometry data were analyzed using FlowJo software version 10.8.2 (Becton
Dickinson).

Bone marrow preparation for imaging

Bone marrow was carefully flushed from the diaphysis of femurs and tibias using a syringe filled
with DPBS with 2% FBS, ensuring gentle handling to obtain the whole, intact bone marrow plugs.
The collected bone marrow was immediately fresh-frozen in tissue freezing medium (Electron
Microscopy Sciences, Cat. #72592) for downstream applications.

Bone marrow imaging

Glass coverslips, 22 mm x 22 mm, were pre-treated with poly-L-lysine (Santa Cruz Biotechnology,
Cat. #sc-286689) to enhance tissue adherence. The fresh-frozen bone marrow samples were
cryosectioned at a thickness of 5 μm and mounted onto the prepared coverslips. Six BM section
samples from six mice were imaged and processed, yielding a total of 245,719 cells identified
through QuPath segmentation. Cells from all six samples were pooled for downstream analyses.
Antibody staining and imaging were performed by Akoya Biosciences personnel using Akoya's
established protocols.

Cell segmentation

Nuclei-based cell segmentation was performed based on DAPI stains in QuPath[35] (version
0.3.0) using the StarDist[49] (version 0.3.0) algorithm, following default settings (provided in our
GitHub, see Code availability). For nuclei-based segmentation, QuPath automatically infers
cytoplasm and membrane compartments by expanding nuclear boundaries using a watershed
approach. Membrane-based segmentation was performed with Cellpose-SAM[58] (version 4.0.6)
using a composite image of DAPI, F4/80, CD45, Endomucin, Sca1, and CD71. To recover
cellular compartments using Cellpose, segmentation was run independently on the composite

(whole cell) and the DAPI channel (nucleus). The cytoplasmic compartment was defined by
subtracting the nuclear area from the whole-cell area. QuantCell performed comparably well
with either membrane-based or nuclei-based cell segmentation (Supplementary Fig. 3). Nuclei-
based segmentation was used for all other analyses.

Conventional annotation and QuPath settings

Qualitative marker annotation was performed in QuPath[35] (version 0.3.0). For each marker,
approximately 20-70 positive and 20-70 negative cells from across the tissue section were
manually annotated and used to classify all cells in the section. This process involved multiple
iterations of model training, manual review, and additional manual refinement of model-
generated annotations, until the resulting cell annotations passed human assessment. A cell
was considered positive for a marker when its signal clearly exceeded the background signal
and exhibited the expected staining pattern for that marker. Signal adjustment and staining
patterns were used to minimize the effects of non-specific binding and background signals.
Mutually exclusive and co-expressing marker combinations were also used for further
verification.

To reduce potential cross-contamination from neighboring cells, our annotation strategy
incorporated the spatial distribution of marker signal within each cell. During manual binary
marker annotation in QuPath, cells were classified as negative when the apparent signal was
confined to regions adjacent to neighboring positive cells. In the QuantCell annotation, spatial
distribution for each marker was quantified using multiple statistical metrics, including the
standard deviation of signal intensity within each cellular compartment.

The resulting marker assignments were independently verified by three field experts, including a
board-certified pathologist, the director of our imaging core facility, and an experienced tissue
biologist. After verification, the protein-expression–by–cell matrix and the binary
positive/negative marker annotations were downloaded from QuPath. The binary annotations
were then used for the conventional cell annotation. Detailed information on the procedure used
to generate the protein-expression-by-cell matrix is available in our GitHub repository (see Code
availability).

Using the binary positive/negative marker annotations generated by QuPath, QuantCell
constructed the training and testing datasets by annotating cells that exactly matched
predefined marker combinations (Supplementary Table 1). Cells with ambiguous profiles, such
as those matching more than one combination or none at all, were not assigned a cell type and
were labeled as "Other". To further verify cell annotation accuracy, single-cell annotations
generated by both the conventional approach and by QuantCell were independently assessed
by three experts as mentioned above.

Data preprocessing

Cells that were successfully annotated by the conventional method were split into an 80/20
train-test dataset using train_test_split() from scikit-learn[59], with stratification based on class

distribution. Features were normalized to have a mean of 0 and a standard deviation of 1 using
StandardScaler() from scikit-learn. The scaler was fitted to the marker distributions in the
training dataset and then applied to both the testing dataset and unannotated cells without
refitting, to prevent data leakage. Cell type labels were encoded using LabelEncoder() from
scikit-learn after excluding any cells that were originally unannotated.

### Model selection

430   Initial model performance was evaluated across eight different base classifiers from scikit-learn
431   with default parameters, using the training dataset and 10-fold cross-validation. The three best-
432   performing models were selected based on macro average precision (AP) score. These models
433   were further optimized using HalvingGridSearchCV from scikit-learn, with 10-fold cross-
434   validation on the training dataset.

### Training size analysis

To evaluate the dependence of model performance on training data availability, we performed a
training size analysis using 5-fold cross-validation. Within each fold, the test set was held
constant to ensure paired comparisons, while the training set was randomly subsampled at
varying proportions (5%, 10%, 25%, 50%, 75%, and 100%). Consequently, the 100% condition
represents the full set of remaining 4 folds and is equivalent to the 80/20 train-test split used
during preprocessing.

### Feature analysis

443   Performance of the model was evaluated by systematically varying the inclusion or exclusion of
444   specific measures or markers across 10-fold cross-validation on the training set. Each fold was
445   treated as an individual data point. F1 scores were calculated without applying a false discovery
446   rate (FDR) threshold. The model using all features was used as the baseline to assess changes
447   in macro AP and F1 scores

### Model thresholding

449   Thresholding was performed independently for each cell type by examining the predicted cell
450   type probabilities for each cell on the testing dataset and selecting a threshold that ensured the
451   FDR was less than or equal to the specified target. For thresholded analyses in this study, the
452   default FDR target was set to 5%. Thresholding was not applied in any analysis involving the AP
453   score, as AP is calculated by evaluating performance across all possible thresholds.

### Benchmarking QuantCell against existing models

For all benchmarking analyses, annotations from the conventional method were used as the
ground truth. For Astir, AnnoSpat, and MAPS, mean expression of each marker across the
entire cell was used as input, following the pipeline described in the corresponding papers.
MAPS was also provided the cell area ($\mu m^2$) as directed.

Astir and AnnoSpat were applied to the entire conventionally annotated dataset. No
preprocessing was performed. The models were run with default parameters. The same marker
combinations used in QuantCell were applied, except negative markers were excluded for Astir,
as they are not supported by this model.

MAPS was trained and tested using the same 80/20 train–test split as QuantCell. Labels were
encoded as in the QuantCell workflow prior to training. No preprocessing was applied, and
default parameters were used, except for "min_epochs" which was reduced to 5 and "patience"
which was reduced to 25 to avoid unnecessary computation.

QuantCell was trained and tested using the same 80/20 train–test split as MAPS. Performance
benchmarking was conducted both with thresholding at 5% FDR**.**

<u>Computational environment</u>

All benchmarking tests and analyses were performed on a dedicated server with Ubuntu
20.04.6 LTS, 2 x Intel(R) Xeon(R) Gold 6136 CPU @ 3.00GHz (24 physical cores, 48 threads
total, 49.5 MB L3 cache), 494.7 GB memory. All code is in Python, except for a Bash script used
to run AnnoSpat.

<u>Statistics and reproducibility</u>

Statistical tests used in data analysis are detailed in the relevant figure legends and "Methods"
sections. No statistical method was used to predetermine sample size. Expression data from
markers with poor staining quality or those not included in any marker combinations
(Supplementary Table 1) were excluded from analysis. Flow cytometry validation was
performed for cell types with gating strategies matching the marker combinations
(Supplementary Table 1). Where applicable, 10-fold cross-validation was employed for model
selection. A fixed random seed was applied during data splitting to ensure reproducibility across
sessions.

**Figures**

484
**Figure 1. Overview of conventional and QuantCell annotation pipelines.** a) Tissue sections
are prepared, stained, and imaged. b) Cell segmentation is performed using QuPath[35], based
on 4′,6-diamidino-2-phenylindole (DAPI) nuclei staining. c) Each protein marker is manually
annotated as positive or negative in a small subset of cells. These annotations are then
propagated to the remaining cells using QuPath. d) Cells are classified based on expected
combinations of positive and negative markers for each cell type. e) Data are scaled and
encoded. 80% of the cells annotated by the conventional method are used for model selection
and training, while the remaining 20% are reserved as a testing dataset. f) Multiple machine
learning models are evaluated using 10-fold cross-validation. g) The top three performing
models are further optimized through hyperparameter tuning to select the most effective model.
h) The final model is evaluated on the reserved 20% testing dataset. i) The final model is
applied to classify cells that are unannotated in d), using a 5% false discovery rate threshold.

**Figure 2. Selection and optimization of machine learning models for cell classification.** a) Precision-recall curve for base classifiers without hyperparameter tuning. Data represent the unweighted average of all cell types. b) Macro average precision (AP) score for each classifier, equivalent to the area under the precision-recall curve (AUPRC). The dashed line indicates the AUPRC corresponding to random prediction performance. c) Comparison of the default and hyperparameter-optimized performance of the top three models based on AP score. d) Comparison of QuantCell performance using varying proportions of the training dataset. Average scores across 5-fold cross-validation are shown as black dots above each bar. e) Evaluation of feature importance using the best-performing model, hyperparameter-optimized random forest. Shown are the prediction AP scores using training data from the indicated

508  measurement(s). f) Feature exclusion analysis to assess the contribution of individual features.
509  Shown are changes in macro AP when the model was trained on all measurements except the
510  indicated one(s). e and f) Standard deviations were calculated from 10-fold cross-validation.
511  "Cell" refers to the average intensity across the entire cell.

512
**Figure 3. Evaluation of QuantCell prediction performance.** a) Column-normalized confusion
matrix comparing conventional cell annotation with QuantCell annotation. Values on the
diagonal are equivalent to precision. b) Fraction of cells classified by the conventional method
(blue) and those unclassified by the conventional method but successfully classified by
QuantCell (green). QuantCell was configured to ensure that the false discovery rate (FDR) for
all cell types are below the specified threshold. c) Comparison of cell type frequencies obtained

from conventional annotation and QuantCell annotation, calculated as fractions of all cells,
including unannotated ones. d) Comparison of cell type frequencies derived from QuantCell
annotations and flow cytometry analyses of the same mice. QuantCell frequencies are
calculated among the annotated cell types, which represent most but not all bone marrow cells,
whereas flow cytometry frequencies are measured relative to all alive bone marrow cells. e)
Benchmarking QuantCell against other spatial proteomics annotation methods. Cell type
abbreviations are listed in Supplementary Table 1. c and d) Two-tailed Pearson's correlation
coefficient (r) and *p*-value are shown. Each dot represents one cell type. The dashed line
represents the best-fit line computed using least squares regression.

**Figure 4. QuantCell prediction performance across cell types.** a) Comparison of F1 scores
without FDR thresholding based on the cell type specific attributes: cell frequency, cell size, cell
circularity, and the number of positive and negative markers used in conventional annotation.
Each dot represents one of 21 distinct cell types. Two-tailed Pearson's correlation coefficients
and corresponding *p*-values are shown. b) Impact of individual protein markers on the prediction
performance for representative cell types. Shown are changes in F1 score resulting from the

exclusion of individual protein markers from the training and testing datasets. Each dot

represents the change in the F1 score relative to the baseline (no exclusion), calculated using

10-fold cross-validation. Red dots indicate the mean change across all 10 folds. Markers with a

mean change exceeding 1.96 standard deviations from the overall distribution (z-score > 1.96)

are annotated.

**Table 1. Comparison of QuantCell with existing annotation frameworks**

| | Method | Input data | Manual annotation required | Marker combos | Uncertainty (allow unlabeled) | Balanced label accuracy |
|---|---|---|---|---|---|---|
| **Astir**[10] | Clustering | Protein-Expression-by-cell matrix & marker combos | No | Positive | Yes | 0.258 |
| **AnnoSpat**[15] | Clustering | Protein-Expression-by-cell matrix & marker combos | No | Positive & negative | Yes | 0.433 |
| **MAPS**[19] | Feed forward neural network | Protein-Expression-by-cell matrix & cell size | Yes | No | No | 0.873 |
| **QuantCell** | Variable classifiers | Protein-Expression-by-cell matrix & marker combos | No | Positive & negative | Yes | 0.928 |

**References**

1. Marx, V. Method of the Year: spatially resolved transcriptomics. *Nat. Methods* **18**, 9–14 (2021).
2. Lundberg, E. & Borner, G. H. H. Spatial proteomics: a powerful discovery tool for cell biology. *Nat. Rev. Mol. Cell Biol.* **20**, 285–302 (2019).
3. Seydel, C. Beyond cell atlases: spatial biology reveals mechanisms behind disease. *Nat. Biotechnol.* **43**, 841–844 (2025).
4. Cheng, X. *et al.* Application of single-cell and spatial omics in deciphering cellular hallmarks of cancer drug response and resistance. *J Hematol Oncol* **18**, 70- (2025).
5. Stadler, C. *et al.* Immunofluorescence and fluorescent-protein tagging show high correlation for protein localization in mammalian cells. *Nat. Methods* **10**, 315–323 (2013).
6. Goltsev, Y. *et al.* Deep Profiling of Mouse Splenic Architecture with CODEX Multiplexed Imaging. *Cell* **174**, 968-981.e15 (2018).
7. Merritt, C. R. *et al.* Multiplex digital spatial profiling of proteins and RNA in fixed tissue. *Nat. Biotechnol.* **38**, 586–599 (2020).
8. Eng, C.-H. L. *et al.* Transcriptome-scale super-resolved imaging in tissues by RNA seqFISH. *Nature* **568**, 235–239 (2019).
9. Chen, A. *et al.* Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA nanoball-patterned arrays. *Cell* **185**, 1777-1792.e21 (2022).
10. Geuenich, M. J. *et al.* Automated assignment of cell identity from single-cell multiplexed imaging and proteomic data. *Cell Syst.* **12**, 1173-1186.e5 (2021).
11. Coleman, K., Hu, J., Schroeder, A., Lee, E. B. & Li, M. SpaDecon: cell-type deconvolution in spatial transcriptomics with semi-supervised learning. *Commun. Biol.* **6**, 378–13 (2023).
12. Shen, R. *et al.* Spatial-ID: a cell typing method for spatially resolved transcriptomics via transfer learning and spatial embedding. *Nat. Commun.* **13**, 7640–17 (2022).
13. Zhou, Y., He, W., Hou, W. & Zhu, Y. Pianno: a probabilistic framework automating semantic annotation for spatial transcriptomics. *Nat. Commun.* **15**, 2848–15 (2024).
14. Ianevski, A., Giri, A. K. & Aittokallio, T. Fully-automated and ultra-fast cell-type identification using specific marker combinations from single-cell transcriptomic data. *Nat. Commun.* **13**, 1246–1246 (2022).
15. Mongia, A. *et al.* AnnoSpat annotates cell types and quantifies cellular arrangements from spatial proteomics. *Nat. Commun.* **15**, 3744–19 (2024).
16. Kiselev, V. Y., Andrews, T. S. & Hemberg, M. Challenges in unsupervised clustering of single-cell RNA-seq data. *Nat. Rev. Genet.* **20**, 273–282 (2019).
17. Kumagai, Y. BootCellNet, a resampling-based procedure, promotes unsupervised identification of cell populations via robust inference of gene regulatory networks. *PLoS Comput. Biol.* **20**, e1012480- (2024).
18. Marghi, Y., Gala, R., Baftizadeh, F. & Sümbül, U. Joint inference of discrete cell types and continuous type-specific variability in single-cell datasets with MMIDAS. *Nat. Comput. Sci.* **4**, 706–722 (2024).
19. Shaban, M. *et al.* MAPS: pathologist-level cell type annotation from tissue images through machine learning. *Nat. Commun.* **15**, 28–11 (2024).

20.    Amitay, Y. *et al.* CellSighter: a neural network to classify cells in highly multiplexed
images. *Nat. Commun.* **14**, 4302–4302 (2023).
21.    Nguyen, A., Khoo, W. H., Moran, I., Croucher, P. I. & Phan, T. G. Single Cell RNA
Sequencing of Rare Immune Cell Populations. *Front. Immunol.* **9**, 1553–1553 (2018).
22.    Patel, S. B., Franceski, A. M., Crown, B. L. & Welner, R. S. Leading Edge Techniques in
the Quest for Characterizing Rare Hematopoietic Stem Cells. *Curr. Stem Cell Rep.* **10**, 108–125
(2024).
23.    Altshuler, A., Wickström, S. A. & Shalom-Feuerstein, R. Spotlighting adult stem cells:
advances, pitfalls, and challenges. *Trends Cell Biol.* **33**, 477–494 (2023).
24.    Abbaszadegan, M. R. *et al.* Isolation, identification, and characterization of cancer stem
cells: A review. *J. Cell. Physiol.* **232**, 2008–2018 (2017).
25.    Wu, M. *et al.* Spatial proteomics: unveiling the multidimensional landscape of protein
localization in human diseases. *Proteome Sci.* **22**, 7–15 (2024).
26.    Bressan, D., Battistoni, G. & Hannon, G. J. The dawn of spatial omics. *Sci. Am. Assoc.*
*Adv. Sci.* **381**, (2023).
27.    Hung, M.-C. & Link, W. Protein localization in disease and therapy. *J. Cell Sci.* **124**,
3381–3392 (2011).
28.    Blise, K. E., Sivagnanam, S., Banik, G. L., Coussens, L. M. & Goecks, J. Single-cell
spatial architectures associated with clinical outcome in head and neck squamous cell
carcinoma. *NPJ Precis. Oncol.* **6**, 10–10 (2022).
29.    Han, J.-D. J. Understanding biological functions through molecular networks. *Cell Res.*
**18**, 224–237 (2008).
30.    Dong, K. & Zhang, S. Deciphering spatial domains from spatially resolved
transcriptomics with an adaptive graph attention auto-encoder. *Nat. Commun.* **13**, 1739–12
(2022).
31.    Bae, S. *et al.* CellDART: cell type inference by domain adaptation of single-cell and
spatial transcriptomic data. *Nucleic Acids Res.* **50**, e57–e57 (2022).
32.    Li, B. *et al.* Benchmarking spatial and single-cell transcriptomics integration methods for
transcript distribution prediction and cell type deconvolution. *Nat. Methods* **19**, 662–670 (2022).
33.    Ge, S., Sun, S., Xu, H., Cheng, Q. & Ren, Z. Deep learning in single-cell and spatial
transcriptomics data analysis: advances and challenges from a data science perspective. *Brief.*
*Bioinform.* **26**, (2025).
34.    Gohil, V. *et al.* The Importance of Generalizability in Machine Learning for Systems.
*IEEE Comput. Archit. Lett.* **23**, 95–98 (2024).
35.    Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis.
*Sci. Rep.* **7**, 16878–7 (2017).
36.    Loughrey, M. B. *et al.* Validation of the systematic scoring of immunohistochemically
stained tumour tissue microarrays using QuPath digital image analysis. *Histopathology* **73**, 327–
338 (2018).
37.    Black, S. *et al.* CODEX multiplexed tissue imaging with DNA-conjugated antibodies. *Nat.*
*Protoc.* **16**, 3802–3835 (2021).
38.    Cesar Nombela-Arrieta & Markus G Manz. Quantification and three-dimensional
microanatomical organization of the bone marrow. *Blood Adv.* **1**, 407–416 (2017).

39.    Baccin, C. *et al.* Combined single-cell and spatial transcriptomics reveal the molecular,
cellular and spatial bone marrow niche organization. *Nat. Cell Biol.* **22**, 38–48 (2020).
40.    Dolgalev, I. & Tikhonova, A. N. Connecting the Dots: Resolving the Bone Marrow Niche
Heterogeneity. *Front. Cell Dev. Biol.* **9**, 622519 (2021).
41.    Ding, L. & Morrison, S. J. Haematopoietic stem cells and early lymphoid progenitors
occupy distinct bone marrow niches. *Nature* **495**, 231–235 (2013).
42.    Bryder, D., Rossi, D. J. & Weissman, I. L. Hematopoietic stem cells: the paradigmatic
tissue-specific stem cell. *Am. J. Pathol.* **169**, 338–346 (2006).
43.    Spangrude, G. J., Heimfeld, S. & Weissman, I. L. Purification and characterization of
mouse hematopoietic stem cells. *Science* **241**, 58–62 (1988).
44.    Purton, L. E. Adult murine hematopoietic stem cells and progenitors: an update on their
identities, functions, and assays. *Exp. Hematol.* **116**, 1–14 (2022).
45.    Wu, Y.-C., Kissner, M. & Momen-Heravi, F. A comprehensive multiparameter flow
cytometry panel for immune profiling and functional studies of frozen tissue, bone marrow, and
spleen. *J. Immunol. Methods* **515**, 113444–113444 (2023).
46.    Breiman, L. Random Forests. *Mach. Learn.* **45**, 5–32 (2001).
47.    Chen, X. & Ishwaran, H. Random forests for genomic data analysis. *Genomics* **99**, 323–
329 (2012).
48.    Boulesteix, A.-L., Janitza, S., Kruppa, J. & König, I. R. Overview of random forest
methodology and practical guidance with emphasis on computational biology and
bioinformatics. *Wiley Interdiscip. Rev. Data Min. Knowl. Discov.* **2**, 493–507 (2012).
49.    Schmidt, U., Weigert, M., Broaddus, C. & Myers, G. Cell Detection with Star-convex
Polygons. *arXiv.org* https://doi.org/10.48550/arxiv.1806.03535 (2018)
doi:10.48550/arxiv.1806.03535.
50.    Weber, L. M. & Robinson, M. D. Comparison of clustering methods for high-dimensional
single-cell flow and mass cytometry data. *Cytom. Part J. Int. Soc. Anal. Cytol.* **89**, 1084–1096
(2016).
51.    Lubeck, E., Coskun, A. F., Zhiyentayev, T., Ahmad, M. & Cai, L. Single-cell in situ RNA
profiling by sequential hybridization. *Nat. Methods* **11**, 360–361 (2014).
52.    Chen, K. H., Boettiger, A. N., Moffitt, J. R., Wang, S. & Zhuang, X. Spatially resolved,
highly multiplexed RNA profiling in single cells. *Sci. Am. Assoc. Adv. Sci.* **348**, 412–412 (2015).
53.    Ståhl, P. L. *et al.* Visualization and analysis of gene expression in tissue sections by
spatial transcriptomics. *Sci. Am. Assoc. Adv. Sci.* **353**, 78–82 (2016).
54.    Rodriques, S. G. *et al.* Slide-seq: A scalable technology for measuring genome-wide
expression at high spatial resolution. *Sci. Am. Assoc. Adv. Sci.* **363**, 1463–1467 (2019).
55.    Jiang, D. *et al.* Quantitative association between gene expression and blood cell
production of individual hematopoietic stem cells in mice. *Sci. Adv.* **10**, eadk2132 (2024).
56.    Nogalska, A. *et al.* Age-associated imbalance in immune cell regeneration varies across
individuals and arises from a distinct subset of stem cells. *Cell. Mol. Immunol.* **21**, 1459–1473
(2024).
57.    Nguyen, L. *et al.* Functional compensation between hematopoietic stem cell clones in
vivo. *EMBO Rep.* **19**, e45702 (2018).

58.      Pachitariu, M., Rariden, M. & Stringer, C. Cellpose-SAM: superhuman generalization for
cellular segmentation. 2025.04.28.651001 Preprint at
https://doi.org/10.1101/2025.04.28.651001 (2025).
59.      Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *J. Mach. Learn. Res.*
https://doi.org/10.5555/1953048.2078195 (2011) doi:10.5555/1953048.2078195.
60.      Shaban, M. et al. Data for MAPS: Pathologist-level Cell Type Annotation from Tissue
Images through Machine Learning. Zenodo https://doi.org/10.5281/zenodo.10067010 (2023).
679
680

**Acknowledgements**
We thank Akoya Biosciences for their assistance in establishing the antibody panel and
acquiring imaging data, and Drs. G. Fudenberg, A. MacLean as well as the members of the Lu
laboratory for their valuable discussions and insights. We also thank Drs S.W. Ruffins and I.
Hajjali for assistance with validating marker and cell annotations. This study is supported by
R35HL150826, R01DK143671, and R01AG080982. W.B. was supported by USC Provost's
Fellowship, QCB Google Alumni Award in Quantitative Biology, and the John Petruska
Memorial Award.

**Author contributions**
W.B. and R.L. conceived and designed the project. B.W. designed the antibody marker panel
and cell type marker combinations, conducted all wet lab experiments including bone marrow
sectioning. B.W. also performed cell segmentation and marker annotation in QuPath. W.B.
analyzed the data and wrote all project-related code. B.W., Z.T., A.N., and R.L. assisted with
data analysis. W.B. and R.L. wrote the manuscript. All authors reviewed and approved the final
manuscript.

**Data availability**
Data supporting the findings of this study are available from the corresponding author upon
request.

**Code availability**
QuantCell code is available at https://github.com/ronglulab/QuantCell. All code used for data
analysis and figure generation is provided in the "paper" directory of the repository.

**Competing interests**
The authors declare no competing interests.