1

Differential pairings of MAT1-1-1 and MAT1-2-1 protein variants with heteromorphic tertiary structures produced by *Ophiocordyceps sinensis* strains

Xiu-Zhang Li [1], Yu-Ling Li [1], Wei Liu [2], Jia-Shi Zhu [1]*

[1] State Key Laboratory of Plateau Ecology and Agriculture, Qinghai Academy of Animal and Veterinary Sciences, Qinghai University, Xining, Qinghai 810016, China

[2] Institute of Immunology, Army Medical University, Chongqing, 400038, China

* Corresponding author

E-mail: zhujosh@163.com (JSZ)

Simple Summary

Cordyceps sinensis insect–fungal complex is a combination of the remains of a Hepialidae moth larva and the *Ophiocordyceps sinensis* fruiting body. For centuries, it has been used in traditional Chinese medicine for health maintenance, disease amelioration, help recover after being sick or having surgery, and even for slowing down the aging process. The sexual reproduction of *O. sinensis* is controlled by the 3D structures of specific functional parts of the paired MAT1-1-1 and MAT1-2-1 proteins. This study examined the structures of paired MAT1-1-1 and MAT1-2-1 protein variants from 20 *O. sinensis* strains. Differentially altered functional parts of some of the mating proteins are shorter than others with divergent replacement of amino acids, resulting in altered 3D structures of the proteins. Together with findings from previous and present studies, the paired mating proteins with divergent alterations of 3D structures prove that *O. sinensis* cannot reproduce on its own (self-fertilize). Instead, *O. sinensis* is "self-sterile" and its sexual reproduction requires mating partner(s). This new understanding of how *O. sinensis* reproduces sexually provides a new perspective to learn about *O. sinensis* and the *C. sinensis* insect–fungal complex and will guide future academic and applied studies.

Abstract

The MATα_HMGbox domain in MAT1-1-1 protein and the HMG-box_ROX1-like domain in MAT1-2-1 protein play essential roles in DNA binding and the transcriptional regulation that control sexual reproduction in *Ophiocordyceps sinensis*. Previous studies have documented differential occurrences, differential transcription, and alternative splicing of the *MAT1-1-1* and *MAT1-2-1* and pheromone receptor genes in *Hirsutella sinensis* (Genotype #1 among the 17 genome-independent *O. sinensis* genotypic fungi). This study further revealed that the DNA-binding domains of the paired MAT1-1-1 and MAT1-2-1 proteins derived from 20 *O. sinensis* strains exhibit heteromorphic tertiary structures, as predicted by AlphaFold 3D structural modeling. The differentially paired mating protein variants, characterized by differential truncations, 1−4 amino acid substitutions at distinct sites, altered hydrophobicity and secondary structures, and heteromorphic tertiary structures indicate divergency in the fungal origins of the mating proteins within the mycologically impure *O. sinensis* strains. These findings support the self-sterility hypothesis for *O. sinensis* and are likely ensure the fidelity and genetic diversity of heterothallic or hybrid reproduction throughout the lifecycle of the *Cordyceps sinensis* insect–fungal complex.

Keywords: MATα_HMGbox domain of MAT1-1-1 protein; HMG-box_ROX1-like domain of MAT1-

2-1 protein; heteromorphic primary, secondary, and tertiary structures of proteins; *Ophiocordyceps sinensis* strain; sexual reproduction; the *Cordyceps sinensis* insect−fungal complex

1. Introduction

Natural *Cordyceps sinensis*, also known as Chinese cordyceps (a non-Latin term) or DōngChóng XiàCǎo in Chinese, is endemic to high-altitude alpine meadows (3,000−5,000 m), which contributes to its natural scarcity. It has a centuries−long history of clinical application in health maintenance, characterized by the dual reinforcement of "*YIN*" and "*YANG*", as well as in disease amelioration, post-illness and postoperative recovery, and antiaging therapy [Zhu *et al*. 1998a, 1998b, 2011; Song *et al*. 2001]. Modern science is increasingly elucidating the complex biochemical mechanisms (*e.g.*, immunomodulation, anti-inflammation, antioxidant activity, organ protection, enhancement of energy-vitality-stamina, etc.) that underpin these traditional applications, attesting to its status as a precious therapeutic resource, albeit the complexity and rarity of this medicinal resource. This natural substance with its unique therapeutic significance embodies a remarkable fusion of nature's ingenuity and traditional medicinal wisdom; however, its high value has led to overharvesting, rendering it a Level-II endangered natural substance [China Ministry of Agriculture and Rural Affairs 2021].

Natural *C. sinensis* comprises the fruiting body of *Ophiocordyceps sinensis* and the remains of a mummified *Hepialidae* moth larva, which retains a thick body wall with numerous bristles, intact larval intestine, head tissues, and fragments of other larval tissues [Ren *et al*. 2013; Zhang *et al*. 2014; Lu & St Leger 2016; Zhu & Li 2017; Li *et al*. 2022, 2023a, 2023d, 2023e]. Dual-fluorescence microscopy examinations have revealed multicellular heterokaryotic microstructures, containing mono-, bi-, tri-, and tetra-nucleate cells [Bushley *et al*. 2013]. Natural *C. sinensis* exhibits apparent genetic heterogeneity, harboring >90 fungal species from at least 37 fungal genera, as well as 17 genome−independent genotypes of *O. sinensis* that differentially co-occur naturally, the abundance of which undergoes dynamic alterations in different compartments during the maturation of natural *C. sinensis* [Li 1988; Dai *et al*. 1989, 2008; Liu *et al*. 1989; Kinjo & Zang 2001; Jiang & Yao 2003; Chen *et al*. 2004, 2011; Stensrud *et al*. 2005, 2007; Leung *et al*. 2006; Zhu *et al*. 2007, 2010; Yang *et al*. 2008, 2021; Xiao *et al*. 2009; Zhang *et al*. 2010a, 2010b, 2018; Li *et al*. 2013, 2016b, 2020b, 2022, 2023d, 2023e; Barseghyan *et al*. 2011; Gao *et al*. 2012; Ren *et al*. 2013; Meng *et al*. 2015; Xia *et al*. 2015, 2016; Guo *et al*. 2017; Zhu & Li 2017; Zhong *et al*. 2018; Kang *et al*. 2024]. Thus, the Chinese Pharmacopoeia defines this medicinal microecosystem as an insect–fungal complex.

Among the numerous intrinsic fungal species [Jiang & Yao 2003; Zhang *et al*. 2010a, 2010b; Xia *et al*. 2015, 2016; Zhang *et al*. 2018], *Hirsutella sinensis* has been postulated to be the sole anamorph of *O. sinensis* [Wei *et al*. 2006], although this hypothetical claim has not yet met all 4 criteria of Koch's postulates. Ten years later, however, Wei *et al*. [2016] reported a successful artificial cultivation project conducted in an industrial product-oriented setting and identified a species contradiction that arose between the anamorphic inoculants of 3 *H. sinensis* strains (GC-biased Genotype #1 of *O. sinensis*) inoculated on *Hepialidae* moth larvae and the genome−independent, sole teleomorphic AT-biased Genotype #4 of *O. sinensis* identified in the fruiting body of cultivated *C. sinensis*.

Since the 1840s, the Latin name *Cordyceps sinensis* has been indiscriminately applied to both the teleomorph/holomorph of the fungus *C. sinensis* and the wild insect–fungal complex, disregarding Edward Doubleday's original identification in 1842 that the insect portion of the natural material as belonging to *Agrotis* [Lu 2014]. This imprecise usage of the Latin-name has blurred the distinction between the fungus, its host, and the insect−fungal complex, resulting in ambiguity in scientific literature, commercial trade, and governmental regulations worldwide. The fungus was renamed *O. sinensis* in 2007, using *H. sinensis* strain EFCC7287 (GC-biased Genotype #1 of *O. sinensis*) as the nomenclatural reference [Sung *et al*. 2007; Zhang *et al*. 2012; Ren *et al*. 2013; Zhu & Li 2017; Wang *et al*. 2018; Li *et al*. 2022, 2026]. Zhang *et al*. [2013] have improperly implemented the "One Fungus=One Name" nomenclature rule set by the International Mycological Association (IMA) and proposed replacing the anamorphic name *H. sinensis* with the teleomorphic name *O. sinensis* according to the international rule [Hawksworth *et al*. 2011; Taylor 2011]. They interpreted *O. sinensis* as a single fungus to fulfill the prerequisite of "One Fungus" of the international rule. However, their interpretation and implementation of the international rule did not fully consider the 17 genotypic *O. sinensis* fungi, which are genome-independent belonging to distinct fungi and differentially co-exist in the compartments of *C. sinensis* insect-fungal complex. In the present paper, we retain the anamorphic name *H. sinensis* for GC-biased Genotype #1 among the 17 genotypes of *O. sinensis* and refer to the genomically independent Genotypes #2–17 fungi as *O. sinensis* following the taxonomic annotations used in several public depositories, including GenBank and AlphaFold, pending the formal determination and differentiation of the systematic taxonomic positions of these GC- and AT-biased Genotypes #2–17. We also maintain the long-standing use of the traditional name *C. sinensis* for both wild and cultivated insect–fungal complex, because the 2007 renaming of *C. sinensis* to *O. sinensis*, based on *H. sinensis* strain EFCC7287 as the nomenclatural reference, did not address the long-standing indiscriminate application of the Latin name

to the insect–fungal complex [Sung *et al*. 2007; Yao & Zhu 2016; Zhu & Li 2017; Li *et al*. 2022, 2026]. Nevertheless, this imperfect and temporarily necessary practice will likely be revised taxonomically in the future, through the differential use of proprietary and exclusive Latin names for the multiple genome–independent *O. sinensis* genotypic fungi and the insect–fungal complex.

The sexual reproductive behavior of ascomycetes is controlled by transcription factors encoded at the mating-type (*MAT*) locus, which constitute the core regulatory mechanism governing mating type recognition and compatibility, and the development of fruiting bodies and sexual reproductive structures such as ascocarps and ascospores [Turgeon & Yoder 2000; Debuchy & Turgeon 2006; Jones & Bennett 2011; Zheng & Wang 2013; Wilson *et al*. 2015; Sun *et al*. 2019; Ramšak *et al*. 2021; Ramšak & Kück 2022]. Similar to other ascomycetous fungi, the reproductive behavior of *O. sinensis* depends on the synergistic interplay of 2 core mating proteins, the *MAT1-1* and *MAT1-2* idiomorphs. The MAT1-1-1 protein harbors a mating-type alpha high mobility group box (MATα_HMGbox) domain, whereas the MAT1-2-1 protein contains the high mobility group box ROX1-like (HMG-box_ROX1-like) domain [Metin *et al*. 2010; Bushley *et al*. 2013; Hu *et al*. 2013; Zheng & Wang 2013; Li *et al*. 2026]. These 2 conserved domains of mating proteins play essential roles in regulating the expression of genes associated with sexual reproduction. Li *et al*. [2019, 2023c, 2024b] documented differential occurrence, differential transcription, and alternative splicing of mating-type and pheromone receptor genes in the genomes of *H. sinensis* strains, challenging the prior hypothesis that *H. sinensis* undergoes self-fertilization *via* homothallism or pseudohomothallism. Instead, these discoveries support self-sterility consistent with a heterothallic or hybrid lifestyle, indicating necessity of compatible sexual partners to complete the sexual cycle within the *Cordyceps sinensis* insect-fungus complex. Moreover, Li *et al*. [2025, 2026] demonstrated heteromorphic tertiary structures of the DNA-binding domains of full-length MAT1-1-1 and MAT1-2-1 proteins in impure wild-type *C. sinensis* isolates, which might be generated by either GC-biased Genotype #1 (*H. sinensis*) or Genotype #3 of *O. sinensis*. These observations further underscore the complexity underlying heterothallic or hybrid sexual reproduction of *O. sinensis*.

Variations in the DNA-binding domains of the 2 mating proteins, arising from diverse amino acid substitutions, induce conformational changes in their tertiary structures as well as synergistic effects of the 2 idiomorphic mating proteins. These proteins are derived from numerous mycologically impure, wild-type *C. sinensis* isolates [Li *et al*. 2026], which were collected and reported by Prof. Zhang Y-J's group [Zhang *et al*. 2009, 2011, 2013, 2014; Zhang & Zhang 2015]. To clarify the protein structural basis of *O. sinensis* sexual reproduction, the present study further analyzes the N- and/or C-terminal

truncations and amino acid substitutions of the reported by Prof. Yao Y-J's group and their collaborators

[Bushley *et al*. 2013; Li *et al*. 2013] and correlates the changes in the primary, secondary, and tertiary

structures and hydrophobic properties of the MATα_HMGbox and the HMG-box_ROX1-like domains

of the MAT1-1-1 and MAT1-2-1 proteins, respectively. Critically, the present study investigates the

differential pairings of MAT1-1-1 and MAT1-2-1 protein variants generated by the *O. sinensis* strains

and the possible heterospecific fungal origins within these heterogeneous *O. sinensis* strains.

2. Materials and Methods

2.1   The MAT1-1-1 and MAT1-2-1 protein sequences of the *O. sinensis* strains

The AlphaFold database provides the AI-predicted 3D structural morphs for 20 truncated MAT1-1-

1 proteins (AGW27517–AGW27536) and 5 MAT1-2-1 proteins (AGW27543, AGW27548, AGW27552,

AGW27554, and AGW27555) [Bushley *et al*. 2013]. As reported by Li *et al*. [2013], 11 of the 20 *O.*

sinensis strains were obtained from fresh caterpillar body specimens collected from various *C. sinensis*

production regions on the Qinghai-Tibet Plateau, marked as "TS" in Table 1. The remaining 9 *O. sinensis*

strains were from the cultures of monoascospores of *C. sinensis* insect–fungal complexes that were

collected from a single production region (Maqên, Guoluo, Qinghai Provence of China), marked as "SS"

in Table 1. The MAT1-1-1 protein sequences of different lengths were encoded by the *MAT1-1-1* cDNAs

that were amplified from total RNA extracted from the *O. sinensis* strains using the primer pair of

m1cDNAF3/m1cDNAR5, while the *MAT1-2-1* cDNAs were amplified using the primer pair of *Mat1-*

2F/Mat1-2R (Figures S1–S2). Li *et al*. [2013] deposited the internal transcribed spacer (ITS) sequences

to GenBank for 15 of the 20 *O. sinensis* strains, including 10 homogeneous strains shown in brown in

Table 1 that reportedly harbor only the GC-biased Group-A ITS sequences and 5 other strains shown in

red in Table 1 that were characterized as heterogeneous, containing both GC-biased Group-A and AT-

biased Group-B ITS sequences. The remaining 5 *O. sinensis* strains shown in black in Table 1 lack ITS

sequence information in GenBank.

Table 1. *O. sinensis* strains and their tissue sources [Li et al. 2013], GenBank accession numbers for the internal
transcribed spacer (ITS) nucleic acid sequences and mating proteins, and AlphaFold UniProt codes for the
truncated and full-length MAT1-1-1 and MAT1-2-1 proteins derived from the *O. sinensis* strains.

O. sinensis strain code	Source of strain	GenBank accession number				AlphaFold UniProt code for the mating protein	
		ITS nucleotide sequence		Protein sequence			
		GC-biased Group-A	AT-biased Group-B	MAT1-1-1	MAT1-2-1	MAT1-1-1	MAT1-2-1
CS71-1219	SS	JQ900151	—	AGW27534	AGW27554	U3NE87 *	U3NEA9
CS91-1291	TS	JQ900166	—	AGW27523	AGW27543	U3N9T9 ℙ	U3N9W0

CS68-2-1229	SS	JQ900158	—	AGW27528	AGW27548	U3N9T9 ℙ	U3N9W5 ⁂
CS37-295	TS	JQ900169	—	AGW27519	AGW27539	U3N9T9 ℙ	D7F2E9 ♦
CS18-266	TS	JQ900168	—	AGW27520	AGW27540	U3N9T9 ℙ	—
CS560-961	TS	JQ900143	—	AGW27522	AGW27542	U3N6U0	D7F2E3
CS71-1218	SS	JQ900150	—	AGW27533	AGW27553	U3N6U8 †	U3N9X0
CS76-1284	TS	JQ900159	—	AGW27525	AGW27545	U3N919	—
CS561-964	TS	JQ900144	—	AGW27526	AGW27546	U3N7G5	—
CS25-273	TS	JQ900167	—	AGW27527	AGW27547	U3N6U4	—
CS71-1220	SS	JQ900152	JQ900175	AGW27535	AGW27555	U3N6U8 †	U3N9W5 ⁂
CS6-251	TS	JQ900163	JQ900180	AGW27517	AGW27537	U3N9T9 ℙ	U3N6V5
CS34-291	TS	JQ900162	JQ900179	AGW27524	AGW27544	U3N9T9 ℙ	—
CS70-1208	SS	JQ900146	JQ900172	AGW27529	AGW27549	U3NE79 ‡	—
CS68-2-1228	SS	JQ900157	JQ900178	AGW27536	AGW27556	U3N7H7	—
CS68-5-1216	SS	—	—	AGW27532	AGW27552	U3N6U8 †	U3N6W6
CS26-277	TS	—	—	AGW27521	AGW27541	U3N9T9 ℙ	D7F2E9 ♦
CS36-1294	TS	—	—	AGW27518	AGW27538	U3N9T9 ℙ	D7F2E9 ♦
CS70-1211	SS	—	—	AGW27530	AGW27550	U3NE79 ‡	—
CS70-1212	SS	—	—	AGW27531	AGW27551	U3NE87 *	—

Note: The *O. sinensis* strains in brown have ITS sequences of Group A; those in red have ITS sequences for both Groups A and B; and those in black do not have ITS sequences in GenBank [Bushley *et al*. 2013; Li *et al*. 2013]. "TS" indicates that the *O. sinensis* strains were isolated from the caterpillar body specimens. "SS" indicates that the *O. sinensis* strains were from the cultures of monoascospores that were collected from mature *C. sinensis* insect–fungal complexes. The ITS sequences in dark blue belong to GC-biased Group A (Genotype #1) of *O. sinensis* and those in light blue belong to AT-biased Group B (Genotype #17) of *O. sinensis* (Tables S1−S2) [Li *et al*. 2013, 2022]. The GenBank protein accession numbers in green refer to the full-length MAT1-2-1 protein sequences, while the remaining protein accession numbers in pink refer to the truncated mating proteins. The AlphaFold codes with the same suffix symbol (*, ℙ, †, ‡, ⁂, or ♦) belong to the same morphs of AlphaFold-predicted 3D structures. Those without a suffix symbol indicate the unique 3D structural morphs.

2.2 Alignment of the DNA-binding domain sequences of the mating proteins

The amino acid sequences of the MATα_HMGbox domains of the MAT1-1-1 proteins and the HMG-box_ROX1-like domains of the MAT1-2-1 proteins of the *O. sinensis* strains were aligned using the GenBank Blastp program (https://blast.ncbi.nlm.nih.gov/ (Bethesda, MD, USA), accessed from 18 June 2025 to 31 January 2026).

2.3 Amino acid properties and scale analysis

The amino acid sequences of the DNA-binding domains of the mating proteins were scaled on the basis of the general properties of their side chains (https://web.expasy.org/protscale/; Basel, Switzerland) (Table S3), accessed from 18 May 2025 to 20 October 2025 [Kyte & Doolittle 1982; Deleage & Roux 1987; Gasteiger *et al*. 2005; Peters & Elofsson 2014; Simm *et al*. 2016; Li *et al*. 2024b, 2025, 2026]. The MATα_HMGbox domain of MAT1-1 proteins (amino acids 51→225 of the reference sequence

AGW27560 derived from the *H. sinensis* strain CS68-2-1229 [Bushley *et al*. 2013]) and the HMG-box_ROX1-like domain of MAT1-2-1 proteins (amino acids 127→197 of the reference sequence AEH27625 [Zhang *et al*. 2009]), with additional 9 amino acid residues extending upstream and downstream of the domains to ensure comprehensive coverage, were plotted sequentially with a window size of 9 amino acid residues using the linear weight variation model of the ExPASy ProtScale algorithms [Kyte & Doolittle 1982; Deleage & Roux 1987; Gasteiger *et al*. 2005; Li *et al*. 2024b, 2025, 2026] to generate ExPASy ProtScale plots to predict the potential changes in hydrophobicity property and secondary structures of proteins related to α-helices, β-sheets, β-turns, and coils. The resulting topological structures and waveforms shown in the ProtScale plots were compared to exploring changes in the hydrophobicity and 2D structural features of the DNA-binding domains of the mating proteins.

2.4   AlphaFold-based prediction of the tertiary structures of the mating proteins

For heteromorphic stereostructure analysis in this study, the 3D structures of the MAT1-1-1 and MAT1-2-1 proteins from *O. sinensis* strains were predicted computationally from their amino acid sequences utilizing the artificial intelligence (AI)-driven machine learning technology AlphaFold (https://alphafold.com/, Cambridgeshire, UK; accessed from 18 June 2025 to 20 October 2025) [Jumper *et al*. 2021; Tunyasuvunakool *et al*. 2021; David *et al*. 2022; Monzon *et al*. 2022; Rettie *et al*. 2023; Xu *et al*. 2023b; Abramson *et al*. 2024; Varadi *et al*. 2024; Wroblewski & Kmiecik 2024; Li *et al*. 2025; Yang *et al*. 2025].

The AlphaFold database provides per-residue model confidence, predicts scores between 0 and 100 in the partial local distance difference test (pLDDT), and assigns a per-residue score to each individual residue [Jumper *et al*. 2021; Tunyasuvunakool *et al*. 2021; David *et al*. 2022; Monzon *et al*. 2022; Xu *et al*. 2023b; Abramson *et al*. 2024; Varadi *et al*. 2024]. To visually represent confidence levels, the 3D structure uses a color-coding scheme based on the pLDDT score: residues with very high confidence (pLDDT>90) are shown in dark blue, those with high confidence (90>pLDDT>70) appear in light blue, residues with low confidence (70>pLDDT>50) are shown in yellow, and residues with very low confidence (pLDDT<50) are shown in orange [Mariani *et al*. 2013; Wroblewski & Kmiecik 2024; Li *et al*. 2025, 2026].

The main reasons for the low-confidence in AlphaFold predictions include the following: (1) Insufficient supporting data, as the AlphaFold-based prediction rely on the quality of multiple sequence alignment (MSA). If few homologous sequences are available in a certain region of the MSA, the model will lack sufficient evolutionary information to infer the structure, resulting in low confidence. (2) Some

protein regions are inherently flexible under physiological conditions, such as the activation domains of

transcription factors, which do not have a fixed 3D structure, leading to low prediction confidence

assigned by AlphaFold. (3) In terms of special structural regions, such as small-molecule binding sites

and artificial linkers in fused proteins, the AlphaFold system may have limited prediction power, leading

to insufficient confidence in the prediction of complex structures. Thus, the low-confidence regions in

AlphaFold predictions are informative flags for highlighting challenges of prediction and should not be

interpreted with excessive confidence.

2.5 Correlations of the primary, secondary, and tertiary structures of the DNA-binding domains of the paired MAT1-1-1 and MAT1-2-1 proteins

Among various AlphaFold 3D structural heteromorphs, the tertiary structural changes around the

mutation sites within the MATα_HMGbox domains of MAT1-1-1 proteins or the HMG-box_ROX1-like

domains of MAT1-2-1 proteins are locally magnified to visualize the subtle 3D variations. Primary

sequence variations in the DNA-binding domains of the mating proteins, such as peptide chain

truncations and amino acid replacements, are correlated with topological structural and waveform

changes in hydrophobicity, α-helix, β-sheet, β-turn, and coil presented in the ExPASy ProtScale plots,

and the locally magnified 3D structures around the mutation sites. The variant MAT1-1-1 and MAT1-2-

1 proteins produced by the *O. sinensis* strains are paired to explore diverse fungal origins.

3. Results

3.1 Primary structures of the truncated MAT1-1-1 proteins in *O. sinensis* strains

As aligned with the representative full-length MAT1-1-1 protein AGW27560 derived from the *H.*

sinensis strain CS68-2-1229 [Bushley *et al*. 2013], the 20 MAT1-1-1 proteins are differentially truncated

at both N- and C-termini (Figure 1). These truncated proteins of varying lengths were encoded by the

cDNAs amplified using the same pair of primers (the forward primer *m1cDNAF3* and the reverse

complementary primer *m1cDNAR5* shown in Figure S1 [Bushley *et al*. 2013]) from the *MAT1-1-1* genes

of *O. sinensis* strains. As represented by the full-length reference protein AGW27560 [Bushley *et al*.

2013], the MATα_HMGbox domain of the MAT1-1-1 proteins locates at amino acids 51→225 (shown

in pink in Figures 1 and S1). The MATα_HMGbox domains of the 20 MAT1-1-1 proteins shown in pink

in Figure 1 are truncated at N-termini and contributed to 9 AlphaFold 3D structure morphs (Figure 2,

Table 1).

247
248 **Figure 1**. Alignment of the sequences of the query MAT1-1-1 protein AGW27560 derived from *O. sinensis* strain

CS68-2-1229 [Bushley et al. 2013] and the 9 truncated MAT1-1-1 proteins (AGW27528, AGW27522, AGW27525, AGW27526, AGW27527, AGW27535, AGW27530, AGW27536, and AGW27534 [Bushley et al. 2013]) that are truncated at both the N- and C-termini. These truncated proteins represent the 20 MAT1-1-1 proteins under 9 AlphaFold 3D structural models (U3N9T9, U3N6U0, U3N919, U3N7G5, U3N6U4, U3N6U8, U3NE79, U3N7H7, and U3NE87). The MATα_HMGbox domain (amino acids 51→225 in the AGW27560 sequence) are shown in pink and those outside the domain are shown in blue. The amino acid substitutions are shown in green. The hyphens indicate identical amino acids, and the spaces denote unmatched sequence gaps.

Figure 2. Sequence distribution of the N-terminally truncated MATα_HMGbox domains of 20 MAT1-1-1 proteins derived from *O. sinensis* strains (Table 1). The sequences are aligned with that of the full-length MATα_HMGbox domain (51→225) of the query MAT1-1-1 protein AGW27560 derived from the *H. sinensis* strain CS68-2-1229.

3.2   Primary structures of the truncated MAT1-2-1 proteins in *O. sinensis* strains

The AlphaFold database lists 5 MAT1-2-1 proteins, which were produced by *O. sinensis* strains and are truncated at the C-termini (Figure 3). The truncated MAT1-2-1 proteins with varying lengths are encoded by the *MAT1-2-1* cDNAs amplified using the same pair of primers (the forward primer *Mat1-2F* and the reverse complementary primer *Mat1-2R* shown in Figure S2 [Bushley *et al*. 2013]) from the total RNA extracted from various *O. sinensis* strains. The full-length authentic MAT1-2-1 protein AEH27625 under AlphaFold code U3N942derived from the *H. sinensis* strain CS2 is taken as the alignment reference, which contains an HMG-box_ROX1-like domain at amino acids 127→197 shown in pink in Figures 3 and S2. As aligned with the HMG-box_ROX1-like domain sequence of the query protein AEH27625 [Zhang *et al*. 2009], the domain sequences of the AMT1-2-1 proteins derived from the 5 *O. sinensis* strains are truncated at the C-termini and contain a Y-to-H substitution in each of the

truncated proteins, contributing to diverse 3D structures under 4 AlphaFold codes (Figure 4). The MAT1-

2-1 protein AGW27554 contains an additional Q-to-T substitution shown in Figure 3.

Figure 3. Alignment of the sequences of 5 truncated AMT1-2-1 proteins derived from the *O. sinensis* strains and
representing query full-length MAT1-2-1 protein AEH27625 of the *H. sinensis* strain CS2 under AlphaFold code
D7F2E9. The amino acids in the HMG-box_ROX1-like domain (amino acids 127→197 in the AEH27625
sequence) are shown in pink and those outside the domain are shown in blue. The amino acid substitutions are
shown in green. The hyphens indicate identical amino acids, and the spaces denote unmatched sequence gaps.

Figure 4. Sequence distribution of the C-terminally truncated HMG-box_ROX1-like domains of 5 truncated
MAT1-2-1 proteins derived from the *O. sinensis* strains (*cf*. Table 1). The sequences are aligned with the HMG-
box_ROX1-like domain (127→197) of the full-length MAT1-2-1 protein AEH27625 of *H. sinensis* strain CS2.

3.3 Heteromorphic tertiary structures of the MATα_HMGbox domains of the truncated MAT1-1-1 proteins derived from *O. sinensis* strains

Section 3.3 reports the correlation analyses between the changes in the hydrophobicity and
structures of the MATα_HMGbox domains of truncated MAT1-1-1 proteins under 9 AlphaFold codes

(Table 1, Figures 1−2) derived from various *O. sinensis* strains, compared to the reference MAT1-1-1

proteins (under the AlphaFold code U3N942) represented by AGW27560 that was derived from the *O.*

sinensis strain CS68-2-1229 (Table S4) [Bushley et al. 2013]. The MATα_HMGbox domain sequences

(amino acid residues 51→225) were extended by 9 amino acids both upstream and downstream of the

domains because of the setting of the sequential ExPASy ProtScale plotting for hydropathy, α-helices,

β-sheets, β-turns, and coils at a window size of 9 amino acid residues (*cf*. Section 2.3).

Figure 5 compares the hydrophobicity property and the structures of the MATα_HMGbox domain

of 2 groups of the truncated MAT1-1-1 protein: (1) AGW27522 (under the AlphaFold code U3N6U0)

derived from the *O. sinensis* strain CS560-961 and (2) the truncated proteins AGW27517, AGW27518,

AGW27519, AGW27520, AGW27521, AGW27523, AGW27524, and AGW27528 (under the

AlphaFold code U3N9T9) derived from the *O. sinensis* strains CS6-251, CS36-1294, CS37-295, CS18-

266, CS26-277, CS91-1291, CS34-291, and CS68-2-1229, respectively (Table 1) with those of the

MATα_HMGbox domain of the query protein AGW27560 (under the AlphaFold code U3N942) derived

from the *H. sinensis* strain CS68-2-1229 [Bushley *et al.* 2013]. Panel (A) of Figure 5 shows 13-residue

truncations at the N-termini of MATα_HMGbox domains of the 2 groups of subject proteins, which

result in changes in topological structure and waveform in the ExPASy ProtScale plots for hydropathy,

α-helices, β-sheets, β-turns, and coils, as illustrated in Panels (B–F), respectively. Compared to the 3D

structure of the full-length protein AGW27560 as the comparison reference shown in Panel (G), where

the core structure of 3 α-helices within the MATα_HMGbox domain form a functional L-shape

stereostructure, Panels (H–I) illustrate that the truncations significantly altered the 3D structures of the

proteins under the heteromorphic AlphaFold codes U3N6U0 and U3N9T9, respectively. The truncation

regions of the variant proteins are located upstream of the core structure of 3 α-helices and may have

steric auxiliary influence on stabilizing the hydrophobic core structures of the MATα_HMGbox domains

[Baxevanis *et al*. 1995; Thapar 2015].

Panel (A): Alignment of amino acid sequences of the MATα_HMGbox domains

```
AGW27560    42    ADVLTPVSTAAASRATRQTKEASCDRAKRPLNAFMAFRSYYLKLFPDVQQKTASGFLTT    100
AGW27522    1     ------------------------------------    37
AGW27517    1     ------------------------------------    37
AGW27518    1     ------------------------------------    37
AGW27519    1     ------------------------------------    37
AGW27520    1     ------------------------------------    37
AGW27521    1     ------------------------------------    37
AGW27523    1     ------------------------------------    37
AGW27524    1     ------------------------------------    37
AGW27528    1     ------------------------------------    37

AGW27560    101   LWHKDPFRNKWALIAKVYSFVRDQIGKDKVSLSYFMSLACPTMTIIEPAAYLNALGWCVQ    160
AGW27522    38    ---------------------------------------------------------    97
AGW27517    38    ---------------------------------------------------------    97
AGW27518    38    ---------------------------------------------------------    97
AGW27519    38    ---------------------------------------------------------    97
AGW27520    38    ---------------------------------------------------------    97
AGW27521    38    ---------------------------------------------------------    97
AGW27523    38    ---------------------------------------------------------    97
AGW27524    38    ---------------------------------------------------------    97
AGW27528    38    ---------------------------------------------------------    97

AGW27560    161   DDDAGSQKLFQDESSANLDQSSLLSAEYPSTEIELLSALVNIGYFPDHGADLVERMGSSH    220
AGW27522    98    ---------------------------------------------------------    157
AGW27517    98    ---------------------------------------------------------    157
AGW27518    98    ---------------------------------------------------------    157
AGW27519    98    ---------------------------------------------------------    157
AGW27520    98    ---------------------------------------------------------    157
AGW27521    98    ---------------------------------------------------------    157
AGW27523    98    ---------------------------------------------------------    157
AGW27524    98    ---------------------------------------------------------    157
AGW27528    98    ---------------------------------------------------------    157

AGW27560    221   SGIMAPRAANCTPP    234
AGW27522    158   --------------    171
AGW27517    158   --------------    171
AGW27518    158   --------------    171
AGW27519    158   --------------    171
AGW27520    158   --------------    171
AGW27521    158   --------------    171
AGW27523    158   --------------    171
AGW27524    158   --------------    171
AGW27528    158   --------------    171
```

Panel (B): Hydropathy   Panel (C): α-Helices   Panel (D): β-Sheets

AGW27560

AGW27560

AGW27560

AGW27522

AGW27517, AGW27518, AGW27519,   AGW27520, AGW27521, AGW27523,   AGW27524, AGW27528

Panel (E): β-Turns

AGW27560

AGW27522
AGW27517, AGW27518, AGW27519,
AGW27524, AGW27528

Panel (F): Coils

AGW27560

AGW27520, AGW27521, AGW27523,

Panel (G): AGW27560 (U3N942); Average pLDDT: 69.25

SCDR

SCDR

Panel (H): AGW27522 (U3N6U0); Average pLDDT: 70.69

SCDR

SCDR

Panel (I): AGW27517, AGW27518, AGW27519, AGW27520, AGW27521,
AGW27523, AGW27524, AGW27528 (U3N9T9); Average pLDDT: 72.62

SCDR

SCDR

Figure 5. Correlations of the changes in the hydrophobicity and the primary, secondary, and tertiary structures of the MATα_HMGbox domains of MAT1-1-1 proteins: the reference protein AGW27560 (under the AlphaFold code U3N942) derived from the *H. sinensis* strain CS68-2-1229 and 2 groups of the truncated protein: (1) AGW27522 shown in brown (under the AlphaFold code U3N6U0) derived from the *O. sinensis* strain CS560-961, and (2) the truncated proteins shown in green (under the AlphaFold code U3N9T9) derived from the *O. sinensis* strains CS6-251, CS36-1294, CS37-295, CS18-266, CS26-277, CS91-1291, CS34-291, and CS68-2-1229, respectively. Panel (A) shows an alignment of the amino acid sequences of the MATα_HMGbox domains of the MAT1-1-1 proteins; whereas the hyphens indicate identical amino acid residues, the spaces denote unmatched sequence gaps. The ExPASy ProtScale plots show the changes in hydrophobicity and the 2D structure in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and coils of the proteins, respectively; the open rectangles in blue highlight the N-terminal truncation region in the ExPASy plots. Panels (G–I) show the 3D structures of the entire proteins on the left and the locally magnified structures surrounding the truncation region on the right. The model confidence for the AlphaFold-predicted 3D structures is as follows: ■ very high (pLDDT>90); ■ high (90>pLDDT>70); ■ low (70>pLDDT>50); and ■ very low (pLDDT<50).

Figure 6 compares the hydrophobicity property and structures of the MATα_HMGbox domain of the truncated subject MAT1-1-1 protein AGW27525 (under the AlphaFold code U3N919) derived from the *O. sinensis* strain CS76-1284 (Table 1) with those of the MATα_HMGbox domain of the reference full-length protein AGW27560 (under the AlphaFold code U3N942) derived from the *H. sinensis* strain CS68-2-1229 [Bushley *et al*. 2013]. Panel (A) of Figure 6 shows a 16-residue truncation at the N-terminus of the MATα_HMGbox domain of the proteins AGW27525, which results in altered topological structure and waveform in the ExPASy ProtScale plots for hydropathy, α-helices, β-sheets, β-turns, and coils, as illustrated in Panels (B–F), respectively. Compared to the 3D structure of the reference protein AGW27560 shown in Panel (G), Panel (H) illustrates the truncation variation and significantly altered 3D structure of the protein AGW27525 under the AlphaFold code U3N919. In particular, the N-terminal amino acids 2–4 [AKR shown in Panel (A)] have hydropathy indices of 1.8, -3.9, and -4.5, respectively (Table S3 [*cf*. Kyte & Doolittle 1982]), featuring hydrophilic feature in combination, and form the beginning of the β-sheet structure in the truncated protein AGW27525 in Panel (H); while the amino acids AKR in the full-length protein AGW27560 are not part of the β-sheet structure and the hydrophobic core of 3 critical α-helices within the MATα_HMGbox domain in Panel (G). The truncation region of the protein AGW27525 is located upstream of the core structure of 3 α-helices within the MATα_HMGbox domain and may not directly participate in stabilizing the hydrophobic core of the MATα_HMGbox domain [Baxevanis *et al*. 1995; Thapar 2015].

Panel (A):   Alignment of the amino acid sequences of the MATα_HMGbox domains

```
AGW27560   42   ADVLTPVSTAAASRATRQTKEASCDRAKRPLNAFMAFRSYYLKLFPDVQQKTASGFLTT   100
AGW27525   1    ----------------------------------                            34

AGW27560   101  LWHKDPFRNKWALIAKVYSFVRDQIGKDKVSLSYFMSLACPTMTIIEPAAYLNALGWCVQ   160
AGW27525   35   -----------------------------------------------------------   94

AGW27560   161  DDDAGSQKLFQDESSANLDQSSLLSAEYPSTEIELLSALVNIGYFPDHGADLVERMGSSH   220
AGW27525   95   -----------------------------------------------------------   154

AGW27560   221  SGIMAPRAANCTPP                                                234
AGW27525   155  --------------                                                168
```

Figure 6. Correlations of the changes in hydrophobicity and the primary, secondary, and tertiary structures of the MATα_HMGbox domains of MAT1-1-1 proteins: the reference protein AGW27560 (under the AlphaFold code U3N942) derived from the *H. sinensis* strain CS68-2-1229 and the variant protein AGW27525 (under the AlphaFold code U3N919) derived from the *O. sinensis* strain CS76-1284. Panel (A) shows an alignment of the amino acid sequences of the MATα_HMGbox domains of the MAT1-1-1 proteins; whereas the hyphens indicate identical amino acid residues, the spaces denote unmatched sequence gaps. The ExPASy ProtScale plots show the changes in hydrophobicity and the 2D structure in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and coils of the protein, respectively; the open rectangles in blue highlight the N-terminal truncation region in the ExPASy plots. Panels (G–H) show the 3D structures of the entire proteins on the left and the locally magnified structures surrounding the variation region on the right. The model confidence for the AlphaFold-predicted 3D structures is as follows: ▮ very high (pLDDT>90); ▮ high (90>pLDDT>70); ▮ low (70>pLDDT>50); and ▮ very low (pLDDT<50).

Figure 7 compares the hydrophobicity property and the structures of the MATα_HMGbox domain of the variant MAT1-1-1 protein AGW27526 (under the AlphaFold code U3N7G5) derived from the *O. sinensis* strain CS561-964 (Table 1) with those of the MATα_HMGbox domain of the full-length protein AGW27560 as the comparison reference (under the AlphaFold code U3N942) derived from the *H. sinensis* strain CS68-2-1229 [Bushley *et al*. 2013]. Panel (A) of Figure 7 shows a 13-residue truncation at the N-terminus of the MATα_HMGbox domain of the proteins AGW27526 and amino acid substitutions of C-to-S (cysteine to serine, both with polar neutral side chains; hydropathy indices changed from -3.5 to -0.8; *cf*. Table S3), D-to-S (acidic aspartic acid to serine with polar neutral side chains; hydropathy indices changed from 2.5 to -0.8), R-to-F (basic arginine to aromatic phenylalanine;

hydropathy indices changed from -4.5 to 2.8), and A-to-T (aliphatic alanine to threonine with polar

neutral side chains; hydropathy indices changed from 1.8 to -0.7) [Kyte & Doolittle 1982]. The

truncation and amino acid substitutions result in changes in hydrophobicity and secondary structures of

the variant protein, as illustrated by the altered topological structure and waveform in the ExPASy

ProtScale plots for hydropathy, α-helices, β-sheets, β-turns, and coils, as illustrated in Panels (B−F),

respectively. Compared with the 3D structure of the reference protein AGW27560 shown in Panel (G),

the truncation and replaced amino acids in the variant protein AGW27526 shown in Panel (H) are located

upstream of the 3 core α-helices of the MATα_HMGbox domain, which caused apparent changs in the

tertiary structure of the variant protein under the AlphaFold code U3N919.

Figure 7. Correlations of the changes in the hydrophobicity and the primary, secondary, and tertiary structures of the MATα_HMGbox domains of MAT1-1-1 proteins: the reference protein AGW27560 (under the AlphaFold code U3N942) derived from the *H. sinensis* strain CS68-2-1229 and the variant protein AGW27526 (under the AlphaFold code U3N7G5) derived from the *O. sinensis* strain CS561-964. Panel (A) shows an alignment of the amino acid sequences of the MATα_HMGbox domains of the MAT1-1-1 proteins; whereas the hyphens indicate identical amino acid residues, the spaces denote unmatched sequence gaps. The ExPASy ProtScale plots show the changes in hydrophobicity and the 2D structure in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and coils of the protein, respectively; the open rectangles in blue highlight the N-terminal truncation regions and those in green highlight the altered topological structure and waveform in the ExPASy plots. Panels (G–H) show the 3D structures of the entire proteins on the left and the locally magnified structures surrounding the variation regions on the right. The model confidence for the AlphaFold-predicted 3D structures is as follows: ▮ very high

(pLDDT>90); ▮ high (90>pLDDT>70); ▮ low (70>pLDDT>50); and ▮ very low (pLDDT<50).

Figure 8 compares the hydrophobicity property and the structures of the MATα_HMGbox domain

of the variant MAT1-1-1 protein AGW27527 (under the AlphaFold code U3N6U4) derived from the *O.*

sinensis strain CS25-273 (Table 1) with those of the MATα_HMGbox domain of the reference protein

AGW27560 (under the AlphaFold code U3N942) derived from the *H. sinensis* strain CS68-2-1229

[Bushley *et al*. 2013]. Panel (A) of Figure 8 shows a 27-residue truncation at the N-terminus of the

MATα_HMGbox domain of the protein AGW27527, which results in changes in topological structure

and waveform in the ExPASy ProtScale plots in Panels (B–F) for hydropathy, α-helices, β-sheets, β-

turns, and coils, respectively. Compared to the 3D structure of the reference protein AGW27560 shown

in Panel (G), Panel (H) shows that the truncation that is located within the first of the 3 core α-helices

and the apparent alteration of the tertiary structure of the MATα_HMGbox domain of the truncated

protein AGW27527 under the AlphaFold code U3N6U4.

Panel (A): Alignment of the amino acid sequences of the MATα_HMGbox domains

```
AGW27560   42   ADVLTPVSTAAASRATRQTKEASCDRAKRPLNAFMAFRSYYLKLFPDVQQKTASGFLTT   100
AGW27527    1            ----------------------------------------------------    23

AGW27560  101   LWHKDPFRNKWALIAKVYSFVRDQIGKDKVSLSYFMSLACPTMTIIEPAAYLNALGWCVQ   160
AGW27527   24   -----------------------------------------------------------    83

AGW27560  161   DDDAGSQKLFQDESSANLDQSSLLSAEYPSTEIELLSALVNIGYFPDHGADLVERMGSSH   220
AGW27527   84   -----------------------------------------------------------   143

AGW27560  221   SGIMAPRAANCTPP                                               234
AGW27527  144   --------------                                               157
```

Figure 8. Correlations of the changes in the hydrophobicity and the primary, secondary, and tertiary structures of the MATα_HMGbox domains of MAT1-1-1 proteins: the reference protein AGW27560 (under the AlphaFold code U3N942) derived from the *H. sinensis* strain CS68-2-1229 and the variant protein AGW27527 (under the AlphaFold code U3N6U4) derived from the *O. sinensis* strain CS25-273. Panel (A) shows an alignment of the amino acid sequences of the MATα_HMGbox domains of the MAT1-1-1 proteins; whereas the hyphens indicate identical amino acid residues, the spaces denote unmatched sequence gaps. The ExPASy ProtScale plots) show the changes in hydrophobicity and the 2D structure in Panels (B−F for hydropathy, α-helices, β-sheets, β-turns, and coils of the protein, respectively; the open rectangles in blue highlight the N-terminal truncation regions in the ExPASy plots. Panels (G−H) show the 3D structures of the entire proteins on the left and the locally magnified structures surrounding the truncation region on the right. The model confidence for the AlphaFold-predicted 3D

structures is as follows: █ very high (pLDDT>90); █ high (90>pLDDT>70); █ low (70>pLDDT>50); and █ very
low (pLDDT<50).

Figure 9 compares the hydrophobicity property and the structures of the MATα_HMGbox domain

of the variant MAT1-1-1 proteins AGW27532, AGW27533, and AGW27535 (under the AlphaFold code

U3N6U8) derived from the *O. sinensis* strains CS68-5-1216, CS71-1218, and CS71-1220 (Table 1) with

those of the MATα_HMGbox domain of the reference protein AGW27560 (under the AlphaFold code

U3N942) derived from the *H. sinensis* strain CS68-2-1229 [Bushley *et al*. 2013]. Panel (A) of Figure 9

shows 47-residue truncations at the N-termini of the MATα_HMGbox domains of the variant proteins;

the truncations result in changes in topological structure and waveform in the ExPASy ProtScale plots

in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and coils, respectively. Comparison of the

tertiary structures of the MAT1-1-1 proteins shows the truncations of the variant proteins, as illustrated

in Panels (G–H). The truncations are located within the first of the 3 core α-helices of the

MATα_HMGbox domains, which results in significant changes in the tertiary structures of the variant

proteins AGW27532, AGW27533, and AGW27535 under the AlphaFold code U3N6U8.

Figure 9. Correlations of the changes in the hydrophobicity and the primary, secondary, and tertiary structures of the MATα_HMGbox domains of MAT1-1-1 proteins: the reference protein AGW27560 (under the AlphaFold code U3N942) derived from the *H. sinensis* strain CS68-2-1229 and the variant protein AGW27532, AGW27533, and AGW27535 (under the AlphaFold code U3N6U8) derived from the *O. sinensis* strains CS68-5-1216, CS71-1218, and CS71-1220, respectively. Panel (A) shows an alignment of the amino acid sequences of the MATα_HMGbox domains of the MAT1-1-1 proteins; whereas the hyphens indicate identical amino acid residues, the spaces denote unmatched sequence gaps. The ExPASy ProtScale plots show the changes in hydrophobicity and the 2D structure in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and coils of the protein, respectively; the open rectangles in blue highlight the N-terminal truncation regions in the ExPASy plots. Panels (G–I) show the 3D structures of the entire proteins on the left and the locally magnified structures surrounding the truncation site on the right. The model confidence for the AlphaFold-predicted 3D structures is as follows: very high (pLDDT>90); high (90>pLDDT>70); low (70>pLDDT>50); and very low (pLDDT<50).

Figure 10 compares the hydrophobicity property and the structures of the MATα_HMGbox domains of the variant MAT1-1-1 proteins AGW27529 and AGW27530 (under the AlphaFold code U3NE79) derived from the *O. sinensis* strain CS70-1208 and CS70-1211, respectively (Table 1), with those of the

MATα_HMGbox domain of the reference protein AGW27560 (under the AlphaFold code U3N942) derived from the *H. sinensis* strain CS68-2-1229 [Bushley *et al*. 2013]. Panel (A) of Figure 10 shows 46-residue truncations at the N-termini of the MATα_HMGbox domains of the proteins AGW27529 and AGW27530 and an F-to-V substitutions (aromatic phenylalanine to aliphatic valine; hydropathy indices changed from 2.8 to 4.2 (Table S3 [*cf*. Kyte & Doolittle 1982]) within the MATα_HMGbox domains, indicating increased hydrophobicity. The truncation and increased hydrophobicity result in altered secondary structure, as illustrated by the changes in topological structure and waveform in the ExPASy ProtScale plots for α-helices, β-sheets, β-turns, and coils shown in Panels (B−F), respectively. Compared the 3D structures of the MAT1-1-1 proteins as illustrated in Panels (G−H), the truncation and replaced amino acids in the proteins AGW27529 and AGW27530 are located within the first of the 3 core α-helices of the MATα_HMGbox domains, resulting in significant change in the tertiary structures of the proteins AGW27529 and AGW27530 under the AlphaFold code U3NE79.

Panel (A): Alignment of the amino acid sequences of the MATα_HMGbox domains

```
AGW27560   42   ADVLTPVSTAAASRATRQTKEASCDRAKRPLNAFMAFRSYYLKLFPDVQQKTASGFLTT   100
AGW27529   1                                                           V---     4
AGW27530   1                                                           V---     4

AGW27560  101   LWHKDPFRNKWALIAKVYSFVRDQIGKDKVSLSYFMSLACPTMTIIEPAAYLNALGWCVQ   160
AGW27529    5   -----------------------------------------------------------    64
AGW27530    5   -----------------------------------------------------------    64

AGW27560  161   DDDAGSQKLFQDESSANLDQSSLLSAEYPSTEIELLSALVNIGYFPDHGADLVERMGSSH   220
AGW27529   65   -----------------------------------------------------------   124
AGW27530   65   -----------------------------------------------------------   124

AGW27560  221   SGIMAPRAANCTPP                                               234
AGW27529  125   --------------                                               138
AGW27530  125   --------------                                               138
```

Panel (B): Hydropathy   Panel (C): α-Helices   Panel (D): β-Sheets

464
Figure 10. Correlations of the changes in the hydrophobicity and the primary, secondary, and tertiary structures
of the MATα_HMGbox domains of MAT1-1-1 proteins: the reference protein AGW27560 (under the AlphaFold
code U3N942) derived from the *H. sinensis* strain CS68-2-1229 and the variant proteins AGW27529 and
AGW27530 (under the AlphaFold code U3NE79) derived from the *O. sinensis* strain CS70-1208 and CS70-1211.
Panel (A) shows an alignment of the amino acid sequences of the MATα_HMGbox domains of the MAT1-1-1
proteins; amino acid substitutions are shown in green, whereas the hyphens indicate identical amino acid residues,
the spaces denote unmatched sequence gaps. The ExPASy ProtScale plots show the changes in hydrophobicity
and the 2D structure in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and coils of the protein,
respectively; the open rectangles in blue highlight the N-terminal truncation regions in the ExPASy plots. Panels
G–I show the 3D structures of the entire proteins on the left and the locally magnified structures surrounding the
variation site on the right. The model confidence for the AlphaFold-predicted 3D structures is as follows: ▮ very

high (pLDDT>90); ▮ high (90>pLDDT>70); ▮ low (70>pLDDT>50); and ▮ very low (pLDDT<50).

Figure 11 compares the hydrophobicity property and the structures of the MATα_HMGbox domain

of the variant MAT1-1-1 protein AGW27536 (under the AlphaFold code U3N7H7) derived from the *O.*

sinensis strain CS68-2-1228 (Table 1) with those of the MATα_HMGbox domain of the reference protein

AGW27560 (under the AlphaFold code U3N942) derived from the *H. sinensis* strain CS68-2-1229

[Bushley *et al*. 2013]. Panel (A) of Figure 11 shows a 44-residue truncation at the N-terminus of the

MATα_HMGbox domain of the protein AGW27536 and an S-to-E substitution (serine with polar neutral

side chains to acidic glutamic acid; hydropathy indices changed from -0.8 to -3.5; *cf*. Table S3 [Kyte &

Doolittle 1982]) and a deletion of leucine (hydropathy index 3.8) within the MATα_HMGbox domain,

indicating largely reduced hydrophobicity. The truncation and reduced hydrophobicity result in altered

secondary structure, as illustrated by the changes in topological structure and waveform in the ExPASy

ProtScale plots in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and coils, respectively.

Compared the 3D structures of the MAT1-1-1 proteins as illustrated in Panels (G–H), the truncation and

replaced amino acids and the deletion of the protein AGW27536 are located within the first of the 3 core

α-helices of the MATα_HMGbox domain, resulting in significant change in the tertiary structure of the

proteins AGW27536 under the AlphaFold code U3N7H7.

Figure 11. Correlations of the changes in the hydrophobicity and the primary, secondary, and tertiary structures
of the MATα_HMGbox domains of MAT1-1-1 proteins: the reference protein AGW27560 (under the AlphaFold
code U3N942) derived from the *H. sinensis* strain CS68-2-1229 and the variant protein AGW27536 (under the
AlphaFold code U3N7H7) derived from the *O. sinensis* strain CS68-2-1228. Panel (A) shows an alignment of the
amino acid sequences of the MATα_HMGbox domains of the MAT1-1-1 proteins; whereas the hyphens indicate
identical amino acid residues, the spaces denote unmatched sequence gaps. The ExPASy ProtScale plots show the
changes in hydrophobicity and the 2D structure in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and
coils of the protein, respectively; the open rectangles in blue highlight the N-terminal truncation regions and those
in green highlight the altered topological structure and waveform in the ExPASy plots. Panels (G–H) show the 3D
structures of the entire proteins on the left and the locally magnified structures surrounding the variation site on
the right. The model confidence for the AlphaFold-predicted 3D structures is as follows: ▌ very high (pLDDT>90);

 high (90>pLDDT>70); low (70>pLDDT>50); and very low (pLDDT<50).

Figure 12 compares the hydrophobicity property and the structures of the MATα_HMGbox domains

of the variant MAT1-1-1 proteins AGW27531 and AGW27534 (under the AlphaFold code U3NE87)

derived from the *O. sinensis* strains CS70-1212 and CS71-1219, respectively (*cf.* Table 1) with those of

the MATα_HMGbox domain of the reference protein AGW27560 (under the AlphaFold code U3N942)

derived from the *H. sinensis* strain CS68-2-1229 [Bushley *et al.* 2013]. Panel (A) of Figure 12 shows

60-residue truncations at the N-termini of the MATα_HMGbox domains of the proteins AGW27531 and

AGW27534. The truncations result in altered hydrophobicity and secondary structures, as illustrated by

the changes in topological structure and waveform in the ExPASy ProtScale plots in Panels (B–F) for

hydropathy, α-helices, β-sheets, β-turns, and coils, respectively. Comparison of the 3D structures of the

MAT1-1-1 proteins as illustrated in Panels (G–H) shows the truncations in the proteins AGW27531 and

AGW27534, which are located within the 3 core α-helices of the MATα_HMGbox domains and result

in the significantly altered tertiary structures of the variant proteins under the AlphaFold code U3NE87.

Panel (A): Alignment of the amino acid sequences of the MATα_HMGbox domains

```
AGW27560    42    ADVLTPVSTAAASRATRQTKEASCDRAKRPLNAFMAFRSYYLKLFPDVQQKTASGFLTT    100
AGW27531
AGW27534

AGW27560   101    LWHKDPFRNKWALIAKVYSFVRDQIGKDKVSLSYFMSLACPTMTIIEPAAYLNALGWCVQ    160
AGW27531     1    --------------------------------------------------           50
AGW27534     1    --------------------------------------------------           50

AGW27560   161    DDDAGSQKLFQDESSANLDQSSLLSAEYPSTEIELLSALVNIGYFPDHGADLVERMGSSH    220
AGW27531    51    --------------------------------------------------          110
AGW27534    51    --------------------------------------------------          110

AGW27560   221    SGIMAPRAANCTPP                                               234
AGW27531   111    --------------                                               124
AGW27534   111    --------------                                               124
```

Figure 12. Correlations of the changes in the hydrophobicity and the primary, secondary, and tertiary structures of the MATα_HMGbox domains of MAT1-1-1 proteins: the reference protein AGW27560 (under the AlphaFold code U3N942) derived from the *H. sinensis strain* CS68-2-1229 and the variant proteins AGW27531 and AGW27534 (under the AlphaFold code U3NE87) derived from the *O. sinensis* strains CS70-1212 and CS71-1219, respectively. Panel (A) shows an alignment of the amino acid sequences of the MATα_HMGbox domains of the MAT1-1-1 proteins; whereas the hyphens indicate identical amino acid residues, the spaces denote unmatched sequence gaps. The ExPASy ProtScale plots show the changes in hydrophobicity and the 2D structure in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and coils of the protein, respectively; the open rectangles in blue highlight the N-terminal truncation regions in the ExPASy plots. Panels (G–H) show the 3D structures of the entire proteins on the left and the locally magnified structures surrounding the truncation site on the right. The model confidence for the AlphaFold-predicted 3D structures is as follows: ▮ very high (pLDDT>90); ▮ high (90>pLDDT>70); ▮ low (70>pLDDT>50); and ▮ very low (pLDDT<50).

3.4 Heteromorphic tertiary structures of the HMG-box_ROX1-like domains of the truncated MAT1-2-1 proteins derived from *O. sinensis* strains

Section 3.4 presents the correlation analyses between the changes in the hydrophobicity property and structures of the HMG-box_ROX1-like domains of variant MAT1-2-1 proteins under 4 AlphaFold codes (*cf.* Table 1) derived from various *O. sinensis* strains, compared to the reference full-length MAT1-2-1 proteins (under the AlphaFold code D7F2E9) represented by AEH27625 derived from the *H. sinensis* strain CS2 (*cf.* Table S5) [Zhang et al. 2009]. The HMG-box_ROX1-like domain sequences (amino acid residues 127→197 of the authentic protein AEH27625) were extended both upstream and downstream by 9 external amino acids because the sequential ExPASy ProtScale plotting for hydropathy, α-helices, β-sheets, β-turns, and coils was set up at a window size of 9 amino acids (*cf.* Section 2.5).

Figure 13 compares the hydrophobicity property and structures of the HMG-box_ROX1-like domain of the MAT1-2-1 protein AGW27543 (under the AlphaFold code U3N9W0) derived from the *O. sinensis* strain CS91-1291 (*cf.* Table 1) [Bushley *et al*. 2013] with those of the reference protein AEH27625 (under the AlphaFold code D7F2E9) derived from the *H. sinensis* strain CS2 [Zhang *et al*. 2009]. Panel (A) of Figure 13 shows a 12-residue truncation at the C-terminus of the HMG-box_ROX1-like domain of the protein AGW27543 and an amino acid substitution of Y-to-H (aromatic tyrosine to basic histidine; hydropathy indices changed from -1.3 to -3.2; *cf.* Table S3 [Kyte & Doolittle 1982]). The truncation and amino acid substitution with reduced hydrophobicity resulted in altered secondary structures surrounding the variation sites within the HMG-box_ROX1-like domain of the protein AGW27543, as illustrated by the changes in topological structure and waveform in the ExPASy ProtScale plots in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and coils, respectively. As shown in Panels (G–H) for 3D structures, the truncation and replaced amino acid residues are located in 2 of the 3 core α-helices and result in altered stereostructure of the HMG-box_ROX1-like domain of the protein AGW27543 under the AlphaFold code U3N9W0.

Panel (A): Alignment of amino acid sequences the HMG-box_ROX1-like domains

```
AEH27625 118 APPSPKDIKIPRPPNAYILYRKERHHYVKDANPGITNNEISQILGKAWNMESNDVRQKYK  177
AGW27543 118 -------------------------H---------------------------------  177

AEH27625 178 DMSQQVKQALLEKHPDYQYKPRRPCERRR                                206
AGW27543 178 --------                                                     185
```

Panel (B): Hydropathy

Panel (C): α-Helices

Panel (D): β-Sheets

Panel (E): β-Turns

Panel (F): Coils

Panel (G): AEH27625 (D7F2E9); Average pLDDT: 67.31

Figure 13. Correlations of the changes in hydrophobicity and the primary, secondary, and tertiary structures of the HMG-box_ROX1-like domains of MAT1-2-1 proteins. The reference protein AEH27625 (under the AlphaFold code D7F2E9) is derived from the *H. sinensis* strain CS2, and the variant MAT1-2-1 protein AGW27543 (under the AlphaFold code U3N9W0) is derived from the *O. sinensis* strain CS91-1291. Panel (A) shows an alignment of the amino acid sequences of the HMG-box_ROX1-like domains of the MAT1-2-1 proteins; the amino acid substitution is shown in green, whereas the hyphens indicate identical amino acid residues, the spaces denote unmatched sequence gaps. The ExPASy ProtScale plots show the changes in hydrophobicity and the 2D structure of the protein in Panels (B–F) for the hydropathy, α-helices, β-sheets, β-turns, and coils, respectively; the open rectangles in blue highlight the C-terminal truncation regions in the ExPASy plots and those in green highlight the changes in topological structure and waveform seen in the plots. Panels (G–H) show representations of the 3D structures of the entire proteins on the left; the locally magnified structures at the variation sites are shown on the right. The model confidence for the AlphaFold-predicted 3D structures is as follows: ▮ very high (pLDDT>90); ▮ high (90>pLDDT>70); ▮ low (70>pLDDT>50); and ▮ very low (pLDDT<50).

Figure 14 compares the hydrophobicity property and structures of the HMG-box_ROX1-like domain of the MAT1-2-1 protein AGW27554 (under the AlphaFold code U3NEA9) derived from the *O. sinensis* strain CS71-1219 [Bushley *et al*. 2013] with those of the reference protein AEH27625 (under the AlphaFold code D7F2E9) derived from the *H. sinensis* strain CS2 [Zhang *et al*. 2009]. Panel (A) of Figure 14 shows a 12-residue truncation at the C-terminus of the HMG-box_ROX1-like domain of the protein AGW27554 and amino acid substitutions: Y-to-H (aromatic tyrosine to basic histidine; hydropathy indices changed from -1.3 to -3.2) and Q-to-T (glutamine to threonine, both with polar neutral side chains; hydropathy indices changed from -3.5 to -0.7; *cf*. Table S3 [Kyte & Doolittle 1982]). The truncation mutation and amino acid substitutions with altered hydrophobicity result in changes in the secondary structure surrounding the mutation sites in the HMG-box_ROX1-like domain of the protein AGW27543, as illustrated by the changes in topological structure and waveform in the ExPASy ProtScale plots in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and coils, respectively. As illustrated in Panels (G–H) for 3D structures, the truncated and replaced residues are located in 2 of the 3 core α-helices and result in altered stereostructure of the HMG-box_ROX1-like domain of the protein AGW27554 under the AlphaFold code U3NEA9.

Panel (A): Alignment of amino acid sequences the HMG-box_ROX1-like domains

```
AEH27625 118 APPSPKDIKIPRPPNAYILYRKERHHYVKDANPGITNNEISQILGKAWNMESNDVRQKYK   177
AGW27554 118 --------------------------H--------------------------------   177

AEH27625 178 DMSQQVKQALLEKHPDYQYKPRRPCERRR                                 206
AGW27554 178 -------T                                                     185
```

Panel (B): Hydropathy

Panel (C): α-Helices

Panel (D): β-Sheets

Panel (E): β-Turns

Panel (F): Coils

Panel (G): AEH27625 (D7F2E9); Average pLDDT: 67.31

592
Figure 14. Correlations of the changes in hydrophobicity and the primary, secondary, and tertiary structures of
the HMG-box_ROX1-like domains of MAT1-2-1 proteins. The reference protein AEH27625 (under the
AlphaFold code D7F2E9) is derived from the *H. sinensis* strain CS2, and the variant MAT1-2-1 protein
AGW27554 (under the AlphaFold code U3NEA9) is derived from the *O. sinensis* strain CS71-1219. Panel (A)
shows an alignment of the amino acid sequences of the HMG-box_ROX1-like domains of the MAT1-2-1 proteins;
amino acid substitutions are shown in green, whereas the hyphens indicate identical amino acid residues, the
spaces denote unmatched sequence gaps. The ExPASy ProtScale plots show the changes in hydrophobicity and
the 2D structure of the protein in Panels (B–F) for hydropathy, α-helices, β-sheets, β-turns, and coils, respectively;
the open rectangles in blue highlight the C-terminal truncation regions in the ExPASy plots and those in green
highlight the changes in topological structure and waveform. Panels (G–H) show representations of the 3D
structures of the entire proteins on the left; the locally magnified structures at the variation sites are shown on the
right. The model confidence for the AlphaFold-predicted 3D structures is as follows: ▮ very high (pLDDT>90);
▮ high (90>pLDDT>70); ▮ low (70>pLDDT>50); and ▮ very low (pLDDT<50).

Figure 15 compares the hydrophobicity and structures of the HMG-box_ROX1-like domains of the

MAT1-2-1 proteins AGW27548 and AGW27555 (under the AlphaFold code U3N9W5) derived from

the *O. sinensis* strains CS68-2-1229 and CS71-1220 (*cf*. Table 1) [Bushley *et al*. 2013] with those of the

reference protein AEH27625 (under the AlphaFold code D7F2E9) derived from the *H. sinensis* strain

CS2 [Zhang *et al*. 2009]. Panel (A) of Figure 15 shows 22-residue truncations at the C-termini of HMG-

box_ROX1-like domains of the MAT1-2-1 proteins AGW27548 and AGW27555 and amino acid

substitutions in both variant proteins: Y-to-H (aromatic tyrosine to basic histidine; hydropathy indices

changed from -1.3 to -3.2; *cf*. Table S3 [Kyte & Doolittle 1982]) within the HMG-box_ROX1-like

domains. The truncation and amino acid substitutions caused decreased hydrophobicity and altered

secondary structures surrounding the variation sites in the HMG-box_ROX1-like domains of the MAT1-

2-1 proteins AGW27548 and AGW27555, as illustrated by the changes in topological structure and

waveform in Panels (B–F) for the hydropathy, α-helices, β-sheets, β-turns, and coils, respectively. As

shown in Panels (G–H) for 3D structures, the truncation and replaced residues are located in 2 of the 3

core α-helices, resulting in altered tertiary structures of the HMG-box_ROX1-like domains of the MAT1-

2-1 proteins AGW27548 and AGW27555 under the AlphaFold code U3N9W5.

Panel (A): Alignment of amino acid sequences the HMG-box_ROX1-like domains

```
AEH27625 118 APPSPKDIKIPRPPNAYILYRKERHHYVKDANPGITNNEISQILGKAWNMESNDVRQKYK  177
AGW27548 118 ------------------------H--------------------------------  175
AGW27555 118 ------------------------H--------------------------------  175

AEH27625 178 DMSQQVKQALLEKHPDYQYKPRRPCERRR                                206
AGW27548 175
AGW27555 175
```

623
Figure 15. Correlations of the changes in hydrophobicity and the primary, secondary, and tertiary structures of
the HMG-box_ROX1-like domains of MAT1-2-1 proteins. The reference protein AEH27625 (under the
AlphaFold code D7F2E9) is derived from the *H. sinensis* strain CS2, and the variant MAT1-2-1 proteins
AGW27548 and AGW27555 (under the AlphaFold code U3N9W5) is derived from the *O. sinensis* strains CS68-
2-1229 and CS71-1220, respectively. Panel (A) shows an alignment of the amino acid sequences of the HMG-
box_ROX1-like domains of the MAT1-2-1 proteins; amino acid substitutions are shown in green, whereas the
hyphens indicate identical amino acid residues, the spaces denote unmatched sequence gaps. The ExPASy
ProtScale plots show the changes in hydrophobicity and the 2D structure in Panels (B−F) for the hydropathy, α-
helices, β-sheets, β-turns, and coils of the proteins, respectively; the open rectangles in blue highlight the C-
terminal truncation regions in the ExPASy plots and in red highlight the changes in topological structure and
waveform seen in the plots. Panels (G−H) show representations of the 3D structures of the entire proteins on the
left; the locally magnified structures at the variation sites are shown on the right. The model confidence for the
AlphaFold-predicted 3D structures is as follows: ■ very high (pLDDT>90); ■ high (90>pLDDT>70); ■ low
(70>pLDDT>50); and ■ very low (pLDDT<50).

Figure 16 compares the hydrophobicity property and structures of the HMG-box_ROX1-like

domain of the MAT1-2-1 protein AGW27552 (under the AlphaFold code U3N6W6) derived from the *O.*

sinensis strain CS68-5-1216 [Bushley *et al*. 2013] with those of the reference protein AEH27625 (under

the AlphaFold code D7F2E9) derived from the *H. sinensis* strain CS2 [Zhang *et al*. 2009]. Panel (A) of

Figure 16 shows a 23-residue truncation at the C-terminus of HMG-box_ROX1-like domain of the

MAT1-2-1 protein AGW27552 and an amino acid substitution: Y-to-H (aromatic tyrosine to basic

histidine; hydropathy indices changed from -1.3 to -3.2; *cf*. Table S3 [Kyte & Doolittle 1982]) within the

HMG-box_ROX1-like domain. The replaced amino acid caused decreased hydrophobicity and altered

secondary structures surrounding the mutation sites in the HMG-box_ROX1-like domain of the MAT1-

2-1 protein AGW27552, as illustrated by the changes in topological structure and waveform in the

ExPASy ProtScale plots for α-helices, β-sheets, β-turns, and coils, respectively. As shown in Panels

(G−H) for 3D structures, the truncation and replaced residues are located in 2 of the 3 core α-helices and

completely altered the stereostructure of the HMG-box_ROX1-like domain of the MAT1-2-1 protein

AGW27552 under the AlphaFold code U3N6W6.

Panel (A): Alignment of amino acid sequences the HMG-box_ROX1-like domains

```
AEH27625 118 APPSPKDIKIPRPPNAYILYRKERHHYVKDANPGITNNEISQILGKAWNMESNDVRQKYK  177
AGW27552 118 --------------------------H---------------------------------  174

AEH27625 178 DMSQQVKQALLEKHPDYQYKPRRPCERRR                                206
AGW27552 174
```

Panel (B): Hydropathy

Panel (C): α-Helices

Panel (D): β-Sheets

Panel (E): β-Turns

Panel (F): Coils

Panel (G): AEH27625 (D7F2E9); Average pLDDT: 67.31

Figure 16. Correlations of the changes in hydrophobicity and the primary, secondary, and tertiary structures of the HMG-box_ROX1-like domains of MAT1-2-1 proteins. The reference protein AEH27625 (under the AlphaFold code D7F2E9) is derived from the *H. sinensis* strain CS2, and the variant MAT1-2-1 protein AGW27552 (under the AlphaFold code U3N6W6) is derived from the *O. sinensis* strain CS68-5-1216. Panel (A) shows an alignment of the amino acid sequences of the HMG-box_ROX1-like domains of the MAT1-2-1 proteins; amino acid substitutions are shown in green, whereas the hyphens indicate identical amino acid residues, the spaces denote unmatched sequence gaps. The ExPASy ProtScale plots show the changes in hydrophobicity and the 2D structure of the protein in Panels (B–F) for the hydropathy, α-helices, β-sheets, β-turns, and coils, respectively; the open rectangles in blue highlight the C-terminal truncation regions in the ExPASy plots and in red highlight the changes in topological structure and waveform seen in the plots. Panels (G–H) show representations of the 3D structures of the entire proteins on the left; the locally magnified structures at the sites of substitutions are shown on the right. The model confidence for the AlphaFold-predicted 3D structures is as follows: ■ very high (pLDDT>90); ■ high (90>pLDDT>70); ■ low (70>pLDDT>50); and ■ very low (pLDDT<50).

3.5 Diverse fungal sources of the MAT1-1-1 and MAT1-2-1 protein variants co-produced in pairs by the 20 *O. sinensis* strains

3.5a Genetic heterogeneity of the *O. sinensis* strains

The ITS sequences for phylogeny analysis are available in GenBank for 15 of the 20 *O. sinensis* strains but lack for the remaining 5 strains (*cf*. Table 1). Li *et al*. [2013] reported the design of PCR experiment including the use of 3 pairs of primers (GAF/GAR, GCF/GCR, and 5.8S-F/R) to amplify the ITS sequences and varying annealing temperatures at 53℃, 55℃, or 60℃. Percent identities and sequence coverages between the sequences of the primers and the representative 17 GC- and AT-biased genotypes of *O. sinensis* are listed in Table S6. Under this study setting, different component *O. sinensis* genotypes in the test samples might exhibit varying PCR amplification rates and the amplicons might be preferentially dominated by the most efficiently amplified fungal components. Li *et al*. [2013] also reported direct sequencing of the PCR amplicons. Apparently, this study did not apply strict amplicon cloning-sequencing techniques with a selection of sufficient 30−50 white colonies for comprehensive profiling without bias, nor use MassARRAY single nucleotide polymorphism (SNP) MALDI-TOF mass spectrum genotyping technique with use of multiple extension primers for phylogenetic analysis. Subsequently, the study would have inevitably overlooked some of cooccurring amplicon sequences,

which might be less-abundant within the amplicon pools due to relatively low-amplification efficiency
under the study settings [Li *et al*. 2013, 2016b].

Li *et al*. [2013] reported that 10 of the 15 *O. sinensis* strains are homogenous, containing only GC-
biased *H. sinensis*, which is Genotype #1 of *O. sinensis* (*cf*. Table 1); 5 other strains are heterogeneous,
containing both GC-biased Genotype #1 and AT-biased Genotype #17 of *O. sinensis* strains (*cf*. Tables
S1−S2) [Li *et al*. 2013, 2016b, 2022, 2023d, 2023e; Zhu & Li 2017]. However, because the research
design adopted by Li *et al*. [2013] was not sufficiently refined, one should maintain a high level of
vigilance whether additional fungal species or *O. sinensis* genotypes coexist in the purportedly
homogenous and heterogeneous strains. Because of the complex genetic background of the *O. sinensis*
strains, it cannot be excluded whether the MAT1-1-1 and/or MAT1-2-1 protein variants with differential
truncations and various amino acid substitutions at varying variation sites originate from different
coexisting fungi.

Notably, the remaining 5 *O. sinensis* strains (CS26-277, CS36-1294, CS68-5-1216, CS70-1211, and
CS70-1212, shown in black in Table 1) lack the ITS sequence reports in the GenBank database with
unknown homo/heterogeneity characteristics. Hopefully, the authors [Bushley *et al*. 2013; Li *et al*. 2013,
2025] who possess these strains will upload their ITS sequences to GenBank to ensure comprehensive
and impartial molecular documentation of these strains.

3.5b Differential pairing of the MAT1-1-1 and MAT1-2-1 proteins simultaneously produced by
the 20 *O. sinensis* strains

The above sections present detailed tertiary structural modeling analyses of the 20 variant MAT1-
1-1 and MAT1-2-1 proteins, which were assessed with various AlphaFold codes. These 20 strains
simultaneously produced the paired mating proteins with differential truncations at the N- and/or C-
termini and amino acid substitutions, as summarized in Tables 1−2, including the 10 "homogenous"
strains shown in blown, 5 heterogeneous strains shown in red, and other 5 strains of unknown
homo/heterogeneity characteristics shown in black. These strains can be divided into 2 groups on the
basis of the pairing of 2 mating proteins (Table 2) [Bushley *et al*. 2013; Li *et al*. 2013]: **Group I** consists
of 5 *O. sinensis* strains, which simultaneously produced both truncated MAT1-1-1 and MAT1-2-1
proteins; while **Group II** is composed of remaining 15 *O. sinensis* strains, which simultaneously
produced the truncated MAT1-1-1 proteins and full-length MAT1-2-1 proteins listed in Table S5. The
MAT1-1-1 and MAT1-2-1 proteins that share the same AlphaFold 3D structural morph are marked with
the same suffix symbol (℗, †, *, ⁂, ‡, or ♦) in Table 2.

none
Table 2. Two groups of the 20 *O. sinensis* strains on basis of differential cooccurrence of the MAT1-1-1 and MAT1-2-1 protein in pairs with differential truncations and amino acid substitutions

Group	*O. sinensis* strains	MAT1-1-1 protein					MAT1-2-1 protein			
		GenBank accession #	Number of residues truncated at N-terminus	Number of residues truncated at C-terminus	Amino acid substitution	AlphaFold code	GenBank accession #	Number of residue(s) truncated at C-terminus	Amino acid substitution	AlphaFold code
I	CS71-1219	AGW27534	110	∞		U3NE87 *	AGW27554	65	Y-to-H, Q-to-T	U3NEA9
	CS68-5-1216	AGW27532	97	∞		U3N6U8 †	AGW27552	74	Y-to-H	U3N6W6
	CS71-1220	AGW27535	97	∞		U3N6U8 †	AGW27555	74	Y-to-H	U3N9W5 **
	CS68-2-1229	AGW27528	63	∞		U3N9T9 ‖	AGW27548	74	Y-to-H	U3N9W5 **
	CS91-1291	AGW27523	63	∞		U3N9T9 ‖	AGW27543	64	Y-to-H	U3N9W0
II	CS37-295	AGW27519	63	∞		U3N9T9 ‖	AGW27539			D7F2E9 ◆
	CS26-277	AGW27521	63	∞		U3N9T9 ‖	AGW27541			D7F2E9 ◆
	CS36-1294	AGW27518	63	∞		U3N9T9 ‖	AGW27538			D7F2E9 ◆
	CS6-251	AGW27517	63	∞		U3N9T9 ‖	AGW27537		V-to-I, Y-to-H	U3N6V5
	CS18-266	AGW27520	63	∞		U3N9T9 ‖	AGW27540		Y-to-H	—
	CS34-291	AGW27524	63	∞		U3N9T9 ‖	AGW27544		Y-to-H	—
	CS71-1218	AGW27533	97	∞		U3N6U8 †	AGW27553	1	Y-to-H, K-to-X	U3N9X0
	CS70-1211	AGW27530	97	∞	F-to-V	U3NE79 ‡	AGW27550		Y-to-H	—
	CS70-1208	AGW27529	97	∞	F-to-V	U3NE79 ‡	AGW27549		Y-to-H	—
	CS70-1212	AGW27531	110	∞		U3NE87 *	AGW27551		Y-to-H	—
	CS560-961	AGW27522	63	∞	P-to-L	U3N6U0	AGW27542		Y-to-H	D7F2E3
	CS76-1284	AGW27525	66	10		U3N919	AGW27545		Y-to-H	—
	CS561-964	AGW27526	63	13	CDRA-to-SSFT	U3N7G5	AGW27546		Y-to-H	—
	CS25-273	AGW27527	77	18		U3N6U4	AGW27547		Y-to-H	—
	CS68-2-1228	AGW27536	95	∞	S-to-E, L-residue deletion	U3N7H7	AGW27556		Y-to-H	—

Note: The *O. sinensis* strains in brown are "homogenous" and were detected only GC-biased Genotype #1 of *O. sinensis*; those in red are heterogeneous and were detected both GC-biased Genotype #1 and AT-biased Genotype #17 of *O. sinensis*; and those in black are unknown for their homo/heterogeneity characteristics (*cf*. Tables 1 and S1−S2) [Bushley *et al*. 2013; Li *et al*. 2013, 2022]. The GenBank protein accession numbers in pink fer to truncated MAT1-1-1 and MAT1-2-1 proteins and those in green indicate full-length MAT1-2-1 proteins. **Group I** strains feature as producing paired truncated MAT1-1-1 and MAT1-2-1 proteins. **Group II** strains character as producing truncated MAT1-1-1 proteins that paired with the full-length MAT1-2-1 proteins. The AlphaFold codes with the same suffix symbol (‖, †, *, ⁂, ‡, or ♦) belong to the same AlphaFold-predicted 3D structural morphs. Those without a suffix symbol indicate the unique 3D structural morphs.

In addition to the analyses of the truncated mating proteins in Sections 3.1−3.4 (*cf*. Figures 1−16), the 15 **Group-II** *O. sinensis* strains simultaneously produced the full length MAT1-2-1 proteins shown in Green in Table 2. Three (AGW27538, AGW27539, AGW27541) of the 15 MAT1-2-1 proteins are 100% identical to the authentic MAT1-2-1 proteins represented by protein AEH27625. The remaining 12 MAT1-2-1 proteins (AGW27537, AGW27540, AGW27542, AGW27544−AGW27547, AGW27549−AGW27551, AGW27553, AGW27556) contain 1−2 amino acid substitutions (Table 2, Figure S3). The substitutions are located in the HMG-box_ROX1-like domains of the MAT1-2-1 proteins, except the V-to-I substitution is located upstream of the HMG-box_ROX1-like domain of MAT1-2-1 protein AGW27537 originated from the *O. sinensis* strain CS6-251.

3.5c *O. sinensis* strains simultaneously produced the MAT1-1-1 proteins with the same tertiary structures and the MAT1-2-1 proteins with various tertiary structures

The sequence distribution of the MAT1-1-1 and MAT1-2-1 proteins simultaneously produced in pairs by each of the 20 *O. sinensis* strains are visualized in Figure 17, in which the panels are arranged according to the AlphaFold 3D structural models for the MAT1-1-1 proteins. Panel A of Figure 17 includes 8 *O. sinensis* strains, including 4 purportedly homogenous strains shown in blown in Table 2, 2 heterogeneous strains in red, and 2 strains in black with unknown homo/heterogeneity characteristics. These strains produced the 63-/8-residue truncated MAT1-1-1 proteins at the N-/C-termini, respectively, exhibiting the same tertiary structures under the AlphaFold code U3N9T9 labeled with a suffix symbol "‖" in Table 2 and analyzed in Figure 5. Two purportedly homogenous strains CS68-2-1229 and CS91-1291 fall under **Group I**, which simultaneously produced the MAT1-2-1 proteins AGW27548 and AGW27543 that are 74- and 64-residue truncated at the C-termini (Table 2, Figures 13,15, and 17), resulting in differently altered tertiary structures under AlphaFold codes U3N9W5 and U3N9W0, respectively.

Figure 17. Alignment and sequence distributions of the paired MAT1-1-1 and MAT1-2-1 proteins derived from the *O. sinensis* strains in **Group I** in blue and **Group II** in green (*cf*. Table 2). The MAT1-1-1 protein AGW27560 (AlphaFold code U3N942) derived from the *H. sinensis* strain CS68-2-1229 was used as query MAT1-1-1 sequence and the MAT1-2-1 protein AEH27625 (AlphaFold code D7F2E9) derived from the *H. sinensis* strain CS2 was used as query MAT1-2-1 sequence in the alignment analysis.

The remaining 6 *O. sinensis* strains in Panel A fall under **Group II**, including 2 purportedly homogenous strains, 2 heterogeneous strains, and 2 other strains with unknown homo/heterogeneity characteristics. In addition to the productions of the same truncated MAT1-1-1 proteins under the AlphaFold code U3N9T9 (labeled with suffix symbol "⊩" in Table 2), these 6 strains simultaneously produced the full-length MAT1-2-1 proteins with heteromorphic 3D structures under different AlphaFold codes or lacking AlphaFold 3D structure codes as follows (*cf*. Table 2 and Panel A of Figure 17):

(1) The co-produced full-length MAT1-2-1 proteins AGW27539, AGW27541, and AGW27538 exhibit

the same 3D structure under AlphaFold 3D structural code **D7F2E9,** with a suffix symbol "♦" in Table 2. These MAT1-2-1 proteins are 100% identical to the authentic MAT1-2-1 protein AEH27625 (*cf*. Figure S3, Table S5). These MAT1-2-1 proteins were generated by the "homogenous" *O. sinensis* strain CS37-295, and by the strains CS26-277 and CS36-1294 that lack the ITS sequence records in GenBank (*cf*. Tables 1−2, Panel A of Figure 17).

(2) The co-produced full-length MAT1-2-1 protein AGW27537 has altered tertiary structure under AlphaFold 3D structural code **U3N6V5** (*cf*. Figure 12 of [Li *et al*. 2026] and Table 5) and was generated by the heterogeneous *O. sinensis* strain CS6-251 (*cf*. Tables 1−2, Panel A of Figure 17). This protein shows 99.2% similarity to the authentic MAT1-2-1 protein AEH27625 and contains V-to-I and Y-to-H substitutions, which are located upstream and within the HMG-box_ROX1-like domain, respectively (*cf*. Figure S3 and 17, Table 2).

(3) The co-produced full-length MAT1-2-1 proteins AGW27540 and AGW27544 were generated by the "homogenous" *O. sinensis* strain CS18-266 and the heterogeneous strain CS34-291 and do not have predicted 3D structures in the AlphaFold database (*cf*. Tables 1−2, Panel A of Figure 17). These MAT1-2-1 proteins contain Y-to-H substitutions within the HMG-box_ROX1-like domains, showing 99.6% identity to the authentic MAT1-2-1 protein AEH27625 (*cf*. Figure S3).

Panel B of Figure 17 and Tables 1−2 list the *O. sinensis* strains CS71-1220 and CS68-5-1216 of **Group-I** and the strain CS71-1218 of **Group-II**. Strain CS71-1218 is "homogenous", Strain CS71-1220 is heterogenous, while Strain CS68-5-1216 is unknown about its homo/heterogeneity characteristics [Bushley *et al*. 2013] (*cf*. Tables 1−2). These 3 strains produced the MAT1-1-1 proteins (AGW27532, AGW27533, and AGW27535; *cf*. Figures 9 and 17) have 97-/8-residue truncations at the N-/C-termini, respectively, under the same AlphaFold 3D structural model (U3N6U8 labeled with a suffix symbol "†" in Table 2). The strains simultaneously generated the MAT1-2-1 proteins AGW27555 and AGW27552 74-residue truncated at their C-termini (*cf*. Figures 3 and 15−17) and the 1-residue truncated MAT1-2-1 protein AGW27553 at its C-terminus (*cf*. Figure S3) with distinct tertiary structures under AlphaFold 3D structural codes U3N9W5, U3N6W6, and U3N9X0, respectively. These 3 MAT1-2-1 proteins contain Y-to-H substitutions but the protein AGW27553 contains additional K-to-X substitution within the HMG-box_ROX1-like domain, showing 99.6%, 99.6% and 99.2% similarities, respectively, to the authentic MAT1-2-1 protein AEH27625 (*cf*. Figure S3).

Panel C of Figure 17 and Tables 1−2 list the "homogenous" *O. sinensis* strain CS71-1219 of **Group-I** and the strain CS70-1212 of **Group-II** with unknown homo/heterogeneity characteristics. These 2

strains produced the MAT1-1-1 proteins AGW27534 and AGW27531 having 110-/8-residue truncations

at their N-/C-termini, respectively (*cf.* Figures 12 and 17), under the same AlphaFold 3D structural code

U3NE87 labeled with a suffix symbol "*" in Tables 1−2. Simultaneously, Strain CS71-1219 produced a

MAT1-2-1 protein AGW27554 having a 65-residue truncation at its C terminus and Y-to-H and Q-to-T

substitutions within the HMG-box_ROX1-like domain, resulting in the altered tertiary structure under

the AlphaFold 3D structural code U3NEA9 (*cf.* Figures 14 and 17). Whereas Strain CS70-1212 produced

the full-length MAT1-2-1 protein AGW27551 with a Y-to-H substitution within the HMG-box_ROX1-

like domain but without Q-to-T substitution (*cf.* Figure S3, Table 2). The MAT1-2-1 protein AGW27551

does not have an AlphaFold predicted 3D structure.

Panel D of Figure 17 and Tables 1−2 list 2 **Group-II** strain CS70-1211 and CS70-1208, the former

has unknown homo/heterogeneity characteristics and the latter is heterogeneous. These strains generated

the 97-/8-residue truncated MAT1-1-1 proteins AGW27530 and AGW27529 at their N-/C-termini and

have F-to-V substitutions within the MATα_HMGbox domains. These 2 MAT1-1-1 proteins are under

the same AlphaFold 3D model U3NE79 labeled with a suffix symbol "‡" in Tables 1−2 (*cf.* Figures 10,

and 17). Simultaneously, these strains co-produced the full-length MAT1-2-1 proteins AGW27550 and

AGW27549, which contain Y-to-H substitutions within the HMG-box_ROX1-like domains (*cf.* Tables

1−2; Figures 17 and S3), showing 99.6% sequence identity to the authentic MAT1-2-1 protein

AEH27625. The AlphaFold database does not contain AlphaFold predicted 3D structural model for these

2 MAT1-2-1 proteins.

3.5e *O. sinensis* strains simultaneously produced the MAT1-2-1 proteins with the same tertiary

structures and the MAT1-1-1 proteins with various tertiary structures

In **Group I** shown in Table 2, the "homogenous" strain CS68-2-1229 and the heterogeneous strain

CS71-1220 produced the 74-residue truncated MAT1-2-1 proteins AGW27548 and AGW27555 with Y-

to-H substitutions within the HMG-box_ROX1-like domains (*cf.* Figures 15 and 17), respectively. These

2 proteins share the same tertiary structures under the AlphaFold code U3N9W5 labeled with a suffix

symbol "⁑" in Tables 1−2. Simultaneously. these strains produced the MAT1-1-1 proteins AGW27528

and AGW27535, which are differentially truncated with deletions of 63/8 and 97/8 residues at the N-

termini of the proteins and have distinct 3D structures under AlphaFold codes U3N9T9 and U3N6U8

labeled with a suffix symbol "℘"or "†" in Tables 1−2 that were fall differently under Panels (A) and (B)

of Figure 17, respectively (*cf.* Figures 5, 9, and 17).

The 5 **Group-II** *O. sinensis* strains (CS560-961, CS76-1284, CS561-964, CS25-273, and CS68-2-

1228; other than those fall under Panels A−D of Figure 17) generated the full-length MAT1-2-1 proteins (AGW27542, AGW27545, AGW27546, AGW27547, and AGW27556; *cf*. Figures 17 and S3), respectively. Although all these full-length MAT1-2-1 proteins contain the same Y-to-H substitutions within the HMG-box_ROX1-like domains, only protein AGW27542 was assigned an AlphaFold 3D structural code D7F2E3. Simultaneously, these strains generated the MAT1-1-1 protein variants (AGW27522, AGW27525, AGW27526, AGW27527, and AGW27536; *cf*. Figures 9,17, Table 2) with the differential truncations (63-/8-residue, 66-/10-residue, 63-/13-residue, 77-/18-residue, and 95-/8-residue) at the N-/C-termini and the various amino acid substitutions under different AlphaFold 3D structural codes (U3N6U0, U3N919, U3N7G5, U3N6U4, and U3N7H7), respectively (*cf*. Table 2, Figures 5−8,11, and 17).

Aggregating the data presented in Section 3.5, the 20 *O. sinensis* strains simultaneously produced the critical MAT1-1-1 and MAT1-2-1 proteins in pairs with differential truncations and amino acid substitutions, indicating the genetic variations within the *MAT* loci in the genomes of impure *O. sinensis* strains, regardless of whether the strains were previously reported as "homogenous" or heterogenous based on the experimental results using less sophisticated study design by Li *et al*. [2013]. Because of no repetitive copies of the *MAT1-1-1* and *MAT1-2-1* genes in the genome of *H. sinensis* [Li *et al*. 2025], the various co-productions of the essential mating proteins in pairs indicate that they are not necessarily the products of GC-biased Genotype #1 *H. sinensis*, instead the possible products of different cooccurring heterospecific or genotypic fungi within the impure *O. sinensis* strains. Table S7 summarizes differentially occurrence and co-occurrence of the MAT1-1-1 and MAT1-2-1 proteins in *O. sinensis* strains, wild-type *C. sinensis* isolates, and *C. sinensis* insect–fungal complexes. Genomically, both the *MAY1-1-1* and *MAT1-2-1* genes are present in the genome assemblies ANOV00000000 and LKHE00000000 of *H. sinensis* strains Co18 and 1229, respectively [Hu *et al*. 2013; Li *et al*. 2016a]. However, the *MAY1-1-1* gene is absent in the genome assemblies LWBQ00000000 and NGJJ00000000 of *H. sinensis* strains ZJB12195 and CC1406-20395, respectively, and the *MAT1-2-1* gene is absent in the genome assembly JAAVMX000000000 of *H. sinensis* strain IOZ07 [Jin *et al*. 2020; Liu *et al*. 2020; Shu *et al*. 2020]. As shown in Figure S1 of [Li *et al*. 2026], the Y-to-M substitutions were found in the MATα_HMGbox domains of MAT1-1-1 proteins encoded by the genome assemblies LKHE00000000 and JAAVMX000000000 of *H. sinensis* strains 1229 and IOZ07, respectively, but not by the genome assembly ANOV00000000 of *H. sinensis* strain Co18 [Hu *et al*. 2013; Li *et al*. 2016a, 2026; Shu *et al*. 2020]. The Y-to-M substitution does not exist in the variant MAT1-1-1 proteins derived from the 20 *O.*

sinensis strains. The genomically encoded MAT1-2-1 proteins contain S-to-A substitutions within the HMG-box_ROX1-like domains (*cf*. Figure S2 of [Li *et al*. 2026]), while the S-to-A substitution does not occur in the MAT1-2-1 proteins derived from the **Group-II** *O. sinensis* strains listed in Table 2. It indicates that the simultaneously detected MAT1-1-1 and MAT1-2-1 proteins in pairs with different truncations and amino acid substitutions shown in Tables 1−2 and Figure 17 might be originated from different cooccurring fungi within the impure *O. sinensis* strains.

4. Discussion

4.1 The altered tertiary structures of the critical DNA-binding domains of mating proteins produced by various *O. sinensis* strains

Previous studies [Li et al. 2023c, 2024b] reported the differential occurrence, differential transcription, and alternative splicing of *MAT1-1-1* and *MAT1-2-1* and pheromone receptor genes in *H. sinensis*, wild-type *C. sinensis* isolates, and *C. sinensis* insect−fungal complex. The full-length MAT1-1-1 and MAT1-2-1 proteins contribute to 15 and 17 stereostructure heteromorphs, belonging to 5 and 5 Bayesian clusters, respectively [Li *et al*. 2025]. Whereas the DNA-binding domains of the MAT1-1-1 and MAT1-2-1 proteins derived from wild-type *C. sinensis* isolates clustered to 5 and 2 branched Bayesian clusters, respectively (*cf*. Tables S6−S9 of [Li *et al*. 2026]). Evidence at genetic, transcriptional, and protein structural levels supports that *O. sinensis* experiences self-sterility and employs a heterothallic hybrid, or even parasexuality strategy to accomplish sexual reproduction during the lifecycle of the *C. sinensis* insect–fungal complex [Li et al. 2023c, 2024b, 2025, 2026]. The research presented in this paper extends the previous investigations by focusing on differential truncations and diverse amino acid substitutions within the MATα_HMGbox domains of MAT1-1-1 proteins and HMG-box_ROX1-like domains of MAT1-2-1 proteins, which are co-produced in pairs from *O. sinensis* strains. The N- and/or C-terminal truncations and 1−4 amino acid substitutions within the DNA-binding domains significantly influence hydrophobicity properties and modify the secondary and tertiary structures of the functional domains of mating proteins—particularly the tertiary structures of hydrophobic cores formed by 3 α-helices inside the DNA-binding domains. These structural variations ultimately modulate the synergistic functions of the key functional domains of mating proteins during the sexual reproduction process of *O. sinensis*.

The HMGbox domains in mating proteins are multifunctional motifs that are central to the transcriptional regulation of mating-related genes, *O. sinensis* sexual reproduction, and host adaptation

within the *C. sinensis* insect–fungal complex [Zheng & Wang 2013; Li *et al.* 2023c, 2024b, 2025, 2026]. In particular, the MATα_HMGbox within the MAT1-1-1 protein represents is inferred to be of conserved descent from the ancestral mating-type HMG lineage *via* conservative continuation model; whereas the HMG-box_ROX1-like domain within the MAT1-2-1 protein arises from evolutionary cooption of a different regulatory lineage (*e.g.*, ROX1-like) through functional convergence—wherein it was repurposed to control mating type specification [Martin *et al.* 2010; Czaja *et al.* 2011; Angelini *et al.* 2016; Poveda-Huertes *et al.* 2020; Ramšak & Kück 2021; Ramšak *et al.* 2022].

The MATα_HMGbox and HMG-box_ROX1-like domains of the MAT1-1-1 and MAT1-2-1 proteins, respectively, are structurally defined DNA-binding domains characterized by a conserved L-shape conformation. This conformation is formed by a hydrophobic core containing 3 asymmetrically arranged α-helices, whose stability is highly dependent on tightly packed, hydrophobic cores located at the interface where these helices converge. The unique tertiary structures of these sequence-specific DNA binders preferentially ensure high−affinity, sequence−dependent binding to short, AT-rich DNA sequences within the promoters of their targets, mating-type-specific genes [Hansen *et al.* 2008; Rajewska *et al.* 2012; Jackson *et al.* 2013; Zheng *et al.* 2013; Kim *et al.* 2015; Ramšak & Kück 2021]. The sequence-specific binding of the HMG-box domains to AT-rich promoter motifs serves as the initial trigger, thereby playing pivotal roles in controlling fungal mating identity and gene expression [Yamamoto *et al.* 1997; Martin et al. 2010; Ait Benkhali et al. 2013; Ramšak *et al.* 2022].

The hydrophobic core, composed of tightly packed amino acid residues, is the foundation of the structures [Kyte & Doolittle 1982; Baxevanis *et al.* 1995; Thapar 2015]. It provides precise conformational stability and constraints that enable the 3 α-helices to form the characteristic L-shaped fold, which creates stable resistance to environmental perturbations that might otherwise unfold or distort the domains. Disruption of this hydrophobic core, whether by amino acid substitutions introducing incompatible residues or by peptide chain truncations removing parts of it, directly impairs structural integrity and abolishes the crucial functions of the domains in DNA binding and the transcription regulation of mating-type-specific genes.

Apparently, the differential truncations and 1−4 amino acid substitutions identified in this study at various mutation sites within the functional domains of paired mating proteins in *O. sinensis* strains directly disrupt key structural folding elements (helices, angles, hydrophobic core) (*cf*. Figures 5−16), and fundamentally impair the DNA-binding specificity in recognizing the target DNA segments. These alterations have significant consequences for the regulation of *O. sinensis* sexual reproduction during

the sexual life of *C. sinensis* insect−fungal complex. Relative to the authentic MAT1-1-1 and MAT1-2-

1 proteins of *O. sinensis* analyzed previously by Li et al. [2025, 2026], the differentially varied DNA-

binding domains clustered into diverse clades/branches in the Bayesian clustering trees [Li *et al*. 2025,

2026].

Although the MATα_HMGbox domain within MAT1-1-1 protein and the HMG-box_ROX1-like

domain within MAT1-2-1 protein both belong structurally to the HMG-box superfamily and share a

conserved L-shaped fold that enables DNA binding and bending, they are functionally distinct in the

context of mating-type regulation in *O. sinensis via* regulating different gene sets, resulting in opposite

regulatory outcomes. The MATα_HMGbox domain of MAT1-1-1 protein activates transcription of genes

encoding MAT1-1-specific receptors or other regulators. Although the HMG-box_ROX1-like domain of

MAT1-2-1 protein binds to specific DNA sequences in the MAT1-2 specific gene promoter encoding the

MAT1-2 specific pheromone receptor, it may overlap differently with the MATα_HMGbox targets and

act through an opposite pathway to counterbalance the activity of MAT1-1-1 protein [Balasubramanian

et al. 1993; Zitomer *et al*. 1997; Zheng *et al*. 2013]. Functionally, the MATα_HMGbox domain primarily

functions as an activator of its cognate mating-type genes (*i.e.,* MAT1-1 genes). In contrast, the HMG-

box_ROX1-like domain acts as a repressor of genes specific to the opposite mating type (MAT1-1 or "a-

specific" genes) and may also serve as an activator for MAT1-2-specific genes. This repressive capacity

represents a key functional hallmark derived from its homology to ROX1 [Kastaniotis & Zitomer 2000].

Thus, despite their structural similarity as DNA-binding and DNA-bending HMG-box domains, the

MATα_HMGbox and HMG-box_ROX1-like domains are functionally specialized to control distinct,

complementary sets of target genes, thereby defining the two different mating types. Their divergent

regulatory outputs (activation *vs*. repression of key target gene sets) constitute the molecular and protein

structural basis of mating-type specification during the sexual reproduction of *O. sinensis* during the

lifecycle of *C. sinensis* insect-fugal complex.

The putative heterodimerization between the MATα_HMGbox domain of MAT1-1-1 protein and

the HMG-box_ROX1-like domain of MAT1-2-1 protein has been proposed as a potential regulatory

mechanism to enforce outbreeding and initiate the sexual cycle in heterothallic fungi [Kües & Casselton

1993; Asante-Owusu *et al*. 1996; Jacobsen *et al*. 2002]. However, because of the lack of direct evidence

in *O. sinensis*, it remains a compelling hypothesis for future research to confirm the occurrence of this

heterodimerization, characterize the interaction interface, and define its precise regulatory targets and

functional impact on mating and fruiting body development in this economically and ecologically

important insect−fungal complex. Nevertheless, the MATα_HMGbox and HMG-box_ROX1-like domains may interact electrostatically and complementarily, providing a plausible physicochemical basis for cooperative synergy in DNA binding. This interaction could coordinate the transition from asexual growth to sexual development distorting target DNA to repress haploid programs and activate mating/meiosis genes of *O. sinensis* during the sexual life of *C. sinensis* insect−fungal complex [Hancock *et al*. 2019]. Despite this, direct experimental validation using protein biochemistry and reproductive physiology approaches is still required to confirm this hypothetical model, clarify its species-specific characteristics, and determine the impact of the protein-level variations on the complementary interactions and mating functional alteration of *O. sinensis* [Li *et al*. 2023c, 2024b, 2025, 2026]—even as the genome-independent, multi-fungal, and multi-genotypical nature of *O. sinensis* remains to be fully elucidated [Kinjo & Zang 2001; Stensrud et al. 2007; Xiao et al. 2009; Zhu et al. 2010; Gao *et al*. 2012; Mao *et al*. 2013; Yao & Zhu 2016; Li et al. 2016b, 2020b, 2022, 2023d, 2023e; Zhu & Li 2017].

4.2 Heterogenous fungal sources of the MAT1-1-1 and MAT1-2-1 proteins simultaneously produced in pairs by the impure *O. sinensis* strains

4.2a Phylogenetic heterogeneity of *O. sinensis* strains

Bushley *et al*. [2013] employed a dual-fluorescence microscopy technique to characterize the multicellular heterokaryotic microstructures of *C. sinensis* ascospores and hyphae, which contain mono-, bi-, tri-, and tetra-nucleate cellular configurations. Zhang & Zhang [2015] subsequently commented on these heterokaryotic microstructures and expressed doubts regarding whether binucleate (as well as trinucleate and tetra-nucleate) cells might inconsistently harbor homogenous or heterogeneous hereditary substances. Molecular biological observations by Li *et al*. [2013, 2016b, 2023d, 2023e], which identified multiple fungal components in the ascospores and ascocarps of *C. sinensis* insect-fungal complex and in "pure" *H. sinensis* strains, provided indisputable evidence supporting the hypothesis proposed by Zhang & Zhang [2015] that the multicellular heterokaryotic cells contain heterogeneous hereditary substances.

Important culture-dependent evidence was reported by Li *et al*. [2013], who deposited the ITS sequences of 15 of the 20 *O. sinensis* strains in GenBank. Among these 15 *O. sinensis* strains, 10 were identified as "homogenous" *O. sinensis* strains, reportedly harboring only the GC-biased Genotype #1. The remaining 5 strains were characterized as heterogeneous, containing both GC-biased Genotype #1,

and AT-biased Genotype #17. For this heterogeneity phenomenon, Li *et al*. [2013, 2017, 2020c] proposed the hypothesis that the AT-biased sequences represent "ITS pseudogenes" or "rRNA pseudogenes" within the genome of GC-biased *H. sinensis*, which are generated through Repeat-Induced Point (RIP) mutation in a single genome. However, subsequent studies have demonstrated that the sequences of Genotypes #2−17 of *O. sinensis* do not reside in the genome of *H. sinensis* but are genomically independent and belong to distinct fungal lineages. These findings directly invalidate the hypotheses that the "RIP" mutation induced the AT-biased "ITS pseudogene" or "rRNA pseudogenes" in the genome of GC-biased *H. sinensis* [Li *et al*. 2016a, 2022, 2023d, 2023e, 2024a; Zhu & Li 2017]. As analyzed in Section 3.5a, unfortunately, the research methodology employed by Li et al. (2013) was less sophisticated. This technical limitation led to an imbalance in the amplification efficiency of coexisting fungal sequences, with only the sequence(s) exhibiting the highest PCR amplification efficiency being captured. which precluded the comprehensive detection of all coexisting fungal taxa in the samples. Notably, the different abundance of amplicons (high *vs*. low)primarily reflects the level of PCR amplification efficiency and velocity under specific research settings and should not be equated with the natural occurrence ratio of coexisting fungi in the tested samples. The natural occurrence ratio of coexisting fungal taxa can be quantified using specialized techniques such as Southern blotting [Zhu *et al*. 2010].

Following the research clue of the inconsistent co-occurrence of GC-biased Genotype #1 and AT-biased Genotype #17 in the cultures of monoascospores identified in the culture-independent study (Tables 1 and S1−S2) [Li *et al*. 2013], Li *et al*. [2016b, 2023d, 2023e] further adopted an exhaustive culture-independent approach, including multi-primers, "touch-down" PCR protocols and selection and sequencing of multiple white colonies, as well as MALDI-TOF SNP mass spectrum genotyping approaches with use of multiple extension primers, to exhaustively investigate the comprehensive fungal composition within *C. sinensis* ascospores and the "pure" *H. sinensis* strains Hs-Ma-4, Hs7.7, Hs7.8, and Hs-Qing2 (gifts from renowned mycologist Prof. Gou YL). The authors reported that *C. sinensis* ascospores and "pure" *H. sinensis* strains harbor cooccurrence of genomically independent fungal taxa, including genome−independent, psychrophilic GC-biased Genotypes #1 and #14, AT-biased Genotypes #5−6 and #16 of *O. sinensis*, as well as the mesophilic *Paecilomyces hepiali* (recently reclassified as *Samsoneilla hepiali* [Wang *et al*. 2020]). These findings clearly indicate that the study by Li *et al*. [2013] overlooked non-culturable fungal components under the particular experimental conditions and coexisting fungal sequences with low PCR amplification efficiency. The origins of paired MAT1-1-1

and MAT1-2-1 proteins with differential truncations and various amino acid substitutions at different mutation sites require further clarification *via* a rigorously designed mycological and molecular protocols by the custodian(s) of the original *O. sinensis* strains.

The paired heteromorphic MAT1-1-1 and MAT1-2-1 proteins are assumed to have distinct fungal origins, derived from the cooccurring GC-biased Genotypes #1 and #3 and AT-biased Genotype #17 of *O. sinensis* in wild-type *C. sinensis* isolates [Li *et al*. 2026] and the *O. sinensis* strains analyzed in this study. Given the limited available information on the diverse coexisting genotypes of *O. sinensis*, the different pairings of the MAT1-1-1 and MAT1-2-1 protein variants that exhibit differential truncations and diverse amino acid substitutions at distinct mutation sites and were co-produced by the differentially co-existing genomically independent fungi in *O. sinensis* strains [Li 1988; Dai *et al*. 1989, 2008; Liu *et al*. 1989; Kinjo & Zang 2001; Jiang & Yao 2003; Chen *et al*. 2004, 2011; Stensrud *et al*. 2005, 2007; Leung *et al*. 2006; Zhu *et al*. 2007, 2010; Yang *et al*. 2008, 2021; Xiao *et al*. 2009; Zhang *et al*. 2010a, 2010b, 2018; Barseghyan *et al*. 2011; Gao *et al*. 2012; Li *et al*. 2013, 2016b, 2020b, 2022, 2023d, 2023e; Ren *et al*. 2013; Meng *et al*. 2015; Xia *et al*. 2015, 2016; Guo *et al*. 2017; Zhu & Li 2017; Zhong *et al*. 2018; Kang *et al*. 2024].

4.2b Differential pairings of the MAT1-1-1 and MAT1-2-1 proteins simultaneously produced by *O. sinensis* strains

As summarized in Table S7, both authentic and variant MAT1-1-1 and MAT1-2-1 proteins display differential occurrence or co-occurrence in the various *O. sinensis* strains (including GC-biased Genotype #1 *H. sinensis* strains), wild-type *C. sinensis* isolates, and *C. sinensis* insect–fungal complexes. The 20 *O. sinensis* strains analyzed in this study are categorized into two distinct groups based on co-productions of their paired mating proteins: Group I produces the truncated MAT1-1-1 and truncated MAT1-2-1 proteins in pairs, while Group II harbors truncated MAT1-1-1 proteins paired with full-length MAT1-2-1 proteins (*cf*. Tables 2 and S5, Figures 5−17 and S3) [Bushley *et al*. 2013; Li *et al*. 2013, 2025]. Most full-length or truncated MAT1-1-1 and MAT1-2-1 proteins contain diverse amino acid substitutions (*cf*. Figures 1−16 and S3).

When addressing the issue of overlooked, low-abundance co-existing fungal sequences that likely arise from inefficient amplification and sequencing bias associated with suboptimal experimental design, the "homogenous" and heterogeneous cultures of *C. sinensis* monoascospores previously reported by Li et al. [2013] presumably contain a greater diversity of uncharacterized fungal sequences. The actual fungal cooccurrence within the tested strains may be far more complex than what reported by Li *et al*.

[2013]. As analyzed in Sections 3.5 and 4.2, the distinct pairings of the highly divergent MAT1-1-1 and MAT1-2-1 protein variants that were simultaneously and differentially produced by the 20 *O. sinensis* strains present a complex, non-uniform pairing pattern, indicating divergent fungal origins from cooccurring fungal taxa in the samples. Further investigations are therefore warranted, employing unbiased and rigorous culture-independent methodologies to comprehensively characterize coexisting genomically independent genotypic *O. sinensis* and heterospecific fungal taxa [Li *et al*. 2016b, 2023d, 2023e].

4.2c Paradoxical dilemmas regarding highly possible heterogeneity of the prior reported "pure"

***O. sinensis* strains**

The first dilemma concerns the purportedly homogenous strain CS68-2-1229 of **Group I** (dark green and underlined in Figures 2 and 4). Only the GC-biased Genotype #1 was detected in this "pure" culture of *Cordyceps sinensis* monoascospores [Bushley *et al*. 2013; Li *et al*. 2013], although the study design adopted by Li *et al*. [2013] was insufficiently rigorous to comprehensively detect all cooccurring fungal taxa (*cf*. Section 3.5a). Two MAT1-1-1 protein sequences from this strain have been deposited in GenBank: a full-length MAT1-1-1 protein AGW27560 containing 372 amino acids and an N-/C-terminally truncated MAT1-1-1 protein AGW27528 containing 301 amino acids (*cf*. Figure 5). These 2 MAT1-1-1 proteins share 100% sequence identity, yet AGW27528 exhibits only 81% query coverage with 63- and 8-residue truncations at the N- and C-termini, respectively, (*cf*. Figures 17 and S3). This truncation pattern was likely attributable to the use of different primers for amplifying the *MAT1-1-1* gene or its transcript from the total RNA pool in the experiments. This strain also produced the truncated MAT1-2-1 protein AGW27548 (*cf*. Figure 15) containing 175 amino acids, which shares 99.4% sequence identity with the authentic full-length MAT1-2-1 protein and carries a Y-to-H substitution within the HMG-box_ROX1-like domain; notably, it shows only 70% sequence coverage due to a 74-residue truncation at the C-terminus relative to the authentic full-length MAT1-2-1 proteins containing 249 amino acids. These 3 mating protein sequences derived from Strain CS68-2-1229 were all released by GenBank on the same date (28-SEP-2013). The truncated MAT1-1-1 AGW27528 and MAT1-2-1 protein AGW27548 were derived from cDNAs that were amplified from the same batch of total RNA pool extracted from the strain CS68-2-1229 using a primer pair *m1cDNAF3*/*m1cDNAR5* to amplify *MAT1-1-1* cDNA and a primer pair *Mat1-2F*/*Mat1-2R* to amplify *MAT1-2-1* cDNA (*cf*. Figures S1−S2) [Bushley *et al*. 2013; Li *et al*. 2013].

As shown in Figures 2 and 5, the MATα_HMGbox domain (162 amino acids) of the MAT1-1-1

protein AGW27528 is among the longest domains across all 9 truncated MAT1-1-1 proteins. In contrast, the HMG-box_ROX1-like domain (49 amino acids) of the MAT1-2-1 protein AGW27548 (*cf.* Figure 15) is 10 residues shorter than the longest truncated HMG-box_ROX1-like domains (59 amino acids) of the MAT1-2-1 protein AGW27543 (*cf.* Figure 4), and 22 residues shorter than the 71-residue HMG-box_ROX1-like domain of the authentic MAT1-2-1 protein AEH27625 [Li *et al*. 2026]. The truncated MAT1-2-1 protein AGW27548, which harbors the Y-to-H substitution, exhibits altered secondary and tertiary structures (*cf.* Figure 15). Li *et al*. [2025] reported no repetitive copies of the *MAT1-1-1* and *MAT1-2-1* genes in the genome of *H. sinensis* strain. These findings suggest that the MAT1-1-1 protein AGW27528 is most likely encoded by the authentic *MAT1-1-1* gene in the *H. sinensis* genome. However, the MAT1-2-1 protein AGW27548 is presumably encoded by a variated *MAT1-2-1* gene within the *H. sinensis* genome or by an independent cooccurring fungal taxon in the actually "impure" *O. sinensis* strain CS68-2-1229. Thus, the true genetic identity of strain CS68-2-1229 requires re-evaluation: it remains to be determined whether this strain is genuinely homogenous, containing only Genotype #1 *H. sinensis*, or it is indeed heterogeneous, harboring 2 or more fungal taxa like the other heterogeneous strains, as shown in Tables 1−2 and S1−S2.

Another paradoxical purportedly homogenous *O. sinensis* strain CS561-964 (**Group II**) was reported to harbor only GC-biased Genotype #1 *H. sinensis* [Bushley *et al*. 2013; Li *et al*. 2013]. In contrast to the "homogenous" strain CS68-2-1229, Strain CS561-964 produced the MAT1-1-1 protein AGW27526, which features 63-/13-residue truncated at the N-/C-termini alongside 4 amino acid substitutions (CDRA-to-SSEF) within the MATα_HMGbox domains (*cf.* Figures 7 and 17, Tables 1−2). The markedly altered MAT1-1-1 protein AGW27526 is clearly encoded by a variated *MAT1-1-1* gene, either within the single genome of *H. sinensis* or, more plausibly, from a genome-independent, cooccurring fungal taxon. Such a fungal taxon is likely overlooked in the study by Li *et al*. [2013] due to the use of suboptimal experimental protocols (*cf.* Section 3.5a),

Analyses of the purportedly homogenous *O. sinensis* strains raise fundamental questions regarding fungal purification and comprehensive detection of all cooccurring fungal taxa in such cultures. Two key questions remain unresolved: first, whether the purportedly homogenous strains listed in Tables 1−2 and S1−S2 truly harbor only GC-biased Genotype #1, or are indeed heterogeneous and contai multiple fungi? Second, whether the characterized heterogeneous strains only contain the GC-biased Genotype #1 and AT-biased Genotype #17, or an even greater diversity of undetected coexisting fungi? It is reasonable to hypothesize that the differentially truncated MAT1-1-1 and MAT1-2-1 protein variants, which form

paired combinations and exhibit diverse amino acid substitutions at distinct mutation sites, are not necessarily from GC-biased Genotype #1 *H. sinensis*. Instead, these protein variants likely originate from AT-biased Genotype #17 (*cf*. Tables S1−S2), or from other uncharacterized cooccurring fungal taxa that were not detected and reported in the study by Li *et al*. [2013].

4.3 The self-sterility behavior of *O. sinensis* and requirement of sexual partners to accomplish heterothallic or hybrid reproduction

Wei *et al*. [2006] proposed that *H. sinensis* (GC-biased Genotype #1 of *O. sinensis*) represents the sole anamorph of *O. sinensis* teleomorph. Zhang *et al*. [2013] proposed implementation of "One Fungus=One Name" nomenclature principle established by the International Mycological Association [Hawksworth *et al*. 2011; Taylor 2011], yet inappropriately replaced the anamorphic name *H. sinensis* with the teleomorphic name *O. sinensis*. This designation was applied uniformly to all 17 genomically independent genotypes, irrespective of their genetic GC- and AT-biased characteristics. These 17 genomically independent genotypes share a common hereditary ancestor and naturally co-exist in divergent combinational patterns. To support the sole anamorph hypothesis, Bushley *et al*. [2013] and Hu *et al*. [2013] proposed the self-fertilization hypothesis for *O. sinensis* to accomplish homothallic or pseudohomothallic reproduction. However, multiple research groups [Holliday & Cleaver 2008; Stone 2010; Hu *et al*. 2013] have reported unsuccessful attempts to artificially cultivate the fruiting bodies and ascospores of *C. sinensis* insect−fungal complex using pure fungal cultures. Zhang *et al*. [2013] summarized the 40-year history of failed attempts, both published and unpublished , to cultivate the insect−fungal complex in academic research-oriented settings. Qin *et al*. [2018] further elaborated on the key obstacles hindering the artificial cultivation of *C. sinensis* fruiting bodies and ascospores. Evidently, the contentious sole anamorph hypothesis and the concomitant self-fertilization hypothesis have not yet been scientifically validated in academia to meet all 4 criteria of Koch's postulations.

Contradictory to the hypothesis that *H. sinensis* represents the sole anamorph of *O. sinensis* [Wei *et al*. 2006], molecular studies have revealed considerable genetic complexity of *O. sinensis*. This Latin name *O. sinensis* encompasses at least 17 distinct genotypes [Stensrud *et al*. 2007; Xiao *et al*. 2009; Zhu *et al*. 2010; Gao *et al*. 2012; Li *et al*. 2022]. Although these *O. sinensis* genotypes may share a common evolutionary ancestor [Stensrud *et al*. 2007], the sequences of Genotypes #2−17 of *O. sinensis* do not reside in the genome of Genotype #1 *H. sinensis*, indicating that they represent divergent, modern fungal species with independent genomes [Xiao *et al*. 2009; Zhu and Li 2017; Li *et al*. 2022]. A medicinal mycology research group led by Prof. S.P. Wasser reported that *H. sinensis* and *Tolypocladium sinensis*

serve as co-anamorphs of *O. sinensis* [Barseghyan *et al.* 2011], supporting the prior detections of *T. sinensis* from natural *C. sinensis* insect-fungal complexes [Li 1988; Jiang & Yao 2003; Chen *et al.* 2004; Leung *et al.* 2006]. Quandt *et al.* [2014] subsequently proposed taxonomic revisions within the genus *Tolypocladium*, subsequently proposed taxonomic revisions within the genus *Tolypocladium*, reassigning several species to the family Ophiocordycipitaceae. To date, the taxonomic positions of 17 evolutionarily related genotypes as independent fungi, as well as the precise anamorph–teleomorph relationship of *O. sinensis*, remains incompletely resolved. Regarding reproductive modes, early hypotheses proposed homothallic or pseudohomothallic reproduction for *O. sinensis*. However, Zhang and Zhang [2015] reported the differential presence of *MAT1-1-1* and *MAT1-2-1* genes across numerous wild-type *C. sinensis* isolates and advanced a facultative hybridization hypothesis for *O. sinensis*. Li *et al.* [2023c, 2024b] further documented differential occurrence, alternative splicing, and differential translation of the *MAT1-1-1* and *MAT1-2-1* genes and the pheromone receptor genes. More recently, Li *et al.* [2025, 2026] and the present study revealed heteromorphic tertiary structures of the MAT1-1-1 and MAT1-2-1 protein variants, in particular variations within the DNA-binding domains of the mating proteins. These structural observations provide direct protein-level evidence for heterogenous fungal origins of mating-type loci in numerous wild-type *C. sinensis* isolates and *O. sinensis* strains. Notably, Li *et al.* [2025] reported that both *MAT1-1-1* and *MAT1-2-1* genes exist as single-copy genes in the *H. sinensis* genome with no repetitive genomic copies. Collectively, evidence from genetics, transcription, and protein structure invalidates the self-fertilization hypothesis for *O. sinensis* and instead support self-sterility, implying that *O. sinensis* relies on heterothallic mating, hybridization, or even parasexual outcrossing to accomplish sexual reproduction [Bennett & Johnson 2003; Sherwood & Bennett 2009; Seervai *et al.* 2013; Nakamura *et al.* 2019; Samarasinghe *et al.* 2020; Mishra *et al.* 2021; Steensels *et al.* 2021; Li *et al.* 2023c, 2023e, 2024b, 2025, 2026]. Similar self-sterile reproductive strategies have been documented in many other *Ophiocordyceps* species, which require compatible mating partners to undergo sexual reproduction [Yokoyama *et al.* 2005; Lee *et al.* 2010; Zheng *et al.* 2013; Araújo *et al.* 2018; Xu *et al.* 2023a; Zou *et al.* 2024].

Wei *et al.* [2016] reported successful industrial cultivation of *C. sinensis* insect−fungal complexes. A puzzling species paradox was observed between the input and output of the cultivated insect–fungal complexes: 3 GC-biased *H. sinensis* strains were used as inoculants, yet only the teleomorph of AT-biased Genotype #4 was detected in the fruiting bodies of the cultivated *C. sinensis* insect–fungal complexes. The sole teleomorphic AT-biased Genotype #4 identified in the cultivated fruiting bodies by

Wei et al. [2016] had apparently replaced the inoculated GC-biased Genotype #1 strains, resulting in a fundamental species contradiction. The sequences of Genotypes #2−17 of *O. sinensis* are not present in the genome assemblies ANOV00000000, JAAVMX000000000, LKHE00000000, LWBQ00000000, or NGJJ00000000 of the pure *H. sinensis* strains Co18, IOZ07, 1229, ZJB12195, and CC1406-20395, respectively [Stensrud *et al*. 2007; Xiao *et al*. 2009; Gao *et al*. 2012; Hu *et al*. 2013; Li *et al*. 2013, 2016a, 2020b, 2022, 2023d, 2023e; 2024a; Mao *et al*. 2013; Zhu & Li 2017; Jin *et al*. 2020; Liu *et al*. 2020; Shu *et al*. 2020]. Instead, these genotypes belong to distinct fungi with independent genomes. These results invalidate the illusory assumptions proposed by Li et al. [2013, 2017, 2020c] that the so called "ITS pseudogene" or "rRNA pseudogenes" within the genome of *H. sinensis* were generated through "RIP" mutation. Furthermore, genomic or transcriptomic analysis provided no supporting evidence for the unfounded speculations [Li *et al*. 2020b. 2024a]. In contrast to the significantly variable maturation-related dynamics of AT-biased genotypes of *O. sinensis*, GC-biased *O. sinensis* genotypes maintain a relatively constant low abundance in the stromata throughout the entire maturation course of *C. sinensis* insect−fungal complex [Zhu *et al*. 2010; Gao *et al*. 2012]. These findings indicate that the 17 genome−independent genotypes of *O. sinensis*, regardless of being GC-biased or AT-biased, represent distinct fungal species [Xiao *et al*. 2009; Li *et al*. 2022, 2023d, 2023e], further reinforcing the invalidation of the RIP mutation/pseudogenes hypotheses proposed by Li *et al*. [2013, 2017, 2020c]. The successful industrial cultivation of *C. sinensis* insect–fungal complexes, coupled with the apparent species paradox, indicates that the sole anamorph hypothesis for *H. sinensis* proposed by Wei *et al*. [2006] failed to satisfy all 4 criteria of Koch's postulates.

Wei *et al*. [2016] identified AT-biased Genotype #4 of *O. sinensis* as the sole teleomorph in the fruiting body of cultivated insect−fungal complexes. Early studies reported that AT-biased Genotype #4 occurred in the stromata throughout all maturation stages of *C. sinensis* insect−fungal complex, including the stromal fertile portion (SFP). However, its abundance exhibited a marked ontogenetic shift: while Genotype #4 was highly abundant in immature stromata during the asexual growth phase, its levels declined substantially in mature stromata and SFP during the sexual reproduction stage or the transition from asexual growth to sexual reproduction [Zhu *et al*. 2010; Gao *et al*. 2012; Li *et al*. 2020b, 2022, 2023d, 2023e]. Li *et al*. [2020a] subsequently clarified that the inoculant used in the industrial inoculation-cultivation project, which is the cultures of the *C. sinensis* ascospores, rather than pure *H. sinensis* strains initially described by Wei *et al*. [2016].

Regrettably, Li *et al*. [2013] failed to detect AT-biased Genotype #4 from cultures of *C. sinensis*

monoascospores using culture–dependent methods but instead identified GC-biased Genotype #1 and AT-biased Genotype #5 (actually Genotype #17; *cf*. Table S2). Used culture–independent approaches, Li *et al*. [2016b, 2023d, 2023e] employed a comprehensive PCR amplicon cloning-sequencing strategy (with the selection of ≥30 white colonies) and SNP MALDI-TOF mass spectrum genotyping using multiple extension primers. These analyses identified GC-biased Genotypes #1 and #14 and AT-biased Genotypes #5−6 and #16, and *S. hepiali* in *C. sinensis* ascospores, yet again failed to detect AT-biased Genotype #4. Collectively, these findings confirm the absence of Genotype #4 of *O. sinensis* in the *C. sinensis* ascospores [Li *et al*. 2013, 2023d, 2023e]. Consequently, the fungal origin of the sole teleomorphic AT-biased Genotype #4 of *O. sinensis* detected in the fruiting body of cultivated *C. sinensis* insect−fungal complex [Wei *et al*. 2016] remains an unresolved mycological enigma.

Li *et al*. [2026] summarized the evidence related to the industrial success of cultivating the insect−fungal complex and postulated that the technical design for the cultivation project adopts a "fungal impurity" strategy, with the synergistic effect of coexisting fungi potentially being the key to success. The supporting evidence includes: (1) Li *et al*. [2020a] reported that the cultivation inoculant was cultures of *C. sinensis* ascospores, which harbor multiple genotypic and heterospecific fungi [Li *et al*. 2013, 2023d, 2023e]; (2) the *C. sinensis* ascospores do not contain AT-biased Genotype #4 of *O. sinensis* [Li *et al*. 2013, 2020b, 2022, 2023d, 2023e]. It is therefore plausible that the origin of the teleomorphic Genotype #4 that was detected by Wei *et al*. [2016] originated from co-cultures of *C. sinensis* ascospores with stroma and/or SFP containing numerous ascocarps, which indeed contain AT-biased Genotype #4 of *O. sinensis*, along with other genotypic and heterospecific fungi; (3) the industrial cultivation system was supplemented with soil collected from the natural *C. sinensis* production areas on the Qinghai–Tibet Plateau [Wei *et al*. 2016].

To date, the purification and genomic sequencing of GC- and AT-biased Genotypes #2−17 of *O. sinensis* have not been reported. Consequently, genomic and transcriptomic information regarding the significantly divergent *MAT1-1-1* and *MAT1-2-1* genes in the genome−independent Genotypes #2−17 of *O. sinensis*, especially the critical DNA-binding domains of the MAT1-1-1 and MAT1-2-1 proteins, remains lacking. Irrespective of the precise biophysical details of their interaction, whether the paired MAT1-1-1 and MAT1-2-1 proteins form heterodimers or interact *via* complementary electrostatic surfaces, the synergy of paired mating proteins serves as the central and indispensable molecular trigger for mating compatibility and the initiation of sexual reproduction in *O. sinensis*, [Debuchy & Turgeo 2006; Jones & Bennett 2011; Zheng & Wang 2013; Wilson *et al*. 2015; Sun *et al*. 2019]. Molecular

evidence regarding differential occurrence, alternative splicing, and differential transcription of mating-type and pheromone receptor genes in *H. sinensis* unequivocally indicates that *O. sinensis* is self-sterile, relying on heterothallism or hybrid outcrossing to complete sexual reproduction, thereby effectively ruling out homothallism (self-fertilization) as a viable mechanism in this species. [Hu *et al*. 2013; Bushley *et al*. 2013; Zhou *et al*. 2013; Wei *et al*. 2006; Li *et al*. 2019, 2023c, 2024b, 2025]. Furthermore, the discovery of heteromorphic tertiary structures of variant MATα_HMGbox domains (in MAT1-1-1 proteins) and HMG-box_ROX1-like domains (in MAT1-2-1 proteins) provides the protein level validation that (1) self-fertilization is biochemically unfeasible and (2) coordinated heterothallic or hybrid reproduction is enforced through structural complementarity between the paired mating proteins, which are likely generated by divergent heterogeneous *O. sinensis* strains. This self-sterility hypothesis resolves longstanding uncertainty in *O. sinensis* cultivation—namely, the failure of single-species cultures to fruit—and underscores the role of multi-fungal symbiotic synergism in its ecological success throughout the lifecycle of the naturally occurring or artificially cultivated *C. sinensis* insect−fungal complex.

Conclusions

This study presents the protein structural-level evidence for self-sterility in *O. sinensis* and further verifies the heterothallic or hybrid mating pattern of this fungus. For the 20 *O. sinensis* strains examined in this study, the diverse pairings of MAT1-1-1 and MAT1-2-1 protein variants exhibit differential truncations and 1−4 amino acid substitutions in their DNA-binding domains. The primary structural variations caused dramatic changes in the hydrophobicity property of the mating proteins and altered the secondary and tertiary structures of the MATα_HMGbox and HMG-box_ROX1-like domains of the differentially paired MAT1-1-1 and MAT1-2-1 proteins, respectively. The structural heterogeneity of differentially paired mating proteins not only indicates the co-occurrence of heterogeneous fungal origins in impure *O. sinensis* strains but also invalidates the hypothesis of homothallic or pseudohomothallic self-fertilization in *O. sinensis*. Complementing the evidence of differential occurrence, differential transcription, and alternative splicing of the *MAT1-1-1* and *MAT1-2-1* genes and pheromone receptor genes, the protein structural variations of these DNA-binding domains impair the homologous interaction required for homothallic or pseudohomothallic elf-fertilization. Collectively, our findings of differential pairings of the mating protein variants in *O. sinensis* strains clarify the sexual reproduction mechanism of *O. sinensis* and confirm that the selection of suitable heterologous mating partners is an

1257 essential natural strategy for self-sterile *O. sinensis* to complete sexual reproduction during the lifecycle

1258 of *Cordyceps sinensis* insect-fungal complexes on the Qinghai–Tibet Plateau.

Acknowledgments

The authors are grateful to Prof. Mu Zang, Prof. Ru-Qin Dai, and Prof. Zong-Qi Liang for their consultation.

Supplementary Materials: The following supporting information can be downloaded at www.xxxxxx.

Figure S1. Alignment of the DNA sequence of the *MAT1-1-1* gene KC437356 (6530→7748) and the encoded sequence of

the MAT1-1-1 protein AGW27560 (1→ 372) derived from *Ophiocordyceps sinensis* strain CS68-2-1229 [Bushley *et al.* 2013].

Residues in pink indicate the MATα_HMGbox domain of the *MAT1-1-1* gene and MAT1-1-1 protein; those in green indicate

Introns I and II. The underlined residues in red indicate the forward primer m1cDNAF3/*MAT1-1-1*

(AGCGAAGCGACCTCTCAAT) and the reverse complementary primer m1cDNAR5/*MAT1-1-1*

(GGAGGACTCGGAGGAAG) reported by [Bushley *et al.* 2013]. The triplet residues (TAG) in red represent the stop codon.

Figure S2. Alignment of the *MAT1-2-1* gene HM212637 (3241→4097) and the encoded sequence of MAT1-2-1 protein

AEH27625 (1→249) by derived from *Ophiocordyceps sinensis* strain CS2 [Zhang *et al.* 2009]. Residues in pink indicate the

HMG-box_ROX1-like domain of the *MAT1-2-1* gene and MAT1-2-1 proteins; those in green indicate Introns I and II. The

underlined residues in red indicate the forward primer Mat1-2F/*MAT1-2-1* (TGGAATGCGACTGACTACGA) and the

reverse complementary primer Mat1-2R/*MAT1-2-1* (CCAGGAGAGCTTGCTTGACT) reported by [Bushley *et al.* 2013].

The triplet residues (TAA) in red represent the stop codon.

Figure S3. Alignment of the sequences of full-length MAT1-2-1 proteins derived from **Group II** *O. sinensis*

strains listed in Table 2 with that of the authentic MAT1-2-1 protein AEH27625 (under the AlphaFold code

D7F2E9) derived from the *H. sinensis* strain CS2. The residues in pink indicate the HMG-box_ROX1-like

domains of the proteins, those in green indicate the variant amino acids; the hyphens indicate identical amino acid

residues; and the space denotes unmatched sequence gap.

Table S1. The *O. sinensis* strains, GenBank accession numbers for the ITS nucleic acid sequences corresponding to the

GenBank accession numbers for the MAT1-1-1 and MAT1-2-1 proteins, and percent similarities *vs*. GC-biased Genotypes

#1–3, #7–10, and #12 of *O. sinensis*.

Table S2. The *O. sinensis* strains, GenBank accession numbers for the ITS nucleic acid sequences corresponding to the

GenBank accession numbers for the MAT1-1-1 and MAT1-2-1 proteins, and percent similarities *vs*. AT-biased Genotypes

#4–6 and #15–17 of *O. sinensis*.

Table S3. Amino acid scales based on the general chemical characteristics of their side chains for ProtScale analysis

(https://web.expasy.org/protscale/) to predict hydrophobicity and secondary structures (α-helices, β-sheets, β-turns, and coils)

of proteins.

Table S4. GenBank accession numbers (in red in parentheses) for the full-length MAT1-1-1 proteins in the AlphaFold

database under the corresponding AlphaFold UniProt codes [Li *et al*. 2025].

Table S5. GenBank accession numbers (in red) for the full-length MAT1-2-1 proteins of 69 *H. sinensis* strains or wild-type

C. sinensis isolates under the corresponding AlphaFold UniProt codes [Li *et al*. 2025].

Table S6. Similarity of the 3 pairs of primers that were used by Li *et al*. [2013] to amplify the ITS sequences of the *O. sinensis*

strains listed in Table 1 that were isolated either from the caterpillar body specimens that were collected from various

production areas on Qinghai-Tibet Plateau or from the cultures of mono-ascospores that were collected from mature *C.

sinensis* insect–fungal complexes obtained from a single production region (Maqên, Guoluo, Qinghai Provence of China).

Table S7. Cooccurrence or differential occurrence of the MAT1-1-1 and MAT1-2-1 proteins in the O. sinensis strains, wild-

type C. sinensis isolates, and C. sinensis insect–fungal complex that are listed in the GenBank database.

Author Contributions: Conceptualization, XZL, YLL, and JSZ; methodology, JSZ; formal analysis, JSZ; investigation, XZL and JSZ; data curation, XZL and JSZ; writing–original draft preparation, JSZ; writing—review and editing, XZL, YLL, and JSZ; project administration, YLL; funding acquisition, YLL. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by (1) the Chinese Academy of Sciences−People's Government of Qinghai Province on Sanjiangyuan National Park (#LHZX-2022-01); (2) Wild *Ophiocordyceps sinensis* Identification and Application Project (QHRD-2025-004); (3) Process Optimization and Application of Antioxidant Performance of Yushu *Cordyceps sinensis* Extract (2025-NK-P45); and (4) the Shaanxi Key Laboratory of Natural Product & Chemical Biology Open Foundation (SXNPCB 2024003).

Institutional Review Board Statement: Not applicable because this paper is an *in silico* reanalysis of public data.

Informed Consent Statement: Not applicable because this paper is a public bioinformatic data reanalysis.

Data Availability Statement: All sequence and 3D structure data are available in the GenBank and AlphaFold databases.

Conflicts of Interest: The authors declare no conflicts of interest.

[REFERENCES]

1. Ait Benkhali, J.; Coppin, E.; Brun, S.; Peraza-Reyes, L.; Martin, T.; Dixelius, C.; Lazar, N.; van Tilbeurgh, H.; Debuchy, R. A Network of HMG-box Transcription Factors Regulates Sexual Cycle in the Fungus Podospora anserina. *PLoS Genet.* **2013**, *9*, e1003642. https://doi.org/10.1371/journal.pgen.1003642.

2. Angelini, R.M.D.M.; Rotolo, C.; Pollastro, S.; Faretra, F. Molecular analysis of the mating type (*MAT1*) locus in strains of the heterothallic ascomycete *Botrytis cinerea*. *Plant Pathol*. **2016**, *65*, 1221−1400. https://doi.org/10.1111/ppa.12509

3. Araújo JPM, Evans HC, Kepler R, Hughes DP. Zombie-ant fungi across continents: 15 new species and new combinations within *Ophiocordyceps*. I. *Myrmecophilous hirsutelloid* species. *Stud Mycol*. **2018**, *90*, 119−160. doi: 10.1016/j.simyco.2017.12.002

4. Asante-Owusu, R.N.; Banham, A.H.; Böhnert, H.U.; Mellor, E.J.C.; Casselton, L.A. Heterodimerization between two classes of homeodomain proteins in the mushroom *Coprinus cinereus* brings together potential DNA-binding and activation domains. *Gene* **1996**, *172*, 25–31. https://doi.org/10.1016/0378-1119(96)00177-1.

5. Balasubramanian, B.; Lowry, C.V.; Zitomer, R.S. The Rox1 repressor of the *Saccharomyces cerevisiae* hypoxic genes is a specific DNA-binding protein with a high-mobility-group motif. *Mol. Cell Biol.* **1993**, *13*, 6071–6078. https://doi.org/10.1128/mcb.13.10.6071-6078.1993.

6. Barseghyan, G.S.; Holliday, J.C.; Price, T.C.; Madison, L.M.; Wasser, S.P. Growth and cultural-morphological characteristics of vegetative mycelia of medicinal caterpillar fungus *Ophiocordyceps sinensis* G.H. Sung et al. (Ascomycetes) Isolates from Tibetan Plateau (P. R. China). *Intl. J. Med. Mushrooms* **2011**, *13*, 565–581. https://doi.org/10.1615/intjmedmushr.v13.i6.90.

7. Baxevanis, A.D.; Bryant, S.H.; Landsman, D. Homology model building of the HMG-1 box structural domain. *Nucleic Acids Res.* **1995**, *23*, 1019–1029. https://doi.org/10.1093/nar/23.6.1019.

8. Bennett, R.J.; Johnson, A.D. Completion of a parasexual cycle in *Candida albicans* by induced chromosome loss in tetraploid strains. *EMBO J.* **2003**, *22*, 2505–2515. https://doi.org/10.1093/emboj/cdg235.

9.   Bushley, K.E.; Li, Y.; Wang, W.-J.; Wang, X.-L.; Jiao, L.; Spatafora, J.W.; Yao, Y.-J. Isolation of the MAT1-1 mating type
idiomorph and evidence for selfing in the Chinese medicinal fungus *Ophiocordyceps sinensis*. *Fungal Biol.* **2013**, *117*, 599–
610. https://doi.org/10.1016/j.funbio.2013.06.001.

10. Chen, C.-S.; Hseu, R.-S.; Huang, C.-T. Quality Control of *Cordyceps sinensis* Teleomorph, Anamorph, and Its Products,
Chapter 12. In *Quality Control of Herbal Medicines and Related Areas*; Shoyama, Y., Ed.; InTech: Rijeka, Croatia, 2011; pp.
223–238. Available online: www.intechopen.com (accessed on 3 May 2024).

11. Chen, Y.-Q.; Hu, B.; Xu, F.; Zhang, W.; Zhou, H.; Qu, L.-H. Genetic variation of *Cordyceps sinensis*, a fruit-body-producing
entomopathogenic species from different geographical regions in China. *FEMS Microbiol. Lett.* **2004**, *230*, 153–158.
https://doi.org/10.1016/S0378-1097(03)00889-9.

12. China Ministry of Agriculture and Rural Affairs. Announcement (No. 15 of 2021) of National Forestry and Grassland
Administration: List of National Key Protected Wild Plants. 7 September 2021. Available online:
https://m.163.com/dy/article/HHCVOJPU055360T7.html (accessed on 3 May 2025).

13. Czaja, W.; Miller, K.Y.; Miller, B.L. Complex mechanisms regulate developmental expression of the matA (HMG) mating
type gene in homothallic Aspergillus nidulans. *Genetics*. **2011**, *189*, 795–808. doi: 10.1534/genetics.111.131458

14. Dai, R.-Q.; Lan, J.-L.; Chen, W.-H.; Li, X.-M.; Chen, Q.-T.; Shen, C.-Y. Discovery of a new fungus *Paecilomyces hepiali* Chen
& Dai. *Acta Agricult. Univ. Pekin.* **1989**, *15*, 221–224.

15. Dai, R.-Q.; Li, X.-M.; Shao, A.-J.; Lin, S.-F.; Lan, J.-L.; Chen, W.-H.; Shen, C.-Y. Nomenclatural validation of *Paecilomyces*
hepiali. *Mycosystema* **2008**, *27*, 641–644.

16. David, A.; Islam, S.; Tankhilevich, E.; Sternberg, M.J.E. The AlphaFold Database of Protein Structures, A Biologist's Guide.
J. Mol. Biol. **2022**, *434*, 167336. https://doi.org/10.1016/j.jmb.2021.167336.

17. Debuchy, R.; Turgeo, B.G. Mating-Type Structure, Evolution, and Function in Euascomycetes. In *Growth, Differentiation*
and Sexuality; Kües, U., Fischer, R., Eds.; Springer: Berlin/Heidelberg, Germany, 2006; pp. 293–323.

18. Deleage, G.; Roux, B. An algorithm for protein secondary structure prediction based on class prediction. *Protein Eng. Des.*
Sel. **1987**, *1*, 289–294. https://doi.org/10.1093/protein/1.4.289.

19. Du, X.-H.; Wu, D.-M.; Kang, H.; Wang, H.-C.; Xu, N.; Li, T.-T.; Chen, K.-L. Heterothallism and potential hybridization
events inferred for twenty-two yellow morel species. *IMA Fungus* **2020**, *11*, 4. https://doi.org/10.1186/s43008-020-0027-1.

20. Gao L, Li X-H, Zhao J-Q, Lu J-H, Zhao J-G, Zhu J-S. Maturation of *Cordyceps sinensis* associates with alterations of fungal
expressions of multiple *Ophiocordyceps sinensis* mutants with transition and transversion point mutations in stroma of
Cordyceps sinensis. *Beijing Da Xue Xue Bao*. **2012**, *44*, 454–463. doi: 10.3969/j.issn.1671-167X.2012.03.025

21. Gasteiger, E.; Hoogland, C.; Gattiker, A.; Duvaud, S.; Wilkins, M.R.; Appel, R.D.; Bairoch, A. Protein Identification and
Analysis Tools on the ExPASy Server, Chapter 52. In *The Proteomics Protocols Handbook*; Walker, J.M., Ed.; Humana Press:
Totowa, NJ, USA, 2005; pp. 571–607.

22. Guo, M.-Y.; Liu, Y.; Gao, Y.-H.; Jin, T.; Zhang, H.-B.; Zhou, X.-W. Identification and bioactive potential of endogenetic
fungi isolated from medicinal caterpillar fungus *Ophiocordyceps sinensis* from Tibetan Plateau. *Int. J. Agric. Biol.* **2017**, *19*,
307–313. https://doi.org/10.17957/IJAB/15.0281.

23. Hancock, S.P.; Cascio, D.; Johnson, R.C. Cooperative DNA binding by proteins through DNA shape complementarity.
Nucleic Acids Res. **2019**, *47*, 8874–8887. https://doi.org/10.1093/nar/gkz642.

24. Hansen FT, Madsen CK, Nordland AM, Grasser M, Merkle T, Grasser KD. A novel family of plant DNA-binding proteins
containing both HMG-box and AT-rich interaction domains. *Biochemistry*. **2008**, *47*, 13207–13214. doi: 10.1021/bi801772k

25. Hawksworth, D.L.; Crous, P.W.; Redhead, S.A.; Reynolds, D.R.; Samson, R.A.; Seifert, K.A.; Taylor, J.W.; Wingfield, M.J.;
Abaci, Ö.; Aime, C.; et al. The Amsterdam declaration on fungal nomenclature. *IMA Fungus* **2011**, *2*, 105–112.
https://doi.org/10.5598/imafungus.2011.02.01.14.

26. Hénault, M.; Marsit, S.; Charron, G.; Landry, C.R. The effect of hybridization on transposable element accumulation in an
undomesticated fungal species. *eLife* **2020**, *9*, e60474. https://doi.org/10.7554/eLife.60474.

27. Holliday, J.; Cleaver, M. Medicinal value of the caterpillar fungi species of the genus *Cordyceps* (Fr.) Link (Ascomycetes). A review. *Int. J. Med. Mushrooms* **2008**, *10*, 219–234. https://doi.org/10.1615/IntJMedMushr.v10.i3.30.

28. Hu, X.; Zhang, Y.-J.; Xiao, G.-H.; Zheng, P.; Xia, Y.-L.; Zhang, X.-Y.; St Leger, R.J.; Liu, X.-Z.; Wang, C.-S. Genome survey uncovers the secrets of sex and lifestyle in caterpillar fungus. *Chin. Sci. Bull.* **2013**, *58*, 2846–2854. https://doi.org/10.1007/s11434-013-5929-5.

29. Jackson, D.; Lawson, T.; Villafane, R.; Gary, L. Modeling the structure of yeast MATα1, An HMG-Box motif with a C-terminal helical extension. *Open J. Biophys.* **2013**, *3*, 1–12. https://doi.org/10.4236/ojbiphy.2013.31001.

30. Jacobsen, S.; Wittig, M.; Pöggeler, S. Interaction Between Mating-Type Proteins from the Homothallic Fungus Sordaria macrospora. *Curr. Genet.* **2002**, *41*, 150–158. https://doi.org/10.1007/s00294-002-0276-0.

31. Jiang, Y.; Yao, Y.-J. A review for the debating studies on the anamorph of *Cordyceps sinensis*. *Mycosistema* **2003**, *22*, 161–176.

32. Jin, L.-Q.; Xu, Z.-W.; Zhang, B.; Yi, M.; Weng, C.-Y.; Lin, S.; Wu, H.; Qin, X.-T.; Xu, F.; Teng, Y.; Yuan, S.-J.; Liu, Z.-Q.; Zheng, Y.-G. Genome sequencing and analysis of fungus *Hirsutella sinensis* isolated from *Ophiocordyceps sinensis*. *AMB Expr.* **2020**, *10*, 105. https://doi.org/10.1186/s13568-020-01039-x.

33. Jones, S.K.; Bennett, R.J. Fungal mating pheromones: Choreographing the dating game. *Fungal Genet. Biol.* **2011**, *48*, 668–676. https://doi.org/10.1016/j.fgb.2011.04.001.

34. Jumper, J.; Evans, R.; Pritzel, A.; Green, T.; Figurnov, M.; Ronneberger, O.; Tunyasuvunakool, K.; Bates, R.; Žídek, A.; Potapenko, A.; Bridgland, A.; Meyer, C.: Kohl, S,A,A,; Ballard, A.J.; Cowie, A.; Romera-Paredes, B.; Nikolov, S.; Jain, R.; Adler, J.; Back, T.; Petersen, S.; Reiman, D.; Clancy, E.; Zielinski, M.; Steinegger, M.; Pacholska, M.; Berghammer, T.; Bodenstein; S.; Silver, D.; Vinyals, O.; Senior, A.W.; Kavukcuoglu, K.; Kohli, P.; Hassabis, D. Highly accurate protein structure prediction with AlphaFold. *Nature* **2021**, *596*, 583–589. https://doi.org/10.1038/s41586-021-03819-2.

35. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. Highly accurate protein structure prediction with AlphaFold. *Nature*. **2021**, *596*, 583–589. https://doi.org/10.1038/s41586-021-03819-2

36. Kang, Q.; Zhang, J.; Chen, F.; Dong, C.; Qin, Q.; Li, X.; Wang, H.; Zhang, H.; Meng, Q. Unveiling mycoviral diversity in *Ophiocordyceps sinensis* through transcriptome analyses. *Front. Microbiol.* **2024**, *15*, 1493365. https://doi.org/10.3389/fmicb.2024.1493365.

37. Kastaniotis, A.J.; Zitomer, R.S. Oxygen Dependent Repression in Yeast. In *Rox1 Mediated Repression*; Advances in Experimental Medicine and Biology; Springer Nature: Cham, Switzerland, 2000; Volume 475, pp. 185–195. https://doi.org/10.1007/0-306-46825-5_18.

38. Kim, H.-K.; Jo, S.-M.; Kim, G.-Y.; Kim, D.-W.; Kim, Y.-K.; Yun, S.-H. A large-scale functional analysis of putative target genes of mating-type loci provides insight into the regulation of sexual development of the cereal pathogen *Fusarium graminearum*. *PLoS Genet.* **2015**, *11*, e1005486. https://doi.org/10.1371/journal.pgen.1005486.

39. Kinjo, N.; Zang, M. Morphological and phylogenetic studies on *Cordyceps sinensis* distributed in southwestern China. *Mycoscience* **2001**, *42*, 567–574. https://doi.org/10.1007/BF02460956.

40. Kück, U.; Bennett, R.J.; Wang, L.; Dyer, P.S. Editorial, Sexual and Parasexual Reproduction of Human Fungal Pathogens. *Front. Cell Infect. Microbiol.* **2022**, *12*, 934267. https://doi.org/10.3389/fcimb.2022.934267.

41. Kües, U.; Casselton, L.A. The origin of multiple mating types in mushrooms. *J. Cell Sci.* **1993**, *104*, 227–230. https://doi.org/10.1242/jcs.104.2.227.

42. Kyte, J.; Doolittle, R.F. A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* **1982**, *157*, 105–132. https://doi.org/10.1016/0022-2836(82)90515-0.

43. Lee SC, Ni M, Li W, Shertz C, Heitman J. The Evolution of Sex: a Perspective from the Fungal Kingdom. Microbiol *Mol Biol Rev*. **2010**, *74*, 298–340. doi: 10.1128/MMBR.00005-10

44. Leung, P.-H.; Zhang, Q.-X.; Wu, J.-Y. Mycelium cultivation, chemical composition and antitumour activity of a *Tolypocladium* sp. fungus isolated from wild *Cordyceps sinensis*. *J. Appl. Microbiol.* **2006**, *101*, 275–283. https://doi.org/10.1111/j.1365-2672.2006.02930.x.

45. Li, C.-L. A study of *Tolypocladium sinense* C.L. Li. sp. nov. and cyclosporin production. *Acta Mycol. Sin.* **1988**, *7*, 93–98.

46. Li, M.-M.; Zhang, J.-H.; Qin, Q.-L.; Zhang, H.; Li, X.; Wang, H.-T.; Meng, Q. Transcriptome and Metabolome Analyses of Thitarodes xiaojinensis in Response to *Ophiocordyceps sinensis* Infection. *Microorganisms* **2023a**, *11*, 2361. https://doi.org/10.3390/microorganisms11092361.

47. Li, W.-J.; Xia, J.-M.; Li, Q.-P.; Zhang, Z.-Y.; Zhang, W.-W.; Dong, C.-H.; Wei, J.-C.; Liu, X.-Z. Developmental recording of the ghost-moth larvae after ex situ infection by *Ophiocordyceps sinensis*. *Sci. China Life Sci.* **2020a**, *63*, 1093–1095. https://doi.org/10.1007/s11427-020-1686-7. Erratum in Sci. China Life Sci. **2023b**, 66, 892. https://doi.org/10.1007/s11427-022-2232-5.

48. Li, X.; Wang, F.; Liu, Q.; Li, Q.-P.; Qian, Z.-M.; Zhang, X.-L.; Li, K.; Li, W.-J.; Dong, C.-H. Developmental transcriptomics of Chinese cordyceps reveals gene regulatory network and expression profiles of sexual development-related genes. *BMC Genom.* **2019**, *20*, 337. https://doi.org/10.1186/s12864-019-5708-z.

49. Li, X.-Z.; Li, Y.-L.; Wang, Y.-N.; Zhu, J.-S. Translations of mutant repetitive genomic sequences in *Hirsutella sinensis* and changes in secondary structures and functional specifications of the encoded proteins. *Int. J. Mol. Sci.* **2024a**, *25*, 11178. https://doi.org/10.3390/ijms252011178.

50. Li X-Z, Li Y-L, Yao Y-S, Xie W-D, Zhu J-S. Further discussion with Li *et al*. (2013, 2019) regarding the "ITS pseudogene hypothesis" for *Ophiocordyceps sinensis*. *Mol. Phylogenet. Evol*. **2020b**, *146*, 106728. https://doi.org/10.1016/j.ympev.2019.106728

51. Li, X.-Z.; Li, Y.-L.; Zhu, J.-S. Differential transcription of mating-type genes during sexual reproduction of natural *Cordyceps sinensis*. *Chin. J. Chin. Mater. Medica* **2023c**, *48*, 2829–2840. https://doi.org/10.19540/j.cnki.cjcmm.20230213.102.

52. Li, X.-Z.; Li, Y.-L.; Liu, W.; Zhu, J.-S. Altered stereostructures of the DNA-binding domains of variant mating proteins of *Ophiocordyceps sinensis* and the wild insect–fungal complex. *Biol. (Basel).* **2026**, *15*, 186. https://doi.org/10.3390/biology15020186

53. Li, X.-Z.; Li, Y.-L.; Zhu, J.-S. Three-dimensional structural heteromorphs of mating-type proteins in *Hirsutella sinensis* and the natural *Cordyceps sinensis* insect–fungal complex. *J. Fungi.* **2025**, *11*, 244. https://doi.org/10.3390/jof11040244.

54. Li, X.-Z.; Xiao, M.-J.; Li, Y.-L.; Gao, L.; Zhu, J.-S. Mutations and differential transcription of mating-type and pheromone receptor genes in *Hirsutella sinensis* and the natural *Cordyceps sinensis* insect–fungi complex. *Biology* **2024b**, *13*, 632. https://doi.org/10.3390/biology13080632.

55. Li, Y.; Hsiang, T.; Yang, R.-H.; Hu, X.-D.; Wang, K.; Wang, W.-J.; Wang, X.-L.; Jiao, L.; Yao, Y.-J. Comparison of different sequencing and assembly strategies for a repeat-rich fungal genome, *Ophiocordyceps sinensis*. *J. Microbiol. Methods* **2016a**, *128*, 1–6. https://doi.org/10.1016/j.mimet.2016.06.025.

56. Li, Y.; Jiang, L.; Wang, K.; Wu, H.-J.; Yang, R.-H.; Yan, Y.-J.; Bushley, K.E.; Hawksworth, D.L.; Wu, Z.-J.; Yao, Y.-J. RIP mutated ITS genes in populations of *Ophiocordyceps sinensis* and their implications for molecular systematics. *IMA Fungus* **2020c**, *11*, 18.

57. Li, Y.; Jiao, L.; Yao, Y.-J. Non-concerted ITS evolution in fungi, as revealed from the important medicinal fungus *Ophiocordyceps sinensis*. *Mol. Phylogenet. Evol.* **2013**, *68*, 373–379. https://doi.org/10.1016/j.ympev.2013.04.010.

58. Li, Y.; Yang, R.-H.; Jiang, L.; Hu, X.-D.; Wu, Z.-J.; Yao, Y.-J. rRNA Pseudogenes in Filamentous Ascomycetes as Revealed by Genome Data. *G3-Genes Genom. Genet.* **2017**, *7*, 2695–2703. https://doi.org/10.1534/g3.117.044016.

59. Li, Y.-L.; Gao, L.; Yao, Y.-S.; Wu, Z.-M.; Lou, Z.-Q.; Xie, W.-D.; Wu, J.-Y.; Zhu, J.-S. Altered GC- and AT-biased genotypes of *Ophiocordyceps sinensis* in the stromal fertile portions and ascospores of natural *Cordyceps sinensis*. *PLoS ONE.* **2023d**, *18*, e0286865. https://doi.org/10.1371/journal.pone.0286865.

60. Li, Y.-L.; Li, X.-Z.; Yao, Y.-S.; Wu, Z.-M.; Gao, L.; Tan, N.-Z.; Lou, Z.-Q.; Xie, W.-D.; Wu, J.-Y.; Zhu, J.-S. Differential cooccurrence of multiple genotypes of *Ophiocordyceps sinensis* in the stromata, stromal fertile portion (ascocarps) and ascospores of natural *Cordyceps sinensis*. *PLoS ONE* **2023e**, *18*, e0270776. https://doi.org/10.1371/journal.pone.0270776.

61. Li, Y.-L.; Li, X.-Z.; Yao, Y.-S.; Xie, W.-D.; Zhu, J.-S. Molecular identification of *Ophiocordyceps sinensis* genotypes and the indiscriminate use of the Latin name for the multiple genotypes and the natural insect–fungi complex. *Am. J. BioMed. Sci.* **2022**, *14*, 115–135. https://doi.org/10.5099/aj220300115.

62. Li, Y.-L.; Yao, Y.-S.; Zhang, Z.-H.; Xu, H.-F.; Liu, X.; Ma, S.-L.; Wu, Z.-M.; Zhu, J.-S. Synergy of fungal complexes isolated from the intestines of *Hepialus lagii* larvae in increasing infection potency. *J. Fungal Res.* **2016b**, *14*, 96–112.

63. Liu, J.; Guo, L.-N.; Li, Z.-W.; Zhou, Z.; Li, Z.; Li, Q.; Bo, X.-C.; Wang, S.-Q.; Wang, J.-L.; Ma, S.-C.; Zheng J, Yang Y. Genomic analyses reveal evolutionary and geologic context for the plateau fungus *Ophiocordyceps sinensis*. *Clin. Med.* **2020**, *15*, 107–119. https://doi.org/10.1186/s13020-020-00365-3.

64. Liu, X.-J.; Guo, Y.-L.; Yu, Y.-X.; Zeng, W. Isolation and identification of the anamorph of *Cordyceps sinensis* fungus. *Acta Mycol. Sin.* **1989**, *8*, 35–40.

65. Liu, Z.-Q.; Lin, S.; Baker, P.J.; Wu, L.-F.; Wang, X.-R.; Wu, H.; Xu, F.; Wang, H.-Y.; Brathwaite, M.E.; Zheng, Y.-G. Transcriptome sequencing and analysis of the entomopathogenic fungus *Hirsutella sinensis* isolated from *Ophiocordyceps sinensis*. *BMC Genom.* **2015**, *16*, 106–123. https://doi.org/10.1186/s12864-015-1269-y.

66. Lu, D. Western records and studies of the Chinese caterpillar fungus to the beginning of the 20th century. *J. Fungal. Res.* **2014**, *12*, 233–244. https://doi.org/10.13341/j.jfr.2014.0043.

67. Lu, H.-L.; St. Leger, R.J. Chapter Seven—Insect Immunity to Entomopathogenic Fungi. In *Advances in Genetics*; Lovett, B., St. Leger, R.J., Eds.; Academic Press: Cambridge, MA, USA, 2016; Volume 94, pp. 251–285.

68. Mao, X.-M.; Zhao, S.-M.; Cao, L.; Yan, X.; Han, R.-C. The morphology observation of *Ophiocordyceps sinensis* from different origins. *J. Environ. Entomol.* **2013**, *35*, 343–353.

69. Mariani, V.; Biasini, M.; Barbato, A.; Schwede, T. lDDT, a local superposition-free score for comparing protein structures and models using distance difference tests. *Bioinformatics* **2013**, *29*, 2722–2728. https://doi.org/10.1093/bioinformatics/btt473.

70. Martin, T.; Lu, S.-W.; van Tilbeurgh, H.; Ripoll, D.R.; Dixelius, C.; Dixelius, C.; Turgeon, B.G.; Debuchy, R. Tracing the Origin of the Fungal a1 Domain Places Its Ancestor in the HMG-Box Superfamily, Implication for Fungal Mating-Type Evolution. *PLoS ONE* **2010**, *5*, e15199. https://doi.org/10.1371/journal.pone.0015199.

71. Meng, Q.; Yu, H.-Y.; Zhang, H.; Zhu, W.; Wang, M.-L.; Zhang, J.-H.; Zhou, G.-L.; Li, X.; Qin, Q.-L.; Hu, S.-N.; et al. Transcriptomic insight into the immune defenses in the ghost moth, Hepialus xiaojinensis, during an *Ophiocordyceps sinensis* fungal infection. Insect Biochem. *Mol. Biol.* **2015**, *64*, 1–15. https://doi.org/10.1016/j.ibmb.2015.06.014.

72. Metin, B.; Findley, K.; Heitman, J. The mating type locus (MAT) and sexual reproduction of *Cryptococcus heveanensis*: Insights into the evolution of sex and sex-determining chromosomal regions in fungi. *PLoS Genet.* **2010**, *6*, e1000961. https://doi.org/10.1371/journal.pgen.1000961.

73. Mishra, A.; Forche, A.; Anderson, M.Z. Parasexuality of Candida Species. *Front. Cell. Infect. Microbiol.* **2021**, *11*, 796929. https://doi.org/10.3389/fcimb.2021.796929.

74. Monzon, V.; Haft, D.H.; Bateman, A. Folding the unfoldable, using AlphaFold to explore spurious proteins. *Bioinform. Adv.* **2022**, *1*, vbab043. https://doi.org/10.1093/bioadv/vbab043.

75. Nakamura, N.; Tanaka, C.; Takeuchi-Kaneko, Y. Transmission of antibiotic-resistance markers by hyphal fusion suggests partial presence of parasexuality in the root endophytic fungus Glutinomyces brunneus. *Mycol. Progress.* **2019**, *18*, 453–462. https://doi.org/10.1007/s11557-018-1455-9.

76. Peters, C.; Elofsson, A. Why is the biological hydrophobicity scale more accurate than earlier experimental hydrophobicity scales? *Proteins* **2014**, *82*, 2190–2198. https://doi.org/10.1002/prot.24582.

77. Poveda-Huertes D, Matic S, Marada A, Habernig L, Licheva M, Myketin L, Gilsbach R, Tosal-Castano S, Papinski D, Mulica P, Kretz O, Kücükköse C, Taskin AA, Hein L, Kraft C, Büttner S, Meisinger C, Vögtle FN. An Early mtUPR: Redistribution of the Nuclear Transcription Factor Rox1 to Mitochondria Protects against Intramitochondrial Proteotoxic

Aggregates. *Mol Cell*. **2020**, *77*, 180–188. doi: 10.1016/j.molcel.2019.09.026

78. Qin, Q.-L.; Zhou, G.-L.; Zhang, H.; Meng, Q.; Zhang, J.-H.; Wang, H.-T.; Miao, L.; Li, X. Obstacles and approaches in artificial cultivation of Chinese cordyceps. *Mycology* **2018**, *9*, 7–9. https://doi.org/10.1080/21501203.2018.1442132.

79. Quandt, C.A., Kepler, R.M., Gams, W., Araújo, J.P.M., Ban, S., Evans, H.C., Hughes, D., Humber, R., Hywel-Jones, N., Li, Z.-Z., Luangsa-ard, J.J., Rehner, S.A., Sanjuan, T., Sato, H., Shrestha, B., Sung, G.-H., Yao, Y.-J., Zare, R., Spatafora, J.W. Phylogenetic-based nomenclatural proposals for Ophiocordycipitaceae (Hypocreales) with new combinations in Tolypocladium. *IMA Fungus*. **5**, *1*, 121–134. doi:10.5598/imafungus.2014.05.01.12.

80. Rajewska M, Wegrzyn K, Konieczny I. AT-rich region and repeated sequences - the essential elements of replication origins of bacterial replicons. *FEMS Microbiol Rev*. **2012**, *36*, 408–434. doi: 10.1111/j.1574-6976.2011.00300.x

81. Ramšak, B.; Kück, U. The *Penicillium chrysogenum* tom1 gene a major target of transcription factor MAT1-1-1 encodes a nuclear protein involved in sporulation. *Front. Fungal Bio.* **2022**, *3*, 937023. https://doi.org/10.3389/ffunb.2022.937023.

82. Ramšak, B.; Kűck, U.; Hofmann, E. The mating type transcription factor MAT1-1-1 from the fungal human pathogen *Aspergillus fumigatus*: Synthesis, purification, and crystallization of the DNA binding domain. *bioRxiv* **2021**. https://doi.org/10.1101/2021.12.13.472399.

83. Ramšak, B.; Markau, J.; Pazen, T.; Dahlmann, T.A.; Krappmann, S.; Kűck, U. The master regulator MAT1-1-1 of fungal mating binds to its targets via a conserved motif in the human pathogen *Aspergillus fumigatus*. *G3 Genes Genom. Genet.* **2020**, *11*, jkaa012. https://doi.org/10.1093/g3journal/jkaa012.

84. Ren, Y.; Wan, D.-G.; Lu, X.-M.; Guo, J.-L. The study of scientific name discussion for TCM Cordyceps. *LisShenzhen Med. Mater. Medica Res.* **2013**, *24*, 2211–2212.

85. Rettie, S.A.; Campbell, K.V.; Bera, A.K.; Kang, A.; Kozlov, S.; De La Cruz, J.; Adebomi, V.; Zhou, G.; DiMaio, F.; Ovchinnikov, S.; et al. Cyclic peptide structure prediction and design using AlphaFold. *bioRxiv* **2023**, 26:2023.02.25.529956. https://doi.org/10.1101/2023.02.25.529956.

86. Samarasinghe, H.; You, M.; Jenkinson, T.S.; Xu, J.-P.; James, T.Y. Hybridization Facilitates Adaptive Evolution in Two Major Fungal Pathogens. *Genes* **2020**, *11*, 101. https://doi.org/10.3390/genes11010101.

87. Seervai, R.N.H.; Jones, S.K.; Hirakawa, M.P.; Porman, A.M.; Bennett, R.J. Parasexuality and ploidy change in *Candida tropicalis*. *Eukaryot. Cell.* **2013**, *12*, 1629–1640. https://doi.org/10.1128/EC.00128-13.

88. Sherwood, R.K.; Bennett, R.J. Fungal meiosis and parasexual reproduction--lessons from pathogenic yeast. *Curr. Opin. Microbiol.* **2009**, *12*, 599–607. https://doi.org/10.1016/j.mib.2009.09.005.

89. Shu, R.-H.; Zhang, J.-H.; Meng, Q.; Zhang, H.; Zhou, G.-L.; Li, M.-M.; Wu, P.-P.; Zhao, Y.-N.; Chen, C.; Qin, Q.-L. A new high-quality draft genome assembly of the Chinese cordyceps *Ophiocordyceps sinensis*. *Genome Biol. Evol.* **2020**, *12*, 1074–1079. https://doi.org/10.1093/gbe/evaa112.

90. Simm, S.; Einloft, J.; Mirus, O.; Schleiff, E. 50 years of amino acid hydrophobicity scales, revisiting the capacity for peptide classification. *Biol. Res.* **2016**, *49*, 31. https://doi.org/10.1186/s40659-016-0092-5.

91. Song, L.-R.; Hong, X.; Ding, X.-L.; Zang, Z.-Y. A comprehensive dictionary of modern pharmacy of Chinese traditional medicine. People's Medical Press, Beijing. 2001; pp 733–737.

92. Steensels, J.; Gallone, B.; Verstrepen, K.J. Interspecific hybridization as a driver of fungal evolution and Adaptation. *Nat. Rev. Microbiol.* **2021**, *19*, 485–500.

93. Stensrud, Ø.; Hywel-Jones, N.L.; Schumacher, T. Towards a phylogenetic classification of Cordyceps: ITS nrDNA sequence data confirm divergent lineages and paraphyly. *Mycol. Res.* **2005**, *109*, 41–56. https://doi.org/10.1017/s095375620400139x.

94. Stensrud, Ø.; Schumacher, T.; Shalchian-Tabrizi, K.; Svegardenib, I.B.; Kauserud, H. Accelerated nrDNA evolution and profound AT bias in the medicinal fungus *Cordyceps sinensis*. *Mycol. Res.* **2007**, *111*, 409–415. https://doi.org/10.1016/j.mycres.2007.01.015.

95. Stone, R. Improbable partners aim to bring biotechnology to a Himalayan kingdom. *Science* **2010**, *327*, 940–941. https://doi.org/10.1126/science.327.5968.940.

96. Sun, S.; Coelho, M.A.; David-Palma, M.; Priest, S.J.; Heitman, J. The evolution of sexual reproduction and the mating-type locus: Links to pathogenesis of *Cryptococcus* human pathogenic fungi. *Annu. Rev. Genet.* **2019**, *53*, 417–444. https://doi.org/10.1146/annurev-genet-120116-024755.

97. Sung, G.-H.; Hywel-Jones, N.L.; Sung, J.-M.; Luangsa-ard, J.J.; Shrestha, B.; Spatafora, J.W. Phylogenetic classification of *Cordyceps* and the clavicipitaceous fungi. *Stud. Mycol.* **2007**, *57*, 5–59. https://doi.org/10.3114/sim.2007.57.01.

98. Taylor JW. One Fungus = One Name: DNA and fungal nomenclature twenty years after PCR. *IMA Fungus*. **2011**, *2*, 113–120. doi:10.5598/imafungus.2011.02.02.01

99. Thapar, R. Structure-specific nucleic acid recognition by L-motifs and their diverse roles in expression and regulation of the genome. *Biochim. Biophys. Acta* **2015**, *1849*, 677–687. https://doi.org/10.1016/j.bbagrm.2015.02.006.

100. Tunyasuvunakool, K.; Adler, J.; Wu, Z.; Green, T.; Zielinski, M.; Žídek, A.; Bridgland, A.; Cowie, A.; Meyer, C.; Laydon, A.; Velankar S, Kleywegt GJ, Bateman A, Evans R, Pritzel A, Figurnov M, Ronneberger O, Bates R, Kohl SAA, Potapenko A, Ballard AJ, Romera-Paredes B, Nikolov S, Jain R, Clancy E, Reiman D, Petersen S, Senior AW, Kavukcuoglu K, Birney E, Kohli P, Jumper J, Hassabis D. Highly accurate protein structure prediction for the human proteome. *Nature* **2021**, *596*, 590–596. https://doi.org/10.1038/s41586-021-03828-1.

101. Turgeon, B.G.; Yoder, O.C. Proposed nomenclature for mating type genes of filamentous ascomycetes. *Fungal Genet. Biol.* **2000**, *31*, 1–5. https://doi.org/10.1006/fgbi.2000.1227.

102. Varadi, M.; Bertoni, D.; Magana, P.; Paramval, U.; Pidruchna, I.; Radhakrishnan, M.; Tsenkov, M.; Nair, S.; Mirdita, M.; Yeo, J.; et al. AlphaFold Protein Structure Database in 2024, providing structure coverage for over 214 million protein sequences. *Nucleic Acids Res.* **2024**, *52*, D368–D375. https://doi.org/10.1093/nar/gkad1011.

103. Wang, Y.; Stata, M.; Wang, W.; Stajich, J.E.; White, M.M.; Moncalvo, J.M. Comparative genomics reveals the core gene toolbox for the fungus-insect symbiosis. *mBio* **2018**, *9*, e00636-e18. https://doi.org/10.1128/mBio.00636-18.

104. Wang, Y.-B., Wang, Y., Fan, Q., Duan, D-E., Zhang, G.-D., Dai6, R.-Q., Dai, Y.-D., Zeng, W.-B. Chen, Z.-H., Li, D.-D., Tang, D.-X., Xu, Z.-H., Sun, T., Nguyen, T.-T., Tran, N.-L., Dao, V.-M., Zhang, C.-M., Huang, L.-D., Liu, Y.-J., Zhang, X.-M., Yang, D.-R., Sanjuan, T., Liu, X.-Z., Yang, Z.-L., Yu, H. Multigene phylogeny of the family Cordycipitaceae (Hypocreales): new taxa and the new systematic position of the Chinese cordycipitoid fungus *Paecilomyces hepiali*. *Fungal Diversity* **2020**, *103*, 1–46. https://doi.org/10.1007/s13225-020-00457-3.

105. Wei, J.-C.; Wei, X.-L.; Zheng, W.-F.; Guo, W.; Liu, R.-D. Species identification and component detection of *Ophiocordyceps sinensis* cultivated by modern industry. *Mycosystema* **2016**, *35*, 404–410.

106. Wei, X.-L.; Yin, X.-C.; Guo, Y.-L.; Shen, N.-Y.; Wei, J.-C. Analyses of molecular systematics on *Cordyceps sinensis* and its related taxa. *Mycosystema* **2006**, *25*, 192–202.

107. Wilson, A.M.; Wilken, P.M.; van der Nest, M.A.; Steenkamp, E.T.; Wingfield, M.J.; Wingfield, B.D. Homothallism: An umbrella term for describing diverse sexual behaviours. *IMA Fungus* **2015**, *6*, 207–214. https://doi.org/10.5598/imafungus.2015.06.01.13.

108. Wroblewski, K.; Kmiecik, S. Integrating AlphaFold pLDDT Scores into CABS-flex for enhanced protein flexibility simulations. *Comput. Struct. Biotechnol. J.* **2024**, *30*, 4350–4356. https://doi.org/10.1016/j.csbj.2024.11.047.

109. Xia, E.-H.; Yang, D.-R.; Jiang, J.-J.; Zhang, Q.-J.; Liu, Y.; Liu, Y.-L.; Zhang, Y.; Zhang, H.-B.; Shi, C.; Tong, Y.; Kim C-H, Chen H, Peng Y-Q, Yu Y, Zhang W, Eichler EE, Gao L-Z. The caterpillar fungus, *Ophiocordyceps sinensis*, genome provides insights into highland adaptation of fungal pathogenicity. *Sci. Rep.* **2017**, *7*, 1806. https://doi.org/10.1038/s41598-017-01869-z.

110. Xia, F.; Liu, Y.; Guo, L.-X.; Shen, G.-L.; Lin, J.; Zhou, X.-W. Pyrosequencing analysis revealed complex endogenetic microorganism community from natural DongChong XiaCao and its microhabitat. *BMC Microbiol.* **2016**, *16*, 196. DOI 10.1186/s12866-016-0813-5

111. Xia, F.; Liu, Y.; Shen, G.-L.; Guo, L.-X.; Zhou, X.-W. Investigation and analysis of microbiological communities in natural *Ophiocordyceps sinensis*. *Can. J. Microbiol.* **2015**, *61*, 104–111. https://doi.org/10.1139/cjm-2014-0610.

112. Xiang, L.; Li, Y.; Zhu, Y.; Luo, H.; Li, C.; Xu, X.; Sun, C.; Song, J.-Y.; Shi, L.-H.; He, L.; Sun W, Chen S-L. Transcriptome analysis of the *Ophiocordyceps sinensis* fruiting body reveals putative genes involved in fruiting body development and cordycepin biosynthesis. *Genomics* **2014**, *103*, 154–159. https://doi.org/10.1016/j.ygeno.2014.01.002.

113. Xiao, W.; Yang, J.-P.; Zhu, P.; Cheng, K.-D.; He, H.-X.; Zhu, H.-X.; Wang, Q. Non-support of species complex hypothesis of *Cordyceps sinensis* by targeted rDNA-ITS sequence analysis. *Mycosystema* **2009**, *28*, 724–730.

114. Xu M, Ashley NA, Vaghefi N, Wilkinson I, Idnurm A. Isolation of strains and their genome sequencing to analyze the mating system of *Ophiocordyceps robertsii*. *PLoS One*. **2023a**, *18*, e0284978. doi: 10.1371/journal.pone.0284978.

115. Xu, T.; Xu, Q.; Li, J.-Y. Toward the appropriate interpretation of Alphafold2. *Front. Artif. Intell.* **2023a**, *6*, 1149748. https://doi.org/10.3389/frai.2023.1149748.

116. Yamamoto, A.; Ando, Y.; Yoshioka, K.; Saito, K.; Tanabe, T.; Shirakawa, H.; Yoshida, M. Difference in affinity for DNA between HMG proteins 1 and 2 determined by surface plasmon resonance measurements. *J. Biochem.* **1997**, *122*, 586–594. https://doi.org/10.1093/oxfordjournals.jbchem.a021793. PMID 9348088.

117. Yang, J.-L.; Xiao, W.; He, H.-X.; Zhu, H.-X.; Wang, S.-F.; Cheng, K.-D.; Zhu, P. Molecular phylogenetic analysis of *Paecilomyces hepiali* and *Cordyceps sinensis*. *Acta Pharmaceut. Sinica.* **2008**, *43*, 421–426.

118. Yang, J.-Y.; Tong, X.-X.; He, C.-Y.; Bai, J.; Wang, F.; Guo, J.-L. Comparison of endogenetic microbial community diversity between wild *Cordyceps sinensis*, artificial *C. sinensis* and habitat soil. *Chin. J. Chin. Mater. Medica* **2021**, *46*, 3106–3115.

119. Yang, X.; Zhu, H.-Q.; Shi, L.-X.; Song, T.-G.; Gong, W.-B.; He, S.-M.; Shan, S.; Xu, C.-F.; Zhou, Z. AlphaFold-guided structural analyses of nucleosome binding proteins. *Nucleic Acids Res.* **2025**, *53*, gkaf735. https://doi.org/10.1093/nar/gkaf735.

120. Yao, Y.-S.; Zhu, J.-S. Indiscriminate use of the Latin name for natural *Cordyceps sinensis* and *Ophiocordyceps sinensis* fungi. *Chin. J. Chin. Mater. Med.* **2016**, *41*, 1316–1366.

121. Yokoyama E, Yamagishi K, Hara A, Heterothallism in *Cordyceps takaomontana*. *FEMS Microbiol Lett*. **2005**, *250*, 145–150. https://doi.org/10.1016/j.femsle.2005.07.004.

122. Zhang, S.; Zhang, Y.-J. Molecular evolution of three protein-coding genes in the Chinese caterpillar fungus *Ophiocordyceps sinensis*. *Microbiol. China.* **2015**, *42*, 1549–1560.

123. Zhang, S.; Zhang, Y.-J.; Liu, X.-Z.; Wen, H.-A.; Wang, M.; Liu, D.-S. Cloning and analysis of the MAT1-2-1 gene from the traditional Chinese medicinal fungus *Ophiocordyceps sinensis*. *Fungal Biol.* **2011**, *115*, 708–714.

124. Zhang, S.; Zhang, Y.-J.; Shrestha, B.; Xu, J.-P.; Wang, C.-S.; Liu, X.-Z. *Ophiocordyceps sinensis* and *Cordyceps militaris*: Research advances, issues and perspectives. *Mycosystema* **2013**, *32*, 577–597.

125. Zhang, S.-W.; Cen, K.; Liu, Y.; Zhou, X.-W.; Wang, C.-S. Metatranscriptomics analysis of the fruiting caterpillar fungus collected from the Qinghai-Tibetan plateau. *Sci. Sinica Vitae* **2018**, *48*, 562–570.

126. Zhang, Y.-J.; Li, E.-W.; Wang, C.-S.; Li, Y.-L.; Liu, X.-Z. *Ophiocordyceps sinensis*, the flagship fungus of China: Terminology, life strategy and ecology. *Mycology* **2012**, *3*, 2–10. https://doi.org/10.1080/21501203.2011.654354.

127. Zhang, Y.-J.; Sun, B.-D.; Zhang, S.; Wàngmǔ Liu, X.-Z.; Gong, W.-F. Mycobiotal investigation of natural *Ophiocordyceps sinensis* based on culture-dependent investigation. *Mycosistema* **2010a**, *29*, 518–527.

128. Zhang, Y.-J.; Xu, L.-L.; Zhang, S.; Liu, X.-Z.; An, Z.-Q.; Wàngmǔ Guo, Y.-L. Genetic diversity of *Ophiocordyceps sinensis*, a medicinal fungus endemic to the Tibetan Plateau: Implications for its evolution and conservation. *BMC Evol. Biol.* **2009**, *9*, 290. doi:10.1186/1471-2148-9-290

129. Zhang, Y.-J.; Zhang, S.; Li, Y.-L.; Ma, S.-L.; Wang, C.-S.; Xiang, M.-C.; Liu, X.; An, Z.-Q.; Xu, J.-P.; Liu, X.-Z. Phylogeography and evolution of a fungal–insect association on the Tibetan Plateau. *Mol. Ecol.* **2014**, *23*, 5337–5355. https://doi.org/10.1111/mec.12940.

130. Zhang, Y.-J.; Zhang, S.; Wàngmǔ; Bai, F.-Y.; Liu, X.-Z. High Diversity of the Fungal Community Structure in Naturally-Occurring *Ophiocordyceps sinensis*. *PLoS ONE* **2010b**, 5(12), e15570. doi:10.1371/journal.pone.0015570.

131. Zheng, P.; Xia, Y.-L.; Zhang, S.-W.; Wang, C.-S. Genetics of Cordyceps and related fungi. *Appl Microbiol Biotechnol*, **2013**, *97*, 2797–2804. doi: 10.1007/s00253-013-4771-7.

132. Zheng, P.; Wang, C.-S. Sexuality Control and Sex Evolution in Fungi. *Sci. Sin. Vitae* **2013**, *43*, 1090–1097.

133. Zheng, Q.; Hou, R.; Zhang, J.-Y.; Ma, J.; Ma, J.-W.; Wu, Z.-S.; Wang, G.-H.; Wang, C.-F.; Xu, J.-R. The MAT locus genes play different roles in sexual reproduction and pathogenesis in *Fusarium graminearum*. *PLoS ONE.* **2013**, *8*, e66980. https://doi.org/10.1371/journal.pone.0066980.

134. Zhong, X.; Gu, L.; Wang, H.-Z.; Lian, D.-H.; Zheng, Y.-M.; Zhou, S.; Zhou, W.; Gu, J.; Zhang, G.; Liu, X. Profile of *Ophiocordyceps sinensis* transcriptome and differentially expressed genes in three different mycelia, sclerotium and fruiting body developmental stages. *Fungal Biol.* **2018**, *122*, 943–951. https://doi.org/10.1016/j.funbio.2018.05.011.

135. Zhou, X.W.; Li, L.J.; Tian, E.W. Advances in research of the artificial cultivation of *Ophiocordyceps sinensis* in China. *Crit. Rev. Biotechnol.* **2013**, *34*, 233–243. https://doi.org/10.3109/07388551.2013.791245.

136. Zhu, J.-S.; Gao, L.; Li, X.-H.; Yao, Y.-S.; Zhou, Y.-J.; Zhao, J.-Q.; Zhou, Y.-J. Maturational alterations of oppositely orientated rDNA and differential proliferations of CG:AT-biased genotypes of *Cordyceps sinensis* fungi and *Paecilomyces hepiali* in natural *C.* sinensis. *Am. J. Biomed. Sci.* **2010**, *2*, 217–238. https://doi.org/10.5099/aj100300217.

137. Zhu J-S, Guo Y-L, Yao Y-S, Zhou Y-J, Lu J-H, Qi Y, Chen W, Zheng T-Y, Zhang L, Wu Z-M, Zhang L-J, Liu X-J, Yin W-T. Maturation of *Cordyceps sinensis* associates with co-existence of *Hirsutella sinensis* and *Paecilomyces hepiali* DNA and dynamic changes in fungal competitive proliferation predominance and chemical profiles. *J. Fungal Res.* **2007**, *5*, 214–224.

138. Zhu, J.-S.; Halpern, G.M.; Jones, K. The scientific rediscovery of a precious ancient Chinese herbal regimen: *Cordyceps sinensis*: Part I. *J. Altern. Complem. Med.* **1998a**, *4*, 289–303. https://doi.org/10.1089/acm.1998.4.3-289.

139. Zhu, J.-S.; Halpern, G.M.; Jones, K. The scientific rediscovery of an ancient Chinese herbal medicine: *Cordyceps sinensis*: Part II. *J. Altern. Complem. Med.* **1998b**, *4*, 429–457. https://doi.org/10.1089/acm.1998.4.429.

140. Zhu, J.-S.; Li, C.-L.; Tan, N.-Z.; Berger, J.L.; Prolla, T.A. Combined use of whole-gene expression profiling technology and mouse lifespan test in anti-aging herbal product study. In Proceedings of the 2011 New TCM Products Innovation and Industrial Development Summit, Hangzhou, China, 27 November 2011; pp. 443–448. Available online: https://xueshu.baidu.com/usercenter/paper/show?paperid=08341c17fa58c8f85584b92572b90f75&site=xueshu_se (accessed on 30 January 2025).

141. Zhu, J.-S.; Li, Y.-L. A Precious Transitional Chinese Medicine, *Cordyceps sinensis*: Multiple heterogeneous *Ophiocordyceps sinensis* in the insect–fungi complex. Lambert Academic Publishing, Saarbrüchen. Germany, 2017.

142. Zitomer, R.S.; Limbach, M.P.; Rodriguez-Torres, A.M.; Balasubramanian, B.; Deckert, J.; Snow, P.M. Approaches to the study of Rox1 repression of the hypoxic genes in the yeast Saccharomyces cerevisiae. *Methods* **1997**, *11*, 279–288. https://doi.org/10.1006/meth.1996.0422.

143. Zou, J.; Zhang, Y.-T.; Luo, Y.; Fu, M.-H.; Sun, B.-L.; Liu, S.-G. MAT1-1 and MAT1-2 *Ophiocordyceps xuefengensis* and comparison of their chemical composition. methods. *Biol. (Basel)* **2024**, *13*, 686; https://doi.org/10.3390/biology13090686.