Signalome-wide mapping of the NFκB pathway in T-cells reveals novel targets for

immunotherapy

Joseph Clarke[1], Haoqi Chen[2], Cristina Tormo Garcia[1], Ewa Basiarz[3], Mateusz Kotowski[1], Ana

Mafalda Santos[1], Mai T Voung[1], Paige Sherman[1], Jia Xin Li[1], Christopher J Tape[3#], Simon J

Davis[1#], Evangelia Petsalaki[2#], Sumana Sharma[1#]

[1]Radcliffe Department of Medicine, John Radcliffe Hospital, University of Oxford, Oxford, OX3

9DS, United Kingdom

[2]European Molecular Biology Laboratory, European Bioinformatics Institute (EMBL-EBI),

Wellcome Genome Campus, Hinxton, Cambridgeshire, CB10 1SD, United Kingdom

[3]Cell Communication Laboratory, Department of Oncology, University College London Cancer

Institute, 72 Huntley Street, London, WC1E 6DD, United Kingdom

[#]Correspondence should be addressed to:

sumana.sharma@rdm.ox.ac.uk

petsalaki@ebi.ac.uk

simon.davis@imm.ox.ac.uk

c.tape@ucl.ac.uk

Abstract

Cell signalling networks govern fundamental cellular processes yet remain incompletely defined. Moreover, what is known is biased toward a limited subset of well-characterised components. Phosphoprotein-based interrogation methods, including mass spectrometry and targeted phosphosite panels, have limited utility in physiological settings dependent on cell–cell interactions because the signalling fluxes can be difficult to detect despite producing robust functional responses. Here we developed a perturbation-based experimental framework that infers signalling pathway architecture using quantitative functional outputs rather than direct measurements of effector state, e.g., phosphorylation levels. Using antigen-specific, NF-κB–GFP reporter-expressing transformed T-cells co-cultured with cellular targets, we performed an arrayed CRISPR–Cas9 screen targeting a curated signalome of kinases, phosphatases, adaptor and scaffolding proteins, totalling 706 genes. Quantitative effect-size profiling recovered canonical T-cell receptor regulators and revealed unequal, family-specific patterns of control over NF-κB activation. Comparing T-cell stimulation with low- and high-affinity antigen uncovered signal-strength-dependent buffering of proximal signalling nodes, exemplified by reduced sensitivity to perturbation of *LCK* under high-intensity stimulation. Targeted perturbation in primary human CD8+ T-cells validated our findings and identified *TRRAP* and *CTDSPL2* as negative regulators of T-cell effector output, whose disruption enhanced cytotoxicity, degranulation, and cytokine production in both polyclonal and TCR-engineered T cells. Together, these results establish a scalable strategy for mapping signalling pathway architecture in the setting of physiological T-cell activation.

Introduction

Cellular signalling is crucial for fundamental processes such as growth, proliferation, differentiation, survival, metabolism, and specialised effector responses. Although powerful technologies such as CyTOF, LC–MS-based phosphoproteomics, and targeted phospho-panels have significantly advanced our ability to interrogate signalling events, our understanding of cellular signalling network structure and regulation remains incomplete. Moreover, current knowledge is disproportionately focused on a limited subset of well-studied components, leaving large portions of the signalling landscape poorly characterised.[1]. This bias is particularly pronounced for kinases and phosphatases, the principal enzymatic drivers of signal transduction, for which an estimated 30-50% of cellular targets remain unknown, indicating that large regions of the signalling landscape remain unexplored [2].

Most contemporary approaches to studying signalling rely on direct measurement of molecular intermediates, most commonly phosphorylation events detected by antibody-based assays or mass spectrometry-based phosphoproteomics. Advances in enrichment-based LC-MS have enabled increasingly global profiling of signalling states [3–8], and these approaches have been extended through integration with genetic or pharmacological perturbations to infer causal relationships, exemplified by resources such as the Connectivity Map (CMap) [9]. Despite these advances, molecular readout-based approaches remain limited in their ability to capture signalling behaviour under physiological conditions. They are biased toward abundant or highly dynamic phosphorylation events, require large numbers of cells, and rely heavily on curated pathway annotations for interpretation, reinforcing existing knowledge rather than revealing new regulatory logic. Consequently, signalling components that exert modest, distributed, or context-dependent effects on pathway output may be systematically overlooked.

These limitations are particularly pronounced in systems where signalling is spatially constrained and dependent on direct cell–cell interactions. Most phosphoproteomic and

kinase-activity-based studies are performed on bulk cell lysates, restricting their applicability

in such contexts. T-cell activation exemplifies this challenge: apart from a small number of

SILAC-based co-culture studies (*10–12*), much of the available signalling data relies on

supraphysiological stimulation applied outside native cellular contexts to generate detectable

responses (*5, 13*). Single-cell approaches such as mass cytometry (CyTOF) partially address

population averaging (*14–17*). However, these approaches also encounter biological limits. In

direct T cell–target cell interactions, single-cell mass cytometric analyses reveal remarkably

limited signalling flux, often indistinguishable from unstimulated states, in contrast to

pharmacological induced T-cell activation (*18*). Notably, despite these modest detectable

fluxes, T cells mount robust functional responses, indicating that phosphoprotein-based

measurements fail to capture physiologically relevant signalling dynamics.

In contrast, functional outputs integrate signalling information over time and across molecular

layers. This suggests an alternative strategy for interrogating signalling networks: rather than

measuring signalling intermediates directly, signalling architecture can be inferred by

systematically perturbing signalling components and quantifying their impact on downstream

functional outputs.

Here, we implement this concept through a perturbation-based framework that reconstructs

signalling dependencies directly from functional readouts. By combining arrayed CRISPR-

Cas9-mediated gene disruption with a quantitative NF-κB reporter assay, we systematically

interrogate the contribution of kinases, phosphatases, and adaptor proteins to signalling

pathway transmission, and subsequent transcription factor activity, in a physiologically

relevant, cell–cell interaction context. This approach enables quantitative assessment of

graded effects across hundreds of signalling components, revealing how control is distributed

across a signalling network rather than identifying only a small number of essential nodes.

Applying this strategy to T-cell receptor signalling, we generate a functional dependency map

of NF-κB activation that reveals hierarchical control, distributed engagement across signalling

families, and context-dependent buffering of proximal signalling nodes as a function of ligand

affinity. Extending these findings to primary human CD8$^+$ T cells, we identify previously

unrecognized negative regulators of T-cell function, including TRRAP and CTDSPL2, whose

disruption enhances cytotoxic activity and effector responses. Together, this work establishes

a generalisable framework for reconstructing signalling architecture from functional outputs in

physiological contexts.

99 Results

A functional 1G4-TCR$^+$ Jurkat NF-kB reporter system enables analysis of T-cell

receptor signalling under physiological stimulation conditions.

To enable systematic interrogation of T-cell signalling through functional outputs, we

generated a Jurkat T-cell reporter line in which NF-κB activity is monitored via GFP expression

downstream of antigen-specific T-cell receptor (TCR) engagement. NF-κB was selected as a

core transcriptional output of T-cell activation, integrating signals from multiple proximal and

distal signalling pathways. Jurkat cells were first engineered to stably express an NF-κB-driven

GFP reporter (**Supplementary Fig. 1A-B)**. To ensure antigen-specific activation, endogenous

TCR expression was ablated by CRISPR targeting of *TRAC* and *TRBC*, and cells were

subsequently transduced with lentivirus encoding the 1G4 TCR (**Supplementary Figure 1C**),

which recognizes the cancer-testis antigen NY-ESO-1$_{157\text{-}165}$ (henceforth referred to simply as

NY-ESO-1) peptide presented by HLA-A*02:01, followed by introduction of constitutive Cas9

expression to facilitate CRISPR-Cas9-mediated genetic perturbation (**Supplementary Fig.**

1D).

This configuration establishes a defined receptor-ligand interaction that mimics physiological

T-cell-target cell engagement. When co-cultured with A375 melanoma cells pulsed with NY-

ESO-1 peptides, 1G4-TCR$^+$ Jurkat reporter cells exhibited robust NF-κB–dependent GFP

induction, whereas TCR-deficient parental cells remained unresponsive (**Fig. 1A–C**). Upon

the use of NY-ESO-1 altered peptide ligands (6T, 3Y, 3I and 9V) to initiate Jurkat T-cell

activation, GFP expression scaled with peptide affinity, demonstrating sensitivity across a

physiologically relevant range of TCR stimulation strengths (**Fig. 1B,C**).

To confirm that reporter activation reflected canonical TCR signalling, we genetically disrupted

key proximal signalling components. CRISPR-mediated targeting of established TCR

signalling mediators, including *LCP2* and *ZAP70*, significantly reduced NF-κB reporter activity

upon stimulation with both high- and low-affinity antigen (**Fig. 1D**). These results confirm that

NF-κB reporter activation depends on intact TCR-proximal signalling.

Together, these data establish a genetically tractable, antigen-specific Jurkat reporter system

that reliably converts physiologically relevant TCR engagement into a quantitative functional

readout of NF-κB activation. This platform provides a robust foundation for systematic

perturbation-based analysis of signalling dependencies in a cell–cell interaction context,

where direct phosphoprotein measurements are known to be challenging.

Figure 1: Establishing a peptide:MHC specific cell-cell interaction system to report functional activity of NFkB in T cells A: Schematic representing the co-culture system between Jurkat-NFkB-GFP reporter cell lines expressing the 1G4-TCR, and A375 melanoma cells pulsed with NY-ESO-1 peptides. **B:** Time course IncuCyte measurements of NFkB-GFP intensities from either 1G4-TCR⁺ Jurkat reporter lines (filled circles) or TCR-KO parental lines (filled squares). All NY-ESO-1 altered peptide ligands were used at final concentrations of 100µg/mL. **C:** Area under the curve (AUC) analysis of the data shown in panel B. **D:** Lentiviral sgRNA targeting of key TCR-signalling components *LCP2* or *ZAP70* in 1G4-TCR⁺ Jurkat reporters prevents GFP expression upon co-culture with NY-ESO-1 peptide pulsed A375 melanoma cells. All peptides were used at final concentrations of 100µg/mL, and time course GFP measurements were performed via IncuCyte imaging. All data shown from a minimum of three independent experiments.

A signalome-wide CRISPR-Cas9 screen reveals a quantitative map of NF-κB signalling in T cells

To enable systematic interrogation of T-cell signalling, we first defined a comprehensive "signalome" comprising kinases, phosphatases, and key adaptor proteins. We combined a curated list of 556 human kinases with 136 protein phosphatases defined by the HUGO Gene

Nomenclature Committee (*19*), and filtered this set for genes expressed across multiple

immune and cancer cell models using bulk RNA-seq data from Cell Model Passports (TPM >

0.3; Jurkat, MOLT-4, Daudi, THP-1, and two colorectal cancer organoids) (*20*). To further

enrich for signalling components relevant to T cells, we included genes annotated in KEGG

pathways associated with TCR signalling, JAK-STAT, calcium signalling, actin regulation,

phosphatidylinositol signalling, and MAPK signalling (*21*), as well as all genes encoding for

SH2 domain-containing proteins, to ensure coverage of adaptor and scaffolding molecules.

This yielded a final library of 706 genes, each targeted with two sgRNAs for CRISPR-Cas9-

mediated perturbation (**Supplementary Table 1**).

To place this signalome in the context of existing knowledge, we integrated transcriptional and

perturbation-based datasets from diverse T-cell contexts. Analysis of published expression

datasets spanning multiple activation timepoints in CD8$^+$ and pan-T cells (*22–25*) revealed

that 446 signalome genes are dynamically regulated at the transcriptional level (**Fig. 2A;**

Supplementary Table 2). In parallel, we mapped hits from published genome-wide CRISPR

loss-of-function and gain-of-function screens in T cells (*26–29*) onto the signalome (**Fig. 2B;**

Supplementary Table 2). Across these datasets, 64 genes were identified in loss-of-function

screens and 49 in gain-of-function screens. Genes implicated by CRISPR perturbation were

significantly enriched for genes encoding SH2-domain-containing proteins (p = 2x10^{-5}, Fisher's

exact test), underscoring their central role in T-cell signalling. Notably, 37 genes that were not

transcriptionally regulated nonetheless exhibited functional effects in CRISPR screens,

including seven core T-cell signalling genes encoding for SH2-domain-containing proteins

(CBLB, GRB2, ZAP70, SLA2, CBL, LCP2, VAV1), whereas many transcriptionally regulated

signalling genes were not detected in pooled CRISPR screens, highlighting potential gaps in

existing perturbation-based maps (**Fig. 2A,B**).

We next performed a signalome-wide, arrayed CRISPR perturbation screen to directly quantify

the contribution of each component to NF-κB activation. 1G4-TCR$^+$ Jurkat reporter cells were

transduced with individual sgRNAs and stimulated through cell–cell co-culture with peptide-

pulsed target cells. In total, ~1,500 sgRNAs targeting ~700 genes were assayed in 96-well

format across 13 experimental batches (**Fig. 2C**). Each batch included internal controls,

including an empty sgRNA vector and sgRNAs targeting known TCR signalling components

(*LCP2, ITK*), enabling normalization of gene-level effects based on GFP signal area under the

curve (AUC) relative to empty controls and an internal positive control to assess the editing

efficiency (see details in **Methods**).

The screen showed high technical robustness. Independent sgRNAs targeting the same gene

exhibited strong concordance, with ~80% of guide pairs showing consistent effect direction

(**Fig. 2D; Supplementary Figure. 2A**). Gene-level effects were highly correlated for T-cell

stimulation with low-affinity (6T) and high-affinity (9V) antigens (Spearman $\rho = 0.93$), indicating

that the core signalling architecture governing NF-κB activation is largely conserved across

stimulation strengths (**Fig. 2E**). Most genes clustered near the identity line, whereas a subset

showed context-dependent deviations. Notably, disruption of *LCK* strongly attenuated NF-κB

activation under weak stimulation but had a markedly reduced effect under high-affinity

stimulation (**Supplementary Figure. 2B**), consistent with partial bypass or compensation at

higher signalling strength (*30, 31*).

To characterise the global structure of signalling control, we ranked genes by their mean effect

size across both conditions. The distribution was strongly skewed toward negative effects,

consistent with depletion-driven signalling dependencies rather than enrichment effects (**Fig.**

2F). Canonical TCR signalling components, including *LCK, ZAP70, ITK, and LCP2*, ranked

among the strongest negative regulators, validating the ability of this approach to recover core

pathway architecture (**Fig. 2F**). TCR-specific signalling was among the highest-ranked

pathways in the enrichment analysis, suggesting that the screen successfully recapitulated

known genes while also assigning quantitative scores to signalome genes involved in

regulating the NF-κB pathway (**Supplementary Figure 2C**).

We next examined how signalling control was distributed across protein families by comparing

effect size distributions within each family (**Fig. 2G**). All families contributed to NF-κB

regulation, but with distinct patterns. SH2-domain-containing proteins exhibited broad effect

size distributions, indicating distributed engagement of multiple family members. In contrast,

MAP kinase contributions were concentrated in a small number of nodes, leading towards

ERK1/2 (*MAPK3/MAPK1*) and p38α (*MAPK14*) (**Fig 2F, H**). These patterns suggest that NF-

κB signalling integrates both distributed adaptor-mediated control and more focal kinase

bottlenecks, rather than being uniformly dominated by single components. Based on these

quantitative dependencies, we reconstructed a data-driven network representation of NF-κB

regulation, capturing hierarchical patterns of control, characterized by a small number of high-

impact nodes alongside broadly distributed contributions from adaptor families (**Fig. 2H;**

Supplementary Table 3).

Figure 2: A signalome-wide, arrayed CRISPR perturbation screen reveals hierarchical and distributed control of NF-κB signalling in T cells. **A:** Overlap between genes that are not transcriptionally regulated across four T-cell–specific expression datasets and genes identified as hits in genome-wide CRISPR screens, revealing 37 signalling genes that are functionally required despite lacking dynamic expression changes. UpSet plot shows intersections across expression studies. **B:** Overlap between genes that are differentially expressed in at least one T-cell expression dataset and genes not identified as hits in genome-wide CRISPR screens, identifying 388 signalling genes that are transcriptionally regulated but not functionally required in pooled CRISPR screens. UpSet plot shows intersections across CRISPR studies. **C:** Schematic of the arrayed CRISPR perturbation strategy. **D:** Concordance between independent sgRNAs targeting the same gene under high-affinity stimulation (9V). Each point represents a gene; axes show $\log_2$(AUC ratio) for sgRNA1 versus sgRNA2. **D:** Comparison of gene-level perturbation effects between low-affinity (6T) and high-affinity (9V) stimulation conditions. Each point represents one gene collapsed across the two sgRNAs. **F.** Genes ranked by overall effect size, defined as the mean gene-level $\log_2$(AUC ratio) across both stimulation conditions. **G:** Distribution of effect sizes in 9V condition stratified by functional protein families **H:** Network representation of selected signalling modules with node colour indicating signed effect size under stimulation with low affinity antigen (6T). Nodes represent genes; edges correspond to curated functional interactions from STRING.

Targeted perturbation of SH2-domain-containing proteins in primary CD8$^+$ T cells

validates functional TCR signalling dependencies

Although large-scale perturbation screens with functional read outs are readily implemented

in cell lines, extending such approaches to primary human T cells remains challenging. To

enable arrayed perturbation in primary CD8$^+$ T cells at scale, we established a workflow based

on lentiviral delivery of sgRNAs followed by Cas9 electroporation, a strategy compatible with

systematic perturbation, while avoiding the cost and scalability constraints of RNP-only

approaches (*27*).

A further barrier to scalable functional assays in primary T cells is antigen specificity, as

individual T cells respond only to their cognate TCR ligand. To circumvent this limitation, we

employed bispecific molecules, which engage CD3 independently of endogenous TCR

specificity (*32, 33*). Here, the bispecific engager consists of an anti-CD3 single-chain variable

fragment fused to a TCR targeting an HLA-A2-bound gp100 peptide (YLEPGPVTV), thereby

triggering TCR signalling and directing T-cell activation against gp100-pulsed A375 melanoma

cells (*18*). This approach enables the use of polyclonal primary CD8⁺ T cells while preserving

physiologically relevant TCR-proximal signalling, thus substantially increasing experimental

scalability.

Upon co-culture with gp100-pulsed target cells, polyclonal CD8⁺ T cells exhibited robust

upregulation of activation markers (CD69, CD25, 4-1BB) and potent cytotoxicity (**Fig. 3A**).

Despite these strong functional responses, phosphoprotein changes measured by CyTOF

remained limited (**Fig. 3B, Supplementary Figure 3**), consistent with the low signalling flux

observed during direct cell–cell interactions and reinforcing the sensitivity of functional

readouts for capturing signalling outcomes (*18*).

To validate the signalling architecture inferred from the Jurkat signalome screen in a

physiological setting, we focused on the SH2-domain-containing protein family, which

exhibited broad functional engagement in NF-κB regulation despite lacking a coherent pattern

of transcriptional regulation during T-cell activation (**Fig. 2G**). This family includes multiple

well-established TCR signalling components, making it a suitable test case for functional

validation. We targeted 66 SH2-domain-containing genes, along with control targets, in

polyclonal primary CD8⁺ T cells from six independent donors using lentiviral sgRNA delivery

followed by Cas9 electroporation (**Supplementary Figure 4A**). Functional consequences

were assessed by quantifying target cell killing in co-culture assays (**Fig. 3C**).

Disruption of canonical TCR signalling components, including *LCK*, *LCP2*, and *ITK*, markedly

impaired CD8⁺ T-cell cytotoxicity (**Fig. 3C,D**) and reduced expression of activation markers 4-

1BB (**Fig. 3E**) and CD25 (**Supplementary Figure 4B**) at 48 hours, validating the sensitivity

and specificity of the assay. After correcting for donor and batch effects (**Methods**), systematic

perturbation of the SH2 gene family revealed a spectrum of functional effects on cytotoxicity

(**Fig. 3F**).

Genes were ranked by their impact on tumour cell killing, with cytotoxicity levels quantified

using the AUC metric (**Methods**). Our analysis revealed profiles that closely mirrored both

known TCR signalling annotations, and the dependencies observed in the Jurkat screen (**Fig.**

3F). Perturbation of key signalling mediators (*ITK*, *LCK*, *LCP2*, *PLCγ*, *PIK3R1-3*), adaptor

proteins (*GRAP*, *GRB2*), and cytoskeletal regulators (*VAV1*, *VAV3*) consistently reduced

CD8$^+$ T-cell cytotoxicity, whereas disruption of *SLA2* and *PTEN*, both known negative

regulator of TCR signalling, enhanced CD8$^+$ T-cell-mediated killing (**Fig. 3G; Supplementary**

Fig. 4C,D). These results demonstrate that the SH2 gene family constitutes a distributed and

functionally essential regulatory module in primary T-cell activation.

Because bispecific-based activation bypasses endogenous antigen specificity, we next asked

whether these dependencies were preserved in antigen-specific TCR-engineered T cells.

Primary CD8$^+$ T cells were transduced with plasmids encoding the 1G4-TCR and an LNGFR

tag, and TCR Vβ13.1$^+$ LNGFR$^+$ cells were sorted and expanded for use *in vitro*

(**Supplementary Fig. 4E**). This system enabled direct comparison of signalling requirements

across different ligand affinities. Consistent with the Jurkat screen, disruption of *LCK* exhibited

affinity-dependent effects: under high-affinity stimulation with the 3Y peptide, *LCK* targeting

had minimal impact on tumour cell killing, whereas under weaker stimulation conditions (6T),

cytotoxicity was strongly impaired (**Fig. 3G**).

Together, these results demonstrate that functional signalling dependencies identified in

Jurkat cells are largely conserved in primary human CD8$^+$ T cells. This validation establishes

the robustness of the perturbation-based framework across experimental systems and

highlights how distributed adaptor-mediated regulation and signal-strength–dependent

buffering shape TCR-driven functional responses.

Figure 3: Targeted perturbation in primary CD8+ T-cells reveals the spectrum of activity of the SH2 gene family on T-cell cytotoxicity. A: Top panel - Expression (mean fluorescence intensity, MFI) of activation markers on CD8- T-cells at 24 hours following co-

culture with gp100 pulsed A375 melanoma cell lines and a bispecific T cell engager. Bottom
panel - CD8$^+$ cytotoxic responses following co-culture with A375 melanoma cell lines as
described for the top panel. Data in both panels from 3 independent donors, representative of
at least 3 experimental repeats. **B:** Phosphorylation of early signalling mediators downstream
of TCR engagement as measured by mass cytometry (CyTOF). Data shown from two
independent donors in one experiment, representative of at least 4 independent donors across
4 experiments. **C:** Representative killing curves from CD8$^+$ T-cell co-cultures with A375
melanoma cells. T-cell activation was initiated through use of the bispecific engager at a final
concentration of 10^{-10}M. Targeting of known TCR-signalling mediators with sgRNAs inhibits
cytotoxic T-cell responses. Data show from one donor and one independent experiment,
representative of at least 6 independent donors across 6 independent experiments. **D:** Area
under the curve (AUC) analysis of killing curves shown in panel C, normalised to SHC4
targeted controls. Data from 6 independent donors across 6 independent experiments.
Statistical analysis performed by paired T-test. **E:** Targeting TCR-signalling mediators with
sgRNAs reduces surface 4-1BB expression, measured at 48 hours post co-culture with A375
melanoma cells as detailed in D. Data shown are the same T-cells from panel D, and are from
6 independent CD8$^+$ donors across 6 independent experiments. Statistical analysis performed
via One-Way ANOVA with Dunnett's multiple comparison tests. **F:** Waterfall plot of normalised
AUCs from killing curves illustrated in panel C, from experiments targeting all genes encoding
SH2-domain containing proteins. Killing curves for each perturbation were first normalised to
donor-specific internal negative control sgRNAs before computing AUC values. Ranked gene
analysis identifies both positive and negative regulators of T-cell cytotoxicity within the SH2
gene family. **G:** Representative killing curves from targeting *LCK*, *LCP2* or *SLA2* in primary
1G4-TCR$^+$ CD8$^+$ T-cells. A375 melanoma cell lines were pulsed with NY-ESO-1 peptides as
indicated to final concentrations of 100µg/mL. Data shown from one independent experiment,
representative of two independent repeats using two individual CD8$^+$ donors.

TRRAP and *CTDSPL2* constrain CD8$^+$ T-cell effector output downstream of TCR

signalling

Although the majority of perturbations in the signalome screen reduced NF-κB activity, a small

number of gene disruptions increased reporter output. Among these, targeting of *TRRAP* and

CTDSPL2 consistently enhanced NF-κB-driven GFP expression in Jurkat cells (**Fig. 4A**).

TRRAP is a pseudokinase that functions as a scaffold within histone acetyltransferase

complexes involved in transcriptional regulation, whereas CTDSPL2 is a poorly characterized

serine/threonine phosphatase. Neither protein has been well defined in the context of T-cell

signalling, prompting further functional validation in primary cells.

To assess their roles in physiological settings, we targeted *TRRAP* or *CTDSPL2* in primary

human CD8$^+$ T cells using lentiviral sgRNA delivery followed by Cas9 electroporation.

Disruption of either gene significantly enhanced CD8$^+$ T-cell-mediated killing of A375

melanoma cells in bispecific-based co-culture assays (**Fig. 4B**). Consistent with increased

effector activity, targeting *TRRAP* or *CTDSPL2* also elevated IFNγ secretion at 24 hours (**Fig.**

4C). In addition, after targeting *TRRAP* or *CTDSPL2* we observed enhanced degranulation,

as measured by surface CD107α expression at 4 hours post A375 co-culture (**Fig. 4D**). These

results indicate that both proteins act to constrain T-cell effector responses downstream of

TCR-proximal signalling.

Because bispecific-based activation bypasses endogenous antigen specificity, we next

examined whether these effects were preserved in antigen-specific TCR-engineered T cells.

Primary CD8$^+$ T cells expressing the 1G4 TCR were electroporated with sgRNA–Cas9

ribonucleoprotein (RNP) complexes targeting *TRRAP* or *CTDSPL2*, and cytotoxic activity was

assessed against A375 melanoma cells presenting endogenous levels of NY-ESO-1.

Importantly, these experiments utilised a distinct, non-overlapping sgRNA sequence

compared to these utilised in our signalome-wide arrayed screen. Under these basal antigen

conditions, disruption of either gene significantly enhanced tumour cell killing relative to

scrambled controls (**Fig. 4E**). This enhancement was maintained across a range of peptide

affinities, again utilising 6T or 3Y NY-ESO-1 altered peptide ligands (**Supplementary Fig. 5A**).

Finally, to assess whether enhanced CD8$^+$ T-cell effector functions arising from *TRRAP* or

CTDSPL2 perturbations were accompanied by an increase in markers of T-cell dysfunction

and exhaustion (e.g., PD-1, LAG-3 and TIM-3), cell surface expression of these molecules

was assessed in polyclonal CD8$^+$ T-cells 24 hours post stimulation with A375 melanoma cells

and the bispecific engager. Phenotypically, *TRRAP* or *CTSPL2* targeted T-cells were similar

to the unedited, empty plasmid controls, where expression of these three inhibitory receptors

displayed an activation-marker-like expression pattern. Expression of surface PD-1, LAG-3

and TIM-3 all correlated with the bispecific concentration utilised to initiate T-cell activation in

these experiments, and reduced expression was observed in the *LCP2* targeted control

(**Supplementary Fig. 5B**).

To investigate how TRRAP and CTDSPL2 depletion enhanced cytotoxicity, we performed

RNA sequencing following CRISPR-mediated perturbation in primary CD8$^+$ T-cells. TRRAP is

a scaffold component of the SAGA and TIP60 histone acetyltransferase complexes and

supports MYC- and E2F-dependent transcriptional programs. Consistent with this role, upon

activation, TRRAP-deficient T cells exhibited coordinated downregulation of genes associated

with cell cycle progression and DNA replication, including CENPA, PLK1, POLE2, and multiple

MCM complex components. In contrast, inflammatory and immune signalling genes, including

IFITM5, IL23R and NR4A family components were upregulated (**Figure 4F, Supplementary**

Table 4). Pathway-level analysis revealed suppression of proliferative programs alongside

enrichment of inflammatory and cytokine signalling modules (**Figure 4G**). These findings

suggest that TRRAP loss shifts T cells away from a proliferative transcriptional state toward

an inflammatory, effector-associated profile. By contrast, CTDSPL2-deficient cells displayed

minimal global transcriptional remodelling. Donor-paired analyses demonstrated that most

genes   remained   near   baseline   at   both   stimulated   and   unstimulated   conditions

(**Supplementary Fig. 5C, Supplementary Table 4**). The increased killing arising from

CTDSPL2 perturbation is, therefore, unlikely to arise from large-scale transcriptional

reprogramming and instead reflects altered signalling dynamics.

Together, these data identify TRRAP and CTDSPL2 as previously unrecognized negative

regulators of CD8$^+$ T cell effector function. Their disruption enhances cytotoxicity, cytokine

production, and degranulation under both polyclonal and antigen-specific activation conditions

without inducing excessive expression of inhibitory receptors. These findings highlight novel

regulatory   nodes   that   may   be   therapeutically   leveraged   to   enhance   T   cell–based

immunotherapies.

Figure 4: Perturbation of novel targets *CTDSPL2* and *TRRAP* enhances primary CD8+ T-cell functional activity. A: NFkB-GFP traces from 1G4-TCR+ Jurkat reporters when transduced with *CTDSPL2* (left) or *TRRAP* targeting sgRNAs (right). Jurkat cells were stimulated with A375 melanoma cell lines pulsed with NY-ESO-1 9V peptide (100µg/mL). **B:** Cytotoxic responses from primary CD8+ T-cells when co-cultured with A375 melanoma cell lines pulsed with gp100 peptide and a bispecific T-cell engager (final concentration 10^{-11}M). Data shown from 9 independent donors across 3 experiments (TRRAP sgRNAs) or a minimum of 4 independent donors across 2 experiments (CTDSPL2 sgRNAs). A375 cell growth was normalised to the empty control for each donor. **C:** IFNγ secretion at 24 hours from primary CD8+ T-cells stimulated as described for panel B. Data shown from 2 independent donors across 1 individual experiment, representative of at least 4 independent donors across 2 experiments. Statistical analysis performed via One-Way ANOVA with Dunnett's multiple comparisons. **D:** CD107α mean fluorescence intensity (MFI) from CD8+ T-cells stimulated as described for panel B, measured at 4 hours post stimulation. CD107a expression was normalised to unstimulated controls (no bispecific) for each CRISPR targeting condition. Data

shown from 4 independent donors across 2 experiments. **E:** Cytotoxic responses of 1G4-TCR[+] primary CD8[+] T-cells when co-cultured with A375 melanoma cell lines expressing basal NY-ESO-1. 1G4-TCR[+] CD8[+] T-cells were electroporated with RNPs targeting either *CTDSPL2* or *TRRAP*, together with a scrambled control (scRNA). Data shown from 6 donors across 3 independent experiments. Growth curves were normalised to the scRNA controls for all donors. **F.** Differential gene expression analysis of *TRRAP*-targeted primary CD8[+] T cells following stimulation. Volcano plot showing $\log_2$ fold change (KO vs WT) versus $-\log_{10}$ adjusted P value from donor-controlled limma-voom analysis (n = 3 donors). Selected representative genes are highlighted. Inset panels show donor-resolved normalized expression for representative upregulated inflammatory genes (e.g., IL2, IL7R, IL23R, TNFSF4) and downregulated proliferative genes (e.g., CDC6, CENPA, CHEK1, POLE2). Lines connect paired donor samples. TRRAP transcript abundance was modestly but consistently reduced across donors (see Fig. 4F, inset). **G**. Pathway enrichment analysis of differentially expressed genes in *TRRAP*-targeted cells. Hallmark gene set enrichment (Enrichr) identified suppression of proliferative programs, including G2/M checkpoint and E2F target pathways, and enrichment of inflammatory and TNFα–NF-κB signalling modules. Combined enrichment scores are shown.

Discussion

In this study, we present a perturbation-based framework to interrogate cellular signalling during physiological cell–cell engagement. Leveraging functional readouts rather than direct measurement of signalling intermediates allowed us to identify key signalling regulators despite remarkably low signalling flux under these conditions (*18*). By quantifying transcription factor activity following systematic perturbation of signalling components, we could capture pathway activity through integrated transcriptional outputs rather than discrete molecular states. This approach is motivated by models of temporal signal integration, as articulated by Purvis and Lahav, and by subsequent optogenetic studies demonstrating that many proximal T-cell signalling intermediates decay rapidly following receptor disengagement (*34, 35*). In contrast, transcriptional outputs integrate signalling over time, accumulating to provide a longer-lived functional record of signalling activity. In the context of T-cell activation, our findings support this framework by demonstrating that robust NF-κB activation and effector responses results from low-amplitude or spatially restricted signalling during direct T-cell-tumour cell engagement.

A key feature of this framework is its compatibility with physiologically relevant stimulation

formats. By interrogating signalling in cell–cell co-culture systems, our approach avoids

reliance on supraphysiological stimulation commonly required to elicit detectable changes in

phosphorylation or kinase activity, and instead captures the integration of complex signal

inputs that arise during direct cellular interactions. Although strong stimulation (most often

achieved using anti-CD3/CD28 bead-based activation) has been instrumental for defining

canonical signalling pathways in global phosphoproteomic studies (*3*, *5*, *11*), it may not reflect

signalling behaviour in physiological settings. This limitation is particularly relevant in studies

of T-cell activation, where signalling outcomes reflect the combined influence of antigen

affinity, co-stimulatory, and inhibitory inputs, such that maximal pathway activation cannot be

assumed to be an appropriate starting point for interrogating authentic signalling processes.

Similar considerations apply to high-dimensional single-cell approaches, including mass

cytometry, which have provided powerful insights when phosphorylation flux is sufficiently

robust, but which typically rely on maximal signalling to infer signalling relationships (*17*, *36*).

Applying this strategy to T-cell receptor signalling revealed that the nodes important for

functional signalling are largely unchanged at different stimulation strengths produced by

differing-quality TCR ligands, whereas there do exist specific nodes that are highly context

dependent. Perturbation of *LCK*, a widely studied SRC-kinase which is known to initiate

receptor signalling from the TCR (31) exhibited a strong dependence on ligand affinity;

disruption of both markedly impaired NF-κB activation under weak stimulation but was

buffered under high-intensity conditions. Quantitative modulation of proximal kinase activity

has been described for LCK. Partial reduction of LCK activity preferentially impairs T cell

responses to low-affinity or low-dose antigen, whereas strong stimulation can partially

compensate for diminished kinase availability (*30*, *37*, *38*). Notably, Cabezas-Caballero and

colleagues demonstrated that altering coreceptor identity (swapping CD8 and CD4) effectively

redistributes and sequesters LCK, thereby tuning TCR sensitivity by modulating the local

availability of active kinase. This work highlights how LCK availability functions as a

quantitative regulator of signalling output. Together, these findings outline how targeting key

proximal signalling components can serve as a mechanism to engineer tuning of cellular

signalling in the context of varied stimulation strengths.

Beyond individual nodes, our data reveal how signalling control is unequally distributed across

molecular families. Rather than treating pathway components as interchangeable, our

perturbation-based analysis quantifies family-level engagement, distinguishing signalling

nodes dominated by a small number of high-impact components from those that rely on

distributed contributions. This is exemplified by the MAPK pathway, in which NF-κB regulation

was driven primarily by a limited set of kinases, consistent with bottlenecked enzymatic

cascades. In contrast, SH2-domain–containing proteins exhibited broad functional

engagement, reflecting their role as proximal adaptors that integrate receptor-derived signals

and connect them to diverse downstream pathways. These patterns are consistent with the

early proposal by Pawson and colleagues, i.e., that signalling networks are assembled from

conserved interaction modules whose combinatorial deployment confers specificity,

robustness, and evolutionary adaptability without reliance on rigid, linear pathways (*39–41*).

Our inferred signalling network therefore supports a modular, hierarchical organisation in

which control is unevenly distributed across molecular layers and funnelled through preferred

pathways under defined conditions.

Notably, many SH2-domain-containing proteins exerted strong functional effects despite

limited transcriptional regulation across T-cell activation states. This suggests that signalling

control at this level operates largely through constitutively expressed components whose

regulatory capacity is governed post-translationally, through phosphorylation, conformational

changes, and dynamic recruitment to signalling complexes. As a result, expression-based

analyses may underestimate this contribution to signalling control, whereas perturbation-

based functional approaches are particularly well suited to reveal their role.

Finally, our experimental framework enabled the identification of previously unrecognised negative regulators of T cell effector function, specifically TRRAP and CTDSPL2. TRRAP functions as a core scaffold for multiple histone acetyltransferase complexes, thereby regulating chromatin accessibility and transcriptional output downstream of diverse signalling pathways (*42*). Consistent with this role, TRRAP depletion resulted in coordinated suppression of proliferative and DNA replication–associated transcriptional programs alongside enrichment of inflammatory and immune signalling modules, indicating a redistribution of transcriptional state rather than global activation. In contrast, CTDSPL2 is a poorly characterised serine/threonine phosphatase implicated in transcriptional regulation and cell-state control, and its role in T cell biology has not previously been examined (*43*). Notably, CTDSPL2 depletion enhanced cytotoxicity with minimal global transcriptional remodelling, suggesting that its effects are mediated primarily through altered signalling dynamics rather than broad transcriptional reprogramming. These findings provide functional validation of nodes identified in the signalome-wide arrayed perturbation screen and illustrate how this approach can uncover regulatory layers that modulate signalling output and effector potential that may be overlooked by pooled genetic screens.

In conclusion, whereas genome-wide perturbation screens and high intensity stimulation-based approaches have been effective for defining essential signalling components, they are less well-suited to dissecting signalling networks dependent on quantitative effects. By emphasising quantitative, time-resolved functional outputs in the setting of physiological stimulation, the framework presented here offers a general strategy for interrogating cell responses as the field moves toward a system-level understanding of signalling.

Materials and Methods

Cell culture

All Jurkat cell lines were cultured in RPMI-1640 medium (Gibco, #21875-034) supplemented with 10% FCS (Gibco, #A5256801, Lot #B3024242RP), 1% Penicillin-Streptomycin-Neomycin (Sigma-Aldrich, #P4083), 1% L-Glutamine (Sigma-Aldrich, #G7513), 1% Sodium Pyruvate (Sigma-Aldrich, #S8636) and 1% HEPES (Gibco, #15630-056). HEK293-T cells for lentivirus production were cultured in DMEM complete medium (Gibco, #41965-039) supplemented as described for the RPMI-1640 Jurkat medium. A375 melanoma cell lines were cultured in DMEM-F12 medium (Gibco, #11320-033) supplemented with 10% FCS, 1% Penicillin-Streptomycin-Neomycin, 1% Sodium Pyruvate and 1% HEPES. Finally, all primary human cells were cultured in RPMI-1640 medium supplemented with 10% human serum, 1% Penicillin-Streptomycin-Neomycin, 1% L-Glutamine, 1% Sodium Pyruvate, 1% non-essential amino acids (NEAA, Gibco, #11140), 1% HEPES and 0.1% B-mercaptoethanol (Gibco, #31350-010).

Production of 1G4$^+$ Jurkat NF-kB GFP reporter cell lines

Jurkat T-cells (ATCC) were first transduced with the Cas9 encoding plasmid pKLV2-EF1a-Cas9Bsd-W, as previously described (26), to produce Cas9 expressing Jurkat cell lines. Next, Cas9$^+$ Jurkat cells were transduced with pSIRV-NF-kB-eGFP construct which was a gift from Peter Steinberger (Addgene plasmid # 118093)(44) prior to single cell cloning and selection of NF-kB GFP expressing clones. Endogenous TCR was then deleted from these lines via electroporation with a TRAC/TRBC sgRNA - Cas9 RNP complex. For the generation of RNPs, sgRNAs targeting TRAC and TRBC1/2 (IDT, TRAC sequence: AGAGTCTCTCAGCTGGTACA, TRBC1/2 sequence: GGAGAATGACGAGTGGACCC) (45) was pre-incubated with Cas9 (IDT, #10007807) at room temperature for 30 minutes. Here, 0.73µL of each sgRNA (400µM) was added to 0.5µL of Cas9 (62µM). Electroporation was

then performed using the Lonza P3 Nucleofector kit (Lonza, # V4SP-3096) as per

manufacturer's instructions. Briefly, one million Jurkat cells were resuspended in 20µL P3

buffer (Lonza, #V4SP-3096), to which the complexed RNP was added. Cells were then moved

to one well of a 16-well electroporation strip for electroporation with a Lonza Nucleofector on

programme #E0115. Cells were immediately recovered from the electroporation strip by the

addition of 80µL of media before returning them to culture. Following deletion of endogenous

TCR, Cas9$^+$ NF-kB GFP$^+$ TCR$^{-/-}$ Jurkats were then transduced with constructs encoding for

the 1G4-TCR (*18*) to produce the final 1G4$^+$ Cas9$^+$ NF-kB GFP$^+$ Jurkat cell lines.

sgRNA design and cloning

To design sgRNA molecules, the top two candidates from CRISPick (Broad Institute) were

selected for each gene. sgRNA sequences were purchased as oligos in both forward and

reverse orientations (Sigma-Aldrich) prior to annealing, and were then cloned into the BbsI

site of the CRISPR sgRNA expression vector pKLV2-U6gRNA5(BbsI)-PGKpuro2ABFPW

(Addgene, #67974) as previously described (*46*). All sgRNA sequences produced for the

signalome library are displayed in supplementary table 1.

Lentivirus production

Lentiviruses were prepared by transfecting HEK293T with the sgRNA expression vector.

Firstly, 500ng sgRNA expression plasmid was combined with 250ng pMD2.G (Addgene,

#12259) and 1.2µg psPAX.2 (Addgene, #12260) in 500µL optiMEM (Gibco, #13985062). To

this, 2µL plus reagent (Thermo Fisher, #15338100) was added prior to mixing by inversion

and incubation at room temperature for 5 minutes. Next, 6µL lipofectamine-LTX (Thermo

Fisher #15338100) was added and the solution was again inverted to mix, prior to incubation

at room temperature for 30 minutes. This optiMEM mixture was then added to HEK293T cells

in a 6 well plate together with 2mL of DMEM medium supplemented as described above.

Lentiviral supernatants were then harvested at 48-72hrs and filtered through a 0.45µm filter.

All lentiviruses were either used immediately or stored at -80°C until use, but were never

freeze-thawed.

Peptide sequences for A375 pulsing

Where indicated, A375 melanoma cell lines were pulsed with the following NY-ESO-1

altered peptide ligands to induce TCR signalling downstream of the 1G4-TCR. Peptide

pulsed A375 cell lines were co-cultured with either Jurkat or primary CD8[+] T-cells as

indicated, and final peptide concentrations are shown throughout the text. All peptides were

ordered from Genscript with a purity >95% and were reconstituted in DMSO upon arrival,

stored at -80°C, and never freeze-thawed more than twice. 9V: SLLMWITQV, 3Y:

SLYMWITQV, 3I: SLIMWITQV, 6T: SLLMWTTQV. Affinities (K_D) for these NY-ESO-1

peptides have been previously described (9V ≅ 7.2µM, 3Y ≅ 30µM, 3I ≅17µM, 6T ≅ 101µM)

(*47*). Finally, for experiments utilising bispecific T-cell engagers, A375 cell lines were pulsed

with the gp100 peptide (10µg/mL, sequence: YLEPGPVTV) as described, which was also

obtained from Genscript at >95% and handled as per the NY-ESO-1 altered peptide ligands.

Arrayed screening of the signalome in 1G4-TCR[+] Jurkat reporter cell lines

1×10^5 1G4-TCR[+] Jurkat NF-kB GFP reporter cells were transduced with 100µL of the arrayed

library lentivirus in a 96-U bottomed plate, in a total of 200µL. Transduction was performed by

centrifugation at 800g, 90min, 32°C. After 48hrs, transduced cells were grown in the presence

of puromycin (1µg/mL final concentration, Gibco, #A11138-03) to select only transduced cells.

At day 7, transduced Jurkat reporter cells were counted via flow cytometry alongside

simultaneously ensuring transduction and selection success via expression of the BFP tag in

the sgRNA plasmid. All cells were 95-100% BFP[+]. Following this counting, all culture volumes

across the arrayed library were normalised to allow for consistent cell numbers to be utilised

in co-cultures with A375 melanoma cells. Here, A375 melanoma cells were treated with

100µg/mL of 6T or 9V NY-ESO-1 peptides (alongside a no peptide control), before plating

5,000 cells in 25µL in a 384 well plate (Falcon, #353961), to which 10,000 Jurkat reporter cells

were added in 25µL. NF-kB signalling pathway activation downstream of TCR engagement

was measured using IncuCyte imaging for 42-48 hours.

596 Arrayed screen analysis in Jurkat cells

Raw IncuCyte time-course data from multiple experiments and plates were aggregated and

annotated with experimental metadata. Condition labels were harmonized, and control wells

were explicitly annotated. For each peptide, signalling output was summarized by calculating

the area under the curve (AUC) using trapezoidal integration. Data was collected for a total of

706 genes with signalome library. AUC values were normalized within each experiment and

condition to the median AUC of empty control wells and log2-transformed for downstream

analysis.

Each gene was targeted by two independent gRNAs. Guide-level effects were computed as

mean log2-normalized AUC ratios across replicates. Gene-level effects were derived using a

rule-based guide collapsing strategy that accounts for guide concordance and effect strength.

When guide effects were concordant (AUC within 1), their median was used; in cases of strong

discordance (particularly when one guide showed a weak effect or when guides had opposite

signs) the guide with the larger absolute effect was retained. Guide concordance was

assessed using Spearman correlation and same-direction agreement.

Gene-level effects were computed independently for each stimulation condition (6T and 9V)

and compared using Spearman correlation. A composite discordance score integrating effect

size differences and sign reversals was used to prioritize condition-specific dependencies.

Genes were annotated with functional families and essentiality, and family-level effect

distributions were visualized to identify pathway-level trends.

Primary CD8$^+$ T-cell transduction

Primary CD8$^+$ T-cells were isolated from healthy donor leukocyte cones obtained from the NHS blood-transfusion service. CD8$^+$ T-cells were isolated using RosetteSep human CD8$^+$ T-cell enrichment cocktail (StemCell, #15023) as per manufacturer's instructions. Isolated CD8$^+$ T-cells were then stimulated for 48 hours prior to transduction with 10µL/mL of CD3/CD28 immunocult (StemCell, #10971) and 50U/mL IL-2 (Biolegend, #589104), with cells at 2x10^6/mL and 2mL/well in a 48 well plate. 250µL of Retronectin solution (Takara Bio, #351178) at 25µg/mL was added to non-treated 24-well tissue culture plates (Costar, #3738) for 24 hours at 4°C prior to transduction. On the day of transduction, Retronectin solution was removed and plates were blocked by addition of 300µL of 2% BSA in PBS (w/v) for 30 minutes at room temperature. The BSA solution was then removed and wells were washed once in 500µL PBS, before addition of 1.5mL of lentiviral supernatant and centrifugation at 2000g for 90 minutes, 32°C, no brake. Following centrifugation, all lentiviral supernatant (except 200-250µL to prevent the well from drying) was removed from the well and 10^6 CD8$^+$ T-cells were seeded into the well in 1mL of complete RPMI with supplementation of IL-2 at 50U/mL. Cells were briefly spun onto the lentiviral coated wells at 2000rpm for 2 minutes, 32°C with the brake applied. Transduction efficiency was assessed 48-72 hours post transduction by measuring BFP expression via flow cytometry. Transduced cells were then selected by the addition of puromycin to 1µg/mL (Gibco, #A11138-03), prior to electroporation with Cas9 protein.

Electroporation of primary CD8$^+$ T-cells

Transduced primary CD8$^+$ T-cells were electroporated with Cas9 protein (IDT, #10007807) once BFP expression had reached >75-80%. Electroporation was performed using the Lonza P3 Primary cell nucleofection kit (Lonza, #V4SP-3096) as per manufacturer's instructions. Here, 10^6 CD8$^+$ T-cells were resuspended in 20µL P3 buffer (Lonza, #V4SP-3096), to which 0.5µL Cas9 (62µM stock) was added. CD8$^+$ T-cells were then transferred to 1 well of the 16-well electroporation strip before electroporation on programme #E0115. Immediately after

electroporation, cells were recovered by the addition of 80µL of complete RPMI-1640 medium supplemented with 50U/mL IL-2. Cells were then plated across 2 wells in a 96 well-flat bottomed tissue culture plate, each in 200µL with IL-2 at 50U/mL. Cells were maintained at 1-$2x10^6$/mL, in the presence of 50U/mL IL-2 and 1µg/mL puromycin, until day 7 post electroporation at which point functional assays were performed. All sgRNA sequences used in the signalome library are displayed in supplementary table 1. Where indicated, perturbations were also performed with RNPs, using independent and non-overlapping guides (IDT) from the signalome library. Guide sequences used for RNP based perturbations are as follows: TRRAP: AGTGGTCTGGTCAACCACCG, CTDSPL2: TTATTCATCAGCCCACGCGG, LCK: GACCCACTGGTTACCTACGA, LCP2: TGAAGAAGTACCACATCGAT, SLA2: GGGACCGGATCAGACACTAC, scRNA: GTATTACTGATATTGGTGGG.

A375 melanoma cytotoxicity and T-cell degranulation assays

At day 7 post Cas9 electroporation, $5x10^4$ A375 mOrange$^+$ melanoma cells were plated in 100µL in a 96 well-flat bottomed tissue culture plate and pulsed with gp100 peptide at a final concentration of 10µg/mL. Cells were incubated at 37°C for at least 4 hours to allow for cell adherence and antigen presentation. A375 cells were then washed once in PBS before addition of $1x10^5$ CD8$^+$ T-cells in 100µL, and 100µL of the gp100-HLA-A2 - CD3 bispecific engager to final concentrations as indicated. Details of this bispecific T-cell engager have been previously described (*18*). Growth of the A375 melanoma cell line was then tracked for approximately 48 hours via IncuCyte imaging, tracking the A375-expressed mOrange fluorescent protein, allowing for quantification of T-cell mediated cytotoxicity (*26*). CD8$^+$ T-cell degranulation was assessed at 4 hours post co-culture by flow cytometry with staining with the following antibodies: CD8 PeCy7, 1:200, Biolegend #344712, CD107α PE, 1:500, Biolegend #328608 and eBioscience fixable viability dye 780, 1:1000, Thermo Fisher #65-0865-18.

RNA sequencing in primary CD8$^+$ T-cells

To assess transcriptional changes with *TRRAP* perturbation, we first targeted *TRRAP* via

lentiviral delivery of sgRNA and electroporation of Cas9 protein as previously described. For

these experiments, three primary CD8$^+$ T-cell donors were transduced with lentiviruses

encoding the TRRAPg1 sequence, or empty plasmid controls. On day 7 post Cas9

electroporation, 5×10^5 CD8$^+$ T-cells were activated on OKT3 coated 48-well tissue culture

plates for 3 hours. OKT3-coated tissue culture plates were prepared in advance, where each

well was treated with 300µL of Ultra-LEAF OKT3 at 1µg/mL (Biolegend, #317326), or PBS

only for unstimulated controls. After 3 hours, cells were recovered, washed in PBS and RNA

was extracted using the Qiagen RNA mini kit (#74104) as per manufacturer's instructions.

Isolated RNA was then sent to Novogene for NGS. FASTQ files were aligned using STAR

version 2.7.3a and quantified using featureCounts version 2.0.6. Ensembl Homo sapiens

GRCh38.86 was used as the reference genome. Lowly expressed genes were filtered out,

and gene expression distributions of each sample were normalised using edgeR version

3.42.2. Differential expression analysis was performed using limma version 3.54.2.

For experiments investigating transcriptional changes associated with *CTDSPL2*

perturbation, primary CD8$^+$ T-cells from two donors were transduced with lentiviruses

encoding for the CTDPSL2g1 or CTDSPL2g2 sequences, alongside empty plasmid controls.

Again, at day 7 post Cas9 electroporation, T-cells were stimulated as described as per

TRRAP perturbations. At 3 hours post stimulation on plate-bound OKT3, T-cells were

collected and resuspended in 50µL DNA/RNA shield (Zymo Research, #R1100-250) and

sent to Plasmidsaurus for NGS. Data handling and analysis was performed by

Plasmidsaurus as follows: Quality of the fastq files was assessed using FastQC v0.12.1.

Reads were then quality filtered using fastp v0.24.0 with poly-X-tail trimming, 3' quality-

based tail trimming, a minimum Phred quality score of 15, and a minimum length

requirement of 50bp. Quality-filtered reads were aligned to the reference genome using

STAR aligner v2.7.11 with non-canonical splice junction removal and output of unmapped reads, followed by coordinate sorting using samtools v1.22.1. PCR and optical duplicates were removed using UMI-based deduplication with UMIcollapse v1.1.0. Alignment quality metrics, strand specificity, and read distribution across genomic features were assessed using RSeQC v5.0.4 and Qualimap v2.3, with results aggregated into a comprehensive quality control report using MultiQC v1.32. Gene-level expression quantification was performed using featureCounts (subread package v2.1.1) with strand specific-counting, multi-mapping read fractional assignment, exons and three prime UTR as the feature identifiers, and grouped by gene_id. Final gene counts were annotated with gene biotype and other metadata extracted from the reference GTF file.

Flow cytometry for assessment of CD8[+] T-cell phenotype

Genetically perturbed CD8[+] T-cells were stimulated on gp100 pulsed A375 melanoma cell lines together with the bispecific T-cell engager as described. At 24 hours post stimulation, T-cells were recovered and transferred to a 96-U bottomed plate before staining with 100µL of antibody mix containing the following antibodies, all used at 1:200 unless otherwise stated: CD8 PE-Cy7, Biolegend, #344750, PD-1 Alexa Fluor 647, Biolegend, #329910, LAG3 FITC, Biolegend, #369308, TIM-3 PE, Biolegend, #345006, 4-1BB BV711, Biolegend, #309832, and eBioscience fixable viability dye 780, 1:1000, Thermo Fisher, #65-0865-18. Surface staining was performed at 4°C for a minimum of 20 minutes, before cells were washed twice and acquired on a BD-LSR Fortessa. Data were then analysed in FlowJo (version 10).

CRISPR-Cas9 mediated targeting of the SH2 domain gene family in primary CD8[+] T-cells

Primary CD8[+] T-cells were isolated from healthy donor leukocyte cones and transduced with lentiviruses encoding sgRNA molecules targeting all SH2 domain containing genes as

previously described (**Supplementary Table 1**). Due to the size of this family, this analysis

was split across 6 independent healthy CD8$^+$ T-cell donors, where each donor was

consistently assayed against an internal control sgRNA panel consisting of guides targeting:

LCK, *LCP2*, *ITK*, *CD8* and *SHC4* together with an empty plasmid control. Following

electroporation of Cas9 to induce perturbation of these genes, CD8$^+$ T-cells were co-cultured

with gp100 pulsed A375 melanoma cell lines, and the bispecific T-cell engager to a final

concentration of 10^{-10}M. Cytotoxic responses of CD8$^+$ T-cells were assessed by measuring

A375 melanoma growth, for a period of 48 hours. After 48 hours, CD8$^+$ T-cells were

harvested and stained with the following antibody cocktail at 4°C for 20 minutes to assess

activation marker expression (CD8 PeCy7, 1:200, Biolegend #344712; CD25 AF647, 1:200,

Biolegend #302618; 4-1BB BV711, 1:200, Biolegend #309832; eBioscience fixable viability

dye 780, 1:1000, Thermo Fisher #65-0865-18).

Donor-donor normalisations in primary CD8$^+$ T-cell killing assays

targeting SH2 domain family genes

In order to account for differing cytotoxic capacities between all CD8$^+$ T-cell donors used to

cover perturbation of the entire SH2 domain gene family, killing curves for each perturbation

were first normalised to their corresponding internal negative control for each individual

donor. This allows for the effect of targeting any given gene within the family to be compared

with other genes in the family that were assayed on separate donors, given the size of the

SH2 family. Next, to produce a single quantitative parameter for each gene, the area under

the curve (AUC) was calculated for all normalised killing curves using the trapz function from

the pracma package in R-studio (version 2024.09.1)(*48*). This analysis produced an

individual value for targeting each gene within the family, which was used to determine

whether this gene behaved as a negative or positive regulator of CD8$^+$ T-cell responses (**Fig**

3F).

Production of primary CD8$^+$ 1G4-TCR$^+$ T-cells and subsequent cytotoxicity assays

CD8$^+$ T-cells isolated from healthy donor leukocyte cones were lentivirally transduced with plasmid encoding for the 1G4-TCR together with a LNGFR tag as previously described (*18*). Next, cells were then stained with TCR Vβ13.1 APC (Biolegend, #362408) and LNGFR FITC (Miltenyi Biotech, #130-113-423) at 1:200 and double positive events were purified by fluorescence-activated cell sorting. At the same time, PBMCs were isolated from three separate healthy donor leukocyte cones, and were subjected to an irradiation dose of 3000rad. Irradiated PBMCs were then centrifuged at 300g for 5 minutes, and resuspended at 2×10^6/mL in complete RPMI-1640 medium. Equal volumes of PBMCs from each of the three donors was then combined, before supplementation of the mixture with 200U/mL IL-2, 1X PHA (500X stock, Invitrogen, #00-4977-03) and 50ng/mL OKT3 (Biolegend, #), to produce T-cell rapid expansion medium (REM). Approximately 1×10^5 sorted LNGFR$^+$ TCR Vβ13.1$^+$ CD8$^+$ T-cells was then added to 10mL REM and cultured for 5 days before counting and re-seeding to 10^6 cells/mL in complete RPMI-1640 medium with 200U/mL IL-2. Cells were then expanded for a further 10-12 days, maintaining IL-2 concentrations at 200U/mL and cell densities at 1-2×10^6/mL throughout. Resultant CD8$^+$ T-cell cultures were checked for purity via flow cytometry and staining with CD8 PeCy7, TCR Vβ13.1 APC and LNGFR FITC. Functional reactivity was then assessed by co-culture with NY-ESO-1 expressing A375 cell lines prior to cryopreservation.

IFN$_γ$ ELISAs

Supernatants were collected at 24 hours post CD8$^+$ T-cell and A375 co-culture, and IFN$_γ$ secretion from CD8$^+$ T-cells into the culture supernatant was assessed by ELISA as per manufacturer's instructions (Biolegend, #430101). Supernatants were either used fresh in ELISAs, or were stored at -20°C prior to analysis but were never freeze-thawed.

CyTOF sample preparation and data analysis

A375 cells were plated in 96-well flat-bottom plates at 5×10^4 cells per well and pulsed with gp100 peptide (10µg/mL) for a minimum of 4 hours. After peptide loading, cells were washed once with PBS and either incubated with the bispecific T-cell engager at a final concentration of 10^{-10}M or left untreated. Where applicable, bispecific binding was allowed to proceed for at least 30 minutes at 37 °C prior to initiation of the experiment.

Isolated CD8$^+$ T cells were subsequently added to either gp100-pulsed bispecific-coated (stimulated) or uncoated (unstimulated) A375 cells and centrifuged (500 g, 1 min) to promote cell contact and initiate activation. At the end of the time course, cells were transferred to 96-well U-bottom plates, pelleted in ice-cold PBS (1800 rpm, 3 min), and fixed in 100 µl of 1.6% PFA. Fixation was carried out at 4 °C for a minimum of 20 minutes, followed by three washes with 200 µl PBS.

Mass cytometric analysis was performed as previously described. Briefly, dead cells were labelled with 0.25 mM ^{194}Cisplatin (Standard BioTools, Cat# 201194), after which cells were incubated overnight at 4 °C with thiol-reactive barcodes. Excess barcodes were quenched using 2 mM GSH and samples were pooled. Cells were then washed in cell staining buffer (CSB; Standard BioTools, Cat# 201068) and stained with extracellular rare-earth metal-conjugated antibodies for 30 minutes at room temperature. Cells were then permeabilised with 0.1% (v/v) Triton X-100/PBS and 50% methanol/PBS prior to staining with intracellular rare-earth metal-conjugated antibodies for 30 minutes at room temperature. Antibodies were cross-linked using 1.6% (v/v) FA/PBS for 10 minutes. Cells were subsequently incubated overnight at 4 °C with 125 nM ^{191}Ir/^{193}Ir DNA intercalator, washed, resuspended in CAS+ supplemented with 2 mM EDTA, and analysed on a CyTOF XT at 200–400 events per second.

Signal intensities were measured by CyTOF at 1, 5, 10, 15, 30, 45, 60, and 120 minutes post-stimulation using a panel of 42 markers. Raw data were preprocessed in FlowJo and

796 subsequently analysed using the CyGNAL python package (github.com/TAPE-
797 Lab/CyGNAL(*14*)) with default settings.

Analysis of published RNA expression and CRISPR dataset

To obtain the lists of signalome genes that were regulated during T cell activation, raw counts

of bulk RNA sequencing dataset were downloaded from Gene Expression Omnibus (GEO).

We filtered the data for CD8+ T cells except for GSE197067, where only pan-T cells were

analyzed. Gene IDs were mapped to gene names by mygene package (Python). Data was

prefiltered by subsetting for genes that have at least 10 counts in more than 20% of samples.

Differential expression analysis was performed using DESeq2 (R)(*49*) using appropriate

model design for study and p values were obtained with the wald test. Log fold change (LFC)

was shrunk with the apeglm method to reduce inflation of lowly expressed genes. Differentially

expressed genes (DEGs) were obtained by thresholding for adjusted p-value < 0.001, and

LFC < 1.5. DEGs of different conditions (e.g. timepoints) of the same dataset were merged by

their union and intersected with our signalome gene list. To obtain the lists of signalome genes

that were identified as hits in pooled CRISPR screening, we downloaded the processed result

files from each study, thresholded the hits following the description of each study and

intersected with our signalome gene list.

References

1. B. M. Invergo, P. Beltrao, Reconstructing phosphorylation signalling networks from quantitative phosphoproteomic data. Essays Biochem. 62, 525–534 (2018).

2. E. J. Needham, B. L. Parker, T. Burykin, D. E. James, S. J. Humphrey, Illuminating the dark phosphoproteome. Sci. Signal. 12, eaau8645 (2019).

3. M. Locard-Paulet, G. Voisinne, C. Froment, M. Goncalves Menoita, Y. Ounoughene, L. Girard, C. Gregoire, D. Mori, M. Martinez, H. Luche, J. Garin, M. Malissen, O. Burlet-Schiltz, B. Malissen, A. Gonzalez de Peredo, R. Roncagalli, LymphoAtlas: a dynamic and integrated phosphoproteomic resource of TCR signaling in primary T cells reveals ITSN2 as a regulator of effector functions. Mol. Syst. Biol. 16, e9524 (2020).

4. R. N. Joshi, N. A. Binai, F. Marabita, Z. Sui, A. Altman, A. J. R. Heck, J. Tegnér, A. Schmidt, Phosphoproteomics reveals regulatory T cell-mediated DEF6 dephosphorylation that affects cytokine expression in human conventional T cells. Front. Immunol. 8, 1163 (2017).

5. P. Ruperez, A. Gago-Martinez, A. L. Burlingame, J. A. Oses-Prieto, Quantitative phosphoproteomic analysis reveals a role for serine and threonine kinases in the cytoskeletal reorganization in early T cell receptor activation in human primary T cells. Mol. Cell. Proteomics 11, 171–186 (2012).

6. R. B. Kitata, W.-K. Choong, C.-F. Tsai, P.-Y. Lin, B.-S. Chen, Y.-C. Chang, A. I. Nesvizhskii, T.-Y. Sung, Y.-J. Chen, A data-independent acquisition-based global phosphoproteomics system enables deep profiling. Nat. Commun. 12, 2539 (2021).

7. P. Mertins, NCI CPTAC, D. R. Mani, K. V. Ruggles, M. A. Gillette, K. R. Clauser, P. Wang, X. Wang, J. W. Qiao, S. Cao, F. Petralia, E. Kawaler, F. Mundt, K. Krug, Z. Tu, J. T. Lei, M. L. Gatza, M. Wilkerson, C. M. Perou, V. Yellapantula, K.-L. Huang, C. Lin, M. D. McLellan, P. Yan, S. R. Davies, R. R. Townsend, S. J. Skates, J. Wang, B. Zhang, C. R. Kinsinger, M. Mesri, H. Rodriguez, L. Ding, A. G. Paulovich, D. Fenyö, M. J. Ellis, S. A. Carr, Proteogenomics connects somatic mutations to signalling in breast cancer. Nature 534, 55–62 (2016).

8. V. Cordo', M. T. Meijer, R. Hagelaar, R. R. de Goeij-de Haas, V. M. Poort, A. A. Henneman, S. R. Piersma, T. V. Pham, K. Oshima, A. A. Ferrando, G. J. R. Zaman, C. R. Jimenez, J. P. P. Meijerink, Phosphoproteomic profiling of T cell acute lymphoblastic leukemia reveals targetable kinases and combination treatment strategies. Nat. Commun. 13, 1048 (2022).

9. A. Subramanian, R. Narayan, S. M. Corsello, D. D. Peck, T. E. Natoli, X. Lu, J. Gould, J. F. Davis, A. A. Tubelli, J. K. Asiedu, D. L. Lahr, J. E. Hirschman, Z. Liu, M. Donahue, B. Julian, M. Khan, D. Wadden, I. C. Smith, D. Lam, A. Liberzon, C. Toder, M. Bagul, M. Orzechowski, O. M. Enache, F. Piccioni, S. A. Johnson, N. J. Lyons, A. H. Berger, A. F. Shamji, A. N. Brooks, A. Vrcic, C. Flynn, J. Rosains, D. Y. Takeda, R. Hu, D. Davison, J. Lamb, K. Ardlie, L. Hogstrom, P. Greenside, N. S. Gray, P. A. Clemons, S. Silver, X. Wu, W.-N. Zhao, W. Read-Button, X. Wu, S. J. Haggarty, L. V. Ronco, J. S. Boehm, S. L. Schreiber, J. G. Doench, J. A. Bittker, D. E. Root, B. Wong, T. R. Golub, A next generation Connectivity Map: L1000 platform and the first 1,000,000 profiles. Cell 171, 1437–1452.e17 (2017).

10. S. Ibáñez-Molero, J. T. M. Pruijs, A. Atmopawiro, F. Wang, A. M. Terry, M. Altelaar,

D. S. Peeper, K. E. Stecker, Phosphoprotein dynamics of interacting T cells and tumor cells by HySic. Cell Rep. 43, 113598 (2024).

11. M. N. Navarro, J. Goebel, C. Feijoo-Carnero, N. Morrice, D. A. Cantrell, Phosphoproteomic analysis reveals an intrinsic pathway for the regulation of histone deacetylase 7 that controls the function of cytotoxic T lymphocytes. Nat. Immunol. 12, 352–361 (2011).

12. G. Franciosa, A. W. P. Jensen, A. Martinez-Val, I. Piga, M. Donia, J. V. Olsen, The proteomics and phosphoproteomics landscape of melanoma under T cell attack, bioRxiv (2025). https://doi.org/10.1101/2025.09.12.675787.

13. A. S. Tocheva, M. Peled, M. Strazza, K. R. Adam, S. Lerrer, S. Nayak, I. Azoulay-Alfaguter, C. J. R. Foster, E. A. Philips, B. G. Neel, B. Ueberheide, A. Mor, Quantitative phosphoproteomic analysis reveals involvement of PD-1 in multiple T cell functions. J. Biol. Chem. 295, 18036–18050 (2020).

14. J. Sufi, X. Qin, F. C. Rodriguez, Y. J. Bu, P. Vlckova, M. R. Zapatero, M. Nitz, C. J. Tape, Multiplexed single-cell analysis of organoid signaling networks. Nat. Protoc. 16, 4897–4918 (2021).

15. C. B. Nattress, R. O'Sullivan, D. Fowler, C. Hutton, P. Vlckova, J. Sufi, M. Buschhaus, E. Basiarz, M. Ramos Zapatero, F. Cardoso Rodriguez, X. Qin, A. Campbell, A. Kanouta, V. S. W. Li, K. Chester, J. Anderson, M. Barisa, J. P. H. Fisher, C. J. Tape, Phenoscaping reveals multimodal γδ T-cell cytotoxicity as a strategy to overcome cancer cell-mediated immunomodulation. Cancer Res. 85, 4415–4432 (2025).

16. X. Qin, F. Cardoso Rodriguez, J. Sufi, P. Vlckova, J. Claus, C. J. Tape, An oncogenic phenoscape of colonic stem cell polarization. Cell 186, 5554–5568.e18 (2023).

17. S. Krishnaswamy, M. H. Spitzer, M. Mingueneau, S. C. Bendall, O. Litvin, E. Stone, D. Pe'er, G. P. Nolan, Systems biology. Conditional density-based analysis of T cell signaling in single-cell data. Science 346, 1250689 (2014).

18. M. Kotowski, D. F. Heraghty, M. Körbel, D. Banik, Z. Zhang, S. Kedia, B. Fu, J. McColl, B. Li, J. Clarke, M. Fellermeyer, Y. Lui, G. Bossi, D. K. Cole, H. Chen, E. Basiarz, S. Sharma, E. Petsalaki, C. J. Tape, B. de Wet, S. F. Lee, A. M. Santos, D. Klenerman, S. J. Davis, T-cell signaling relies on partial CD45-exclusion at sub-micron sized cellular contacts, bioRxiv (2025). https://www.biorxiv.org/content/10.1101/2025.08.29.673015v1.

19. N. Moret, C. Liu, B. M. Gyori, J. A. Bachman, A. Steppi, C. Hug, R. Taujale, L.-C. Huang, M. E. Berginski, S. M. Gomez, N. Kannan, P. K. Sorger, Exploring the understudied human kinome for research and therapeutic opportunities, bioRxiv (2020); https://doi.org/10.37473/fic/10.1101/2020.04.02.022277

20. D. van der Meer, S. Barthorpe, W. Yang, H. Lightfoot, C. Hall, J. Gilbert, H. E. Francies, M. J. Garnett, Cell Model Passports-a hub for clinical, genetic and functional datasets of preclinical cancer models. Nucleic Acids Res. 47, D923–D929 (2019).

21. M. Kanehisa, M. Furumichi, Y. Sato, Y. Matsuura, M. Ishiguro-Watanabe, KEGG: biological systems database as a model of the real world. Nucleic Acids Res. 53, D672–D677 (2025).

22. N. G. Bediaga, H. D. Coughlan, T. M. Johanson, A. L. Garnham, G. Naselli, J. Schröder, L. G. Fearnley, E. Bandala-Sanchez, R. S. Allan, G. K. Smyth, L. C. Harrison, Multi-level remodelling of chromatin underlying activation of human T

cells. Sci. Rep. 11, 528 (2021).

23.   J. Ogando, M. E. Sáez, J. Santos, C. Nuevo-Tapioles, M. Gut, A. Esteve-Codina,
S. Heath, A. González-Pérez, J. M. Cuezva, R. A. Lacalle, S. Mañes, PD-1
signaling affects cristae morphology and leads to mitochondrial dysfunction in
human CD8+ T lymphocytes. J. Immunother. Cancer 7, 151 (2019).

24.   M. Rade, S. Böhlen, V. Neuhaus, D. Löffler, C. Blumert, M. Merz, U. Köhl, S.
Dehmel, K. Sewald, K. Reiche, A time-resolved meta-analysis of consensus gene
expression profiles during human T-cell activation. Genome Biol. 24, 287 (2023).

25.   Y. Lui, E. Jenkins, E. Z. Q. Ng, M. Kotowski, S. J. Mullin, J. Clarke, S. J. Davis, A.
M. Santos, S. Sharma, Cell state and transcription factor modulation during
extended ex-vivo CD8+ T-cell expansion, bioRxiv. (2024).
https://www.biorxiv.org/content/10.1101/2024.07.17.603780v1.full.

26.   S. Sharma, T. Whitehead, M. Kotowski, E. Z. Q. Ng, J. Clarke, J. Leitner, Y.-L.
Chen, A. M. Santos, P. Steinberger, S. J. Davis, A high-throughput two-cell assay
for interrogating inhibitory signaling pathways in T cells. Life Sci Alliance 7 (2024).

27.   E. Shifrut, J. Carnevale, V. Tobin, T. L. Roth, J. M. Woo, C. T. Bui, P. J. Li, M. E.
Diolaiti, A. Ashworth, A. Marson, Genome-wide CRISPR Screens in Primary
Human T Cells Reveal Key Regulators of Immune Function. Cell 175, 1958–
1971.e15 (2018)

28.   J. Carnevale, E. Shifrut, N. Kale, W. A. Nyberg, F. Blaeschke, Y. Y. Chen, Z. Li, S.
P. Bapat, M. E. Diolaiti, P. O'Leary, S. Vedova, J. Belk, B. Daniel, T. L. Roth, S.
Bachl, A. A. Anido, B. Prinzing, J. Ibañez-Vega, S. Lange, D. Haydar, M. Luetke-
Eversloh, M. Born-Bony, B. Hegde, S. Kogan, T. Feuchtinger, H. Okada, A. T.
Satpathy, K. Shannon, S. Gottschalk, J. Eyquem, G. Krenciute, A. Ashworth, A.
Marson, RASA2 ablation in T cells boosts antigen sensitivity and long-term
function. Nature 609, 174–182 (2022)

29.   R. Schmidt, Z. Steinhart, M. Layeghi, J. W. Freimer, R. Bueno, V. Q. Nguyen, F.
Blaeschke, C. J. Ye, A. Marson, CRISPR activation and interference screens
decode stimulation responses in primary human T cells. Science 375, eabj4008
(2022).

30.   J. Cabezas-Caballero, A. Huhn, M. A. Kutuzov, V. Andre, A. Shomuradova, B. W.
A. Peeters, G. M. Gillespie, P. A. van der Merwe, O. Dushek, Generation of T cells
with reduced off-target cross-reactivities by engineering co-signalling receptors.
Nat. Biomed. Eng., doi: 10.1038/s41551-025-01563-w (2026)

31.   S. J. Davis, P. A. van der Merwe, Lck and the nature of the T cell receptor trigger.
Trends Immunol. 32, 1–5 (2011).

32.   E. Güç, A. Treveil, E. Leach, A. Broomfield, A. Camera, J. Clubley, P. Nieto Garcia,
A. Kazachenka, R. Khanolkar, L. Del Carpio, H. Heyn, J. C. Hassel, J. J. Sacco, S.
Stanhope, L. Collins, J. M. Piulats, K. Ranade, A. Benlahrech, Tebentafusp, a T
cell engager, promotes macrophage reprogramming and in combination with IL-2
overcomes macrophage immunosuppression in cancer. Nat. Commun. 16, 2374
(2025).

33.   J. Harper, K. J. Adams, G. Bossi, D. E. Wright, A. R. Stacey, N. Bedke, R. Martinez-
Hague, D. Blat, L. Humbert, H. Buchanan, G. S. Le Provost, Z. Donnellan, R. J.
Carreira, S. J. Paston, L. U. Weigand, M. Canestraro, J. P. Sanderson, S. Botta
Gordon-Smith, K. L. Lowe, K. A. Rygiel, A. S. Powlesland, A. Vuidepot, N. J.
Hassan, B. J. Cameron, B. K. Jakobsen, J. Dukes, An approved in vitro approach
to preclinical safety and efficacy evaluation of engineered T cell receptor anti-CD3

bispecific (ImmTAC) molecules. PLoS One 13, e0205491–e0205491 (2018).

34.   J. E. Purvis, G. Lahav, Encoding and decoding cellular information through
signaling dynamics. Cell 152, 945–956 (2013).

35.   M. J. Harris, M. Fuyal, J. R. James, Quantifying persistence in the T-cell signaling
network using an optically controllable antigen receptor. Mol. Syst. Biol. 17, e10091
(2021).

36.   S. C. Bendall, E. F. Simonds, P. Qiu, E.-A. D. Amir, P. O. Krutzik, R. Finck, R. V.
Bruggner, R. Melamed, A. Trejo, O. I. Ornatsky, R. S. Balderas, S. K. Plevritis, K.
Sachs, D. Pe'er, S. D. Tanner, G. P. Nolan, Single-cell mass cytometry of
differential immune and drug responses across a human hematopoietic continuum.
Science 332, 687–696 (2011).

37.   H. Dada, M. L. Dustin, Goldilocks and the three TILs. J. Exp. Med. 219 (2022).

38.   V. Horkova, A. Drobek, D. Paprckova, V. Niederlova, A. Prasai, V. Uleri, D.
Glatzova, M. Kraller, M. Cesnekova, S. Janusova, E. Salyova, O. Tsyklauri, T. A.
Kadlecek, K. Krizova, R. Platzer, K. Schober, D. H. Busch, A. Weiss, J. B. Huppa,
O. Stepanek, Unique roles of co-receptor-bound LCK in helper and cytotoxic T
cells. Nat. Immunol. 24, 174–185 (2023).

39.   T. Pawson, J. D. Scott, Signaling through scaffold, anchoring, and adaptor
proteins. Science 278, 2075–2080 (1997).

40.   T. Pawson, P. Nash, Assembly of cell regulatory systems through protein
interaction domains. Science 300, 445–452 (2003).

41.   W. A. Lim, T. Pawson, Phosphotyrosine signaling: evolving a new cellular
communication system. Cell 142, 661–667 (2010).

42.   D. Detilleux, P. Raynaud, B. Pradet-Balade, D. Helmlinger, The TRRAP
transcription cofactor represses interferon-stimulated genes in colorectal cancer
cells. Elife 11, e69705 (2022).

43.   M. Li, L. Chen, F. Yu, H. Mei, X. Ma, K. Ding, Y. Yang, Z. Rong, CTDSPL2
promotes the progression of non-small lung cancer through PI3K/AKT signaling via
JAK1. Cell Death Discov. 10, 389 (2024).

44.   S. Jutz, A. Hennig, W. Paster, Ö. Asrak, D. Dijanovic, F. Kellner, W. F. Pickl, J. B.
Huppa, J. Leitner, P. Steinberger, A cellular platform for the evaluation of immune
checkpoint molecules. Oncotarget 8, 64892–64906 (2017).

45.   Y.-L. Chen, J. S. W. Ng, R. Ottakandathil Babu, J. Woo, J. Nahler, C. S. Hardman,
P. Kurupati, L. Nussbaum, F. Gao, T. Dong, K. Ladell, D. A. Price, D. A. Duncan,
D. Johnson, U. Gileadi, H. Koohy, G. S. Ogg, Group A Streptococcus induces
CD1a-autoreactive T cells and promotes psoriatic inflammation. Sci. Immunol. 8,
eadd9232 (2023).

46.   S. Sharma, G. J. Wright, Cell Surface Receptor Identification Using Genome-Scale
CRISPR/Cas9 Genetic Screens (2020; https://www.jove.com/t/60803/cell-surface-
receptor-identification-using-genome-scale-crisprcas9).

47.   M. Aleksic, O. Dushek, H. Zhang, E. Shenderov, J.-L. Chen, V. Cerundolo, D.
Coombs, P. A. van der Merwe, Dependence of T cell antigen recognition on T cell
receptor-peptide MHC confinement time. Immunity 32, 163–174 (2010).

48.   H. W. Borchers, Pracma: Practical Numerical Math Functions (https://cran.r-
project.org/web/packages/pracma/index.html).

49.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2, bioRxiv (2014).
https://doi.org/10.1101/002832.

Author contribution:

The experiments were conceived by S.S, E.P, S.J.D, and C.J.T, and the methodology developed by J.C and S.S. The experimental work was undertaken by J.C, S.S, C.T.G, E.B, M.K, A.M.S, M.T. V, P.S, J.X.L. Data analysis was performed by J.C, H.C, and S.S. The work was supervised by S.S, E.P, S.J.D, C.J.T, and funding was acquired by J.C, S.S, E.P, S.J.D, C.J.T. The original manuscript was written by J.C, S.S, and E.P and revised by H.C, M.K, A.M.S, C.J.T and S.J.D.

Acknowledgements

This work was funded by MRC (grant MR/W025507/1 to E.P., S.S., C.T., and S.J.D.) and Wellcome grant to S.S (317519/Z/24/Z), and MSIF pump priming grant to J.C (BRDNB420).