1

Restricted amino acid diversity alters enzymatic phosphoryl-transfer catalysis

Sota Yagi[1,2], Subrata Dasgupta[2], Shusuke Tagami[2], Satoshi Akanuma[1]

[1]Faculty of Human Sciences, Waseda University, 2-579-15 Mikajima, Tokorozawa, Saitama

359-1192, Japan

[2]RIKEN Center for Integrative Medical Sciences, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama,

Kanagawa, 230-0045, Japan.

***Corresponding author:** Satoshi Akanuma

Phone: +81-4-2947-6727

Email: akanuma@waseda.jp

Classification: BIOLOGICAL SCIENCES – Evolution

Keywords: amino acid composition, enzyme evolution, origin of life
22

This file includes:

Main Text

Tables 1 and 2

Figures 1 to 4

Significant Statement

The first enzymes arose on early Earth, where only a limited set of amino acids was likely

available. Whether such limitations reduce catalytic efficiency or reshape enzyme function is

still unclear. We reconstructed a model of an ancestral enzyme using a simplified amino acid

set approximating those available on early Earth and found that restricting amino acid

diversity replaced its original catalytic activity with an alternative function that produces ATP

from two ADP molecules. Structural and mutational analyses indicate that this shift involves a

reorganization of the active site. These results provide experimental evidence that limiting

amino acid diversity can reshape catalytic function and may have influenced the emergence of

catalytic activities during enzyme evolution.

Abstract

How amino acid composition shapes enzymatic function remains a central question in

molecular evolution. Early proteins were likely composed of a limited set of amino acids, yet

the catalytic properties of proteins under such compositional constraints are still poorly

understood. Here we show that restricting amino acid repertoires can replace native enzymatic

activity with an alternative phosphoryl-transfer reaction that disproportionates ADP into ATP

and AMP. Using a reconstructed ancestral nucleoside diphosphate kinase (NDK) variant

composed of a restricted amino acid set, we demonstrate this reaction, which is not observed

in extant NDKs but is chemically analogous to that catalyzed by adenylate kinase. Although

modest in catalytic rate, X-ray crystallography, molecular dynamics, and mutational analyses

reveal a distinct active-site organization in which aspartate and arginine residues

cooperatively coordinate Mg^{2+} and support phosphoryl transfer through a noncanonical

mechanism. Lysine can substitute for arginine under these constraints while retaining activity.

Together, these findings show that restricting amino acid diversity can remodel active sites

and promote alternative phosphoryl-transfer reactions, illustrating how limitations in amino

acid availability could influence catalytic functions in early enzyme evolution.

Introduction

Understanding how proteins first emerged is a central issue in elucidating the transition

from prebiotic chemistry to biological evolution[1-3]. One of the key challenges in this context

is determining how the building blocks of proteins were initially acquired[4,5]. While modern

proteins are generally composed of 20 L-amino acids, and it is believed that the last universal

common ancestor (LUCA) could already synthesize proteins using 20 types of amino acids[6-8],

the amino acid composition of proteins prior to LUCA remains unclear[9-11]. To our best

knowledge, there is no experimental evidence indicating that all 20 of the current

proteinogenic amino acids were available during the earliest stages of life. More likely, these

amino acids were instead gradually incorporated through the evolution of biosynthetic

pathways[12-14]. Indeed, this proposal is supported by several lines of evidence. For example,

ten amino acids were synthesized in Miller's spark discharge experiment[15-17], eight were

identified in the Murchison meteorite[18,19], and recent analyses of samples from asteroid Ryugu

detected 11 proteinogenic amino acids, ten of which overlapped with those from the Miller

experiment[20]. These findings suggest that primitive proteins were synthesized using only a

limited subset of amino acids that were available on the early Earth, prior to the emergence of

endogenous biosynthetic pathways. Specifically, the set of ten amino acids synthesized in

Miller's spark discharge experiment are often referred to as "prebiotic amino acids," because

they plausibly represent the amino acid repertoire available on early Earth[12,21,22]. However,

modern enzymes rely heavily on the chemical diversity of side chains for substrate binding,

transition-state stabilization, and catalysis. This raises a fundamental question of how

primitive polypeptides, composed of a limited set of amino acids, could have folded into

tertiary structures that exhibited enzymatic functions. Resolving this question is essential for

reconstructing the molecular steps that led from simple prebiotic chemistry to the complex

metabolic networks found in all known life.

The plausibility of primitive protein synthesis using the prebiotic set of amino acids has

been supported by multiple lines of evidence. The set of prebiotic amino acids have been

shown to associate with self-assembled fatty acid membranes, which enhances membrane

stability. The localized concentration of these amino acids on the membrane surfaces could

have facilitated peptide synthesis[21]. Computational and theoretical studies have further

predicted that hypothetical proteins composed of the ten prebiotic amino acids would exhibit

folding preferences and structural characteristics similar to those of modern proteins[23].

Moreover, another study suggested that among all possible combinations of ten amino acids

drawn from the twenty used in modern proteins, the prebiotic set is particularly favorable for

forming robust protein backbones[14]. Collectively, these findings indicate that the prebiotic

amino acid subset was not only abundant in the early Earth environment but also inherently

suited to forming stable, protein-like structures.

Nucleoside diphosphate kinase (NDK) catalyzes the transfer of the γ-phosphate from a

nucleoside triphosphate to a nucleoside diphosphate, thereby maintaining the intracellular

balance of nucleoside triphosphates[24]. Previously, we performed ancestral sequence

reconstruction on NDK to produce several ancestral NDKs that might have existed in the last

common ancestors of Archaea and Bacteria[25, 26]. Next, we systematically restricted the amino

acid composition of Arc1, one of the archaeal ancestral NDKs, while preserving its catalytic

function[27]. This approach yielded simplified proteins composed of only 13 amino acid types

that retained detectable enzymatic activity and were thermostable above 70°C. Notably, the

majority of amino acids in these simplified proteins overlapped with the aforementioned

prebiotic amino acids. Furthermore, when histidine, asparagine, and tyrosine, which are not

listed as prebiotic amino acids, were removed from a 13-amino acid variant, its high thermal

stability was retained, although its catalytic activity was completely lost.

In this study, we extended our previous work by designing new variants with distinct amino

acid compositions. Interestingly, one variant that was constructed using a 12-amino-acid set

consisting of prebiotic amino acids supplemented with lysine and arginine exhibited a

noncanonical catalytic activity: the synthesis of ATP and AMP from two ADP molecules,

distinct from the canonical NDK reaction. This unexpected gain-of-function highlights a

fundamental insight; namely, constraining amino acid composition can result in the

emergence of alternative catalytic functions. ATP synthesis is thought to be crucial in early

bioenergetics[28]. Thus, the emergence of ATP synthesis activity may reflect the role of early

proteins composed of a restricted set of amino acids in primitive energy metabolism. Our

findings provide a compelling example that reduced chemical complexity may have shaped

the functional landscape of early proteins during the origin and early evolution of life.

Results

A 12-amino acid ancestral NDK variant catalyzes noncanonical ATP synthesis

To investigate the relationship between amino acid composition and catalytic function in

ancestral proteins, we first constructed two new variants of the reconstructed ancestral NDK

Arc1. One of these variants, designated Arc1-16, was designed with a unique 16-amino acid

alphabet that excluded the residues histidine, asparagine, tyrosine and cysteine. Histidine,

asparagine and tyrosine residues are thought to play important roles in the catalytic activity of

Arc1[27], and cysteine is a residue not present in original Arc1 (Table 1; Fig. S1; Data S1). In

addition, we constructed a 12-amino acid variant, Arc1-12KR, which was composed of the ten

prebiotic amino acids supplemented with two basic residues, lysine and arginine (Table 1; Fig.

S1; Data S1). Note, the N-terminal Met was excluded from the compositional definition.

Unless otherwise noted, all protein variants were expressed with an N-terminal His$_6$-tag.

For Arc1-12KR, we confirmed that the His-tag did not affect folding or enzymatic activity by

comparing the tagged and tag-free forms using circular dichroism (CD) spectroscopy, and

ATP-synthesizing assays (Fig. S2). In addition, the constructs lacking aromatic amino acids

were appended with a Gly-Gly-Trp tag at the C-terminus to enable spectroscopic detection of

these proteins.

NDK catalyzes the transfer of the γ-phosphate from a nucleoside triphosphate to a

nucleoside diphosphate (Fig. 1A). To assess whether the new variants retained this canonical

activity, enzyme assays were performed in which ATP production was monitored using GTP

and ADP as substrates. Because both variants lacked histidine, a residue shown to be essential

for catalysis, none were expected to retain the original enzymatic activity. Consistent with this

expectation, Arc1-16 did not display detectable catalytic activity under these conditions (Fig.

1C; Data S2). Intriguingly, however, ATP synthesis activity was detected for Arc1-12KR (Fig.

1C inset). More surprisingly, ATP formation was also observed when GTP, one of the

substrates, was omitted from the reaction mixture (Fig. 1D; Table 1; Data S2). This result

prompted us to hypothesize that Arc1-12KR was catalyzing an alternative reaction, namely,

the formation of ATP and AMP from two molecules of ADP (Fig. 1B). To test this hypothesis,

we analyzed the reaction products using high-performance liquid chromatography (HPLC).

Prior to incubation, only a single peak corresponding to ADP was observed in the reaction

mixture (Fig. 1E; Fig. S3). After incubation with Arc1-12KR at 60 °C for 10 min, two

additional peaks emerged, corresponding to ATP and AMP (Fig. 1E; Fig. S3). These results

confirmed that Arc1-12KR catalyzes the formation of ATP and AMP from two molecules of

ADP.

To examine whether this unexpected activity was specific to Arc1-12KR, we tested the

original Arc1 and two 13-amino acid-alphabet variants, Arc1-13FKR and Arc1-13KMR, in

which all phenylalanine residues or all methionine residues present in the original Arc1

sequence were restored to the Arc1-12KR scaffold, respectively (Table 1; Fig. S1; Data S1).

When the activity was measured with ADP as the only substrate, no detectable ATP

production was observed above background levels for these constructs (Fig. 1D; Table 1; Data

S2), indicating that the ability to catalyze ADP disproportionation is unique to Arc1-12KR

under the conditions used in this study.

Next, we examined the rate of ATP formation as a function of ADP concentration (Fig. 1F;

Data S2). In the low ADP concentration range (0.3–1.0 mM), the reaction rate determined

from the linear phase of ATP formation exhibited an approximately second-order dependence

on ADP concentration (Fig. 1G). This nonlinear relationship cannot be readily explained by a

canonical ping-pong mechanism and is more consistent with a mechanism involving the direct

transfer of a phosphate between two ADP molecules.

We also examined whether Arc1-12KR is able to catalyze the reverse reaction. Upon

incubating the enzyme with ATP and AMP under the standard assay conditions (2 mM Mg^{2+}),

no detectable formation of ADP was observed (Fig. S4A). However, increasing the Mg^{2+}

concentration to 10 mM resulted in detectable ADP formation from ATP and AMP (Fig. S4B),

suggesting that the reverse reaction depends on Mg^{2+} availability relative to nucleotide

concentration. Furthermore, when GDP was used in place of ADP, no reaction products were

detected (Fig. S4C), indicating that the observed activity is specific to adenylates and does not

reflect a generalized NDK activity.

To assess the oligomeric states of Arc1-12KR and its related variants, we performed

analytical gel filtration (Fig. S5). Arc1-13FKR and Arc1-13KMR eluted at volumes

corresponding to their respective hexameric assemblies, similar to the oligomeric state of

Arc1 and many extant NDKs[29-31]. In contrast, Arc1-12KR eluted as a single peak

corresponding to a dimer. Notably, Arc1-12KR, which uniquely exhibited the altered catalytic

activity, existed predominantly as a dimer, whereas the hexameric variants did not display this

activity. Structural analysis indicates that the interfaces involved in hexamer formation do not

directly contribute to the active site. Therefore, while the change in oligomeric state correlates

with the altered activity, a causal relationship between oligomerization and functional

divergence cannot be established.

Dissecting the roles of arginine and lysine in noncanonical ATP synthesis

To further dissect the contribution of basic amino acids in the newly observed catalytic

activity of Arc1-12KR, we designed three additional variants: Arc1-11K, in which all arginine

residues were replaced with lysine (Table 1; Fig. S1; Data S1); Arc1-11R, in which all lysine

residues were replaced with arginine (Fig. S1; Data S1); and Arc1-10, in which both lysine

and arginine residues were replaced with prebiotic amino acids selected from the remaining

ten used in Arc1-12KR (Fig. S1; Data S1). The corresponding genes were codon-optimized

for expression in *Escherichia coli* and artificially synthesized. Despite codon optimization, no

expression was detected for either Arc1-11R or Arc1-10 in *E. coli* under our standard

expression conditions, suggesting that these variants did not fold into an adequate tertiary

structure and were readily degraded. Due to this poor expression, Arc1-11R and Arc1-10 were

not purified and structurally characterized. In contrast, Arc1-11K was expressed at high levels

and the recombinant protein successfully purified for further analysis.

Gel filtration chromatography revealed that Arc1-11K, like Arc1-12KR, exists

predominantly as a dimer (Table 1; Fig. S5). Far-UV circular dichroism spectroscopy

demonstrated that Arc1-11K exhibits a secondary structure content highly reminiscent of

Arc1-12KR (Fig. S6). These findings indicated that the lysine residues could be substituted

for arginine without disrupting the overall protein fold. Catalytic activity assays using ADP as

the sole substrate showed that Arc1-11K retained the ability to catalyze the conversion of two

ADP molecules into ATP and AMP (Fig. 1D; Table 1; Data S2). However, the rate of reaction

was approximately half that of Arc1-12KR, suggesting arginine contributes more effectively

than lysine to this noncanonical catalytic function.

Structural basis of histidine-independent ATP synthesis mediated by Arc1-12KR

To gain insight into the structural basis of the unexpected catalytic activity observed in

Arc1-12KR, we determined the crystal structure of the apo form of this variant at 3.3 Å

resolution by X-ray crystallography (Table S1). The asymmetric unit contained six

independent dimeric assemblies (Fig. S7A), consistent with gel filtration data indicating that

the protein predominantly exists as a dimer in solution. Superposition of the six dimers

revealed notable conformational heterogeneity within a loop region (residues 49–59) located

adjacent to the putative active site (Figs. S7B and S7C). This heterogeneity suggests inherent

flexibility at the loop region, which may be relevant to the catalytic mechanism. Except for

this flexible motif, the six protomers exhibited highly conserved backbone conformations,

indicating that the overall fold is well maintained despite the reduced amino acid alphabet.

Next, the crystal structure of Arc1-12KR was compared with that of the ancestral Arc1 protein

(Fig. 2A). The two structures aligned closely, with the majority of backbone atoms showing

excellent overlap. However significant deviation was observed in the loop formed by residues

49-59, which is located near the active-site and likely involved in substrate recognition and

positioning (Fig. S8).

To directly visualize substrate binding, we attempted co-crystallization of Arc1-12KR with

ADP. However, electron density maps revealed no interpretable density corresponding to ADP

in any of the datasets obtained. Instead, a structural model of the Arc1-12KR–Mg^{2+}–ADP

complex was generated using molecular dynamics (MD) simulations (Fig. S9). Starting from

the most "opened" apo conformation observed among the six crystallographically independent

dimers, ADP molecules were docked into the putative active site. In addition, MD simulations

were performed for the apo form of Arc1-12KR to provide a reference for comparison with

the ADP-bound model.

The modeled complex structure revealed that ADP is located in the canonical binding

pocket, with its phosphate groups coordinated by a central Mg^{2+} ion (Figs. 2B and 2C). In the

modeled active site, the side chain guanidinium groups of Arg85 and Arg102 directly interact

with the β-phosphate group of ADP through hydrogen bonds, indicating that these two

arginine residues play important roles in catalysis. During the MD simulation, three water

molecules (W1, W2, and W3) exhibited nearly 100% occupancy and contributed to ADP

binding through hydrogen-bond networks (Fig. 2D). The hydroxyl group of Thr91 forms a

hydrogen bond with ADP via W1. The side chain amino group of Lys55 forms a hydrogen

bond with the α-phosphate of ADP via W2. In addition, the side chain guanidinium group of

Arg102 interacts with ADP through W3. In the apo form, the Mg^{2+} ion adopts a distorted

octahedral coordination geometry, involving the side chain carboxylate groups of Asp115 and Asp118, the main-chain carbonyl oxygen of Ala116, and one water molecule (W5) in the equatorial plane, while two additional water molecules (W4 and W6) occupy the axial positions. In the ADP-bound form, these water molecules are displaced by ADP, and the Mg^{2+} ion is coordinated by the side chain carboxylate groups of Asp115 and Asp118 as well as the α- and β-phosphate groups of ADP.

The active sites of the modeled Arc1-12KR and the *Pyrococcus horikoshii* OT3 NDK-ADP-Mg^{2+} complex (PDB ID: 2DY9) are compared in Fig. 3 and superimposed in Fig. S10. Several critical differences between the two active sites were observed. The bound ADP adopts different positions in the active site and, therefore, the phosphorus atom of the β-phosphate is shifted by ~1.3 Å, and the position of the coordinated Mg^{2+} ion differs by ~2.2 Å between the two structures. Residue 115, which is a catalytically essential histidine in all known NDKs, is replaced by an aspartate in Arc1-12KR. In canonical NDKs, including that from *P. horikoshii*, the conserved histidine residue functions as a transient phosphoryl acceptor during catalysis. In Arc1-12KR, however, Asp115 is positioned in close proximity to the catalytic Mg^{2+} ion, suggesting that it contributes primarily to metal coordination rather than directly participating in phosphoryl transfer. The carboxylate moiety of Asp118 in Arc1-12KR is also positioned close to the Mg^{2+} ion, whereas the corresponding aspartate residue in *P. horikoshii* NDK is oriented away from the active site. In contrast, Arg85 and Arg102 in Arc1-12KR and the corresponding arginine residues in *P. horikoshii* NDK are similarly positioned and oriented in the active site. This conservation of relative spatial positioning and orientation suggests that these arginine side chains play comparable roles in phosphate binding. The structural analyses suggest that Arc1-12KR adopts a distinct catalytic strategy primarily derived from alterations to acidic residues and metal coordination, while preserving the canonical phosphate-binding arginine residues. The active site architecture in

Arc1-12KR reflects adaptation resulting from a restricted amino acid alphabet. These findings

demonstrate how simplified sequence information can generate functional enzymatic folds

with novel mechanistic features.

Mutational analysis identifies residues critical for noncanonical ATP synthesis

Next, we aimed to identify crucial residues for the newly acquired ATP synthesis activity.

Mutants of four candidate residues suggested from analysis of the crystal structure of

Arc1-12KR and its MD-generated ADP-bound model were generated. Asp115 and Asp118,

both oriented toward the catalytic Mg^{2+} ion, were individually substituted with alanine. The

resultant D115A and D118A mutants retained 60% and 68% of the original ATP synthesizing

activity, respectively (Table 2; Fig. S11; Data S2). However, simultaneous substitution of both

residues drastically reduced this activity to ~6% (Table 2; Fig. S11; Data S2), suggesting that

the presence of at least one negatively charged side chain is important for coordinating Mg^{2+}

and for electrostatic contributions to ADP binding.

Arg85 and Arg102, predicted to interact with the phosphate moieties of ADP, were

individually replaced with methionine. R102M maintained approximately 50% of the original

ATP synthesizing activity, but R85M resulted in a significant reduction in activity to ~13%

(Table 2; Fig. S11; Data S2). The double mutant lacking both arginine residues displayed only

~4% activity (Table 2; Fig. S11; Data S2), indicating the cooperative role of Arg85 and

Arg105 in positioning the substrate and stabilizing the transition state. Notably, Arg85 appears

to play a more central role than Arg102 in this variant, despite the fact that, in extant NDKs,

both Arg85 and Arg102 are identified as catalytically important residues[32].

Intriguingly, the variant Arc1-11K, in which all arginine residues including Arg85 and

Arg102 were replaced with lysine, retained 50% of the ATP-synthesizing activity of

Arc1-12KR (Fig. 1D; Table 1; Data S2). Therefore, the catalytic roles of Arg85 and Arg102

can be compensated by lysine, although with reduced efficiency. These findings likely reflect

a compensatory redistribution of catalytic roles that emerged under the constraints of being

restricted to a prebiotically plausible set of amino acids.

Discussion

In this study, we show that ATP-synthesizing activity emerges in the reconstructed kinase-like

enzymes Arc1-12KR and Arc1-11K even though they are built only from prebiotic and basic

amino acids. As such, these enzymes operate under severe compositional constraints. The

reaction converting two molecules of ADP into ATP and AMP is, among known enzymes,

catalyzed by adenylate kinases[33], but it has not been observed in extant NDKs or in the

reconstructed ancestral NDK Arc1. Thus, within the NDK family, this ATP-synthesizing

activity represents a highly unusual and genuinely novel function. From an evolutionary

perspective, this activity may reflect a primitive mode of phosphoryl transfer that predates the

appearance of histidine-containing active sites. A restricted amino-acid alphabet lacking

histidine, asparagine, tyrosine and other late-emerging residues[34] would have constrained

conventional catalytic strategies and forced a repurposing of residue side chains such as

arginine and aspartate for substrate binding, charge compensation and possibly phosphate

transfer. Our results therefore indicate that changes in biochemical function can arise not only

from increasing molecular complexity but also from the adaptive exploitation of

physicochemical constraints.

The ability of Arc1-12KR to catalyze a reaction that is absent from its modern homologs

suggests that structural plasticity and chemical adaptability may have played key roles in the

functional diversification of primordial enzymes. Rather than representing an entirely novel

catalytic activity, the observed enzymatic function may instead reveal a latent capacity that

was accessible on early protein scaffolds but was subsequently lost as more efficient and

specialized mechanisms evolved. Our findings demonstrate that compositional simplification

of biocatalysts can unmask alternative activities that are no longer retained by modern, fully

optimized enzymes.

Consistent with this notion, Arc1-12KR and Arc1-11K catalyze the same overall reaction as

modern adenylate kinases despite having no detectable homology and entirely distinct tertiary

structures. This functional convergence implies that phosphoryl transfer reactions may have

arisen independently multiple times during early evolution from nonhomologous protein

scaffolds. The canonical NDK reaction, which involves transfer of the γ-phosphate from NTP

to NDP, is nearly universal across extant life. By contrast, the ADP disproportionation

reaction characterized here may reflect an older stage of molecular evolution. In environments

where NTPs were scarce or absent, a mechanism that uses ADP as both phosphate donor and

acceptor could have provided a plausible route for transient ATP formation and energy

recycling in prebiotic systems. As illustrated in Fig. 4, we envisage a prebiotic world in which

kinase-like proteins resembling Arc1-12KR/Arc1-11K with a limited amino acid repertoire

generated ATP via ADP disproportionation. With the subsequent expansion of the amino acid

alphabet, this chemistry was later superseded by modern NDKs, adenylate kinases and rotary

ATPases.

Although the ADP disproportionation reaction is thermodynamically close to neutral, its

evolutionary and metabolic implications could be substantial. Local or transient increases in

ATP concentration would have facilitated downstream biosynthetic reactions and early

nucleotide-dependent processes[28]. At the same time, the co-produced AMP could have

contributed to nucleotide signaling functions in primitive biochemical networks[33]. In

proto-cellular environments, such reactions, when combined with compartmentalization or

ion gradients, may have helped to create and sustain non-equilibrium states, thereby

supporting the emergence of proto-metabolic organization before dedicated ATP synthase

systems evolved[35,36].

Canonical NDKs catalyze γ-phosphate transfer from NTP to NDP via a ping-pong

mechanism involving a transient phosphohistidine intermediate[37]. This mechanism depends

on a conserved histidine residue (His115 in Arc1), which serves as the nucleophile and is

transiently phosphorylated during catalysis. By contrast, Arc1-12KR entirely lacks histidine

and catalyzes disproportionation between two ADP molecules to yield ATP and AMP.

Although aspartate residues have been reported to form transient phosphoaspartate

intermediates in other enzymatic contexts[38,39], mutational analyses indicate that neither

Asp115 nor the nearby Asp118 is essential for ATP synthesis in Arc1-12KR, as alanine

substitutions at these positions retain substantial activity (Table 2). Indeed, even the

simultaneous substitution of both residues results in reduced but detectable catalytic activity.

These observations argue against the formation of a covalent phosphoaspartate intermediate in

Arc1-12KR and instead are consistent with a histidine-independent phosphoryl-transfer

mechanism distinct from that of canonical NDKs. This interpretation is further supported by

kinetic analyses showing that, at low ADP concentrations, the rate of ATP formation exhibits

an approximately second-order dependence on ADP concentration (Fig. 1G). This kinetic data

cannot be readily explained by a ping-pong mechanism and is more consistent with direct

transfer of a phosphate moiety between two molecules of ADP.

Under the conditions used in this study (2 mM Mg^{2+}), the ADP disproportionation reaction

approached saturation at ~1.6 mM ADP (Fig. 1F), and no detectable reverse reaction (ATP +

AMP → 2 ADP) was observed (Fig. S4A). However, increasing the Mg^{2+} concentration to 10

mM resulted in detectable ADP formation from ATP and AMP (Fig. S4B), indicating that the

apparent irreversibility reflects Mg^{2+} availability relative to nucleotide concentration. Given

that nucleotide substrates are likely to enter the active site as Mg–ADP complexes, depletion

of free Mg^{2+} at high ADP concentrations could account for the apparent saturation behavior.

Moreover, preferential coordination of Mg^{2+} by ATP may reduce formation of Mg–AMP

complexes under Mg^{2+}-limiting conditions, thereby suppressing the reverse reaction.

Our structural and mutational analyses suggest that the noncanonical activity relies on

cooperative interactions among Mg^{2+}-coordinating aspartate and phosphate-binding arginine

residues. Together, these residues may partially substitute for the catalytic role normally

played by histidine in canonical NDKs. This occurs by organizing the Mg^{2+}-ADP complex,

stabilizing the negative charge on the phosphate groups, and facilitating phosphoryl transfer,

which does not involve a covalent intermediate. These observations indicate that, when the

amino acid alphabet is constrained, alternative catalytic mechanisms can emerge that diverge

from those employed by modern enzymes.

The variant completely lacking basic amino acids (Arc1-10) failed to fold into a stable

structure, highlighting a paradox; namely, although basic residues were probably rare under

prebiotic conditions[40], they seem to be essential for proper protein folding and activity.

Previous work has shown that primitive proteins can be stabilized in high-salt environments[22],

and that proteins devoid of basic residues can interact with RNA via metal-ion mediation[41].

Future experiments should clarify how environmental factors, particularly the availability of

mono and divalent cations, might have compensated for the absence of basic residues and

enabled enzymatic activity in the early Earth environment.

However, we cannot rule out the possibility that lysine, arginine, or both these amino acids

were in fact synthesized abiotically and already present in early protein synthesis[42,43]. Tawfik

and co-workers proposed that the first nucleic acid-binding proteins may have arisen from

short, simple sequences containing ornithine, which is not used in extant proteins[44]. By

contrast, Thoma and Powner showed that ornithine and diaminobutyric acid rapidly cyclize

intramolecularly, blocking further peptide extension, whereas the shortest basic α-amino acid

diaminopropionic acid supports peptide formation and may therefore have substituted for or

complemented lysine and arginine during early evolution[45].

In conclusion, our work demonstrates that changes in enzyme function can arise not only

through diversification and increasing structural complexity, but also through constraint and

simplification of amino acid composition. By experimentally reconstructing conditions that

approximate early evolution, we show that latent or lost catalytic potentials can be reactivated

under restricted amino acid alphabets. Arc1-12KR and Arc1-11K may therefore exemplify

how functional innovation, or reversion to ancient catalytic modes, can emerge when proteins

are forced to operate within a limited repertoire of building blocks. The findings of this study

support the view that the first enzymatic activities exploited prebiotic constraints to bridge the

gap between nonenzymatic chemistry and genetically encoded biocatalysis, illuminating a

plausible route toward the emergence of modern energy metabolism.

Methods

Design of Simplified NDK Sequences

To design amino acid sequences for the simplified NDK variants, residues targeted for

elimination were substituted with the most frequent amino acids found at the corresponding

positions in a multiple sequence alignment of 309 extant NDKs, excluding the residues

chosen for removal (Data S3). For positions that were completely conserved, chemically

similar amino acids were used instead. The N-terminal methionine residue as well as the tag

sequences appended to the N- or C-termini were not included in these considerations.

Expression and purification of simplified proteins

The simplified amino acid sequences were reverse-translated into nucleotide sequences

optimized for *E. coli* codon usage. To enable spectroscopic detection of the expressed proteins,

especially in the absence of aromatic residues such as tryptophan and tyrosine, all constructs

were designed to include a Gly-Gly-Trp tripeptide tag at the C-terminus. In addition, we

prepared both His-tagged and untagged forms of Arc1-12KR to assess the potential effect of

the N-terminal His_6-tag on protein structure and activity (Fig. S2). The genes were

synthesized by Eurofins Genomics (Tokyo, Japan) and cloned into the NdeI–BamHI

restriction recognition sites of either pET15b (N-terminal His_6-tagged form) or pET23a(+)

(untagged form) vectors (Merck, Tokyo, Japan). The resulting expression plasmids were used

to transform *E. coli* Rosetta 2(DE3) cells (Merck), which were plated on Luria–Bertani (LB)

agar containing 150 μg/mL ampicillin and incubated at 37 °C overnight. Several colonies

were then inoculated into 200 mL of LB medium supplemented with ampicillin (150 μg/mL)

and grown overnight at 37 °C. Protein overexpression was induced using the Overnight

Express Autoinduction System 1 (Merck).

The cultured cells were harvested by centrifugation (5,000 × g, 10 min, 4 °C) and then

disrupted by sonication. Lysates were heat-treated at 60, 65, or 70 °C for 20 min to precipitate

host *E. coli* proteins. After centrifugation (20,000 × g, 30 min, 4 °C), His$_6$-tagged proteins

were purified using a HisTrap HP column followed by gel filtration using a Superdex 200

Increase 10/300 GL column (GE Healthcare Japan, Tokyo, Japan). Untagged Arc1-12KR was

purified from the supernatant by sequential chromatography using HiTrap Q, Resource Q

anion exchange columns, and finally the Superdex 200 Increase gel filtration column.

Homogeneity of the protein preparation was confirmed by performing sodium dodecyl

sulfate-polyacrylamide gel electrophoresis and subsequently staining the gel with Coomassie

Brilliant Blue. Protein concentrations were determined by measuring absorbance at 280 nm

using the method of Pace et al.[46], a refinement of the procedure originally described by Gill

and von Hippel[47].

Physicochemical analyses

Analytical gel filtration chromatography was conducted using the Superdex 200 Increase

10/300 GL column (Cytiva, Tokyo, Japan) equilibrated with 20 mM Tris-HCl buffer (pH 8.0)

containing 100 mM NaCl and 1 mM EDTA. Protein samples (0.1 mL) were applied to the

column and eluted at 25 °C using a flow rate of 0.7 mL/min. Elution profiles were monitored

by absorbance at 280 nm. Apparent molecular masses were estimated from elution volumes

by referencing a calibration curve generated with protein standards.

Far-UV CD spectra were recorded using a J-1100 spectropolarimeter (Jasco, Hachioji,

Japan) equipped with a programmable temperature control unit. Protein samples were

prepared at a concentration of 20 μM in 20 mM Tris-HCl buffer (pH 8.0) containing 100 mM

NaCl and 1 mM EDTA. Spectra were acquired from 200 to 250 nm using a 0.1-cm path

length quartz cuvette at 25 °C.

Enzyme assays

Canonical NDK activity was assessed at 60 °C by measuring the transfer of the γ-phosphate group from GTP to ADP, resulting in the production of GDP and ATP. The assay buffer consisted of 50 mM HEPES (pH 8.0), 25 mM KCl, 10 mM $(NH_4)_2SO_4$, 2.0 mM $(CH_3COO)_2Mg$, 1.0 mM DTT, 5 mM ADP, and 5 mM GTP. Disproportionation activity of ADP was similarly quantified at 60 °C by monitoring ATP production, using an assay buffer containing 50 mM HEPES (pH 8.0), 25 mM KCl, 10 mM $(NH_4)_2SO_4$, 2.0 mM $(CH_3COO)_2Mg$, 1.0 mM DTT, and 5 mM ADP. In both cases, the amount of ATP produced was quantified using the luminescent Kinase-Glo Plus assay kit (Promega Japan, Tokyo, Japan), as described previously[48]. One unit of enzyme activity was defined as the amount of protein required to generate 1 μmol of ATP per minute under the assay conditions. To examine the dependence of ATP formation on ADP concentration, the reaction was performed at 60 °C using ADP as the sole nucleotide substrate. ATP production was monitored using the same assay buffer with varying concentrations of ADP from 0.3 to 5 mM. Initial velocities were determined from the linear phase of ATP formation.

HPLC analysis

The ability of Arc1-12KR to catalyze the disproportionation of ADP into ATP and AMP was confirmed by HPLC analysis. The reaction mixture contained 50 mM HEPES (pH 8.0), 25 mM KCl, 10 mM $(NH_4)_2SO_4$, 2.0 mM $(CH_3COO)_2Mg$, 1.0 mM DTT, and 5 mM ADP. Immediately after the addition of Arc1-12KR, the reaction solution was passed through a Millipore filter unit (0.45 μm pore size) and injected into the HPLC system (e-HPLC Kotori type A2, Uniflows, Tokyo, Japan) equipped with a PEGASIL ODS SP100 AQ column (Φ6×30 mm, Senshu Kagaku, Tokyo, Japan). Bound material was subsequently eluted from the column with 200 mM potassium phosphate (pH 6.0) at a flow rate of 1 mL/min.

Nucleotides were detected by UV absorbance at 265 nm. A second sample, in which the

Arc1-12KR reaction was incubated at 60 °C for 10 min, was also analyzed in the same manner.

For reference, a chromatogram of standard solution of ATP and AMP mixture was also

obtained (Fig. S3).

To assess the reverse reaction, an assay buffer containing 50 mM HEPES (pH 8.0),

25 mM KCl, 10 mM $(NH_4)_2SO_4$, either 2 or 10 mM $(CH_3COO)_2Mg$, 1.0 mM DTT, plus 5 mM

each of AMP and ATP was used. In addition, a reaction mixture containing 5 mM GDP as a

potential substrate was analyzed under the same experimental conditions used for the ADP

disproportionation assay.

X-ray crystallography

To obtain crystals of the simplified NDK Arc1-12KR, 200 nL of protein solution

(15.4 mg/mL), supplemented with 5 mM ADP, was mixed with an equal volume of reservoir

solution consisting of 20% (w/v) PEG3350 and 100 mM CHES-NaOH buffer (pH 9.5), using

the sitting-drop vapor diffusion method. Crystallization drops were incubated at 20 °C.

Crystals were harvested and briefly soaked in cryoprotectant solution containing 20% (w/v)

PEG3350, 100 mM CHES-NaOH (pH 9.5), and 20% glycerol, followed by flash-cooling in

liquid nitrogen. X-ray diffraction data were collected at beamline BL32XU of SPring-8 using

the fully automated ZOO system[49-52]. Multiple small-wedge datasets (30° per wedge) were

acquired from single crystals and automatically processed and merged using KAMO with the

XDS suite[53, 54].

Although ADP was included in the crystallization conditions to potentially stabilize the

active site conformation, no clear electron density corresponding to ADP was observed in the

final structure. The crystal structure was solved by molecular replacement with Phaser

(v2.8.2) in the PHENIX suite[55], using a ColabFold-predicted structure (v1.3.0)[56] as the search

model. Manual model building was performed in Coot (v0.8.9.1)[57], followed by iterative

refinement with phenix.refine (v1.14_3260) and validation with MolProbity. The final model

has been deposited in the Protein Data Bank under accession code 9XEJ. Crystallographic

data collection and refinement statistics are summarized in Table S1.

Molecular docking and MD simulations

The ligand structure of ADP was built and energy-minimized using Avogadro[58], followed by

further optimization with the B3LYP/def2-SVP level of theory in ORCA 6.0[59] to obtain

accurate geometries and atomic charges. The crystal structure of Arc1-12KR was used as the

receptor model, and the ligand position was initially guided by superimposing the Mg^{2+}-bound

NDK structure (PDB ID: 2DY9). Molecular docking was performed using an integrative

modeling approach with HADDOCK 2.4[60]. This procedure included rigid-body, semi-flexible,

and final flexible refinement steps. The resulting complexes were scored using HADDOCK's

integrated scoring function, and the top-ranked model was selected for MD simulations.

MD simulations were conducted for both the apo and ADP-bound forms of Arc1-12KR

using GROMACS 2024.3[61] with the CHARMM36[62] force field. Each system was solvated in

a TIP3P water box, neutralized with Na^+ and Cl^- ions, and energy-minimized using the

steepest descent method. Equilibration was performed for 5 ns each under NVT and NPT

ensembles at 310 K and 1 bar, followed by a 1 μs production run without restraints. Bond

constraints were applied using the LINCS algorithm[63], and long-range electrostatics were

treated with the PME method[64] (1.2 nm cutoff). Simulations were repeated to ensure

reproducibility.

Hydrogen-bonding interactions and water occupancy near the ligand-binding pocket were

analyzed over the trajectory. Water molecules were considered conserved if they maintained

interactions for more than 90 % of the simulation frames[65-67]. Spatial clustering of the water

molecules was visualized using PyMOL (http://www.pymol.org) and VMD[68].

Synthesis of Arc1-12KR mutant genes

The genes for four single mutants (R85M, R102M, D115A, D118A) and two double mutants

(R85M/R102M, D115A/D118A) were artificially synthesized by Eurofins Genomics. NdeI

and BamHI restriction recognition sites were added upstream and downstream of the

synthesized genes, respectively. The synthetic genes were digested with NdeI and BamHI

(New England Biolabs, Tokyo, Japan) and ligated into the corresponding restriction

recognition sites of plasmid pET15b. Expression, purification, and subsequent enzymatic

analysis of each mutant protein, carrying an N-terminal His$_6$-tag, were performed as described

above.

References

1. Frenkel-Pinter, M., Samanta, M., Ashkenasy, G. & Leman, L. J. (2020) Prebiotic Peptides:
Molecular Hubs in the Origin of Life. *Chemical Reviews*. **120**, 4707-4765.

2. Hlouchová, K. (2024) Peptides En Route from Prebiotic to Biotic Catalysis. *Acc Chem Res*.
57, 2027-2037.

3. Ezerzer, Y., Frenkel-Pinter, M., Kolodny, R. & Ben-Tal, N. (2025) A building blocks
perspective on protein emergence and evolution. *Curr Opin Struct Biol*. **91**, 102996.

4. Kitadai, N. & Maruyama, S. (2018) Origins of building blocks of life: A review.
Geoscience Frontiers. **9**, 1117-1153.

5. Cleaves, H. J., 2nd (2010) The origin of the biologically coded amino acids. *J Theor Biol*.
263, 490-498.

6. Mat, W. K., Xue, H. & Wong, J. T. (2008) The genomics of LUCA. *Front Biosci*. **13**,
5605-5613.

7. Hendrickson, T. L., Wood, W. N. & Rathnayake, U. M. (2021) Did Amino Acid Side Chain
Reactivity Dictate the Composition and Timing of Aminoacyl-tRNA Synthetase
Evolution? *Genes (Basel)*. **12**, 409.

8. Ilardo, M., Bose, R., Meringer, M., Rasulev, B., Grefenstette, N., Stephenson, J., Freeland,
S., Gillams, R. J., Butch, C. J. & Cleaves, H. J., 2nd (2019) Adaptive Properties of the
Genetically Encoded Amino Acid Alphabet Are Inherited from Its Subsets. *Sci Rep*. **9**,
12468.

9. Wehbi, S., Wheeler, A., Morel, B., Manepalli, N., Minh, B. Q., Lauretta, D. S. & Masel, J.
(2024). Order of amino acid recruitment into the genetic code resolved by last
universal common ancestor's protein domains. *Proc Natl Acad Sci U S A*. **121**,
e2410311121.

10. Philip, G. K. & Freeland, S. J. (2011) Did evolution select a nonrandom "alphabet" of
amino acids? *Astrobiology*. **11**, 235-240.

11. Kirschning, A. (2022) On the Evolutionary History of the Twenty Encoded Amino Acids.
Chemistry. **28**, e202201419.

12. Higgs, P. G. & Pudritz, R. E. (2009) A thermodynamic basis for prebiotic amino acid
synthesis and the nature of the first genetic code. *Astrobiology*. **9**, 483-490.

13. Tretyachenko, V., Vymětal, J., Neuwirthová, T., Vondrášek, J., Fujishima, K. & Hlouchová,
K. (2022) Modern and prebiotic amino acids support distinct structural profiles in
proteins. *Open Biol*. **12**, 220040.

14. Solis, A. D. (2019) Reduced alphabet of prebiotic amino acids optimally encodes the
conformational space of diverse extant protein folds. *BMC Evol Biol*. **19**, 158.

15. Ferus, M., Pietrucci, F., Saitta, A. M., Knížek, A., Kubelík, P., Ivanek, O., Shestivska, V. &

Civiš, S. (2017) Formation of nucleobases in a Miller-Urey reducing atmosphere. *Proc Natl Acad Sci USA*. **114**, 4306-4311.

16. Weber, A. L. & Miller, S. L. (1981) Reasons for the occurrence of the twenty coded protein amino acids. *J Mol Evol*. **17**, 273-284.

17. Miller, S. L. (1953) A production of amino acids under possible primitive earth conditions. *Science*. **117**, 528-529.

18. Cronin, J. R. (1989) Origin of organic compounds in carbonaceous chondrites. *Adv Space Res*. **9**, 59-64.

19. Cronin, J. R. & Pizzarello, S. (1983) Amino acids in meteorites. *Adv Space Res*. **3**, 5-18.

20. Nakamura, E., et al. *Proceedings of the Japan Academy, Series B*. **98**, 227-282.

21. Cornell, C. E., et al. (2019) Prebiotic amino acids bind to and stabilize prebiotic fatty acid membranes. *Proc Natl Acad Sci USA*. **116**, 17239-17244.

22. Longo, L. M., Lee, J. & Blaber, M. (2013) Simplified protein design biased for prebiotic amino acids yields a foldable, halophilic protein. *Proc Natl Acad Sci U S A*. **110**, 2135-2139.

23. Angyan, A. F., Ortutay, C. & Gaspari, Z. (2014) Are proposed early genetic codes capable of encoding viable proteins? *J Mol Evol*. **78**, 263-274.

24. Deville-Bonne, D., Sellam, O., Merola, F., Lascu, I., Desmadril, M. & Veron, M. (1996) Phosphorylation of nucleoside diphosphate kinase at the active site studied by steady-state and time-resolved fluorescence. *Biochemistry*. **35**, 14643-14650.

25. Akanuma, S., Yokobori, S., Nakajima, Y., Bessho, M. & Yamagishi, A. (2015) Robustness of predictions of extremely thermally stable proteins in ancient organisms. *Evolution*. **69**, 2954-2962.

26. Akanuma, S., Yokobori, S. & Yamagishi, A. (2013) Comparative Genomics of Thermophilic Bacteria and Archaea. *Thermophilic Microbes in Environmental and Industrial Biotechnology" (Eds. Satyanarayana, T., Litterchild J., Kawarabayasi, Y). Springer Science, .* pp. 331-349.

27. Shibue, R., Sasamoto, T., Shimada, M., Zhang, B., Yamagishi, A. & Akanuma, S. (2018) Comprehensive reduction of amino acid set in a protein suggests the importance of prebiotic amino acids for stable proteins. *Sci Rep*. **8**, 1227.

28. Pinna, S., Kunz, C., Halpern, A., Harrison, S. A., Jordan, S. F., Ward, J., Werner, F. & Lane, N. (2022) A prebiotic basis for ATP as the universal energy currency. *PLoS Biol*. **20**, e3001437.

29. Misra, G., Aggarwal, A., Dube, D., Zaman, M. S., Singh, Y. & Ramachandran, R. (2009) Crystal structure of the Bacillus anthracis nucleoside diphosphate kinase and its characterization reveals an enzyme adapted to perform under stress conditions. *Proteins*. **76**, 496-506.

30. Morera, S., LeBras, G., Lascu, I., Lacombe, M. L., Veron, M. & Janin, J. (1994) Refined X-ray structure of Dictyostelium discoideum nucleoside diphosphate kinase at 1.8 A

resolution. *J Mol Biol*. **243**, 873-890.

31. Webb, P. A., Perisic, O., Mendola, C. E., Backer, J. M. & Williams, R. L. (1995) The

crystal structure of a human nucleoside diphosphate kinase, NM23-H2. *J Mol Biol*.

251, 574-587.

32. Moréra, S., Chiadmi, M., LeBras, G., Lascu, I. & Janin, J. (1995) Mechanism of

phosphate transfer by nucleoside diphosphate kinase: X-ray structures of the

phosphohistidine intermediate of the enzymes from Drosophila and Dictyostelium.

Biochemistry. **34**, 11062-11070.

33. Dzeja, P. & Terzic, A. (2009) Adenylate kinase and AMP signaling networks: metabolic

monitoring, signal communication and body energy sensing. *Int J Mol Sci*. **10**,

1729-1772.

34. Trifonov, E. N. (2000) Consensus temporal order of amino acids and evolution of the

triplet code. *Gene*. **261**, 139-151.

35. Lane, N. & Martin, W. F. (2012) The origin of membrane bioenergetics. *Cell*. **151**,

1406-1416.

36. Lane, N. (2017) Proton gradients at the origin of life. *Bioessays*. **39**.

37. Schaertl, S., Konrad, M. & Geeves, M. A. (1998) Substrate specificity of human

nucleoside-diphosphate kinase revealed by transient kinetic analysis. *J Biol Chem*. **273**,

5662-5669.

38. Piserchio, A., Isiorho, E. A., Abzalimov, R., Dalby, K. N. & Ghose, R. (2026)

Phosphorylation of a conserved aspartate in the catalytic site of eukaryotic elongation

factor 2 kinase. *Protein Sci*. **35**, e70442.

39. Shin, D. H., Roberts, A., Jancarik, J., Yokota, H., Kim, R., Wemmer, D. E. & Kim, S. H.

(2003) Crystal structure of a phosphatase with a unique substrate binding domain from

Thermotoga maritima. *Protein Sci*. **12**, 1464-1472.

40. McDonald, G. D. & Storrie-Lombardi, M. C. (2010) Biochemical constraints in a

protobiotic earth devoid of basic amino acids: the "BAA(-) world". *Astrobiology*. **10**,

989-1000.

41. Giacobelli, V. G., Fujishima, K., Lepšík, M., Tretyachenko, V., Kadavá, T., Makarov, M.,

Bednárová, L., Novák, P. & Hlouchová, K. (2022) In Vitro Evolution Reveals

Noncationic Protein-RNA Interaction Mediated by Metal Ions. *Mol Biol Evol*. **39**,

msac032.

42. Zhao, M., Ding, R., Liu, Y., Ji, Z. & Zhao, Y. (2022) Determination of the Amino Acid

Recruitment Order in Early Life by Genome-Wide Analysis of Amino Acid Usage

Bias. *Biomolecules*. **12**, 171.

43. Patel, B. H., Percivalle, C., Ritson, D. J., Duffy, C. D. & Sutherland, J. D. (2015)

Common origins of RNA, protein and lipid precursors in a cyanosulfidic

protometabolism. *Nat Chem*. **7**, 301-307.

44. Longo, L. M., Despotović, D., Weil-Ktorza, O., Walker, M. J., Jabłońska, J.,

Fridmann-Sirkis, Y., Varani, G., Metanis, N. & Tawfik, D. S. (2020) Primordial emergence of a nucleic acid-binding protein via phase separation and statistical ornithine-to-arginine conversion. *Proc Natl Acad Sci USA*. **117**, 15731-15739.

45. Thoma, B. & Powner, M. W. (2023) Selective Synthesis of Lysine Peptides and the Prebiotically Plausible Synthesis of Catalytically Active Diaminopropionic Acid Peptide Nitriles in Water. *J Am Chem Soc*. **145**, 3121-3130.

46. Pace, C. N., Vajdos, F., Fee, L., Grimsley, G. & Gray, T. (1995) How to measure and predict the molar absorption coefficient of a protein. *Protein Sci*. **4**, 2411-2423.

47. Gill, S. C. & von Hippel, P. H. (1989) Calculation of protein extinction coefficients from amino acid sequence data. *Anal Biochem*. **182**, 319-326.

48. Akanuma, S., Nakajima, Y., Yokobori, S., Kimura, M., Nemoto, N., Mase, T., Miyazono, K., Tanokura, M. & Yamagishi, A. (2013) Experimental evidence for the thermophilicity of ancestral life. *Proc Natl Acad Sci USA*. **110**, 11067-11072.

49. Hirata, K., Yamashita, K., Ueno, G., Kawano, Y., Hasegawa, K., Kumasaka, T. & Yamamoto, M. (2019) ZOO: an automatic data-collection system for high-throughput structure analysis in protein microcrystallography. *Acta Crystallogr D Struct Biol*. **75**, 138-150.

50. Ito, S., Ueno, G. & Yamamoto, M. (2019) DeepCentering: fully automated crystal centering using deep learning for macromolecular crystallography. *J Synchrotron Radiat*. **26**, 1361-1366.

51. Nakamura, Y., Baba, S., Mizuno, N., Irie, T., Ueno, G., Hirata, K., Ito, S., Hasegawa, K., Yamamoto, M. & Kumasaka, T. (2020) Computer-controlled liquid-nitrogen drizzling device for removing frost from cryopreserved crystals. *Acta Crystallogr F Struct Biol Commun*. **76**, 616-622.

52. Okazaki, N., Hasegawa, K., Ueno, G., Murakami, H., Kumasaka, T. & Yamamoto, M. (2008) Mail-in data collection at SPring-8 protein crystallography beamlines. *J Synchrotron Radiat*. **15**, 288-291.

53. Yamashita, K., Hirata, K. & Yamamoto, M. (2018) KAMO: towards automated data processing for microcrystals. *Acta Crystallogr D Struct Biol*. **74**, 441-449.

54. Kabsch, W. (2010) XDS. *Acta Crystallogr D Biol Crystallogr*. **66**, 125-132.

55. Adams, P. D., et al. (2010) PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr*. **66**, 213-221.

56. Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S. & Steinegger, M. (2022) ColabFold: making protein folding accessible to all. *Nat Methods*. **19**, 679-682.

57. Emsley, P. & Cowtan, K. (2004) Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr*. **60**, 2126-2132.

58. Snyder, H. D. & Kucukkal, T. G. (2021) Computational Chemistry Activities with Avogadro and ORCA. *J Chem Educ*. **98**, 1335-1341.

59. Neese, F., Wennmohs, F., Becker, U. & Riplinger, C. (2020) The ORCA quantum

chemistry program package. *J Chem Phys*. **152**, 224108.

60. Honorato, R. V., et al. (2024) The HADDOCK2.4 web server for integrative modeling of

biomolecular complexes. *Nat Protoc*. **19**, 3219-3241.

61. Van Der Spoel, D., Lindahl, E., Hess, B., Groenhof, G., Mark, A. E. & Berendsen, H. J.

(2005) GROMACS: fast, flexible, and free. *J Comput Chem*. **26**, 1701-1718.

62. Brooks, B. R., Bruccoleri, R. E., Olafson, B. D., States, D. J., Swaminathan, S. & Karplus,

M. (1983) CHARMM: A program for macromolecular energy, minimization, and

dynamics calculations. *J Comput Chem* **4**, 187-217.

63. Hess, B., Bekker, H., Berendsen, H. J. C. & Fraaije, J. G. E. M. (1997) LINCS: A linear

constraint solver for molecular simulations. *J Comput Chem*. **18**, 1463-1472.

64. Harvey, M. J. & De Fabritiis, G. (2009) An Implementation of the Smooth Particle Mesh

Ewald Method on GPU Hardware. *J Chem Theory Comput*. **5**, 2371-2377.

65. Dasgupta, S. & Bhaumik, P. (2025) Unraveling the Molecular Architecture of Mosquito

D1-Like Dopamine Receptors: Insights Into Ligand Binding and Structural Dynamics

for Insecticide Development. *Proteins*. **93**, 1157-1170.

66. Mukherjee, S., Dasgupta, S., Panja, S. S. & Adhikari, U. (2023) Structural insight to

human Retinoid X receptor alpha-Thyroid hormone receptor beta heterodimer by

molecular modelling and MD-simulation studies: role of conserved water molecules. *J*

Biomol Struct Dyn. **41**, 9828-9839.

67. Dasgupta, S., Mukherjee, S., Mukhopadhyay, B. P., Banerjee, A. & Mishra, D. K. (2018)

Recognition dynamics of dopamine to human Monoamine oxidase B: role of

Leu171/Gln206 and conserved water molecules in the active site cavity. *J Biomol*

Struct Dyn. **36**, 1439-1462.

68. Humphrey, W., Dalke, A. & Schulten, K. (1996) VMD: visual molecular dynamics. *J Mol*

Graph. **14**, 33-38, 27-38.

Acknowledgments: This work was supported by JSPS KAKENHI (Grant numbers

19K21903, 21H01200 and 23K20903) to SA; and by the Astrobiology Center Program of the

National Institutes of Natural Sciences (Grant Numbers AB022003, AB031007 and AB0609)

to SA.

Author Contributions: S.A. designed research; S.Y., S.D., and S.A. performed research; S.Y.,

S.D., and S.A. analyzed data and wrote the manuscript; and all authors discussed the results

and reviewed the final manuscript.

Competing Interest Statement: The authors declare no competing interest.

Data Availability: The atomic coordinate file of Arc1-12KR is deposited in PDB under the

accession code 9XEJ. Experimental data of biochemical characterization and the sequences of

all constructs used in this study can be found in Supplementary Information. Requests for

materials should be directed to the corresponding author.

Table 1. Specific activities at 60 °C, oligomeric structure and amino acid composition of Arc1

and its reduced amino acid set variants.

	Arc1	Arc1-13FKR	Arc1-13KMR	Arc1-12KR	Arc1-11K
Specific activity (canonical reaction) (U/mg)	1900 ± 200	< 0.10 [a]	< 0.10 [a]	n/a [b]	n/a [b]
Specific activity (ADP disproportionation) (U/mg)	< 0.10 [a]	< 0.10 [a]	< 0.10 [a]	5.0 ± 0.3	2.5 ± 0.0
Oligomeric structure	Hexamer	Hexamer	Hexamer	Dimer	Dimer
Ala	11	12	12	12	12
Asp	6	9	9	9	9
Glu	16	16	16	16	16
Gly	11	11	11	11	11
Ile	10	10	10	10	10
Leu	7	19	19	26	26
Pro	6	6	6	6	6
Ser	8	8	8	8	8
Thr	4	4	4	4	4
Val	14	14	14	14	14
Lys	11	11	11	11	22
Arg	10	11	11	11	0
Phe	7	7	0	0	0
Met [c]	7	0	7	0	0
His	3	0	0	0	0
Asn	3	0	0	0	0
Gln	1	0	0	0	0
Trp	1	0	0	0	0
Tyr	2	0	0	0	0
Cys	0	0	0	0	0

[a] The catalytic activities were below the limit of detection.

[b] The assay used does not distinguish ATP production via the canonical reaction from that via

ADP disproportionation.

[c] The N-terminal residue was not considered.

Table 2. Specific ATP-synthesizing activity via ADP disproportionation for Arc1-12KR and
its variants at 60 °C.

Mutants	Specific activity (U/mg)
Arc1-12KR	5.0 ± 0.3
D115A	3.0 ± 0.2
D118A	3.4 ± 0.4
D115A/D118A	0.30 ± 0.01
R85M	0.67 ± 0.04
R102M	2.4 ± 0.1
R85M/R102M	0.21 ± 0.01

Fig. 1. Reaction schemes and catalytic activities of canonical NDKs and reduced amino acid

set variants. **(A)** Canonical NDK-catalyzing reaction for the transfer of γ-phosphate of a

nucleoside triphosphate (e.g., GTP) to a nucleoside diphosphate (e.g., ADP). **(B)** ADP

disproportionation reaction catalyzed by Arc1-12KR and its variants, producing ATP and

AMP from two molecules of ADP. **(C)** Time-course of ATP formation catalyzed by Arc1,

Arc1-16, and Arc1-12KR using ADP and GTP as substrates, corresponding to the canonical

NDK-catalyzing reaction shown in panel (A). The inset shows an expanded view highlighting

the activities of Arc1-16 and Arc1-12KR. **(D)** Time-course of ATP production from ADP as

the sole substrate by Arc1 and its reduced amino acid set variants. **(E)** HPLC analysis of

reaction products obtained after incubating ADP with Arc1-12KR at 60 °C for 10 min,

demonstrating the formation of ATP and AMP as predicted by the disproportionation reaction

shown in panel (B). **(F)** Initial reaction velocity (*v*) of ATP formation as a function of ADP

concentration for the ADP disproportionation reaction catalyzed by Arc1-12KR (0.3–5.0 mM

ADP). **(G)** Expanded view of the low ADP concentration range (0.3–1.0 mM) shown in panel

(F). Power-law fitting yielded an exponent of approximately 1.96. This result is consistent

with an apparent second-order dependence on ADP concentration.

Fig. 2. Crystal structure of Arc1-12KR and its MD-generated ADP–Mg²⁺ complex model.

(A) Superposition of the Arc1-12KR dimer (yellow) with the crystal structure of Arc1 (black

and gray, PDB code; 3VVT). **(B)** Model of the Arc1-12KR–ADP–Mg²⁺ complex generated by

MD simulation. **(C)** Direct and water-mediated hydrogen bonds involving W1, W2, and W3

stabilize ADP binding in the catalytic pocket of the Arc1-12KR–ADP–Mg^{2+} complex model.

(D) Comparison of Mg^{2+} coordination in the apo and holo forms. In the apo form (cyan), the

Mg^{2+} ion is coordinated by Asp115, the main-chain carbonyl oxygen of Ala116, Asp118, and

three conserved water molecules (W4, W5, and W6). In the holo form (yellow), these water

molecules are displaced by ADP, and the Mg^{2+} ion is coordinated by Asp115, Asp118, and the

α- and β-phosphate groups of ADP.

Fig. 3. Comparison of the ADP-binding pockets of the Arc1-12KR-ADP-Mg^{2+} complex

model (left) and the crystal structure of *P. horikoshii* NDK bound to ADP and Mg^{2+} (right).

The side chains of the four key residues in each structure are also shown.

Fig. 4. Conceptual model illustrating a possible evolutionary trajectory of ATP-generating

enzymes. Under prebiotic conditions with a limited amino acid repertoire, kinase-like proteins

resembling Arc1-12KR or Arc1-11K may have generated ATP through ADP

disproportionation ($2ADP \rightarrow ATP + AMP$). As the amino acid alphabet expanded toward the

LUCA era, this chemistry was likely superseded by the emergence of modern NDKs, which

catalyze phosphoryl transfer between NTPs and NDPs, and by adenylate kinases, which

catalyze the reversible reaction $2ADP \rightleftharpoons ATP + AMP$. Ultimately, the evolution of rotary ATP

synthases enabled efficient ATP production from ADP and inorganic phosphate.

805