1

4 **Hippocampal trace coding dominates and disrupts place coding**

7 Hannah S Wirtshafter[1*], Mayank R Mehta[2], Sara A Solla[1], John F Disterhoft[1]

10 *Lead contact: hsw@northwestern.edu (HSW)

12   1. Department of Neuroscience, Northwestern University Feinberg School of Medicine,
13   Chicago, IL, USA.
14
15   2. W. M. Keck Center for Neurophysics, Departments of: Physics & Astronomy,
16   Neurobiology, Neurology, University of California, Los Angeles, Los Angeles, CA 90095,
17   USA.
18
19

Abstract

The hippocampus is widely viewed as a spatial mapping system because many CA1 neurons show

location-specific activity during exploration. However, the hippocampus is also required for non-spatial

learning, including trace eye-blink conditioning. Because most prior reports of non-spatial signals were

obtained in immobile animals, it has been proposed that the hippocampus encodes space during

locomotion and non-spatial variables during immobility. To test this directly, we used calcium imaging to

record thousands of CA1 neurons while rats performed trace eye-blink conditioning during free

exploration of an open field. Across more than 6,000 neurons from five rats, mean firing rates during

trace-conditioning periods were ~1.5-fold higher than during non-trial periods, and this difference

persisted after controlling for locomotor speed. At the single-cell level, task-related modulation was

widespread and strongly biased toward increased firing. Task-enhanced neurons outnumbered spatially

selective neurons by more than threefold, indicating that trace coding predominated over place coding.

Although trace-conditioning events occurred at random spatial locations and during continuous

locomotion, trace-related activity remained robust at both single-cell and population levels. In contrast,

spatial coding was reduced during trace periods, with lower spatial information and decreased similarity

between task and non-task rate maps. These findings show that during active behavior, trace coding

dominates and disrupts place coding, challenging the view that the hippocampus functions primarily as a

stable spatial map.

Introduction

The hippocampus is widely regarded as a spatial mapping system because many CA1 neurons exhibit

location-specific activity during exploration[1-8]. However, the hippocampus is also required for a range of

non-spatial cognitive tasks[9-13] and exhibits task-related activity during these behaviors[14-28], including

trace eyeblink conditioning (the 'trace task')[15,16,18,29-34]. Because most reports of non-spatial hippocampal

signals were obtained in immobile or head-fixed animals[14,15,35,36], it has been proposed that spatial

variables dominate during locomotion, whereas non-spatial variables are represented primarily during

immobility[6,7,37,38]. Many studies of hippocampal involvement in non-spatial tasks similarly constrain

movement or tightly couple task structure to spatial position[30,39], limiting the ability to dissociate spatial

and non-spatial variables. Few studies have directly tested this idea, yielding mixed and sometimes

conflicting results[12,37]. Whether non-spatial signals reflect genuine task encoding during active

exploration rather than indirect modulation of spatial firing remains unresolved.

To address this question, we recorded large populations of CA1 neurons using calcium imaging while

rats performed trace eyeblink conditioning during free exploration of an open field. This experimental

design allowed us to dissociate spatial position, movement, and task structure while monitoring

hippocampal activity at both the single-cell and population levels.

Results

We trained a cohort of five freely moving rats on the trace task while recording neural activity from the

dorsal CA1 region using miniature microscopes and the calcium indicator GCaMP8m. In this task, each

trial consists of a 250ms auditory tone (conditioned stimulus, CS), followed by a 500 ms trace interval

with no stimuli, and then a 100ms eyelid shock (unconditioned stimulus, US) (Fig. 1a-b). Inter-trial

intervals were randomly varied between 30 and 60 seconds (mean 45s). Blink responses and shock

delivery were monitored using EMG wires implanted in the eyelid muscle (see Methods). Throughout

training and testing, animals were allowed to move freely within the behavioral arena (78.7cmx50.8cm

box), and they did not exhibit freezing or pausing during trials (Fig. 1c). Data were analyzed only in those

sessions where the rats performed the trace task on at least 70% of trials across three consecutive

sessions (See Methods, Fig. S1).

We recorded and analyzed a total of 6192 dorsal CA1 cells from five rats across 15 sessions, three

sessions per rat (See Methods). Many neurons showed clear spatial modulation, consistent with

extensive literature on place cells[4,40,41](Fig. 1d,e, upper-left panels, S2). Simultaneously, recorded

neurons also showed distinct responses during different phases of the task, with their activity aligned to

either the tone, trace interval, eyelid shock, or post-shock period, and these responses were quite stable

across trials (Fig. 1d-e, S2). The rats were at diverse locations throughout the maze during these trials,

yet the neurons were active selectively and reliably during specific epochs of the trace task, showing

that the trace-responses were not merely spatial responses. In fact, neural activity during the trace task

was so reliable that the neuron's spatial selectivity was substantially degraded and differed compared to

the non-task periods. We did several analyses and controls to test these observations, as follows.

Fig 1

Figure 1. Schematic of task and cell responses

a. Schematic of the trace eyeblink conditioning (trace task) environment. After one day of exploration in the conditioning environment, rats were trained daily in the trace task until reaching behavioral criterion

b. Schematic of the trace eyeblink conditioning task. Trial timeline showing a 250 ms tone (auditory conditioned stimulus, CS), a 500 ms trace interval, and a 100 ms eyelid shock (unconditioned stimulus, US). EMG recordings from the orbicularis oculi muscle illustrate untrained (top) and trained (bottom) animals (note that the shock period saturates the EMG trace and is truncated here for visualization). Conditioned responses (CRs) are evident in trained animals prior to US onset.

c. Representative movement trajectories during conditioning sessions. Top: Gray traces depict the trajectory across the session during times where the animal is not performing the trace task. Bottom: Blue trajectories depict movement during the tone (CS), green during the trace, red during the shock (US), and black in the post shock period.

d. Example neuron showing rate maps during non-trace task (top left) and task (bottom left) periods. Task period is also represented as raster plots (top right) and peri-stimulus histograms (PSTH, bottom left) aligned to tone (CS) onset. Boxed graphs indicate figures corresponding to the trace task.

e. Same as d but for a different cell.

Hippocampal neurons encode distinct aspects of the trace task

To assess how hippocampal activity during the trace task differs from ongoing activity during free behavior, we quantified calcium event rates separately during trace task and non-trace task periods. At the population level, calcium event rates during the trace task were positively correlated with non-task rates (Fig. 2a), indicating that baseline firing tendencies were preserved across behavioral contexts. However, activity during trace task periods was systematically elevated, with most cells falling above the unity line. This increase was not driven by a small subset of highly active neurons but reflected a broad shift across the population (Fig. 2a; Fig. S3–4).

We next asked how many neurons exhibited significant task-related increases in activity. . Comparing calcium event rates during the entire trace task window to intertrial periods revealed that 34.2 ± 8.2% of CA1 neurons exhibited significant task-enhanced activity relative to a shuffled null distribution (one-sided permutation test, $\alpha = 0.05$; Fig. 2b–c, Fig. S5–6). A smaller fraction (4.2% ± 2.8%) showed significant task suppression. This directional bias was evident at the population level, consistent with net recruitment of hippocampal cells during conditioning.

We then examined the temporal profile of task-modulated neurons. Among cells that were significantly modulated across the full trial window, responses were not uniformly distributed across the trial. Instead, most neurons showed peak modulation during a specific epoch (CS, trace, US, or post-shock), indicating temporally specific encoding rather than diffuse task engagement (Fig. 2c, Fig. S7). This demonstrates that neurons encoded specific epochs of the trace task rather than non-specific mechanisms such as altered arousal state.

Indeed, trace task-modulated neurons exhibited pronounced temporal heterogeneity. Distinct subsets of cells were preferentially active during the tone, trace, shock, or post-shock epochs, with relatively limited overlap across epochs (Fig. 2c–f, Fig. S7–S10). 85.7% of significantly modulated neurons were selective for one or two epochs, rather than being broadly active throughout the trial, indicating that task is neuronally structured in time. Although the post-shock period contained the largest absolute fraction of modulated cells, this epoch was also substantially longer than the others. After normalizing modulation by epoch duration, the density of post-epoch modulation was comparable to that observed during the tone and trace intervals and was exceeded by modulation during the brief shock epoch (Fig. 2c). Thus, the apparent dominance of post-epoch responses reflects window length rather than preferential recruitment.

Overlap analyses further showed that trace task-modulated activity was not restricted to a distinct cell

class (Fig. 2d). Task-selective and place-selective populations partially overlapped, consistent with the

idea that the trace task recruits neurons that may also participate in spatial coding. However, because

spatial tuning is examined in detail later (Fig. 4), we do not interpret this overlap here beyond noting

that task modulation is not confined to non-place cells.

Single-cell examples illustrate the heterogeneity of trace task-related responses (Fig. 2f). Some neurons

showed increased activity tightly aligned to the tone or shock, while others ramped during the trace

interval, and others exhibited delayed or prolonged post-US responses. Importantly, these trace task-

aligned calcium events occurred across a wide range of running speeds, suggesting that firing changes

were not trivially explained by locomotion.

Rats ran significantly faster during the trace task interval than during the pre-trial baseline (Fig. 2e–f; Fig.

S10–11), raising the possibility that increased activity was due to increased speed. However, speed–

firing correlations at both the trial level and within trials were near zero and indistinguishable from

shuffled controls (Fig. S12–13). Critically, when neural activity during trials was compared directly to

speed-matched intertrial periods, population-level calcium event rates remained significantly elevated

during the task (Fig. 2g). This effect was consistent across all animals and significant relative to a shuffled

null distribution. The same conclusion held at the single-cell level and was actually more pronounced:

most neurons fired 1.68x more during trials than during speed-matched non-task periods (Fig. 2h), an

effect that also persisted after controlling for acceleration (Fig. S14).

Together, these findings reveal that hippocampal neurons exhibit robust, heterogeneous task-related

increases in activity that occur independent of movement variables. The trace task engages a broad

subset of CA1 cells whose activity increases across multiple trace task epochs, with distinct temporal

profiles spanning the tone, trace, shock, and post-shock periods. These responses were not attributable

to locomotor speed and exceeded activity observed during speed-matched intertrial periods.

Thus, the trace task recruits a hippocampal ensemble in which task firing is not explained by motor

variables.

Fig 2.

a. Positive correlation between non-trial and trial calcium event rate pooled across all animals. Top figure shows raw calcium event rate in Hz. The dashed diagonal is unity (y=x), and the overlaid text gives Pearson's r and p-value. Note two outlier points are not on top figure for visualization purposes but are included in statistical analysis. See also figure S3 and S4.

b. Distribution of event rate change (trial / nontrial) across all recorded cells (blue) compared to a shuffled null distribution (gray). The dotted red line indicates no change (fold-change = 1). Modulation was biased toward increases, with 34.2 ± 8.2% of cells exhibiting significant task-enhanced activity (one-sided permutation test for increases, α

178		= 0.05). Plot ranges are set to the 99.5th percentile of pooled observed and shuffled
179		values for visualization. See Fig. S5 for individual animals and Fig. S6 for the symmetric
180		difference metric.
181	**c.**	Among task-modulated cells (full-window significant), fraction with peak in each epoch;
182		stacks show how many also have substantial modulation in +1/+2/+3 other epochs (by
183		effect size). Top: For each task epoch (Tone/CS, Trace, Shock/US, Post) and for place,
184		bars show the percent of cells significantly modulated (data normalized so that each rat
185		contributes equally). Stacks partition each epoch's modulated population into cells
186		modulated only in that epoch (blue) or additionally in +1 (green), +2 (gray), or +3
187		(purple) other epochs. This visualizes how strongly modulation is restricted to a specific
188		epoch versus shared across multiple task periods. Right: The same counts normalized by
189		epoch duration (cells per second of the epoch window). This compensates for large
190		differences in window length (e.g., the short 100 ms shock epoch vs. the longer 1.15s
191		post epoch) and reflects the density of modulation per unit time. Taller bars therefore
192		indicate higher modulation rates independent of how long the epoch lasts. Also see figure
193		S7 for same figure for place cells only; there was no significant difference between
194		proportions for all cells and place cells (paired ttest, p>0.05, ks test p>0.05). See figure
195		S8 for further breakdown by epoch using different methods.
196	**d.**	Venn diagrams showing overlap of trace task selectivity and spatial selectivity (data is
197		normalized so that each rat contributes equally). Left: Venn diagram shows percentage of
198		cells modulated by trace task versus place. Right: Cell selectivity when
199		'tone+trace+shock' are treated as one epoch and 'post' as a separate epoch.
200	**e.**	Top: Trial-aligned averages of running speed (blue, left y axis) and EMG activity (purple,
201		right y axis) for across all sessions and all rats. EMG activity is purple and purple shaded
202		areas denote ±1 standard deviation. Rats maintained locomotion throughout the trace
203		period, with EMG responses reliably occurring before US delivery. See also figures S10
204		and S11. Middle, large peri-stimulus time histogram (PSTH): PSTH averaged across all
205		rat, showing the average firing dynamics of CA1 neurons during trace conditioning. Each
206		panel illustrates the mean firing rate (solid bars) in successive 133ms bins from 1s before
207		to 2s after CS (tone) onset; error bars denote the SEM across cells. See figure S10 for
208		individual animals and figure S11 for comparison of task vs pre task running speed.
209	**f.**	Top row: Trial-by-trial raster plots (black dots) overlaid on trial-aligned speed heatmaps
210		for three example neurons. Red dashed line indicates tone onset. Middle row: Average
211		speeds across all trials for the represented day. Bottom row: PSTHs for the three pictured
212		cells. Calcium events occurred across a wide range of speeds and cells were turned to
213		different aspects of the task.
214	**g.**	Comparison of population level mean calcium event rates during trace task periods versus
215		speed-matched non trace task periods. Left: Fold change in firing rate (trace task / non-
216		trace task, speed-matched) for each rat, shown as mean ± SEM for the population. The
217		red line marks the across-animal mean. At the population level, there was on average
218		1.16x more calcium events during trials than during speed-matched controls, with
219		individual rat means±std of 1.10 ± 0.04 1.14 ± 0.05 1.13 ± 0.05, 1.27 ± 0.06, and
220		1.15 ± 0.05. Middle: Per-rat mean calcium event rates during non-trial and trial periods;
221		gray lines connect matched values from the same rat, and the red line indicates the
222		across-animal mean. Across rats, calcium event rates were significantly higher during the
223		trace task than in speed matched non-trial times (paired t-test, ***p<0.001). Right:
224		Permutation test of the population mean Δ-rate (trial − non-trial). Histogram shows the
225		null distribution obtained by random sign-flips across cells, and the blue line marks the
226		observed mean (0.014hz), which lay far outside the null range (p=0). See figure S14 for
227		analogous method used on acceleration data.
228	**h.**	Comparison of calcium event rates for individual cells during trial periods versus speed-
229		matched intertrial periods. Left: Fold change in firing rate (trace task / non-trace task) for
230		each rat, shown as mean ± SEM across tested cells. The red line marks the across-animal

mean. Neurons had on average 1.68x more calcium events during trials than during
speed-matched controls, with individual means±std of 1.39 ± 0.10, 1.59± 0.05,
1.41 ± 0.04, 1.91 ± 0.05, and 1.62 ± 0.05. Middle: Per-rat mean firing rates during non-
trial and trial periods; gray lines connect matched values from the same rat, and the red
line indicates the across-animal mean. Across rats, calcium event rates were significantly
higher during the trace task than in speed matched non-trial times (paired t-test,
***p<0.001). Right top: Permutation test of the population mean Δ-rate (trial − non-trial).
Histogram shows the null distribution obtained by random sign-flips across cells, and the
blue line marks the observed mean (0.029hz), which lay far outside the null range (p=0).
Right bottom: scatter plot comparing the trace task calcium event rate with the rate during
speed-matched non-task periods. The majority of points fall above the unity line,
indicating an overall higher rate during the trace task. See figure S15 for analogous
method used on acceleration data.

246
247
248

The trace task evokes a temporally structured, trajectory-like population code

We next asked how the task organizes population activity. Comparing population vectors (PVs) across

behavioral epochs showed that activity during the trace period was highly consistent from trial to trial

and clearly distinct from non-task activity (Fig. 3a). Population similarity within the trace period was

substantially higher than similarity between trace and non-task periods, or within non-task periods. This

separation remained when non-task windows were selected to match the running speed profile of the

trace period, indicating that differences in locomotor state alone do not explain the distinct population

pattern observed during the task. Notably, speed-matched non-task activity was itself internally

consistent, suggesting that both task and non-task behavioral states can support stable ensemble

structure. However, the trace period was associated with a population configuration that was

systematically different from non-task activity, reflecting a reproducible, trial-associated ensemble

pattern.

To examine the organization within the trial, we divided the 0–2 s window into 15 temporal bins and

computed similarity across split-averaged PVs. The across-rat similarity matrix showed a strong diagonal

with banded off-diagonals, and lag–similarity curves decayed monotonically for every rat (Fig. 3b, S16-

17). Population activity was also segmented by task epoch: within-epoch similarities exceeded between-

epoch similarities across animals (Fig. 3b, S16-S17). These results indicate that the trial engages a

conserved, temporally organized trajectory through population space.

We next asked whether this temporal trajectory mirrors the distance-dependent structure of spatial

coding. Using equal-occupancy spatial bins, spatial PV similarity was highest for adjacent locations and

declined with spatial distance, consistent with place coding (Fig. 3c, S18-S19). Strikingly, temporal and

spatial lag curves exhibited nearly identical shapes across a range of bin numbers (4-14 bins; Fig. 3d),

and split-half analyses confirmed the reliability of both structures (S17, S19). This alignment shows that

population similarity falls off with "distance" along the relevant axis—time during the task; space during

exploration.

Together, these results reveal that trace conditioning engages diverse temporal coding strategies,

including early CS-driven responses, trace-interval modulations, US rate increases, and sustained post-

US changes. Despite this heterogeneity, population activity during the trial is highly stereotyped and

distinct from non-trial periods, indicating the emergence of a structured ensemble code during the task.

Together, these findings reveal that trace conditioning recruits a robust, temporally structured

ensemble code that is distinct from non-trial activity yet geometrically analogous to the canonical

distance code in space. This temporal–spatial alignment indicates a shared organizing principle for

spatial and temporal hippocampal codes. In summary, the trace task restructures CA1 activity into a

robust, temporally organized population code that is distinct from place coding.

Fig 3

287
288
289

Figure 3. Heterogeneous but distinct neuronal dynamics and coding across trials

a. TT = trace task, NTT = non–trace task, smNTT = speed-matched non–trace task. Top,
pooled bootstrap distributions across sessions. Bottom, session means ± SD. For each
session (n = 15), per-trial 2-s trace task windows were compared with (i) randomly
selected non-task windows and (ii) speed-matched non-task windows drawn strictly from
outside the task window. For each bootstrap draw, trials were split into disjoint groups to
form population vectors (PVs), and similarity was quantified as the Pearson correlation
across cells. Trace task similarity (TT vs TT; 0.750±0.093) was substantially higher than
trace task–non-task (TT vs NTT; 0.427±0.101), non-task–non-task (NTT vs NTT;
0.507±0.107), and trace task–speed-matched non-task (TT vs smNTT; 0.580±0.100)
similarity (paired tests across sessions, ***p < 0.0001 for all comparisons). In contrast,

speed-matched non-task activity was internally reliable (smNTT vs smNTT;
0.736±0.057) and did not differ significantly from trace task similarity, indicating that the
trace task organizes CA1 population activity into a consistent, trial-specific ensemble
state distinct from non-task activity.
b. Population-vector similarity is strongly time-structured and shared across rats. Left, heat
map: Across-rat mean similarity matrix (0–2 s after CS onset divided into 15 equal
"splits"), population vectors were built from split-averaged event rates centered around
their means and compared with Pearson correlation. Bright diagonal and banded off-
diagonals indicate that population activity is most similar for nearby time bins and
progressively less similar with temporal distance. Top right: within vs between epochs.
Bins were grouped into CS, trace, US, and post-US epochs. Bars show across-rat mean (±
SEM) similarity for all within-epoch pairs vs between-epoch pairs; colored lines are
individual rats. Within-epoch similarity exceeded between-epoch similarity (Paired t test,
p*** = <0.001 across rats and for each rat, see figure S16), indicating that activity is not
only temporally smooth but also clustered by task epoch. Bottom, lag curves: For each rat
(colored lines), mean similarity is plotted as a function of lag (difference in split index).
Thick black line shows the across-rat mean ± SEM. All rats show a monotonic decay of
similarity with lag, consistent with a temporally organized population code. (Fig. S16).
Split-half analyses confirmed that lag structures were reliable within session (Fig. S17)
c. Spatial population similarity is structured by physical proximity. Left, heat map: Across-
rat mean similarity matrix. Population vectors (PVs) were built from 15 equal-occupancy
spatial bins computed from non-trial running epochs (speed ≥ 4 cm/s). PVs were built
from per-cell mean event rates in each bin and centered around the mean per cell and
compared with Pearson correlation. The bright diagonal and banded off-diagonals
indicate strongest similarity for nearby bins that share borders. Top right: spatially
adjacent bins vs non-adjacent bins (adjacency defined on the 2-D grid by rook/edge
sharing). Bars show across-rat mean ± SEM; colored lines are individual rats. Adjacent
bins were more similar than non-adjacent bins (Paired t test, p*** = <0.001 across rats
and for each rat, see figure S18). Bottom, lag curves: For each rat (colored lines), mean
similarity is plotted as a function of lag (distance of bin centers in cm.). Similarity
generally decreases with physical distance, consistent with a distance-structured spatial
code. Split-half analyses confirmed lag structures were reliable within session (Fig. S19)
d. Temporal and spatial population-vector similarity show a shared distance–decay profile.
Population-vector (PV) similarity is plotted versus temporal lag during the task period
(0–2 s; temporal, dark purple) and versus spatial separation during non-trial running
(spatial, dark blue). Because spatial position is two-dimensional, spatial "lag" is
expressed as distance between arena bins rather than the number of bin steps (for
reference, one temporal lag bin corresponds to 0.13 s). Each panel shows a different
discretization of time/space (5, 8, 12, or 15 bins; top-left to bottom-right). Solid lines
show across-rat means. Faint purple and blue bands indicate shuffle/null references for
temporal and spatial curves, respectively (mean of 500 bin-order permutations of the PV
similarity matrix). In every binning scheme the temporal and spatial curves decline with
lag and closely overlap in shape (Pearson r on curves: 0.990, 0.981, 0.975, 0.967 for
5/8/12/15 bins; RMSE_z = 0.123–0.246). The integrated magnitudes of the temporal and
spatial curves did not differ from each other (ΔAUC permutation test, p = 0.866, 0.732,
0.624, 0.714 for 5/8/12/15 bins). Together, these analyses show that CA1 ensemble states
obey a similar distance–decay rule in time during the trial and in space during
exploration.
348
349
350   **The trace task disrupts hippocampal spatial coding**
351

To determine how task-related activity interacts with spatial representations in hippocampal CA1, we

examined the impact of the trace task on spatial coding during free exploration. As expected, a subset of

neurons exhibited spatially structured firing during non-task running, as quantified by spatial mutual

information (MI) computed outside trial epochs (Fig. S20). Across animals, approximately 9% of cells

exceeded a shuffled MI threshold, consistent with previous estimates of spatial tuning reported in

calcium imaging studies during open-field exploration.

Including trace task activity altered spatial structure. Excluding task-period calcium events increased

spatial mutual information across animals, whereas removing comparable activity from non-task periods

had a smaller effect (Fig. S21). This indicates that task-locked activity disproportionately degrades spatial

information, consistent with these events being temporally organized and only weakly related to the

animal's position. Furthermore, this disruption persisted beyond the nominal trace interval. Spatial

information remained reduced for several seconds following the unconditioned stimulus (Fig. S22), and

trace-period activity was characterized by elevated trial-to-trial variability and reduced reliability relative

to non-task running, despite higher mean firing rates (Fig. S23). Alternative information-theoretic

measures yielded qualitatively similar results (Figs. S24–S25).

We next asked whether task-related responses were constrained by spatial position at the level of

individual neurons. Trace-period firing rates did not differ between trials whose occupancy centroids fell

inside versus outside a cell's non-task-defined place field (Fig. S26). Likewise, neither distance to the

place-field center nor spatial separation between trials meaningfully predicted firing variability during

the task (Figs. S27–S28). These results indicate that task-locked responses are largely uncoupled from

spatial location.

Direct comparison of spatial firing patterns revealed a redistribution of activity during the trace task.

Individual neurons frequently expressed calcium events outside their non-task-defined place fields

during trials, as reflected by increased activity in task-defined spatial masks and, in some animals,

reduced activity within non-task-defined place-field zones (Fig. S29). These results indicate that the task

reshapes where neurons are active in space, rather than simply scaling pre-existing place fields.

At the population level, these effects exposed a pronounced asymmetry between task and spatial

representations. Trial–trial population-vector (PV) similarity was only weakly influenced by spatial

location: PV similarity was modestly higher for trials occurring at the same location than for those at

different locations, and this effect was small and inconsistent across animals (Fig. 4b). A parallel single-

cell split-half analysis yielded a similar result, with only weak spatial specificity of temporal task-

response profiles (Fig. 4c). Together, these analyses indicate that the task ensemble itself is not strongly

organized by space.

In contrast, spatial coding was robustly disrupted during the trace task. Spatial PV similarity for matched

locations was markedly reduced during the trace interval compared to non-task running, even when

both were referenced to the same non-task spatial template (Fig. 4e). This reduction was large,

consistent across animals and days, and highly significant relative to within-day permutation controls

(Fig. S30). A corresponding effect was evident at the single-cell level: correlations of per-cell spatial firing

patterns across spatial bins were strongly reduced during the task (Fig. 4f; Fig. S31), indicating a loss of

stable, location-specific firing within individual neurons. Additionally, averaging single-cell spatial

stability within each animal revealed a robust reduction during the trace task (Fig. 4g), indicating a loss

of stable, location-specific firing at the level of individual neurons; a corresponding reduction in spatial

stability was also observed in population-vector analyses (Fig. S32).

Together, these findings reveal a pronounced asymmetry between task and spatial representations in

CA1. Trace task activity is only weakly constrained by spatial location, whereas the trace task

substantially degrades spatial firing structure at both the single-cell and population levels. During trials,

neurons increase their firing and frequently have calcium events outside their established place fields,

leading to a degradation of spatial population structure. These results indicate that the task recruits a

mode of hippocampal processing that is largely distinct from classical spatial coding and can transiently

override it.

Fig 4

Figure 4

a. Example single-cell temporal activity profiles underlying split-half spatial similarity
analyses. Each row shows one hippocampal neuron from a representative recording day. Left
and middle: mean calcium event rate across temporal bins within the trace interval, computed
separately for trials in which the animal occupied the same spatial bin (left) versus different
spatial bins (middle). Right: mean across-trial correlation of temporal activity profiles
(Pearson r) for same-place and different-place comparisons. These examples illustrate how
trial-to-trial consistency of temporally structured activity can differ depending on spatial
context, motivating the population-level split-half similarity analyses shown in panels e–f.
b. Trial–trial population-vector (PV) similarity is compared for pairs of trials whose time-bin-
matched occupancy centroids occurred in the same vs different spatial locations. Bars show
across-day means (n=15 days from 5 rats); thin lines connect paired day means colored by
rat. PV similarity was not signigicantly higher for trials occurring in the same location
compared with trials occurring in different locations when assessed across animals (paired t
test in Fisher z space, t(4) = 2.37, p>0.05). PV similarity was modestly but significantly
higher for trials occurring in the same location compared with trials occurring in different
locations when assessed across days (paired t test in Fisher z space: t(14)=2.30, p=0.04).
c. Single-cell split-half spatial specificity during the task period. Trial-to-trial similarity of
single-cell temporal activity profiles was compared for pairs of trials occurring in the same
spatial bin versus different spatial bins. Similarity was quantified as the Pearson correlation
of per-cell temporal activity vectors across split halves, Fisher-z transformed and averaged
within day. Across animals (n = 5), same spatial bin correlations were higher than different
bin correlations but did not reach statistical significance (paired t test in Fisher z space,
t(4)=1.96, p>0.05). Across days (n = 15), same spatial bin similarity was modestly but
significantly higher than OUT–OUT (paired t test in Fisher z space, t(14)=2.44, p<0.05).
Bars show means; thin lines indicate paired day values colored by animal.

d. Single-cell example illustrating the spatial-stability metric used in population analyses. Left:
Spatial firing-rate maps from one neuron during non-task exploration (top; reference map)
and during the trace task (bottom). Each square corresponds to one spatial quadrant of the
arena; color indicates mean firing rate (Hz) on a shared scale. Right: Quadrant-averaged
firing rates (top) and spatial similarity (bottom), computed as the Pearson correlation across
quadrants between activity vectors. In the absence of the task, the neuron exhibited a
spatially structured firing pattern with high within-map similarity. During the trace task,
firing became more uniformly distributed across space, yielding reduced spatial similarity.
This single-cell example mirrors the correlation-based spatial-stability measures computed
across neurons and days in the subsequent analyses.
e. Spatial PV similarity for the same locations was compared between the trace task and non-
task running. For each day, population vectors were computed as mean activity within coarse
spatial bins. Without-task stability was defined as the correlation between PVs from two
independent halves of non-task data, while with-task similarity was computed by correlating
trace-interval PVs to a non-task spatial template from the same day. Values reflect averages
across Monte Carlo resamples computed in Fisher-z space and converted back to r for
visualization. Spatial similarity was significantly lower during the task trace interval than
during non-task periods (mean r: 0.050 task vs 0.54 non-task). This reduction was robust
across animals (paired t-test in z space: t(4)=−10.10, p<0.00) and across days (paired t-test in
z space: t(14)=−13.81, p<0.001). Within each animal, significance was confirmed using a
within-day sign-flip permutation test on paired bin-wise differences (two-sided p≤0.001 for
all animals; see Fig. S30).
f. Single-cell spatial reliability was compared between the trace task and non-task running. For
each neuron, spatial firing maps were constructed across coarse spatial bins. Without-task
reliability was defined as the correlation of each cell's spatial map across independent non-
task subsets, whereas with-task reliability was quantified by correlating each cell's task-
epoch spatial map with its non-task spatial template from the same day. Correlations were
Fisher-z transformed and averaged within day and animal before visualization. Single-cell
spatial similarity was markedly reduced during the trace task compared to non-task periods
(mean r: 0.050 task vs 0.530 non-task), indicating disruption of location-specific coding at
the level of individual neurons. This effect was significant across animals (paired t-test in z
space: t(4)=−7.96, p***=0.001) and across days (paired t-test in z space: t(14)=−12.07,
p<0.001). Significance was further supported by within-animal, day-aware sign-flip
permutation tests (all rats p≈0; see Fig. S31).
g. Single-cell spatial stability is reduced during the trace task. Spatial stability was quantified
for each neuron as the Pearson correlation of its spatial firing pattern across arena bins,
computed separately during non-task and task epochs. For each animal, single-cell
correlations were averaged across neurons and days. Points show per-animal means (bold
colors), lighter lines indicate individual days, and bars show across-animal means. Single-cell
spatial similarity was significantly lower during task compared to non-task epochs (mean r:
0.314 vs 0.557; rat-averaged Fisher z: 0.325 vs 0.628), indicating task-associated disruption
of spatial firing structure at the level of individual neurons. This effect was significant across
animals (paired t-test in Fisher z space: t(4)=4.392, p**=0.01, n=5 rats) and across pooled
day pairs (t(14)=4.805, p<0.001 n=15 days). See Fig. S32 for the corresponding population-
vector analysis.

Discussion

The trace task reorganized hippocampal CA1 activity into a dominant, temporally structured population response that was less constrained by spatial location. Across animals, task-related population structure exceeded spatial coding in magnitude, and individual neurons exhibited heterogeneous, epoch-specific responses across the tone, trace, shock, and post-shock periods. Despite this single-cell diversity, population activity cohered into a consistent trajectory that was distinct from non-task activity and internally organized by task time. At the same time, spatial coding was transiently disrupted: place-related firing became less reliable, and spatial similarity decreased at both the single-cell and population levels. Together, these findings show that during active behavior, hippocampal coding can be reorganized by task demands into a non-spatial population response that disrupts classical place coding.

Reports that hippocampal task responses appear gated by place fields have been interpreted as evidence that non-spatial signals in CA1 are secondary to spatial coding[6,7,37]. Our results instead show that such place-bound task responses arise under restricted behavioral conditions. When animals are immobile or task events occur at fixed sites or within a limited set of locations, trials are sampled from a narrow spatial range, causing task-related activity to appear place-gated because spatial variation is not tested. In contrast, when animals remain mobile and task events are sampled across the environment, the trace task recruits a temporally organized population response that reduces spatial reliability rather than reinforcing it. Notably, even on trials preceded by brief immobility, we did not observe stable place-specific firing during task periods. Thus, whether task responses appear spatially gated or spatially disruptive reflects behavioral state and spatial sampling, rather than a fixed or obligatory relationship between task responses and spatial coding (Box 1).

Additionally, despite their distinct organizing variables, task-related and spatial population structure obeyed a similar distance–decay relationship. Population similarity decreased with increasing temporal separation during the task and with increasing spatial separation during exploration, and the shapes of these decay functions closely overlapped (Fig 3d). This correspondence indicates that task and spatial coding share a common population-level geometry, even as different dimensions dominate under different behavioral demands. The trace task therefore recruits an existing population structure rather than introducing a separate representational mode.

Taken together, these results reveal three core features of hippocampal coding during trace

conditioning. First, task-related population structure exceeds spatial coding in magnitude

during active behavior. Second, the trace task reduces spatial reliability, demonstrating that

task coding is not simply layered onto stable place representations. Third, task and spatial

coding share a common distance–decay structure, indicating a shared population geometry.

Apparent disagreements in the literature may therefor arise from differences in

whichdimension (space or task time) dominates in a behavioral task. Our findings extend the

classical view of the hippocampus as a primarily spatial map toward a flexible population code

that prioritizes behaviorally relevant structure across time and space.

Prior task paradigms	This study
Fixed stimulus location or trials tend to happen at specific locations such as endpoints[37]	Random trial locations (Fig 1)
Long CS or repeated US[38]	Brief CS-trace-US structure (Fig 1)
Reward linked endpoints[37] or rewarded field[38]	No reward (Fig 1)
Non-hippocampally dependent task (e.g. no trace period)[38]	Hippocampally dependent task[29]
Freezing at trials[38]	No freezing at trials (Fig 2)
Electrical artifact during US limits reliable recording of hippocampal activity during shock[37]	Calcium imaging permits continuous measurement through CS, trace, US, and post-US epochs
Observed outcome: Task responses appear gated by place fields when spatial and task signal align	**Observed outcome:** Temporally structured task ensemble (Figs. 2–3) accompanied by reduced spatial stability (Fig. 4)
Whether hippocampal task responses appear spatially gated or spatially disruptive depends on behavioral state and spatial sampling, reflecting flexible coding rather than a fixed place-only representation[11].	

Box 1 | Behavioral and methodological factors that determine whether hippocampal task
responses appear place-bound.
Prior reports of place-gated task responses can arise from restricted spatial sampling,
immobility, reward structure, or limited access to brief task epochs. When these constraints are
removed, the task recruits a temporally structured population response that reduces spatial
stability

Methods

DATA AVAILABILITY

The data that support the findings of this study are available from the corresponding author. All code
produced for data analysis is available on https://github.com/hsw28/ca_imaging

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All procedures were performed within Northwestern Institutional Animal Care and Use
Committee and NIH guidelines. Five male Long Evans rats (275–325 g) were sourced from Charles River
Laboratories, injected with AAV9-GCaMP8m, implanted with a 2-mm GRIN lens, and trained and tested
on eyeblink conditioning in a rectangular apparatus (Fig. 1). Animals were individually housed in an
animal facility with a 12/12 h light/dark cycle.

METHOD DETAILS

GCaMP8 injection, lens implantation, EMG implantation

GCaMP8 injection and lens implantation were completed as reported in Wirtshafter and
Disterhoft, 2022 and Wirtshafter and Disterhoft, 2023, and Wirtshafter, Solla, and Disterhoft, 2024[17,42,43].
Briefly, rats were anesthetized with isoflurane (induction 4%, maintenance 1-2%) and a craniotomy was
performed at stereotaxic coordinates Bregma AP −4.00mm, ML 3.00mm. 0.06uL of GCaMP8m (obtained
from AddGene, packaged AAV9 of pGP-AAV-syn-jGCaMP8m-WPRE, lot v175525, titer 1.3E+13 GC/mL)
was injected over 12 minutes (approximate coordinates Bregma AP −4.00mm, ML 3mm, DV 2.95mm
relative to skull); then the syringe was raised 0.2mm and an additional 0.6ul of GCaMP7 was injected.
We repeated this process once more and at slightly different coordinates in the craniotomy hole,
resulting in 4 total injections.

We then aspirated tissue from the craniotomy site using a vacuum pump and 25 gauge needle.
Tissue was aspirated up to and including the horizontal striations of the corpus collosum. A 2mm GRIN
lens (obtained from Go!Foton, CLH lens, 2.00mm diameter, 0.448 pitch, working distance 0.30mm,
550nm wavelength) was then inserted into the craniotomy hole and cemented in place using dental
acrylic. Animals were given buprenorphine (0.05mg/kg) and 20mL saline, taken off anesthesia, and
allowed to recover in a clean cage placed upon a heat pad.

Six to eight weeks after surgery, animals were again anesthetized with isoflurane and checked
for GCaMP expression. If expression was seen, baseplates were attached using UV-curing epoxy and

dental acrylic. Electrode implantation to record obicularis oculi electromyographic (EMG) activity

occurred in the same surgery as baseplate attachment, as described previously[44,45]. Briefly, a connector

containing 5 wires was cemented on the front of the animal's head: 4 wires were implanted directly

above the eye in the surrounding muscle (2 for recording, 2 for electrical stimulation). An additional wire

was attached to a connector attached to a ground screw located above the cerebellum; this screw was

implanted during lens implantation surgery.

Behavioral environment and training

A single behavioral apparatus was used in these experiments. The environment was a 78.7cm x

50.8cm unscented rectangular enclosure with wire floor and walls and white lighting. A tether

containing a plug to relay the EMG activity and to deliver a shock to the rat's eye was attached to a the

eyeblink connector on the rat's head. The miniscope was plugged into the cemented baseplate. The

miniscope and EMG cords were all attached to a commutator for ease of animal movement.

The CS was a 250ms, 85dB free-field tone (5ms rise-fall time). The US was a 100ms shock

directed to the left eye. Shock amount varied per session per animal and was calibrated, if needed, at

the end of a training session for the next session's training. Shock level was deemed appropriate when a

shock was met with a firm shake of the animal's head.

The trace interval was 500ms and the intertrial interval (ITI) was randomized between 30s and

60s, with a 45s average. EMG signal output was amplified (5000×) and filtered (100 Hz to 5 kHz), then

digitized at 3 kHz and stored by computer.

A conditioned response (CR) was identified as an increase in integrated EMG activity that

exceeded the baseline mean amplitude by more than four standard deviations, sustained for a minimum

duration of 15ms. Baseline mean amplitude was calculated during the 500ms preceding CS onset.

Additionally, the response had to commence at least 50ms after the conditioned stimulus (CS) onset and

before the unconditioned stimulus (US) onset.

The animal's first exposure to each environment was a 38min exploration session, in which the

animal was able to freely move and explore the environment without any conditioning (Figure 1a).

Animals were then trained in one environment per session, with no more than one session per day, and

were considered to have learned the task after reaching criterion (70% CRs in 50 trials) on three

consecutive training sessions (termed 'criterion sessions').

Calcium imaging

Calcium imaging was completed as reported in Wirtshafter and Disterhoft, 2022 and Wirtshafter

and Disterhoft, 2023, and Wirtshafer, Solla, and Disterhoft 2024[17,42,43]. Briefly, calcium imaging was done

using UCLA V4 Miniscopes[46,47], assembled with two 3mm diameter, 6mm FL achromat lens used in the

objective module and one 4mm diameter, 10mm FL achromat lens used in the emission module.

QUANTIFICATION AND STATISTICAL ANALYSIS

Unless otherwise noted, means are reported as mean ± standard deviation. All analyses exclude cells with fewer than five calcium events across all trial periods. For multi-animal comparisons, averages were computed by first calculating the relevant metric per animal and then averaging across animals, rather than pooling across sessions or cells. The exception is population-level analyses in Fig. 3, which were computed across all cells. All statistical tests are two sided. All analysis and figure-generation code is available at https://github.com/hsw28/ca_imaging.

Position and speed analysis

Position was sampled by an overhead camera at 30Hz. Position tracking was done post-recording using DeepLabCut[48]. Position was then converted from pixels to cm. Position was smoothed using a Gaussian filter with standard deviation of 2cm. Speed was calculated by taking the hypotenuse of the coordinates one before and after the time of interest.

Video pre-processing and cell identification

Video pre-processing and cell identification were performed as reported in Wirtshafter and Disterhoft, 2022, Wirtshafter and Disterhoft, 2023, and Wirtshafter, Solla, and Disterhoft, 2024[17,42,43]. In brief, videos were recorded with Miniscope software at 15frames/second. Video processing was done using CIATAH software[49]. Videos were down sampled in space and normalized by subtracting the mean value of each frame from the frame. Each frame was then normalized using a bandpass FFT filter (70-100cycles/pixel) and motion corrected to a using TurboReg[50]. Videos were then converted to relative florescence (dF/F_0); F_0 was the mean over the entire video.

Cells were automatically identified using CIATAH[49] using CNMF-E[51]. Images were filtered with a gaussian kernel of width 2 pixels and neuron diameter was set at a pixel size of 8. The threshold for merging neurons was set at a calcium trace correlation of 0.65; neurons were merged if their distances were smaller than 4 pixels and they had highly correlated spatial shapes (correlation>0.8) and small temporal correlations (correlation <0.4).

All cells identified using CNMF-E were then scored as neurons or not by a human scorer. Scoring was also done within CIATAH software in a Matlab GUI. Scoring was done while visualizing and considering a calcium activity trace, average waveform, a montage of the candidate cell's Ca2+ events, and a maximum projection of all cells on which the candidate cell was highlighted. The relative fluorescence (ΔF/F_0) local maxima of each identified cell were considered calcium event times.

We recorded a total of 6,664 cells across 15 sessions, three sessions per rat (average of 205 ± 51, 339 ± 169, 242 ± 21, 777 ± 121, and 658 ± 213 cells per rat). All analyses included only cells with either a calcium rate in non-trial periods of at least 0.05hz or a trial period calcium event rate of at

least 0.05h. This resulted in a final dataset of 6192 cells, with an average of 179±34, 328±163, 220±2,

732±130, and 603±203 cells per animal.

Task modulation analysis (conditioning vs intertrial).

Trial-period activity was defined as the 0–2 s window following CS onset. All other time points

were considered non-trial periods unless otherwise noted. Task modulation was quantified as the ratio

of mean activity during the conditioning window to mean activity during the intertrial interval.

Significance was assessed by comparing the observed modulation value for each neuron to a null

distribution generated by permuting trial and intertrial labels 500 times.

In a complementary analysis, symmetric-difference modulation was computed as:

654
$$\frac{r_{task} - r_{basline}}{r_{task} + r_{basline}}$$

655   Where

656
$$r_{task} = mean\ event\ rate\ during\ trial\ periods$$

657
$$r_{basline} = mean\ reate\ during\ nontrial\ periods$$

Firing rate comparisons.

We first computed per-neuron calcium event rates during the trace window and during

randomly sampled non-task windows of equal duration. We compared the distribution of firing rates

across conditions using a two-sample Kolmogorov–Smirnov (KS) test and a paired t-test. Population-

level differences in mean event rate were visualized using histograms and bar plots of mean ± SEM.

Modulation Clustering by Epoch

To identify neurons with selective modulation during different phases of the trace-conditioning

trial, we computed firing-rate changes relative to baseline in four task epochs: conditioned stimulus (CS;

0–250 ms after tone onset), trace interval (250–750 ms), unconditioned stimulus (US; 750–850 ms), and

a post-US period (850–2000 ms). Baseline activity was calculated from a pre-trial window (−2–0 s

relative to CS onset) for each neuron and trial. For each neuron, we computed modulation indices for

each epoch as:

672
$$\Delta = \frac{r_{epoch} - r_{baseline}}{r_{epoch} + r_{baseline} + \epsilon}$$

673   Where:

674
$$r_{epoch} = mean\ calcium\ event\ rate\ during\ epoch\ period$$

675
$$r_{baseline} = mean\ calcium\ event\ rate\ during\ baseline\ period$$

676
$$\epsilon = 10^{-5}$$

Because CS (250 ms) and US (100 ms) are short relative to our 133 ms frame, very low counts

inflate variance and can saturate delta. To ensure well-estimated rates, we included cells with ≥5 total

events across all trials in the 0–2 s task window OR ≥5 baseline events (−2–0 s). This criterion is

direction-agnostic (does not favor excitation) and was chosen based on stability diagnostics. As a

robustness check, we repeated the analysis using PSTHs with data centered around the mean (and

obtained the same directional effects (Fig. S10).

The resulting four-dimensional modulation vectors (Δ values) from all neurons underwent k-

means clustering (k = 4, 10 replicates, MATLAB Statistics Toolbox) to identify functional response classes.

Clusters were reordered such that those with maximal CS modulation were plotted first, followed by

trace, US, and post-US groups. Heat maps display individual neuron modulation indices sorted first by

dominant epoch and then by response sign. For each cluster, mean ± SEM Δ values were plotted, and

cluster prevalence was calculated separately for each animal and as an across-animal mean. For the

epoch defining each cluster, we also quantified the proportion of neurons with positive versus negative

modulation.

Sliding-Window Latency Analysis

To determine the latency of task-related neuronal recruitment, we performed a sliding-window

analysis of peristimulus activity. For each neuron, spike times were aligned to CS onset and binned into

non-overlapping windows of 133 ms (reflecting the calcium imaging sampling rate of 7.5 Hz) from 1 s

before CS to 2 s after CS onset. For each trial, we computed baseline activity as the mean firing rate

during the pre-CS window (−5–0 s). Each post-CS bin was compared to the baseline distribution across

trials using a paired t-test (α = 0.05). False discovery rate (FDR) correction was applied within each

neuron to control for multiple comparisons. The latency of recruitment was defined as the first post-CS

bin showing significant deviation from baseline.

Histograms of recruitment latencies were generated for each rat as well as across the full

dataset. Cumulative recruitment curves quantified the fraction of significantly modulated neurons

recruited as a function of time, enabling comparison of fast CS-locked responses and slower trace-

interval responses.

Population state shift analysis

To compare task-period activity with post-trial activity relative to baseline running, we

quantified population-vector (PV) similarity to a running-only template. For each animal, we identified

the last up to three pre-conditioning days and analyzed each day separately before combining them

with trial-count weighting. Calcium event trains were converted to instantaneous firing-rate vectors

aligned to the position timestamps. Periods of running were identified by thresholding the

instantaneous speed (≥4 cm/s). For each day, a "running template" PV was obtained by averaging

activity across all running epochs that did not overlap with any CS-to-US window. For each trial, PVs

were computed in two windows: a task epoch (0–0.85 s after CS) and a post-trial epoch (0.85–2.0 s after

CS). Trials with insufficient running within either window were discarded. PV similarity to the running

template was measured using cosine similarity (default) or Pearson correlation. Pre- and post-trial

similarities were averaged within day and then combined across days using weights proportional to the

number of included trials. For sliding-state analyses, PVs were computed in 25 contiguous bins spanning

−2 to +4 s relative to CS onset, with similarity to the running template computed for each bin when

running duration exceeded a minimum threshold. This produced a time-resolved trajectory of state

similarity surrounding the CS. Group statistics were calculated across animals using paired tests on per-

animal similarity estimates.

Post-epoch peak-time analysis

To determine whether post-epoch–modulated neurons exhibited temporally anchored responses at trial

offset, we quantified the timing of each cell's maximal post-epoch activity using a trial-averaged peri-

stimulus time histogram (PSTH). For each rat, calcium event timestamps were converted into continuous

firing-rate estimates (7.5 Hz sampling). For every trial, firing rates were extracted during the post-epoch

window (0.85–2.0 s after CS onset), and times were converted to a relative coordinate spanning 0→1.15

s from post-window onset. These relative-time segments were binned at 0.133hz and averaged across

trials to obtain a PSTH for each cell. PSTHs were smoothed with a Gaussian kernel (3 bins). Peak times

were defined as the location of the PSTH maximum; when the maximum occurred away from bin edges,

the peak time was refined by a three-point quadratic fit. Peak times were computed only for cells

classified as post-modulated by the epoch-labeling procedure and were expressed in seconds from post-

window onset. Within each rat, an empirical cumulative distribution function (CDF) of peak times was

computed, and a cross-rat mean CDF was obtained by interpolating each rat's CDF onto a shared

temporal grid. This analysis returns the temporal distribution of peak firing within the post-epoch and

tests whether responses are concentrated immediately after trial termination or distributed throughout

the window.

Speed Modulation Across Trials

To test whether calcium event rates were modulated by running speed at the trial level, we

calculated the Pearson correlation between mean speed and mean calcium event rate across trials

(defined over the 0–2 s window following CS onset) for each neuron. Mean speed was computed within

the 0–2 s window following CS onset, matching the trace interval. Neurons with fewer than five total

calcium events across all trials were excluded. To assess significance, we applied a false discovery rate

(FDR) correction (q = 0.05) to each animal's set of correlation p-values. Additionally, we compared the

observed correlations to a shuffled null distribution in which trial speeds were randomly permuted 500

times, recalculating the correlation for each permutation. Neurons whose observed correlation

exceeded the 95th percentile of the shuffled distribution were considered significantly speed-

modulated.

Speed Modulation Within Trials

To assess whether calcium activity was modulated by moment-to-moment speed fluctuations

within trials, we binned the 0–2 s post-CS window into 7.5 Hz time bins and computed the Pearson

correlation between binned speed and calcium event rate within each trial. Each neuron's per-trial

correlations were then averaged across trials. Neurons with fewer than five total events were excluded.

Statistical significance was assessed using an FDR correction (q = 0.05) across the population. As a

control, we generated shuffled speed vectors by randomly permuting the speed values within each trial

and recomputed the correlations 500 times; the fraction of neurons exceeding the 95th percentile of

this null distribution was used to estimate the false positive rate.

Speed-Matched and Acceleration-Matched Comparison of Task and Non-Task Activity

We quantified whether hippocampal neurons changed their firing during the task window

relative to non-trial periods while explicitly controlling for running speed. "Trial" time was defined as the

window from CS onset to 2 s post-CS (0–2 s). "Non-trial" comprised all other times in the session. For

speed matching, non-trial periods were selected such that their instantaneous speeds closely matched

the speed distribution observed during trials, using fixed-width speed bins. Acceleration was computed

as the first derivative of speed and used analogously for acceleration-matched controls.

To match firing rates across comparable speeds, we partitioned time into speed bins with fixed

width 2 cm/s, wherein the bin edges spanned the min–max speed observed within the trial window, and

those same edges were applied to both trial and non-trial data. For each speed bin, we formed the

union of contiguous time segments belonging to that bin separately for trial and non-trial, and

computed the duration-matched firing rates as spike counts divided by the total time in those segments.

Cells were tested if they had at least one eligible paired bin, and bins with zero rate on both sides were

excluded; parameters were selected to include the maximum number of cells (n=6192). A paired test

across all cells was then used to compare each cell's mean trial versus mean non-trial rate.

As a population-level control, we implemented a shuffle test: for each cell's per-bin rate

differences (trial − non-trial), we randomly flipped the sign of each bin difference and recomputed the

mean across bins. Repeating this procedure 500 times yielded a null distribution of mean Δ-rates under

the hypothesis of no systematic difference. The observed mean Δ-rate was compared to this distribution

to obtain a permutation p-value.

The same steps were used for acceleration matching except acceleration was binned in $3cm/s^2$

bins rather than speed.

Population vector similarity

We quantified the structure of population activity during the trace window using a bootstrap

grouping analysis of population firing-rate vectors. For each session, firing-rate matrices were

constructed for all trace task trials by computing each neuron's event rate within the 0–2 s post-CS

window. To assess population similarity, we repeatedly drew two non-overlapping groups of trials (25

trials per group) and averaged firing rates across trials within each group to obtain two population

vectors. Similarity between group-averaged vectors was quantified using Pearson correlation across

neurons.

For non-task comparisons, we sampled 2 s windows drawn from periods that did not overlap

any trace windows. For analyses controlling for locomotor confounds, speed-matched non-task windows

were additionally selected by matching the within-window speed profile of each trace trial to candidate

non-task windows using a binned speed-profile distance metric.

This procedure yielded five within-session similarity measures: trace–trace, non-task–non-task,

trace–non-task, trace–speed-matched non-task, and speed-matched–speed-matched. For each metric,

bootstrap resampling was repeated 500 times per session to generate similarity distributions, and

session means were used for statistical comparisons. All similarity analyses were performed within

session, and sessions were weighted equally in pooled summaries (Fig. 3a).

Temporal structure during the trial: For temporal structure analyses (Fig. 3b, S17), the 0–2 s trial

window was divided into 15 equal bins aligned to imaging frames. For each bin, PVs were computed by

averaging event rates across included frames. Pairwise Pearson similarity across bins produced a

temporal similarity matrix per day. Lag–similarity curves were generated by averaging entries along

successive matrix diagonals. Task-epoch segmentation was assessed by grouping bins corresponding to

CS, trace, US, and post-US epochs (Fig. 3b, S16–S17).

Spatial structure during non-trial running: For spatial analyses (Fig. 3c, S18–S19), non-trial

running frames (speed ≥ 4 cm/s) were partitioned into equal-occupancy spatial bins. PVs were

computed for each spatial bin using the same normalization as temporal PVs. Pairwise Pearson

similarities across spatial bins formed a spatial similarity matrix, and spatial lag–similarity curves were

derived analogously.

Reliability of temporal and spatial structure was assessed using even–odd split-half analyses (Fig.

S17, S19). Lag curves from each split were correlated to confirm within-session consistency.

Lag–similarity analysis

For the lag–similarity analysis (Fig. 3d), temporal and spatial similarity matrices were reduced to

lag–similarity curves by averaging entries along their diagonals. To visualize a locality-free baseline, we

generated null curves by permuting the joint row/column order of each similarity matrix 500 times and

recomputing lag curves.

To compare temporal and spatial organization directly, lag–similarity curves were centered

around the mean and their correspondence quantified using Pearson correlation and RMSE across bins.

Differences in overall magnitude between temporal and spatial curves were tested using permutation

tests on the difference in area under the curve (ΔAUC). Analyses were repeated across multiple bin

numbers (4–14 bins) to verify robustness.

Place cell identification, computing spatial mutual information, and computing bits per second

Place cells were identified using mutual information computed when the animals were running

at speeds greater than or equal to 4cm/s (. MI was computed for all cells; there was no calcium event

rate criterion for included cells. To be considered significant, the computed mutual information (MI)

must be greater than 95% of MI scores computed 500 times from shuffled positions[52]. To compute the

MI for each cell, the training environments were divided into 2.5cm x 2.5cm bins. The calcium event rate

of each cell and the occupancy of the animal were found for each bin. Rate and occupancy were

smoothed with a Gaussian kernel with filter width of 3cm and Sigma of 0.5cm. Mutual information was

computed during periods of movement as follows[43,52,53]:

$$p = \frac{P_s}{P_o}$$

$$M_s = \sum P_s$$

$$M_o = \sum P_o$$

$$MI = \sum p * \log_2(\frac{p}{M_s * M_o})$$

where:

P_s = calcium event probability in each bin

P_o = occupancy probability at each bin

For all spatial MI analyses reported in the Results, MI was computed using only non-trial data,

excluding the 0–2 s window following each CS onset.

To quantify how task-period calcium events influence spatial information, we recomputed MI

including the 0–2 s trial window. Including trial periods consistently reduced MI in every animal,

indicating that trial-locked activity tended to be non-spatial.

To ensure this reduction was not driven solely by loss of event count, we performed a spike-

matched control: for each neuron we removed the same number of calcium events as occurred in the 0–

2 s window but sampled them at random from all high-velocity non-trial events. MI was then

recomputed using this randomly reduced spike train.

To measure the duration of post-trial, task-locked disruption of spatial information, we

performed a rolling post-CS/US mutual-information analysis (Fig. S22). Beginning at CS/US offset, we

stepped through successive time windows and, for each window, removed only the calcium events
within that window before recomputing spatial MI. For each window we also generated a spike-
matched control distribution by removing an equal number of events selected randomly from the entire
non-trial spike pool..
860
861 **Computing bits per spike and bits per second**
862 Bits per spike for spatial and trial periods was computed as follows[54]:

$$\text{bits per spike} = \sum_i \left(P_i \left(\frac{R_i}{R}\right) \log_2 \left(\frac{R_i}{R}\right) \right)$$

Where:

$i = spatial\ bin\ number$ or CSUS bin number

$P_i = occupancy\ probability\ for\ bin\ i$

$R_i = mean\ firing\ rate\ at\ bin\ i$

$R = overall\ mean\ firing\ rate$

Bits per second for spatial and trial periods was computed as follows[54]:

$$\text{bits per second} = \sum_i \left(P_i\ (R_i) \log_2 \left(\frac{R_i}{R}\right) \right)$$

871
872   Bits per spike and bits per second were computed for both spatial (non-trial) and trial epochs
873 (0–2 s post-CS) using the same occupancy-normalized bin structure as the MI analysis. These measures
874 identified a similar proportion of spatially and trial-modulated neurons, and trial-period bits-per-second
875 values were positively associated with spatial bits-per-second across neurons.
876
Cross-epoch information comparisons
To determine whether spatial and trial-related information were correlated beyond shared
firing-rate structure, we performed firing-rate–residualized comparisons (Fig. S4). For each neuron, we
regressed spatial MI, trial-period MI, bits-per-spike, and bits-per-second against mean firing rate and
computed residuals. Correlations were then computed between residual spatial and residual trial-period
metrics. Residual MI and residual bits-per-spike remained significantly positively associated, indicating
shared information beyond rate alone, whereas residual bits-per-second showed no consistent
relationship.
885
Rate Map Masking and Task vs. Non-Task Firing Comparison
To assess whether calcium events during task periods occurred outside of typical place fields, we
created spatial masks based on rate maps computed separately for task and non-task periods. First, task
and non-task spike times were defined based on a 2-second window following each CS onset. We then

computed 2D firing rate maps using smoothed occupancy-normalized spatial histograms (bin size = 2.5
cm, event count and occupancy were each smoothed with a 2D gaussian kernel with σ = 0.5 bins) For
each condition (task or non-task), we defined a binary spatial mask by thresholding the firing rate map
at one standard deviations above the mean rate across bins. The non-task high-rate mask represents a
putative place fields during each condition. Next, we computed the fraction of calcium events that
occurred within each mask during both task and non-task periods. This yielded four firing rate values per
neuron: task-period firing in task-defined mask, non-task-period firing in task-defined mask, task-period
firing in non-task-defined mask, and non-task-period firing in non-task-defined mask. These values were
compared using paired two-sided t-tests to assess whether activity in each spatial mask was significantly
modulated by behavioral state.
To further quantify task-related changes in spatial firing, spatial masks were generated
separately for task and non-task periods by thresholding occupancy-normalized rate maps at one
standard deviation above the mean rate (Fig. S29). For each neuron we computed event rates within
both masks during trial and non-trial periods. Comparisons of mask-restricted firing assessed whether
task-period calcium events occurred preferentially in task-defined high-activity zones or non-task (place-
field) zones.
906
Firing variability by location in place field and distance
For each animal and session, spatial place fields (PFs) were defined from non-task running
(speed ≥ 4 cm/s; CS→CS+2 s excluded), using standard rate maps and thresholding contiguous regions
with activity >1 SD above the mean. Task-related firing rates were computed on a trial-by-trial basis
during the CS→2 s post-CS window.
To test whether spatial position influenced trial-to-trial variability of task-related firing, each
trial was assigned an occupancy-mode centroid (highest-dwell spatial bin during the 0–2 s trial window).
For each neuron, trial firing rates were compared between trials whose centroids fell inside versus
outside the cell's non-trial place field (S26). Additional analyses examined the relationship between
firing rate and centroid distance from the place-field center (Fig. S27) and the relationship between
pairwise centroid proximity and pairwise firing-rate differences (Fig. S28). These analyses assessed
whether place-field membership or spatial proximity modulated trial-period firing reliability.
To summarize position per trial, we used an occupancy-mode centroid: within the CS→2 s
window we accumulated a 2-D histogram of the animal's x–y positions (bin size ≈2.5 cm) weighted by
dwell time between samples, and selected the bin with maximal weighted occupancy; the bin center
served as the trial's centroid. The histogram grid was aligned to the non-task rate-map grid so that
centroid lookup into the PF mask was consistent. (This avoids arithmetic-mean centroids drifting into
unvisited space, e.g., for perimeter trajectories.)

Each trial was then assigned both a firing rate (per cell) and a spatial centroid. For each cell, we

formed all pairwise comparisons of task firing rates across trials. Pairs were categorized as "in-field" or

"out-of-field" depending on whether both trials' centroids fell within or outside the cell's non-task-

defined PF. We computed absolute and normalized differences in firing rate across trial pairs and

compared in-field vs out-of-field variability using paired, two-sided t-tests.

In parallel, we examined the relationship between spatial proximity and variability irrespective

of PF membership: pairwise Euclidean distances between trial centroids were regressed against firing-

rate differences to obtain Pearson correlation coefficients (r) and p-values for each cell. At the

population level, we pooled across cells and animals, and also aggregated results into rat-level

summaries before pooling.

Spatial interference in task coding: population-vector (PV) and single cell analysis

For each recording day, calcium event activity was aligned to imaging frames and analyzed

within a fixed cue-aligned trace window (CS to CS + 2 s). Animal position was discretized into a coarse

two-dimensional spatial grid (2 × 2), and neural activity was converted to framewise per-cell event rates.

Prior to downstream analyses, activity was centered around the mean across time so that similarity

metrics reflected relative modulation patterns rather than global firing-rate differences. The trace

window was subdivided into 15 equal temporal bins. For each trial, temporal bin, and spatial bin, a

population vector (PV) was computed as the mean activity across cells for frames in the corresponding

temporal–spatial intersection.

To quantify spatial structure during the trace task, PV similarity was assessed using a split-half

procedure. Within each temporal bin and spatial bin, trials were randomly divided into two non-

overlapping halves. Mean PV templates were computed separately for each half. SAME-space similarity

was defined as the Pearson correlation between templates from the same spatial bin across halves, with

DIFFERENT-space was the Pearson correlation between templates from different spatial bins across

halves. Correlations were Fisher-z transformed and averaged across spatial bins and temporal bins to

yield day-level similarity measures.

Split-half resampling was repeated 50 times per day, and results were averaged across

resamples to obtain stable day-level estimates. Day-level values were then averaged across days within

each animal. Statistical significance was assessed using paired comparisons in Fisher-z space at both the

day level (n = 15 days) and the animal level (n = 5 rats). Values shown in figures are inverse Fisher-

transformed (tanh) for interpretability.

For single-cell analyses, the same temporal and spatial binning and split-half procedure were

used. For each neuron, temporal activity vectors across bins were computed separately for each spatial

bin and split half. SAME-place correlations were defined as correlations between temporal vectors from

the same spatial bin across halves, with OUT–OUT and IN–OUT controls defined analogously. Fisher-z–

transformed correlations were averaged across neurons to yield day-level and animal-level similarity

measures.

Task interference in spatial coding: population-vector (PV) and single cell analysis

For each recording day, neural activity was aligned to imaging frames and assigned to a 2x2

spatial grid. "Control" (non-task) frames were defined by excluding all frames within a cue-aligned task

window (CS:CS+2s) plus an optional post-window buffer; analyses optionally applied a running-speed

threshold. Control frames were randomly split into two halves. For each spatial bin, population vectors

(PVs) were computed as the mean per-cell activity across frames in that bin. Prior to constructing rate

maps and population vectors, activity was centered around the mean per cell to remove cell-specific

baseline rate offsets so that similarity metrics reflected relative spatial modulation patterns rather than

global firing-rate differences (although results were consistent when not centered around the mean).

One half of the control data was used to construct a spatial template PV for each bin, and

without-task spatial reliability was quantified as the Pearson correlation between the template PV and

the PV computed from the held-out control half (split-half reliability).

To quantify spatial coding during the task, PVs were computed from frames within the task

window and grouped by spatial bin. For each bin, the task PV was correlated with the corresponding

control template PV from the same day. Correlations were Fisher-z transformed and averaged across

bins to yield a day-level spatial similarity measure, which was then averaged across days within each

animal (Fig. 4e).

For single-cell analyses (Fig. 4f), the same framework was applied at the level of individual

neurons: spatial firing maps were constructed per cell across bins, and reliability was quantified as the

correlation of each cell's spatial map across conditions or split halves, followed by Fisher-z averaging

across cells.

To reduce sampling variability, the control split and template construction were repeated 50

times with random resampling, and Fisher-z values were averaged across resamples prior to statistical

testing. Task effects were assessed by comparing with-task versus without-task Fisher-z values using a

within-day sign-flip permutation test applied to bin-wise differences, yielding two-sided p-values that

preserve spatial structure while randomizing task labels. Values shown in figures reflect the inverse

Fisher transform (tanh) of mean z values for interpretability.

For spatial stability analysis (Fig. 4g), for each animal and recording day, we quantified the

stability of coarse spatial firing patterns on a fixed 2×2 spatial grid. To reduce across-cell rate offsets

unrelated to spatial patterning, rate maps were optionally normalized per day by centering each cell

around its mean using pooled finite map entries across all bins and windows, applied identically to task

and non-task maps.

We estimated spatial stability using split-half correlations to reduce the influence of single-

window noise. For each day and condition (task, non-task), windows were randomly split into two halves

and averaged within each half to form independent estimates of the spatial firing pattern; this

procedure was repeated 50 times with different random splits. We report two complementary stability

metrics, each computed as Pearson correlation followed by Fisher z-transform and averaging in z-space.

Single-cell map stability (SC) was computed per cell as the correlation between its 4-bin mean rate

vector from half A and half B, using pairwise-valid bins (minimum 2 valid bins), and then averaged across

cells and split-half repetitions. Population-vector stability (PV) was computed within each spatial bin as

the correlation across cells between the half-A and half-B population vectors for that bin, using pairwise-

valid cells (minimum 10 valid cells), and then averaged across bins and repetitions. Pearson correlation

coefficients were Fisher z-transformed prior to averaging and statistical testing. Values shown in figures

reflect the inverse Fisher transform (tanh) of mean z values for interpretability. Task vs non-task

differences were assessed with paired t-tests performed on Fisher z values either at the animal level

(mean across days within each animal) or pooled across day pairs.

1009

1010

1011

1012

1013

1014

1015

1016

1017

References

1018

1   Penfield, W. & Milner, B. Memory deficit produced by bilateral lesions in the
hippocampal zone. *A.M.A. archives of neurology and psychiatry* **79**, 475-497 (1958).
2   Scoville, W. B. & Milner, B. Loss of recent memory after bilateral hippocampal lesions.
Journal of neurology, neurosurgery, and psychiatry **20**, 11-21 (1957).
3   O'Keefe, J. & Dostrovsky, J. The hippocampus as a spatial map. Preliminary evidence
from unit activity in the freely-moving rat. *Brain Res* **34**, 171-175 (1971).
4   O'Keefe, J. Place units in the hippocampus of the freely moving rat. *Exp Neurol* **51**, 78-
109 (1976).
5   Knierim, J. J. From the GPS to HM: Place cells, grid cells, and memory. *Hippocampus* **25**,
719-725 (2015). https://doi.org:10.1002/hipo.22453
6   O'Keefe, J. & Krupic, J. Do hippocampal pyramidal cells respond to nonspatial stimuli?
Physiol Rev **101**, 1427-1456 (2021). https://doi.org:10.1152/physrev.00014.2020
7   O'Keefe, J. Do hippocampal pyramidal cells signal non-spatial as well as spatial
information? *Hippocampus* **9**, 352-364 (1999). https://doi.org:10.1002/(SICI)1098-
1063(1999)9:4<352::AID-HIPO3>3.0.CO;2-1
8   Williams, D. A., Frame, K. A. & LoLordo, V. M. Discrete signals for the unconditioned
stimulus fail to overshadow contextual or temporal conditioning. *Journal of
Experimental Psychology: Animal Behavior Processes* **18**, 41 (1992).
9   Eichenbaum, H. & Fortin, N. J. Bridging the gap between brain and behavior: cognitive
and neural mechanisms of episodic memory. *Journal of the experimental analysis of
behavior* **84**, 619-629 (2005).
10   Fortin, N. J. in *Learning and memory a comprehensive reference* (ed John H. Byrne) 385-
417 (Academic,, London, 2008).
11   Wallenstein, G. V., Hasselmo, M. E. & Eichenbaum, H. The hippocampus as an associator
of discontiguous events. *Trends in neurosciences* **21**, 317-323 (1998).
12   Mau, W. *et al.* The Same Hippocampal CA1 Population Simultaneously Codes Temporal
Information over Multiple Timescales. *Current biology : CB* **28**, 1499-1508 e1494 (2018).
https://doi.org:10.1016/j.cub.2018.03.051
13   Crystal, J. D. Evaluating evidence from animal models of episodic memory. *J Exp Psychol
Anim Learn Cogn* **47**, 337-356 (2021). https://doi.org:10.1037/xan0000294
14   Berger, T. W., Rinaldi, P. C., Weisz, D. J. & Thompson, R. F. Single-unit analysis of
different hippocampal cell types during classical conditioning of rabbit nictitating
membrane response. *J Neurophysiol* **50**, 1197-1219 (1983).
https://doi.org:10.1152/jn.1983.50.5.1197
15   McEchron, M. D. & Disterhoft, J. F. Sequence of single neuron changes in CA1
hippocampus of rabbits during acquisition of trace eyeblink conditioned responses. *J
Neurophysiol* **78**, 1030-1044 (1997). https://doi.org:10.1152/jn.1997.78.2.1030
16   Wilmot, J. H., Puhger, K. & Wiltgen, B. J. Acute Disruption of the Dorsal Hippocampus
Impairs the Encoding and Retrieval of Trace Fear Memories. *Front Behav Neurosci* **13**,
116 (2019). https://doi.org:10.3389/fnbeh.2019.00116
17   Wirtshafter, H. S., Solla, S. A. & Disterhoft, J. F. A universal hippocampal memory code
across animals and environments. *bioRxiv*, 2024.2010.2024.620127 (2024).
https://doi.org:10.1101/2024.10.24.620127
18   Woodruff-Pak, D. S. & Disterhoft, J. F. Where is the trace in trace conditioning? *Trends
Neurosci* **31**, 105-112 (2008). https://doi.org:10.1016/j.tins.2007.11.006
19   Zuniga, A. *et al.* Extinction training suppresses activity of fear memory ensembles across
the hippocampus and alters transcriptomes of fear-encoding cells.

Neuropsychopharmacology : official publication of the American College of
Neuropsychopharmacology (2024). https://doi.org:10.1038/s41386-024-01897-0

20 Gauthier, J. L. & Tank, D. W. A Dedicated Population for Reward Coding in the
Hippocampus. *Neuron* **99**, 179-193 e177 (2018).
https://doi.org:10.1016/j.neuron.2018.06.008

21 Meshulam, L., Gauthier, J. L., Brody, C. D., Tank, D. W. & Bialek, W. Collective Behavior
of Place and Non-place Neurons in the Hippocampal Network. *Neuron* **96**, 1178-1191
e1174 (2017). https://doi.org:10.1016/j.neuron.2017.10.027

22 Panoz-Brown, D. *et al.* Replay of Episodic Memories in the Rat. *Curr Biol* **28**, 1628-1634
e1627 (2018). https://doi.org:10.1016/j.cub.2018.04.006

23 Sheridan, C. L. *et al.* Replay of incidentally encoded episodic memories in the rat.
Current biology : CB **34**, 641-647 e645 (2024). https://doi.org:10.1016/j.cub.2023.12.043

24 Allen, T. A., Salz, D. M., McKenzie, S. & Fortin, N. J. Nonspatial Sequence Coding in CA1
Neurons. *J Neurosci* **36**, 1547-1563 (2016). https://doi.org:10.1523/JNEUROSCI.2874-
15.2016

25 Shahbaba, B. *et al.* Hippocampal ensembles represent sequential relationships among
an extended sequence of nonspatial events. *Nat Commun* **13**, 787 (2022).
https://doi.org:10.1038/s41467-022-28057-6

26 Ng, C. W., Elias, G. A., Asem, J. S. A., Allen, T. A. & Fortin, N. J. Nonspatial sequence
coding varies along the CA1 transverse axis. *Behav Brain Res* **354**, 39-47 (2018).
https://doi.org:10.1016/j.bbr.2017.10.015

27 Dong, S. *et al.* Decoding memory features from hippocampal spiking activities using
sparse classification models. *Annu Int Conf IEEE Eng Med Biol Soc* **2016**, 1620-1623
(2016). https://doi.org:10.1109/EMBC.2016.7591023

28 Deadwyler, S. A., Berger, T. W., Opris, I., Song, D. & Hampson, R. E. Neurons and
networks organizing and sequencing memories. *Brain Res* **1621**, 335-344 (2015).
https://doi.org:10.1016/j.brainres.2014.12.037

29 Moyer, J. R., Jr., Deyo, R. A. & Disterhoft, J. F. Hippocampectomy disrupts trace eye-blink
conditioning in rabbits. *Behav Neurosci* **104**, 243-252 (1990).

30 McEchron, M. D. & Disterhoft, J. F. Hippocampal encoding of non-spatial trace
conditioning. *Hippocampus* **9**, 385-396 (1999). https://doi.org:10.1002/(SICI)1098-
1063(1999)9:4<385::AID-HIPO5>3.0.CO;2-K

31 Weiss, C. *et al.* Trace eyeblink conditioning in the freely moving rat: optimizing the
conditioning parameters. *Behav Neurosci* **113**, 1100-1105 (1999).
https://doi.org:10.1037//0735-7044.113.5.1100

32 Thompson, R. F. The neurobiology of learning and memory. *Science* **233**, 941-947
(1986).

33 Berger, T. W., Alger, B. & Thompson, R. F. Neuronal substrate of classical conditioning in
the hippocampus. *Science* **192**, 483-485 (1976).

34 Weiss, C. & Disterhoft, J. F. The impact of hippocampal lesions on trace-eyeblink
conditioning and forebrain-cerebellar interactions. *Behav Neurosci* **129**, 512-522 (2015).
https://doi.org:10.1037/bne0000061

35 Larra, M. F., Behrje, A., Finke, J. B., Blumenthal, T. D. & Schachinger, H. Filling the gap:
Evidence for a spatial differentiation in trace eyeblink conditioning. *Neurosci Lett* **654**,
33-37 (2017). https://doi.org:10.1016/j.neulet.2017.06.011

36 Kehoe, E. J. Repeated acquisitions and extinctions in classical conditioning of the rabbit
nictitating membrane response. *Learn Mem* **13**, 366-375 (2006).
https://doi.org:10.1101/lm.169306

Shan, K. Q., Lubenov, E. V., Papadopoulou, M. & Siapas, A. G. Spatial tuning and brain state account for dorsal hippocampal CA1 activity in a non-spatial learning task. *eLife* **5** (2016). https://doi.org:10.7554/eLife.14321

Moita, M. A., Rosis, S., Zhou, Y., LeDoux, J. E. & Blair, H. T. Hippocampal place cells acquire location-specific responses to the conditioned stimulus during auditory fear conditioning. *Neuron* **37**, 485-497 (2003). https://doi.org:10.1016/s0896-6273(03)00033-3

MacDonald, C. J., Carrow, S., Place, R. & Eichenbaum, H. Distinct hippocampal time cell sequences represent odor memories in immobilized rats. *J Neurosci* **33**, 14607-14616 (2013). https://doi.org:10.1523/JNEUROSCI.1537-13.2013

Moser, E. I., Moser, M.-B. & McNaughton, B. L. Spatial representation in the hippocampal formation: a history. *Nature neuroscience* **20**, 1448-1464 (2017).

Skaggs, W. E., McNaughton, B. L., Wilson, M. A. & Barnes, C. A. Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus* **6**, 149-172 (1996). https://doi.org:10.1002/(SICI)1098-1063(1996)6:2<149::AID-HIPO6>3.0.CO;2-K

Wirtshafter, H. S. & Disterhoft, J. F. Place cells are nonrandomly clustered by field location in CA1 hippocampus. *Hippocampus* **33**, 65-84 (2023). https://doi.org:10.1002/hipo.23489

Wirtshafter, H. S. & Disterhoft, J. F. In Vivo Multi-Day Calcium Imaging of CA1 Hippocampus in Freely Moving Rats Reveals a High Preponderance of Place Cells with Consistent Place Fields. *J Neurosci* (2022). https://doi.org:doi: 10.1523/JNEUROSCI.1750-21.2022

Weiss, C. & Thompson, R. F. The effects of age on eyeblink conditioning in the freely moving Fischer-344 rat. *Neurobiol Aging* **12**, 249-254 (1991). https://doi.org:10.1016/0197-4580(91)90105-s

Skelton, R. W. Bilateral cerebellar lesions disrupt conditioned eyelid responses in unrestrained rats. *Behav Neurosci* **102**, 586-590 (1988). https://doi.org:10.1037//0735-7044.102.4.586

Aharoni, D., Khakh, B. S., Silva, A. J. & Golshani, P. All the light that we can see: a new era in miniaturized microscopy. *Nat Methods* **16**, 11-13 (2019). https://doi.org:10.1038/s41592-018-0266-x

Silva, A. J. Miniaturized two-photon microscope: seeing clearer and deeper into the brain. *Light: Science & Applications* **6**, e17104-e17104 (2017).

Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nature neuroscience* **21**, 1281-1289 (2018).

Corder, G. *et al.* An amygdalar neural ensemble that encodes the unpleasantness of pain. *Science* **363**, 276-281 (2019). https://doi.org:10.1126/science.aap8586

Thevenaz, P., Ruttimann, U. E. & Unser, M. A pyramid approach to subpixel registration based on intensity. *IEEE Trans Image Process* **7**, 27-41 (1998). https://doi.org:10.1109/83.650848

Zhou, P. *et al.* Efficient and accurate extraction of in vivo calcium signals from microendoscopic video data. *eLife* **7** (2018). https://doi.org:10.7554/eLife.28728

Kinsky, N. R., Sullivan, D. W., Mau, W., Hasselmo, M. E. & Eichenbaum, H. B. Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long Timescales. *Current biology : CB* **28**, 3578-3588 e3576 (2018). https://doi.org:10.1016/j.cub.2018.09.037

Olypher, A. V., Lansky, P., Muller, R. U. & Fenton, A. A. Quantifying location-specific information in the discharge of rat hippocampal place cells. *J Neurosci Methods* **127**, 123-135 (2003). https://doi.org:10.1016/s0165-0270(03)00123-7

1164   54   Wirtshafter, H. S. & Wilson, M. A. Differences in reward biased spatial representations in
1165   the lateral septum and hippocampus. *eLife* **9** (2020).
1166   https://doi.org:10.7554/eLife.55252
1167

1168

1169   **Acknowledgements**: This work was supported by an NIA T32 (T32-AG020506/AG/NIA), an NIA

1170   R37 (R37-AG008796/AG/NIA), an NINDS R01 (R01 NS113804/NS/NINDS), and a K99 award (K99

1171   MH135062). We would like to thank all members of the Disterhoft lab, especially Mackenzie

1172   Kneisly.

1173

1174   **Author contributions:** Investigation, H.S.W.; Formal Analysis, H.S.W.; Writing – Original Draft,

1175   H.S.W.; Writing – Review & Editing, H.S.W., M.R.M, S.A.S., J.F.D.; Funding: H.S.W, J.F.D.,

1176   Supervision, M.R.M, S.A.S., J.F.D.

1177

1179