A mycobacterial ligand coordinates multi-receptor signaling to reprogram macrophage lipid metabolism

Dhrubajyoti Nag[1], Joycelyn Radeny[1,2], Jinyan Cui[4], Omair Vehra [1,2], Yan Yu[4], Jérôme Nigou[5], Samantha L. Bell[1,2,3], Maria L Gennaro[1,6]*.

[1] Public Health Research Institute, Rutgers New Jersey Medical School, Newark, NJ, USA

[2] Center for Emerging and Re-emerging Pathogens, Rutgers New Jersey Medical School, Newark, NJ, USA

[3] Department of Microbiology, Biochemistry & Molecular Genetics, Rutgers New Jersey Medical School, Newark, NJ, USA

[4] Department of Chemistry, Department of Biomedical Engineering, Washington University in St. Louis, St. Louis, MO, USA

[5] Université de Toulouse, CNRS, Institut de Pharmacologie et de Biologie Structurale, Toulouse, France.

[6] Department of Medicine, Rutgers New Jersey Medical School, Newark, NJ, USA

* To whom correspondence may be addressed. Email: gennarma@njms.rutgers.edu

Keywords: Dectin-2; TLR2; innate immunity; tuberculosis; lipoarabinomannan; signal integration; lipid droplets; foam cells.

Abstract. How innate immune receptors integrate signals from complex microbial ligands remains poorly understood. Here, we show that a single microbial component can encode coordinated engagement of two pattern-recognition receptors to reprogram macrophage lipid metabolism and drive foam cell formation, a hallmark of necrotizing tuberculosis lesions. These lipid-laden macrophages are defined by the accumulation of cytosolic lipid droplets enriched in neutral lipids. The mycobacterial lipoglycan mannose-capped lipoarabinomannan (ManLAM) drives macrophage lipid droplet accumulation through combined activation of Toll-like receptor 2 (TLR2) and Dectin-2, with distinct structural moieties selectively mediating recognition by each receptor. Dual receptor engagement induces lipid metabolic reprogramming and enhances NF-kB-mediated inflammatory signaling, with lipid accumulation proceeding through an mTORC1-PPARγ-dependent pathway that is largely independent of NF-kB activation. ManLAM-induced lipid metabolic changes closely mirror those elicited during *Mycobacterium tuberculosis* infection, both in neutral lipid composition and in dependence on the mTORC1-PPARγ axis. These findings identify ManLAM as a major mycobacterial driver of foam cell-associated lipid metabolism and suggest that the architecture of complex microbial ligands can organize multi-receptor signaling.

Introduction

Tuberculosis (TB) remains the leading cause of death worldwide from a single infectious agent [1]. Infection with *Mycobacterium tuberculosis* drives the formation of lung granulomas, which, under sustained immune stimulation, can progress from protective structures to necrotizing lesions that underlie lung tissue destruction and transmission [2-4]. A defining feature of this transition is the emergence of lipid-droplet-laden macrophage foam cells, which exhibit impaired antimicrobial activity, amplify pro-inflammatory and pro-necrotic signaling, and directly contribute to the necrotic core [4,5]. Foam cells may also provide a lipid-rich environment supporting *M. tuberculosis* persistence and sequester lipophilic antibiotics such as bedaquiline [5,6]. Despite their central role at the interface of immunity, metabolism, and pathology, the mechanisms driving foam cell biogenesis in TB remain poorly defined.

Several mycobacterial components, most of them associated with the cell envelope, have been linked to macrophage lipid droplet accumulation and engage multiple pattern-recognition receptors. How signals from these receptors are coordinated to drive macrophage lipid remodeling remains unclear. Mannose-capped lipoarabinomannan (ManLAM) is the best characterized of these factors. ManLAM is a ~17-kDa structurally heterogeneous lipoglycan composed of a mannosyl-phosphatidylinositol anchor, a mannan core, and an arabinan domain that is capped with terminal mannose residues in pathogenic mycobacteria, including the *M. tuberculosis* complex [7-9]. In addition to its structural role in cell wall integrity [10], ManLAM is a major determinant of mycobacterial virulence, disrupting macrophage phagosome maturation and modulating adaptive immune responses, including CD1-restricted T cell activation, CD4$^+$ T cell function, and cytokine and antibody production by B cells [8,9]. Such pleiotropic effects reflect ManLAM's capacity to engage multiple pattern-recognition receptors on macrophages, including C-type lectin receptors (CLRs) such as the mannose receptor and Dectin-2, as well as Toll-like receptor (TLR) 2. Notably, ManLAM binds Dectin-2 with high avidity [11,12], whereas its interaction with TLR2 occurs with comparatively low affinity [13,14]. Seminal work by D'Avila et al. [15] demonstrated that TLR2 is required for ManLAM-induced lipid droplet accumulation, yet activation of TLR2 alone by a potent agonist fails to induce this response [16]. These observations indicate that TLR2 signaling is necessary but not sufficient to promote foam cell formation, leaving unresolved the receptor mechanism by which ManLAM drives lipid droplet biogenesis.

We identify coordinated engagement of TLR2 and Dectin-2 as the mechanistic principle underlying ManLAM-driven foam cell formation. Distinct structural moieties within ManLAM engage these pattern-recognition receptors to activate an mTORC1-PPARγ axis that drives

triglyceride synthesis and lipid droplet biogenesis in macrophages. These same structural

determinants are also required for full inflammatory activation. Together, these findings establish

a mechanistic framework for foam cell formation in TB and identify host pathways that represent

potential points of vulnerability for host-directed therapeutic intervention.

Results

***M. tuberculosis* ManLAM induces macrophage lipid droplet accumulation through TLR2**

and Dectin-2.

Previous studies showed that the vaccine strain *Mycobacterium bovis* Bacillus Calmette-

Guérin (BCG), a member of the *M. tuberculosis* complex, induces lipid droplet (LD)

accumulation in murine macrophages in a TLR2-dependent manner, yet activation of TLR2

alone by the synthetic agonist Pam3CSK4 is insufficient to induce LD formation [15,16]. Similar

results were obtained with BCG-derived ManLAM [15,16]. These observations left unresolved how

BCG bacilli – and specifically ManLAM – promote LD biogenesis in macrophages. Because

ManLAM is a relatively weak TLR2 agonist compared to bacterial lipoproteins (EC_{50} is in the

µg/ml *vs* ng/ml range) [13,17], it was important to exclude the possibility that the observed TLR2-

stimulating activity of the ManLAM preparation resulted from trace contamination with highly

active lipopeptides [18]. To address this, purified ManLAM was treated with 1% H_2O_2, which

inactivates lipoproteins by converting the N-terminal cysteine-thioether substructure required for

TLR2 activation into sulfoxide derivatives [19]. Untreated and H_2O_2-treated ManLAM were tested

with a TLR2 reporter HEK293 cell line engineered to express alkaline phosphatase activity as a

read-out of TLR2-dependent NF-κB activation (see *Methods*). H_2O_2 treatment had little effect on

the TLR2 agonist activity of *M. tuberculosis* ManLAM (**Fig. 1A**) while completely abrogating

TLR2 activation by the sentinel lipoprotein LrpG (**SI Fig. 1**). This result demonstrates that our

ManLAM preparations are highly purified and express bona fide TLR2 agonism.

Next, to investigate the mechanisms underlying ManLAM-induced LD accumulation in

macrophages, we used immortalized murine bone marrow-derived macrophages (iBMDMs),

which are well suited for genetic manipulations and closely recapitulate primary macrophage

phenotypes [20]. Gene knockdown in iBMDMs was achieved by CRISPR-interference (CRISPRi),

which employs a nuclease-inactivated Cas9 (dCas9) and promoter-targeting guide RNAs

(gRNAs) to block transcription of target genes. We generated TLR2 knockdown (KD) iBMDMs

by stably expressing dCas9 and two different gRNAs targeting the *Tlr2* promoter (see *Methods*).

Control cells expressed gRNAs against an irrelevant target gene (bacterial luciferase, *lux*A).

Efficient TLR2 knockdown was confirmed by reduced *Tlr2* transcript levels measured by RT-

qPCR (**SI Fig. 2A**) and by impaired induction of TLR2-responsive genes *Il6* and *Tnfa* following

stimulation with the TLR2 agonist Pam3CSK4 (**SI Fig. 2B,C**). ManLAM and H_2O_2-treated

ManLAM induced LD accumulation in control iBMDMs but failed to do so in TLR2 KD cells (**Fig.**

1B; representative imaging flow cytometry data in **Fig. 1C**). Moreover, in agreement with

previous reports [16], stimulation of control iBMDMs with Pam3CSK4 alone failed to induce LD

accumulation at any of the doses tested (**Fig. 1B-E**). The observed insufficiency of TLR2

signaling prompted us to investigate whether ManLAM engages other host receptors to drive

macrophage LD accumulation.

Previous work has shown that *M. tuberculosis* ManLAM is also an agonist of Dectin-2 [11,12], a

surface CLR that recognizes microbial carbohydrates and has been studied primarily in the

context of fungal infections [21]. Consistent with these reports, *M. tuberculosis* ManLAM activated

Dectin-2 signaling in a reporter HEK293 cell line engineered to express alkaline phosphatase

activity as a read-out of Dectin-2-dependent NF-κB activation (see *Methods*) (**Fig. 2A**). H_2O_2

treatment had no significant effect on the Dectin-2 agonist activity of *M. tuberculosis* ManLAM

(**Fig. 2A**). To assess whether Dectin-2 signaling contributes to ManLAM-induced LD

accumulation, we generated Dectin-2 knockdown (KD) and control cells in iBMDM using the

same strategy described for TLR2. Efficient Dectin-2 knockdown was validated by reduced

Dectin2 transcript levels measured by RT-qPCR (**SI Fig. 2D**) and by impaired induction of the

downstream genes *Il6* and *Tnfa* following stimulation with the Dectin-2 agonist furfurman (**SI**

Fig. 2E,F). When LD induction was assessed, both ManLAM and H_2O_2-treated ManLAM

induced LD accumulation in control iBMDM but failed to do so in Dectin-2 KD cells (**Fig. 2B**;

representative imaging flow cytometry data are in **Fig. 2C**). In contrast, knockdown of other

CLRs, including Dectin-1 and Mincle, had no effect on ManLAM-induced LD accumulation (**SI**

Fig. 3A). Moreover, treating parental iBMDMs with increasing doses of furfurman up to 10 μg/ml

failed to induce LD accumulation (**Fig. 2D**; representative imaging flow cytometry data are in

Fig. 2E). Thus, as observed with TLR2, Dectin-2 activation is necessary but not sufficient to

drive ManLAM-induced LD accumulation.

The data above strongly suggest that *M. tuberculosis* ManLAM engages both TLR2 and

Dectin-2 signaling to induce LD accumulation in macrophages. To determine whether co-

engagement of these two receptors is sufficient, we treated iBMDMs with the TLR2 agonist

Pam3CSK4 and the Dectin-2 agonist furfurman, alone and in combination. While neither agonist

induced LD accumulation when applied alone, they did so when used in combination (**Fig. 3A**;

representative imaging flow cytometry data are in **Fig. 3B**). These results demonstrate that co-

engagement of TLR2 and Dectin-2 is required to induce macrophage lipid droplet accumulation

and raise the possibility that ManLAM functions as a single ligand that coordinates signaling

through both receptors to drive macrophage lipid reprogramming.

We next asked whether specific ManLAM structural features are required for lipid droplet

accumulation mediated by TLR2 and Dectin-2. Prior work showed that acylation of the

phosphatidyl-myo-inositol anchor of mycobacterial lipoglycans is required for TLR2 activation

[14,22], whereas the mannose caps of ManLAM are required for recognition by Dectin-2 [11,12]. To

selectively disrupt these interactions, we generated deacylated ManLAM by NaOH treatment

(dManLAM, see *Methods*) and mannose-cap-deficient ManLAM by enzymatic removal with α-

mannosidase (αtManLAM, see *Methods*). When tested in TLR2 and Dectin-2 reporter HEK293

cell lines, dManLAM failed to activate TLR2 while retaining Dectin-2 activation ability (**Fig. 3C**).

In contrast, αtManLAM retained TLR2 activation but lost the ability to activate Dectin-2 (**Fig.**

3C). We next assessed the ability of these ManLAM derivatives to induce LD accumulation in

iBMDMs. In contrast to intact ManLAM, neither dManLAM nor αtManLAM induced LD

accumulation (**Fig. 3D**; representative imaging flow cytometry data are in **Fig. 3E**). These

findings were recapitulated in human monocyte-derived macrophages (MDM) (**SI Fig. 3B**).

Together, these results demonstrate that engagement of both TLR2 and Dectin-2 by ManLAM is

required for macrophage LD accumulation, identify the structural moieties that mediate this dual

receptor recognition, and establish its relevance in human macrophages.

Structural integrity of ManLAM is required for full NF-κB activation.

Because ManLAM is also known to activate the transcription factor NF-κB [11], we next used

NF-κB activation as a quantitative readout to compare signaling elicited by intact ManLAM with

that induced by the structural derivatives used in the lipid droplet assays. NF-κB activation

occurs downstream of multiple innate immune receptors, including TLRs and CLRs [23]. Upon

receptor engagement, a series of signaling events results in the translocation of NF-κB from the

cytoplasm to the nucleus (schematic in **Fig. 4A**), where it induces expression of

proinflammatory cytokines such as TNF-α and IL-1β [24]. We therefore quantified NF-κB

activation by measuring nuclear translocation of the RelA (p65) subunit of NF-κB and

downstream cytokine gene expression in macrophages treated with ManLAM or its structural

derivatives.

Nuclear translocation was calculated as the ratio of nuclear to cytoplasmic RelA

fluorescence intensity in RAW 264.7 macrophages stably expressing green-fluorescence-

protein-tagged RelA. Stimulation with ManLAM induced a gradual increase in nuclear RelA

accumulation over time relative to unstimulated control cells (**Fig. 4B,C**). As expected,

ManLAM-induced NF-κB activation was substantially less potent than that triggered by soluble

Pam3CSK4, a benchmark agonist for robust TLR2-mediated activation.

ManLAM-induced RelA translocation exhibited substantial cell-to-cell variability, consistent

with the intrinsic heterogeneity of immune cell activation [25]. To quantify this response, we

defined a threshold that classified 95% of unstimulated cells as nonresponsive. Using this

criterion, the macrophage population segregated into two groups: a responsive subpopulation

displaying robust nuclear RelA accumulation and a nonresponsive subpopulation remaining

below the threshold (**SI Fig. 4**). Compared with intact ManLAM, stimulation with αtManLAM and

dManLAM -- which selectively disrupt recognition by Dectin-2 and TLR2, respectively -- reduced

both the fraction of responsive cells by approximately 50-60% (**Fig. 4D,E**) and the magnitude of

RelA translocation within the responsive cell population (**Fig. 4F**), indicating attenuated NF-κB

activation at both the population and single-cell levels. Since these structural modifications

selectively disrupt Dectin-2 and TLR2 recognition, these findings indicate that full NF-κB

activation by ManLAM requires coordinated engagement of both receptors. Consistent with this

conclusion, intact ManLAM induced markedly greater expression of downstream cytokine genes

in RAW 264.7 macrophages, including *Tnfa* and *Il1b*, with induction ~25-fold and ~6-fold higher,

respectively, than that elicited by either structural derivative (**Fig. 4G**). Comparable results were

obtained in iBMDM (**SI Fig. 5**), indicating similar responses across murine macrophage

systems.

TLR2 and Dectin-2 contribute to macrophage lipid droplet accumulation during *M.*

***tuberculosis* infection**

To assess the relevance of our findings with purified ManLAM to live *M. tuberculosis*

infection, we tested whether TLR2 and Dectin-2 signaling contribute to macrophage LD

accumulation during infection. Macrophages infected with *M. tuberculosis* and subjected to

knockdown of TLR2 or Dectin-2 exhibited a ~25% reduction in LD content compared with

parental cells (**Fig. 5A,B**). This reduction indicates that both receptors contribute to maximal LD

accumulation during infection. In light of our findings with purified ManLAM, these results also

suggest that ManLAM represents a substantial, though not exclusive, driver of the LD-inducing

capacity of the pathogen.

ManLAM induces triglyceride-rich lipid droplets through an mTORC1-PPARγ pathway

independent of NF-κB.

We next investigated whether ManLAM-induced macrophage LD accumulation resembles

that observed during live *M. tuberculosis* infection. To define the chemical nature of the

ManLAM-induced neutral lipids, untreated and ManLAM-treated iBMDMs were exposed to

chemical inhibitors of diacylglycerol acyltransferase (DGAT), which converts di- into tri-

glycerides, or acyl-CoA:cholesterol acyltransferase (ACAT), which esterifies free cholesterol.

Inhibition of DGAT, but not ACAT, reduced ManLAM-induced LD accumulation in a dose-

dependent manner (**Fig. 5C,D**), indicating that ManLAM induces triglyceride-enriched LDs,

consistent with our prior findings with *M. tuberculosis* infection [26].

We next examined the signaling pathways underlying ManLAM-induced LD formation.

Previous work showed that LD accumulation during tuberculous mycobacterial infection requires

the protein kinase Mechanistic Target of Rapamycin Complex 1 (mTORC1) and the downstream

transcription factor peroxisome proliferator-activated receptor gamma (PPARγ) [16,26,27]. Both are

key regulators of cellular lipid metabolism [28-30]. Inhibition of mTORC1 with rapamycin or of

PPARγ with GW9662 markedly reduced ManLAM-induced LD accumulation in a dose-

dependent manner (**Fig. 5E**), recapitulating findings in *M. tuberculosis* infection [16,26,27]. In

contrast, although ManLAM activates NF-κB, LD accumulation was unaffected by inhibition of

RelA (p65) nuclear translocation by JSH-23 (**Fig. 5E**). Similar results were obtained with a

second NF-κB inhibitor, QNZ (**SI Fig. 6A**), indicating that Man-LAM-induced LD formation is

largely independent of NF-kB activity.

To validate this pathway separation at the transcriptional level, we examined how inhibition

of mTORC1 or NF-κB affected expression of representative target genes following ManLAM

stimulation. Rapamycin selectively inhibited ManLAM-mediated upregulation of the PPARγ

target gene *Plin2*, which encodes a lipid droplet-associated protein [31], but did not affect

induction of the NF-κB-regulated genes *Tnfa* and *Il1b* [32] (**Fig. 5F**). Conversely, the NF-κB

inhibitor JSH-23 suppressed ManLAM-mediated induction of *Tnfa* and *Il1b* but not *Plin2*. Both

rapamycin and JSH-23 reduced ManLAM-induced upregulation of *Hif1a*, a gene controlled by

both NF-κB and mTORC1 [33,34]. Comparable results were obtained with the NF-κB inhibitor QNZ

(**Fig. S6B**). Together, these findings indicate that Man-LAM-induced LD accumulation proceeds

through an mTORC1-PPARγ-dependent metabolic program that operates independently of NF-

κB signaling, mirroring the pathway architecture observed during *M. tuberculosis* infection [26,35].

Discussion

Foam cells are the cellular hallmark of necrotizing TB lesions [26], yet how mycobacterial signals integrate to reprogram macrophage lipid metabolism remains unresolved. Here, we identify ManLAM as a key structural determinant of tuberculous foam cell formation and show that it drives this process by coordinating engagement of TLR2 and Dectin-2. Distinct structural features of ManLAM -- acylation and mannose capping -- selectively mediate activation of TLR2 and Dectin-2, respectively [11,12,14,22], and are jointly required for triglyceride-rich lipid droplet accumulation**.** These findings explain earlier observations that TLR2 signaling is necessary but insufficient for lipid droplet formation. Lipid remodeling proceeds through an mTORC1-PPARγ axis that is largely independent of inflammatory signaling. Together, these results support a model in which ManLAM co-engages two macrophage surface receptors through distinct molecular moieties, reprogramming lipid metabolism and achieving full inflammatory activation via separable pathways (**Fig. 6**). We further show that, as the only *M. tuberculosis*-derived agonist of Dectin-2 [11], ManLAM contributes to full lipid droplet accumulation during infection and that the lipid droplets it induces mirror those elicited by live bacilli in both triglyceride composition and dependence on the mTORC1-PPARγ axis. Collectively, these findings link ManLAM architecture to dual receptor recognition and macrophage lipid metabolic reprogramming during infection, providing a molecular basis for coordinated receptor engagement.

Activation of multiple receptors by distinct structural features within a microbial ligand defines a mode of signal integration at the ligand level. In contrast, prevailing models of innate immune signaling emphasize receptor crosstalk, in which distinct microbial ligands engage separate receptors and signal integration occurs downstream of receptor activation – for example, cooperative signaling between CLRs and TLRs in fungal infections [36]. Embedding ligands for multiple receptors within a single microbial component may confer an intrinsic advantage by promoting spatial proximity of engaged receptors. This idea is supported by prior studies showing that productive crosstalk between Dectin-1 and TLR2 depends on spatial coordination of their ligands, often requiring scaffolded presentation at nanometer-scale proximity [37,38]. Such spatial organization promotes receptor clustering, increases effective avidity, and converts weak interactions into productive signaling [39]. By analogy, recognition of ManLAM by Dectin-2 and TLR2 may facilitate integration of parallel signaling cascades, enabling synergistic signaling even when individual receptor-ligand interactions are of low

affinity. Ligand-level coordination may therefore represent a strategy by which microbes with

complex, lipoglycan-rich cell envelopes -- including mycobacteria and fungi – integrate host

sensing pathways.

ManLAM-driven lipogenic and inflammatory programs in macrophages appear to be

functionally separable. This is consistent with established signaling biology, as TLR2 and

Dectin-2, in addition to activating NF-kB [23], can independently engage the mTORC1-PPARγ

axis through PI3K/Akt signaling downstream of MYD88 or Syk, respectively [40,41]. At the same

time, the synergistic effects of TLR2 and Dectin-2 co-engagement on both lipogenic and

inflammatory outputs suggest convergence at shared regulatory nodes. One candidate is the

PI3K/Akt axis, which lies upstream of both NF-kB and mTORC1–PPARγ signaling [42]. ManLAM

may therefore promote foam cell formation by activating parallel signaling pathways that

convergence at common control points while preserving functional independence of

downstream outputs.

The lipid-inducing activity of ManLAM closely mirrors, but does not fully recapitulate, the

response elicited by live *M. tuberculosis*. ManLAM drives triglyceride-rich lipid droplet

accumulation in macrophages through an mTORC1-PPARγ-dependent pathway that is largely

insensitive to NF-κB inhibition, a signaling profile also observed during infection [26]. Lipid droplet

accumulation during infection is reduced in Dectin-2- or TLR2-deficient macrophages,

suggesting that ManLAM contributes to this response in vivo, particularly given that it is the only

known *M. tuberculosis*-derived agonist of Dectin-2 [11]. However, whereas TLR2 or Dectin-2

deficiency fully abrogates ManLAM-induced lipid droplet formation, it only partially reduces lipid

accumulation during infection, indicating that ManLAM contributes substantially -- but not

exclusively -- to foam cell formation. Additional microbial and host-derived inputs, including

trehalose dimycolate and mycolic acids [43,44], cGAS–STING activation [45], and TNF signaling [26],

likely cooperate in this process. Together, these observations support a hierarchical model in

which ManLAM-dependent surface receptor signaling acts as an upstream driver of lipid

metabolic reprogramming, while additional pathways shape the magnitude and persistence of

foam cell formation during infection.

By identifying mTORC1 and PPARγ as downstream effectors of dual receptor engagement, we

define key nodal control points within the lipid metabolic program that may be targeted

pharmacologically to disrupt pathogenic foam cell formation. Because these regulators lie

downstream of receptor engagement, their modulation may selectively constrain lipid

remodeling while preserving upstream innate immune recognition. Thus, elucidating how ligand

architecture encodes receptor integration not only advances our understanding of host–

pathogen signaling logic but also highlights mechanistically grounded opportunities for host-

directed therapy.

Materials and Methods

Mycobacterial strains and purified components

Mycobacterium tuberculosis Erdman (TMC#107) was cultured in complete 7H9 medium,

which consists of Middlebrook 7H9 base (BD Difco, Franklin Lakes, NJ) supplemented with 10%

(v/v) oleic acid-albumin-dextrose-catalase (OADC) obtained from oleic acid (0.5 g/L, Thermo

Fisher Scientific, Waltham, MA), albumin (50 g/L), dextrose (20 g/L), and catalase (0.04 g/L all

from Fisher Scientific, Hampton, NH). The medium also contained sodium chloride (8 g/L),

glycerol (0.5% v/v), and was supplemented with 0.05% tyloxapol (all from Thermo Fisher

Scientific, Hampton, NH).

Mannosylated lipoarabinomannan (ManLAM) was extracted from *M. tuberculosis* Erdman

and purified by a previously described method [46,47]. Deacylated ManLAM (dManLAM) was

prepared as described previously [46]. Briefly, 200 µg of ManLAM was incubated in 200 µl of

NaOH 0.1 M for 2 h at 37°C. After neutralization with 200 µl of HCl 0.1 M, the reaction products

were dialyzed against water. ManLAM devoid of mannose caps (αtManLAM) was prepared as

described previously [48]. Briefly, 200 µg of ManLAM was incubated for 6 h at 37°C in 30 µl of α-

mannosidase solution (2 mg/ml, 0.1 M sodium acetate buffer, pH 4.5, 1 mM $ZnSO_4$) followed by

a second addition of 50 µl of the same enzyme solution for overnight incubation at 37°C. The

reaction products were dialyzed against 50 mM NH_4CO_3, pH 7.6. Elimination of α-mannosidase

was achieved by denaturation (2 min at 110°C) followed by overnight tryptic digestion with

trypsin/α-mannosidase (2% by weight) at 37°C. αtManLAM was recovered after dialysis against

water. Removal of mannose caps was controlled by capillary electrophoresis as described [48].

dManLAM and αtManLAM were suspended in 1x phosphate buffer saline (PBS) pH 7.4 and

stored at -20°C. To inactivate putative lipoprotein contaminating ManLAM and its derivatives,

samples were incubated with 1% H_2O_2 for 3h at 37°C, a treatment that converts the N-terminal

cysteine-thioether substructure of lipoproteins into TLR-2-inactive sulfoxide derivatives [19]. At the

end of the incubation period, H_2O_2 was evaporated under N_2 gas and the resulting pellets were

reconstituted in distilled H_2O.

Cell lines

Murine bone marrow marrow-derived macrophages obtained from C57BL/6 mice and immortalized with J2 retrovirus (iBMDMs) [49] were kindly provided by Dr. Phillip West (Jackson Laboratory, Bar Harbor, ME). iBMDMs and RAW 264.7 cells (ATCC TIB-71) were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with heat-inactivated 10% FBS, 1x penicillin-streptomycin solution (100 IU/mL penicillin and 100 μg/ml Streptomycin; Corning, Manassas, VA) and 20 mM HEPES (Millipore, Billerica, MA) and incubated at 37°C in a humidified atmosphere consisting of 5% CO_2. RAW 264.7 cells carrying an enhanced green fluorescence protein (EGFP)-RelA reporter [25] were cultured in DMEM containing 10% FBS, 2 mM l-glutamine, 1x penicillin-streptomycin solution (Corning, Manassas, VA), and 10 mM HEPES (Millipore, Billerica, MA) at 37°C and 5% CO_2. The HEK-Blue mDectin-2 and HEK-Blue hTLR2 (InvivoGen, San Diego, CA) are derivatives of HEK293 cells that stably express the murine Dectin-2 and the human TLR2 genes, respectively, along with a NF-κB-inducible reporter system (secreted alkaline phosphatase). These cells were maintained in DMEM containing 10% FBS, 4.5 g/l glucose, 2 mM L-glutamine, 1x penicillin-streptomycin solution, 100 μg/ml zeocin, 200 μg/ml hygromycin, 10 μg/ml blasticidin, 1 μg/ml puromycin, and 1x TLR and CLR Selection solution (all from InvivoGen, San Diego, CA). Reporter cells were cultured for two passages without selection before use.

Gene knock-down (KD) in iBMDMs

iBMDMs were transduced with lentivirus carrying nuclease-dead FLAG-tagged Cas9 (dCas9) and selected for using blasticin (5-10 μg/ml) (InvivoGen, San Diego, CA). The Lenti-dCas9-KRAB-blast plasmid was a gift from Gary Hon (U. Texas South Western) (Addgene plasmid # 89567; http://n2t.net/addgene:89567) [50]. For each gene of interest, 5-6 small guide RNAs (sgRNAs) were selected using the CRISPick tool (Broad Institute, https://portals.broadinstitute.org/gppx/crispick/public) [51,52]. sgRNAs were cloned into the pLentiGuide-Puro backbone, which was a gift from Paul Khavari (Stanford U.) (Addgene plasmid # 117986; http://n2t.net/addgene:117986). dCas9-containing cells were transduced with lentivirus carrying gene-specific sgRNAs and selected for by using puromycin (5-10 μg/ml) (InvivoGen, San Diego, CA). Non-targeting sgRNAs (targeting *luxA*) were used as negative controls: *lux1*: GGCAATGAAACGCTACGCTC, *lux2*: ATAAAGAGCGCGCCCAACAC.

Knockdown cells were validated by RT-qPCR, and the cell lines carrying the two gene-specific sgRNAs with the highest knockdown efficiency were used for subsequent experiments.

The gene-specific sgRNAs used in this study were: TLR2-1: TGGGTGTCCCTCTTCCTGCA,

TLR2-2: AAGCTGATCCGCCCGGCTGG; Dectin-2-1: GGACCTGGCTTCTGTCAAAG, Dectin-2-

2: TGAGTTAAATGCCACAGAGC. For both steps of transduction, lentivirus was produced using

HEK293T-Lenti-X cells (Takara) and psPAX2 (Addgene plasmid #12260;

http://n2t.net/addgene:12260) and pMD2.G/VSV-G (Addgene plasmid #12259;

http://n2t.net/addgene:12259) packaging plasmids, which were gifts from Didier Trono (EPFL,

Swiss Federal Technology Institute of Lausanne).

Human monocyte-derived macrophages

Human leukopaks were obtained from the New York Blood Center (Long Island City, NY,

USA). Peripheral blood mononuclear cells (PBMC) were prepared by Ficoll density gradient

centrifugation (Ficoll-Paque, GE Healthcare, Uppsala, Sweden) and human monocyte-derived

macrophages (hMDM) were cultivated as described in previous work {30161232}. Treatments

were performed on day 7 of culture.

M. tuberculosis **infection**

M. tuberculosis strain mc^{2}6206 (H$_{37}$Rv *ΔpanCD ΔleuCD*) was kindly provided by Dr. William

Jacobs Jr. (Albert Einstein College of Medicine) [53]. The strain was confirmed positive for

phtiocerol dimycocerosates (PDIM) both initially by the Jacobs lab via thin layer chromatography

of lipid extracts and during its successive culturing by using vancomycin susceptibility (VAN10)

assays [54]. To prevent PDIM loss, bacteria were cultured in media containing propionate [54]. Prior

to infection experiments, the strain was grown in Middlebrook 7H9 supplemented with 10%

OADC, 0.5% glycerol, 0.05% Tween-80, 24 ug/ml pantothenate, 50 ug/ml leucine, and 0.1 mM

propionate. Macrophage infections were performed as previously described [55,56]. Briefly, *M.*

tuberculosis cultures were grown to an OD$_{600}$ of 0.6-1.0, washed three times with PBS, and

centrifuged at 500 x *g* for 5 min to remove clumps. Bacteria were diluted in DMEM plus 10%

horse serum and added to iBMDMs plated at 3-4x10^5 cells/well in a 12-well dish in iBMDM

media containing 24 ug/ml pantothenate and 50 ug/ml leucine. A diluted bacterial inoculum was

added to cultured iBMDM (MOI = 15), and cells were centrifuged at 1,000 x *g* for 10 min to

synchronize the infection and resuspended in fresh iBMDM media containing 24 ug/ml

pantothenate and 50 ug/ml leucine; culture media was changed daily. After 48 h, iBMDMs were

detached from tissue culture plates by resuspending in 1 x PBS plus 4mM EDTA by gentle

scraping. An equal volume of 8% paraformaldehyde was added to fix and inactivate the

samples. Cells were washed twice with 1 x PBS and processed for LD formation as described

below.

Chemical reagents and antibodies

TLR2 ligand Pam3CSK4 and Dectin-2 ligand furfurman were purchased from Invivogen

(San Diego, CA). Inhibitors of diacylglycerol transferase (A922500) and acyl-coenzyme

A:cholesterol acyltransferase (CAS 615264-52-3), both from Santa Cruz Biotechnology, Dallas,

TX, were stored as per manufacturer's instructions. Inhibitors of Mechanistic Target of

Rapamycin Complex 1 (Rapamycin), peroxisome proliferator-activated receptor gamma

(GW9662), NF-κB (JSH-23 and QNZ) were all purchased from Selleckchem, Houston, TX.

Inhibitor doses were selected at and around published EC_{50} data and toxicity profiles in

macrophages [26,57]. Only inhibitor doses yielding >85% cell viability were utilized by using trypan

blue (Cytiva, Marlborough, MA) and counting with an automated cell counter (LUNA-FX7, Logos

Biosystem, Annandale, VA). Mouse Fc receptor blocking solution, anti-mouse CD16/CD32,

BV421 rat anti-mouse F4/80, BV421 mouse anti-human CD11c (BD Biosciences, San Diego,

CA) and Bodipy 493/503 (Life Technologies, Carlsbad, CA) were used at the dilutions

suggested by the manufacturers and as per previous work [26].

TLR2 and Dectin-2 reporter cell experiments

The HEK-Blue reporter cell experiments were performed following manufacturer's protocols

(13), with minor modifications. For the HEK-Blue hTLR2 reporter assay, 20 µl of ManLAM,

dManLAM, αtManLAM, and Pam3CSK4 (TLR2 ligand), all suspended in 1x PBS, were added to

each well in 96-well plates (in triplicate) to obtain a final concentration of 5 µg/ml of each

reagent. HEK-Blue hTLR2 cells were harvested, washed in 1x PBS, and resuspended in

Quanti-Blue media (InvivoGen, San Diego, CA) at a final density of 280,000 cells/ml. 180 µl of

cell suspension was added to each well and incubated for 24 h at 37°C in a humidified

atmosphere containing 5% CO_2. For the HEK-Blue mDectin-2 reporter assay, 10 µl of ManLAM,

dManLAM, and αtManLAM, and furfurman (Dectin-2 ligand) were suspended in 1x PBS and

added to each well in 96-well plates (in triplicate) to obtain a final concentration of 10 µg/ml for

ManLAM and derivatives and 200 µg/ml for furfurman. The plates were dried at 37 °C until the

solvent evaporated (~3 h). HEK-Blue mDectin-2 cells were harvested, washed in 1x PBS, and

resuspended in Quanti-Blue media at a final density of 280,000 cells/ml. 200 µl of cell

suspension was added to each well and incubated for 24 h at 37°C in a humidified atmosphere

containing 5% CO_2. For both reporter cell lines, the colorimetric reading was taken at 650 nm

(OD$_{650}$). The data were expressed as background-subtracted OD$_{650}$ or OD$_{650}$ ratio between

treated cells and vehicle control cells.

Lipid droplet measurements

One million murine iBMDM cells or one million human PBMCs in 1 ml culture medium were

added to each well of 12-well tissue culture plates. Incubation was 2 h for murine cells and 7

days at 37°C in a humidified atmosphere containing 5% CO$_2$. *M. tuberculosis* components and

control agonists were diluted in the same medium and 1 ml of each suspension or solution was

added to each well to obtain final concentrations of 500 ng/ml for ManLAM, ManLAM

derivatives, and Pam3CSK4, and 10 µg/ml for furfurman. Treated and vehicle control cells were

incubated for 24 h at 37°C in a humidified atmosphere containing 5% CO$_2$. When appropriate,

chemical inhibitors were added at the indicated doses in a 10 µl solution, concurrently with the

treatments above.

After incubation, cells were subjected to imaging flow cytometry as described [26]. Briefly,

medium was discarded and 1 ml 1x PBS was added to each well. Macrophages were detached

from culture plates by gentle scraping, pelleted by centrifugation at 350g for 5 min, and fixed

with 4% paraformaldehyde in 1x PBS for 45 min at RT. Cells were washed with 1x PBS

containing 0.1% bovine serum albumin (PBS-BSA), resuspended in 50 µl of PBS-BSA

containing 5 µl of mouse Fc-receptor blocking solution, and incubated at RT for 7 min. After

incubation, 50 µl of PBS-BSA containing 5 µl of mouse F4/80-BV421 or human CD11c-BV421

antibody were added to each tube, and samples were incubated for 30 min at 4°C. After

washing with PBS-BSA, cells were stained with 0.3 µg/ml Bodipy 493/503 in 1x PBS for 15 min.

For each condition, data were acquired from 5,000-10,000 F4/80-positive cells utilizing an

ImageStreamXMark II imaging flow cytometer (Amnis Corporation, Seattle, WA), using 60x

magnification. Image data were analyzed by IDEAS software version 6.0 (Amnis Corporation,

Seattle, WA) after applying a compensation matrix and selecting the region of interest (lipid

droplets) with the Spot Mask tool. Mean fluorescence intensity (MFI) and spots per cell were

extracted and expressed as ratio of treated vs its vehicle control cells.

Real Time PCR analysis

RNA was isolated using RNeasy Mini Kit (Qiagen, Valencia, CA, USA). Isolated RNA was

subjected to cDNA synthesis using the SuperScriptTM First-Strand Synthesis System (Bio-Rad,

Hercules, CA). Real time PCR was performed in a QuantStudio 7 Flex Real-Time time PCR

machine (Thermo Fisher Scientific, Waltham, MA, USA) using the ABsolute QPCR SYBER

Green Mix (ABgene, Rochester, NY, USA), according to the standard ABgene protocol. The

sequence of the primers used for real time PCR are listed in **SI Table 1**. The internal control

gene β-Actin was amplified simultaneously in a separate reaction. The threshold cycle number

(CT) was determined by qPCR for triplicate reactions, and the mean CT of the triplicate

reactions was calculated. Levels of expression of the genes of interest were normalized to β-

Actin using the formula $2^{-\Delta DCT}$, where $-\Delta DCT = \Delta CT$ (treated) $- \Delta CT$ (control), and ΔCT is the

CT of the target gene subtracted from the CT of the housekeeping gene β-Actin.

Measurements of RelA nuclear translocation (NF-κB activation)

For the Rel-A nuclear translocation assay, RAW264.7 cells carrying a EGFP-RelA reporter

gene fusion were plated at 4×10^5 cells per well in a 96-well glass-bottom plate overnight and

then serum starved for 3 h. Cells were stimulated with the same treatments used for lipid

accumulation experiments, and incubated for 0.5, 1.0, 1.5 and 2.0 h at 37°C. For fixed cell

imaging, cells were fixed after stimulation with 2% PFA on ice for 5 min. Nuclei were

counterstained with 2 µg/mL Hoechst stain at RT for 15 min. Fluorescence images were

acquired using a Nikon Eclipse Ti microscope equipped with an Andor iXon3 EMCCD camera

and a Nikon Plan Apo 40×/0.95 objective. For each sample, 20 images were taken at

different locations. Images were analyzed using ImageJ following a previously reported method

[25].

Statistical analysis

All values are presented as means ± standard deviation (SD) for technical triplicates.

Comparisons between two groups were performed using a two-tailed Student's t-test in the

Prism graphpad software or by one-way ANOVA. The criterion for statistical significance was $p <$

0.05.

Acknowledgements

We thank John Chan and Karl Drlica for critical comments on the manuscript, and staff of

the Flow Cytometry and Immunology Core Laboratory at Rutgers New Jersey Medical School

for assistance with imaging flow cytometry. This work was funded in part by AHA fellowship

24PRE1198252 (J.R.), NIH grants R01HL149450, R01AI158911, NCATS UM1TR004789

(M.L.G.), R35GM124918 (Y. Y.), NIH DP2AI154429 (S.B.), and a grant from the Fondation pour

la Recherche Médicale Equipes FRM DEQ20180339208 (J.N.).

Author contributions

D.N., M.G., S.B., J.N., and Y.Y. contributed to the experimental design; D.N., J.R., O.V.,

S.B., J.C., and J.N. performed experiments and analyzed data; D.N. and M.G. wrote the

manuscript. All authors reviewed and edited the manuscript.

References

1.   WorldHealthOrganization. Global tuberculosis report. (World Health Organization,
Geneva, Switzerland, 2025).

2.   Flynn, J.L., Chan, J. & Lin, P.L. Macrophages and control of granulomatous inflammation
in tuberculosis. *Mucosal immunology* **4**, 271-278 (2011).

3.   Ramakrishnan, L. Revisiting the role of the granuloma in tuberculosis. *Nat Rev Immunol*
12, 352-366 (2012).

4.   Guerrini, V. & Gennaro, M.L. Foam Cells: One Size Doesn't Fit All. *Trends Immunol* **40**,
1163-1179 (2019).

5.   Peyron, P*., et al.* Foamy macrophages from tuberculous patients' granulomas constitute
a nutrient-rich reservoir for M. tuberculosis persistence. *PLoS Pathog* **4**, e1000204
(2008).

6.   Greenwood, D.J*., et al.* Subcellular antibiotic visualization reveals a dynamic drug
reservoir in infected macrophages. *Science* **364**, 1279-1282 (2019).

7.   Nigou, J., Gilleron, M. & Puzo, G. Lipoarabinomannans: from structure to biosynthesis.
Biochimie **85**, 153-166 (2003).

8.   Mishra, A.K., Driessen, N.N., Appelmelk, B.J. & Besra, G.S. Lipoarabinomannan and
related glycoconjugates: structure, biogenesis and role in Mycobacterium tuberculosis
physiology and host-pathogen interaction. *FEMS Microbiol Rev* **35**, 1126-1157 (2011).

9.   Torrelles, J.B. & Chatterjee, D. Collected Thoughts on Mycobacterial
Lipoarabinomannan, a Cell Envelope Lipoglycan. *Pathogens* **12**(2023).

10.   Sparks, I.L*., et al.* Lipoarabinomannan mediates localized cell wall integrity during
division in mycobacteria. *Nat Commun* **15**, 2191 (2024).

11.   Decout, A*., et al.* Deciphering the molecular basis of mycobacteria and lipoglycan
recognition by the C-type lectin Dectin-2. *Sci Rep* **8**, 16840 (2018).

12.   Yonekawa, A*., et al.* Dectin-2 is a direct receptor for mannose-capped
lipoarabinomannan of mycobacteria. *Immunity* **41**, 402-413 (2014).

13.   Nigou, J*., et al.* Mannan chain length controls lipoglycans signaling via and binding to
TLR2. *J Immunol* **180**, 6696-6702 (2008).

14.   Ray, A., Cot, M., Puzo, G., Gilleron, M. & Nigou, J. Bacterial cell wall macroamphiphiles:
pathogen-/microbe-associated molecular patterns detected by mammalian innate
immune system. *Biochimie* **95**, 33-42 (2013).

15.   D'Avila, H.*, et al.* Mycobacterium bovis bacillus Calmette-Guerin induces TLR2-mediated
577   formation of lipid bodies: intracellular domains for eicosanoid synthesis in vivo. *J*
578   *Immunol* **176**, 3087-3097 (2006).

579   16.   Almeida, P.E.*, et al.* Mycobacterium bovis bacillus Calmette-Guerin infection induces
580   TLR2-dependent peroxisome proliferator-activated receptor gamma expression and
581   activation: functions in inflammation, lipid metabolism, and pathogenesis. *J Immunol*
582   **183**, 1337-1345 (2009).

583   17.   Shukla, S., Richardson, E.T., Drage, M.G., Boom, W.H. & Harding, C.V. Mycobacterium
584   tuberculosis Lipoprotein and Lipoglycan Binding to Toll-Like Receptor 2 Correlates with
585   Agonist Activity and Functional Outcomes. *Infect Immun* **86**(2018).

586   18.   Krishna, S.*, et al.* Lipoglycans contribute to innate immune detection of mycobacteria.
587   *PLoS One* **6**, e28476 (2011).

588   19.   Zahringer, U., Lindner, B., Inamura, S., Heine, H. & Alexander, C. TLR2 - promiscuous or
589   specific? A critical re-evaluation of a receptor expressing apparent broad specificity.
590   *Immunobiology* **213**, 205-224 (2008).

591   20.   Spera, I.*, et al.* The J2-Immortalized Murine Macrophage Cell Line Displays
592   Phenotypical and Metabolic Features of Primary BMDMs in Their M1 and M2
593   Polarization State. *Cancers (Basel)* **13**(2021).

594   21.   Sato, K.*, et al.* Dectin-2 is a pattern recognition receptor for fungi that couples with the
595   Fc receptor gamma chain to induce innate immune responses. *J Biol Chem* **281**, 38854-
596   38866 (2006).

597   22.   Gilleron, M., Nigou, J., Nicolle, D., Quesniaux, V. & Puzo, G. The acylation state of
598   mycobacterial lipomannans modulates innate immunity response through toll-like
599   receptor 2. *Chem Biol* **13**, 39-47 (2006).

600   23.   Liu, T., Zhang, L., Joo, D. & Sun, S.-C. NF-κB signaling in inflammation. *Signal*
601   *Transduction and Targeted Therapy* **2**, 17023 (2017).

602   24.   Hayden, M.S. & Ghosh, S. Shared Principles in NF-κB Signaling. *Cell* **132**, 344-362
603   (2008).

604   25.   Li, W., Li, M., Anthony, S.M. & Yu, Y. Spatial organization of FcγR and TLR2/1 on
605   phagosome membranes differentially regulates their synergistic and inhibitory receptor
606   crosstalk. *Scientific Reports* **11**, 13430 (2021).

607   26.   Guerrini, V.*, et al.* Storage lipid studies in tuberculosis reveal that foam cell biogenesis is
608   disease-specific. *PLoS Pathog* **14**, e1007223 (2018).

27. Salamon, H.*, et al.* Cutting Edge: Vitamin D Regulates Lipid Metabolism in
Mycobacterium tuberculosis Infection. *J Immunol* (2014).

28. Laplante, M. & Sabatini, D.M. Regulation of mTORC1 and its impact on gene expression
at a glance. *J Cell Sci* **126**, 1713-1719 (2013).

29. Ricoult, S.J. & Manning, B.D. The multifaceted role of mTORC1 in the control of lipid
metabolism. *EMBO Rep* **14**, 242-251 (2013).

30. Angela, M.*, et al.* Fatty acid metabolic reprogramming via mTOR-mediated inductions of
PPARgamma directs early activation of T cells. *Nat Commun* **7**, 13683 (2016).

31. Arimura, N., Horiba, T., Imagawa, M., Shimizu, M. & Sato, R. The Peroxisome
Proliferator-activated Receptor γ Regulates Expression of the Perilipin Gene in
Adipocytes *. *Journal of Biological Chemistry* **279**, 10070-10076 (2004).

32. Cogswell, J.P.*, et al.* NF-kappa B regulates IL-1 beta transcription through a consensus
NF-kappa B binding site and a nonconsensus CRE-like site. *J Immunol* **153**, 712-723
(1994).

33. Land, S.C. & Tee, A.R. Hypoxia-inducible Factor 1α Is Regulated by the Mammalian
Target of Rapamycin (mTOR) via an mTOR Signaling Motif*. *Journal of Biological
Chemistry* **282**, 20534-20543 (2007).

34. van Uden, P., Kenneth, N.S. & Rocha, S. Regulation of hypoxia-inducible factor-1alpha
by NF-kappaB. *Biochem J* **412**, 477-484 (2008).

35. Rajaram, M.V.*, et al.* Mycobacterium tuberculosis activates human macrophage
peroxisome proliferator-activated receptor gamma linking mannose receptor recognition
to regulation of immune responses. *J Immunol* **185**, 929-942 (2010).

36. Plato, A., Hardison, S.E. & Brown, G.D. Pattern recognition receptors in antifungal
immunity. *Semin Immunopathol* **37**, 97-106 (2015).

37. Li, W., Yan, J. & Yu, Y. Geometrical reorganization of Dectin-1 and TLR2 on single
phagosomes alters their synergistic immune signaling. *Proceedings of the National
Academy of Sciences* **116**, 25106-25114 (2019).

38. Li, M., Vultorius, C., Bethi, M. & Yu, Y. Spatial Organization of Dectin-1 and TLR2 during
Synergistic Crosstalk Revealed by Super-resolution Imaging. *The Journal of Physical
Chemistry B* **126**, 5781-5792 (2022).

39. Li, M. & Yu, Y. Innate immune receptor clustering and its role in immune regulation. *J
Cell Sci* **134**(2021).

40. Laird, M.H.*, et al.* TLR4/MyD88/PI3K interactions regulate TLR4 signaling. *J Leukoc Biol*
85, 966-977 (2009).

41.   Hatton, O.*, et al.* Syk activation of phosphatidylinositol 3-kinase/Akt prevents HtrA2-
644   dependent loss of X-linked inhibitor of apoptosis protein (XIAP) to promote survival of
645   Epstein-Barr virus+ (EBV+) B cell lymphomas. *J Biol Chem* **286**, 37368-37378 (2011).

646   42.   Manning, B.D. & Cantley, L.C. AKT/PKB signaling: navigating downstream. *Cell* **129**,
647   1261-1274 (2007).

648   43.   Kim, M.J.*, et al.* Caseation of human tuberculosis granulomas correlates with elevated
649   host lipid metabolism. *EMBO Mol Med* **2**, 258-274 (2010).

650   44.   Dkhar, H.K.*, et al.* Mycobacterium tuberculosis keto-mycolic acid and macrophage
651   nuclear receptor TR4 modulate foamy biogenesis in granulomas: a case of a
652   heterologous and noncanonical ligand-receptor pair. *J Immunol* **193**, 295-305 (2014).

653   45.   Liu, S.*, et al.* Mycobacterium tuberculosis suppresses host DNA repair to boost its
654   intracellular survival. *Cell Host Microbe* **31**, 1820-1836 e1810 (2023).

655   46.   Nigou, J.*, et al.* The phosphatidyl-myo-inositol anchor of the lipoarabinomannans from
656   Mycobacterium bovis bacillus Calmette Guerin. Heterogeneity, structure, and role in the
657   regulation of cytokine secretion. *J Biol Chem* **272**, 23094-23103 (1997).

658   47.   Lanéelle, M.A., Spina, L., Nigou, J., Lemassu, A. & Daffé, M. Lipid and
659   Lipoarabinomannan Isolation and Characterization. *Methods Mol Biol* **2314**, 109-150
660   (2021).

661   48.   Sidobre, S., Nigou, J., Puzo, G. & Riviere, M. Lipoglycans are putative ligands for the
662   human pulmonary surfactant protein A attachment to mycobacteria. Critical role of the
663   lipids for lectin-carbohydrate recognition. *J Biol Chem* **275**, 2415-2422 (2000).

664   49.   De Nardo, D., Kalvakolanu, D.V. & Latz, E. Immortalization of Murine Bone Marrow-
665   Derived Macrophages. *Methods Mol Biol* **1784**, 35-49 (2018).

666   50.   Xie, S., Duan, J., Li, B., Zhou, P. & Hon, G.C. Multiplexed Engineering and Analysis of
667   Combinatorial Enhancer Activity in Single Cells. *Mol Cell* **66**, 285-299.e285 (2017).

668   51.   Doench, J.G.*, et al.* Optimized sgRNA design to maximize activity and minimize off-
669   target effects of CRISPR-Cas9. *Nature Biotechnology* **34**, 184-191 (2016).

670   52.   Sanson, K.R.*, et al.* Optimized libraries for CRISPR-Cas9 genetic screens with multiple
671   modalities. *Nature Communications* **9**, 5416 (2018).

672   53.   Vilcheze, C.*, et al.* Rational Design of Biosafety Level 2-Approved, Multidrug-Resistant
673   Strains of Mycobacterium tuberculosis through Nutrient Auxotrophy. *mBio* **9**(2018).

674   54.   Mulholland, C.V.*, et al.* Propionate prevents loss of the PDIM virulence lipid in
675   Mycobacterium tuberculosis. *Nat Microbiol* **9**, 1607-1618 (2024).

55.   Bell, S.L., Lopez, K.L., Cox, J.S., Patrick, K.L. & Watson, R.O. Galectin-8 Senses
Phagosomal Damage and Recruits Selective Autophagy Adapter TAX1BP1 To Control
Mycobacterium tuberculosis Infection in Macrophages. *mBio* **12**, e0187120 (2021).
56.   Watson, R.O*., et al.* The Cytosolic Sensor cGAS Detects Mycobacterium tuberculosis
DNA to Induce Type I Interferons and Activate Autophagy. *Cell Host Microbe* **17**, 811-819
(2015).
57.   Shin, H.M.*, et al.* Inhibitory action of novel aromatic diamine compound on
lipopolysaccharide-induced nuclear translocation of NF-kappaB without affecting
IkappaB degradation. *FEBS Lett* **571**, 50-54 (2004).

Figure legends

Figure 1: Involvement of TLR2 signaling in lipid droplet induction by *M. tuberculosis* lipoarabinomannan. A. *M. tuberculosis* ManLAM (500 ng/ml), H_2O_2-treated ManLAM (ManL+H; 500 ng/ml), Pam3CSK4 (PAM, synthetic TLR2 agonist; 500 ng/ml) and vehicle control (C) were tested for ability to induce NF-κB activation in HEK-TLR2 reporter cells. Reporter cells were treated for 24h and TLR2 activity was measured as a function of inducible secreted embryonic alkaline phosphatase in a colorimetric assay at 650nm. The Y-axis shows background-subtracted OD_{650} values. **B.** CRISPRi technology was used to generate TLR2 knock-down (KD) murine iBMDM. Control (non-targeting guide RNAs) and TLR2 KD clones were treated for 24h with 500ng/ml ManLAM preparations and Pam3CSK4. Abbreviations are as in panel A. Macrophages were stained with Bodipy 493/503 and antibodies to the macrophage marker F4/80. Data were obtained by imaging flow cytometry in triplicate and expressed as ratio of Bodipy mean fluorescence intensity (MFI) of treated vs. vehicle-treated cells. The horizontal dotted line marks the vehicle-treated cell reading. Comparison for statistical significance was between treated and vehicle-treated cells. **C**. Representative microscopy images (60x magnification) of vehicle-control (C) and treated iBMDM in the experiment shown in panel B. Fluorescence stain: Bodipy 493/503 (neutral lipid dye, green fluorescence). **D.** iBMDM were treated with ManLAM (500 ng/ml) and increasing doses of Pam3CSK4 (10, 50, 100, 500 ng/ml) for 24h. Data were generated and expressed as in panel B. **E.** Representative images (60x magnification) of the experiment shown in panel D. Description of the images as in panel C. *, $p <0.05$ by unpaired *t* test in all relevant panels.

Figure 2: Involvement of Dectin-2 signaling in lipid droplet induction by *M. tuberculosis* lipoarabinomannan. A. *M. tuberculosis* ManLAM (500 ng/ml), H_2O_2 treated ManLAM (ManL+H, 500 ng/ml), furfurman (FUR; commercial Dectin-2 agonist; 10 µg/ml) and vehicle control (C) were tested were tested for ability to induce NF-κB activation in HEK-Dectin-2 reporter cells. Data were generated and analyzed as in Fig. 1A. **B.** Control and Dectin-2 knockdown (KD) iBMDM were treated 24h with 500ng/ml ManLAM preparations. Abbreviations are as in panel A. Imaging flow cytometry data were generated and expressed as in Fig. 1B. **C.** Representative microscopy images (60x magnification) of vehicle-control (C) and treated iBMDM in the experiment shown in panel B. Panel description as in Fig. 1C. **D.** iBMDM were treated with ManLAM (500 ng/ml) doses of furfurman (FUR; 2.5, 5 and 10 µg/ml). Imaging flow cytometry data were generated and expressed as in Fig. 1B. **E.** Representative microscopy

images (60x magnification) of vehicle-control (C) and treated iBMDM in the experiment shown in

panel D. Panel description as in Fig. 1C. *, $p < 0.05$ by unpaired t test in all relevant panels.

Figure 3: Involvement of *M. tuberculosis* lipoarabinomannan moieties in lipid droplet

induction. A. iBMDM were treated with Pam3CSK4 (PAM; 500 ng/ml) and furfurman (FUR; 10

µg/ml), singly or in combination (P+R), and corresponding vehicle controls. Imaging flow

cytometry data were generated and expressed as in Fig. 1B. **B.** Representative microscopy

images (60x magnification) of vehicle-control (C) and treated iBMDM in the experiment shown in

panel A. Panel description as in Fig. 1C. **C.** *M. tuberculosis* ManLAM, demannosylated ManLAM

(αtManLAM), and deacylated ManLAM (dManLAM) (all H_2O_2 pre-treated, 500 ng/ml) and

corresponding vehicle controls were tested for ability to induce NF-κB activation in HEK-TLR2

and HEK-Dectin 2 reporter cells. Treatment was for 24h as in preceding figures. Data are

presented as ratio of OD_{650} values obtained with treated vs vehicle control cells. The horizontal

dotted line marks the corresponding vehicle-control reading. **D.** ManLAM and derivatives, as in

panel C, were used to treat iBMDM (all 500 ng/ml for 24h). Imaging flow cytometry data were

generated and expressed as in Fig. 1B. **E.** Representative microscopy images (60x

magnification) of vehicle-control (C) and treated iBMDM in the experiment shown in panel D.

Panel description as in Fig. 1C. *, $p < 0.05$ by unpaired t test in all relevant panels.

Figure 4: Ligand-dependent NF-κB RelA nuclear translocation dynamics in EGFP-RelA

RAW264.7 macrophages. **A.** Schematic illustrating cytoplasm-to-nucleus translocation of NF-

κB RelA (p65) upon macrophage cell stimulation. Grey circles, ManLAM and derivatives; red

rods, RelA. **B.** (Top row) Time-lapse live-cell epifluorescence images showing EGFP-RelA

translocation following ManLAM stimulation. (Bottom row) Same cell images shown with EGFP-

RelA fluorescence intensity encoded in pseudo-color, as indicated by the color bar. Scale bars:

5 µm. **C.** Quantification of RelA nuclear translocation over time for the representative cell shown

in panel B and for an unstimulated control cell, expressed as the fluorescence intensity ratio of

EGFP-RelA in the nucleus relative to the cytoplasm. **D.** Representative epifluorescence images

of fixed EGFP-RelA RAW264.7 macrophages stimulated with *M. tuberculosis* ManLAM,

demannosylated ManLAM (αtManLAM), and deacylated ManLAM (dManLAM) or furfurman

(FUR) for 240 min, or with Pam3CSK4 for 60 min, illustrating ligand-induced RelA nuclear

translocation. Scale bars: 10 µm. **E.** Percentage of responsive and non-responsive cells under

various ligand stimulation conditions, as indicated. Responsive cells were defined as those with

a RelA nucleus-to-cytosol ratio greater than 1.089, a threshold set to achieve 95% of

unstimulated cells being classified as non-responsive. Approximately 200 cells per condition

were analyzed; the fraction of responsive cells was 40% for ManLAM, 21% for αtManLAM, 16%

for dManLAM, 28% for furfurman (FUR), and 80% for Pam3CSK4. **F.** Single-cell scatter plots of

NF-κB activation, expressed as the nucleus-to-cytosol EGFP-RelA fluorescence intensity ratio,

in all responsive cells after 240 min ligand stimulation (for ManLAM and derivatives, and

furfurman) and 60 min for Pam3CSK4. Each dot represents one cell. Box plots show the

interquartile range and the median line; whiskers indicate ±1.5× SD. **G.** Expression levels of

Tnfa and *Il1b* in RAW 264.7 cells treated with ManLAM, αtManLAM and dManLAM for 6 hours

were measured by RTqPCR. Measurements were performed on triplicate samples and

expressed as $\log_2$ fold change relative to the housekeeping actin gene. Statistical significance

was assessed by one-way ANOVA: ***$P < 0.001$; **$P < 0.01$; *$P < 0.05$.

Figure 5: Effects on lipid droplet content of knocking down TLR2 and Dectin-2 in *M.*

***tuberculosis*-infected macrophages and treatment with neutral lipid biosynthesis**

inhibitors in lipoarabinomannan-treated macrophages. A. iBMDM, control (C, non-targeting

guide RNAs), TLR2 knock-down (KD), and Dectin-2 (Dec-2) KD iBMDM clones were left

uninfected or infected with *M. tuberculosis* mc^{2}6206 (MOI= 15) for 48h in triplicate wells,

washed, and stained with Bodipy 493/503 and antibodies to the macrophage marker F4/80 for

imaging flow cytometry. Data were generated and expressed as in Fig. 1B. **B.** Representative

microscopy images (60x magnification) of uninfected (U) and infected (Mtb) iBMDM in the

experiment shown in panel A. Panel description as in Fig. 1C. **C, D.** iBMDM were treated with

ManLAM (500ng/ml) and vehicle-treated or treated with increasing doses of the diacylglycerol

transferase inhibitor, A922500 (DGAT-i; 60, 90, 120ng/ml) and the acyl-coenzyme A:cholesterol

acyltransferase inhibitor, CAS 615264-52-3 (ACAT-i; 5, 10, 15µg/ml). Imaging flow cytometry

data for lipid droplet content were generated and expressed as in Fig. 1B. *, $p < 0.05$ by

unpaired *t* test in all relevant panels. **E.** iBMDM were treated with ManLAM (500ng/ml) and

vehicle-treated or treated with increasing doses of the PPARγ inhibitor GW9662 (0.5, 1, 2µM),

mTORC1 inhibitor rapamycin (0.2, 0.4, 0.8nM), and the NF-κB inhibitor JSH-23 (3.5, 7, 14µM).

Imaging flow cytometry data for lipid droplet content were generated and expressed as in Fig.

1B. **F.** iBMDM were treated with ManLAM (500ng/ml) plus vehicle-control (ManL), the mTORC1

inhibitor rapamycin (ManL+R, 0.4nM), and the NF-κB inhibitor JSH-23 (ManL+J, 7µM) for 6

hours. Expression levels of *Plin2*, *Tnfa, Il1b, and Hif1a* were measured by RTqPCR in triplicate

and expressed as $\log_2$ fold change, relative to the housekeeping actin gene. *, $p < 0.05$ by

unpaired *t* test in all relevant panels.

Figure 6: Dual receptor recognition of mannosylated lipoarabinomannan integrates

macrophage signaling. Mannosylated lipoarabinomannan (ManLAM) engages TLR2 and

Dectin-2 through distinct structural features. Lipid acyl groups interact with TLR2, whereas

terminal mannose caps are recognized by Dectin-2. Signals from both receptors converge on

shared intracellular pathways and diverge to activate two responses: mTORC1/PPARγ–

dependent lipid droplet formation and NF-κB–dependent TNFα production. Illustration

generated with AI-assisted tools and edited by the authors.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6