Spatial and temporal localization of *Serratia ureilytica* causing cucurbit yellow vine disease
in cucurbits indicates phloem-associated colonization and systemic movement

Kensy D. Rodriguez-Herrera[1*], Elise Boisvert[,2*], Margaret H Frank[2], and Christine D. Smart[1, †]

[1] Plant Pathology and Plant-Microbe Biology Section, School of Integrative Plant Science,

Cornell AgriTech, Cornell University, Geneva, NY, 14456

[2] Plant Biology Section, School of Integrative Plant Science, Cornell University, Ithaca,

NY 14853, USA

*These authors contributed equally to this work

[†] Corresponding author: C. D. Smart; Email: cds14@cornell.edu

Abstract (180 words)

Cucurbit yellow vine disease (CYVD), caused by the bacterium *Serratia ureilytica*, is a phloem-

associated disease of cucurbits. This study characterized the spatial and temporal distribution of

S. ureilytica in *Cucurbita pepo* cultivar 'Delicata' plants under greenhouse conditions using a

GFP-tagged isolate (P01). Seedlings were sampled weekly for four weeks. Transverse sections

from the stem, petiole, leaf, shoot apex, and root were imaged by laser scanning confocal and

fluorescent dissecting microscopy. In parallel, bacterial abundance in each plant tissue was

assessed by quantifying colony-forming units (CFU) via droplet plating over a 4-week time

course. Across plant tissues and time points, *S. ureilytica* fluorescent signal was primarily

concentrated in the inner and outer periphery of the bicollateral vascular bundles, with higher

magnification images revealing mainly symplastic localization within phloem-associated cells.

Consistent with the imaging results, bacterial quantification data showed a high abundance of

CFUs in the main stem across weeks, with an irregular pattern of presence in the distal tissues at

later time points. These results suggest that *S. ureilytica* is predominantly localized within

phloem-associated cells and spreads both acropetally and basipetally during infection.

Introduction

Serratia ureilytica (formerly known as *Serratia marcescens*) is a relatively understudied vascular-based plant pathogen that infects cucurbit crops, causing Cucurbit Yellow Vine Disease (CYVD) (Mphande et al. 2025b; Rascoe et al. 2003). *S. ureilytica* is transmitted by the squash bug (*Anasa tristis*), a common pest of cucurbits (Bextine et al. 2001; Bonjour and Fargo 1989; Pair et al. 2004), as well as cucumber beetles (Mphande et al. 2025b). Symptoms of CYVD include yellowing, leaf scorching, stunting, and honey-brown discoloration of the phloem (Bruton et al. 2003, 1998a). CYVD was first reported in 1988 in Texas and Oklahoma and has since spread to several states in the US, including recent reports from New York and Iowa (Mphande et al., 2024a; Rodriguez-Herrera et al., 2023). Losses due to CYVD have been reported from 5%-100% (Zhang et al. 2005). Previous work proposes that *S. ureilytica* is present in phloem sieve tubes, the intercellular space of parenchyma, and xylem vessels using scanning electron micrographs and fluorescence microscopy (Bruton et al. 1998b; Luo 2006).

Bacterial colonization of plant vascular tissue systems is a well-documented route for plant pathogens. One such example is *Erwinia tracheiphila*, the causative agent of bacterial wilt in cucurbits (Rojas et al. 2015). This pathogen colonizes the xylem, where it clogs vessel elements(Rojas et al. 2015). The infection ultimately blocks water transport, causing rapid wilting and, often, death (Rojas et al. 2015). Huanglongbing (HLB), caused by '*Candidatus Liberibacter asiaticus*', is another vasculature-localized disease that causes tremendous crop loss in citrus groves (Gottwald 2010; Perilla-Henao and Casteel 2016). This pathogen resides in

phloem sieve tube cells, where it consumes host carbon sources, replicates, and clogs phloem

transport (Ding et al. 2015). Another example of a phloem colonizer is *Candidatus*

Arsenophonus phytopathogenicus, which causes low-sugar syndrome in sugar beets and

marginal chlorosis in strawberries. This pathogen can metabolize sugars from phloem sap as its

sole source of nutrition (Mahillon et al. 2025). Phytoplasmas are another, well-described class of

pathogens that colonize host sieve elements. Phytoplasmas can adhere to cell surfaces, move

passively through the phloem, and, in some cases, such as *Candidatus Phytoplasma asteris*,

which causes aster yellows disease, manipulate plant development to induce distinctive

phenotypes through secreted effector proteins (Bendix and Lewis 2018). The majority of

phloem-colonizing pathogens share commonalities: they are found in phloem sieve tube

elements and are generally unculturable in artificial media (Bendix and Lewis 2018; Haider et al.

2024; Mahillon et al. 2025). Although several examples of phloem-colonizing pathogens are

known, they all infect sieve elements; little is known about pathogens that colonize other

phloem-associated cells.

Vascular plants transport water and nutrients via xylem and phloem (Agustí and Blázquez 2020).

Xylem transports water and solutes from the soil upward, whereas phloem transports

photoassimilates and signaling molecules throughout the plant, from source to sink tissues

(Taylor et al. 2009). In flowering plants, the phloem is composed of enucleate sieve elements

through which phloem sap travels, companion cells that support sieve tubes in phloem sap

transport, and which are extensively connected with sieve tube elements via specialized

plasmodesmatal pores to form sieve element-companion cell complexes (Van Bel 2003; Haider

et al. 2024; Reidel et al. 2009; De Schepper et al. 2013; Turgeon 2017). Companion cells are

relatively small and contain the regulatory and metabolic mechanisms for the transport of

phloem sap (Braun 2022; Perilla-Henao and Casteel 2016). Phloem parenchyma are

metabolically active cells within the vascular bundle that can function as transient or long-term

storage for carbohydrates and other solutes (Esau 1969; De Schepper et al. 2013).

Phloem physiology can be broken down into three functions: collection, transport, and release

(de Schepper et al., 2013). In source tissues (i.e., mature leaves) where photosynthesis occurs,

sugars are produced and loaded into the sieve element-companion cell complexes of collection

phloem, passed via transport phloem throughout the plant, and finally offloaded into sink tissues

via the release phloem. Sink tissues change throughout development and can include meristems,

new leaves, and fruits, but most sugars are directed to the roots. According to the long-accepted

Münch theory, this mass flow is driven by a pressure gradient, with high solute concentrations at

the source and low concentrations at the sink providing the necessary force (Münch, 1930; de

Schepper et al., 2013). One key modification to this theory is the "leakage retrieval mechanism,"

in which loading and unloading occur dynamically between sieve element-companion cell

complexes and phloem parenchyma throughout the transport phloem (Van Bel 2003; De

Schepper et al. 2013; Thorpe et al. 2005). This model, which helps to explain how the pressure

gradient can be tightly regulated, also provides a framework for understanding how bacteria

utilizing the phloem can colonize not only terminal sink tissues, but also lateral exchange sites

along the transport pathway (De Schepper et al. 2013). A notable feature of Cucurbit vasculature

is its bicollateral vascular bundles, in which phloem occurs on both the internal and external

sides of the xylem, rather than only externally as in a typical vascular bundle (Turgeon 2017;

Zhang et al. 2012).

There is very little research on the localization of *S. ureilytica* in cucurbits. Using electron

microscopy, Bruton et al. 1998 reported that, once in the plant, S. *ureilytica* moves through the

phloem sieve tube elements. Recent studies have examined this bacterium in the main stem at a

single time point by looking at the number of bacteria in different tissues (Mphande et al. 2025a).

However, it is still unclear how the bacteria colonize the plant over time. By using fluorescence

microscopy and sampling at multiple developmental stages, this study provides valuable new

insights into *S. ureilytica* localization and movement as the infection spreads throughout the

plant. This project had the following objectives: i) Determine the cellular and anatomical

localization of *S. ureilytica* in squash plants at weekly intervals after inoculation using dissecting

and confocal microscopy, and ii) quantify the distribution of *S. ureilytica* in different plant tissues

throughout time using bacterial quantification methods.

Materials and Methods

Bacterial growth and storage. Two isolates were used for this experiment: Isolate 22212 was

recovered from Ontario County, New York, at the AgriTech Research Station (Geneva, NY) in

2021, and is known to cause typical CYVD symptoms following greenhouse inoculation of

plants (Rodriguez-Herrera et al. 2023). The green fluorescent protein (GFP)-tagged isolate (P01)

was received from Elizabeth Little, University of Georgia, and was isolated in Georgia in 2015

(Besler 2014; Besler and Little 2017). The GFP insertion was performed in the Gerardo Lab at

Emory University, as described by Mendiola et al. (2022). Isolates were stored at -80 °C in

Microbank tubes (Pro-Lab Diagnostics, Ontario, Canada). To grow them from storage, three to

four beads from each Microbank tube were plated onto Luria Broth (LB) agar (Bertani 1951)

supplemented with tetracycline (20 μg/ml) (Mphande et al. 2025b) and incubated overnight.

Plant cultivation. Seeds (n=100) of winter squash (*Cucurbita pepo* cv Delicata; Johnny's

Selected Seeds, Winslow, ME, USA) were germinated in deep-well 50-cell flats (T.O. Plastics,

Clearwater, MN, USA) filled with soilless potting mix (Lambert's LM-3 All Purpose Mix;

Quebec, Canada). Once the seedlings had emerged (approximately 7 days after planting), they

were transplanted into two-gallon pots (Grower's Solution One Gallon Black Trade Pots;

Cookeville, TN, USA). All plants were grown in Cornell University's greenhouse facilities under

a 16-hour photoperiod and day/night temperatures of 23 °C/21 °C.

Comparison of isolates. In all experiments, isolate 22212 was included as a positive control due

to its known pathogenicity on cucurbits. Isolates P01-GFP and 22212 were compared to confirm

their pathogenicity and inoculation success rate.

Inoculations: Four treatments were established: Isolate P01-GFP, isolates 22212, mock-

inoculated control, and an uninoculated control. Isolates P01-GFP and 22212 were resuspended

in phosphate-buffered saline (PBS) to a 1.0 optical density at 600 nm (OD600) and amended

with Silwet L-77 silicone surfactant (Momentive Performance Materials, Niskayuna, NY) at a

concentration of 0.02% (Mphande et al. 2024). Approximately 10 days after planting, inoculation

was performed by withdrawing 300 μL of inoculum using a 26 ½ gauge needle attached to a 1 ml

syringe (Becton, Dickinson & Co., New Jersey, US) and injecting it into the stem three times at a

45° angle, releasing the inoculum gently (Mphande et al. 2024). A new syringe was used for each

treatment, and the needle was sterilized with 70% ethanol for 1 minute and rinsed in deionized

water for 5 seconds after each plant inoculation. Each experiment was repeated once.

Sampling: For each experiment, plants were destructively sampled at zero, seven, fourteen,

twenty-one, and twenty-eight days post-treatment. Samples were collected from six locations on

each plant: stem 1 cm below the soil line, 1 cm above the soil line, 2 cm above the soil line, and

from the first petiole, the first leaf, and the shoot apex. Root tissue was only collected for

fluorescence-based image localization. To confirm *S. ureilytica* presence in every plant sample, a

direct PCR was conducted using *S. ureilytica*-specific primers (A79F-A79R) (Zhang et al. 2005).

Bacteria localization using imaging. Samples for imaging were taken from plants inoculated

with *S. ureilytica* isolate P01 (GFP), and from mock-inoculated controls. Plants were inoculated

as previously described, and samples were collected for imaging at the specified time points.

Transverse sections of the stem were taken from **t**he six tissue types described above, with an

additional sample collected from the root, approximately 2 cm below the root-stem junction.

Samples were sectioned by hand and mounted in water on a slide with a 0.17 mm coverslip

(Avantor, Radnor, PA, USA). To assess the spatial distribution of *S. ureilytica* within tissue cross

sections, samples were imaged using a Leica, M205 FCA fluorescence dissecting microscope

equipped with an EL6000 light source and a K5 camera, with LAS X software (Leica

Microsystems, Wetzlar, Germany). Exposure time and detector gain were adjusted to avoid

signal saturation. Cellular localization was characterized using an LSM880 upraight Axio

Examiner Z1 laser-scanning confocal microscope, with Zen image acquisition software (Zeiss,

Oberkocehn, Germany). GFP fluorescence was excited using a 23 mW ArgonRemote laser and

MBS 488 dichroic mirror (Zeiss, MBS-488) with an excitation peak at 488nm and an emissions

peak at 545.5nm. Laser power was set to 20% of maximum output, and detector gain was set to

450, except where explicitly noted in images specifically depicting autofluorescence of plant-

endogenous compounds (Supplementary Fig. 1).

To distinguish apoplastic versus symplastic compartmentalization, plants were grown and

inoculated with *S. ureilytica* isolate P01 (GFP) or mock-inoculated as described above. One

sample type (1 cm below the soil line) was selected because it had consistently shown a distinct

GFP signal across all time points. Samples were harvested 21-days post inoculation and stained

with Calcofluor White (Biotium, Fremont, CA, USA; 5mM in water), a fluorescent dye that

binds to β-glucans in cell walls, for visual cell segmentation. Samples were hand-sectioned as

described above, then incubated with the dye for 30 seconds, and rinsed with deionized water

immediately before imaging. Samples stained with Calcofluor White were simultaneously

imaged using two channels: the GFP channel (described above, but with laser power reduced to

19%), and a Calcofluor White-specific channel, using a 405-nm diode laser at 30% power, with

an excitation peak at 405 nm and emission peak at 467 nm. Detector gain was adjusted at

different magnifications (10x, 20x, and 63x) to optimize signal-to-noise ratios and avoid

saturation. Brightfield images were also collected for all samples.

Image processing for all confocal and dissecting-scope images was performed in Fiji (ImageJ v.

1.54; Schindelin et al. 2012). Image stacks were imported, and channels (GFP, Calcofluor White,

and brightfield) were merged using the 'Merge Channels' function for visualization and ROI

identification.

Quantification of *S. ureilytica*: Three plants for each of the four treatments were destructively

sampled at the same time points and sampling locations, with exception for root samples, which

were excluded from CFU quantification. Plant tissue samples (0.5 g) were surface-sterilized with

70% ethanol for 1 minute, followed by a 20-second rinse with sterile distilled water. The

sterilized tissue was macerated in 500 µl of sterile distilled water. A serial dilution series was

performed using the steps described by (Herigstad et al. 2001; Slack et al. 2021) resulting in five

10-fold serial dilutions. The droplet plating method was employed using LB agar plates

supplemented with tetracycline (20 µg/ml), (Mphande et al. 2025b). For each dilution, 10 drops

each with 10 µL were distributed across two plates (5 per dilution per plate, n=30 droplets per

plate). Plates were incubated at 28°C and checked for bacterial growth after 24 hours. The

dilution that gave 3-30 colonies per drop was used for counts (Herigstad et al. 2001). All colonies

in the 10 drops were counted (5 drops × 2 plates) at the countable dilution. Bacterial colony

forming units (CFU/ml) were calculated by multiplying the total number of colonies counted on

the plate by the dilution factor, then dividing the result by the volume of the plated sample.

Data analysis for bacterial colonization. Data were analyzed using a generalized linear mixed

model (PROC GLIMMIX, SAS Institute, 2013) with adaptive quadrature (colony forming units

per ml data) methods (Gbur et al., 2012). A log model was used to represent the data. The best fit

for the model was based on the lowest Akaike information criterion (AIC) value. All models

included Plant ID, Treatment, Isolate, Sample Location, Week, and their interaction as fixed

effects, and replication as a random effect. When a significant interaction effect was detected, the

SLICEDIFF function was used to compare treatments. The model Kenward–Roger method to

estimate denominator degrees of freedom (Kenward and Roger 1997). Mean separations were

calculated using Tukey's Honest Significant Difference [HSD] test at $\alpha = 0.05$. For this analysis,

if colony growth was not observed in any tissue of a plant, the plant was excluded from the data

analysis.

Results

Comparison of isolates: Isolations were obtained from all plants inoculated for both imaging

and bacterial density experiments from six plant sampling locations. *S. ureilytica* was

consistently isolated from the main stem (1 cm below, 1 cm above, and 2 cm above the

inoculation site), and it was also found, although less frequently, in the petiole, leaf base, and

apex (**Table 1**). Isolations were successful in recovering *S. ureilytica* even when inoculated

plants showed no symptoms. For example, while 22% of plants inoculated with isolate P01

(GFP-tagged) showed CYVD-like symptoms (yellowing, leaf scorching, and stunting), the

isolate was recovered from 91% of inoculated plants. Likewise, 41% of the plants inoculated

with the positive control (isolate 22212) showed symptoms, and the isolate was recovered from

100% of the inoculated plants. No *S. ureilytica* was recovered from isolations of the uninoculated

and/or mock-inoculated plants.

Bacterial localization using imaging. Transverse sections from the seven established locations

were destructively sampled from inoculated and control cucurbit plants (var. *Delicata*) each

week for 28 days. In inoculated plants, *S. ureilytica* localization was consistently observed in

phloem-associated cells across the main stem and in distal tissues (**Table 1**). Dissecting scope

images showed that *S. ureilytica* P01-GFP was concentrated in both the interior and exterior

regions of each vascular bundle, consistent with the bicollateral anatomy of phloem tissue in

cucurbits (**Fig. 2**). GFP signal was detectable to some degree in all sample locations across time

points. Notably, the signal at 1 cm below soil level and in the root was consistently stronger than

in distal shoot samples (**Fig. 2**). In these belowground sample locations the tagged *S. ureilytica*

inoculate appeared to colonize a larger proportion of the host cells at three and four weeks post-

inoculation relative to earlier timepoints (**Fig. 2**). Due to the destructive nature of sampling, the

dynamics of disease progression could not be tracked in individuals over time. Fluorescence in

control or mock-inoculated plants was only observed around the cell walls of vessel elements,

which have known autofluorescent properties (Donaldson 2020) (**Supplementary Fig. 1**). This

autofluorescence was clearly distinguishable from GFP signal based on its localization and faint

emission (**Supplementary Fig. 1**).

Dissecting scope imaging enabled visualization of *S. ureilytica* colonization in entire transverse

sections, whereas confocal microscopy resolved signal localization at the cellular level. Confocal

images taken at 10x magnification confirmed *S. ureilytica* localization within the vascular

bundles and provided additional context that the signal localized to a subset of cells, scattered

throughout phloem tissue, rather than colonizing in aggregated host cell clusters (**Fig. 3**). Images

taken at 10x, 20x, and 63x of the lower stem (1 cm below the soil line) and stained with

Calcofluor White to distinguish cell walls, revealed that the *S. ureilytica* GFP signal was

predominantly symplastic, with some signal putatively localized to apoplastic regions (**Fig. 4**).

Colonized cells were frequently irregular in shape and typically smaller than surrounding cells**.**

Brightfield images also showed that colonized cells along with neighboring uncolonized cells

have a darker, shadow appearance **(Fig. 4).**

Quantification of *S. ureilytica*. For the GFP P01 isolate, quantification results yielded plants in

which *S. ureilytica* was isolated from one or more, but not all, sample locations, as well as plants

where the pathogen was isolated from all sampling locations. Therefore, data analyses of

bacterial colonization were performed using two datasets. First, analyses were performed on

plants with *S. ureilytica* growth in at least one plant sampling location (**Fig 5**). A second analysis

was performed comparing only plants with growth from all plant locations (**Fig. 6**). In the first

analysis, results showed that at seven days post-inoculation (DPI), colonization by the GFP-

tagged *S. ureilytica* P01 isolate was successful but more localized to the inoculation site (**Fig. 5**).

There were high bacterial counts from the stem sections (1 cm below, 1 cm above, and 2 cm

above the inoculation point), but no significant differences in log CFU/mL (**Fig. 5**). At 14 DPI,

S. ureilytica was again isolated from stem sections, but the bacterium was also isolated from

petioles. (**Fig. 5**). At 21 and 28 DPI, there was a higher detection rate from distal tissue, with

stem sections at 1 and 2 cm above the inoculation point having higher log CFU/ml compared to

the petiole, leaf, or apex tissue (**Fig. 5**). No *S. ureilytica* growth was seen in the mock-inoculated

and uninoculated controls.

Analysis limited to plants with successful growth in all parts showed that *S. ureilytica* colony

counts were higher at the crown level than at other sampled locations (**Fig. 6**). At 21 DPI, *S.*

ureilytica colony counts were significantly higher (log CFU/ml) in the three stem sections

compared to all other plant tissues, with distal tissues having the lowest log CFU/ml (**Fig. 6**). At

28 DPI, bacterial distribution was not significantly different across sample locations, and the

mean log CFU/ml showed a more even bacterial distribution across the plant with no significant

differences among plant tissues (**Fig. 6**).

Discussion

The present study aimed to understand the *in planta* localization and temporal distribution of *S.*

ureilytica, the causal organism of CYVD. To achieve this, cucurbit plants (*C. pepo*) were

inoculated at the cotyledon stage using a GFP-tagged *S. ureilytica* isolate. Dissection and

confocal laser scanning microscopy revealed that *S. ureilytica* colonized specific cells at the

interior and exterior peripheries of the vascular bundles of cucurbits, suggesting that bacterial

colonization of *C. pepo* plants is restricted to phloem-associated tissues as previously

hypothesized (Bruton et al. 1998a). This localization differs from *E. tracheiphila*, which is found

in the xylem vessels of cucurbit plants, causing rapid water deprivation and wilt, in contrast to

the slow symptom development caused by CYVD (Bruton et al. 1998a; Rojas et al. 2015).

The presence of *S. ureilytica* was observed across time and location. We found strong evidence

of symplastic localization, with putative GFP signal also localized to the apoplast.

Quantitative analysis of *S. ureilytica* showed that this bacterium moves within the plant over

time. Results also suggest that the bacterium travels bidirectionally through the phloem sieve

elements from the main stem to the root, up to the crown, and eventually throughout the plant.

This is consistent with previous reports, including in vitro studies showing that Type I fimbriae

present in *S. ureilytica* impedes this pathogen's biofilm formation, thereby allowing a larger

population of bacteria to travel through the plant's phloem and colonize more tissue (Mphande et

al.,2025a). Results from bacterial quantification in six plant tissues indicated that the pathogen

was mainly found in stem tissue near the inoculation site for the first 14 days, then moved to

more distal locations by 21 and 28 DPI, with titers that are not significantly different between

sampling locations at 28 DPI. This might be explained by *S. ureilytica* fully colonizing phloem

cells as it moves through the plant, which can also lead to increased symptom development, as

higher populations of this bacterium are required to see CYVD symptoms (Mphande et al.

2025a).

Based on the results presented here, we hypothesize that *S. ureilytica* utilizes the sieve tubes of

the phloem for transport, but colonizes other cells within the phloem. These findings are

consistent with the leakage retrieval mechanism of phloem transport in which solutes in transport

sieve tubes can "leak" out of sieve tubes, either into the apoplast or, in some conditions, through

PD into companion cells (De Schepper et al., 2013).

Although we do not observe a GFP signal in sieve tube cells, bacterial presence in those cells

may be transient and at concentrations too low to be detected. Fluorescence in our images is

mainly symplastic, scattered throughout a subset of small to medium, dark, irregularly shaped

cells, within the phloem-associated regions of the vascular bundle. We hypothesize that *S.*

ureilytica could be colonizing one of two cell types: phloem parenchyma or companion cells.

Companion cells are typically small and have dense cytoplasm, which can make them appear

dark in brightfield images (Esau 1969). Companion cells also tend to be scattered throughout the

phloem region of the vascular bundle, similar to what we see in our dissecting scope images.

However, companion cells are always directly associated with a sieve cell due to shared lineage,

and it is not clear that our cells consistently show this phenotype. These cells may appear dark

and irregularly shaped as a result of the infection, making them difficult to identify based on

morphology alone. This localization differs from that of other known phloem colonizing

bacteria, such as *Candidatus Liberibacter, asiaticus* in citrus, which is restricted to the sieve tube

elements of the phloem (Ding et al. 2015; Tiwari 2025). Other examples include *Candidatus*

Arsenophonus phytopathogenicus and *Candidatus Phytoplasma asteris,* which are also found in

phloem sieve tube elements and are unculturable. This resistance to culturing may be explained

by the specialization required for the bacteria to occupy sieve tubes, an environment with an

extremely high nutrient concentration. This distinction may also explain why *S. ureilytica,* can be

isolated on artificial media when other phloem colonizers cannot. These results contrast with

previous studies that showed the presence of *S. ureilytica* only in phloem sieve tube elements

(Bruton et al. 1998b).The fact that *S. ureilytica* is localized on phloem-associated cells raises the

questions as to how both squash bug and cucumber beetles are able to deliver the bacterium into

these cells. It is well known that the squash bug is able to feed from phloem tissue (Neal

1993).On the other hand, cucumber beetles use a stercorarian transmission, where they acquire

the bacterium through feeding and deposit the infected frass into open chewing wounds when

feeding (Mphande et al. 2025b). This infection process could involve using the polymer trapping

loading system to deliver *S. ureilytica* from the leaves to the main stem which utilizes a

concentration gradient to move sugars cell-to-cell via plasmodesmata into the phloem transport

stream, using intermediary cells (Reidel et al. 2009; Turgeon 2017; Turgeon and Hepler 1989).

Furthermore, previous research has shown that *Serratia* species that colonize insects proliferate

at higher rates within the host when fed a high-sucrose diet in fruit flies (*Drosophila*

melanogaster)(Darby et al. 2024). Given the high sucrose content in both companion cells and

phloem parenchyma and in the apoplast compared to other plant tissues, this could drive the

pathogen's localization to these cells. Since sucrose is an important carbon source for nitrogen

assimilation in plants (Huang et al. 2022), hijacking sucrose may explain CYVD symptoms such

as stunting and yellowing of older leaves (Besler 2014; Bruton et al. 1998a; Pair et al. 2004).

In both the imaging and quantification experiments, we consistently observe greater bacterial

abundance in the lower tissues of the main stem and root. This is consistent with the dynamics of

bulk flow in the phloem. Bulk flow, or the movement of sap through the phloem, is powered by

the pressure gradient, driven by osmotic forces due to varying concentrations of solute in the sap

(high concentrations in mature leaves where photosynthesis occurs and low concentrations in

storage tissues like roots) (Fellows and Geiger 1974). The direction of flow in any given sieve

tube changes throughout development, with architectural modifications such as the initiation and

maturation of new leaves and the growth of the root system, but during vegetative growth stages

apical-to-basal flow dominates. Furthermore, it is established that some phloem-occupying

pathogens can modify metabolic activity in colonized cells, thereby altering transport direction

and increasing nutrient delivery to infected organs (Malinowski et al., 2024). These dynamics are

consistent with the localization of *S. ureilytica* observed in our plants, as distal tissues are more

likely to be occupied at higher densities at later time points, with the majority of bacterial growth

occurring in the main stem and root. This work provides novel evidence of *S. ureilytica*

localization in cucurbit plants and advances our knowledge of this emergent disease in the US.

Building on these findings, microscopy methods, including histology and immunolocalization

can be used to confidently establish cell-type localization within the phloem. Given *S.*

ureilytica's preference to colonize phloem companion cells or phloem parenchyma cells and

move systemically throughout the plant, further research should prioritize elucidating the

molecular mechanisms driving this localization. Future studies could investigate environmental

cues, such as sucrose gradients, to determine if they act as chemoattractants for the bacterium

within the phloem. Moreover, further investigation into the interaction between *S. ureilytica* and

cucurbit plasmodesmata could clarify if the bacterium actively manipulates symplastic transport

pathways or relies on passive movement.

Acknowledgements

We thank Elizabeth Little and Nicole Gerardo for sharing the GFP *S. ureilytica* isolate used in

this study.

Funding

This work was made possible by the Schmittau-Novak Integrative Plant Science Small Grants

Program. The Federal Capacity Hatch program, project award number 7005329, from the

U.S. Department of Agriculture's National Institute of Food and Agriculture. Imaging data were

acquired through the Cornell Institute of Biotechnology's Imaging Facility, with NIH

S10OD018516 funding for the shared Zeiss LSM880 confocal/multiphoton microscope.

References

Agustí, J., and Blázquez, M. A. 2020. Plant vascular development: mechanisms and
environmental regulation. Cellular and Molecular Life Sciences 77:3711–3728.
https://doi.org/10.1007/s00018-020-03496-w.

Van Bel, A. J. E. 2003. The phloem, a miracle of ingenuity. Plant Cell Environ. 26:125–149.
https://doi.org/10.1046/j.1365-3040.2003.00963.x.

Bendix, C., and Lewis, J. D. 2018. The enemy within: Phloem-limited pathogens. Mol. Plant Pathol. 19:238–254. https://doi.org/10.1111/mpp.12526.

Bertani, G. 1951. Studies on lysogenesis I: the mode of phage liberation by lysogenic *Escherichia coli*. J Bacterio 62:293–300.

Besler, K. R. 2014. Epidemiology and management of cucurbit yellow vine disease, and characterization of the causal agent *Serratia marcescens*. Master's thesis. University of Georgia. .

Besler, K. R., and Little, E. L. 2017. Diversity of *Serratia marcescens* strains associated with cucurbit yellow vine disease in Georgia. Plant Dis. 101:129–136. https://doi.org/10.1094/PDIS-05-16-0618-RE.

Bextine, B., Wayadande, A., Bruton, B. D., Pair, S. D., Mitchell, F., and Fletcher, J. 2001. Effect of insect exclusion on the incidence of yellow vine disease and of the associated bacterium in squash. Plant Dis. 85:875–878. https://doi.org/10.1094/PDIS.2001.85.8.875.

Bonjour, E. L., and Fargo, W. S. 1989. Host effects on the survival and development of *Anasa tristis* (Heteroptera: Coreidae). Environ. Entomol 18:1083–1085.

Braun, D. M. 2022. Phloem Loading and Unloading of Sucrose: What a Long, Strange Trip from Source to Sink. Annual Review of Plant Biology Downloaded from www.annualreviews.org. Guest (guest 22:39. https://doi.org/10.1146/annurev-arplant-070721.

Bruton, B. D. , Fletcher, J., Pair, S. D., Shaw, M., and Sittertz-Bhatkar, H. 1998a. Association of a phloem-limited bacterium with yellow vine disease in cucurbits. Plant Dis. 82:512.

Bruton, B. D. , Pair, S. D. , Popham, T. W. , and Cartwright, B. O. 1998b. Occurrence of yellow vine, a new disease of squash and pumpkin, in relation to insect pests, mulches, and soil fumigation. Plant Sci. 47.

Bruton, B. D., Mitchell, F., Fletcher, J., Pair, S. D., Wayadande, A., Melcher, U., Brady, J., Bextine, B., and Popham, T. W. 2003. *Serratia marcescens*, a phloem-colonizing, squash bug-transmitted bacterium: causal agent of cucurbit yellow vine disease. Plant Dis. 937–944.

Darby, A. M., Okoro, D. O., Aredas, S., Frank, A. M., Pearson, W. H., Dionne, M. S., and Lazzaro, B. P. 2024. High sugar diets can increase susceptibility to bacterial infection in *Drosophila melanogaster*. PLoS Pathog. 20. https://doi.org/10.1371/journal.ppat.1012447.

Ding, F., Duan, Y., Paul, C., Brlansky, R. H., and Hartung, J. S. 2015. Localization and Distribution of *"Candidatus Liberibacter asiaticus"* in citrus and periwinkle by direct tissue blot immuno assay with an anti-ompa polyclonal antibody. PLoS One 10. https://doi.org/10.1371/journal.pone.0123939.

Donaldson, L. 2020. Autofluorescence in plants. Molecules 25. https://doi.org/10.3390/molecules25102393.

Esau, K. 1969. *The Phloem*. 2nd ed. ed. Gebr. Borntraeger.

Fellows, R. J., and Geiger, D. R. 1974. Structural and Physiological Changes in Sugar Beet
Leaves during Sink to Source Conversion. Plant Physiol. 54:877–885.
https://doi.org/10.1104/pp.54.6.877.

Gottwald, T. R. 2010. Current epidemiological understanding of citrus huanglongbing. Annu.
Rev. Phytopathol. 48:119–139. https://doi.org/10.1146/annurev-phyto-073009-114418.

Haider, M. W., Sharma, A., Majumdar, A., Fayaz, F., Bromand, F., Rani, U., Singh, V. K.,
Saharan, M. S., Tiwari, R. K., Lal, M. K., and Kumar, R. 2024. Unveiling the phloem: a
battleground for plant pathogens. Phytopathology Research 6. https://doi.org/10.1186/s42483-
024-00286-1.

Herigstad, B., Hamilton, M., and Heersink, J. 2001. How to optimize the drop plate method for
enumerating bacteria. J. Microbiol. Methods 44:121–129.

Huang, X. J., Jian, S. F., Chen, D. L., Zhong, C., and Miao, J. H. 2022. Concentration-dependent
dual effects of exogenous sucrose on nitrogen metabolism in Andrographis paniculata. Sci. Rep.
12. https://doi.org/10.1038/s41598-022-08971-x.

Luo, G. 2006. Phytopathogenicity, virulence factors, and in planta movement of *Serratia*
marcescens. Doctoral Dissertation. Oklahoma State University 1–139.

Mahillon, M., Debonneville, C., Groux, R., Roquis, D., Brodard, J., Faoro, F., Foissac, X.,
Schumpp, O., and Dittmer, J. 2025. From insect endosymbiont to phloem colonizer: comparative
genomics unveils the lifestyle transition of phytopathogenic Arsenophonus strains . mSystems
10. https://doi.org/10.1128/msystems.01496-24.

Mendiola, S. Y., Stoy, K. S., DiSalvo, S., Wynn, C. L., Civitello, D. J., and Gerardo, N. M. 2022.
Competitive Exclusion of Phytopathogenic Serratia marcescens from Squash Bug Vectors by the
Gut Endosymbiont Caballeronia. Appl. Environ. Microbiol. 88.
https://doi.org/10.1128/AEM.01550-21.

Mphande, K., Gleason, M. L., and Beattie, G. A. 2024. A bioassay that yields quantifiable
symptoms of cucurbit yellow vine disease caused by *Serratia marcescens*. Plant Dis.
https://doi.org/10.1094/pdis-07-24-1401-re.

Mphande, K., LaSarre, B., Gleason, M. L., and Beattie, G. A. 2025a. Type I Fimbriae Promote
the Virulence of Serratia ureilytica Strains That Cause Cucurbit Yellow Vine Disease While
Impeding Biofilm Formation In Vitro . Phytopathology. https://doi.org/10.1094/phyto-05-25-
0165-r.

Mphande, K., LaSarre, B., Paulsen, A. A., Hartung, R., Badilla-Arias, S., Gleason, M. L., and
Beattie, G. A. 2025b. Bacteria that Cause Cucurbit Yellow Vine Disease Fall Within the *Serratia*
ureilytica Species of the *S. marcescens* Complex and Can Be Vectored by Cucumber Beetles.
Phytopathology 115:606–617. https://doi.org/10.1094/PHYTO-10-24-0321-R.

Neal, J. J. 1993. Xylem transport interruption by *Anasa tristis* feeding causes *Cucurbita pepo* to
wilt. Entomol. Exp. Appl. 69:195–200. https://doi.org/10.1111/j.1570-7458.1993.tb01741.x.

Pair, S. D., Bruton, B. D., Mitchell, F., Fletcher, J., Wayadande, A., and Melcher, A. U. 2004.
Overwintering squash bugs harbor and transmit the causal agent of cucurbit yellow vine disease.
J. Econ. Entomol. 97:74–78.

Perilla-Henao, L. M., and Casteel, C. L. 2016. Vector-borne bacterial plant pathogens:
Interactions with hemipteran insects and plants. Front. Plant Sci. 7.
https://doi.org/10.3389/fpls.2016.01163.

Rascoe, J., Berg, M., Melcher, U., Mitchell, F. L., Bruton, B. D., Pair, S. D., and Fletcher, J.
2003. Identification, Phylogenetic Analysis, and Biological Characterization of *Serratia*
marcescens Strains Causing Cucurbit Yellow Vine Disease. Phytopathology 93:1233–1239.

Reidel, E. J., Rennie, E. A., Amiard, V., Cheng, L., and Turgeon, R. 2009. Phloem loading
strategies in three plant species that transport sugar alcohols. Plant Physiol. 149:1601–1608.
https://doi.org/10.1104/pp.108.134791.

Rodriguez-Herrera, K. D., Ma, X., Swingle, B., Pethybridge, S. J., Gonzalez-Giron, J. L.,
Herrmann, T. Q., Damann, K., and Smart, C. D. 2023. First report of cucurbit yellow vine
disease caused by *Serratia marcescens* on cucurbits in New York. Plant Dis. 107:3276.
https://doi.org/10.1094/PDIS-06-23-1051-PDN.

Rojas, E. S., Batzer, J. C., Beattie, G. A., Fleischer, S. J., Shapiro, L. R., Williams, M. A., Bessin,
R., Bruton, B. D., Boucher, T. J., Jesse, L. C. H., and Gleason, M. L. 2015. Bacterial Wilt of
Cucurbits: Resurrecting a Classic Pathosystem. Plant Dis. 99:564–574.
https://doi.org/10.1094/PDIS-10-14-1068-FE.

De Schepper, V., De Swaef, T., Bauweraerts, I., and Steppe, K. 2013. Phloem transport: A review
of mechanisms and controls. J. Exp. Bot. 64:4839–4850. https://doi.org/10.1093/jxb/ert302.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J. Y., White, D. J., Hartenstein, V., Eliceiri, K.,
Tomancak, P., and Cardona, A. 2012. Fiji: An open-source platform for biological-image
analysis. Nat. Methods 9:676–682. https://doi.org/10.1038/nmeth.2019.

Slack, S. M., Walters, K. J., Outwater, C. A., and Sundin, G. W. 2021. Effect of kasugamycin,
oxytetracycline, and streptomycin on in-orchard population dynamics of *Erwinia amylovora* on
apple flower stigmas. Plant Dis. 105. https://doi.org/10.1094/PDIS-07-20-1469-RE.

Taylor, T. N., Taylor, E. L., and Krings, M. 2009. Introduction to Vascular Plant Morphology and
Anatomy. In *Biology and Evolution of Fossil Plants*, eds. Thomas N Taylor, Edith L Taylor, and
Michael Krings. Elsevier, pp. 201–222. https://doi.org/10.1016/B978-0-12-373972-8.00007-3.

Thorpe, M., Minchin, P., Gould, N., and McQueen, J. 2005. The Stem Apoplast. In *Vascular*
Transport in Plants, Elsevier, pp. 201–220. https://doi.org/10.1016/B978-012088457-5/50012-5.

Tiwari, R. 2025. Liberibacter Turning Citrus into Its Hideout. Molecular Plant-Microbe
Interactions 38:4–5. https://doi.org/10.1094/MPMI-01-25-0007-CM.

Turgeon, R. 2017. Phloem biology of the Cucurbitaceae. In *Genetics and Genomics of*
Cucurbitaceae, Plant Genetics and Genomics: Crops and Models, eds. Rebecca Grumet, Nurit
Katzir, and Jordi Garcia-Mas. Cham: Springer International Publishing.
https://doi.org/10.1007/978-3-319-49332-9.

Turgeon, R., and Hepler, P. K. 1989. Symplastic continuity between mesophyll and companion
cells in minor veins of mature Cucurbita pepo L. leaves. Springer-Verlag 179:24–31.

Zhang, C., Yu, X., Ayre, B. G., and Turgeon, R. 2012. The origin and composition of cucurbit
"Phloem" exudate. Plant Physiol. 158:1873–1882. https://doi.org/10.1104/pp.112.194431.

Zhang, Q., Melcher, U., Zhou, L., Najar, F. Z., Roe, B. A., and Fletcher, J. 2005. Genomic
Comparison of Plant Pathogenic and Nonpathogenic *Serratia marcescens* Strains by suppressive
subtractive hybridization. Appl. Environ. Microbiol. 71:7716–7723.
https://doi.org/10.1128/AEM.71.12.7716-7723.2005.

Fig 1. Description of methodology for bacteria quantification using the dilution plating method.

Created in https://BioRender.com

564   S

572   Fig 2. Dissecting scope images showing the localization of the green fluorescence protein (GFP)-

573   tagged *Serratia ureilytica* in *Cucurbita pepo* plants. In six tissue types across 4 weeks. Plants

574   were destructively sampled. Scale bar 1000 μm.

Fig. 3 Confocal images show the localization of GFP-tagged *Serratia ureilytica* isolates across six tissue types. Representative images were selected for each tissue to illustrate bacterial localization within the anatomical context of the vascular bundle. Images were acquired at 10× magnification and are shown as merged brightfield and GFP channels. Brightness and contrast were adjusted on a per-image basis for visualization to facilitate detection of low-intensity GFP signals, where present; control images were displayed with intensity scaling matched to the most dimly fluorescent GFP-positive sample. Some images were cropped to highlight the anatomical region of interest. Post-acquisition adjustments were limited to display scaling and were not used for quantitative analysis.

*white arrow indicates xylem

Fig. 4 Confocal images of the same sample (1 cm below soil line) of *Cucurbita pepo* var.

Delicata, acquired at 10×, 20×, and 63× magnifications to demonstrate symplastic and

apoplastic (63X (B)) localization of GFP-tagged *Serratia ureilytica* relative to host cell walls.

Tissue was collected 2 cm below the soil surface at 4 weeks post-inoculation and stained with

calcofluor white immediately prior to imaging. From left to right, panels show brightfield,

calcofluor white-stained cell walls (blue), GFP fluorescence (green), and merged composite

images of all three channels.

*white arrow in 10x indicates xylem

614   Supplementary Fig. 1: autofluorescence in control images using confocal microscopy at 10x. Fig.

615   A. Autofluorescence in the chloroplast at 20X. Fig. B-D Autofluorescence in xylem tissue. GFP

616   Gain A:985 B: 559. C: 718. D: 705

632 Fig 5. Log CFU/ml of *Serratia ureilytica* (CFU/ml) isolated from *Cucurbita pepo* plants

633 inoculated at two weeks old and sampled weekly for four weeks. Each plant was sampled in one

634 of six locations shown on the y-axis. Bars with the same letter are not significantly different

635 from each other ($P > 0.05$) according to Tukey's honestly significant difference. Error bars

636 indicate standard error.

Fig. 6 Log CFU/ml of *Serratia ureilytica* isolated from *Cucurbita pepo* plants inoculated at two

weeks old, and sampled weekly for four weeks, including only those plants in which *S. ureilytica*

colonies were detected in all six plant parts. Each plant was sampled in one of six locations

shown on the y-axis. Bars with the same letter are not significantly different from each other (P >

0.05) according to Tukey's honestly significant difference. Error bars indicate standard error.

650   Table 1. *Serratia ureilytica* isolations from inoculated plants with isolate P01 from one to four weeks post-inoculation in different

651   plant sampling locations of *Cucurbita pepo* plants (var. Delicata)

Plant Sampling location	Isolations Confocal trials				Isolation bacteria counting trials				Total Isolations (n=45 plants)
	7 DPI (n=4)	14 DPI (n=7)	21 DPI (n=5)	28 DPI (n=6)	7 DPI (n=5)	14 DPI (n=6)	21 DPI (n=6)	28 DPI (n=6)	
1 cm below	3	6	4	6	5	5	4	6	39
1 cm above	3	6	4	6	5	6	5	6	41
2 cm above	4	6	3	6	5	6	5	6	41
Petiole	2	3	1	1	3	3	3	3	19
leaf base	1	1	1	1	0	0	2	2	8
apex	1	1	1	1	0	0	2	3	9

655