TSUMUGI: a platform for phenotype-driven gene network identification from comprehensive knockout mouse phenotyping data

Akihiro Kuno[1,2,3*], Kinari Matsumoto[4], Taito Taki[4], Satoru Takahashi[2,3] and Seiya Mizuno[3]

1. Tsukuba Institute for Advanced Research (TIAR), University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki, 305-8577, Japan.
2. Department of Anatomy and Embryology, Institute of Medicine, University of Tsukuba, Tsukuba, Ibaraki, 305-8575, Japan.
3. Laboratory Animal Resource Center, Transborder Medical Research Center, Institute of Medicine, University of Tsukuba, Tsukuba, Ibaraki, 305-8575, Japan.
4. Ph.D. Program in Human Biology, Graduate School of Comprehensive Human Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki, 305-8575, Japan.

*Corresponding author. Akihiro Kuno, Tsukuba Institute for Advanced Research (TIAR), University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki, 305-8577, Japan. E-mail: akuno@gwe.md.tsukuba.ac.jp

Abstract

Summary: Deciphering complex organismal phenotypes requires elucidation of coordinated functions among multiple genes, yet this remains a fundamental challenge in functional genetics. The International Mouse Phenotyping Consortium (IMPC) has recently established a comprehensive phenotypic atlas based on systematic single-gene knockout mouse lines, providing an unprecedented resource for gene-phenotype associations in mammals. However, extracting phenotype-associated multiple gene relationships remains challenging. Here, we present TSUMUGI, a platform that identifies a phenotype-driven gene network by leveraging gene-phenotype associations from the IMPC. TSUMUGI enables users to initiate analyses from a user-specified phenotype or gene list and interactively explore a gene network. In addition, TSUMUGI highlights human disease-associated genes, supports flexible network filtering, and allows seamless export of a gene network for downstream analyses. By linking shared phenotypic signatures to putative functional gene modules, TSUMUGI provides a framework for systematic interpretation and hypothesis generation of complex organismal phenotypes.

Availability and implementation: The web application is available online at https://larc-tsukuba.github.io/tsumugi/. The command-line tools are distributed via PyPI

35   (https://pypi.org/project/TSUMUGI) and Bioconda

36   (https://bioconda.github.io/recipes/tsumugi/README.html). Source code and documentation

37   are hosted at GitHub (https://github.com/akikuno/TSUMUGI-dev) and archived on Zenodo

38   (https://doi.org/10.5281/zenodo.18464478) under the MIT license.

1. Introduction

Complex organismal phenotypes are often driven by coordinated interactions among

multiple genes (Costanzo et al., 2019). Although multiple gene associations have been

extensively investigated using protein-protein interactions (PPI), biological pathways, and

gene co-expression networks, these approaches primarily focus on molecular and

intracellular relationships and may not fully reflect gene interactions at the organismal level

(Allayee et al., 2023). Similarly, genome-wide association studies and polygenic risk score

approaches identify genetic risk factors for complex traits but do not explicitly capture

coordinated gene interactions that give rise to organismal phenotypes (Lewis and Vassos.,

2020).

The International Mouse Phenotyping Consortium (IMPC) has generated comprehensive

phenotypic profiles for more than 9,000 single-gene knockout (KO) mouse lines, providing a

large-scale standardized resource for systematic gene-phenotype analyses (Dickinson et al.,

2016; Wilson et al., 2025). This dataset enables direct comparison of organism-wide

phenotypic consequences resulting from individual gene disruptions. However, because the

IMPC dataset is fundamentally organized around single-gene knockout lines, extracting

phenotypic relationships across multiple genes remains challenging.

Here, we present TSUMUGI (Trait-driven Surveillance for Mutation-based Gene module

Identification), a platform designed to identify a phenotype-driven gene network from the

IMPC dataset. TSUMUGI is based on the working hypothesis that genes whose knockout

mice exhibit similar phenotypes participate in shared biological systems, and accordingly

quantifies phenotypic similarity between KO mouse lines. By converting gene-phenotype

associations into a phenotype-driven gene network, TSUMUGI enables systematic

identification of gene associations underlying organism-level phenotypes.

TSUMUGI supports flexible analysis workflows initiated from a user-defined phenotype or a

gene list. Users can interactively explore a phenotype-driven gene network, apply

customizable filtering criteria, highlight human disease-associated genes, and perform

centrality analyses. The extracted gene network can be readily exported for downstream

analyses and integration with external tools.

TSUMUGI is available as both a web application and a command-line interface. The web
application (https://larc-tsukuba.github.io/tsumugi) provides an intuitive environment for
exploratory analysis, whereas the command-line interface, distributed via PyPI and
Bioconda, supports reproducible, large-scale analyses with advanced filtering options.

2. Methods

2.1 Data Sources

Gene-phenotype association dataset derived from single-gene knockout mouse lines was
obtained from the IMPC statistical summary table
(https://ftp.ebi.ac.uk/pub/databases/impc/all-data-releases/release-23.0/results/, statistical-
results-ALL.csv.gz, release: 23.0). From this dataset, the MGI marker symbol, Mammalian
Phenotype Ontology (MP) term, effect size, P-value, sex-specific P-value, zygosity, pipeline
name, and procedure name were extracted for each gene-phenotype record.

Life-stage annotations were assigned to each gene-phenotype record as follows. An
"Embryo" label was assigned when the procedure included embryonic screening (E9.5,
E10.5, E12.5, Embryo LacZ, E14.5, E14.5-E15.5, or E18.5). An "Interval" (17 weeks to 48
weeks) or "Late" (49 weeks or later) label was assigned when the pipeline name contained
"interval" or "late", respectively; otherwise, an "Early" label was assigned.

Mouse orthologues of human disease-associated genes were obtained from the IMPC
Disease Models Portal (https://diseasemodels.research.its.qmul.ac.uk/,
impc_phenodigm.csv, release: 2025-05-15) (Cacheiro et al., 2024). In the downloaded table,
the description column contains mouse allele symbols, zygosity, and life-stage information.
This field was normalized to match the phenotype dataset and appended accordingly.

MP annotation files were downloaded from OBO Foundry
(https://obofoundry.org/ontology/mp.html, mp.obo, release: 2025-08-27) (Jackson et al.,
2021) and used to calculate phenotype similarity.

2.2 Extraction of abnormal phenotypes

94 In the IMPC statistical summary table, an MP term is assigned to a gene-phenotype record
95 when either the P-value or the sex-specific P-value is below 0.00001. We therefore extracted
96 records with an assigned MP term as abnormal phenotypes for downstream analyses. Sex
97 specificity was defined as "Female," "Male," or "None" according to the significance of sex-
98 specific P-values.

2.3 Phenotype similarity metrics

Phenotypic similarity between KO mouse lines was calculated based on the Phenodigm algorithm (Smedley et al., 2013).

Term similarity for each pair of MP terms was defined as the geometric mean of the Resnik score and the Jaccard index. Information content (IC) was calculated from the number of descendant terms and used to compute the Resnik score. Terms with IC below the 5th percentile were considered overly general, and their Resnik scores were set to zero. The Jaccard index was calculated as the overlap ratio between the parent term sets of the MP terms.

A weighted similarity matrix was then constructed for all pairwise combinations of MP terms associated with two KO mouse lines by multiplying the term similarity score by a metadata concordance score. The metadata concordance score was defined as 1.0 when genotype, sex, and life stage were fully matched, and as 0.75, 0.5, or 0.25 depending on the number of mismatched metadata attributes.

Finally, Phenodigm-based scaling was applied to the weighted similarity matrix, yielding a phenotype similarity score between 0 and 100 for each gene pair.

2.4 Network construction and visualization in the web application

Gene pairs sharing at least three abnormal phenotypes were extracted, with each gene represented as a node in an undirected graph. The network was visualized using Cytoscape.js (version 3.30.4) (Franz et al., 2015).

3. Results

3.1 Web application

TSUMUGI provides three types of entry points: Phenotype, Gene, and Gene List, through which users can submit a MP term or gene symbols of interest (Fig. 1A). Upon submission, a dashboard comprising a network view with control panels is displayed (Fig. 1B).

On the top page, when the Phenotype entry is selected, TSUMUGI identifies genes whose KO mice exhibit the user-specified phenotype and groups them based on phenotype similarity. In contrast, the Gene entry identifies genes whose KO phenotypes resemble those of the user-specified gene. The Gene List entry groups genes within the submitted gene list (Fig. S1).

In the network view, nodes represent genes and edges represent phenotype similarity. Node
colour intensity indicates phenotype severity, while edge thickness reflects the phenotype
similarity score. The subnetworks are displayed as modules corresponding to connected
components, where genes within a module share high phenotypic similarity, while similarity
between modules is low. By selecting a module, users can inspect a list of phenotypes
enriched within that module. This feature allows users to assess which phenotypes each
module contributes to and to iteratively narrow down modules of interest (Fig. S2).

The left and right control panels provide interactive filtering based on similarity scores,
severity, and metadata (zygosity, sex, and life stage), as well as highlighting functions based
on human disease-association, phenotypes or genes, and network centrality. All interactions
are immediately reflected on the client side.

The gene network can be exported as static images (PNG, JPG) and vector graphics (SVG).
In addition, module-wise gene lists and associated phenotype information can be exported in
CSV and GraphML formats, enabling downstream analyses using external tools.

3.2 Command-line interface

In addition to the web application, TSUMUGI provides a command-line interface. This allows
researchers to process IMPC phenotype data in a local environment and construct a custom
network. Compared with the web tool, the command-line interface enables more flexible
extraction of a phenotype similarity network based on phenotypes, genes, and metadata.

Using the command-line tool, users can extract or exclude specific gene pairs based on
phenotype information and metadata (Table 1). For example, the 'tsumugi mp' command
enables extraction or exclusion of gene pairs associated with specified MP terms. This
functionality is useful, for instance, when extracting the gene network that is associated with
diabetes-related phenotypes but not with reproductive system phenotypes.

In particular, the 'tsumugi mp --exclude' command allows exclusion of gene pairs that are not
associated with a specified MP term and its descendant terms. Conventionally, the absence
of phenotype annotations for a gene made it unclear whether the phenotype was
unmeasured or not significant. By leveraging IMPC-provided information indicating that a
phenotype was measured but not significant, TSUMUGI enables exclusion of non-significant
phenotypes from the analysis.

Furthermore, as in the web application, the command-line tool supports export in GraphML
format, facilitating straightforward integration of TSUMUGI outputs into other analysis
pipelines and workflows.

4. Discussion

TSUMUGI enables the identification of putative gene interactions associated with organism-
level phenotypes. Through flexible filtering options, users can extract gene modules tailored
to specific research questions. For example, by restricting the analysis to female- or male-
specific phenotypes, users can identify sex-dependent gene modules underlying a
phenotype of interest, thereby facilitating the exploration of sexually dimorphic gene
interactions. Similarly, by focusing on phenotypes observed at late life stages (49 weeks or
later), TSUMUGI allows the identification of gene modules potentially involved in aging-
related traits. Together, these features position TSUMUGI as a hypothesis-generating
framework for linking shared phenotypic signatures to coordinated gene functions.

A representative study related to TSUMUGI integrated gene and phenotype networks based
on pre-defined interactions such as PPI and OMIM to prioritize candidate gene-phenotype
associations (Li and Patra., 2010). In contrast, TSUMUGI constructs a gene network directly
from large-scale KO mouse phenotype data without relying on pre-defined interaction maps.
Thus, TSUMUGI can uncover functional gene modules that may remain undetected by
knowledge-based interaction networks.

From a future perspective, TSUMUGI will incorporate additional network layers, including
PPI and biological pathways, to facilitate mechanistic context for interpreting phenotype-
associated gene modules. Furthermore, the inclusion of gene expression profiles will enable
the investigation of tissue specificity and inter-organ communication contributing to
organism-level traits. Temporal expression dynamics and sex-dependent gene expression
will further allow the characterization of life-stage- and sex-specific regulatory programs that
drive phenotypic diversity. In addition, leveraging genome-wide association studies and
polygenic risk scores may provide complementary insights into the genetic architecture of
human diseases. Collectively, these enhancements will expand the scope of TSUMUGI and
enhance its utility for dissecting complex traits.

5. Conclusion

189 Here, we present TSUMUGI, a platform for identifying a phenotype-driven gene network
190 using comprehensive KO mouse phenotype data from the IMPC. TSUMUGI enables intuitive

exploration of gene relationships associated with organism-level traits through an interactive web interface, alongside flexible and customizable filtering via a command-line interface. By converting gene-phenotype associations into a network representation, TSUMUGI facilitates the discovery of gene interactions underlying organismal phenotypes and human diseases.

Acknowledgements

We thank Dr. Atsushi Yoshiki, Dr. Shinya Ayabe, Dr. Hiroshi Masuya and Dr. Tatsuya Kushida for their valuable insights and discussions regarding this study. We also thank Dr. Haruka Kuno for her advice on designing the web interface.

Author contributions

Akihiro Kuno (Conceptualization [lead], Data curation [lead], Formal analysis [lead], Investigation [lead], Methodology [lead], Software [lead], Visualization [lead], Funding acquisition [lead], Writing—original draft [lead], Writing—review & editing [lead]), Kinari Matsumoto (Conceptualization [supporting], Data curation [supporting], Software [supporting], Writing—review & editing [equal]), Taito Taki (Conceptualization [supporting], Software [supporting], Writing—review & editing [equal]), Satoru Takahashi (Conceptualization [supporting], Funding acquisition [supporting], Writing—review & editing [equal]), and Seiya Mizuno (Conceptualization [supporting], Funding acquisition [lead], Writing—review & editing [equal])

Funding

This work was supported by Research Support Project for Life Science and Drug Discovery (Basis for Supporting Innovative Drug Discovery and Life Science Research) from AMED under grant number JP26ama121047 (to A.K., S.T., and S.M.), JSPS KAKENHI Grant-in-Aid for Scientific Research (A) under grant number 25H00959 (to A.K. and S.M.), Grant-in-Aid for Young Investigators under grant number 24K18045 (to A.K.), Uehara Memorial Foundation (to A.K.) and JST FOREST Program under grant number JPMJFR221H (to S.M.).

References

Allayee H, Farber CR, Seldin MM et al. Systems genetics approaches for understanding complex traits with relevance for human disease. eLife 2023;12:e91004. https://doi.org/10.7554/eLife.91004

Cacheiro P, Pava D, Parkinson H et al. Computational identification of disease models
through cross-species phenotype comparison. Dis Model Mech 2024;17:dmm050604.
https://doi.org/10.1242/dmm.050604

Costanzo M, Kuzmin E, Van Leeuwen J et al. Global genetic networks and the genotype-to-
phenotype relationship. Cell 2019;177:85-100. https://doi.org/10.1016/j.cell.2019.01.033

Dickinson ME, Flenniken AM, Ji X et al. High-throughput discovery of novel developmental
phenotypes. Nature 2016;537:508-514. https://doi.org/10.1038/nature19356

Franz M, Lopes CT, Huck G et al. Cytoscape.js: a graph theory library for visualisation and
analysis. Bioinformatics 2016;32:309-311. https://doi.org/10.1093/bioinformatics/btv557

Jackson R, Matentzoglu N, Overton JA et al. OBO Foundry in 2021: operationalizing open
data principles to evaluate ontologies. Database (Oxford) 2021;2021:baab069.
https://doi.org/10.1093/database/baab069

Lewis CM, Vassos E. Polygenic risk scores: from research tools to clinical instruments.
Genome Med 2020;12:44. https://doi.org/10.1186/s13073-020-00742-5

Li Y, Patra JC. Genome-wide inferring gene-phenotype relationship by walking on the
heterogeneous network. Bioinformatics 2010;26:1219-1224.
https://doi.org/10.1093/bioinformatics/btq108

Smedley D, Oellrich A, Kohler S et al. PhenoDigm: analyzing curated annotations to
associate animal models with human diseases. Database (Oxford) 2013;2013:bat025.
https://doi.org/10.1093/database/bat025

Wilson R, Bülbül Ataç T, Cheng TK et al. International Mouse Phenotyping Consortium
Portal: facilitating investigation of gene function and providing insights into human disease.
Nucleic Acids Res 2026;54:D1133-D1142. https://doi.org/10.1093/nar/gkaf1148

Figure 1. TSUMUGI web interface. (A) Top page. TSUMUGI accepts a phenotype term,

gene symbol, or gene list as input. (B) Network page. The central panel displays the

phenotype-driven gene network. The left panel provides network filtering options, the right

panel provides highlighting and visualization options, and the bottom panel provides export

functions.

Utility	Description
run	Build the TSUMUGI network locally from the IMPC dataset
mp --include/--exclude	Retain or remove gene pairs that include or do not exhibit the specified MP term and its descendant terms
zygosity --keep/--drop	Retain or remove gene pairs associated with specified zygosity types (Homo/Hetero/Hemi)
sex --keep/--drop	Retain or remove gene pairs associated with specified sex categories (Male/Female/None)
life-stage --keep/--drop	Retain or remove gene pairs associated with specified life stages (Embryo/Early/Interval/Late)
count --min/--max	Filter gene pairs or genes by the number of associated phenotypes
score --min/--max	Filter gene pairs by phenotype similarity score
genes --keep/--drop	Retain or remove specified genes
build-graphml	Export the network as GraphML format
build-webapp	Export the network as TSUMUGI web application assets

252   **Table 1.** Summary of supported operations available in the TSUMUGI suite