__Trace gas oxidation supports sub-surface microbial communities across__

__Namib Desert fog and aridity gradients__

Dana Z. Tribbia[1,2], Pedro H. Lebre[3], Xabier Vázquez-Campos[1], Angelique E. Ray[1,4], Thomas

Laird[1], Náthali Machado de Lima[1], Gillian Maggs-Kölling[5], Don A. Cowan[3], Belinda C.

Ferrari[1,2]#.

[1]School of Biotechnology and Biomolecular Sciences, UNSW Sydney, NSW, Australia.

[2]Evolution and Ecology Research Centre, UNSW Sydney, NSW, Australia.

[3]Centre for Microbial Ecology and Genomics, Department of Biochemistry, Genetics and

Microbiology, University of Pretoria, Pretoria, South Africa.

[4]NASA Ames Research Center, Moffett Field, California, USA.

[5]Gobabeb-Namib Research Institute, Walvis Bay, Namibia.

#Corresponding author: Belinda C. Ferrari, b.ferrari@unsw.edu.au

Keywords

Hydrogen oxidation, drylands, soil, precipitation, hyperarid, microbial resuscitation,

chemotrophy, hydro-genesis.

Abstract

Widely accepted climate predictions indicate that drylands will expand to cover more than

half of the Earth's terrestrial surface by the end of the 21st century. In these environments,

harsh conditions including nutrient and water limitations restrict plant and animal life,

thereby increasing the importance of soil microbial communities in nutrient cycling and

ecosystem functioning. The Namib Desert is a distinctive dryland ecosystem characterised by

a steep natural aridity gradient, transitioning from a coastal hyperarid zone influenced by

frequent fog deposition to an inland arid region receiving seasonal rainfall. This study

investigates the impact of water availability and moisture regime on microbial trace gas

oxidation and community composition across this aridity gradient. Quantitative analyses

revealed that total microbial abundance and activity indicators, including ATP concentrations

and respiration rates, were significantly ($p < 0.005$) reduced in hyperarid soils compared to

their arid counterparts. In contrast, hyperarid fog-dominated soils exhibited significantly ($p <$

0.0005) elevated rates of atmospheric hydrogen oxidation, even in the absence of water

inputs. We propose that sustained high-affinity hydrogen oxidation, coupled with rapid

microbial resuscitation following wetting events, supports shallow sub-surface microbial

communities in the Namib Desert, particularly in the coastal hyperarid zone. Together, these

findings challenge current understanding of the lower limits of microbial activity and reveal

alternate metabolic pathways that enable microbial persistence in hyperarid hot desert soils.

Importance

Drylands are expanding globally, yet the mechanisms that allow microbial life to persist

under extreme and sustained water limitation remain poorly understood. This study

demonstrates that atmospheric trace gas oxidation, particularly high-affinity hydrogen

oxidation, supports active and resilient microbial communities in hyperarid soils of the

Namib Desert, even in the absence of liquid water inputs. By revealing how microbes may

couple trace gas metabolism to energy and water generation, our findings provide new insight

into the lower limits of microbial activity in dry hot desert soils and highlight the need to

investigate how microbes persist and sustain soil ecosystem functioning.

Introduction

Water availability across a natural xeric gradient in the Namib Desert

The Namib Desert is a distinctive dryland ecosystem situated along a steep natural aridity

gradient on the west coast of Namibia [1, 2]. Low intensity rainfall events occur

intermittently (< 18 days/year) along this gradient [3, 4]. Precipitation frequency and volume

increase progressively from west to east, with the coastal region receiving the lowest average

annual rainfall (~10 mm/year) [4]. These sporadic rainfall events are separated by prolonged

dry periods, leaving much of the soil sub-surface desiccated for lengthy periods of time [3, 4].

The Aridity Index (AI), defined as the ratio of precipitation to potential evapotranspiration,

divides the Namib Desert into two primary zones: a hyperarid coastal region (AI < 0.05) and

a central arid region ($0.05 \leq AI < 0.2$) [5, 6] (Figure 1A). Despite receiving extremely low

annual rainfall, the hyperarid coastal zone experiences frequent fog events, approximately 3–

9 days/month, which decrease in frequency and duration beyond 60 km inland [2, 4, 7].

Considering the minimal precipitation along the hyperarid coast, fog-derived moisture has

been proposed as the dominant source of bioavailable water for these soil microbial

communities [3, 8-10]. Fog inputs are estimated to contribute up to 183 mm deposited

moisture annually [2, 4, 7], and contribute to slight increases in sub-surface (0–5 cm) soil

relative humidity, compared to soils from rainfall-dominated zones [3]. However, their

limited capacity to penetrate the desert pavement and enter the root zone restricts microbial

and plant productivity [1, 7]. Consequently, soil organic matter, nitrogen and carbon content

are lower in the hyperarid compared to the arid inland region [3, 11, 12]. High soil ionic

content from marine aerosol deposition in the coastal region may further limit the

bioavailability of fog-derived moisture [12]. While it is clear that fog is a strong driver of

lichen and biological soil crust development [8, 13, 14], there has been extensive debate over

the role of fog-derived water in supporting the sub-surface soil microbiome [10, 12, 15, 16].

Microbial trace gas oxidation and alternative survival strategies in dryland environments

It is well-established that the majority of microbial growth and metabolic processes cease at

water activities (a_w) < 0.9, whilst limited function at water activities as low as 0.3 a_w occurs

in some specialist taxa [17-20]. The Namib Desert is an environment under severe water

stress, with average immediate sub-surface (1–2 cm) soil water activities < 0.5 a_w [3]. Despite

this stress, diverse and metabolically active microbial communities persist within both

surface and shallow sub-surface soils [12, 20-23]. It is estimated that over 95% of soil

microorganisms exist in slow-growing or dormant states [24, 25], particularly during

desiccation stress [19]. Microbial dormancy followed by rapid resuscitation during wetting

has been proposed as a key survival strategy in dryland ecosystems including the Namib

Desert [22, 26-28].

Microbial trace gas oxidation is increasingly recognised as a generalist strategy to support

persistence in both hot and cold desert ecosystems [27, 29-32]. In particular, molecular

hydrogen (H_2) is a ubiquitous and energetically favourable substrate present at ~530 ppbv

throughout the atmosphere, capable of readily diffusing through soil surface layers and cell

membranes [27, 30, 33]. In some hyperarid environments, rapid H_2 uptake and the presence

of abundant high-affinity hydrogenases has led to the hypothesis that energy derived from

hydrogen oxidation may exceed requirements for cell maintenance, supporting microbial

growth via carbon fixation pathways [34-37]. This process, termed atmospheric

chemosynthesis, may complement or even surpass photoautotrophic primary production in

water-restricted environments with low phototroph abundances. Atmospheric chemosynthesis

has been established as a significant contributor to microbial carbon fixation in polar soils

[34, 36]. However, microbial trace gas oxidation and its potential contribution to carbon

fixation is poorly characterised in hot deserts, particularly in desiccated soils [31, 38].

In this pilot study, the impact of water limitation and regime on microbial atmospheric

hydrogen oxidation activity was investigated in desiccated and wetted soils, across a

longitudinal aridity gradient in the Namib Desert. The inverse distribution of fog and rainfall-

dominated water regimes across the coastal and inland regions of this gradient are proposed

to drive divergent microbial community composition and function. Trace gas chemotrophy is

predicted to support shallow sub-surface microbial communities in these severely water-

limited soils, particularly in the hyperarid coastal zone receiving low-volume fog inputs.

Materials and Methods

Soil sampling across the Namib Desert C14 aridity transect

Ten sites were sampled across a 200 km longitudinal aridity transect, referred to henceforth

as the C14 transect, spaced at 20 km intervals across a well-defined west-east aridity gradient

from the coastal to inland margins of the Namib Desert (Figure 1B). The C14 transect and

surrounding soils are divided into three well-defined xeric zones: the "Fog zone" (coastal

sites 2–6), "Hyperarid zone" (central sites 8–14) and "Arid zone" (inland sites 16–20). These

zones, proposed by Scola et al. (2018) [12], are supported by extensive research on soil

physicochemical parameters [11, 12, 16, 39, 40], rainfall, fog and humidity records [2-4, 11,

12, 16, 39], and microbial community analysis [12, 39, 40].

Duplicate (approx. 45 g) shallow sub-surface (1–5 cm) soil samples were aseptically

collected into sterile 50 mL tubes from the ten C14 transect sites in April 2023 (Figure 1B,

Table 1). Samples were immediately stored at 4 °C after sampling and transported to the

Centre for Microbial Ecology and Genomics (CMEG, Pretoria, South Africa), where they

were stored at 4 °C until shipment. Samples were imported to the University of New South

Wales (NSW, Australia), where half of the bulk soil from each site was stored at -80 °C and

the remaining soil stored at 4 °C until use. Soil samples used for hydrogen oxidation and

respiratory burst microcosm assays were stored at 4 °C, while DNA extraction and all other

analysis was conducted on soils stored at -80 °C.

Soil moisture analysis

Soil moisture content (SMC%) was calculated using the gravimetric oven-drying method

[41]. Triplicate 1.5 g soil subsamples from each site were placed into pre-weighed 20 mL

glass vials and oven-dried at 105 °C for 48 h. Sample vials were removed from the oven and

cooled to RT in a gas-tight container with silica beads (Ajax Finechem) before re-weighing

using an analytical balance (Sartorius AG). SMC% was recorded by calculating the mass of

water loss after drying as a percentage of the dry soil weight.

Aridity Index (AI) data for each site was retrieved from the Global Aridity Index and

Potential Evapotranspiration Database (Version 3) [6]. Soil water potential and water activity

was calculated from previously published datasets that used iButton remote sensing data to

record sub-surface soil temperature and relative humidity along the C14 transect [3]. Sub-

surface (0–10 cm) soil moisture content was retrieved from the FLDAS Noah Land Surface

Model L4 Global Monthly $0.1° × 0.1°$ dataset [42], subset to a 30-year monthly climatic

mean using NASA GES DISC by selecting the SoilMoi00_10cm_tavg variable between

1994–2024. Average annual rainfall data from each site was retrieved from a previously

published dataset across the C14 transect [12].

 Soil physicochemical parameters

Total soil carbon (TC), organic carbon (TOC), inorganic carbon (TIC), nitrogen (N),

hydrogen (H) and sulfur (S) analysis was performed at the Mark Wainwright Analytical

Centre (MWAC, UNSW Sydney, Australia). Duplicate 0.25 g soil samples from each transect

site were first passed through a 0.15 mm sieve, and soil particles between 0.15–2 mm were

finely ground to < 0.15 mm using a mortar and pestle. Ground soil samples were then oven-

dried at 45 °C until constant mass (> 72 h). To quantify TOC, two additional soil samples per

site were dried and ground according to the same procedure. Inorganic carbonates were

removed by acidification with 1 M HCl, added gradually in excess, and incubated in glass

vials at RT for 12 h [43]. Acidified samples were washed with 90 mL sterile deionised water

and dried at 45 °C until constant mass (> 120 h). Soil mass loss after acidification was

recorded using an analytical balance (Sartorius AG), and TOC was calculated as a percentage

of total soil mass. CHNS analysis was performed on all samples via combustion at 1150 °C

using a varioMACRO CUBE elemental analyser (Elementar, Germany). TIC was calculated

as the difference between TC and TOC content.

Soil pH was measured using 2.5 g soil slurries at a 1:2.5 soil:deionised water ratio, with a pH

meter (HI12303, Hanna Instruments). Additional physicochemical parameters at C14 transect

sites have been investigated previously [12, 40].

Microbial community activity metrics

Intracellular ATP luminescence assay

The BacTiter-Glo™ Microbial Viability Assay kit (Promega) was used to measure

intracellular ATP concentrations in C14 transect soil samples using a modified method

optimised for soils [44]. In triplicate, 1 g soil subsamples from each site were added to 4 mL

0.85% NaCl in 15 mL tubes and vortexed at maximum speed for 5 min to produce a soil

slurry. Slurry samples were then centrifuged at 180 rcf for 5 min and the supernatant was

collected. Triplicate 100 µL volumes of supernatant were used to perform the assay in 96-

well white luminescence plates (Greiner), according to manufacturer's instructions. A

standard curve was produced by serial dilution of 100 mM ATP solution (Thermo Fisher

Scientific). Background adjusted luminescence values were recorded using a CLARIOstar

Plus Microplate reader (BMG Labtech), and calculations were performed in MARS Data

Analysis Software v5.02 R3 (BMG Labtech).

Respiratory activity assay in dry and wetted soils

Basal respiration in dry soils and respiratory burst after wetting were recorded over 14 days

using a custom-built CO_2 and O_2 respirometry system (Qubit Systems, Canada). Triplicate 1

g subsamples of C14 transect soils stored at 4 °C were placed into sterile 114 mL glass vials

(Glass Vials Australia, NSW) and equilibrated at RT for three days. Following equilibration,

soil microcosms were sealed using gas-tight butyl rubber stoppers (Glass Vials Australia,

NSW) and incubated in the dark at 30 °C. Heat-killed controls were prepared from each site

using 1 g soil autoclaved at 121 °C for 15 min.

Headspace CO_2 and O_2 was measured at five timepoints (0, 24, 56, 120, 144 and 168 h) over

the one-week basal incubation period. A gas-tight syringe (Trajan Scientific) was used to

subsample 1 mL headspace gas from serum vial microcosms. Headspace subsamples were

injected through a gas-tight sample port at approximately 100 µL/s into 99.999% N_2 carrier

gas (Coregas, Australia). Using a Q-P103 Gas Pump (Qubit Systems), carrier and sample

gases were pumped through a Q-S151 Infrared CO2 analyser and Q-S102 O2 analyser (Qubit

Systems) at a constant rate. Flow rate was recorded via a Q-G266 Flow Monitor (Qubit

Systems). All readings were recorded using LabQuest Mini sensor data interfaces (Qubit

Systems) and headspace CO_2 and O_2 concentrations were calculated via the Logger Pro

v3.16.2 software.

After one week, each soil microcosm was wetted to approximately 50% water holding

capacity, estimated using the filter paper method [45]. Starting headspace CO_2 concentrations

were recorded immediately before wetting, as above, and wetted microcosms were incubated

under the same conditions. Repeat measurements were taken after 0.5, 1, 3, 5, 9, 24, 48 and

168 h. The cumulative respiratory burst was calculated for each site and timepoint by

normalising against the initial headspace CO_2 concentration.

Hydrogen oxidation rates in wet and dry soil microcosms

For dry microcosm H_2 oxidation assays, five biological replicate soil subsamples (2 g) and a

heat-killed control (2 g soil autoclaved at 121 °C for 15 min) were prepared from each site,

along with an empty serum vial negative control. Microcosm H_2 oxidation activity was

recorded using gas chromatography as described previously [34, 36]. All soil samples and

controls were placed into sterile 114 mL serum vials sealed with gas-tight butyl rubber

stoppers and incubated in the dark at 30 °C. Sterile H_2 gas in synthetic air-balance (BOC,

Australia) was injected into the headspace of each serum vial microcosm to achieve a final

headspace concentration of 10,000 ppbv. Headspace gas from each microcosm was

subsampled in 1 mL volumes at regular intervals using a gas-tight syringe (Trajan Scientific)

and H_2 concentration was measured using a Peak Performer 1 Gas Analyser (Peak

Laboratories, USA). For H_2 oxidation analysis in wet soil microcosms, the same procedure

was repeated using five additional 2 g soil subsamples from each site, wetted immediately

prior to gas addition using 400 µL sterile ddH_2O equilibrated to 30 °C.

Sub-atmospheric hydrogen oxidation was confirmed after two consecutive headspace H_2

readings < 530 ppbv. First order rate constants (k) were calculated using linear regression (R^2

> 0.9). Atmospheric hydrogen oxidation rates were calculated at headspace H_2 = 530 ppbv

with a total microcosm headspace volume of 114 mL at 30 °C and 1 atm. Rates were

normalised against total 16S rRNA gene copy numbers per gram soil.

DNA extraction and quantitative PCR

Genomic DNA was extracted from quadruplicate 0.5 g bulk soil subsamples stored at -80 °C

from each site, using the FastDNA™ Spin Kit for Soil (MP Biomedicals), according to

manufacturer's instructions. Extracted DNA was eluted in 70 µL warm (60 °C) UltraPure™

DNase/RNase-free distilled water (Invitrogen) and stored at -80 °C. Quantity and purity of

extracted DNA was assessed using both Nanodrop (Thermo Fisher Scientific) and the

Qubit™ dsDNA High Sensitivity Assay (Thermo Fisher Scientific). To reduce PCR

inhibition, gDNA was further diluted in UltraPure™ DNase/RNase-Free distilled water

(Invitrogen) to produce final concentrations between 1.76–20.00 ng/µL.

Extracted gDNA was used for qPCR quantification of total bacterial abundance using the

universal 16S rRNA gene degenerate primer set Eub1048f/Eub1194r (Eub1048f; 5'-

GTGSTGCAYGGYTGTCGTCA, Eub1194r; 5'ACGTCRTCCMCACCTTCCTC) [46], as

described previously [47]. QPCR reaction mixtures were prepared using 10 µL QuantiNova

SYBR Green PCR Master Mix (Qiagen, Australia), 0.5 µL of each 40 µM forward and

reverse primer (Integrated DNA Technologies), 7 µL UltraPure™ DNase/RNase-free

distilled water (Invitrogen), 1 µL of 5 µg/mL T4 Gene 32 (New England Biolabs), and 1 µL

diluted template gDNA or positive control. A synthetically designed gene fragment (gBlocks;

Integrated DNA Technologies, Australia) containing a representative 16S rRNA gene

sequence (MF689012.1) was used to generate a standard curve over five orders of magnitude

[47]. All standards, samples and negative template control reactions were performed in

triplicate.

Reaction mixtures were added to 96-well thin wall PCR plates (Bio-Rad Laboratories) and

sealed using optically clear plate seals (Bio-Rad Laboratories). The thermocycling protocol

was completed using a CFX96 Touch™ Real-Time PCR Detection System (Bio-Rad

Laboratories). Plates were incubated at 95 °C for 5 min, followed by 38 cycles of 95 °C for

20 s and 60 °C for 50 s. Quantitative fluorescence data was spectrophotometrically collected

during the combined extension and annealing step. Following PCR amplification, a melt

curve was performed from 50–95 °C, with fluorescence data collected at 0.5 °C steps

throughout. Data analysis was conducted using CFX Manager software (Bio-Rad

Laboratories). Amplification specificity was confirmed by visual inspection of melt peaks.

16S rRNA gene amplicon sequencing of soil microbial communities

Paired-end amplicon sequencing was performed on gDNA samples from all transect sites at

the Ramaciotti Centre for Genomics (UNSW, Australia). The barcoded primer pair 515F-

Y/806RB (515F-Y; 5'-GTGYCAGCMGCCGCGGTAA, 806RB; 5'-

GGACTACNVGGGTWTCTAAT) was used to amplify the V4 region of the 16S rRNA gene

using the Illumina MiSeq v2 platform [48, 49]. Raw reads were processed into amplicon

sequence variants (ASVs) with DADA2 v1.30.0 [50] in R v4.3.1. Reads with ambiguous

bases were discarded, and residual primer sequences were removed with cutadapt v4.3 [51].

Forward and reverse reads were quality filtered using the following parameters: reads

matching phiX removed, 2 maximum expected errors allowed, truncation length of 220 and

215 nt, minimum read length of 150 nt, and a read truncation quality threshold of 2. Paired

reads were merged with mergePairs() using default parameters and chimeric reads were

removed with the removeBimeraDenovo() function, using the consensus method. Taxonomy

was assigned to merged reads with the assignTaxonomy() command, implementing the RDP

Naive Bayesian Classifier algorithm [52] using SILVA database v138.1 [53]. Processed

ASVs assigned to mitochondria or chloroplasts were removed.

ASV sequences were aligned using MAFFT v7.526 [54], and an unrooted phylogenetic tree

was built using FastTree v2.1.11 [55] under the GTR+Gamma model [56]. The resulting tree

was utilised to calculate weighted UniFrac distance matrices [57] using phyloseq v1.46 [58].

Raw ASV abundance tables were rarefied to the minimum read count using phyloseq v1.46

[58]. Microbial community composition across C14 transect sites was analysed in vegan

v2.6-10 [59] using principal coordinates analysis (PCoA) based on weighted UniFrac

distances and the rarefied count table. Permutational ANOVA (PERMANOVA) was

performed using the adonis2 function in vegan v2.6-10 [59] with 999 permutations, where

each explanatory variable was tested individually against the weighted UniFrac dissimilarity

matrix.

Differentially abundant taxa were identified using Maaslin2 v1.16.0 [60] from rarefied data

aggregated to taxonomic level, excluding unclassified taxa. Linear regression models

(analysis method = "LM") were used with default standardisation settings, and no

normalisation was applied. The Fog zone was specified as the reference group for

comparisons across xeric zones. The minimum non-zero prevalence threshold was set at 30%

and nominal p-values were adjusted via the Benjamini-Hochberg correction method to a

significance threshold of qval $\leq$ 0.20.

Data analysis and visualisation

All community activity and aridity datasets were analysed in RStudio [61] using R v4.3.1.

Data processing was performed using reshape2 v1.4.4 [62], stringr v1.5.1 [63] and readr

v2.1.5 [64]. Aridity Index datasets were processed and visualised using the terra v1.8.21 [65]

and tidyterra v0.7.0 [66] packages. Statistical analysis was performed using dplyr v1.1.4 [67]

and ggpubr v0.6.0 [68]. Correlation analysis was performed using corrplot v0.92 [69].

Mantel correlogram was calculated with vegan's v2.6-10 [59] mantel.correlog() function

between the weighted UniFrac distance matrix and the geographical distances between

sampling points. Geographical distances between sampling points were calculated from GPS

coordinates using the Haversine method [70] as implemented in the geosphere package v1.5-

20 [71]. Optimal number of bins for Mantel correlogram was calculated based on the

Freedman-Diaconis rule [72] with the hist.FD() function from the MASS package v7.3-60

[73].

All other data visualisation was performed using ggplot2 v3.5.1 [74], patchwork v1.3.0 [75],

pals v1.10 [76], tidyverse v2.0.0 [77] and ggtext v0.1.2 [78].

Results

Water availability and soil properties across the C14 transect

The Aridity Index (AI) of sites across the C14 transect increased from west to east, with the lowest recorded at Fog zone site C14-2 (AI = 0.0072), and the highest at Arid zone site C14-20 (AI = 0.0780) (Table 1). Sites C14-2 to C14-14, spanning the Fog and Hyperarid zones, were classed as hyperarid (AI < 0.05), and Arid zone sites C14-16–20 were classed as arid ($0.05 \leq AI < 0.2$) [5].

Table 1: Soil moisture metrics and physicochemical properties of C14 transect sampling sites.

Site	Xeric zone [a]	Lat	Lon	Aridity Index (AI)	Gravimetric SMC%	Annual rainfall (mm)	Water potential (MPa)	Water activity (a_w)	Sub-surface SMC (m^3/m^3) [b]	TC%	TIC%	TOC%	N%	H%	S% [c]	pH
C14-2	F	-23.001	14.672	0.0072	1.53	16.7	-100.81	0.498	0.173	0.27	0.19	0.08	0.05	0.41	2.32	6.70
C14-4	F	-23.018	14.860	0.0112	0.20	21.5	-123.47	0.432	0.182	0.18	0.14	0.04	0.04	0.06	0.08	7.19
C14-6	F	-23.066	15.040	0.0196	0.82	40	-144.38	0.384	0.169	0.63	0.55	0.08	0.03	0.26	0.39	6.91
C14-8	HA	-23.143	15.209	0.0233	0.23	59	-162.43	0.333	0.163	0.16	0.11	0.05	0.04	0.09	0.07	7.03
C14-10	HA	-23.246	15.360	0.0308	0.81	62	-162.02	0.346	0.163	1.03	0.89	0.14	0.05	0.30	0.01	8.05
C14-12	HA	-23.311	15.534	0.0426	0.42	78.8	-184.07	0.312	0.162	0.38	0.18	0.20	0.07	0.26	nd	7.87
C14-14	HA	-23.325	15.715	0.0456	0.49	100.5	-145.68	0.384	0.163	1.05	0.83	0.22	0.05	0.27	nd	7.53
C14-16	A	-23.321	15.862	0.0532	0.35	133	-207.39	0.269	0.166	1.13	0.87	0.25	0.08	0.24	nd	7.71
C14-18	A	-23.345	16.011	0.0644	0.27	169.3	-231.39	0.230	0.170	0.22	0.04	0.18	0.06	0.17	nd	8.45
C14-20	A	-23.245	16.143	0.0780	0.31	184.8	-204.08	0.288	0.178	0.74	0.10	0.64	0.09	0.33	0.18	7.45

[a] *Xeric zonation is denoted as F = Fog zone, HA = Hyperarid zone, A = Arid zone*
[b] *Sub-surface (0–10 cm) soil moisture content*
[c] *nd denotes sulfur content not detected*

Gravimetric SMC% varied across the transect, ranging from 0.20–1.53% (0.54% ± 0.42%,

mean ± SD) (Table 1, Table S1). Average SMC% was highest in the Fog zone (0.85% ±

0.59%) followed by the Hyperarid (0.49% ± 0.25%) and Arid (0.31% ± 0.10%) zones.

Average water potential and a_w were highest in the Fog zone (water potential = -122.89 ±

17.79 MPa; a_w = 0.438 ± 0.047), followed by the Hyperarid (water potential = -163.55 ±

13.64 MPa; a_w = 0.344 ± 0.026) and Arid (water potential = -214.28 ± 12.17 MPa; a_w = 0.262

± 0.024) zones. AI and rainfall were both negatively correlated with water potential ($p \leq$

0.001) and a_w ($p \leq 0.01$) (Figure S1). Gravimetric SMC% was positively correlated with a_w (ρ

= 0.67, $p \leq 0.05$). Sub-surface (0-10 cm) soil moisture content was not significantly

correlated with any other moisture metric ($p > 0.05$).

Soil pH ranged from slightly acidic to alkaline (6.70–8.45) and was highest in the Arid zone

(7.87 ± 0.42, mean ± SD), followed by the Hyperarid (7.62 ± 0.39) and Fog (6.93 ± 0.20)

zones (Table 1). Total carbon (TC) content was low (0.16–1.13%) across the transect, and

inorganic carbon (TIC) represented the majority of soil carbon in all sites except C14-12, 18

and 20. Total nitrogen (N) and organic carbon (TOC) content were lowest in the Fog zone (N

= 0.04% ± 0.01%, TOC = 0.06% ± 0.02%) and increased in the Hyperarid (N = 0.05% ±

0.01%, TOC = 0.15% ± 0.07%) and Arid (N = 0.07% ± 0.01%, TOC = 0.36% ± 0.20%)

zones. TOC, N and pH were significantly ($p \leq 0.05$) lower in the Fog zone compared to the

rest of the transect and were significantly positively correlated with AI (Figure S1).

Microbial activity and abundance biomarkers across xeric zones

Microbial activity and 16S rRNA gene abundances increased from west to east across the

Fog, Hyperarid and Arid zones of the C14 transect (Figure 2). Total 16S rRNA gene copy

numbers ranged from 3.82×10^5 g^{-1} soil at Fog site C14-2 to 3.10×10^7 g^{-1} soil at Arid site

C14-16 (Figure 2A, Table S2). Average 16S rRNA gene copy number increased significantly

($p \leq 0.01$) between each xeric zone as AI and rainfall increased (Figure S2A). The lowest

average 16S rRNA gene abundance was observed in the Fog zone (1.04×10^6 copies g^{-1}),

increasing 9-fold in the Hyperarid zone (9.15×10^6 copies g^{-1}) and by an order of magnitude

in the Arid zone (2.36×10^7 copies g^{-1}).

Similarly, cumulative 24-hour respiratory burst values ranged from 23.6 ppmv CO_2 g^{-1} soil

(Hyperarid site C14-8) to 2891.0 ppmv CO_2 g^{-1} soil (Arid site C14-20) (Figure 2B, Table S3-

4) and increased significantly ($p \leq 0.05$) between each xeric zone (Figure S2B). The average

24-hour cumulative respiratory burst across the Fog, Hyperarid and Arid zones was 213.2,

532.1 and 2181.9 ppmv CO_2 g^{-1} soil, respectively. Intracellular ATP concentrations ranged

from 0.002 nmol g^{-1} soil at Fog site C14-6 to 0.071 nmol g^{-1} soil at Arid site C14-18 (Figure

2C, Table S5). Whilst Arid zone sites recorded significantly ($p \leq 0.01$) higher ATP

concentrations than Fog and Hyperarid zone sites, average ATP concentrations between the

Hyperarid and Fog zones were not significantly different ($p > 0.05$) (Figure S2C).

Within the Fog zone, average ATP concentrations in sites C14-2 and C14-4 were 15- and 4-

fold higher, respectively, compared to site C14-6. Aridity Index was significantly positively

correlated with total 16S rRNA gene copy numbers ($\rho = 0.83$, $p \leq 0.01$), intracellular ATP (ρ

$= 0.76$, $p \leq 0.05$) and 24 h respiratory burst ($\rho = 0.90$, $p \leq 0.001$) (Figure S1). Total 16S copy

numbers were significantly positively correlated with cumulative 24 h respiratory burst ($\rho =$

0.92, $p \leq 0.001$) but not intracellular ATP content ($p > 0.05$).

Trace gas oxidation rates in Namib Desert soil microcosms

In dry soil microcosms, two Fog zone sites (C14-2 and C14-4) exhibited rapid H_2 oxidation

activity to sub-atmospheric (< 530 ppbv) levels (Figure 3A). The remaining dry microcosms

demonstrated slow H_2 oxidation that did not reach sub-atmospheric levels after 1,500 h (~2

months) (Figure S3A). In contrast, soil wetting resulted in immediate (< 30 min) and rapid H_2

oxidation activity across all ten transect sites (Figure S3B), with sub-atmospheric hydrogen

oxidation activity observed in all moisture-stimulated microcosms (Figure 3B).

At atmospheric headspace H_2 concentrations, dry soil microcosms from Fog sites C14-2 and

C14-4 oxidised hydrogen at average rates of 2.27×10^{-7} and 4.27×10^{-8} nmol h^{-1} 16S copy^{-1},

respectively (Table S6). Both C14-2 and C14-4 microcosms carried out significantly ($p \leq$

0.0001) accelerated atmospheric H_2 oxidation rates following moisture stimulation,

increasing over 30-fold compared to dry microcosm rates (Figure S4). In contrast, all wet soil

microcosms demonstrated rapid atmospheric H_2 oxidation rates, ranging from 1.59×10^{-8}

nmol h^{-1} 16S copy^{-1} at Arid site C14-18 to 9.39×10^{-6} nmol h^{-1} 16S copy^{-1} at Fog site C14-2

(Figure 3B, Table S7). In moisture-stimulated microcosms, the fastest H_2 oxidation rates were

observed in Fog zone soils (average $= 3.55 \times 10^{-6}$ nmol h^{-1} 16S copy^{-1}), while atmospheric H_2

oxidation rates decreased significantly ($p \leq 0.0001$) in the Hyperarid (3.84×10^{-8} nmol h^{-1}

16S copy^{-1}) and Arid (2.28×10^{-8} nmol h^{-1} 16S copy^{-1}) zones (Figure 3B).

Microbial community composition along the Namib Desert C14 transect

Soils were dominated by the bacterial phyla *Actinobacteriota*, *Proteobacteria*, *Chloroflexi*

and *Bacteroidota*, and the archaeal phylum *Crenarchaeota*, together accounting for > 77.3%

of the microbial community (Figure 4A, Table S8). The relative abundance of

Actinobacteriota decreased significantly ($p \leq 0.001$) across the transect as AI increased

(Figure S6), decreasing from 42.3% in the Fog zone to 22.1% in the Arid zone (Figure 4A).

Conversely, the relative abundance of *Proteobacteria* increased significantly ($p \leq 0.005$)

from 11.5% in the Fog zone to 26.7% as the most abundant phylum in the Arid zone. The

five most abundant families were *Rubrobacteriaceae* (13.4%, *Actinobacteriota*),

Nitrososphaeraceae (8.3%, *Crenarchaeota*), *Beijerinckiaceae* (7.7%, *Proteobacteria*),

Chloroflexi family AKIW781 (4.9%) and *Geodermatophilaceae* (3.9%, *Actinobacteriota*)

(Figure 4B).

PCoA ordination analysis explained 67.51% of total variance across the first two axes and

showed clear seperation of microbial communities by xeric zone across the first axis (Figure

4C). Arid sites C14-16–20 clustered tightly, whereas Hyperarid sites C14-8–14 showed

greater dispersal. Fog zone sites C14-2–6 exhibited the highest dispersion across both axes,

indicating relatively higher compositional variability compared to the other zones. AI was the

most significant ($p \leq 0.001$) explanatory variable, explaining 34.78% of variance in

community structure (Table S9). Community composition was also significantly associated

with rainfall ($p \leq 0.005$, $R^2 = 0.35$), nitrogen ($p \leq 0.01$, $R^2 = 0.29$), water potential ($p \leq 0.01$,

$R^2 = 0.29$), pH ($p \leq 0.05$, $R^2 = 0.25$) and water activity ($p \leq 0.05$, $R^2 = 0.27$). The Mantel

correlogram showed significant positive spatial autocorrelation (r = 0.37, $p < 0.01$) between

sites up to approximately 20 km apart (i.e. immediately consecutive sites) (Figure S5).

Between 20-60 km, correlations were weak (r < 0.2) and non-significant ($p > 0.05$),

indicating that geographical distance alone does not explain community differences. At

distances of ~110 km, sites were more dissimilar than expected by chance (r = -0.33, p <

0.05), while non-significant ($p > 0.5$) negative correlations were observed at all other

distances.

In total, 129 taxa (13 phyla, 4 classes, 26 orders, 37 families and 49 genera) were

significantly (qval $\leq$ 0.20) differentially abundant across xeric zones (Figure 5, Figure S7,

Table S10). At the phylum level, the relative abundance of *Acidobacteriota*, *Abditibacteriota*,

Firmicutes, *Proteobacteria, Bacteroidota* and *Crenarchaeota* increased significantly (qval $\leq$

0.20) in the Arid compared to the Fog zone. Similarly, seven phyla were enriched in the

Hyperarid zone relative to the Fog zone: *Entotheonellaeota, Firmicutes*, *Nitrospirota*,

Acidobacteriota, *Abditibacteriota*, *Armatimonadota* and *Crenarchaeota*. Conversely, the Fog

zone contained significantly (qval ≤ 0.20) greater relative abundances of *Gemmatimonadota*,

Actinobacteriota, *Deinococcota* and *Bdellovibrionota*.

Eight of the 12 families with total relative abundance ≥ 1.5% were significantly (qval ≤ 0.20)

differentially abundant between xeric zones (Figure 5). Of these, relative abundances of

Chitinophagaceae, Geodermatophilaceae, Beijerinckiaceae and *Pyrinomonadaceae* were

significantly (qval ≤ 0.20) greater within both the Arid and Hyperarid zones relative to the

Fog zone. In contrast, *Nitrososphaeraceae* were significantly more abundant in the Hyperarid

but not the Arid zone. Within the Fog zone, *Trueperaceae* were present at a greater relative

abundance compared to both other zones*, while *Longimicrobiaceae* and *Rubrobacteriaceae*

had a greater relative abundance compared to the Hyperarid or Arid zone, respectively.

Across the transect, > 43.5% of ASVs remained unclassified at the genus level, with over half

of total ASVs unclassified in sites C14-4, C14-6 and C14-20 (Table S8). Seven classified

genera were significantly (qval ≤ 0.20) enriched in the Fog zone (Figure S7, Table S10). The

relative abundance of *Truepera, Oxalicibacterium* and *Amaricoccus* was greater in the Fog

zone relative to both other zones, while *Conexibacter,* Ellin6055 and *Lysobacter* were

enriched compared to the Hyperarid zone only, and *Rubrobacter* relative to the Arid zone. An

additional five genera were strongly differentially abundant (coef > 5.0), all enriched within

the Arid zone: *Flaviaesturariibacter, Planococcus, Edaphobaculum, Blastocatella* and

Massilia.

Discussion

Differences in moisture regime and aridity across xeric zones of the Namib Desert are well-

established drivers of microbial community composition and function [12, 39, 40]. In this

pilot study, high-affinity H_2 oxidation and rapid resuscitation following soil wetting were

identified as key mechanisms likely to be supporting sub-surface soil microbial communities

across this aridity gradient. Dry soil microcosms from the hyperarid Fog zone exhibited rapid

hydrogen oxidation to sub-atmospheric concentrations without moisture stimulation (Figure

3A). In low-abundance Fog zone sites capable of H_2 oxidation in dry conditions, elevated

intracellular ATP concentrations compared to Fog zone site C14-6 were also observed,

providing evidence of a continuously active and strongly desiccation-resistant microbial

community.

Trace gas oxidation supports microbial communities in hyperarid Namib Desert soils

Across the Namib Desert C14 transect, AI and average annual precipitation were highest in

the Arid zone (Table 1), with significantly ($p \leq 0.005$) higher total microbial abundances,

intracellular ATP and respiratory burst observed compared to the Hyperarid and Fog zones

(Figure 2, Figure S2). Despite this reduced biomass, dry soils from Fog zone sites C14-2 and

C14-4 demonstrated a remarkable capacity for rapid, high-affinity (530 ppbv) hydrogen

oxidation without water addition, at rates of 2.27×10^{-7} and 4.27×10^{-8} nmol h^{-1} 16S copy^{-1},

respectively (Figure 3A). These rates exceed biomass-normalised H_2 oxidation rates recorded

in hyperarid Negev Desert soils by over two orders of magnitude [38], and contrast with

previous findings from hot desert soil studies where sub-atmospheric H_2 oxidation activity

was strongly limited in the absence of soil wetting [29, 31]. Although even faster atmospheric

H_2 oxidation rates have been recorded in Antarctic desert soils [34, 36], oxidation rates in

these Fog zone sites were comparable to those in cold, arid environments such as the high

Arctic [36] and Andean Altiplano [79], as well low-temperature Antarctic microcosms [35].

In these areas, energy derived from atmospheric hydrogen oxidation is predicted to sustain

microbial community biomass and potentially support hydrogenotrophic growth. Given the

extremely limited organic carbon availability (0.04–0.08%) and infrequent rainfall within the

Fog zone (Table 1) [3, 12], trace H_2 oxidation is likely to be a key contributor to cellular

energy requirements.

Higher soil moisture content and water potential within the Fog zone due to regular fog

inputs and humid coastal airflow [3] may have contributed to this unusual capacity for high-

affinity hydrogen oxidation in dry soils. However, these soils were severely water-restricted,

with average annual water availability < 0.5 a_w [3] and gravimetric SMC% as low as 0.20%

at site C14-4 (Table 1). Prior transcriptomic investigations on Namib Desert Fog zone soils

have revealed active transcription of genes associated with major metabolic pathways

including carbon and nitrogen assimilation, resuscitation and replication, even at

exceptionally low water activities (< 0.28 a_w) [20]. Furthermore, both Fog zone sites capable

of sub-atmospheric H_2 oxidation under dry conditions also exhibited elevated intracellular

ATP concentrations (Figure 2C), a known byproduct of hydrogen oxidation [80], compared

to Fog site C14-6. These findings support the growing body of evidence that some microbial

cells in hot desert soils retain significant metabolic functionality, including trace gas

oxidation, during severe desiccation [19].

Soil wetting results in rapid microbial resuscitation and hydrogen oxidation

Rapid resuscitation following re-wetting is a dominant microbial survival strategy in global

dryland soils, with the majority of microbial growth restricted to short windows during

intermittent rain events [3, 22, 26, 28, 81]. In this study, moisture stimulation resulted in

immediate (< 30 min) H_2 uptake in soil microcosms from all Namib Desert C14 transect sites

(Figure S3B). In similar desiccated dryland samples, soil wetting has prompted rapid (< 15

min) resuscitation and upregulation of metabolic activity, including H_2 oxidation [26, 28, 31].

Consistent with observations of Australian and Negev Desert microcosms [26, 31, 38], soil

wetting significantly ($p < 0.0001$) increased hydrogen oxidation rates over 30-fold (Figure

S4) and resulted in sub-atmospheric H_2 oxidation activity in all microcosms (Figure 3B). In

wetted soil microcosms, biomass-adjusted H_2 oxidation rates were highest in Fog zone sites

and decreased significantly ($p < 0.01$) across xeric zones as aridity decreased (Figure 3B).

These findings support the hypothesis that trace gas oxidation is a survival strategy associated

with aridity and organic carbon limitation, complementing phototrophic and heterotrophic

growth in moisture-limited environments [38, 82]. Many *Actinobacteriota* regulate dormancy

and resuscitation of diverse bacterial lineages within the soil microbiome through the

production of extracellular resuscitation promoting factors (RPFs) [83, 84], and are also

implicated in trace gas oxidation [32]. The high relative abundance of *Actinobacteriota* taxa

across the transect (18.5–56.7%), particularly within the Fog zone (Figure 4A), may have

contributed to this rapid increase in H_2 oxidation activity after wetting.

Impact of water regime and availability on Namib Desert microbial community

composition

Water availability is a major driver of microbial abundance, diversity, and activity in

terrestrial environments, with the diversity and abundance of soil bacteria and fungi

decreasing as aridity increases [15, 19, 85, 86]. Photosynthetic growth is strongly dependent

on bioavailable water at minimum thresholds of 0.8–0.9 a_w, with *Cyanobacteria* particularly

inhibited by desiccation [9, 17]. Although *Cyanobacteria* have been identified as key taxa in

the hypolithic communities that colonise the underside of quartz rocks within the Namib

Desert, the relative abundances of these taxa are markedly reduced in open soils and biocrusts

[9, 23, 87]. This trend was reflected across the C14 transect, where *Cyanobacteria* comprised

0.2–4.7% of the total community (Figure 4A). Instead, sites were dominated by

Actinobacteriota (18.5–56.7%), *Proteobacteria* (8.2–30.6%) and *Chloroflexi* (6.5–27.5%)

lineages (Figure 4A). These metabolically flexible phyla have been linked with trace gas

oxidation and the genetic capacity for atmospheric chemosynthesis, and are proposed to

support primary production in water-limited environments [34-36, 38, 47].

Across the transect, the ratio between *Actinobacteriota* and *Proteobacteria* varied by xeric

zone, with *Actinobacteriota* significantly (qval ≤ 0.20) decreasing in relative abundance as AI

increased (Figure S6). In the Fog zone, *Actinobacteriota* were dominant (42.3% relative

abundance; Figure 4A), with *Rubrobacteriaceae* highly abundant across the transect (3.7–

21.7%, Figure 4B) and significantly enriched in the Fog zone at both the family and genus

(*Rubrobacter*) level (Figure 5, Figure S7, Table S10). *Rubrobacteriaceae* are associated with

strong resistance against UV radiation, desiccation, heat and oxidative stress, as well as

extensive DNA repair potential and mixotrophic growth strategies [26, 81, 88]. Furthermore,

Rubrobacteriaceae are key active taxa in global dryland soil microbial communities,

including in the Namib [20, 28], Negev [26, 81] and Atacama Deserts [86, 89]. High-affinity

oxygen-tolerant [NiFe]-hydrogenases have been identified in the genomes of dryland-

associated *Rubrobacteriaceae* [26, 36, 38, 81], and hydrogenase mRNA transcripts assigned

to *Rubrobacteriaceae* MAGs have been recovered from microcosms capable of trace H_2

oxidation under wet and dry conditions [26, 31]. In polar and hot deserts, *Rubrobacteriaceae*

are associated with mixotrophic growth due to the genetic capacity to use energy derived

from trace hydrogen oxidation and rhodopsin-based light harvesting to support persistence or

growth during starvation [35, 81]. Several other differentially abundant Fog zone taxa present

at lower relative abundances are also associated with hydrogen chemotrophy, including

Conexibacter (*Actinobacteria*) [36, 90], *Euzebyaceae* (*Actinobacteria*) [36, 91],

Xanthomonadaceae (*Proteobacteria*) [35, 36] and *Trueperaceae* (*Deinococcota*) [36]. These

hydrogenotrophic and desiccation resistant taxa, particularly the highly abundant

Rubrobacteriaceae, likely contribute to the significantly faster rates of H_2 oxidation observed

within the Fog zone.

In the Arid zone, the differentially abundant families *Beijerinckiaceae* (*Proteobacteria*) and

Chitinophagaceae (*Bacteroidota*) constituted > 20.9% of the microbial community (Figure 5,

Figure 4B). Within the dominant *Beijerinckiaceae* family, > 98.5% of classified ASVs were

assigned to two genera: *Microvirga* and *Psychroglaciecola* (Table S8). These heterotrophic

nitrogen-cycling genera are associated with bryophytes and rhizosphere root nodules across a

variety of arid environments, including the Namib Desert [21, 92-96]. Many *Microvirga*

species have the capacity to utilise plant-derived substrates for growth and synthesise plant-

growth promoting compounds [97-99] and have been identified as highly abundant in the

rhizosphere of Namib Desert plants such as *Sesuvium sesuvioides* and *Stipograstis* species

[21, 93]. Heterotrophic growth strategies may be more prevalent in the Arid zone compared

to fog-dominated areas due to the capacity of rainfall inputs to saturate soil and facilitate

dispersion of organic matter, as well as the higher TOC (0.18–0.64%) and N (0.06–0.09%)

content in the Arid zone (Table 1) [15]. The production of plant-growth promoting

compounds has also been reported in the differentially enriched Arid zone *Chitinophagaceae*

family, which are abundant within the rhizosphere of Namib Desert *Acanthosicyos horridus*

shrubs and are associated with degradation of plant-derived cellulose and chitin [100-102].

These microorganisms may contribute to supporting plant growth in the nutrient and moisture

limited Arid region of the Namib Desert.

Across the transect, a high proportion of ASVs belonged to unclassified clades, with > 26.9%

unclassified at the family level and > 43.6% at the genus level (Table S8). This high relative

abundance of "microbial dark matter" has been observed across global drylands [103],

representing a significant and underexplored microbial resource. Further research into these

uncharacterised taxa is recommended to support a greater understanding of the complex

microbially mediated processes within desert environments.

Microbial hydro-genesis in Namib Desert sub-surface soils

Hydrogen oxidation, in addition to supporting energy requirements for basal cellular

metabolism [27, 33, 104], has been proposed to contribute to intracellular water production

through a process termed microbial hydro-genesis [15, 19, 35]. In this study, the

determination of substantial H_2 oxidation rates in dry soil microcosms, normalised against

cell numbers (estimated as 16S gene copies), allows for preliminary quantitative assessment

of intracellular water input derived from trace hydrogen oxidation.

Using the fastest H_2 oxidation rate determined in unwetted soil microcosm experiments (2.27

$\times 10^{-7}$ nmol h^{-1} 16S copy^{-1}, soil sample C14-2) (Figure 3A) and making the assumption that

cells contain single genomes and a single 16S gene copy per genome, we estimate H_2

oxidation (equimolar with H_2O production) to be 2.27×10^{-7} nmol h^{-1} cell^{-1}, equivalent to

4.09 fg H_2O produced h^{-1} cell^{-1}. Given a typical saturated cellular water content of 516 fg (for

E. coli, [105]), this contributes approximately 0.79% of total (saturated) cell water per hour.

In desiccated cells, such as those from site C14-4 (SMC% = 0.20%, a_w = 0.43; Table 1), this

contribution would be proportionally higher. While we acknowledge the multiple

assumptions built into this calculation, particularly those related to 16S gene copies and

cellular water contents, we argue that this calculated value of hydro-genesis derived from

atmospheric hydrogen oxidation in unwetted, low-moisture soils is significant, providing

support to the suggestion that this process contributes to cellular water budgets. This hydro-

genic contribution, while unlikely to be functionally significant in water replete cells, may be

vitally important to cellular function in moisture-limited environments [15, 19, 37].

Conclusions

In this study, the critical role of trace gas oxidation, particularly high-affinity hydrogen

oxidation, in sustaining microbial communities in hyperarid Namib Desert soils is

highlighted. Rapid sub-atmospheric H_2 oxidation activity in unwetted moisture-limited soils

is proposed to contribute to cellular ATP and water budgets, supporting a continuously active

desiccation resistant microbial community. In hyperarid, fog-dominated soils, taxa implicated

in trace gas oxidation dominated, while heterotrophic taxa were differentially abundant

within the Arid zone. Emerging evidence of interconnected phototrophic, heterotrophic and

hydrogenotrophic growth strategies in hot and polar desert soils supports the need for broader

studies that link water regime to functional capacity and primary production dynamics in

these complex ecosystems.

Acknowledgements

This research was supported by an Australian Government Research Training Program (RTP)

Scholarship awarded to DZT, the Australian Research Council (ARC) Discovery Project

(DP240102658) awarded to BCF and a National Research Foundation of South Africa grant

(137954) awarded to DAC. We thank the Gobabeb-Namib Research Institute

(http://gobabeb.org/) in Namibia for their hospitality and field assistance. Field research was

conducted under a research permit RPIV00112018, issued by the National Commission on

Research, Science and Technology, and supported by the Ministry of Environment, Forestry

and Tourism in Namibia. This research includes computations using the computational

cluster Katana supported by Research Technology Services at UNSW Sydney (DOI:

10.26190/669X-A286).

<u>CRediT author contribution statement</u>

DZT (Formal analysis, Investigation, Methodology, Visualisation, Writing – original draft,

Writing – review & editing), **PHL** (Conceptualisation, Writing – review & editing), **XVC**

(Formal analysis, Data curation, Visualisation, Writing – review & editing), **AER**

(Methodology, Writing – review & editing), **TL** (Formal analysis, Methodology), **NMDL**

(Formal analysis, Writing – review & editing), **GMK** (Project administration, Resources),

DAC (Conceptualisation, Funding acquisition, Project administration, Resources,

Supervision, Writing – review & editing), **BCF** (Conceptualisation, Funding acquisition,

Project administration, Resources, Supervision, Writing – review & editing). All authors

contributed to the final manuscript.

<u>Conflicts of interest</u>

The authors declare that they have no competing interests.

<u>Data availability</u>

Raw amplicon sequencing data are deposited upon publication in ENA under BioProject

accession PRJEB97070. All remaining data generated or analysed during this study are

included in this published article and its supplementary information files.

<u>References</u>

1.   Seely MK. The namib dune desert: An unusual ecosystem. *Journal of Arid*
Environments. 1978;**1**:117-28 https://doi.org/10.1016/S0140-1963(18)31736-1
2.   Lancaster J, Lancaster, N., Seely M. Climate of the central namib desert. *Madoqua*.
1984;**1984**:5-61

3.   Bosch J, Marais E, Maggs-Kölling G *et al.* Water inputs across the namib desert:
Implications for dryland edaphic microbiology. *Frontiers of Biogeography*. 2022;**14**
https://doi.org/10.21425/F5FBG55302
4.   Eckardt FD, Soderberg K, Coop LJ *et al.* The nature of moisture at gobabeb, in the
central namib desert. *Journal of Arid Environments*. 2013;**93**:7-19
https://doi.org/https://doi.org/10.1016/j.jaridenv.2012.01.011
5.   United Nations Environment Programme. *World atlas of desertification: Second*
edition, 1997.
6.   Zomer RJ, Xu J, Trabucco A. Version 3 of the global aridity index and potential
evapotranspiration database. *Sci Data*. 2022;**9**:409 https://doi.org/10.1038/s41597-
022-01493-1
7.   Mitchell D, Henschel JR, Hetem RS *et al.* Fog and fauna of the namib desert: Past and
future. *Ecosphere*. 2020;**11** https://doi.org/10.1002/ecs2.2996
8.   Büdel B, Darienko T, Deutschewitz K *et al.* Southern african biological soil crusts are
ubiquitous and highly diverse in drylands, being restricted by rainfall frequency.
Microbial Ecology. 2009;**57**:229-47 https://doi.org/10.1007/s00248-008-9449-9
9.   Stomeo F, Valverde A, Pointing SB *et al.* Hypolithic and soil microbial community
assembly along an aridity gradient in the namib desert. *Extremophiles*. 2013;**17**:329-
37 https://doi.org/10.1007/s00792-013-0519-7
10.   Warren-Rhodes KA, McKay CP, Boyle LN *et al.* Physical ecology of hypolithic
communities in the central namib desert: The role of fog, rain, rock habitat, and light.
J Geophys Res-Biogeo. 2013;**118(4)**:1451-60 https://doi.org/10.1002/jgrg.20117
11.   Ramond JB, Woodborne S, Hall G *et al.* Namib desert primary productivity is driven
by cryptic microbial community n-fixation. *Sci Rep-Uk*. 2018;**8**
https://doi.org/10.1038/s41598-018-25078-4
12.   Scola V, Ramond JB, Frossard A *et al.* Namib desert soil microbial community
diversity, assembly, and function along a natural xeric gradient. *Microb Ecol*.
2017;**75**:193-203 https://doi.org/10.1007/s00248-017-1009-8
13.   Lange OL, Green TGA, Meyer A *et al.* Water relations and carbon dioxide exchange
of epiphytic lichens in the namib fog desert. *Flora*. 2007;**202**:479-87
14.   Hinchliffe G, Bollard-Breen B, Cowan DA *et al.* Advanced photogrammetry to assess
lichen colonization in the hyper-arid namib desert. *Front Microbiol*. 2017;**8**:2083
https://doi.org/10.3389/fmicb.2017.02083
15.   Cowan DA, Cary SC, DiRuggiero J *et al.* 'Follow the water': Microbial water
acquisition in desert soils. *Microorganisms*. 2023;**11**
https://doi.org/10.3390/microorganisms11071670
16.   Frossard A, Ramond JB, Seely M *et al.* Water regime history drives responses of soil
namib desert microbial communities to wetting events. *Sci Rep*. 2015;**5**:12263
https://doi.org/10.1038/srep12263
17.   Stevenson A, Burkhardt J, Cockell CS *et al.* Multiplication of microbes below 0.690
water activity: Implications for terrestrial and extraterrestrial life. *Environmental*
Microbiology. 2015;**17**:257-77 https://doi.org/10.1111/1462-2920.12598
18.   Williams JP, Hallsworth JE. Limits of life in hostile environments: No barriers to
biosphere function? *Environ Microbiol*. 2009;**11**:3292-308
https://doi.org/10.1111/j.1462-2920.2009.02079.x
19.   Bosch J, Varliero G, Hallsworth JE *et al.* Microbial anhydrobiosis. *Environ*
Microbiol. 2021;**23**:6377-90 https://doi.org/10.1111/1462-2920.15699
20.   León-Sobrino C, Ramond JB, Maggs-Kölling G *et al.* Nutrient acquisition, rather than
stress response over diel cycles, drives microbial transcription in a hyper-arid namib

671   desert soil. *Frontiers in Microbiology*. 2019;**10**
672   https://doi.org/10.3389/fmicb.2019.01054

673 21. Maphosa S, Steyn M, Lebre PH *et al.* Rhizosphere bacterial communities of namib
674   desert plant species: Evidence of specialised plant-microbe associations. *Microbiol*
675   *Res*. 2025;**293**:128076 https://doi.org/10.1016/j.micres.2025.128076

676 22. Armstrong A, Valverde A, Ramond JB *et al.* Temporal dynamics of hot desert
677   microbial communities reveal structural and functional responses to water input. *Sci*
678   *Rep-Uk*. 2016;**6**

679 23. Monus BD, Nghalipo EN, Marufu VJ *et al.* Contributions of hypolithic communities
680   to surface soil organic carbon across a hyperarid-to-arid climate gradient. *Geoderma*.
681   2023;**433** https://doi.org/10.1016/j.geoderma.2023.116428

682 24. Blagodatskaya E, Kuzyakov Y. Active microorganisms in soil: Critical review of
683   estimation criteria and approaches. *Soil Biology & Biochemistry*. 2013;**67**:192-211

684 25. Lennon JT, Jones SE. Microbial seed banks: The ecological and evolutionary
685   implications of dormancy. *Nature reviews microbiology*. 2011;**9**:119-30

686 26. Imminger S, Meier DV, Schintlmeister A *et al.* Survival and rapid resuscitation
687   permit limited productivity in desert microbial communities. *Nat Commun*.
688   2024;**15**:3056 https://doi.org/10.1038/s41467-024-46920-6

689 27. Leung PM, Bay SK, Meier DV *et al.* Energetic basis of microbial growth and
690   persistence in desert ecosystems. *mSystems*. 2020;**5**
691   https://doi.org/10.1128/mSystems.00495-19

692 28. Leon-Sobrino C, Ramond JB, Coclet C *et al.* Temporal dynamics of microbial
693   transcription in wetted hyperarid desert soils. *FEMS Microbiol Ecol*. 2024;**100**
694   https://doi.org/10.1093/femsec/fiae009

695 29. Bay SK, Dong X, Bradley JA *et al.* Trace gas oxidizers are widespread and active
696   members of soil microbial communities. *Nat Microbiol*. 2021;**6**:246-56
697   https://doi.org/10.1038/s41564-020-00811-w

698 30. Greening C, Biswas A, Carere CR *et al.* Genomic and metagenomic surveys of
699   hydrogenase distribution indicate h2 is a widely utilised energy source for microbial
700   growth and survival. *ISME J*. 2016;**10**:761-77 https://doi.org/10.1038/ismej.2015.153

701 31. Jordaan K, Lappan R, Dong X *et al.* Hydrogen-oxidizing bacteria are abundant in
702   desert soils and strongly stimulated by hydration. *mSystems*. 2020;**5**
703   https://doi.org/10.1128/mSystems.01131-20

704 32. Lynch RC, Darcy JL, Kane NC *et al.* Metagenomic evidence for metabolism of trace
705   atmospheric gases by high-elevation desert actinobacteria. *Frontiers in Microbiology*.
706   2014;**5** https://doi.org/10.3389/fmicb.2014.00698

707 33. Morita RY. Is h(2) the universal energy source for long-term survival? *Microb Ecol*.
708   1999;**38**:307-20 https://doi.org/10.1007/s002489901002

709 34. Ji M, Greening C, Vanwonterghem I *et al.* Atmospheric trace gases support primary
710   production in antarctic desert surface soil. *Nature*. 2017;**552**:400-03
711   https://doi.org/10.1038/nature25014

712 35. Ortiz M, Leung PM, Shelley G *et al.* Multiple energy sources and metabolic strategies
713   sustain microbial diversity in antarctic desert soils. *Proc Natl Acad Sci U S A*.
714   2021;**118** https://doi.org/10.1073/pnas.2025322118

715 36. Ray AE, Zaugg J, Benaud N *et al.* Atmospheric chemosynthesis is phylogenetically
716   and geographically widespread and contributes significantly to carbon fixation
717   throughout cold deserts. *Isme Journal*. 2022;**16**:2547-60
718   https://doi.org/10.1038/s41396-022-01298-5

37.   Ray AE, Tribbia DZ, Cowan DA *et al.* Clearing the air: Unraveling past and guiding
future research in atmospheric chemosynthesis. *Microbiol Mol Biol Rev*.
2023;**87**:e0004823 https://doi.org/10.1128/mmbr.00048-23
38.   Bay SK, Waite DW, Dong X *et al.* Chemosynthetic and photosynthetic bacteria
contribute differentially to primary production across a steep desert aridity gradient.
ISME J. 2021;**15**:3339-56 https://doi.org/10.1038/s41396-021-01001-0
39.   Naidoo Y, Valverde A, Pierneef RE *et al.* Differences in precipitation regime shape
microbial community composition and functional potential in namib desert soils.
Microbial Ecology. 2022;**83**:689-701 https://doi.org/10.1007/s00248-021-01785-w
40.   Vikram S, Ramond JB, Ortiz M *et al.* Soil fungal diversity and assembly along a xeric
stress gradient in the central namib desert. *Fungal Biol*. 2023;**127**:997-1003
https://doi.org/10.1016/j.funbio.2023.03.001
41.   FAO. *Standard operating procedure for soil moisture content by gravimetric method*,
Rome, 2023.
42.   McNally A. Fldas noah land surface model l4 global monthly 0.1× 0.1 degree (merra-
2 and chirps). Greenbelt, MD, USA.
43.   He SF, Yang HY, Cui XQ. Technical note: Assessing pretreatment approaches for
serial pyrolysis-oxidation analysis of sedimentary organic carbon. *Biogeosciences*.
2025;**22**:6243-53 https://doi.org/10.5194/bg-22-6243-2025
44.   Dragone NB, Diaz MA, Hogg I *et al.* Exploring the boundaries of microbial
habitability in soil. *J Geophys Res-Biogeo*. 2021;**126**:6052-52
45.   Robertson GP. *Standard soil methods for long-term ecological research*, vol. 2:
Oxford University Press, 1999.
46.   Maeda H, Fujimoto C, Haruki Y *et al.* Quantitative real-time pcr using taqman and
sybr green for actinobacillus actinomycetemcomitans, porphyromonas gingivalis,
prevotella intermedia, tetq gene and total bacteria. *FEMS Immunology & Medical*
Microbiology. 2003;**39**:81-86 https://doi.org/10.1016/s0928-8244(03)00224-4
47.   Ray AE, Zhang E, Terauds A *et al.* Soil microbiomes with the genetic capacity for
atmospheric chemosynthesis are widespread across the poles and are associated with
moisture, carbon, and nitrogen limitation. *Front Microbiol*. 2020;**11**:1936
https://doi.org/10.3389/fmicb.2020.01936
48.   Apprill A, McNally S, Parsons R *et al.* Minor revision to v4 region ssu rrna 806r gene
primer greatly increases detection of sar11 bacterioplankton. *Aquatic Microbial*
Ecology. 2015;**75**:129-37
49.   Parada AE, Needham DM, Fuhrman JA. Every base matters: Assessing small subunit
rrna primers for marine microbiomes with mock communities, time series and global
field samples. *Environ Microbiol*. 2016;**18**:1403-14 https://doi.org/10.1111/1462-
2920.13023
50.   Callahan BJ, McMurdie PJ, Rosen MJ *et al.* Dada2: High-resolution sample inference
from illumina amplicon data. *Nat Methods*. 2016;**13**:581-3
https://doi.org/10.1038/nmeth.3869
51.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *2011*. 2011;**17**:3 https://doi.org/10.14806/ej.17.1.200
52.   Wang Q, Garrity GM, Tiedje JM *et al.* Naive bayesian classifier for rapid assignment
of rrna sequences into the new bacterial taxonomy. *Appl Environ Microbiol*.
2007;**73**:5261-7 https://doi.org/10.1128/AEM.00062-07
53.   Quast C, Pruesse E, Yilmaz P *et al.* The silva ribosomal rna gene database project:
Improved data processing and web-based tools. *Nucleic Acids Res*. 2013;**41**:D590-6
https://doi.org/10.1093/nar/gks1219

54. Katoh K, Standley DM. Mafft multiple sequence alignment software version 7:
Improvements in performance and usability. *Molecular Biology and Evolution*.
2013;**30**:772-80 https://doi.org/10.1093/molbev/mst010
55. Price MN, Dehal PS, Arkin AP. Fasttree 2-approximately maximum-likelihood trees
for large alignments. *Plos One*. 2010;**5** https://doi.org/10.1371/journal.pone.0009490
56. Tavaré S. Some probabilistic and statistical problems on the analysis of DNA
sequences. *Lectures on Mathematics in the Life Sciences*. 1986;**17**:57-86
57. Lozupone CA, Hamady M, Kelley ST *et al.* Quantitative and qualitative β diversity
measures lead to different insights into factors that structure microbial communities.
Appl Environ Microb. 2007;**73**:1576-85 https://doi.org/10.1128/Aem.01996-06
58. McMurdie PJ, Holmes S. Phyloseq: An r package for reproducible interactive analysis
and graphics of microbiome census data. *PLoS One*. 2013;**8**:e61217
https://doi.org/10.1371/journal.pone.0061217
59. Oksanen J, Simpson GL, Blanchet FG *et al.* Vegan: Community ecology package
https://vegandevs.github.io/vegan/. 2025.
60. Mallick H, Rahnavard A, McIver LJ Multivariable association in population-scale
meta-omics studies.
61. RStudio Team. Rstudio: Integrated development for r. RStudio, PBC, Boston, MA.
http://www.rstudio.com/. 2020.
62. Wickham H. Reshaping data with the reshape package Journal of Statistical Software.
Version 1.4.4 http://www.jstatsoft.org/v21/i12/. 2007.
63. Wickham H. Stringr: Simple, consistent wrappers for common string operations
https://CRAN.R-project.org/package=stringr. 2023.
64. Wickham H, Hester J, Bryan J. Readr: Read rectangular text data https://CRAN.R-
project.org/package=readr. 2024.
65. Hijmans RJ. Terra: Spatial data analysis https://rspatial.org/. 2024.
66. Hernangómez D. Using the tidyverse with terra objects: The tidyterra package.
Journal of Open Source Software. 2023;**8**:5751 https://doi.org/10.21105/joss.05751
67. Wickham H, François R, Henry L *et al.* Dplyr: A grammar of data manipulation
https://dplyr.tidyverse.org. 2023.
68. Kassambara A. Ggpubr: 'Ggplot2' based publication ready plots
https://rpkgs.datanovia.com/ggpubr/. 2023.
69. Wei T, Simko V. R package 'corrplot': Visualisation of a correlation matrix Version
Version 0.95 https://github.com/taiyun/corrplot. 2024.
70. Sinnott RW. Virtues of the haversine. *Sky Telescope*. 1984;**68**:159-59
71. Hijmans RJ. Geosphere: Spherical trigonometry https://github.com/rspatial/geosphere.
2024.
72. Freedman D, Diaconis P. On the histogram as a density estimator - l2 theory. *Z
Wahrscheinlichkeit*. 1981;**57**:453-76 https://doi.org/Doi 10.1007/Bf01025868
73. Venables WN, Ripley BD. *Modern applied statistics with s*: Springer Science &
Business Media, 2013.
74. Wickham H. Ggplot2: Elegant graphics for data analysis Springer-Verlag New York.
https://ggplot2.tidyverse.org. 2016.
75. Lin Pedersen T. Patchwork: The composer of plots https://patchwork.data-
imaginist.com. 2024.
76. Wright K. Pals: Color palettes, colormaps, and tools to evaluate them Version R
package version 1.10 https://CRAN.R-project.org/package=pals. 2025.
77. Wickham H, Averick M, Bryan J *et al.* Welcome to the tidyverse. Journal of Open
Source Software. 1686.

78.   Wilke CO, Wiernik BM. Ggtext: Improved text rendering support for 'ggplot2'
https://github.com/wilkelab/ggtext. 2025.

79.   Garvin ZK, Abades SR, Trefault N *et al.* Prevalence of trace gas-oxidizing soil
bacteria increases with radial distance from polloquere hot spring within a high-
elevation andean cold desert. *ISME J.* 2024;**18**
https://doi.org/10.1093/ismejo/wrae062

80.   Soom S, Moning SU, Cook GM *et al.* Atp synthesis driven by atmospheric hydrogen
concentrations. *Proc Natl Acad Sci U S A.* 2025;**122**:e2506353122
https://doi.org/10.1073/pnas.2506353122

81.   Meier DV, Imminger S, Gillor O *et al.* Distribution of mixotrophy and desiccation
survival mechanisms across microbial genomes in an arid biological soil crust
community. *Msystems.* 2021;**6**:e00786-20 https://doi.org/10.1128/mSystems.00786-
20

82.   Li SY, Yang SS, Wei XM *et al.* Reduced trace gas oxidizers as a response to organic
carbon availability linked to oligotrophs in desert fertile islands. *The ISME Journal.*
2023;**17**:1257-66 https://doi.org/10.1038/s41396-023-01437-6

83.   Nikitushkin VD, Demina GR, Kaprelyants AS. Rpf proteins are the factors of
reactivation of the dormant forms of actinobacteria. *Biochemistry (Mosc).*
2016;**81**:1719-34 https://doi.org/10.1134/S0006297916130095

84.   Lennon JT, Lehmkuhl BK, Chen L *et al.* Resuscitation-promoting factor (rpf)
terminates dormancy among diverse soil bacteria. *mSystems.* 2025;**0**:e01517-24
https://doi.org/doi:10.1128/msystems.01517-24

85.   Maestre FT, Delgado-Baquerizo M, Jeffries TC *et al.* Increasing aridity reduces soil
microbial diversity and abundance in global drylands. *Proc Natl Acad Sci USA.*
2015;**112**:15684-9 https://doi.org/10.1073/pnas.1516684112

86.   Schulze-Makuch D, Wagner D, Kounaves SP *et al.* Transitory microbial habitat in the
hyperarid atacama desert. *P Natl Acad Sci USA.* 2018;**115**:2670-75
https://doi.org/10.1073/pnas.1714341115

87.   Van Goethem MW, Makhalanyane TP, Cowan DA *et al.* Cyanobacteria and
alphaproteobacteria may facilitate cooperative interactions in niche communities.
Front Microbiol. 2017;**8**:2099 https://doi.org/10.3389/fmicb.2017.02099

88.   Albuquerque L, Johnson MM, Schumann P *et al.* Description of two new
thermophilic species of the genus rubrobacter, rubrobacter calidifluminis sp. Nov.
And rubrobacter naiadicus sp. Nov., and emended description of the genus
rubrobacter and the species rubrobacter bracarensis. *Syst Appl Microbiol.*
2014;**37**:235-43 https://doi.org/10.1016/j.syapm.2014.03.001

89.   Bartholomaus A, Genderjahn S, Mangelsdorf K *et al.* Inside the atacama desert:
Uncovering the living microbiome of an extreme environment. *Appl Environ*
Microbiol. 2024;**90**:e0144324 https://doi.org/10.1128/aem.01443-24

90.   Ortiz M, Leung PM, Shelley G *et al.* A genome compendium reveals diverse
metabolic adaptations of antarctic soil microorganisms. *Preprint.* 2020
https://doi.org/10.1101/2020.08.06.239558

91.   Mashamaite L, Lebre PH, Varliero G *et al.* Microbial diversity in antarctic dry valley
soils across an altitudinal gradient. *Front Microbiol.* 2023;**14**:1203216
https://doi.org/10.3389/fmicb.2023.1203216

92.   Cao W, Xiong Y, Zhao D *et al.* Bryophytes and the symbiotic microorganisms, the
pioneers of vegetation restoration in karst rocky desertification areas in southwestern
china. *Appl Microbiol Biotechnol.* 2020;**104**:873-91 https://doi.org/10.1007/s00253-
019-10235-0

93. Marasco R, Mosqueira MJ, Fusi M *et al.* Rhizosheath microbial community assembly
of sympatric desert speargrasses is independent of the plant host. *Microbiome.*
2018;**6**:215 https://doi.org/10.1186/s40168-018-0597-y
94. Yang Y, Liu WG, Huang TW *et al.* Similarities and differences in the rhizosphere
biota among different ephemeral desert plants in gurbantunggut desert. *Environ Sci*
Eur. 2023;**35** https://doi.org/10.1186/s12302-023-00724-5
95. Jara-Servin A, Silva A, Barajas H *et al.* Root microbiome diversity and structure of
the sonoran desert buffelgrass (pennisetum ciliare l.). *Plos One.* 2023;**18**
https://doi.org/10.1371/journal.pone.0285978
96. Lebre PH, Fouche J, Bosch J *et al.* Effects of fire intensity on soil microbial ecology
in a grassland ecosystem. *Environmental Microbiology.* 2025;**27**
https://doi.org/10.1111/1462-2920.70124
97. Msaddak A, Duran D, Rejili M *et al.* Diverse bacteria affiliated with the genera
microvirga, phyllobacterium, and bradyrhizobium nodulate lupinus micranthus
growing in soils of northern tunisia. *Appl Environ Microbiol.* 2017;**83**
https://doi.org/10.1128/AEM.02820-16
98. Jimenez-Gomez A, Saati-Santamaria Z, Igual JM *et al.* Genome insights into the
novel species microvirga brassicacearum, a rapeseed endophyte with biotechnological
potential. *Microorganisms.* 2019;**7** https://doi.org/10.3390/microorganisms7090354
99. Ardley JK, Parker MA, De Meyer SE *et al.* Microvirga lupini sp. Nov., microvirga
lotononidis sp. Nov. And microvirga zambiensis sp. Nov. Are alphaproteobacterial
root-nodule bacteria that specifically nodulate and fix nitrogen with geographically
and taxonomically separate legume hosts. *Int J Syst Evol Microbiol.* 2012;**62**:2579-88
https://doi.org/10.1099/ijs.0.035097-0
100. Unc A, Maggs-Kölling G, Marais E *et al.* Soil bacterial community associated with
the dioecious acanthosicyos horridus in the namib desert. *Biol Fert Soils.*
2019;**55**:393-403 https://doi.org/10.1007/s00374-019-01358-7
101. Rosenberg E The family chitinophagaceae. In: Rosenberg E, DeLong EF, Lory S *et*
al. (eds.). *The prokaryotes: Other major lineages of bacteria and the archaea*, Berlin,
Heidelberg: Springer Berlin Heidelberg. 493-95. Retreived from
https://doi.org/10.1007/978-3-642-38954-2_137
102. Kim HS, Kim JS, Suh MK *et al.* A novel plant growth-promoting rhizobacterium,
rhizosphaericola mali gen. Nov., sp. Nov., isolated from healthy apple tree soil. *Sci*
Rep-Uk. 2024;**14**:1038 https://doi.org/10.1038/s41598-024-51492-y
103. Li S, Lian WH, Han JR *et al.* Capturing the microbial dark matter in desert soils using
culturomics-based metagenomics and high-resolution analysis. *NPJ Biofilms*
Microbiomes. 2023;**9**:67 https://doi.org/10.1038/s41522-023-00439-8
104. Greening C, Carere CR, Rushton-Green R *et al.* Persistence of the dominant soil
phylumacidobacteriaby trace gas scavenging. *Proceedings of the National Academy of*
Sciences. 2015;**112**:10497-502 https://doi.org/10.1073/pnas.1508385112
105. Feijo Delgado F, Cermak N, Hecht VC *et al.* Intracellular water exchange for
measuring the dry mass, water mass and changes in chemical composition of living
cells. *PLoS One.* 2013;**8**:e67590 https://doi.org/10.1371/journal.pone.0067590

Figure 1: Aridity gradients of the Namib Desert and the C14 sampling transect *A) Xeric zonation of Namibia according to Aridity Index; transect location represented by the black star. B) GPS locations of ten sampling sites spaced 20 km apart along the 200 km C14 transect, located across a steep hyperarid-arid aridity gradient. The Fog zone is marked by the dotted grey line, and the hyperarid-arid boundary is marked by the solid black line. Fog, Hyperarid and Arid zone sites are indicated at circular, square and triangular points, respectively.*

Figure 2: Soil microbial activity and abundances across the Namib Desert C14 transect. *Data points represent biological replicates (n=4 for A, n=3 for B and C) and error bars represent standard error. A) Total 16S rRNA gene copy numbers per site. Total copy number increased significantly (p < 0.01) between each xeric zone and was lowest in the Fog zone. B) Cumulative CO2 respiratory burst 24 h after moisture addition per site. Cumulative respiratory burst increased significantly (p < 0.05) between each xeric zone. B) Intracellular ATP concentrations per site. Average ATP was not significantly (p > 0.05) different between the Fog and Hyperarid zone.*

Figure 3: Atmospheric hydrogen oxidation rates in dry and wetted soil microcosms. *Rates were calculated at atmospheric hydrogen concentration = 530 ppbv and normalised to average 16S rRNA gene copy numbers per site. Significance thresholds were assigned as: $p \leq 0.05 = *$, $p \leq 0.01 = **$, $p \leq 0.001 = ***$, $p \leq 0.0001 = ****$. Data points shown represent n=5 biological replicates. A) Atmospheric H2 oxidation rates in dry soil microcosms from two Fog zone sites. All remaining dry soil microcosms did not oxidise hydrogen to below 530 ppbv after 1,500 hours. B) Atmospheric H2 oxidation rates in wetted microcosms.*

Figure 4: Bacterial and archaeal community composition and drivers across the Namib Desert C14 transect. *A) Relative community composition per site aggregated to phylum level. B) Relative community composition per site aggregated to family level. C) Principal coordinates analysis (PCoA) of microbial community composition based on weighted UniFrac distances. PCo1 and 2 explain 43.27% and 24.24% of total variance, respectively. Xeric zone is indicated by shape and transect sites are coloured according to Aridity Index (Table 1).*

Figure 5: Differentially abundant taxa across xeric zones of a Namib Desert aridity transect, relative to the Fog zone. *Coefficient is a measure of the difference in relative abundance of taxa in Hyperarid and Arid zone sites, relative to Fog zone sites. Negative values indicate greater relative abundance within the Fog zone. Only coefficients with adjusted q-values ≤ 0.20 are shown. Error bars represent standard error. Taxonomic levels are represented as follows: (P) = phylum, (F) = family.*