Intron Retention Controls Localization of lncRNAs *PURPL* and *MALAT1* to Promote Cell Proliferation and Migration

Ioannis Grammatikakis[1], Chosita Norkaew[1], You Jin Song[2], Amit K. Behera[3], Erica C. Pehrsson[4,5], Corrine Corrina R. Hartford[1], Shreya Kordale[3], Rishabh Prasanth[2], Yongmei Zhao[6], Biraj Shrethsa[6], Xiao Ling Li[1], Ravi Kumar[1], Ragini Singh[1], Tayvia Brownmiller[7], Xinyu Wen[4,5], Natasha J. Caplen[7], Pablo Perez-Pinera[8], Kannanganattu V. Prasanth[2*], Thomas Gonatopoulos-Pournatzis[3*], Ashish Lal[1*]

[1]Regulatory RNAs and Cancer Section, Genetics Branch, Center for Cancer Research (CCR), National Cancer Institute (NCI), National Institutes of Health (NIH), Bethesda, Maryland, USA

[2]Department of Cell and Developmental Biology, Cancer Center at Illinois, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA

[3]Functional Transcriptomics Section, RNA Biology Laboratory, NCI, NIH, Frederick, Maryland, USA

[4]OMICS Technology Facility, Bioinformatics, Genetics Branch, CCR, NCI, NIH, Bethesda, Maryland, USA

[5]Advanced Biomedical Computational Science, Frederick National Laboratory for Cancer Research, Frederick, Maryland, USA

[6]Sequencing Facility Bioinformatics Group, Bioinformatics and Computational Science Directorate, Frederick National Laboratory for Cancer Research, NCI, NIH, Frederick, Maryland, USA

[7]Functional Genetics Section, Genetics Branch, CCR, NCI, NIH, Bethesda, Maryland, USA

[8]Department of Bioengineering, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA

24   ***Corresponding authors:** ashish.lal@nih.gov; kumarp@illinois.edu;

25   thomas.gonatopoulos@nih.gov

26   **Classification:** Biological Sciences

27   **Keywords:** U2AF2, lncRNA, *PURPL*, *MALAT1*, intron retention, splicing, CRISPR/Cas9, nuclear

28   speckles, proliferation, cell migration

Abstract

Intron retention (IR) is increasingly recognized as a feature of long noncoding RNAs (lncRNAs),

yet the mechanisms that shape IR in lncRNAs and the functional consequences of this process

remain largely unexplored. To investigate how IR contributes to lncRNA regulation, we performed

a genome-wide screen to identify factors controlling IR in the lncRNA *PURPL*. This approach

uncovered a prominent role for U2AF2, which promotes retention of a specific intron in *PURPL*

through a weak polypyrimidine tract. IR of this intron drives nuclear enrichment of *PURPL* and

enhances cell proliferation, revealing biological relevance. Transcriptome-wide analyses showed

that although U2AF2 broadly supports canonical splicing consistent with its well-established

function in promoting splicing, it also facilitates IR within a distinct subset of RNAs, including

the nuclear speckle–associated lncRNA *MALAT1*. Loss of U2AF2 disrupts *MALAT1* speckle

localization and using *MALAT1* knockout cells reconstituted with wild-type or intron deleted

variants, we identified a single intron critical for *MALAT1*'s speckle localization. Deletion of this

intron from endogenous *MALAT1* impaired speckle localization and reduced cell migration,

phenocopying the loss of *MALAT1*. Together, these findings reveal IR as a key regulatory

mechanism governing lncRNA localization and function and uncover an unexpected role for

U2AF2 in promoting IR within specific lncRNA contexts.

Introduction

Long non-coding RNAs (lncRNAs) are RNA molecules >200 nucleotides (nt) long that lack coding potential. Similar to protein-coding genes, lncRNAs are transcribed by RNA Polymerase II, are 5′-capped, spliced and are often polyadenylated (1). Several characteristics distinguish lncRNAs from protein-coding genes, including higher tissue-specific expression, lower abundance and stability, predominant nuclear localization, and having fewer but longer exons (2-6). One of the most distinguished features of lncRNAs is that they are less efficiently spliced compared to mRNAs (4,7-9). Interestingly, lncRNAs contain introns with much slower intron-excision kinetics compared to protein-coding genes (10). However, the factors that regulate lncRNA splicing and their impact on lncRNA biology remain largely unclear. We previously reported that the lncRNA *PURPL* exhibits prominent splicing defects, including frequent retention of its second intron. Transcriptomic analyses using both long-read and short-read sequencing revealed multiple *PURPL* isoforms, several of which retain intron 2 (11). While *PURPL* is also known to be transcriptionally induced by p53 and to contribute to cancer cell growth through multiple mechanisms, the regulatory factors governing intron 2 splicing and the functional significance of its retention have not been defined. This gap provides an opportunity to uncover broader principles of IR regulation in lncRNAs.

Splicing is a highly regulated process where *cis*- and *trans*-acting factors coordinate proper and efficient spliceosomal recruitment (12). Several consensus sequences in the RNA are crucial for splicing such as the polypyrimidine (Py)-tract, a stretch of pyrimidines close to the 3′ splice site (ss) (13). The splicing factor U2AF1 (U2-auxiliary factor 1) binds to the 3′ ss and U2AF2 binds to the Py-tract. The U2AF heterodimer is recruited to the branch point, and this complex serves as a critical intermediate for the subsequent assembly of the spliceosome and splicing

catalysis (14-18). The vast majority of human genes encode pre-mRNAs that contain introns that

are spliced out and most nascent transcripts undergo alternative splicing, which leads to the

combinatorial joining of exons. Alternative splicing is a major contributor to protein isoform

diversity, enabling the generation of multiple transcript variants from a single gene and frequently

shaping measurable cellular phenotypes (19-22). One of the least studied forms of alternative

splicing is intron retention (IR). Lately, it has drawn more attention since it has an important role

in post-transcriptional gene expression regulation, and it is believed that more than half of human

multi-exonic genes are affected by IR (23). IR frequently leads to nuclear detention of transcripts

and eventually degradation which may serve as a defense mechanism of the cell against deleterious

products. The extra sequence added by the retained intron may change the open reading frame

(ORF) of protein-coding genes and lead to nonsense-mediated decay (NMD) (24-26) or initiate

the use of an upstream ORF (27,28). In addition, they can contain binding sites for microRNAs

(miRNAs) or RNA-binding proteins (RBPs), or they may form structural changes that impact the

function of the transcript (29). Interestingly, IR has been associated with several diseases such as

cancer where tumor cells show higher levels of IR compared to normal tissues (30). Studies on IR

have shown that retained introns have been associated with weaker (non-consensus) splice sites,

higher GC content, various lengths of introns, and higher intronic sequence conservation (23,31).

lncRNAs have been shown to participate in splicing regulation, with *MALAT1* as the most

prominent example. *MALAT1* is a highly abundant nuclear lncRNA which has oncogenic activity

and is one of the first lncRNAs associated with cancer (32-34). It also has the ability to regulate

alternative splicing because it resides in the nuclear speckles and can interact with several SR

(Serine/Arginine rich) proteins to regulate their phosphorylation state and distribution between

transcription sites and speckles (33,35-37). Nuclear speckles are sub-nuclear compartments that

act as hubs in coordinating transcription, RNA processing and mRNA export (38). Several RBPs

reside in nuclear speckles such as the protein SON which is a key component and marker of nuclear

speckles (39). However, *MALAT1* itself is not essential for the structural integrity of nuclear

speckles (40,41). Nevertheless, it remains unclear whether *MALAT1* undergoes alternative splicing

and how this process impacts its localization to nuclear speckles.

Here, we show that the intron 2-retaining *PURPL* isoform, which localizes to the nucleus,

has a shorter half-life than the spliced isoform and promotes cell proliferation. To identify

regulatory factors controlling the splicing of this functionally relevant intron, we utilized our

recently developed CRISPR-based genome-wide screening strategy using a pooled guide RNA

library with a *PURPL* intron 2 minigene reporter containing its flanking exons. Unexpectedly, this

approach revealed a role of the splicing activator U2AF2 in promoting *PURPL* IR. U2AF2 was

further found to promote a broader intron retention program that includes two introns within

MALAT1. Notably, depletion of U2AF2 resulted in *MALAT1* exclusion from nuclear speckles.

MALAT1 knockout cells exogenously expressing *MALAT1* truncation mutants lacking specific

introns identified one intron essential for the nuclear speckle association of the *MALAT1* transcript.

Breast cancer cells where this intron is deleted showed reduced cell migration demonstrating a

functional role of the intronic sequences. These findings provide insights into the splicing

regulation of *PURPL* and *MALAT1*, uncovering a previously unrecognized functional role of intron

retention in lncRNAs.

Results

A CRISPR-based screen identifies U2AF2 as an activator of *PURPL* intron 2 retention

We previously identified *PURPL* as a p53-regulated lncRNA that promotes the growth of

colorectal and liver cancer cells (11,42). In those studies, we reported that *PURPL* undergoes

alternative splicing, with the ~1.9 kb intron 2 either retained or spliced in the mature transcript

(11). To identify regulators of *PURPL* intron 2 retention, we applied our recently developed

screening platform, CRISPR-based identification of Regulators of Alternative Splicing with

Phenotypic Sequencing (CRASP-seq) (43). This approach utilizes a genome-wide knockout

lentiviral library targeting ~18,500 protein-coding genes using a dual hybrid guide (hg)RNA

system that combines *Streptococcus pyogenes* (*Sp*)Cas9 and optimized *Acidaminococcus sp.*

(op)Cas12a nucleases. (22,44-46). The library also includes ~2,400 pairs and ~500 curated genetic

interaction pairs, allowing combinatorial perturbations from a single construct. Each gene or gene

pair is targeted by four independent hgRNAs alongside ~500 intergenic and non-targeting controls,

totaling ~ 91,000 hgRNAs. The lentiviral library incorporates a doxycycline-inducible *PURPL* IR

reporter comprised of the full native intron 2 sequences (~1.9 kb) flanked by the adjacent exons 2

(161 bp) and 3 (163 bp) (Figure 1A).

We performed the CRASP-screen in two cell lines: HAP1 and RPE1. Following puromycin

selection for 48 h and doxycycline induction for an additional 24 h, RNA was extracted, and poly-

A RNA was purified using oligo(dT) beads to exclude transcripts still undergoing processing.

cDNA was synthesized using customized primers containing unique molecular identifiers (UMIs),

facilitating the removal of PCR duplicates during downstream analysis. The synthesized cDNA

was then utilized to construct Illumina sequencing libraries for paired-end sequencing. This

sequencing approach enables simultaneous quantification of transcripts that either retain or splice

out an intron of interest from one read, while the corresponding hgRNA sequences were captured

in the other read to pinpoint specific genetic perturbations. By directly linking *PURPL* splicing

outcomes with genetic modifications introduced by each hgRNA in the library, this platform

allowed the genome-wide identification of *PURPL* splicing regulators (Figure 1B).

We employed an analytical pipeline that prioritized candidate genes based on multiple

hgRNAs showing significant deviations from intergenic controls, with consistent identification

across two technical replicates. This analysis identified 62 genes that influence *PURPL* intron 2

splicing, the majority of which act as suppressors of intron 2 splicing (Figure 1C and Table S1).

Gene ontology analysis of biological processes revealed a strong enrichment for terms related to

mRNA processing and RNA splicing. Additionally, analysis of cellular components indicated that

the identified hits are predominantly associated with Spliceosomal Complex, including the U2-

type Spliceosomal Complex and the Catalytic step 2 Spliceosome (Figure 1D). These findings

validate the effectiveness of our screening approach.

Consistent with the endogenous *PURPL* transcript isoforms, the basal levels of IR of the

reporter were very high, with the majority of amplified transcripts corresponding to intron-

containing isoforms, as expected based on the intensity of the Illumina libraries before sequencing

(Figure S1A). Thus, the detection of genes that suppress IR might be below the detection levels of

this screen. Therefore, we focused on the genes that promote intron retention. By calculating the

change in Percent Intron Retention ($\Delta PIR = PIR_{guide} - PIR_{intergenic}$) across the two cell lines, we

identified 5 unique genes as regulators of *PURPL* intron 2 retention, all of which act to suppress

intron 2 splicing (Figure 1E and Table S1). This highly significant overlap between the two cell

lines (Figure 1E; odds ratio = 204.4, p=2.81xe^{-10}; Fisher's exact test) indicated a shared regulatory

mechanism governing *PURPL* IR, although cell-type-specific effects may also contribute.

Surprisingly, the gene with the strongest impact based on the average ΔPIR metric and

identified as a hit in both cell lines, was U2AF2 (Figure 1F). This was followed by Small Nuclear

Ribonucleoprotein U1 Subunit 70 (SNRNP70), a core component of the U1 snRNP, and

Elongation Factor Tu GTP Binding Domain, which is associated with U5 snRNP (Figure 1F). Our

finding that U2AF2 promotes IR of *PURPL* intron 2 was unexpected because U2AF2 is a

component of the spliceosome machinery, and its canonical function is to promote intron removal

(14,18).

U2AF2 binds to the *PURPL* intron 2 transcript and promotes intron retention

U2AF2, as a heterodimer with U2AF1, binds to the Py-tract and the 3′ ss, respectively, to promote

splicing (14,17). To elucidate its role in *PURPL* splicing, we investigated whether U2AF2 binds

to intron 2. Enhanced Crosslinking and Immunoprecipitation sequencing (eCLIP-seq) (47), from

the ENCODE project (48,49) from HepG2 cells revealed multiple U2AF2 and U2AF1 binding

sites not only proximal to the 3′ ss of *PURPL* intron 2 but also broadly distributed across intron 2

(Figure 2A). Compared to eCLIP data for other RBPs, U2AF1 and U2AF2 showed the highest

density of binding sites and the most prominent peaks. Specific binding of U2AF2 to intron 2 was

confirmed by performing RNA immunoprecipitation (RNA-IP) using a U2AF2-specific antibody

and primer pairs designed to detect *PURPL* transcripts. U2AF2 was found to interact with

transcripts containing *PURPL* intron 2, but not with the spliced *PURPL* transcripts (Figure 2B and

S1B).

To functionally validate the role of U2AF2 in *PURPL* intron 2 retention, we performed

RT-PCR assays from HAP1 and RPE1 cells transduced with *PURPL* minigene reporter constructs

and control or *U2AF2*-targeting sgRNAs. In both cell lines, we observed a striking decrease in

intron 2 retention upon *U2AF2* depletion as compared to control cells expressing sgRNAs targeting

intergenic regions (Figure 2C). As an alternative approach, we knocked down *U2AF2* using two

independent siRNAs and conducted semi-quantitative RT-PCR assays from SKHEP1, HepG2, and

normal human diploid fibroblasts W138 cells, using a primer triplet (Figures 2D-2F; Figures S2A

and S2B). In all tested cell lines, U2AF2 depletion increased *PURPL* intron 2 splicing, from ~20%

to ~40-60%, more than doubling the abundance of the spliced isoform. Given that U2AF2

functions as a heterodimer with U2AF1, we also knocked down *U2AF1* in SKHEP1 cells and

observed an effect similar to *U2AF2* knockdown on *PURPL* intron 2 retention regulation (Figure

S2C). We expected that U2AF1 would also come as one of the top hits in the screen but *U2AF1*

sgRNAs were not included in the library due to high off-target scores of the guides. These results

demonstrate that the U2AF complex promotes *PURPL* intron 2 retention and that both U2AF1 and

U2AF2 promote intron retention in *PURPL*.

The intron-retained *PURPL* transcript is unstable and localizes to the nucleus

We previously observed an enrichment of the intron 2-retaining *PURPL* transcript in the nucleus

in liver cancer cells (11). To validate these findings, we conducted single molecule RNA-FISH in

HCT116 cells using probes hybridizing to *PURPL* intron 2 (Figure 3A). Because *PURPL* is

upregulated in response to DNA damage (42), we treated the cells with hydroxyurea for 24 h to

induce *PURPL* expression. RNA-FISH revealed that the *PURPL* intron 2 containing transcript was

exclusively localized to the nucleus. To assay for subnuclear compartmentalization, we also

performed RNA-FISH to detect *MALAT1*, which resides in nuclear speckles (36). Intron 2-retained

PURPL transcripts did not colocalize with *MALAT1* indicating that *PURPL* is excluded from

nuclear speckles (Figure 3A). We further validated this result by conducting nuclear/cytoplasmic

fractionation assays followed by RT-qPCR and found that *PURPL* intron 2-containing transcripts

were almost exclusively nuclear (Figure S3A).

Intron retention can affect transcript stability, with longer RNA species often being more

vulnerable to degradation. To test if *PURPL* intron 2 effects transcript turnover, we conducted

RNA stability assays using the transcription inhibitor Actinomycin D (ActD) and observed that

the intron 2-retained *PURPL* transcript was less stable and had a significantly lower half-life than

the *PURPL* transcript in which intron 2 was spliced out (Figure 3B). To determine the functional

impact of *PURPL* intron 2 retention, we used CRISPRi to generate *PURPL*-depleted SKHEP1

cells by targeting 3 independent sgRNAs to the p53-response element of the *PURPL* promoter that

we had used in our previous study (11). We observed >80% reduction in the levels of endogenous

PURPL in these experiments (Figure 3C). We next stably transduced these *PURPL*-depleted cells

with a lentivirus to express exogenous *PURPL* transcript specifically containing intron 2 under the

control of a doxycycline (doxy)-inducible promoter and observed ~2-3-fold induction of the

exogenous *PURPL* intron 2 containing transcript (Figure 3D). Because we and others have

previously shown that *PURPL* is overexpressed in some cancers and *PURPL* promotes

proliferation (11,42,50,51), we next performed cell proliferation assays from the doxy-inducible

cells. Our results showed that cells expressing intron 2-contaning *PURPL* proliferate faster (~2-

fold) as compared to the no-doxy cells after 6 days of doxy treatment (Figure 2E). Induction of

intron 2-contaning *PURPL* transcripts did not have any effect on live cell percentage which

remained at ~100% (Figure S3B). Importantly, as a negative control, there was no effect of doxy

treatment on the proliferation of parental SKHEP1 cells (Figure S3C). To confirm that

proliferation increase induced by exogenous *PURPL* transcript containing intron 2 was not due to

the spliced isoform, we performed semi-quantitative RT-PCR. As shown in Figure 3F, doxy

treatment resulted exclusively in the detection of the intron-retained *PURPL* transcript. These data

suggest that intron 2 retention modulates *PURPL* function by sequestering the transcript in the

nucleus and reducing its stability, while simultaneously promoting cell proliferation, highlighting

a critical regulatory role for intron 2.

A weak polypyrimidine tract is critical for *PURPL* intron 2 retention

U2AF2 binds to the Py-tract to initiate spliceosomal assembly leading to splicing and intron

removal (14). Suboptimal splicing efficiency may be the result of multiple factors, including splice

site strength and the Py-tract sequence composition. The Py-tract is a stretch of 15-20 pyrimidines

(mostly Us) immediately upstream of the 3′ ss. We analyzed the Py-tract of *PURPL* intron 2 and

noticed that it deviates from the consensus, containing numerous purines (Figure S4A). To

investigate whether U2AF2 regulation on intron 2 retention relies on the Py-tract sequence, we

used the *PURPL* intron 2 minigene reporter system (Figure 1A). We mutated the Py-tract,

replacing it either with a strong Py-tract from *PURPL* intron 1 or with a stretch of Us, simulating

a strong Py-tract. *U2AF2* was knocked down in SKHEP1 cells stably expressing the mutated Py-

tract constructs and RT-PCR was conducted to monitor intron 2 splicing. Interestingly, the basal

levels of intron retention were very different with different mutants; ~90% in intron 2 WT (wild-

type), ~30% in *PURPL* intron 1 and ~20% in U-rich construct (Figure S4B and S4C). This data

demonstrates that the weak Py-tract is responsible for the inefficient intron 2 removal in *PURPL*

(Figure S4C). *U2AF2* knockdown resulted in modest increase in splicing of the mutant constructs

as well. Because the minigene reporter contained the mutant Py-tract along with the remaining

endogenous sequences from *PURPL* intron 2, the modest increase in splicing observed for the

mutants after *U2AF2* knockdown may reflect U2AF2 inhibitory binding at additional sites within

PURPL intron 2 beyond the Py-tract (Figure 2A). These data indicate that the endogenous intron

2 Py-tract in *PURPL* is a weak Py-tract that controls the high intron retention observed in *PURPL*.

U2AF2 directly regulates multiple IR events including *MALAT1*

To determine if U2AF2 promotes IR in additional genes, we next performed total RNA-seq and

PacBio Iso-seq from SKHEP1 cells transfected with siCTRL or siU2AF2. Consistent with our

data, *U2AF2* knockdown resulted in a marked increase in the splicing efficiency of *PURPL* intron

(Figures S5A and S5B). Additionally, *U2AF2* depletion increased overall *PURPL* expression, with increased read coverage across all exons (Figures S5A and S5B; Tables S3 and S4). To identify IR events regulated by U2AF2 at a transcriptome-wide level, we utilized IRFinder, an algorithm specifically designed to calculate changes in IR (52). This analysis revealed ~1000 statistically significant IR events regulated by U2AF2. For majority of these events (732), there was an increase in the IR ratio (Intronic abundance/(Intronic abundance + exonic abundance)) after *U2AF2* depletion, indicating that U2AF2 predominantly promotes splicing, as expected. However, in 269 events, *U2AF2* knockdown resulted in decreased IR, indicating that U2AF2 promotes IR in a subset of genes (Figure 4A and Table S2). We next analyzed ENCODE U2AF2 eCLIP-seq data from HepG2 cells to determine if U2AF2 directly binds to the regulated transcripts. U2AF2 binding was detected in 96 out of the 269 IR events supporting a direct role for U2AF2 in promoting IR (Figure S5C and Table S2).

To validate these findings, we selected three of the identified introns from *MDM1*, *RETREG2*, and *TMEM41B* (Figure S6A). We observed multiple U2AF2 binding peaks within the regulated introns from *MDM1* (Figure S6B). We next assessed their splicing in SKHEP1 cells following *U2AF2* knockdown using two independent siRNAs. In all three cases, *U2AF2* depletion significantly enhanced intron removal (Figures S6C-S6E). IRFinder validated that *PURPL* intron 2 was one of the events showing a significant decrease (~2-fold) in IR ratio upon *U2AF2* depletion (Figure 4B).

Among the lncRNAs in which IR is promoted by U2AF2, was the lncRNA *MALAT1*. IRFinder identified two introns in *MALAT1* that have weak Py-tract sequences similar to *PURPL* intron 2 (Figure S7A-S7C). For both introns, *U2AF2* knockdown resulted in a significant decrease

(~2-fold) in the IR ratio compared to siCTRL (Figures 4A and 4B). This result was confirmed by

RT-PCR from HCT116 and SKHEP1 cells upon *U2AF2* knockdown (Figures 4C-4F).

Intron 2 of *MALAT1* drives its localization to nuclear speckles

MALAT1 regulates splicing by interacting with splicing factors in nuclear speckles.(36) To

determine the impact of IR on *MALAT1* localization to nuclear speckles, we performed RNA-FISH

for *MALAT1* upon *U2AF2* depletion in SKHEP1 and HCT116 cells. Remarkably, *U2AF2*

knockdown resulted in a significant decrease in the ratio of *MALAT1* signal intensity in speckles

relative to nucleoplasm (Figures 5A and 5B; Figures S8A and S8B). U2AF2 depletion did not

affect total *MALAT1* levels as observed in our RNA-seq (Table S3). This reduction in *MALAT1*

levels in nuclear speckles did not compromise the structural integrity of nuclear speckles as

indicated by immunostaining for the RNA-binding protein SON, a well-established nuclear

speckle marker (39) (Figures 5A and S8A) and is consistent with previous reports showing that

MALAT1 is not essential for the integrity of nuclear speckles (40,41). These data suggest that

U2AF2 plays an important role in promoting *MALAT1* localization to nuclear speckles.

To determine whether it is the intron retention in the transcript *per se* which directs

MALAT1 localization or if U2AF2 indirectly regulates its localization, we assessed the nuclear

speckle localization of wild-type (WT) and intron-deleted *MALAT1* transcripts. To do this, we

used CRISPR/Cas9 to generate two *MALAT1* knockout (KO) clones in HCT116 by inserting ~100

bp downstream of the transcription start site, a puromycin resistance gene in one allele and a

hygromycin resistance gene in the other allele. In both KO clones, >95% reduction in *MALAT1*

RNA levels was observed as compared to a wild-type clone (Figure S9A).

Next, we transduced the *MALAT1* KO clones with an empty vector or doxycycline-

inducible lentiviral constructs that exogenously express full-length *MALAT1*-WT in which both

introns are intact, intron 1 deleted *MALAT1* (del-1), intron 2 deleted *MALAT1* (del-2) or both

introns were deleted (del-1+2). After 48 h of doxy-treatment to induce *MALAT1* expression, which

was confirmed with RT-qPCR (Figure S9B), RNA-FISH combined with immunofluorescence for

the nuclear speckle marker SON, revealed distinct localization patterns. As expected, exogenously

expressed *MALAT1*-WT predominantly (~80%) localized to nuclear speckles (Figures 6A and

S10A). Similarly, *MALAT1*-del-1 also predominantly (~80%) localized to nuclear speckles. In

contrast, *MALAT1* lacking intron 2 (del-2) or both introns (del-1+2) exhibited a marked reduction

(<15%) in nuclear speckle localization and instead displayed diffused localization in the nucleus

(Figures 6A and S10A). Importantly, this diffused localization upon intron 2 deletion was observed

in both *MALAT1* knockout clones and was further confirmed through quantitative analyses

(Figures 6B and S10B).

Deletion of *MALAT1* Intron 2 using CRISPR/Cas9 impairs cell migration

MALAT1 has been shown to promote cell migration and metastasis (32,53) and its expression has

been correlated with high tumor progression and metastasis in various tumors (54). We therefore

tested whether the intronic sequences that regulate *MALAT1* nuclear speckle localization are

related to the ability of *MALAT1* to promote cell migration. To investigate this, we used the

metastatic breast cancer cell line MDA-MB-231 and first performed RNA-FISH for *MALAT1* after

knocking down *U2AF2*. Similar to SKHEP1 and HCT116 cells, *U2AF2* knockdown resulted in

decreased localization of *MALAT1* to speckles in MDA-MB-231 cells (Figures 7A and 7B). We

next utilized CRISPR/Cas9 to generate KO clones in MDA-MB-231 cells where either the whole

MALAT1 locus or intron 2 of *MALAT1* were deleted. *MALAT1* depletion in the *MALAT1* KO clone

was confirmed with RT-qPCR (Figure S11A). RNA-FISH in 2 separate clones showed that intron

2 deletion reduced *MALAT1* localization to nuclear speckles compared to the WT clone (Figures

7C-7E). In the intron-2 deletion clones, the transcripts containing intron 2 were, as expected,

undetected with primers specifically amplifying intron 2. Interestingly, there was a modest

increase in total *MALAT1* expression suggesting a compensatory mechanism for the loss of intron

2 (Figure S11B). Functionally, when we tested the migration potential of these cells in transwell

cell migration assays, we found significantly reduced cell migration in the intron 2-deleted clones,

phenocopying the complete loss of *MALAT1* locus (Figures 7F and S11C). These data suggest that

the retention of intron 2 is essential for *MALAT1* localization to nuclear speckles and it plays a key

role in promoting cell migration. Together, our data uncovers a U2AF2-driven intron retention

program in lncRNAs that regulates cell proliferation and migration.

Discussion

The goal of this study was to examine the causes and consequences of IR in lncRNAs, focusing

on *PURPL* and *MALAT1*. Intron retention is a regulated process that shapes gene expression

networks through mechanisms such as NMD and influences alternative protein isoform generation

during differentiation, development, and disease (25,55). We previously identified a p53-induced

lncRNA, *PURPL*, which exhibits growth-promoting functions, and that intron 2 of *PURPL* is

retained in multiple cell lines (11). To accomplish our goal, we employed an innovative RNA-seq

coupled genome-wide CRISPR screen on a reporter containing *PURPL* intron 2 and unexpectedly

identified U2AF2 as a major inhibitor of intron 2 removal in *PURPL*. U2AF2 forms a heterodimer

with U2AF1, binding to the Py-tract and the 3′ ss, respectively, to promote splicing by recruiting

core spliceosomal components (17,18,56). Therefore, given the established role of U2AF2 as a

critical positive regulator of splicing catalysis, we were surprised that U2AF2 inhibits the removal

of *PURPL* intron 2. While U2AF1 was not identified as a hit in our screen, follow-up experiments

revealed that U2AF1 also suppresses *PURPL* intron 2 removal, suggesting a previously

unrecognized role for the U2AF complex as a negative regulator *PURPL* IR.

Beyond IR in *PURPL*, we sought to determine whether U2AF2 promotes IR in other

transcripts. Using IR Finder (52), we identified other genes where U2AF2 promoted IR. One of

the genes that drew our attention was the lncRNA *MALAT1* which showed two intron retention

events regulated and bound by U2AF2. These introns contain weak splice sites suggesting that this

may be the reason that they are regulated, and they are not constitutively spliced. Because *MALAT1*

interacts with splicing factors within nuclear speckles (36), we examined whether U2AF2 affects

MALAT1 localization to speckles. Indeed, our results establish *MALAT1* intron 2 as a critical

determinant of its nuclear speckle localization. Future studies involving mapping of the intronic

sequences and targeted mutagenesis of the splice sites within *MALAT1* intron 2 could help

distinguish whether speckle localization of *MALAT1* is inhibited by the splicing process itself or

depends on a discrete element within the intron.

For *PURPL*, previous work established its context-dependent oncogenic activity (11,42),

but those studies depleted all transcripts from the *PURPL* locus despite the presence of multiple

alternatively spliced isoforms. In the current study, overexpression of the intron 2-retaining

PURPL isoform in CRISPRi-mediated *PURPL*-depleted cells increased proliferation,

demonstrating isoform-specific, growth-promoting function. Our findings reveal that intron

retention in both *MALAT1* and *PURPL* is a regulatory mechanism with direct consequences on

lncRNA localization and cancer-related phenotypes.

The surprising finding of U2AF2 promoted IR promoted us to determine the underlying

mechanism. We first asked whether U2AF2 influences transcription initiation or elongation at the

PURPL locus, given the tight coupling between transcription and splicing (57,58). Transcription

kinetics can modulate IR (23,59) and lncRNAs often lack proximal RNA-pol II phosphorylation

over 5′ ss (10). We therefore assayed for PolII-pSer5 (transcription initiation) and PolII-pSer2

(transcription elongation) occupancy after U2AF2 knockdown using CUT&RUN-Seq but there

was no significant difference around intron 2 or elsewhere across the *PURPL* locus (data not

shown). We also examined the chromatin mark H3K36me3 which has been linked to IR (60) and

again observed no effect after U2AF2 depletion (data not shown).

U2AF2 functions by binding to the Py-tract, and the Py-tract of *PURPL* intron 2 and

MALAT1 introns 1 and 2 deviate from the consensus, containing a mix of pyrimidines and purines.

These atypical sequences likely contribute to splicing inhibition as revealed by our data from the

mini-gene reporter for *PURPL*. Given that U2AF1 and U2AF2 bind to multiple intronic sites, a

plausible model is that their docking obstructs subsequent spliceosomal assembly. lncRNAs

generally undergo less efficient splicing and consequently, they have higher levels of IR compared

to protein coding genes (8,10). A study showed that lncRNA splicing strength correlates with 5′ ss

strength and lncRNA Py-tract has higher thymidine content compared to protein coding genes

(61). The reduced evolutionary pressure to maintain efficient splicing as they are not used as

templates for protein synthesis, likely contributes to this pattern. This is also supported by their

predominant nuclear localization (3,4).

Our data shows that *PURPL* transcripts can be either nuclear or cytoplasmic, and U2AF2

regulation of IR offers a mechanism of nuclear detention. Similar to this notion, the lncRNA *TUG1*

remains in the nucleus due to intron retention (62), and the mouse lncRNA *pCharme* undergoes

an IR event which renders it chromatin-bound through its association with MATR3 and PTBP1

(63). Another aspect of the effects of IR lies in the turnover of the transcripts where additional

sequences added by the introns may cause lower transcript stability. In support of this, nuclear

lncRNAs have been shown to be less stable than cytoplasmic ones (5), and *PURPL* also follows

this rule. Notably, the human lncRNA *hFAST* is cytoplasmic and promotes pluripotency through

WNT signaling, but the mouse homolog does not because of its nuclear localization (64).

Introns have been proposed to act as docking elements that tether lncRNAs to chromatin

(4,63), and a recent study showed that nuclear speckles are associated with distinct classes of

retained introns (65). This raises the question of what makes intron 2 important for speckle

localization of *MALAT1*. One possibility is that this region recruits specific RBPs or adopts a

structural configuration required for speckle association. Our data further indicates that intron 2

retention contributes to *MALAT1's* role in cell migration, potentially because speckle localization

is necessary to sustain this phenotype. Future studies are warranted to dig deeper into the

underlying mechanism and to determine how intron retention shapes the function of *PURPL* and

MALAT1.

Collectively, our findings reveal an expected function of U2AF2 in promoting intron

retention that governs the subcellular localization and function of the lncRNAs *PURPL* and

MALAT1. As intron retention continues to emerge as a regulated form of alternative splicing, the

evidence presented here expands the framework for the mechanisms and biological consequences

of intron retention.

Materials and Methods

Cell culture and treatment

HAP1, HCT116, HEK293T, HepG2, RPE1, SKHEP1, U2OS, WI38, and MDA-MB-231 cells

were purchased from ATCC and maintained in Dulbecco's Modified Eagle Medium (DMEM)

(Gibco) supplemented with 10% (v/v) fetal bovine serum (Gibco) and 1% (v/v) PenStrep (Gibco)

in a 5% CO_2 atmosphere at 37°C. All cell lines were routinely confirmed to be free of mycoplasma

using Venor GeM Mycoplasma Detection Kit (Millipore Sigma-Aldrich).

436 CRASP-Seq *PURPL* screen

The *PURPL* minigene reporter, composed of exons 2 and 3 with the full native intronic sequence,

was cloned into the CRASP-Seq genome-wide library as described previously (43). Reporter

fragments were synthesized by TWIST Biosciences and assembled into the Eco32I-digested

CRASP-Seq knockout library using NEBuilder (46). Library coverage (>250-fold) was maintained

through parallel cloning reactions and electroporation into Endura competent cells followed by

plasmid amplification and extraction.

Lentivirus was produced following established protocols (46). The pooled library was used

to infect HAP1 and RPE1 cells expressing *Sp*Cas9 and opCas12a nucleases (46). at a multiplicity

of infection (MOI) of ~0.2. After puromycin selection (2 µg/mL for 48 hours), cells were pooled,

induced with doxycycline to activate reporter expression and harvested 24 hours later for RNA

analysis.

Total RNA was extracted, poly(A)+ RNA isolated, and cDNA synthesized using a U6-

annealing primer. Spliced and intron-retaining reporter transcripts were amplified using exon- or

intron-specific primers incorporating Illumina barcodes. Libraries were sequenced on the NovaSeq

6000 platform with a 25% PhiX spike-in and a 300-cycle kit. The sequencing strategy consisted

of the following configuration: Read 1 (210 cycles), Index Read 1 (8 cycles), Index Read 2 (8

cycles) and Read 2 (85 cycles). Each sample yielded over 500 million total reads.

454 CRASP-seq data analysis

CRASP-Seq data were processed using a custom pipeline (43). Guides, cell barcodes, and UMIs

were retrieved using STAR, and alignment were managed with SAMTools. Reads containing

complete hybrid guide, barcodes, UMI, and splicing outcomes were deduplicated prior to analysis.

Splicing outcomes were quantified as percent intron retention (PIR): $PIR = [IE / (IE + EE)] \times 100$,

where IE represents intron retention reads, and EE represents exon-exon junction reads.

Differential PIR (ΔPIR) was calculated as: $\Delta PIR = PIR_{guide} - PIR_{intergenic}$

The ΔPIR for each gene and for each guide separately is shown in Table S1. Guides with

fewer than 10 reads were excluded. A ΔPIR threshold corresponding to 10% FDR (based on

intergenic controls) defined significant hits. At the gene level, hits required $FPKM \geq 0.1$, at least

two qualifying guides, and $\geq 50\%$ of guides exceeding the ΔPIR threshold in both technical

replicates. Genes identified through combinatorial targeting that met these criteria were included

as hits (Table S1). The heatmap and Venn diagram in Figure 1C and 1E display single-gene hits

as well as collapsed gene hits resulting from combinatorial targeting.

eCLIP-seq data and functional enrichment analysis

eCLIP-seq data for U2AF2, U2AF1, PTBP1, PRPF8, and SRSF1 were obtained from the

ENCODE database (encodeproject.org) with the following identifiers: ENCFF989JBA (CTRL),

ENCFF566CFJ (CTRL), ENCFF536AFD (U2AF2), ENCFF913WRH (U2AF2), ENCFF159SPZ

(U2AF2), ENCFF368XEI (U2AF2), ENCFF542NOA (U2AF1), ENCFF651VRQ (U2AF1),

ENCFF363UDO (PTBP1), ENCFF044RKN (PRPF8), and ENCFF327NVE (SRSF1). For

visualization of RNA-seq data, the Integrated Genome Viewer (IGV) version 17 was used. The

functional enrichment analysis was performed using g:Profiler (66) and the results are shown in

the 'gProfiler_Results' tab in Table S1.

siRNA transfections

For siRNA transfections, we used AllStars Negative Control siRNA (QIAGEN). U2AF2 siRNAs

were purchased from Dharmacon (J-012380-18 and J-012380-19). U2AF1 siRNA was also

purchased from Dharmacon (J-012325-10). Cells were reverse transfected with siRNA at a final

concentration of 20 nM using Lipofectamine RNAiMAX (Invitrogen) in Opti-MEM I Reduced

Serum Medium (Gibco) according to the manufacturer's protocol. SKHEP1 cells were transfected

for 72 hr before RNA harvesting. U2OS, HepG2 and WI38 cells were transfected for two rounds

of 48 h + 72 h or 72 h + 72 h. For RNA FISH, SKHEP1, HCT116, and MDA-MB-231 cells were

transfected for 72 hr. SKHEP1 expressing the *PURPL* Py-tract mutants were transfected for two

rounds of 48 h and 24 h before harvesting, the cells were treated with 1 µg/mL doxycycline to

induce the transgene expression. For RNA-seq, the cells were transfected for 72 h before harvested.

For RNA stability assay, the cells were reseeded in 12-well plates 48 h after siRNA transfection

and 24 h later they were treated with Act D for the indicated time points at 5 µg/mL before RNA

extraction and subsequent assays. For RNA-IP, the cells were transfected for 72 h before harvested

in 10 cm-plates.

RNA extraction, RT-qPCR, semi-quantitative RT-PCR, and splicing gels

Total RNA was extracted using TRIzol (Invitrogen) according to the manufacturer's protocol.

cDNA was then made using iScript™ Reverse Transcription Supermix (Bio-Rad). For real-time

qPCR, all reactions were carried out on StepOnePlus real-time PCR System (Applied Biosystems)

using FastStart SYBR Green Master Mix (Millipore Sigma). *GAPDH* mRNA was used to

normalize expression except for RNA-IPs in Figure 2B where *18S* was used for normalization, and

the relative expression/enrichment of RNAs was calculated using the $2^{-\Delta\Delta Ct}$ method. RT-qPCR

primer sequences for each gene are indicated in Table S3. For splicing gels, cDNA was used to

amplify the region flanking the retained intron with primers in the flanking exons with the addition

of one primer inside the intron as indicated in Figure S1B. In the case of *MALAT1*, primers in the

flanking exons were utilized due to the short length of introns. For the assay of the effect of U2AF2

on the Py-tract mutants of *PURPL*, primers on the transgene were used instead of the endogenous

PURPL exon 2 and 3. The sequences of the primers can be found in Table S5.

RNA-IPs

For RNA-IP, two 10 cm-plates of SKHEP1 cells were lysed with RIPA buffer (Thermo Scientific)

for 10 min on ice before spin down for 10 min at 4°C. The lysate was incubated with 1 μg of anti-

U2AF2 (CST) or rabbit IgG (CST) in 500 μL total volume overnight at 4°C. The next day, 50 μL

of Protein A/G Dynabeads (Pierce) were added to the mix and incubated for 2 h at 4°C. Beads

were then washed 4 times with RIPA buffer and RNA was isolated using TRIzol. Equal volumes

of extracted RNA were used for cDNA synthesis and RT-qPCR as described above.

Immunoblotting

For immunoblotting, total cell lysate was prepared using RIPA buffer (Thermo Scientific)

containing protease inhibitor cocktail (Roche). Protein concentration was determined using the

BCA protein quantitation kit (Thermo Scientific). Cell lysate was loaded onto SDS-PAGE gel and

transferred to a PVDF membrane (Thermo Scientific) using a semi-dry apparatus (Bio-Rad). The

membrane was blocked in 5% skim milk (Millipore) in TBS containing 0.05% Tween 20. The

following antibodies were used: anti-U2AF2 (1:1000, rabbit) (CST, 70471S), anti-GAPDH

(1:10,000, rabbit) (CST, 2118L) and secondary anti-rabbit (1:5000) (CST, 7074S).

Nuclear cytoplasmic fractionation

For nuclear cytoplasmic fractionation, SKHEP1 cells were harvested after trypsinization at 80-

90% confluence. Cells were pelleted and resuspended in 500 μL RSB buffer (10 mM Tris-HCl,

100 mM NaCl, and 2.5 mM MgCl2) containing 40 μg/mL digitonin (Invitrogen), and 100 U of

RNase Out. After incubation on ice for 10 min, cell viability was tested with trypan blue to make

sure that the cells were permeabilized. The cytoplasmic fraction was harvested as supernatant after

centrifugation at $1,500 \times g$ for 5 min at 4°C. The nuclear pellet was resuspended in 100 µL RSB

buffer. 100 µL of the cytoplasmic fractionation were transferred to a fresh tube and 900 µL of

TRIzol was added as well as to the nuclear extracts for RNA extraction and subsequent cDNA

synthesis and RT-qPCR.

Cell proliferation assay

531 The cell proliferation assay was conducted as previously described (67). To measure proliferation,

532 parental SKHEP1 cells and *PURPL*-CRISPRi cells overexpressing *PURPL* containing intron 2

533 were seeded at $1x10^{\wedge 5}$ cells per well in 6-well plates in the presence of 1 µg/mL doxycycline. After

534 3 days, the cell numbers and viability were measured. The cells were then passaged at $1x10^5$ and

535 measured again at 6 days, and cumulative cell numbers were plotted.

RNA-seq

The TruSeq Stranded mRNA libraries were pooled and sequenced on one NovaSeq6000 S1

flowcell using 2x101 cycles for pair-end run. The Real Time Analysis software (RTA v3.4.4) was

used for processing raw data files, the Illumina bcl2fastq v2.20 was used to demultiplex and

convert binary base calls and quality scores to fastq format. The sequencing reads were trimmed

of adapters and low-quality bases using Cutadapt (v1.18). The trimmed reads were mapped to

human reference genome (hg38) and Gencode annotation GENCODE_v30 GTF using STAR

aligner (version 2.7.0f) with two-pass alignment option. RSEM (v1.3.1) was used for gene and

transcript quantification based on GENCODE annotation file. Palantir Foundry was used for

analysis of Differential Expression of Genes inside the secure NIH Integrated Data Analysis

Platform (NIDAP). The DEG Analysis results are shown in Table S3.

Iso-Seq

The libraries were prepared using PacBio standard Iso-seq library prep using SMRTbell prep kit

3.0. The libraries were sequenced on PacBio Sequel II platform using PacBio v2.0 chemistry.

PacBio SMRTlink raw subreads were converted into HiFi circular consensus sequences (CCS)

with minimum 3 passes. PacBio IsoSeq v3 pipeline was used to demultiplex the barcodes and

remove primers. Additional refine steps included trimming poly-A tails and removing concatemers

to generate Full Length Non-Concatemer (FLNC) reads. The FLNC reads were used to map to

human reference (hg38) using Minimap2 software to generate alignment bam files. Hierarchical

clustering and merging steps were performed to obtain consensus isoforms and the full-length (FL)

consensus sequences. The high-quality FL transcripts were mapped to the reference genome by

using the minimap2 software. Isoform classification and quality control were done using

SQANTI3 software and Illumina short-read gene expression data was combined with PacBio Iso-

seq for transcript quantification. The Iso-Seq Analysis results are shown in Table S4. The resulting

bam files from the Iso-seq analysis were used for plotting and visualization with the R package

Gviz (https://bioconductor.org/packages/release/bioc/html/Gviz.html).

IRFinder

IRFinder (52) was used to identify intron retention events that changed significantly on U2AF2

knockdown. A docker image was downloaded from Docker Hub (docker://cloxd/irfinder:2.0.1).

The reference was built using pre-downloaded resources (IRFinder BuildRefProcess), using

Ensembl release 92 and the 100 bp mappability file provided by IRFinder. The Build-BED-refs.sh

script was adjusted so that transcript biotype was not restricted to "processed_transcript" or

"protein_coding". Intron retention was quantified using IRFinder in FastQ mode with adapter

trimming. Differential intron retention events between siU2AF2 vs. siCTRL were identified with

IRFinder Diff using DESeq2 with the default IR ratio ≥ 0.05 in at least one sample and no warning

filters, which profiled 16,859 unique introns.

Construct generation

For Py-tract mutants, gene fragments were purchased from Twist Biosciences and cloned into the

pLCHKO-puro vector containing the WT sequences of *PURPL* (Figure 1A), between sites for

NseI and AfeI. The *PURPL* transcript containing intron 2 was cloned into the pCW57 vector

(Hygro) (Addgene #80922) in SalI and MluI sites. *MALAT1* transcripts were cloned into the

pCW57 vector (BSD) (Addgene #80921) in SalI and MluI sites. The *PURPL* insert was generated

by Twist Biosciences and the *MALAT1* sequence was amplified using Phusion DNA polymerase

(NEB). For deletion mutants, we used the Q5® Site-Directed Mutagenesis Kit (NEB). The primers

used for cloning are listed in Table S5.

Generation of *MALAT1* KO cell lines

For HCT116 *MALAT1* knockout, we used integration by non-homologous end joining, which was

accomplished by introducing a simultaneous double-strand break in genomic DNA and in the

targeting vectors (pCMV-Puro and pCMV-Hygro) (68,69). Plasmids encoding spCas9 and

sgRNAs were obtained from Addgene (Plasmids #41815 and #47108). Oligonucleotides for

construction of sgRNAs were obtained from Integrated DNA Technologies, hybridized,

phosphorylated and cloned into the sgRNA plasmid or targeting vector using BbsI sites (70).

Target sequences for sgRNAs are provided in Table S5. These vectors were transfected into

HCT116 using Lipofectamine 2000 following recommendations from the manufacturer. Three

days after transfection, the cells were selected with Puromycin (0.5 μg/mL) and Hygromycin (100

μg/mL) to generate clonal populations. Genomic DNA from each clone was isolated using DNEasy

Blood and Tissue Kit (Qiagen). PCRs to detect integration of the targeting vector at the target site

were performed using KAPA2G Robust PCR kits (Kapa Biosystems) according to the

manufacturer's instructions. A typical reaction contained 20-100 ng of genomic DNA in Buffer A

(5 µL), Enhancer (5 µL), dNTPs (0.5 µL), primers forward (PINCR Det FP, 1.25 µL) and reverse

(Targeting vector Det RP, 1.25 µL) and KAPA2G Robust DNA Polymerase (0.5 U). The DNA

sequences of the primers for each target are provided in Table S5. PCR products were visualized

in 2% agarose gels and images were captured using a ChemiDoc-It2 (UVP).

To generate MDA-MB-231 whole-locus and intron 2 *MALAT1* knockout clones, we used

the CRIPSR/Cas9 system by transfecting the cells with gRNAs and purified Cas9. 2 µL of each

30 µM gRNA was mixed with 1 µL (~5 µg) of purified Cas9 and 100 µL of 4D-Nucleofector® X

SE buffer SE from the Cell Line Kit (Lonza) supplemented with 18.5 µL of SE supplement per

100 µL total buffer volume. The mixture was incubated at room temperature for 10 min and

1×10^6 cells were resuspended in nucleofection cuvettes. Nucleofections were carried out using

the 4D-Nucleofector® System (Lonza). The cells were transferred to a 6-well plate, and they were

single-cell sorted 48 h later. The clones were grown for 3-4 weeks before genomic DNA extraction

and genotyping with PCR using the primers in Table S5.

PURPL and *MALAT1* overexpression and transduction

PURPL intron 2 constructs were transduced in previously generated *PURPL*-CRISPRi SKHEP1

cells (11). *MALAT1*-WT and deletion mutants were transduced in the two HCT116-*MALAT1* KO

clones. For lentivirus generation, HEK293T cells were seeded at 2×10^5 cells/well onto 6-well

plates, and 1,200 ng of each construct was transfected with lentiviral package vectors using

Lipofectamine 2000 (Life Technologies, Invitrogen) according to the manufacturer's protocol.

Virus-containing media were harvested at 48 and 72 h post-transfection and added to cells for 24

h. Transduced cells were selected with 500 μg/mL hygromycin for 2 weeks (SKHEP1) or 10

μg/mL Blasticidin (HCT116).

RNA-FISH, IF and Quantification of FISH

The *PURPL* intron probe set was custom-designed using the Stellaris Probe Designer, tagged with

Quasar 570 dye and purchased from Stellaris. The *MALAT1* smFISH probe set with Quasar 670

dye was purchased from Stellaris (Cat# VSMF-2211-5). For *PURPL* intron smRNA-FISH,

SKHEP1 cells were seeded on coverslips and cultured for 24 hours at 37°C with 5% CO_2 before

inducing DNA damage by incubating the cells with 2 mM hydroxyurea for 24 hours (or regular

growth media for control). For *MALAT1* smRNA-FISH, HCT116 and MDA-MB-231 *MALAT1*

KO cells were seeded on coverslips and cultured for 24 hours at 37°C with 5% CO_2. *MALAT1* WT,

MALAT1 del-1, *MALAT1* del-2, *MALAT1* del1+2, or empty vector expression was induced by

incubating the cells with 1 μg/mL doxycycline for 48 hours. The cells that were transfected with

siRNA for U2AF2 were seeded on coverslips at the time of transfection. Cells were fixed with 4%

PFA for 15 min at room temperature, permeabilized with 0.5% Triton X-100, and washed with

washing buffer (10% formamide, 2XSSC) for 5 min. Probe was added to hybridization buffer

(10% formamide in Stellaris hybridization buffer Cat# SMF-HB1-10) at a final concentration of

125 nM. Hybridization was done in a humidified chamber in the dark overnight at 37°C. After

hybridization, the coverslips were washed twice with wash buffer for 30 min at 37°C and then

post-fixed with 4% PFA for 10 min at room temperature. The coverslips were blocked with 5%

normal goat serum and then incubated with SON antibody (Sigma Cat# HPA023535). Primary

and secondary antibodies were diluted in 1% normal goat serum. Antibody incubations were done

at room temperature for 1 h and washed with PBS. DNA was counterstained by DAPI. The

coverslips were then washed with 4XSSC for 5 min at room temperature and mounted in

VectaShield Antifade Mounting Medium (Vector Laboratories). Images were taken using

DeltaVision microscope (GE Healthcare) equipped with 60 X/1.42 NA oil immersion objective

(Olympus) and CoolSNAP-HQ2 camera. To quantify the speckle to nucleoplasm signal intensity

ratio of *MALAT1* in Figures 5B, 7B, 7D, and S8B ImageJ was used to create nucleus masks based

on DAPI and speckles mask based on SON. The signal intensity of *MALAT1* in the speckles mask

was divided by the *MALAT1* signal intensity in the nucleus-minus-speckles area. For the

quantification of Figures 6B, 7E and S10B, the cells with *MALAT1* FISH signal showing

prominent speckle patterns similar to SON were classified as "Speckle" pattern, and the cells with

a diffused *MALAT1* FISH signal were classified as "Diffused". Only *MALAT1*-positive cells were

counted for quantification.

Transwell cell migration assays

For the transwell assay, WT and KO MDA-MB-231 clones were seeded at 10×10^5 cells per well

in 12-well plates. 24 h later, the media were removed and replaced with media containing 1% FBS.

After 5 h, the cells were trypsinized and seeded in the top chamber of the transwell plates (Corning,

Catalog#354578) at 25,000 cells per well at a volume of 100 μL. The lower chamber of the

transwell is filled with 700 μL of media containing 10% FBS. The cells were incubated for 24 h

and the transwell were removed from the plates. The non-migrated cells were scraped with a cotton

swab, and the migrated cells were fixed with ice-cold methanol for 5 min and stained with 0.05%

crystal violet blue. The migrated cells were counted under a light microscope.

Statistics

Statistics were performed using the Student's t-test.

Data availability

The Iso-Seq data files have been deposited at GEO with accession number GSE288684 and

reviewer token: ktcnasayrbgnjev (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi). The RNA-

Seq data files have been deposited at GEO with accession number GSE288685 and reviewer token:

qvwhuawixxqjnml (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi).

Acknowledgments

This research was supported by the Intramural Research Program (A.L.) of the National Cancer

Institute (NCI), Center for Cancer Research (CCR) (project numbers ZIA BC011646, ZIA

BC012019), NIH. KVP laboratory is supported by grants from NIH (R01-GM132458), Cancer

center at Illinois seed grant, ARPA-H, and National Science Foundation (NSF) center for

quantitative cell biology. YJS is supported by NIH T32EB019944. We thank the CCR Genomics

Core, CCR, NCI, Bethesda, MD for its valuable assistance with Sanger and Nanopore sequencing.

We also thank the Sequencing Facility, CCR, NCI, Frederick, MD for performing the RNA

sequencing assays. The contributions of the NIH author(s) were made as part of their official duties

as NIH federal employees, are in compliance with agency policy requirements, and are considered

Works of the United States Government. However, the findings and conclusions presented in this

paper are those of the author(s) and do not necessarily reflect the views of the NIH or the U.S.

Department of Health and Human Services.

Author Contributions

IG, CN, YJS, CCH, SK, RP, XLL, RK, RS, and PPZ conducted experiments. IG, AKB, EP, YZ,

BS, TB, XW, TGP, KP, and AL did data analysis. IG and AL wrote the manuscript. NC, KP, and

TGP edited the manuscript.

Declaration of interests

The authors declare they have no competing interest.

References

1.　Quinn, J.J. and Chang, H.Y. (2016) Unique features of long non-coding RNA biogenesis and
function. *Nat Rev Genet*, **17**, 47-62.
2.　Ulitsky, I. and Bartel, D.P. (2013) lincRNAs: genomics, evolution, and mechanisms. *Cell*, **154**, 26-
46.
3.　Guo, C.J., Xu, G. and Chen, L.L. (2020) Mechanisms of Long Noncoding RNA Nuclear Retention.
Trends Biochem Sci, **45**, 947-960.
4.　Zuckerman, B. and Ulitsky, I. (2019) Predictive models of subcellular localization of long RNAs.
RNA, **25**, 557-572.
5.　Clark, M.B., Johnston, R.L., Inostroza-Ponta, M., Fox, A.H., Fortini, E., Moscato, P., Dinger, M.E.
and Mattick, J.S. (2012) Genome-wide analysis of long noncoding RNA stability. *Genome Res*, **22**,
885-898.
6.　Cabili, M.N., Dunagin, M.C., McClanahan, P.D., Biaesch, A., Padovan-Merhar, O., Regev, A., Rinn,
J.L. and Raj, A. (2015) Localization and abundance analysis of human lncRNAs at single-cell and
single-molecule resolution. *Genome Biol*, **16**, 20.
7.　Tilgner, H., Knowles, D.G., Johnson, R., Davis, C.A., Chakrabortty, S., Djebali, S., Curado, J.,
Snyder, M., Gingeras, T.R. and Guigo, R. (2012) Deep sequencing of subcellular RNA fractions
shows splicing to be predominantly co-transcriptional in the human genome but inefficient for
lncRNAs. *Genome Res*, **22**, 1616-1625.
8.　Mele, M., Mattioli, K., Mallard, W., Shechner, D.M., Gerhardinger, C. and Rinn, J.L. (2017)
Chromatin environment, transcriptional regulation, and splicing distinguish lincRNAs and
mRNAs. *Genome Res*, **27**, 27-37.
9.　Kumar, M. and Grammatikakis, I. (2025) Causes, Consequences and Challenges of Intron
Retention in lncRNAs. *Mol Cell Biol*, **45**, 576-584.
10.　Mukherjee, N., Calviello, L., Hirsekorn, A., de Pretis, S., Pelizzola, M. and Ohler, U. (2017)
Integrative classification of human coding and noncoding genes through RNA metabolism
profiles. *Nat Struct Mol Biol*, **24**, 86-96.
11.　Hartford, C.C.R., Shrestha, R.L., Pongor, L., Zhao, Y., Chen, X., Fromont, C., Chaudhary, R., Li, X.L.,
Pasterczyk, K.R., Kumar, R. *et al.* (2022) Context-Dependent Function of Long Noncoding RNA
PURPL in Transcriptome Regulation during p53 Activation. *Mol Cell Biol*, **42**, e0028922.
12.　Black, D.L. (2003) Mechanisms of alternative pre-messenger RNA splicing. *Annu Rev Biochem*,
72, 291-336.
13.　Wilkinson, M.E., Charenton, C. and Nagai, K. (2020) RNA Splicing by the Spliceosome. *Annu Rev*
Biochem, **89**, 359-388.
14.　Ruskin, B., Zamore, P.D. and Green, M.R. (1988) A factor, U2AF, is required for U2 snRNP binding
and splicing complex assembly. *Cell*, **52**, 207-219.
15.　Berglund, J.A., Chua, K., Abovich, N., Reed, R. and Rosbash, M. (1997) The splicing factor BBP
interacts specifically with the pre-mRNA branchpoint sequence UACUAAC. *Cell*, **89**, 781-787.
16.　Berglund, J.A., Abovich, N. and Rosbash, M. (1998) A cooperative interaction between U2AF65
and mBBP/SF1 facilitates branchpoint region recognition. *Genes Dev*, **12**, 858-867.
17.　Wu, S., Romfo, C.M., Nilsen, T.W. and Green, M.R. (1999) Functional recognition of the 3' splice
site AG by the splicing factor U2AF35. *Nature*, **402**, 832-835.
18.　Zorio, D.A. and Blumenthal, T. (1999) Both subunits of U2AF recognize the 3' splice site in
Caenorhabditis elegans. *Nature*, **402**, 835-838.

19. Wang, E.T., Sandberg, R., Luo, S., Khrebtukova, I., Zhang, L., Mayr, C., Kingsmore, S.F., Schroth,
G.P. and Burge, C.B. (2008) Alternative isoform regulation in human tissue transcriptomes.
Nature, **456**, 470-476.
20. Nilsen, T.W. and Graveley, B.R. (2010) Expansion of the eukaryotic proteome by alternative
splicing. *Nature*, **463**, 457-463.
21. Pan, Q., Shai, O., Lee, L.J., Frey, B.J. and Blencowe, B.J. (2008) Deep surveying of alternative
splicing complexity in the human transcriptome by high-throughput sequencing. *Nat Genet*, **40**,
1413-1415.
22. Gier, R.A., Budinich, K.A., Evitt, N.H., Cao, Z., Freilich, E.S., Chen, Q., Qi, J., Lan, Y., Kohli, R.M. and
Shi, J. (2020) High-performance CRISPR-Cas12a genome editing for combinatorial genetic
screening. *Nat Commun*, **11**, 3455.
23. Braunschweig, U., Barbosa-Morais, N.L., Pan, Q., Nachman, E.N., Alipanahi, B., Gonatopoulos-
Pournatzis, T., Frey, B., Irimia, M. and Blencowe, B.J. (2014) Widespread intron retention in
mammals functionally tunes transcriptomes. *Genome Res*, **24**, 1774-1786.
24. Wong, J.J., Ritchie, W., Ebner, O.A., Selbach, M., Wong, J.W., Huang, Y., Gao, D., Pinello, N.,
Gonzalez, M., Baidya, K. *et al.* (2013) Orchestrated intron retention regulates normal
granulocyte differentiation. *Cell*, **154**, 583-595.
25. Monteuuis, G., Wong, J.J.L., Bailey, C.G., Schmitz, U. and Rasko, J.E.J. (2019) The changing
paradigm of intron retention: regulation, ramifications and recipes. *Nucleic Acids Res*, **47**, 11497-
11513.
26. Eom, T., Zhang, C., Wang, H., Lay, K., Fak, J., Noebels, J.L. and Darnell, R.B. (2013) NOVA-
dependent regulation of cryptic NMD exons controls synaptic protein levels after seizure. *Elife*,
2, e00178.
27. Hossain, M.A., Claggett, J.M., Edwards, S.R., Shi, A., Pennebaker, S.L., Cheng, M.Y., Hasty, J. and
Johnson, T.L. (2016) Posttranscriptional Regulation of Gcr1 Expression and Activity Is Crucial for
Metabolic Adjustment in Response to Glucose Availability. *Mol Cell*, **62**, 346-358.
28. Cha, S., Hong, C.P., Kang, H.A. and Hahn, J.S. (2021) Differential activation mechanisms of two
isoforms of Gcr1 transcription factor generated from spliced and un-spliced transcripts in
Saccharomyces cerevisiae. *Nucleic Acids Res*, **49**, 745-759.
29. Schmitz, U., Pinello, N., Jia, F., Alasmari, S., Ritchie, W., Keightley, M.C., Shini, S., Lieschke, G.J.,
Wong, J.J. and Rasko, J.E.J. (2017) Intron retention enhances gene regulatory complexity in
vertebrates. *Genome Biol*, **18**, 216.
30. Dvinge, H. and Bradley, R.K. (2015) Widespread intron retention diversifies most cancer
transcriptomes. *Genome Med*, **7**, 45.
31. Boutz, P.L., Bhutkar, A. and Sharp, P.A. (2015) Detained introns are a novel, widespread class of
post-transcriptionally spliced introns. *Genes Dev*, **29**, 63-80.
32. Gutschner, T., Hammerle, M., Eissmann, M., Hsu, J., Kim, Y., Hung, G., Revenko, A., Arun, G.,
Stentrup, M., Gross, M. *et al.* (2013) The noncoding RNA MALAT1 is a critical regulator of the
metastasis phenotype of lung cancer cells. *Cancer Res*, **73**, 1180-1189.
33. Hutchinson, J.N., Ensminger, A.W., Clemson, C.M., Lynch, C.R., Lawrence, J.B. and Chess, A.
(2007) A screen for nuclear transcripts identifies two linked noncoding RNAs associated with
SC35 splicing domains. *BMC Genomics*, **8**, 39.
34. Ji, P., Diederichs, S., Wang, W., Boing, S., Metzger, R., Schneider, P.M., Tidow, N., Brandt, B.,
Buerger, H., Bulk, E. *et al.* (2003) MALAT-1, a novel noncoding RNA, and thymosin beta4 predict
metastasis and survival in early-stage non-small cell lung cancer. *Oncogene*, **22**, 8031-8041.
35. Bernard, D., Prasanth, K.V., Tripathi, V., Colasse, S., Nakamura, T., Xuan, Z., Zhang, M.Q., Sedel,
F., Jourdren, L., Coulpier, F. *et al.* (2010) A long nuclear-retained non-coding RNA regulates
synaptogenesis by modulating gene expression. *EMBO J*, **29**, 3082-3093.

36.   Tripathi, V., Ellis, J.D., Shen, Z., Song, D.Y., Pan, Q., Watt, A.T., Freier, S.M., Bennett, C.F., Sharma,
A., Bubulya, P.A. *et al.* (2010) The nuclear-retained noncoding RNA MALAT1 regulates
alternative splicing by modulating SR splicing factor phosphorylation. *Mol Cell*, **39**, 925-938.
37.   Shinn, M.K., Tomares, D.T., Liu, V., Pant, A., Qiu, Y., Vitalis, A., Song, Y.J., Ayala, Y., Ruff, K.M.,
Strout, G.W. *et al.* (2026) Nuclear speckle proteins form intrinsic and MALAT1-dependent
microphases. *Cell*, **189**, 832-852 e824.
38.   Spector, D.L. and Lamond, A.I. (2011) Nuclear speckles. *Cold Spring Harb Perspect Biol*, **3**.
39.   Sharma, A., Takata, H., Shibahara, K., Bubulya, A. and Bubulya, P.A. (2010) Son is essential for
nuclear speckle organization and cell cycle progression. *Mol Biol Cell*, **21**, 650-663.
40.   Nakagawa, S., Ip, J.Y., Shioi, G., Tripathi, V., Zong, X., Hirose, T. and Prasanth, K.V. (2012) Malat1
is not an essential component of nuclear speckles in mice. *RNA*, **18**, 1487-1499.
41.   Zhang, B., Arun, G., Mao, Y.S., Lazar, Z., Hung, G., Bhattacharjee, G., Xiao, X., Booth, C.J., Wu, J.,
Zhang, C. *et al.* (2012) The lncRNA Malat1 is dispensable for mouse development but its
transcription plays a cis-regulatory role in the adult. *Cell Rep*, **2**, 111-123.
42.   Li, X.L., Subramanian, M., Jones, M.F., Chaudhary, R., Singh, D.K., Zong, X., Gryder, B., Sindri, S.,
Mo, M., Schetter, A. *et al.* (2017) Long Noncoding RNA PURPL Suppresses Basal p53 Levels and
Promotes Tumorigenicity in Colorectal Cancer. *Cell Rep*, **20**, 2408-2423.
43.   Behera, A.K., Kim, J.J., Kordale, S., Pekovic, F., Damodaran, A.P., Kumari, B., Vidak, S., Dickson, E.,
Xiao, M.S., Duncan, G. *et al.* (2026) RNA-coupled CRISPR screens reveal ZNF207 as a regulator of
LMNA aberrant splicing in progeria. *Mol Cell*, **86**, 41-59 e15.
44.   Aregger, M., Xing, K. and Gonatopoulos-Pournatzis, T. (2021) Application of CHyMErA Cas9-
Cas12a combinatorial genome-editing platform for genetic interaction mapping and gene
fragment deletion screening. *Nat Protoc*, **16**, 4722-4765.
45.   Gonatopoulos-Pournatzis, T., Aregger, M., Brown, K.R., Farhangmehr, S., Braunschweig, U.,
Ward, H.N., Ha, K.C.H., Weiss, A., Billmann, M., Durbic, T. *et al.* (2020) Genetic interaction
mapping and exon-resolution functional genomics with a hybrid Cas9-Cas12a platform. *Nat
Biotechnol*, **38**, 638-648.
46.   Xiao, M.S., Damodaran, A.P., Kumari, B., Dickson, E., Xing, K., On, T.A., Parab, N., King, H.E.,
Perez, A.R., Guiblet, W.M. *et al.* (2024) Genome-scale exon perturbation screens uncover exons
critical for cell fitness. *Mol Cell*, **84**, 2553-2572 e2519.
47.   Van Nostrand, E.L., Pratt, G.A., Shishkin, A.A., Gelboin-Burkhart, C., Fang, M.Y., Sundararaman,
B., Blue, S.M., Nguyen, T.B., Surka, C., Elkins, K. *et al.* (2016) Robust transcriptome-wide
discovery of RNA-binding protein binding sites with enhanced CLIP (eCLIP). *Nat Methods*, **13**,
508-514.
48.   Consortium, E.P. (2012) An integrated encyclopedia of DNA elements in the human genome.
Nature, **489**, 57-74.
49.   Luo, Y., Hitz, B.C., Gabdank, I., Hilton, J.A., Kagda, M.S., Lam, B., Myers, Z., Sud, P., Jou, J., Lin, K.
et al. (2020) New developments on the Encyclopedia of DNA Elements (ENCODE) data portal.
Nucleic Acids Res, **48**, D882-D889.
50.   Berhane, T., Holm, A., Karstensen, K.T., Petri, A., Ilieva, M.S., Krarup, H., Vyberg, M., Lovendorf,
M.B. and Kauppinen, S. (2022) Knockdown of the long noncoding RNA PURPL induces apoptosis
and sensitizes liver cancer cells to doxorubicin. *Sci Rep*, **12**, 19502.
51.   Han, S., Li, X., Wang, K., Zhu, D., Meng, B., Liu, J., Liang, X., Jin, Y., Liu, X., Wen, Q. *et al.* (2021)
PURPL represses autophagic cell death to promote cutaneous melanoma by modulating ULK1
phosphorylation. *Cell Death Dis*, **12**, 1070.
52.   Middleton, R., Gao, D., Thomas, A., Singh, B., Au, A., Wong, J.J., Bomane, A., Cosson, B., Eyras, E.,
Rasko, J.E. *et al.* (2017) IRFinder: assessing the impact of intron retention on mammalian gene
expression. *Genome Biol*, **18**, 51.

53. Martinez-Terroba, E., Plasek-Hegde, L.M., Chiotakakos, I., Li, V., de Miguel, F.J., Robles-Oteiza,
C., Tyagi, A., Politi, K., Zamudio, J.R. and Dimitrova, N. (2024) Overexpression of Malat1 drives
metastasis through inflammatory reprogramming of the tumor microenvironment. *Sci Immunol*,
9, eadh5462.
54. Arun, G., Aggarwal, D. and Spector, D.L. (2020) MALAT1 Long Non-Coding RNA: Functional
Implications. *Noncoding RNA*, **6**.
55. Jacob, A.G. and Smith, C.W.J. (2017) Intron retention as a component of regulated gene
expression programs. *Hum Genet*, **136**, 1043-1057.
56. Shao, C., Yang, B., Wu, T., Huang, J., Tang, P., Zhou, Y., Zhou, J., Qiu, J., Jiang, L., Li, H. *et al.*
(2014) Mechanisms for U2AF to define 3' splice sites and regulate alternative splicing in the
human genome. *Nat Struct Mol Biol*, **21**, 997-1005.
57. Naftelberg, S., Schor, I.E., Ast, G. and Kornblihtt, A.R. (2015) Regulation of alternative splicing
through coupling with transcription and chromatin structure. *Annu Rev Biochem*, **84**, 165-198.
58. Herzel, L., Ottoz, D.S.M., Alpert, T. and Neugebauer, K.M. (2017) Splicing and transcription touch
base: co-transcriptional spliceosome assembly and function. *Nat Rev Mol Cell Biol*, **18**, 637-650.
59. Wong, J.J., Gao, D., Nguyen, T.V., Kwok, C.T., van Geldermalsen, M., Middleton, R., Pinello, N.,
Thoeng, A., Nagarajah, R., Holst, J. *et al.* (2017) Intron retention is regulated by altered MeCP2-
mediated splicing factor recruitment. *Nat Commun*, **8**, 15134.
60. Guo, R., Zheng, L., Park, J.W., Lv, R., Chen, H., Jiao, F., Xu, W., Mu, S., Wen, H., Qiu, J. *et al.* (2014)
BS69/ZMYND11 reads and connects histone H3.3 lysine 36 trimethylation-decorated chromatin
to regulated pre-mRNA processing. *Mol Cell*, **56**, 298-310.
61. Krchnakova, Z., Thakur, P.K., Krausova, M., Bieberstein, N., Haberman, N., Muller-McNicoll, M.
and Stanek, D. (2019) Splicing of long non-coding RNAs primarily depends on polypyrimidine
tract and 5' splice-site sequences due to weak interactions with SR proteins. *Nucleic Acids Res*,
47, 911-928.
62. Dumbovic, G., Braunschweig, U., Langner, H.K., Smallegan, M., Biayna, J., Hass, E.P., Jastrzebska,
K., Blencowe, B., Cech, T.R., Caruthers, M.H. *et al.* (2021) Nuclear compartmentalization of TERT
mRNA and TUG1 lncRNA is driven by intron retention. *Nat Commun*, **12**, 3308.
63. Desideri, F., Cipriano, A., Petrezselyova, S., Buonaiuto, G., Santini, T., Kasparek, P., Prochazka, J.,
Janson, G., Paiardini, A., Calicchio, A. *et al.* (2020) Intronic Determinants Coordinate Charme
lncRNA Nuclear Activity through the Interaction with MATR3 and PTBP1. *Cell Rep*, **33**, 108548.
64. Guo, C.J., Ma, X.K., Xing, Y.H., Zheng, C.C., Xu, Y.F., Shan, L., Zhang, J., Wang, S., Wang, Y.,
Carmichael, G.G. *et al.* (2020) Distinct Processing of lncRNAs Contributes to Non-conserved
Functions in Stem Cells. *Cell*, **181**, 621-636 e622.
65. Barutcu, A.R., Wu, M., Braunschweig, U., Dyakov, B.J.A., Luo, Z., Turner, K.M., Durbic, T., Lin,
Z.Y., Weatheritt, R.J., Maass, P.G. *et al.* (2022) Systematic mapping of nuclear domain-associated
transcripts reveals speckles and lamina as hubs of functionally distinct retained introns. *Mol Cell*,
82, 1035-1052 e1039.
66. Kolberg, L., Raudvere, U., Kuzmin, I., Adler, P., Vilo, J. and Peterson, H. (2023) g:Profiler-
interoperable web service for functional enrichment analysis and gene identifier mapping (2023
update). *Nucleic Acids Res*, **51**, W207-W212.
67. Lee, J.S., Dan, T., Zhang, H., Cheng, Y., Rehfeld, F., Brugarolas, J. and Mendell, J.T. (2025) An
ultraconserved snoRNA-like element in long noncoding RNA CRNDE promotes ribosome
biogenesis and cell proliferation. *Mol Cell*, **85**, 1543-1560 e1510.
68. Brown, A., Winter, J., Gapinske, M., Tague, N., Woods, W.S. and Perez-Pinera, P. (2019)
Multiplexed and tunable transcriptional activation by promoter insertion using nuclease-assisted
vector integration. *Nucleic Acids Res*, **47**, e67.

69. Brown, A., Woods, W.S. and Perez-Pinera, P. (2016) Multiplexed Targeted Genome Engineering
Using a Universal Nuclease-Assisted Vector Integration System. *ACS Synth Biol*, **5**, 582-588.
70. Brown, A., Woods, W.S. and Perez-Pinera, P. (2017) Targeted Gene Activation Using RNA-Guided
Nucleases. *Methods Mol Biol*, **1468**, 235-250.

Figure legends

Figure 1. CRISPR-based screen identifies U2AF2 as a promoter of intron retention. (A)

Schematic of the *PURPL* IR reporter used in the screen to identify IR regulators. The pLCHKO

vector contains the puromycin resistance gene used for selection. The transgene also contains the

genomic sequences of exon 2 (161 bp), intron 2 (1,943 bp), and exon 3 (163 bp) of the *PURPL*

gene under a Doxycycline-inducible promoter. A hybrid guide (hg)RNA for guiding *Sp*Cas9 and

*Lb*Cas12a is expressed under a U6 promoter. Top arrow: The hgRNA is cleaved by Cas12a to

produce 2 guide RNAs for dual targeting of each protein coding gene, paralog targeting, and

genetic interaction mapping. Bottom arrow: Upon intron splicing or intron retention, two different

reads are generated. **(B)** Two different cell lines, stably expressing Cas9 and Cas12a, are used:

HAP1, and RPE1. The CRASP-seq pipeline until Illumina paired-end sequencing of CRISPR

libraries is depicted. **(C)** Heatmap showing single genes only (after removing gene pairs), whether

identified individually or as part of a gene pair. The hits are ordered by the average ΔPIR values

across HAP1 and RPE1, from lowest to highest. **(D)** Left: Gene ontology enrichment analysis of

biological processes regulated by the hits of the screen. The most significant pathways involve

mRNA processing and RNA splicing. Right: Gene ontology enrichment analysis of cellular

components regulated by the hits of the screen. These components include mainly Spliceosomal

Complex, small nuclear ribonucleoprotein complex and the Catalytic step 2 Spliceosome. **(E)**

Venn diagram showing that there are 5 common hits which cause a reduction in the percentage of

intron retention (ΔPIR) of the splicing reporter in the two cell lines. **(F)** Table showing the top 5

hits of the screen with a ΔPIR for each hit in each cell line and their full names.

Figure 2. U2AF2 binds to intron 2-containing *PURPL* transcripts and promotes intron 2

retention. (A) IGV snapshot of eCLIP-seq data showing binding sites and enrichment of U2AF2

and U2AF1 on the *PURPL* transcripts around intron 2. eCLIP-seq data for PTBP1, PRPF8, and

SRSF1 are indicated. The annotated locus by RefSeq is also indicated. eCLIP data-seq was

downloaded from encodeproject.org. **(B)** RT-qPCR after RNA-IPs using a U2AF2 antibody with

primer pairs specifically detecting *PURPL* transcripts as indicated in Figure S1B. U2AF2 binds to

transcripts containing intron 2 but not the ones with intron 1, intron 3, or spliced exons 2 and 3.

Samples were normalized to IgG-IP. 18S was used as a loading control. **(C)** Validation of hgRNAs

using the *PURPL* minigene in HAP1 and RPE1 cells. The gels show RT-PCR products for *PURPL*

transcripts with specific primers for the minigene. The hgRNAs used were targeting intergenic

region, or U2AF2. The schematics next to the gel indicate the expected products of the intron-

retained and spliced isoforms. **(D)** Western blot for U2AF2 showing successful knockdown of

U2AF2 protein in SKHEP1 cells. GAPDH was used as a loading control (lower panel). **(E)**

Schematic of the PCR primer triplet used to detect intron 2 retention (red) or splicing (purple). The

length for each PCR product is indicated. **(F)** Gel with RT-PCR products for *PURPL* upon

knockdown of U2AF2 with 2 different siRNAs in SKHEP1 cells. The schematics next to the gel

indicate the expected products of the intron-retained and spliced isoforms. Between the two

expected PCR products, we observed an extra band corresponding to the inclusion of an alternative

exon inside *PURPL* intron 2 as observed in RefSeq, the inclusion of which is not affected by

U2AF2. **(G)** Graph with quantitation of the gel bands in **(F)**. Error bars represent standard

deviations from 3 **(G)** experiments. ***$p<0.001$.

Figure 3. Expression of Intron 2 Retaining *PURPL* leads to high proliferation. (A) RNA-FISH images for *PURPL* with intron 2 retention and *MALAT1* in HCT116 cells without treatment or after 24 h of 2 mM of Hydorxyurea (HU) to induce *PURPL* expression. Scale bar is 10 μm. **(B)** RNA stability assays were performed for *PURPL* transcripts by measuring their levels by RT-qPCR following 0, 2, and 4 h of ActD treatment in SKHEP1 cells. *MYC* was used as a positive control for unstable RNA. **(C)** RT-qPCR showing *PURPL* depletion in in SKHEP1 cells using 3 different gRNAs (sg2, sg4, sg5) compared to a Non-Targeting Control gRNA. **(D)** RT-qPCR for intron 2-containing PURPL transcript after 48 h of 1 μg/mL doxycycline treatment in comparison to no treatment in SKHEP1 *PURPL*-CRISPRi populations using the 3 gRNAs as in **(C)**. **(E)** Proliferation assay showing the effect of overexpression of intron 2-containing *PURPL* transcript in the proliferation of SKHEP1 cells where the endogenous *PURPL* is knocked down with CRISPRi. Error bars represent standard deviation from 3 populations with different gRNAs. The cells were treated with 1 μg/mL doxycycline to induce intron 2-retained *PURPL* expression and cell proliferation was monitored at 3 and 6 days. **(F)** Gel with RT-PCR products for transcripts using primers as indicated in Fig. 2E upon doxycycline treatment of SKHEP1 CRIPSRi cells. The three repeats represent 3 different clones of cells. The last lane is an RT- control. The schematics next to the gel indicate the expected products of the intron-retained and spliced isoforms. NS indicates a non-specific band. Error bars in **(B)**, **(C)**, and **(D)** represent standard deviations from 2 experiments. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

Figure 4. U2AF2 directly promotes Intron Retention of *MALAT1*. (A) U2AF2 was knocked down in SKHEP1 cells and 72 h later, RNA was extracted and RNA-seq was performed. *Left*: Number of decreased (blue) and increased (red) IR events at various p-values after U2AF2 knockdown as analyzed with the IR Finder algorithm. The purple arrow indicates the *PURPL* IR

event and green arrows indicate *MALAT1* IR events *Right*: Pie chart of the numbers of increased

and decreased IR events upon U2AF2 knockdown. **(B)** Floating bar plots showing the IR ratio of

PURPL intron 2 and the IR ratio of intron 1 (middle) and intron 2 (right) of *MALAT1* in siCTRL

and siU2AF2#1 samples as analyzed with the IRFinder algorithm. **(C)** and **(E)** RT-PCR for

MALAT1 using a primer pair flanking the regulated intron 1 **(C)** or intron 2 **(E)** upon knockdown

of U2AF2 with 2 different siRNAs in HCT116 and SKHEP1 cells. The schematics next to the gel

indicate the expected products of the intron-retained and spliced isoforms. **(D)** and **(F)** Bar graph

with quantitation of the gel bands from **(C)** and **(E)** in SKHEP1 cells. Error bars represent standard

deviations from 2 independent experiments. *p<0.05, **p<0.01, ***p<0.001.

Figure 5. U2AF2 promotes localization of *MALAT1* to nuclear speckles. (A) RNA-FISH

images for *MALAT1* and Immunofluorescence images for SON is shown upon transfection of

SKHEP1 cells with siCTRL or siU2AF2 with two different siRNAs. *MALAT1* is enriched in

nuclear speckles in the siCTRL but not upon U2AF2 knockdown. **(B)** Quantitation of the speckle

to nuclear plasma *MALAT1* signal ratio in the three replicates in panel **(A)**. ****p<0.0001.

Figure 6. Intron 2 of *MALAT1* dictates its localization to nuclear speckles. (A) RNA-FISH

images for *MALAT1* and Immunofluorescence images for SON in a HCT116 *MALAT1*-KO clone

(clone 1) where Empty Vector (EV) or constructs that exogenously express *MALAT1* transcripts

were re-introduced using a doxy-inducible lentivirus. Expressed *MALAT1* transcripts were full-

length (WT) or intron 1 deleted (del-1) or intron 2 deleted (del-2) or both introns deleted

simultaneously (del-1+2). The cells were first treated with 1 μg/mL doxycycline for 48 h to induce

MALAT1 expression. *MALAT1* without intron 2 does not enrich in speckles as it is shown with

SON protein. Scale bar is 10 μm. **(B)** Graph showing quantification of the percentage of cells

where *MALAT1* transcripts are enriched in speckles or being diffused in HCT116 cells clone 1.

Error bars represent standard deviations from 2 independent experiments.

Figure 7. Intron 2 regulates the migration capacity of *MALAT1* in breast cancer cells. (A)

RNA-FISH images for *MALAT1* and Immunofluorescence images for SON are shown upon

transfection of MDA-MB-231 cells with siCTRL or siU2AF2. *MALAT1* is enriched in nuclear

speckles in the siCTRL but not upon U2AF2 knockdown. **(B)** Quantitation of the speckle to nuclear

plasma *MALAT1* signal ratio in the three replicates in panel **(A)**. **(C)** RNA-FISH images for

MALAT1 and Immunofluorescence images for SON in MDA-MB-231 clones either WT, or whole

locus *MALAT1* deletion, or intron 2 deletion. When intron 2 is deleted, *MALAT1* does not enrich

in speckles as it is shown with SON protein. Scale bar is 10 μm. **(D)** Quantitation of the speckle to

nuclear plasma *MALAT1* signal ratio in the three replicates in panel **(C)**. **(E)** Graph showing

quantification of the percentage of *MALAT1* transcripts enriched in speckles or being diffused in

MDA-MB-231 cells in panel **(C)**, N=2. **(F)** Bar graph showing the number of cells that have

migrated in transwell migration assays conducted in MDA-MB-231 WT, *MALAT1* KO and

MALAT1 – intron 2 deletion clones. Intron 2 deletion leads to decreased migration potential. The

number of cells are the sum of cells from 5 different fields, N=4. *p<0.05, **p<0.01, ***p<0.001,

****p<0.0001.

Figure 1

A

- Dual targeting of each protein coding gene
- Paralogue targeting
- GI mapping

combinatorial genome editing

Cas9 guide | Cas12a guide
hybrid guide (hg)RNA

B

HAP1   RPE1

1. Total RNA extraction

2. oligo-dT RNA purification

3. cDNA synthesis (UMI incorporation)

4. PCR amplification (Dual indexes incorporation)

5. RNA-sequencing and analysis

C

ΔPIR
−20 −10 0 10 20

RPE1
HAP1

D

Gene Ontology Biological Process

mRNA processing
RNA splicing
mRNA splicing, via spliceosome
protein-RNA complex organization
spliceosomal complex assembly
U2-type prespliceosome assembly
ribonucleoprotein complex biogenesis
regulation of RNA splicing
protein-RNA complex assembly
RNA 3'-end processing

-log10 (FDR)

Gene Ontology Cellular Component

spliceosomal complex
small nuclear ribonucleoprotein complex
catalytic step 2 spliceosome
U2-type spliceosomal complex
spliceosomal snRNP complex
nuclear speckles
precatalytic spliceosome
U2 snRNP
mRNA cleavage factor complex

-log10 (FDR)

E

HAP1 **45**   **5**   RPE1 **12**

F

HIT	ΔPIR HAP1	ΔPIR RPE1	Name
U2AF2	-28.42	-23.98	U2 Small Nuclear RNA Auxiliary Factor 2
SNRNP70	-16.05	-3.69	Small Nuclear Ribonucleoprotein U1 Subunit 70
EFTUD2	-14.64	-12.84	Elongation Factor Tu GTP Binding Domain Containing 2
DDX23	-9.49	-6.28	DEAD-Box Helicase 23
SART3	-2.01	-14.01	Spliceosome Associated Factor 3, U4/U6 Recycling Protein

Figure 2

Figure 3

Figure 4

A

SKHEP1

B

Figure 7